1 Do epidemiologically impactful vector control interventions disrupt mosquito
2 population structure?
3

4 Tristan P. W. Dennis[1,2*], W. Moussa Guelbeogo[3,4], Heather M. Ferguson[1], Steve Lindsay[5],
5 Sagnon N'Fale[3], Patricia Pignatelli[2], Hilary Ranson[2], Antoine Sanou[3,6], Alfred Tiono[3,7], David
6 Weetman[2] & Mafalda Viana[1]
7
[1]School of Biodiversity, One Health, and Veterinary Medicine, University of Glasgow, Glasgow,
UK
[2]Dept. Vector Biology, Liverpool School of Tropical Medicine, Liverpool, UK
[3]Centre National de Recherche et de Formation sur le Paludisme, Burkina Faso
[4]Université Joseph KI-ZERBO, Ouagadougou, Burkina Faso
[5]Department of Biosciences, Durham University, Durham, UK
[6]Université Yembila Abdoulaye Toguyeni, Fada N'Gourma, Burkina Faso
[7]Groupe de Recherche Action en Santé, Burkina Faso
16
17 *Corresponding author

18
19
Abstract

Large epidemiological impacts resulting from vector control interventions are typically
associated with significant perturbation of vector populations. While vector density is a
frequently measured response, impacts on population demography and connectivity are
suspected but rarely quantified. As population fragmentation can compromise metapopulation
viability, the type of intervention and effort required to control or eliminate vector populations will
fundamentally change. We analysed whole-genome sequences of 893 malaria *Anopheles*
gambiae mosquitoes collected at 7 time points between 2024-2015 during a clustered
randomized control trial (cRCT) in Burkina Faso. This cRCT compared a pyrethroid-only net
(ITN) with a pyrethroid-pyriproxyfen (ITN-PPF) net. Although the ITN-PPF reduced clinical
malaria by 12% and vector density by 22% compared to ITN, our results revealed no significant
changes in *An. gambiae* population genetic structure or diversity. Analysis showed remarkably
low population differentiation (mean *Fst*: -0.00073, 95% CI: -0.00109, 0.00013) and no
discernible clustering by treatment, time, or space. Metrics of population diversity (nucleotide
diversity and inbreeding coefficient) remained stable between treatments, and genome-wide
scans showed no putative signatures of selection between trial arms. Together, these results
show that ITN-PPF did not alter *An. gambiae* genetic structure, possibly due to dispersal within
this highly mobile vector species known to have large population sizes across West Africa. More
widely, our findings provide the first evidence that reductions in vector density that are
epidemiologically meaningful may not impact genetic diversity or connectivity and challenges
what constitutes adequate vector control and the drivers of population heterogeneity in large
populations.
42
43 **Keywords:** Vector populations, genetics, disease vectors, malaria, cluster-randomised
44 controlled trial, epidemiology

Introduction

Vector control is the cornerstone of the public health response to many vector-borne diseases (VBD)[1]. Effectiveness is typically assessed by the epidemiological impacts vector control generates, such as reduction in the prevalence or incidence of infection[2,3]. These epidemiological impacts are tied to reductions in human exposure to vector populations[4], which can arise through reductions in vector population size, adult mosquito survival or infection rates. Notable examples of vector control strategies that have large epidemiological impacts include the deployment of spatial repellents that reduced arboviral infection by 34.1% and *Aedes* mosquito abundance by 38.6% in Peru[5] and release of *Wolbachia*-infected *Aedes aegypti* that resulted in a 78% reduction in dengue incidence in a trial in Indonesia[6]. For malaria, the deadliest VBD, responsible for over 600,000 deaths annually[7], the effects of vector control have been profound, with a 40% reduction in clinical disease incidence across sub-Saharan Africa between 2000 and 2015[8] as a result of mass insecticide treated bednet (ITN) distribution. Successful malaria vector control has been associated with broader changes in mosquito vector demography and ecology, including changes in biting and resting behaviour[9,10], species composition[11] and the emergence of insecticide resistance[12]. For example, vector control has reduced *Anopheles gambiae* populations over time in East Africa, leading to *An. arabiensis* and *An. funestus* replacing it as the dominant vectors of malaria[13,14]. These examples suggest that that large epidemiological impacts are linked to, and may typically require, significant perturbation of mosquito vector populations, including reducing population size[15] or fragmenting population structure[16]. However, the demographic effects of interventions on vector populations are difficult to measure directly and hence remain elusive and rarely recorded.

Insect populations, such as butterflies[17–20] and gypsy moths[21,22], have provided notable case-studies of how populations are regulated, maintained and structured as well as how population fragmentation can lead to population declines by breaking connections among subpopulations[18,23–25]. However, the impact of vector control on mosquito population dynamics, in particular population structure, remains poorly understood. Vector control may change mosquito population structure by reducing local population size below the threshold for persistence, and in doing so compromise metapopulation viability, or population fitness by driving down diversity. In contrast, in the presence of migration or strong density-dependent regulation, local mosquito populations may persist even in the face of substantial reduction due to vector control[26,27]. It is thus important to understand the mechanisms underlying population

resilience, and the magnitude of reduction required to achieve an epidemiological benefit, to
improve the design of vector control programmes.

Changes in population size and structure leave genomic signatures in populations[28].
Reductions in nucleotide diversity (π and θw) and increases in individual inbreeding (F_{IS}) are
often associated with population decline, which in turn is linked to a greater risk of stochastic
extinction[23,24,29]. Furthermore, changes in rates of dispersal between subpopulations may affect
metapopulation viability. Inference of close-kin relationships using genetic techniques (close-kin
mark-recapture (CKMR)) has been used to quantify *Aedes* mosquito dispersal[30,31], but are
currently untested in estimating *Anopheles* taxa. Population genetics has also been used
previously to estimate vector control impacts. Notable examples include a study in Kilifi, Kenya,
which documented genetic signatures of population decline in *An. gambiae* following large-scale
ITN deployment from 2006. Reductions in nucleotide diversity (~5 % lower π and ~15 % lower
θw), and a deficit in the number of low-frequency sites (negative Tajima's D) were accompanied
by changes in vector species composition, whereby the presence of the primary malaria vector
species *An. gambiae sensu stricto* declined from 79% of *An. gambiae* species complex samples
between 1997-1998 to an undetectable level in 2007-2008[32]. In a separate study in western
Kenya, a shift in *An. gambiae* from 85% to 1% of collected adults in Kenya between 1999 and
2010 was coincident with high ITN coverage[13], with genomic signatures of population decline
such as increased inbreeding, decreased genetic diversity, and allele-frequency shifts, evident
in samples sequenced in a separate study[33]. There are also reported instances where vector
control was not associated with significant signatures of population disruption. In Cameroon,
ITN implementation was associated with insignificant impacts on *An. arabiensis* genetic diversity
and population structure[34], though no clinical or entomological endpoints were reported with
these data. Evidence of changes to population structure in the face of successful vector control
is currently limited and variable.

Significant epidemiological impacts in cluster-randomised controlled trials (cRCT) of
vector control are often accompanied by declines in vector population densities. However, the
impact on mosquito population structure and the extent to which epidemiological impact
requires population disruption, remains unclear. This lack of clarity could stem from the lack, to
our knowledge, of published studies with systematic comparisons between treatment versus
control groups that also incorporate population genetic estimations of changes in structuring or
diversity. This knowledge gap can be addressed by applying population genomic techniques to
dense vector sampling from clinical trials, especially field cRCTs comparing control and
treatment groups. Low-coverage whole-genome-sequencing (WGS) approaches (e.g. a depth-

of-coverage of <5X) are particularly attractive for capture of accurate population allele

frequencies and fine-scale population structure from large samples sizes at much reduced per-

sample cost than medium-full (e.g. 10-30X) coverage approaches[35,36].

Validation of new vector control strategies, such as gene-drive approaches, spatial

repellents and tools using novel insecticide classes with different entomological modes of action

often require evaluation in cRCTs to determine their public health value. While standard

guidelines are in place for measurement of entomological and epidemiological endpoints[37],

methodologies to understand how these interventions impact vector populations more

holistically are missing. Population genetic analysis of vectors coupled with cRCTs can provide

more information on the effect of vector control on mosquito population dynamics and in turn the

extent to which epidemiological outcomes are coupled with substantial demographic impacts on

vector populations.

Here, we use the case study of the AvecNet cRCT performed in Burkina Faso[38], in

which a standard pyrethroid insecticide-treated net (Olyset ITN) was compared with a novel net

that combined pyrethroid with an insect growth regulator, pyriproxyfen (Olyset DUO net also

known as ITN-PPF). The ITN-PPF reduced the incidence of clinical malaria in children by 12%,

and vector numbers by 22%, compared to the standard ITN[38]. Our objective was to analyse

population genetic data from *An. gambiae* samples collected during and after the trial to assess

the impacts on: 1) the spatial genetic structure; 2) genetic diversity and inbreeding (as general,

although imperfect[39] proxies for population size); 3) putative signatures of selection in the

genomes of mosquitoes collected from treated vs control clusters (see **Methods** for more

detailed justification). We provide a framework to detect changes in population structure in the

context of field trials, to give insight into the impacts of control on fragmentation and elimination

of vector populations.

Methods

Study design and entomological sample collection

Briefly, the AvecNet stepwise cRCT was a two-arm trial undertaken in 2014–15 in Banfora,

Burkina Faso[38]. In this area, malaria is endemic and largely transmitted by species of the

morphologically-identical *An. gambiae s.l.* complex (*An. arabiensis*, *An. gambiae* and *An.*

coluzzii). The mosquito population, and consequently malaria incidence, is highly seasonal with

peaks in abundance generally concentrated during the rainy season, which occurs between July

and November.

The cRCT aimed to compare the effectiveness against clinical malaria of the Olyset DUO net
(hereafter ITN-PPF or treated), which in addition to the pyrethroid permethrin contains
pyriproxyfen, an insect growth regulator, to the standard Olyset pyrethroid-only ITN (hereafter
ITN or control), impregnated with permethrin only. The trial included 81 consenting villages that
were grouped into 40 clusters, each including an average of 50 children under 5 years old. At
the start of the malaria transmission season in June 2014, five randomly-chosen clusters were
provided with ITN-PPF bednets, while the remaining clusters were given ITN bednets. Five
clusters were randomly chosen for further ITN-PPF delivery each month from July-Nov 2014 so
that by the end of 2014, each trial arm had an equal number of clusters. In 2015, ITN-PPF were
distributed in a similar fashion starting in June 2015. By the end of the trial in December 2015,
all clusters had the ITN-PPF nets (**See Figure 1** for more information on net and cluster
distribution). During the cRCT, mosquitoes were collected using CDC Light Traps in six
randomly-chosen households per cluster, every four weeks between May-December in 2014
and May-November in 2015. All mosquitoes were identified using microscopy, and a random
subset of approximate 30% of *An. gambiae s.l.* was typed by PCR.[38] Full details of the study
protocol and collections are available in Tiono et al[38].

Sequencing
A total of 16,785 mosquitoes were collected during this trial using CDC light traps, of which
14,489 (86%) were *An gambiae sensu strico* (hereafter *An. gambiae*). From these, we
sequenced the whole genomes of 893 *An. gambiae*. We chose mosquitoes from clusters pre
(ITN) and post-treatment (ITN-PPF) from around early (July/June 2014 and 2015), peak
(September 2014 and 2015) and late (November 2014 and 2015) rainy season; while also
ensuring we had good representation of space to capture the largest possible distances among
clusters (**Figure 2**). To ensure sufficient sample sizes for population genetic analysis, we
primarily restricted our choice to clusters x monthly timepoints for which >10 samples were
available. The clusters selected were an average of 29km apart from each other (minimum of
3.53km, and maximum of 74.45km). In total we sequenced 240 *An. gambiae* specimens from
ITN-PPF clusters, and 174 from ITN clusters in 2014, and 418 and 61 *An. gambiae* sampled
from ITN-PPF and ITN clusters in 2015, respectively. Genomic DNA was extracted using a
modified low salt Proteinase K buffer to maximise DNA yield[40], and Illumina-sequenced by
Novogene UK. Mean per-sample sequencing coverage was 3.43X (for per-sample sequencing
data see **Table S1**). Samples were grouped for analysis by cluster, and by year/month of
collection. Each sample grouping is hereafter referred to as a cluster:timepoint.

Figure 1. Clinical trial and collection setup. A) Map of Burkina Faso highlighting the Banfora region (top panel) and a zoom in on the centroid of the 40 clusters enrolled in the cRCT (bottom panel). Maroon circles indicate clusters from which mosquitoes were sequenced. B) Timeline of the implementation of ITN (yellow, control) and ITN-PPF bednets (blue, treated) in each cluster, by year and month, for clusters where mosquitoes were sequenced. Numbers within a cell indicate the number of *An. gambiae* specimens that were sequenced in that particular cluster:timepoint with the total numbers per cluster given in the right hand column.

Population genetic analyses

We investigated the impact of ITN-PPF on the spatial genetic structure of *An. gambiae* during the study in multiple ways. First we estimated kinship between all mosquito samples with the aim of building a dispersal kernel from geographic distance between close kin[41]. This analysis aimed to determine whether kinship decayed significantly with geographic distance (i.e. displayed isolation-by-distance) and whether isolation by distance was greater between samples from treated/treated and treated/control vs control/control clusters[30,31] (e.g. if dispersal occurred less frequently from treated sites). Second we estimated F_{ST} between groups of samples from each cluster:timepoint (year-month), and all other cluster:timepoints, to determine whether treated sites had higher mean genetic differentiation, and therefore potential population fragmentation[42,43] (possibly due to reduced dispersal or diversity in smaller populations). Finally

we performed a principal component analysis of genotype-likelihoods from all mosquito samples

to test whether those collected from treated (ITN-PPF) clusters have more discrete genetic

clusters than those from control (ITN) clusters[42,43].

We investigated the impact of the ITN-PPF intervention on the genetic diversity of *An.*

gambiae using two metrics: nucleotide diversity (π), and individual inbreeding coefficient (F_{IS}).

We estimated F_{IS} per individual: an increase in per-individual inbreeding is a result of mating

between related individuals, and of low genetic diversity, in each cluster:timepoint. Finally, we

performed genome-wide scans of F_{ST} between treated and control sites to identify potential

regions of the genome subject to selection in candidate insecticide-resistance regions[44]. The

total sample size (893), and mean sequence coverage level (3.43), is sufficient to accurately

estimate individual inbreeding coefficients[45], kinship coefficients[46], the site-frequency spectrum

for the entire sequenced population (for PCA), and for treatment/year subdivisions (for F_{ST}

between years and treatments for genome scans)[47]. Sequenced sample sizes within

cluster:timepoints ranged from 3-20, with sample sizes <10 excluded from per-cluster:timepoint

based analysis such as F_{ST}, based on the expectation that low coverage sequencing of sample

sizes between 10 and 20 are able to recover similar results as higher coverage and sample

sizes[35].

Results

No discernible clustering pattern was present in an all-sample PCA (**Figure 2A**), which revealed

two groups along PC1 (**Figure 2A**) but these did not correspond to any specific cluster,

timepoint, or treatment arm. Instead, the small group beyond the value of 0.2 of PC1 are from

mosquitoes of a mix of clusters and timepoints (**Figure S1, Figure S2**) indicating possible

cryptic population structure associated with currently unknown factors. We were also not able to

detect any spatial structure and consequently no change in spatial structure in mosquito

populations in association with the intervention. The mean F_{ST} between a single

cluster:timepoint, and all other cluster:timepoints, was very low (mean -0.00073), with a

negative F_{ST} value denoting greater genetic diversity within populations than between them. We

estimated slightly lower mean differentiation (F_{ST}) between ITN-PPF clusters (-0.000748)

compared to ITN clusters (-0.000680) and between 2015 (-0.000841) and 2014 (-0.000597)

(**Figure 2B**), though neither difference was significant. These results suggest that the mosquito

population did not detectably fragment or change in structure as a result of treatment as the trial

progressed. We found no obvious potential dispersal events. Kinship coefficient values between

0.125 and 0.5 indicate close-kin relationships[44,48]. We found the mean kinship coefficient for our

data was -0.66 [min= -2.95; max: -0.26] suggesting no close-kin relationships in any of the
samples. Moreover, we found no significant change in between-sample relatedness in pre- or
post- treatment clusters from 2014 or 2015.

The nucleotide diversity (π) of *An. gambiae* from ITN and ITN-PPF clusters in 2014 and
2015 was between 0.13 and 0.21, with a mean of 0.15, comparable to previous estimates of
West African *An. gambiae*[33,49]. Nucleotide diversity was slightly lower in *An, gambiae* from
treated vs control clusters in 2015 (**Figure 2C**) and inbreeding coefficient (*Fis*) slightly lower in
individuals from ITN-PPF vs ITN clusters in 2014 (**Figure 2D**). However, in neither case was
this statistically significant. Finally, a genome scan of F_{ST} between treated and control clusters
did not show any evidence of peaks in regions containing insecticide resistance genes between
ITN-PPF and ITN clusters (**Fig S3**), indicating a lack of differentiation in allele frequencies at
these loci. Taken together, the results indicate that the previously estimated 22% reduction in
mosquito population size caused by the ITN-PPF nets was not associated with alterations in the
spatial genetic structure and diversity of *An. gambiae* in the trial region, or to produce signs of
selection (linked to insecticide resistance, for example), during the time span of the trial.

252
253   **Figure 2. Analysis of spatial population structure and diversity.** (A) Principal component
254   analysis of *An. gambiae* samples, with variance explained by each PC in brackets. Panels B, C
255   and D indicate between-cluster:timepoint F_{ST}, within cluster:timepoint π, and per-individual *Fis*
256   by treatment and year, respectively. Yellow and blue points indicate control (0) and treated (1)
257   sites and individuals respectively.

Discussion

During the AvecNet cRCT, the ITN-PPF nets produced a 12% reduction in clinical malaria in children, and a 22% reduction in mosquito density[38] compared to pyrethroid-only ITNs. These epidemiological and entomological impacts did not coincide with detectable changes in mosquito genetic population structure between treated and control clusters as seen by our analysis of whole genome sequence data of *An. gambiae* mosquitoes, densely sampled over a fine spatial scale. Our findings suggest that an epidemiological benefit from a successful vector control intervention is not necessarily associated with detectable changes in the vector population itself. To our knowledge, this is the first study reporting population genetic data, coupled with corresponding entomological and epidemiological endpoints in a cRCT setting.

Depending on the magnitude and modes of action of the intervention, epidemiological impacts may arise in the absence of substantial changes in vector population size and structure. For example, small reductions in mosquito life span might not necessarily impact recruitment or population size but could be sufficient to reduce the entomological inoculation rate (EIR) and consequently human infection risk[50-52]. Similarly, interventions that work by changing mosquito behaviour (e.g. impeding their ability to find or feed on a host) might substantially reduce human exposure to biting without reducing vector population size. For example, spatial repellents that impede vector ability to locate or feed on a human host might increase the time required for mosquitoes to find a blood meal and their probability of becoming infected (fewer host contacts), but may not diminish mosquito population size if other non-human hosts are available to feed on (e.g. livestock). Such mechanisms reduce the EIR and cut malaria risk even in the absence of substantial vector population size reduction or local extinction. In the case of the AvecNet trial, the additional sterilising effect of pyriproxyfen was sufficient to reduce mosquito population sizes by 22% and the likelihood of clinical malaria in children by 12% in the ITN-PPF cohort without altering mosquito vector population structure or diversity.

Previously published work examining the impact of vector control on population genetic structure and diversity in concert with entomological or epidemiological indicators (albeit none in a cRCT setting)[42,53–55], reveals no consistent pattern of entomological or epidemiological effect required to achieve a population genetic impact. In Dielmo, Senegal, ITN-associated reductions of malaria prevalence from 86% to 0.3% over 22 years, and near-elimination of *An. gambiae* in favour of *An. arabiensis*, was not accompanied by any changes in diversity or inbreeding[56]. By contrast, ITN administration in Papua New Guinea was associated with changes in *An. hinesorum* and *An. farauti* spatial population structure, as well as reductions of 38% in caught *Anopheles* and declines in malaria transmission intensity of 11.1 to 0.9%[42,53]. It is possible that

the reduction in vector population density reported in the ITN-PPF clusters of the AvecNet trial,
and any underlying transient impact on population allele frequencies, might not have been
strong enough to be detectable in the face of homogenising gene flow. Even if ITN-PPF nets
caused temporary local extinction of malaria vectors in some clusters, recolonisation through
migration might occur quickly enough to be indistinguishable from population persistence.
These data suggest using population genetic data as an indicator of a successful vector control
outcome may be of limited use when the impact of vector control on entomological indicators is
modest. This might also be true in some cases where an intervention elicits large
epidemiological and entomological impacts, but large mosquito population sizes and
immigrations from outside the trial area make it difficult to detect changes in in vector population
genetic analyses. In the latter case, application of our approach to situations where evaluations
are conducted at a larger spatial scale than a typical cRCT[57–59] may have more utility.

We show that changes to population structure are not necessarily associated with
significant epidemiological or entomological impacts of vector control, but population genetic
techniques may become more reliably useful when the goal of an intervention is to eliminate
malaria vector populations rather than just reducing human exposure. At low population sizes,
acquiring adequate numbers of mosquitoes to robustly estimate population size may be
challenging; whereas a small number of specimens may be sufficient to elucidate genomic
signatures of vector population decline (e.g. inbreeding, differentiation). As discussions progress
around the deployment of gene-drive technology, as well as other biological control tools that
may eliminate mosquito vector populations [60,61], it is also likely that genomics-based approaches
such as those developed here for estimating mosquito population diversity, structure and sizes
will become crucial for measuring the impact of vector control.

Acknowledgements
This research was funded by the European Research Council, European Union's Horizon 2020
Research and Innovation Programme grant 852957.

Data availability
Full details of laboratory and bioinformatic workflows are available in the *SI Methods.* Analysis
code   and   bioinformatic   workflows   are   available   at:
https://github.com/tristanpwdennis/avecnet_popgen/ . Raw reads are deposited in the European
Nucleotide Archive under accession PRJEB62110.

Supplementary Material

Supplementary Methods

We sequenced the whole genomes of 893 *An. gambiae s.s.* to a median depth-of-coverage of 3.58X with 2x150bp paired-end Illumina reads. Raw reads were trimmed using fastp[62], and aligned to the *An. gambiae* PEST reference genome[63], using bwa-mem[64]. Alignments were sorted, with duplicates marked, using samtools[65]. BAM alignments were grouped into cohorts for analysis, consisting of all 893 samples, and samples grouped by cluster and timepoint (year-month, see **Figure 1** for cohort sizes). Genotype-likelihoods (GLs) and site allele frequencies (SAFs) were inferred using ANGSD[66]. All-sample GL calling was performed for principal-component analysis (PCA) and inbreeding coefficient (Fis) estimation. Per- cluster per-timepoint site allele frequency estimation (see **Figure 1 for cohort sizes**) was performed on the whole genome for the estimation of per-cluster per-timepoint F_{ST} and π. For the all-sample callsets, GLs were called with the reference genome as the major allele (-doMajorMinor 4), filtering GLs based on a a minor allele (maf) frequency of 0.01 (-minmaf 0.01), maximum SNP p value 0.05 (-snp pval 0.05), covered only by properly paired reads (-only proper pairs 1), and uniquely mapped reads with a mapping quality of 20 (-uniqueOnly 1 -minMapQ 20). For the per-cluster per-timepoint SAF estimation, we used the parameters above, but omitted depth, SNP p-value, and maf filters. Filtering that biases the SFS towards more frequent sites can substantially distort estimates of F_{ST} and π. Full parameters are detailed in the avecnet_popgen repository detailed in the main text, under bin/angsd scripts. Principal component analysis and per-sample Fis estimation was performed on chromo- some arm 3L GLs using PCAngsd[67]. Folded site-frequency spectra (SFS) were calculated from SAF files for all samples, as well as per-cluster and per-timepoint, using realSFS[66]. For estimating differentiation of each cluster and timepoint with respect to the wider metapopulation, two-dimensional(2d)-SFS were generated between the allsample SFS, and per-cluster per-timepoint SFS. Each 2d-SFS was used to infer Hudson's F_{ST} between each pair of cluster:timepoints. Hudson's F_{ST} was chosen as it is more robust to sample size differences between populations[68]. Per-cluster/timepoint SFS were used to estimate per-population π.   PCA, F_{ST}, π, and F_{IS} data were analysed in R v4.2.2[69]. GLMs assessing the effects of treatment and year on F_{ST}, π, and Fis were performed using glm. Negative binomial GLMMs were performed using lme4[70]. Residuals were examined using DHARMa v0.4.6[71]. Plotting was performed with ggplot2[72] library. To assess the effect of DUO on per-individual F_{IS}, and per-cluster-, per-timepoint F_{ST}, and π, Gaussian GLMs using the three

parameters prior as a function of DUO administration and time, were parameterised. All three

GLMs returned insignificant results.

Supplementary Figures

Figure S1: Principal component analysis (PCA) of sequenced *An. gambiae* from the AvecNet trial. Panels

A, B and C indicate PCs 1/2, 3/4 , and 5/6 respectively, indicated on the X and Y axes. Point colour

indicates cluster.

Figure S2: Principal component analysis (PCA) of sequenced *An. gambiae* from the AvecNet trial. Panels

A, B and C indicate PCs 1/2, 3/4 , and 5/6 respectively, indicated on the X and Y axes. Point colour

indicates Year:Month.

Figure S3: Genome-wide *Fst* between treated and untreated samples. X axis indicates chromosomal

position, Y axis indicates *Fst* in 50Kb windows. Plot is faceted by chromosome

380
381
382 **References**
383

1. Wilson, A. L. *et al.* The importance of vector control for the control and elimination of vector-

borne diseases. *PLoS Negl. Trop. Dis.* **14**, e0007831 (2020).

2. Ashton, R. A., Prosnitz, D., Andrada, A., Herrera, S. & Yé, Y. Evaluating malaria

programmes in moderate- and low-transmission settings: practical ways to generate robust

evidence. *Malar. J.* **19**, 75 (2020).

3. Yé, Y. *et al.* Framework for Evaluating the Health Impact of the Scale-Up of Malaria Control

Interventions on All-Cause Child Mortality in Sub-Saharan Africa. *Am. J. Trop. Med. Hyg.*

97, 9–19 (2017).

4. Ross, R. Some Quantitative Studies in Epidemiology. *Nature* **87**, 466–467 (1911).

5. Morrison, A. C. *et al.* Efficacy of a spatial repellent for control of Aedes-borne virus

transmission: A cluster-randomized trial in Iquitos, Peru. *Proc. Natl. Acad. Sci.* **119**,

e2118283119 (2022).

6. Indriani, C. *et al.* Impact of randomised wmel Wolbachia deployments on notified dengue

cases and insecticide fogging for dengue control in Yogyakarta City. *Glob. Health Action* **16**,

2166650 (2023).

7. World Health Organisation, C. *World Malaria Report 2023*. (World Health Organisation,

Geneva, Switzerland).

8. Bhatt, S. *et al.* The effect of malaria control on Plasmodium falciparum in Africa between

2000 and 2015. *Nature* **526**, 207–211 (2015).

9. Gatton, M. L. *et al.* The importance of mosquito behavioural adaptations to malaria control in

Africa. *Evol. Int. J. Org. Evol.* **67**, 1218–1230 (2013).

10. Sanou, A. *et al.* Insecticide resistance and behavioural adaptation as a response to long-

lasting insecticidal net deployment in malaria vectors in the Cascades region of Burkina

Faso. *Sci. Rep.* **11**, 17569 (2021).

11. Russell, T. L. *et al.* Increased proportions of outdoor feeding among residual malaria vector

populations following increased use of insecticide-treated nets in rural Tanzania. *Malar. J.*

10, 80 (2011).

12. Hemingway, J. & Ranson, H. Insecticide Resistance in Insect Vectors of Human Disease.

Annu. Rev. Entomol. **45**, 371–391 (2000).

13. Bayoh, M. N. *et al.* Anopheles gambiae: historical population decline associated with

regional distribution of insecticide-treated bed nets in western Nyanza Province, Kenya.

Malar. J. **9**, 62 (2010).

14. Msugupakulya, B. J. *et al.* Changes in contributions of different Anopheles vector species to

malaria transmission in east and southern Africa from 2000 to 2022. *Parasit. Vectors* **16**,

408 (2023).

15. Magesa, S. M. *et al.* Trial of pyrethroid impregnated bednets in an area of Tanzania

holoendemic for malaria Part 2. Effects on the malaria vector population. *Acta Trop.* **49**, 97–

108 (1991).

16. Gimnig, J. E. *et al.* Effect of permethrin-treated bed nets on the spatial distribution of malaria

vectors in western Kenya. *Am. J. Trop. Med. Hyg.* **68**, 115–120 (2003).

17. Hanski, I. Metapopulation dynamics. *Nature* **396**, 41–49 (1998).

18. Fountain, T. *et al.* Predictable allele frequency changes due to habitat fragmentation in the

Glanville fritillary butterfly. *Proc. Natl. Acad. Sci.* **113**, 2678–2683 (2016).

19. Hanski, I. *et al.* Ecological and genetic basis of metapopulation persistence of the Glanville

fritillary butterfly in fragmented landscapes. *Nat. Commun.* **8**, 14504 (2017).

20. Hanski, I. A. Eco-evolutionary spatial dynamics in the Glanville fritillary butterfly. *Proc. Natl.*

Acad. Sci. **108**, 14397–14404 (2011).

21. Elkinton, J. S. & Liebhold, A. M. Population Dynamics of Gypsy Moth in North America.

Annu. Rev. Entomol. **35**, 571–596 (1990).

22. Kuussaari, M., Saccheri, I., Camara, M. & Hanski, I. Allee Effect and Population Dynamics

in the Glanville Fritillary Butterfly. *Oikos* **82**, 384–392 (1998).

23. Nieminen, M., Singer, M. C., Fortelius, W., Schöps, K. & Hanski, I. Experimental

Confirmation that Inbreeding Depression Increases Extinction Risk in Butterfly Populations.

Am. Nat. **157**, 237–244 (2001).

24. Saccheri, I. *et al.* Inbreeding and extinction in a butterfly metapopulation. *Nature* **392**, 491–

494 (1998).

25. Benedick, S. *et al.* Impacts of habitat fragmentation on genetic diversity in a tropical forest

butterfly on Borneo. *J. Trop. Ecol.* **23**, 623–634 (2007).

26. Brady, O. J. *et al.* Adult vector control, mosquito ecology and malaria transmission. *Int.*

Health **7**, 121–129 (2015).

27. Killeen, G. F. *et al.* Eliminating malaria vectors. *Parasit. Vectors* **6**, 172 (2013).

28. Bradburd, G. S. & Ralph, P. L. Spatial Population Genetics: It's About Time. *Annu. Rev.*

Ecol. Evol. Syst. **50**, 427–449 (2019).

29. Söderquist, L., Broberg, A., Rosenberg, V. & Sletvold, N. Predicting heterosis and

inbreeding depression from population size and density to inform management efforts. *J.*

Appl. Ecol. **57**, 1459–1468 (2020).

30. Jasper, M. E., Hoffmann, A. A. & Schmidt, T. L. Estimating dispersal using close kin dyads:

The kindisperse R package. *Mol. Ecol. Resour.* **22**, 1200–1212 (2022).

31. Filipović, I. *et al.* Using spatial genetics to quantify mosquito dispersal for control programs.

BMC Biol. **18**, 104 (2020).

32. O'Loughlin, S. M. *et al.* Genomic signatures of population decline in the malaria mosquito

Anopheles gambiae. *Malar. J.* **15**, 182 (2016).

33. Miles, A. *et al.* Genetic diversity of the African malaria vector Anopheles gambiae. *Nature*

552, 96–100 (2017).

34. Wondji, C. *et al.* Impact of insecticide-treated bed nets implementation on the genetic

structure of Anopheles arabiensis in an area of irrigated rice fields in the Sahelian region of

Cameroon. *Mol. Ecol.* **14**, 3683–3693 (2005).

35. Lou, R. N., Jacobs, A., Wilder, A. P. & Therkildsen, N. O. A beginner's guide to low-

coverage whole genome sequencing for population genomics. *Mol. Ecol.* **30**, 5966–5993

(2021).

36. Therkildsen, N. O. & Palumbi, S. R. Practical low-coverage genomewide sequencing of

hundreds of individually barcoded samples for population and evolutionary genomics in

nonmodel species. *Mol. Ecol. Resour.* **17**, 194–208 (2017).

37. *Data Requirements and Protocol for Determining Comparative Efficacy of Vector Control*

Products. (World Health Organisation, Geneva, Switzerland, 2024).

38. Tiono, A. B. *et al.* Efficacy of Olyset Duo, a bednet containing pyriproxyfen and permethrin,

versus a permethrin-only net against clinical malaria in an area with highly pyrethroid-

resistant vectors in rural Burkina Faso: a cluster-randomised controlled trial. *The Lancet*

392, 569–580 (2018).

39. Buffalo, V. Quantifying the relationship between genetic diversity and population size

suggests natural selection cannot explain Lewontin's Paradox. *eLife* **10**, e67509 (2021).

40. Korlević, P. *et al.* A Minimally Morphologically Destructive Approach for DNA Retrieval and

Whole-Genome Shotgun Sequencing of Pinned Historic Dipteran Vector Species. *Genome*

Biol. Evol. **13**, evab226 (2021).

41. Sharma, Y., Bennett, J. B., Rašić, G. & Marshall, J. M. Close-kin mark-recapture methods to

estimate demographic parameters of mosquitoes. *PLOS Comput. Biol.* **18**, e1010755

(2022).

481 42. Keven, J. B. *et al.* Genetic differentiation and bottleneck effects in the malaria vectors

482   Anopheles farauti and Anopheles punctulatus after an LLIN-based vector control program in

483   Papua New Guinea. *Ecol. Evol.* **14**, e10917 (2024).

484 43. Lamy, T., Pointier, J. P., Jarne, P. & David, P. Testing metapopulation dynamics using

485   genetic, demographic and ecological data. *Mol. Ecol.* **21**, 1394–1410 (2012).

486 44. Lucas, E. R. *et al.* Genome-wide association studies reveal novel loci associated with

487   pyrethroid and organophosphate resistance in Anopheles gambiae and Anopheles coluzzii.

488   *Nat. Commun.* **14**, 4946 (2023).

489 45. Vieira, F. G., Fumagalli, M., Albrechtsen, A. & Nielsen, R. Estimating inbreeding coefficients

490   from NGS data: Impact on genotype calling and allele frequency estimation. *Genome Res.*

491   **23**, 1852–1861 (2013).

492 46. Hanghøj, K., Moltke, I., Andersen, P. A., Manica, A. & Korneliussen, T. S. Fast and accurate

493   relatedness estimation from high-throughput sequencing data in the presence of inbreeding.

494   *GigaScience* **8**, (2019).

495 47. Alex Buerkle, C. & Gompert, Z. Population genomics based on low coverage sequencing:

496   how low should we go? *Mol. Ecol.* **22**, 3028–3035 (2013).

497 48. Manichaikul, A. *et al.* Robust relationship inference in genome-wide association studies.

498   *Bioinformatics* **26**, 2867–2873 (2010).

499 49. Consortium, T. A. gambiae 1000 G. *et al.* Genome variation and population structure among

500   1142 mosquitoes of the African malaria vector species Anopheles gambiae and Anopheles

501   coluzzii. *Genome Res.* **30**, 1533–1546 (2020).

502 50. Smith, D. L., Dushoff, J., Snow, R. W. & Hay, S. I. The entomological inoculation rate and

503   Plasmodium falciparum infection in African children. *Nature* **438**, 492–495 (2005).

504 51. Shaukat, A. M., Breman, J. G. & McKenzie, F. E. Using the entomological inoculation rate to

505   assess the impact of vector control on malaria parasite transmission and elimination. *Malar.*

506   *J.* **9**, 122 (2010).

52. Grisales, N. *et al.* Pyriproxyfen-treated bed nets reduce reproductive fitness and longevity of

pyrethroid-resistant Anopheles gambiae under laboratory and field conditions. *Malar. J.* **20**,

273 (2021).

53. Hetzel, M. W. *et al.* Insecticide-treated nets and malaria prevalence, Papua New Guinea,

2008–2014. *Bull. World Health Organ.* **95**, 695-705B (2017).

54. Trape, J.-F. *et al.* The rise and fall of malaria in a west African rural community, Dielmo,

Senegal, from 1990 to 2012: a 22 year longitudinal study. *Lancet Infect. Dis.* **14**, 476–488

(2014).

55. Sougoufara, S., Harry, M., Doucouré, S., Sembène, P. M. & Sokhna, C. Shift in species

composition in the Anopheles gambiae complex after implementation of long-lasting

insecticidal nets in Dielmo, Senegal. *Med. Vet. Entomol.* **30**, 365–368 (2016).

56. Sougoufara, S. *et al.* The implementation of long-lasting insecticidal bed nets has differential

effects on the genetic structure of the African malaria vectors in the Anopheles gambiae

complex in Dielmo, Senegal. *Malar. J.* **16**, 337 (2017).

57. Gansané, A. *et al.* Design and methods for a quasi-experimental pilot study to evaluate the

impact of dual active ingredient insecticide-treated nets on malaria burden in five regions in

sub-Saharan Africa. *Malar. J.* **21**, 19 (2022).

58. Gonahasa, S. *et al.* LLIN Evaluation in Uganda Project (LLINEUP2) - Effect of long-lasting

insecticidal nets (LLINs) treated with pyrethroid plus pyriproxyfen vs LLINs treated with

pyrethroid plus piperonyl butoxide in Uganda: A cluster-randomised trial. *PLOS Glob. Public*

Health **5**, e0003558 (2025).

59. Maiteki-Sebuguzi, C. *et al.* Effect of long-lasting insecticidal nets with and without piperonyl

butoxide on malaria indicators in Uganda (LLINEUP): final results of a cluster-randomised

trial embedded in a national distribution campaign. *Lancet Infect. Dis.* **23**, 247–258 (2023).

60. Bier, E. Gene drives gaining speed. *Nat. Rev. Genet.* **23**, 5–22 (2022).

61. Nolan, T. Control of malaria-transmitting mosquitoes using gene drives. *Philos. Trans. R.*

Soc. B Biol. Sci. **376**, 20190803 (2020).

62. Chen, S., Zhou, Y., Chen, Y. & Gu, J. fastp: an ultra-fast all-in-one FASTQ preprocessor.

Bioinformatics **34**, i884–i890 (2018).

63. Holt, R. A. *et al.* The Genome Sequence of the Malaria Mosquito Anopheles gambiae.

Science **298**, 129–149 (2002).

64. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows–Wheeler transform.

Bioinformatics **25**, 1754–1760 (2009).

65. Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *GigaScience* **10**, giab008

(2021).

66. Korneliussen, T. S., Albrechtsen, A. & Nielsen, R. ANGSD: Analysis of Next Generation

Sequencing Data. *BMC Bioinformatics* **15**, 356 (2014).

67. Meisner, J. & Albrechtsen, A. Inferring Population Structure and Admixture Proportions in

Low-Depth NGS Data. *Genetics* **210**, 719–731 (2018).

68. Bhatia, G., Patterson, N., Sankararaman, S. & Price, A. L. Estimating and interpreting FST:

The impact of rare variants. *Genome Res.* **23**, 1514–1521 (2013).

69. R Core Team. R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria. https://www.eea.europa.eu/data-and-

maps/indicators/oxygen-consuming-substances-in-rivers/r-development-core-team-2006.

70. Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting Linear Mixed-Effects Models Using

lme4. *J. Stat. Softw.* **67**, 1–48 (2015).

71. Hartig, F. DHARMa: residual diagnostics for hierarchical (multi-level/mixed) regression

models. (2022).

72. Wickham, H. *Create Elegant Data Visualisations Using the Grammar of Graphics.* (Springer-

Verlag New York, 2016).