A comprehensive pipeline for genome annotation across species: a case study on *Schmidtea*

mediterranea

Anastasiia Zaremba, Małgorzata Marszałek-Zeńczak, Annasha Dutta, Anna Samelak-Czajka,

Paulina Jackowiak#

Laboratory of Single Cell Analyses; Institute of Bioorganic Chemistry Polish Academy of

Sciences; Zygmunta Noskowskiego str. 12/14, 61-704 Poznań; Poland

azaremba@ibch.poznan.pl,   marszalekgosia@gmail.com,   adutta@ibch.poznan.pl,

asczajka@ibch.poznan.pl,

#Corresponding author: paulinaj@ibch.poznan.pl

Abstract

Background

Despite advancements in genome annotation tools, challenges persist for non-classical

model organisms with limited genomic resources, such as *Schmidtea mediterranea*.

Findings

To address these challenges, we developed a flexible and scalable genome annotation

pipeline that integrates short-read (Illumina) and long-read (PacBio) sequencing technologies. The

pipeline combines reference-based and *de novo* assembly methods, effectively handling genomic

variability and alternative splicing events. Functional annotation is conducted to filter out low-

confidence transcripts and ensure biological relevance. Applying this pipeline to the asexual strain

of *S. mediterranea* revealed thousands of previously undescribed putative genes and transcripts,

and improved the existing gene models, highlighting its utility in annotating complex,

underexplored genomes.

Conclusions

26   The modularity and comprehensiveness of our pipeline ensure its adaptability for genome

27 annotation across diverse species, making it a valuable tool for annotating genomes of non-model

28 organisms and supporting broader genomic research. The source code and implementation details

29 are available at https://github.com/Norreanea/SmedAnno.

Keywords

32 Genome annotation; Transcriptome assembly; *Schmidtea mediterranea*; Non-model organisms;

33 Functional annotation

Background

Understanding the genetic makeup of organisms is crucial for unraveling the complexities

of their biology, making genome annotation a foundational aspect of molecular and genomic

research. By precisely identifying coding regions, non-coding elements, and regulatory sequences,

genome annotation transforms raw genomic data into a practical blueprint that opens up

possibilities for studying living organisms at the molecular level.

Existing genome annotation methods can be broadly categorized into *ab initio* [1],

homology-based [2], evidence-based (reference-based) [3], and hybrid approaches [4–6], each

leveraging different strategies and tools to predict and validate genomic features. Each category

has its unique advantages and challenges. *Ab initio* methods, which predict genes solely from

intrinsic sequence signals without external evidence, are capable of identifying novel genes;

however, they tend to produce a higher rate of false positives and inaccurate gene models—such

as overpredicted exons, missed genes, and incorrect exon–intron boundaries—especially when

applied to complex or poorly characterized genomes. Homology-based methods use evolutionary

conservation to achieve high accuracy for well-conserved genes but often miss novel or rapidly

evolving genes that lack close homologs. Evidence-based approaches utilize transcriptomic or

proteomic data to support gene models, thereby enhancing annotation quality for expressed genes,

yet they are inherently limited to regions where experimental data exist and are as robust as the

underlying evidence. Hybrid approaches combine intrinsic signals with external evidence in an

effort to balance sensitivity and specificity.

To streamline the annotation process, several genome annotation platforms and pipelines

have been developed. For instance, the RefSeq eukaryotic annotation pipeline at NCBI [7] employs

highly curated alignment-based evidence and gene prediction models, while another recently

reported pipeline [8] aims to integrate evidence from multiple data sources for eukaryotic genome

annotation. Web-based platforms such as Galaxy Genome Annotation (GGA) [9] provide an

accessible interface for constructing annotation workflows, and tools like Prokka [10] offer rapid

prokaryotic annotation that automates the integration of gene prediction and functional annotation,

generating standardized output formats compatible with GenBank. Additionally, PAGIT (Post

Assembly Genome Improvement Toolkit) [11], the community-driven Nextflow-based approach

nf-core GenomeAnnotator [12], and other pipelines [13–15] expand the available toolkit.

Despite these advancements, applying genome annotation methods to non-model or non-

classical model organisms presents significant challenges. Due to limited reference data, tools

must rely more heavily on ab initio methods, which—as noted above— are prone to errors,

particularly for complex genomes, and often lack sufficient training data. Repetitive sequences and

the presence of multiple isoforms complicate the annotation process. Genetic diversity and

evolutionary distance from well-studied species can impede the identification of orthologous genes

and conserved pathways, while limited experimental data often constrain the ability to validate or

refine gene models. Such constraints are further compounded by annotation artifacts—including

fragmented and chimeric transcripts—that can obscure true biological functions and introduce

errors into downstream analyses.

These challenges are particularly relevant to research on the sexual (S2F2 clonal line) and

asexual (CIW4) strains of *Schmidtea mediterranea*, a highly regenerative flatworm model

organism [16,17]. Currently, reference genome sequences and comprehensive annotations are

available primarily for the sexual strain, including contig-level genome assemblies such as

SmedGD_c1.3 [18,19], dd_Smes_g4 [20,21], chromosome-scale genome assemblies schMedS2

[22] and recent schMedS3 [23], as well as corresponding annotations like SmedGD_c1.3, high-

confidence SMESG, schMedS1, schMedS2, schMedS3_1 and schMedS3_2. Chromosome-scale

assemblies of the sexual strain indicate that sexual and asexual *Schmidtea mediterranea* share high

genomic similarity—they are both diploid with four pairs of chromosomes. Although the asexual

strain possesses a chromosomal translocation between chromosomes 1 and 3, these reference

sequences and annotations offer a strong starting point for transcriptome analysis. Nonetheless,

given potential strain-specific transcripts and structural variations, further refinement of the

asexual strain's transcript set is essential [24]. Additionally, the asexual strain exhibits a simplified

reproductive strategy [25–28] and it has been widely used in numerous studies, with extensive data

deposited in public repositories. As of January 2, 2025, there are 122 DNA and 1,694 RNA datasets

on NCBI [29] for this strain, the majority of which are short-read data. In comparison, for the

sexual strain, there are only 18 DNA and 87 RNA datasets. The predominance of RNA-seq data

highlights the extensive transcriptomic resources available for the asexual strain, which offers a

strong foundation for gene expression and functional studies and underscores the potential for

leveraging RNA sequencing to enhance genome annotation efforts. However, limited detailed

reference resources for asexual strain remain a significant challenge. Currently, the scientific

community must rely mostly on reference transcriptomic data—such as

dd_Smed_v6.pcf.contigs.fasta [21] (a transcriptome assembly generated from RNA-Seq data) and

a contig-based genome assembly and annotation SmedAsxl_genome_v1.1 [19]. Beyond that,

existing transcriptome assemblies are fragmented and incomplete, lacking the high-resolution

insights offered by newer sequencing technologies. To bridge this gap, our study aims to

implement a sophisticated bioinformatics pipeline to improve annotated genes and transcripts in

asexual *Schmidtea mediterranea* using the existing chromosome-scale genome of the sexual strain

(schMedS2) as a reference and in-house generated comprehensive collection of transcriptomic data

for asexual worms. Our dataset includes both short-read and long-read data, providing a significant

advantage in resolving complex isoforms. Addressing these challenges requires the integration of

various sequencing technologies and advanced bioinformatics tools. Short-read sequencing

platforms, such as Illumina, offer high accuracy and depth of coverage, a low error rate, which is

essential for identifying single-nucleotide polymorphisms, small indels, and importantly, splice

sites—ensuring accurate distinction of exon-intron boundaries [30]. Conversely, long-read

technologies like PacBio provide valuable information on structural variants and complex

transcript isoforms, enhancing the resolution of genome assemblies [31,32]. Combining these

technologies benefits from their complementary strengths, enabling a more comprehensive and

accurate genome annotation [4].

Here we present a universal, comprehensive, and adaptable pipeline for genome annotation

to process and interpret the vast amounts of sequencing data generated, that integrates both short-

and long-read sequencing data, employs state-of-the-art bioinformatics tools for transcriptome

assembly and genome characterization, and incorporates robust methods for functional and

homology annotation. By doing so, we aim to advance genome annotation efforts within *Schmidtea*

mediterranea and to provide a scalable framework applicable to other non-model organisms,

fostering broader genomic research and discovery.

Methods

Planarian culturing and sample collection

Asexual *Schmidtea mediterranea* were cultured as described in [33]. Animals were

collected, immediately flash-frozen in liquid nitrogen to preserve RNA integrity, and stored at

−80 °C until RNA isolation.

Dataset details

Full dataset-specific details including experimental treatments, RNA extraction, quality

control, are provided in Table S1 and in the associated GEO submissions.

Validation with simulated data

Validation with simulated data

A synthetic genome was created using a custom Python script, which constructed a single

chromosome spanning 2 mega base pairs (bp). This genome included 1,000 genes randomly

distributed along the chromosome, each comprising between 2 and 20 exons with lengths ranging

from 100 to 1,000 nucleotides (nt) and introns of 100 to 1,000 nt. Paired-end short-read RNA-Seq

data were simulated using ART Illumina tool, generating 100 bp reads at 10× coverage across six

samples. Additionally, long-read sequencing data were produced using PBSIM tool with 150 bp

reads at 10× coverage for one sample. To evaluate the pipeline's ability to handle incomplete

annotations, a random subset of genes was removed from the annotation file using the custom bash

script, resulting in a GTF file with missing gene information.

Reference annotation polishing

The reference genome SchMedS2 [21] was utilized for this study. High-confidence

annotations for this genome were obtained from two sources: SmedSxl smed_chr_ref_v1 UCLA

(SIMRBase (https://simrbase.stowers.org)) and schMedS2_PlanMine_high-confidence_liftover

(PlanMine [21]). Both annotations are compatible with the genome sequence; however, due to

minor differences between them, the genes were first unified, and the annotation was carefully

reviewed and corrected. Through detailed analysis and comparison, 27,676 SMEST transcripts and

21,332 SMESG genes were identified as the most confident and consistent across both sources.

Next, 169 transcripts with identical exon structures using the AGAT tool [34] were removed to

eliminate redundancy and enhance annotation quality.

Assembly evaluation and benchmarking

The transcriptome assembly was evaluated using the SIRV-set 4 RNA spike-in controls to assess its accuracy, with specific StringTie2 parameters (-c, the minimum reads per bp coverage to consider for multi-exon transcript, and -f, the minimum isoform fraction) being tested. Completeness was assessed with BUSCO [35] (v5.7.1) with a Metazoa benchmark of 954 conserved genes, while assembly quality metrics were obtained using rnaQUAST [36] (v2.3.0) and Transrate [37] (v1.0.3). Benchmarking procedures were conducted by comparing the performance of the pipeline against existing annotations, with metrics such as the number of correctly identified transcripts, assembly scores, and completeness indicators being analyzed.

ncRNA identification

All ncRNA sequences were downloaded from RNAcentral [38] and supplemented with 457 tRNA sequences from an external publication. The sequences were linearized and deduplicated using seqkit rmdup. The initial dataset contained 27,390,440 sequences, which was reduced to 27,317,511 after deduplication. A BLASTn search was conducted to identify homologous sequences within the *Schmidtea mediterranea* nuclear and mitochondrial genomes. The following parameters were used: -query: 27,317,511 ncRNA sequences; -db: *Schmidtea mediterranea* genome (nuclear + mtDNA), -evalue: 1e-20, -word_size: 15, -qcov_hsp_perc: 90, -dust: no. The output was then processed to extract the original sequence length of hits for further filtering. A custom Python script was used to classify ncRNA types based on query ID patterns. Hits were retained if alignment coverage met predefined thresholds (95% for tRNA, 90% for miRNA, 70% for lncRNA, 90% for rRNA, 80% for other ncRNA). Then a custom R script was used for converting to gtf format, refining and merging overlapping regions to ensure only non-redundant ncRNA annotations were retained.

Analyses

Selection and validation of transcriptome assembly approach

One of the key steps in genome annotation is the assembly of the transcriptome. In our efforts to enhance genome characterization for the asexual strain of *Schmidtea mediterranea*, we employed StringTie2 [4], a widely recognized tool for transcriptome assembly. StringTie2 is versatile, supporting both short-read and long-read sequencing datasets, which makes it well-suited for comprehensive transcriptome analysis. However, StringTie2 has multiple versions, each offering distinct features and performance enhancements. Therefore, before integrating StringTie2 into our pipeline, we thoroughly optimized its parameters to ensure the highest accuracy and reliability in transcript predictions. To this end, we employed two complementary strategies.

In the first strategy, we generated an *in-silico* dataset to simulate genomic sequences with corresponding full and incomplete annotations and short-read RNA-Seq data (see Methods section). This synthetic data allowed us to create scenarios with known simple transcript structures, enabling a comprehensive evaluation of StringTie2 performance. Reads were aligned to the *in-silico* reference genome using STAR. We then compared two commonly available versions of StringTie2, 2.1.1 and 2.2.1, which allowed us to assess version-specific performance improvements. Our results indicated that StringTie2 version 2.1.1 provides a higher identification rate of short-read data in reference-based mode than its predecessor (Figure 1A). Although the StringTie2 documentation does not explicitly state that version 2.1.1 is more sensitive for short-read data, the combination of release note details and our experimental observation suggests that less stringent filtering of version 2.1.1 results in a higher count of assembled isoforms. Meanwhile,

version 2.2.1 has been optimized for hybrid-read assembly (with the --mix option) and incorporates

more conservative measures to reduce false positives—which can lower the total transcript count

when only short-read data are used. This trade-off between sensitivity and precision is common in

transcript assemblers. When maximizing the number of detected isoforms from short reads is a

priority, 2.1.1 may be advantageous, whereas 2.2.1's improvements (including its ability to process

mixed-read data) make it preferable for datasets that include long reads. It should be noted that

these version choices are based on their prevalence at the time of our study, and although our

benchmarking indicates reliable performance, further testing in diverse datasets may be beneficial.

In the second strategy, we utilized Spike-In RNA Variants (SIRVs), a commercially

available collection of RNA spike-in controls, that we added to the RNA isolated from planarians

(see Table S1) prior to the preparation of sequencing libraries. These SIRVs include three distinct

modules — ERCC Mix 1 (a set of 92 synthetic RNA molecules ranging between 250 and 2,000 nt

in length to resemble natural eukaryotic transcripts), Isoform Mix E0 (referred here as multi-exon

SIRVs), and long SIRVs (referred here as mono-exon SIRVs) — each designed to mimic specific

aspects of transcriptome complexity, including transcript abundance, isoform diversity, and

transcript length. RNA-Seq reads were aligned to the reference SIRVs sequences using STAR for

short reads and Minimap2 for long reads. We then employed StringTie2 using the "insufficient

annotation" as the reference provided by Lexogen (SIRV_I), where some SIRVs that are actually

present in the mixes are not annotated, and compared the prediction to the correct annotation of all

SIRVs (SIRV_C). Such a comparative analysis demonstrated how the pipeline could substantially

improve genome annotations through a proper differentiation between known and novel transcript

isoforms. Then, focusing on the mixture of synthetic RNA isoforms Iso Mix E0 category, we

systematically assessed the performances of seven assembly approaches for transcript variant

detection, implementing different combinations of read types—short and mixed—and assembly

modes—reference-based and *de novo*. These approaches included short reads in reference-based

mode (SR_RB), short reads in *de novo* mode (SR_DN), mixed reads in reference-based mode

(MR_RB), mixed reads in *de novo* mode (MR_DN), merged reads in *de novo* mode (MeR_DN),

merged reads in reference-based mode (MeR_RB), and merged reads in both modes

(MeR_RB_DN). For each approach, StringTie2 performance (Table S2) was quantified in terms

of accuracy metrics, reflecting the ability to correctly identify transcript isoforms (Figure 1B).

SR_DN approach achieved the lowest accuracy, ranging from 35.1% to 45.1% with its maximum

at -c 1.5 and -f 0.02. Surprisingly, generally, the highest accuracy rate of 81.1% was obtained when

predictions from both short and mixed reads in both modes were merged (MeR_RB_DN). This

outcome underscores the synergistic benefits of integrating multiple read types and assembly

strategies.

Consequently, our pipeline employs a dual-version strategy—by default, StringTie2

version 2.1.1 is used for assembling short-read data and version 2.2.1 for mixed-read data, ensuring

optimal performance across a range of sequencing data types. Users should note that while these

versions provided robust performance in our tests, alternative versions may be considered

depending on specific experimental contexts and can be adjusted via command-line. We set -c 1.5

and -f 0.02 based on pilot experiments with SIRVs, which balanced sensitivity and precision in

isoform detection (see Table S2). After completing individual assemblies, we incorporated the

MeR_RB_DN approach to merge reference-based and de novo transcripts, resulting in a

comprehensive and unified transcriptome annotation that captures a diverse array of transcript

isoforms.

Identification of artifactual transcripts

In our tests with *in-silico* data, we identified several types of artifactual transcripts,

including those with unusually long exons exceeding 100,000 nt, reversed duplicates (transcripts

overlapping the same genomic regions as the reference ones but on the opposite strand), as well as

fragmented and chimeric RNA. Fragmented transcripts are individual RNAs that have been split

into multiple predicted gene models due to incomplete transcript reconstruction or assembly errors.

Chimeric transcripts, on the other hand, erroneously combine sequences from multiple distinct

biological genes. In eukaryotic genomes, exons generally range in size from approximately 50 to

5,000 nt, with the majority falling between 100 and 1,000 nt [39]. Instances of exons exceeding

10,000 nt are exceptionally rare and are usually associated with specific organismal features, such

as immune-related genes or tandem duplications. Even in these rare cases, such lengths remain an

order of magnitude below 100,000 nt [40]. While highly effective in many contexts, StringTie2

may incorrectly model transcripts, especially in regions with overlapping genes or incomplete stop

codons. Such inaccuracies can extend exon boundaries far beyond biologically realistic lengths

[41].

To address the identified challenges and eliminate artifacts, we added a custom function to

the pipeline that ensured that transcripts with exons exceeding 10,000 nt and/or transcripts that

surpassed 100,000 nt of the genomic region were removed, thus refining the annotation and

preventing the propagation of erroneous data. Additionally, we included a function to identify and

filter out reversed duplicates without inadvertently removing legitimate transcripts. We also

implemented criteria to flag potential fragmented and chimeric RNA based on exon length,

genomic proximity, and functional annotations.

In our pipeline, indicators for fragmentation include multiple predicted genes in close

genomic proximity (within 1,000 nt) on the same strand and adjacent or overlapping genomic

coordinates. The exon length and genomic proximity thresholds are based on general eukaryotic

genome characteristics [40]. These parameters should be adjusted as needed based on the specific

organism or dataset characteristics. Genes and transcripts are flagged as chimeric if they fulfill at

least one of the following criteria: genes have multiple, unrelated functional annotations; display

structural inconsistencies such as exons that are widely spaced or non-contiguous; span multiple

distinct genomic regions. As part of our automated pipeline, we only added information indicating

whether a gene or transcript is possibly fragmented or chimeric. The decision to remove

corresponding genes or transcripts should be made by the user based on specialized evaluation.

Incorporating functional annotations into the pipeline as a part of manual curation can significantly

improve the accuracy of detecting artifacts.

Pipeline overview and workflow

Having resolved the challenges outlined above, we developed a versatile genome

annotation pipeline and demonstrated its performance by applying it to the asexual strain of

Schmidtea mediterranea. This pipeline integrates multiple sequencing technologies and

bioinformatics tools to achieve comprehensive and accurate gene and transcript annotations across

different organisms. The pipeline is divided into two main parts, each structured into distinct

sequential steps.

Part I: Pre-processing and assembly

The first part of the pipeline is fully automated and encompasses pre-processing,

alignment, assembly, and functional annotation. The input data for the pipeline can be either short-

read or mixed, the latter including both short and long reads. While the pipeline can process a

single sample when only short reads are available, incorporating multiple samples is recommended

to achieve more reliable and comprehensive results. Mixed-read datasets require at least one

sample with short reads and one with long reads. Additionally, the automated pipeline is optimized

for paired-end reads in short-read mode. Mapping and transcript assembly are performed

independently for each sample. The pipeline begins by pre-processing the RNA-Seq data, focusing

in particular on the removal of reads corresponding to ribosomal RNA (rRNA), based on their

alignment to user-provided rRNA reference sequences (Figure 2). This step was found to be very

important in the case of organisms with unusually high A/T content, such as *Schmidtea*

mediterranea. Library preparation methods typically involve rRNA depletion or polyA enrichment

to facilitate the study of protein-coding gene expression. However, commercially available rRNA

depletion kits are mostly optimized for model species, and polyA enrichment is not effective in

organisms with a high A/T content. Consequently, reads corresponding to rRNA can be abundant

in sequencing data for non-model organisms. In scenarios where rRNA reference sequences are

unavailable (not provided by the user), the pipeline bypasses the rRNA removal step, continuing

with the whole RNA-Seq dataset. The pipeline then aligns the processed RNA-Seq reads to the

reference genome using STAR for short reads and Minimap2 for long reads. The transcript

assembly phase utilizes StringTie2: by default, version 2.1.1 is applied to short-read data, while

version 2.2.1 is used for mixed-read data (which integrates both short and long reads). Users can

specify alternative versions if desired. After assembly, the pipeline merges transcripts across all

samples using StringTie2 to create unified sets of transcripts. This merging process is performed

separately for reference-based (if reference annotation was provided) and *de novo* assemblies

before combining them into the final complete annotation. This dual approach by performing both

reference-based and *de novo* annotation, provides comprehensive assembly, thus capturing a wide

range of transcript isoforms. Subsequently, transcripts with excessively long exons or that cover

unusually large genomic regions are filtered out. Next, we utilized gffcompare to compare the

predicted annotations against the reference annotations, assessing the completeness and accuracy

of the assembled transcripts. This comparison generates a report that includes data summaries and

accuracy estimations. Additionally, it enriches the predicted annotations with detailed information

about their overlaps with known reference transcripts and genes. However, the generated

annotations may contain issues that can compromise the quality and usability of the annotation

files. To address these issues, we employed AGAT (Another GTF/GFF Tool), which

systematically identifies and corrects common syntax errors, ensuring that essential attributes such

as gene_id and transcript_id are present for each feature, preventing incomplete annotations,

correcting improper use of delimiters in attribute fields (such as replacing spaces with a

semicolon), detecting and removing duplicate entries, and ensuring correct key-value pair syntax.

This correction phase is crucial for maintaining high-quality genome annotation. The obtained set

of transcripts is then subjected to functional annotation in a process that involves the sequential

application of several tools, and which includes homology analysis and the identification of protein

domains. To this end, first, the ORFs are predicted using TransDecoder. Following that, BLASTp

and BLASTx are used for homology search through the SwissProt database. Finally, protein

domains are identified using hmmscan through the PFAM database. This multi-faceted approach

filters out low-confidence annotations, retaining only the transcripts supported by multiple lines

of evidence. We utilized the SwissProt and PFAM databases as a part of the automated pipeline

specifically for their low memory requirements. However, we recommend more sophisticated

resources, for example eggNOG, NCBI, UniRef, KEGG, InterPro, RefSeq, COG, Reactome, and

BioCyc for conducting more comprehensive and in-depth homology analyses, as these databases

provide broader coverage of protein functions, evolutionary relationships, and diverse protein

characteristics. Next, the functional annotations are integrated into the GTF file. This integration

applies specific thresholds: for PFAM predictions, a bit score greater than 20, a conditional E-

value below 1e-5, and coverage exceeding 50%; for SwissProt predictions, a BLAST e-value

threshold of 1e-5 and an identity threshold of 25%. These criteria ensure that low-confidence

functional annotations are discarded. To this end, the pipeline adds new columns to the GTF

annotation   file,   including   has_ORF   (TRUE/FALSE),   best_SwissProt_blastp_hit,

best_SwissProt_blastx_hit, and best_Pfam_hit domain information. Subsequently, users have the

option to remove redundant transcripts and eliminate those lacking functional information,

ensuring that each retained transcript is supported by at least one entry. The next step is the

identification of overlapping genes and transcripts, which detects genes that share genomic regions

and checks if each transcript in a gene has at least one overlapping exonic region with any other

transcript in the same gene. Lastly, the pipeline identifies the aforementioned reversed duplicates

and addresses fragmented and chimeric genes both with and without reference annotations. All

these steps (Figure 2) are implemented in the SmedAnno tool, forming Part I of the entire pipeline.

The source code can be accessed on GitHub (https://github.com/Norreanea/SmedAnno).

Part II: Manual curation and refinement

The second part of the pipeline involves user-driven inspection and refinement of the

annotated data. Despite the implementation of several automated steps aimed at optimizing

transcript assembly and obtaining biologically relevant transcripts that encode functional protein

domains, resolving overlapping or fragmented transcripts remains a significant challenge that

requires specialized approaches. While the primary focus is on transcript assembly, gene features

are subsequently derived from the assembled transcripts. Fragmented and chimeric transcripts,

which primarily affect RNA-level annotations, are addressed by prioritizing the longest isoform to

define each gene, thereby reducing redundancy and improving annotation accuracy. To accurately

determine whether genes should be merged after being marked as potentially fragmented and/or

overlapped, promoter and terminator sequences should be analyzed using Promoter 2.0 and

ARNold tools (Figure 3). This analysis verifies the coherence of regulatory regions, ensuring that

merged gene models maintain consistent promoter and terminator elements. These tools are

accessible via their respective web pages and are not included in the automated pipeline. Users

must extract sequences between inspected genes to check for the presence of promoters and

terminators. After potential merges, manual adjustments are advised to finalize gene models,

ensuring accurate gene length prediction and overall annotation reliability.

While genome annotation typically focuses on protein-coding genes, our pipeline also

includes a step for annotating non-coding RNAs (ncRNAs), which have significant functional

relevance. This step is optional, as it requires downloading the extensive (>20 GB) RNAcentral

database. The ncRNA sequences are first linearized and deduplicated using seqkit rmdup to

remove redundant entries. Homologous regions are then identified in the reference genome using

BLASTn. After the identification step, ncRNA types are classified based on the sequence

identifier. The matching results are then converted to GTF format. A multi-step filtering and

refinement process is used to improve the accuracy of the annotations. This process addresses

several common issues, including multiple alignments at identical genomic positions, inconsistent

naming conventions, and clustering.

The final step integrates all gathered information into a unified genome annotation file,

ready for downstream applications, including both coding and non-coding elements with

comprehensive functional insights. One additional step as part of the genome annotation extension

involves incorporating Gene Ontology (GO) annotations based on the functional annotations

obtained in Part I. However, this requires downloading the large goa_uniprot_all.gaf file (>200 GB

after gunzip). While we provide the necessary code for this integration, it is not included as part

of the automated Part I pipeline.

The bioinformatics tools utilized in each pipeline step, along with their respective versions are

summarized in Table 1.

Table 1. Bioinformatics tools and pipeline steps

Step	Tool	Version
Step 1: Remove rRNA	STAR Minimap2 samtools	2.7.9a
Step 2: Align reads to the reference genome	STAR Minimap2	2.7.9a
Step 3: Assemble genes & transcripts	StringTie2	2.1.1 (short reads) 2.2.1 (mixed reads)
Step 4: Merge reference-based assemblies	StringTie2	2.2.1
Step 5: Merge *de novo* assemblies & create pre-final annotation	StringTie2	2.2.1
Step 6: Filter out transcripts artifacts	bash	
Step 7: Compare isoforms	gffcompare	0.12.6
Step 8: Correct & enhance annotation	AGAT	1.4.1
Step 9: Determine functional attributes	TransDecoder blastp blastx hmmscan	5.7.1 2.15.0+ 2.15.0+ 3.3.2
Step 10: Integrate functional attributes	R	4.4.1

Step 11: Identify overlapping genes & transcripts	R	4.4.1
Step 12: Identify reversed duplicates	R	4.4.1
Step 13: Identify fragmented & chimeric genes	R	4.4.1
Step 14: Analyse promoter & terminator sequences	Promoter 2.0 ARNold	No official version number No official version number
Step 15: Annotate non-coding RNA (ncRNA)	blastn seqkit	2.15.1+ 2.8.2
Step 16: Integrate collected information into a unified genome annotation file	R	4.4.1
Step 17: Incorporate gene ontology (GO)	R	4.4.1
Step 18: Perform final review and quality assurance	R	4.4.1

Identifying fragmented and chimeric artifacts

In the context of genomic annotation, artifacts refer to erroneous gene or transcript models that arise from limitations in the annotation process rather than representing true biological entities. Specifically, we refer here to fragmented artifacts as those that occur when a single gene is incorrectly split into multiple, separate gene models or transcripts. This fragmentation can result from incomplete transcript assembly or sequencing errors, leading to discontinuous or truncated representations of what is biologically a single gene. On the other hand, chimerism typically arises from misassembly during transcriptome reconstruction, especially in regions with high gene density or overlapping gene structures.

Our analysis employs flags such as: *is_potentially_fragmented_de_novo*, *is_potentially_chimeric_due_to_structure_de_novo*, and *is_potentially_chimeric_due_to_function_de_novo* to identify these artifacts based on functional

and structural criteria. Below, we discuss three illustrative examples (depicted in Figure 4) that

highlight the complexities and implications of these annotation challenges. SMESGs here are

reference genes and MSTRGs are putative genes.

Fragmented gene prediction in proximal gene regions

In one genomic region, three genes—SMESG000004200.1, SMESG000004201.1, and

MSTRG.967—are located in close proximity on chromosome 1 (smed_chr1) (Figure 4A).

Specifically, gene SMESG000004200.1 spans position 14,973,263 to 15,003,452 on the positive

strand, while SMESG000004201.1 extends from 15,003,453 to 15,011,292 on the same strand.

StringTie2, predicted an additional transcript, MSTRG.967, which encompasses the regions of

both genes (14,973,256 to 15,011,292). Both SMESG000004200.1 and SMESG000004201.1 were

independently annotated as "Epidermal growth factor receptors." However, the prediction by

StringTie2 that merges these two genes into a single transcript is more biologically plausible. This

is because the close genomic proximity and shared functional annotations suggest a possible gene

fusion or a complex gene structure that is better represented by a single, contiguous transcript.

Flagging these genes as potentially fragmented (is_potentially_fragmented_de_novo) highlights

the necessity to consider merged annotations in densely packed genomic regions to avoid

misinterpretation of gene boundaries.

Chimeric transcript prediction due to excessive intron length

The   gene   SMESG000065897.1   initially   comprised   a   single   transcript,

SMEST065897001.1, spanning positions 325,241,809 to 325,242,781 on chromosome 1 (Figure

4B). However, StringTie2 predicted an additional transcript, MSTRG.16892.1, which extends

dramatically from 325,241,809 to 325,347,749, introducing an intron exceeding 100,000 nt. This

unusually large intron length is atypical and suggests a misassembly, classifying MSTRG.16892.1

as potentially chimeric (is_potentially_chimeric_structure_de_novo).

The presence of such a large intron likely indicates that MSTRG.16892.1 erroneously

combines sequences from multiple genomic loci or fails to accurately represent the true transcript

structure of SMESG000065897.1. Consequently, the chimeric transcript should be disregarded,

and the gene localization should remain based solely on the original transcript

SMEST065897001.1.

Functionally chimeric gene due to divergent transcript annotations

The gene SMESG000077853.1 exhibits a complex transcriptional landscape with multiple

transcripts (Figure 4C). While most transcripts, such as MSTRG.31576.1, MSTRG.31576.5,

MSTRG.31576.6, and all reference SMEST transcripts are annotated as "ankyrin repeat proteins,"

one transcript, MSTRG.31576.11, is annotated as "Squamous cell carcinoma antigen recognized

by T-cells 3 (SART-3)". This divergence in functional annotation across transcripts results in the

gene being flagged as chimeric (is_potentially_chimeric_function_de_novo). These inconsistent

functional annotations suggest that MSTRG.31576.11 may have been incorrectly assembled,

possibly incorporating sequences from a different gene or representing a mispredicted splice

variant. Given that "SART-3" functions are unrelated to "ankyrin repeat proteins," it is evident that

MSTRG.31576.11 does not belong to SMESG000077853.1. Consequently, MSTRG.31576.11

was filtered out.

Detecting missing genes through functional annotation and homology assignments

Functional annotations and homology assignments were then carried out to explain the

biological function of the assembled transcripts and identify orthologous genes by using

established databases and computational tools. This approach played a key role in detecting

missing genes thus highlighting potential gaps and guiding further refinement of the transcriptome

assembly.

To incorporate functional data into our newly developed *Schmidtea mediterranea* genome

annotation (schMed_Pn), we utilized the following databases: Pfam, SwissProt, and non-

redundant (nr) protein sequences from NCBI. Our predicted annotation comprises 25,265 genes,

of which 4,011 are novel and 21,254 are identical to the reference genes. Each gene possesses at

least one function-related entry, either through a database hit or by being classified as having an

open reading frame (ORF). For 18,552 genes, we successfully identified Gene Ontology (GO)

terms using homology-based strategies, which included the pfam2go package, UniProt GOA

conversion, and a custom R script to retrieve GO annotations based on NCBI IDs. Notably, 2,427

of these GO-annotated genes are putative. To evaluate the similarity of GO terms between novel

and reference genes, Venn diagrams were constructed, highlighting shared and unique functional

categories across the gene sets. We assigned a total of 23,909 GO terms to genes within our

predicted annotation. Specifically, 18,586 GO terms were assigned to SMESG (reference) genes,

and 5,323 GO terms were assigned to MSTRG (putative) genes. Of these, 5,180 GO terms were

shared between both gene categories, while 143 GO terms were uniquely assigned to MSTRG

genes (Figure 5A). To visualize the functional distribution of the gene sets, we generated word

clouds representing the most frequent unique GO terms for each group: MSTRG (Figure 5D) and

SMESG (Figure 5B), as well as those that were common for two groups (Figure 5C).

Insights into planarian ncRNA diversity

Beyond refining the annotations of protein-coding genes, we also performed a preliminary

assessment of non-coding RNAs (ncRNAs) in *Schmidtea mediterranea*. Although not an

exhaustive characterization, our analysis reveals a diverse repertoire of regulatory RNAs,

highlighting significant differences between publicly available annotations and newly generated

data. Publicly available datasets from RNAcentral assigned a total of 154 rRNAs, 578 miRNAs

(including 269 precursors), 61 tRNAs, 2 snoRNAs, 12 snRNAs, and 2 SRP RNA hits to *Schmidtea*

mediterranea. After deduplication, these numbers were refined to 129 rRNA, 571 miRNAs (with

156 precursors), 60 tRNAs, 2 snoRNAs, 12 snRNAs, and 2 SRP RNA hits. Additionally, according

to recent literature [42], *Schmidtea* genome contains 457 tRNA hits, which we reduced to 434 after

deduplication. In contrast, our study, utilizing our annotation pipeline, identified 166 rRNA hits

across 7,335 genomic positions, 182 miRNA hits across 1,390 genomic positions, 487 tRNA hits

distributed over 4,408 genomic positions, 2 snoRNA hits across 69 genomic positions, 18 snRNA

hits across 407 genomic positions, 8 miscellaneous RNAs spread across 5,585 genomic positions,

RNase P ncRNA components across 679 genomic positions, 28 lncRNA hits across 326 genomic

positions and 2 sRNA hits across 617 genomic positions. The discrepancy between published

records assigned to *Schmidtea mediterranea* and our annotated RNAs may be due to the fact that

some sequences listed in RNAcentral may be partial or might simply be artifacts, inaccurate

submissions, or too divergent to map unambiguously. Thus, not mapping to a reference genome

does not necessarily imply these entries are universally spurious, but it does highlight the ongoing

need for careful re-annotation and experimental validation in underexplored organisms like

planarians.

In addition, we uncovered 40 copies of signal recognition particle RNAs (SRP_RNAs)

situated on chromosome 3 within an interval flanked by the genes SMESG000020110.1 (BLAST-

predicted as a lysine-specific demethylase) and MSTRG.33721 (BLAST-predicted as containing

a thrombospondin type 3 repeat). Intriguingly, the two protein-coding genes are oriented in the

5′→3′ direction, whereas the SRP_RNAs are oriented 3′→5′. While this arrangement might

suggest a localized regulatory cluster, SRP RNAs known to be evolutionarily conserved and often

found in multiple genomic locations [43]. Their presence in compact regions does not necessarily

imply a functional relationship with adjacent genes. Without further evidence, it is difficult to draw

functional conclusions from their location alone. Experiments measuring SRP_RNA expression

levels, interactions with neighboring genes, or chromatin accessibility in these regions would be

needed to confirm whether these SRP_RNAs play a role in local regulation or are simply part of a

repeated structural element.

Moreover, we identified 10 copies of TERC_RNA located on smed_chr4, which align with

the *Mus musculus* (house mouse) partial telomerase RNA component (TERC). The presence of

TERC_RNA homologous sequences suggests that *Schmidtea mediterranea* may have a functional

TERC. However, *TERT* gene (SMESG000061912.1) is located on smed_chr3, indicating that

TERT and *TERC* are on separate chromosomes. Yet, further analysis is needed to confirm if

telomerase enzyme complex (TERT + TERC) in planarians interact in a functional context.

Benchmarking and genome annotation performance

To ensure the reliability of schMed_Pn genome annotation, we performed thorough

performance benchmarking against existing genome annotations of *Schmidtea mediterranea*.

To assess the completeness and quality of the existing and generated in this study genome

annotations, we analyzed transcript length distribution. It revealed that all annotations

predominantly consist of shorter transcripts, as reflected by a sharp decline in cumulative transcript

counts with increasing transcript length (Figure 6A). Specifically, dd_Smed_v6 showed the

steepest decline in cumulative transcript counts with increasing length, suggesting a higher

fragmentation rate; however, caution is advised in interpreting these distributions, as there is no

universally accepted "standard" shape for planarian transcript length. In contrast, schMed_Pn

demonstrated a more gradual decline, which indicates a broader distribution of transcript lengths.

Assemblies such as schMedS1, schMedS2, schMedS3_1, and schMedS3_2 exhibited moderate

slopes with slight differences, indicating varying transcript length distributions among them. The

SmedAsxl genome_v1.1 displayed a narrower range of longer transcripts. In contrast, SmedGD_c1

and SMESG.1, distributions similar to schMed_Pn, demonstrated relatively smooth distributions,

signifying balanced transcript coverage with a significant proportion of longer transcripts.

The completeness of the transcriptome assembly was evaluated using BUSCO, which

assesses the presence of conserved orthologs, by comparing a transcriptome of interest to those

within the Metazoa lineage (Figure 6B). dd_Smed_v6 exhibited the highest proportion of complete

(single-copy) orthologs, indicating a robust assembly, but also showed moderate levels of

duplicated and missing orthologs, suggesting some redundancy and gaps. schMed_Pn maintained

moderate proportions of both complete single-copy and complete duplicated orthologs while

demonstrating low levels of fragmented orthologs, with missing orthologs comparable to

dd_Smed_v6. Assemblies such as schMedS1, schMedS2, schMedS3_1, and schMedS3_2

generally displayed lower proportions of complete orthologs and higher duplication rates,

particularly in schMedS3_1 and schMedS3_2, indicating possible redundancy. Furthermore,

SMESG.1 achieved high complete orthologs and the lowest fragmentation rates, whereas

SmedAsxl genome_v1.1 had fewer complete orthologs and higher proportions of fragmented and

missing orthologs, highlighting areas that may require further refinement. From a BUSCO

standpoint, SMESG.1 and dd_Smed_v6 both perform very well (albeit with different trade-offs in

duplication and missing orthologs). schMed_Pn also ranks among the stronger annotations,

demonstrating moderate duplication and missing ortholog levels while retaining high

completeness.

In addition, we employed rnaQUAST and Transrate to generate detailed assembly quality

metrics (Table 2). Notably, schMed_Pn exhibited a high number of transcripts over 1,000 nt and

maintained a competitive N50 value (the length of the shortest contig or scaffold covering 50% of

the total assembly), reflecting effective transcript coverage and assembly quality.

Table 2. Comparative metrics for selected transcriptome assemblies of *Schmidtea*

mediterranea

Metric	dd_Smed_v6	schMed_Pn	schMedS1	schMedS2	schMedS3_1	schMedS3_2	SmedAsxl_v1.1	SmedGD_c1.3	SMESG.1
No. of Transcripts	41,745	41,396	22,348	27,79	41,358	41,036	22,855	29,85	31,102
Min transcript length (nt)	201	200	267	267	145	131	75	87	267
Max transcript length (nt)	19,524	48,466	86,373	1,022,601	76,733	94,201	43,226	12,11	48,121
Total Bases (M)	53.23	97.09	40.12	57.93	72.60	71.81	25.55	23.15	65.55

Mean transcript length (nt)	1,275	2,345	1,795	2,084	1,755	1,75	1,114	770	2,108
No. of transcripts >1k	18,487	31,199	16,082	20,727	29,402	28,932	8,61	7,218	23,648
No. of transcripts >10k	94	702	67	293	116	119	47	2	346
No. of transcripts with ORF	20,461	32,798	18,694	23,854	31,503	31,147	12,937	17,567	26,665
N50 (nt)	1,931	3,070	2,159	2,535	2,133	2,141	1,653	1,011	2,570
GC (%)	0.34	0.33	0.33	0.33	0.33	0.33	0.34	0.36	0.33

Performance benchmarking enabled the identification of improvements and areas needing

enhancement while also positioning schMed_Pn within the context of current resources. These

analyses collectively demonstrate that our approach offers a balanced and complete annotation,

effectively capturing a wide range of transcript lengths and maintaining high completeness.

Discussion

Numerous pipelines and tools have been developed to tackle genome annotation across

diverse organisms. For well-studied species—such as *Drosophila melanogaster* or *Arabidopsis*

thaliana—community-maintained platforms (FlyBase [44] and TAIR [45,46] respectively)

provide robust resources that streamline annotation. In prokaryotes, automated tools like Prokka

[10] efficiently handle smaller, less complex genomes, while eukaryotic model organisms often

benefit from RefSeq [7] or large consortia efforts, which incorporate extensive experimental

validation. However, non-model eukaryotes, including many invertebrates and plants, typically

face an incomplete reference, novel repetitive elements, and limited training data for *ab initio*

predictions. Pipelines such as Maker [5,47], Braker [48,49], GGA [9], nf-core GenomeAnnotator

[12], and others can automate workflow portions—especially when reference data or orthologous

gene sets are available. However, in certain species with high genomic complexity or less

functional annotation data, these tools may still underperform or require extensive manual

curation.

In this study, we developed and validated a comprehensive genome annotation pipeline

designed to address the challenges of working with non-classical model organisms, while

generating high-quality annotations applicable to broader genomic research. Our pipeline uniquely

combines: (i) the integration of multiple data types, (ii) automated artifact detection, (iii) dual

approach assembly —reference-based plus *de novo*, and (iv) broad functional integration.

Our analysis identified 4,011 novel genes and multiple previously uncharacterized

isoforms and ncRNAs in the asexual strain, enriching the genomic landscape of *Schmidtea*

mediterranea. Notably, our GO term analysis of putative MSTRG genes revealed enrichment in

diverse biological functions. Among these, "chondroitin 4-sulfotransferase activity" and

"diterpenoid biosynthetic process" suggest that these novel genes might contribute to unique

regulatory pathways or stress responses in the asexual strain. Although many of these assignments

are based on low-frequency observations, the presence of functions related to RNA binding (box

C/D and H/ACA snoRNP complex binding) further underscores the potential regulatory

complexity.

While our transcriptomic data offers strain-specific insights, relying on the genome of the

sexual strain as the reference introduces challenges. Differences between strains, such as

chromosomal translocation and other genetic variations, may compromise the accuracy of gene

prediction and transcript assembly, leading to potential inconsistencies in annotation quality.

Variations in gene expression, alternative splicing events, and structural genomic differences

necessitate meticulous bioinformatics strategies to ensure reliable annotations.

A deeper understanding of the genomic arrangement highlights further complexities that

demand careful interpretation. Genes located in close proximity can reflect various biological

scenarios. One possibility is the presence of functionally independent genes that happen to be

closely spaced, as observed in other genomes where gene density is high [50,51]. For example,

two planarian genes—SMESG000043996.1 and SMESG000043998.1—encode kynureninase and

synaptotagmin XVII, respectively, are unrelated but positioned close to each other, supporting

their independence while reflecting high gene density. Alternatively, such arrangements may arise

from evolutionary events such as gene duplication followed by functional divergence, where both

copies remain in close physical proximity. Another scenario includes operon-like gene structures

commonly seen in prokaryotes but also identified in some eukaryotic contexts [52]. These

structures allow the co-regulation of adjacent genes through shared promoters or other regulatory

elements. Bidirectional promoters, enabling two genes to be transcribed in opposite directions

from the same regulatory region, represent another mechanism driving close gene arrangements

[53].

Non-contiguous transcripts, where exons are unusually spaced or appear fragmented, often

arise from structural variations, alternative splicing, or sequencing artifacts. Biologically, large

introns or dispersed exonic regions can be linked to long-range regulatory elements or structural

genome features such as transposon insertions. Moreover, such patterns may indicate complex

gene models, as seen in genes encoding for proteins with repetitive domains [54,55] or in cases

where alternative promoters and polyadenylation sites lead to diverse transcript forms [56,57].

Consequently, accurately capturing these intricate gene architectures often requires more

comprehensive strategies than those employed in simpler genomes, underscoring the need for user

scrutiny and flexibility in annotation pipelines. The primary advantage of our pipeline lies in its

capacity to address these complexities. By combining reference-based and *de novo* methods and

incorporating both coding and non-coding RNA annotations it addresses the challenges posed by

limited reference genomes. However, the pipeline's reliance on user-defined thresholds and

manual curation introduces subjectivity, potentially impacting reproducibility.

The existing, planarian-specific repositories—such as PlanMine [21], PLANAtools [58],

WomBase [59], PlanoSphere [60], PlanNET [61], and various planaria single-cell atlas datasets

[62–65]—provide essential resources for gene model comparisons, single-cell data, and functional

information in *S. mediterranea*. Despite their value, differences in naming conventions, annotation

updates, and coverage across sexual and asexual strains can lead to inconsistencies. These

discrepancies highlight the importance of robust, well-documented pipelines that can reconcile or

supplement these external databases, thus ensuring more accurate gene models.

Additionally, while our automated steps effectively identify fragmented and chimeric

transcripts, resolving these artifacts requires specialized expertise and is not fully automated.

Another limitation concerns functional annotations, which can lead to false positives in identifying

chimeric genes. For instance, the same protein from different species or proteins representing

different subunits of a complex may be annotated as distinct entries in functional databases. This

can lead to genes being incorrectly flagged as chimeric if the functional entries do not exactly

match. As a result, refined algorithms that account for near-identical annotations and minor

terminological differences are vital.

Future enhancements could include the integration of machine learning algorithms to

automate artifact resolution and improve annotation consistency. Expanding the pipeline to

support multi-omics data integration, such as proteomics and epigenomics, would further enhance

its utility. Community-driven efforts to standardize annotation protocols and databases would

ensure broader applicability and reproducibility. Finally, experimental validation of novel genes

and isoforms identified through this pipeline will be critical to confirm their biological relevance.

Compared to pipelines that excel in either prokaryotic or model vertebrate annotation –

where reference data is abundant and well-curated–, our approach fills the gap for non-classical

model systems with more limited resources. In doing so, it advances genome annotation efforts for

Schmidtea mediterranea and provides a robust framework for addressing similar challenges in

other species, fostering broader genomic discoveries.

Availability of Source Code and Requirements

Project name: SmedAnno

Project homepage: https://github.com/Norreanea/SmedAnno

Operating system: Linux

Programming languages: shell, R, Python, awk

License: MIT License

Data Availability

The *in-silico* datasets supporting the results and all other code used in this article are available in

the GitHub repository archived in Zenodo https://doi.org/10.5281/zenodo.15130394. RNA-seq

data can be accessed from the following GEO repositories: GSE293999 and GSE293997 from

NCBI BioProject PRJNA1243063; and GSE294569 from NCBI BioProject PRJNA1246370. The

final genome annotation file is available in GSE294569.

List of Abbreviations

BP: base pairs; NT: nucleotide; RNA: ribonucleic acid; rRNA: ribosomal RNA; ncRNA: non-

coding RNA; SIRV: Spike-In RNA Variants; ORF: open reading frame; GTF: Gene Transfer

Format; BLAST: Basic Local Alignment Search Tool; STAR: Spliced Transcripts Alignment to a

Reference; BUSCO: Benchmarking Universal Single-Copy Orthologs; GO: Gene Ontology;

cDNA: complementary DNA; DNA: deoxyribonucleic acid; RIN: RNA Integrity Number; GGA:

Galaxy Genome Annotation.

Competing Interests

Nothing to declare

Declaration of generative AI and AI-assisted technologies in the writing process

During the preparation of this work, the authors used the free version of OpenAI-ChatGPT c2022 (v.GPT-o3-mini, February 2025) in order to improve manuscript readability and language. After using this tool, the authors thoroughly reviewed and modified the manuscript and took full responsibility for the content of the publication. The literature survey was done only by the authors, with no input from the AI tools.

Funding

This work was financed by the National Science Centre, Poland, through a grant no. 2019/35/B/NZ2/02658 to PJ.

Authors' Contributions

AZ: Conceptualization, Methodology, Software, Visualization, Writing – Original Draft, Writing – Review & Editing; MMZ: Conceptualization, Methodology, Writing – Review & Editing;

AD: Investigation, Writing – Review & Editing; ASCz: Investigation, Writing – Review & Editing; PJ: Conceptualization, Supervision, Project administration, Funding Acquisition, Writing – Review & Editing.

References

1. Stanke M, Keller O, Gunduz I, Hayes A, Waack S, Morgenstern B. AUGUSTUS: A b initio

prediction of alternative transcripts. *Nucleic Acids Res*. 2006; doi: 10.1093/nar/gkl200.

2. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al.. BLAST+:

Architecture and applications. *BMC Bioinformatics*. 2009; doi: 10.1186/1471-2105-10-421.

3. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al.. STAR: Ultrafast

universal RNA-seq aligner. *Bioinformatics*. 2013; doi: 10.1093/bioinformatics/bts635.

4. Shumate A, Wong B, Pertea G, Pertea M. Improved transcriptome assembly using a hybrid of

long and short reads with StringTie. *PLoS Comput Biol*. Public Library of Science; 2022; doi:

10.1371/journal.pcbi.1009730.

5. Cantarel BL, Korf I, Robb SMC, Parra G, Ross E, Moore B, et al.. MAKER: An easy-to-use

annotation pipeline designed for emerging model organism genomes. *Genome Res*. 2008; doi:

10.1101/gr.6743907.

6. Haas BJ, Salzberg SL, Zhu W, Pertea M, Allen JE, Orvis J, et al.. Automated eukaryotic gene

structure annotation using EVidenceModeler and the Program to Assemble Spliced Alignments.

Genome Biol. 2008; doi: 10.1186/gb-2008-9-1-r7.

7. O'Leary NA, Wright MW, Brister JR, Ciufo S, Haddad D, McVeigh R, et al.. Reference

sequence (RefSeq) database at NCBI: Current status, taxonomic expansion, and functional

annotation. *Nucleic Acids Res*. Oxford University Press; 2016; doi: 10.1093/nar/gkv1189.

8. Kurylo C, Guyomar C, Foissac S, Djebali S. TAGADA: a scalable pipeline to improve genome

annotations with RNA-seq data. *NAR Genom Bioinform*. Oxford University Press; 2023; doi:

10.1093/nargab/lqad089.

9. Hiltemann S, Rasche H, Gladman S, Hotz HR, Larivière D, Blankenberg D, et al.. Galaxy

Training: A powerful framework for teaching! *PLoS Comput Biol*. Public Library of Science;

2023; doi: 10.1371/journal.pcbi.1010752.

10. Seemann T. Prokka: Rapid prokaryotic genome annotation. *Bioinformatics*. Oxford University

Press; 2014; doi: 10.1093/bioinformatics/btu153.

11. Swain MT, Tsai IJ, Assefa SA, Newbold C, Berriman M, Otto TD. A post-assembly genome-

improvement toolkit (PAGIT) to obtain annotated genomes from contigs. *Nat Protoc*. 2012; doi:

10.1038/nprot.2012.068.

12. Ewels PA, Peltzer A, Fillinger S, Patel H, Alneberg J, Wilm A, et al.. The nf-core framework

for community-curated bioinformatics pipelines. *Nat Biotechnol*. 2020; doi: 10.1038/s41587-020-

0439-x.

13. Wu S, Yuan S, Su Z. User-friendly genome assembly and gene annotation pipelines for

vertebrates. bioRxiv. 2024; doi: 10.1101/2024.05.21.595213

14. Maia GA, Filho VB, Kawagoe EK, Teixeira Soratto TA, Moreira RS, Grisard EC, et al..

AnnotaPipeline: An integrated tool to annotate eukaryotic proteins using multi-omics data. *Front*

Genet. 2022; doi: 10.3389/fgene.2022.1020100.

15. Humann JL, Lee T, Ficklin S, Main D. Structural and Functional Annotation of Eukaryotic

Genomes with GenSAS. *Gene Prediction*. 2019; doi: 10.1007/978-1-4939-9173-0_3

16. Sánchez Alvarado A. The freshwater planarian Schmidtea mediterranea: Embryogenesis, stem

cells and regeneration. *Curr Opin Genet Dev*. Elsevier Ltd; 2003; doi: 10.1016/S0959-

437X(03)00082-0

17. Baguñà J, Carranza S, Pala M, Ribera C, Giribet G, Arnedo MA, et al.. From morphology and

karyology to molecules. New methods for taxonomical identification of asexual populations of

freshwater planarians. A tribute to Professor Mario Benazzi. *Italian Journal of Zoology*. Societa

Paleontologica Italiana; 1999; doi: 10.1080/11250009909356258.

18. Robb SMC, Ross E, Alvarado AS. SmedGD: The Schmidtea mediterranea genome database.

Nucleic Acids Res. 2008; doi: 10.1093/nar/gkm684.

19. Robb SMC, Gotting K, Ross E, Sánchez Alvarado A. SmedGD 2.0: The Schmidtea

mediterranea genome database. *Genesis*. John Wiley and Sons Inc.; 2015; doi: 10.1002/dvg.22872.

20. Grohme MA, Schloissnig S, Rozanski A, Pippel M, Young GR, Winkler S, et al.. The genome

of Schmidtea mediterranea and the evolution of core cellular mechanisms. *Nature*. Nature

Publishing Group; 2018; doi: 10.1038/nature25473.

21. Rozanski A, Moon HK, Brandl H, Martín-Durán JM, Grohme MA, Hüttner K, et al.. PlanMine

3.0 - improvements to a mineable resource of flatworm biology and biodiversity. *Nucleic Acids

Res*. Oxford University Press; 2019; doi: 10.1093/nar/gky1070.

22. Guo L, Bloom JS, Dols-Serrate D, Boocock J, Ben-David E, Schubert OT, et al.. Island-specific

evolution of a sex-primed autosome in a sexual planarian. *Nature*. Nature Research; 2022; doi:

10.1038/s41586-022-04757-3.

23. Ivanković M, Brand JN, Pandolfini L, Brown T, Pippel M, Rozanski A, et al.. A comparative

analysis of planarian genomes reveals regulatory conservation in the face of rapid structural

divergence. *Nat Commun*. 2024; doi: 10.1038/s41467-024-52380-9.

24. Rabiasz A, Ziętkiewicz E. Schmidtea mediterranea as a Model Organism to Study the

Molecular Background of Human Motile Ciliopathies. Int J Mol Sci. Multidisciplinary Digital

Publishing Institute (MDPI); *Int J Mol Sci* . 2023; doi: 10.3390/ijms24054472

25. Quinodoz S, Thomas MA, Dunkel J, Schötz EM. The More the Merrier?: Entropy and Statistics

of Asexual Reproduction in Freshwater Planarians. *J Stat Phys*. Springer New York LLC; 2011;

doi: 10.1007/s10955-011-0157-3.

26. Reddien PW. The Cellular and Molecular Basis for Planarian Regeneration. *Cell*. Cell Press;

2018; doi: 10.1016/j.cell.2018.09.021

27. Reddien PW, Sánchez Alvarado A. Fundamentals of planarian regeneration. *Annu Rev Cell*

Dev Biol. 2004; doi: 10.1146/annurev.cellbio.20.010403.095114

28. Benham-Pyle BW, Mann FG, Brewster CE, Dewars ER, Vuu DM, Nowotarski SH, et al..

Planarians employ diverse and dynamic stem cell microenvironments to support whole-body

regeneration. bioRxiv. 2023; doi: 10.1101/2022.03.20.485025

29. Sayers EW, Bolton EE, Brister JR, Canese K, Chan J, Comeau DC, et al.. Database resources

of the national center for biotechnology information. *Nucleic Acids Res*. Oxford University Press;

2022; doi: 10.1093/nar/gkab1112.

30. Palmieri N, Schlötterer C. Mapping accuracy of short reads from massively parallel sequencing

and the implications for quantitative expression profiling. *PLoS One*. 2009; doi:

10.1371/journal.pone.0006323.

31. Sedlazeck FJ, Lee H, Darby CA, Schatz MC. Piercing the dark matter: bioinformatics of long-

range sequencing and mapping. *Nat Rev Genet*. 2018; doi: 10.1038/s41576-018-0003-4.

32. Chen Z, Ain NU, Zhao Q, Zhang X. From tradition to innovation: conventional and deep

learning frameworks in genome annotation. *Brief Bioinformatics*. Oxford University Press; 2024;

doi: 10.1093/bib/bbae138

33. Newmark PA, Sánchez Alvarado A. Bromodeoxyuridine specifically labels the regenerative

stem cells of planarians. *Dev Biol*. Academic Press Inc.; 2000; doi: 10.1006/dbio.2000.9645.

34. Dainat J, Hereñú D, Dr. Murray KD, Davis E, Ugrin I, Crouch K, et al.. NBISweden/AGAT:

AGAT-v1.4.1 (v1.4.1). Zenodo;

35. Simão FA, Waterhouse RM, Ioannidis P, Kriventseva E V., Zdobnov EM. BUSCO: Assessing

genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics*.

Oxford University Press; 2015; doi: 10.1093/bioinformatics/btv351.

36. Bushmanova E, Antipov D, Lapidus A, Suvorov V, Prjibelski AD. RnaQUAST: A quality

assessment tool for de novo transcriptome assemblies. *Bioinformatics*. Oxford University Press;

2016; doi: 10.1093/bioinformatics/btw218.

37. Smith-Unna R, Boursnell C, Patro R, Hibberd JM, Kelly S. TransRate: Reference-free quality

assessment of de novo transcriptome assemblies. *Genome Res*. Cold Spring Harbor Laboratory

Press; 2016; doi: 10.1101/gr.196469.115.

38. Sweeney BA, Petrov AI, Ribas CE, Finn RD, Bateman A, Szymanski M, et al.. RNAcentral

2021: secondary structure integration, improved sequence search and new member databases.

Nucleic Acids Res. 2021; doi: 10.1093/nar/gkaa921.

39. Francis WR, Wörheide G. Similar ratios of introns to intergenic sequence across animal

genomes. *Genome Biol Evol*. Oxford University Press; 2017; doi: 10.1093/gbe/evx103.

40. Gudlaugsdottir S, Boswell DR, Wood GR, Ma J. Exon size distribution and the origin of

introns. *Genetica*. 2007; doi: 10.1007/s10709-007-9139-4.

41. Bhasi A, Philip P, Manikandan V, Senapathy P. ExDom: An integrated database for

comparative analysis of the exon-intron structures of protein domains in eukaryotes. *Nucleic Acids*

Res. 2009; doi: 10.1093/nar/gkn746.

42. Lakshmanan V, Sujith TN, Bansal D, Shivaprasad P V., Palakodeti D, Krishna S.

Comprehensive annotation and characterization of planarian tRNA and tRNA-derived fragments

(tRFs). *RNA*. 2021; doi: 10.1261/rna.077701.120.

43. Larsen N, Zwieb C. SRP-RNA sequence alignment and secondary structure. *Nucleic Acids*

Res. 1991; doi: 10.1093/nar/19.2.209.

44. Öztürk-Çolak A, Marygold SJ, Antonazzo G, Attrill H, Goutte-Gattat D, Jenkins VK, et al..

FlyBase: updates to the *Drosophila* genes and genomes database. *Genetics*. 2024; doi:

10.1093/genetics/iyad211.

45. Berardini TZ, Reiser L, Li D, Mezheritsky Y, Muller R, Strait E, et al.. The arabidopsis

information resource: Making and mining the "gold standard" annotated reference plant genome.

genesis. 2015; doi: 10.1002/dvg.22877.

46. Cheng C, Krishnakumar V, Chan AP, Thibaud-Nissen F, Schobel S, Town CD. Araport11: a

complete reannotation of the *Arabidopsis thaliana* reference genome. *The Plant Journal*. 2017;

doi: 10.1111/tpj.13415.

47. Holt C, Yandell M. MAKER2: an annotation pipeline and genome-database management tool

for second-generation genome projects. *BMC Bioinformatics*. 2011; doi: 10.1186/1471-2105-12-

491.

48. Hoff KJ, Lomsadze A, Borodovsky M, Stanke M. Whole-Genome Annotation with BRAKER.

Methods Mol Biol . 2019; doi: 10.1007/978-1-4939-9173-0_5

49. Brůna T, Hoff KJ, Lomsadze A, Stanke M, Borodovsky M. BRAKER2: automatic eukaryotic

genome annotation with GeneMark-EP+ and AUGUSTUS supported by a protein database. *NAR*

Genom Bioinform. 2021; doi: 10.1093/nargab/lqaa108.

50. Arnone JT, Robbins-Pianka A, Arace JR, Kass-Gergi S, McAlear MA. The adjacent

positioning of co-regulated gene pairs is widely conserved across eukaryotes. *BMC Genomics*.

2012; doi: 10.1186/1471-2164-13-546.

51. Gherman A, Wang R, Avramopoulos D. Orientation, distance, regulation and function of

neighbouring genes. *Hum Genomics*. 2009; doi: 10.1186/1479-7364-3-2-143.

52. Kominek J, Doering DT, Opulente DA, Shen XX, Zhou X, DeVirgilio J, et al.. Eukaryotic

Acquisition of a Bacterial Operon. *Cell*. Cell Press; 2019; doi: 10.1016/j.cell.2019.01.034.

53. Ahmad SS, Samia NSN, Khan AS, Turjya RR, Khan MAAK. Bidirectional promoters: an

enigmatic genome architecture and their roles in cancers. *Mol Biol Rep*. Springer Science and

Business Media B.V.; 2021; doi: 10.1007/s11033-021-06612-6.

54. Hoock TC, Peters LL, Lux SE. Isoforms of Ankyrin-3 That Lack the NH2-terminal Repeats

Associate with Mouse Macrophage Lysosomes. *J Cell Biol*. 1997; doi: 10.1083/jcb.136.5.1059.

55. Yamankurt G, Wu HC, McCarthy M, Cunha SR. Exon Organization and Novel Alternative

Splicing of Ank3 in Mouse Heart. *PLoS One*. 2015; doi: 10.1371/journal.pone.0128177.

56. Aísa-Marín I, García-Arroyo R, Mirra S, Marfany G. The Alter Retina: Alternative Splicing

of Retinal Genes in Health and Disease. *Int J Mol Sci*. 2021; doi: 10.3390/ijms22041855.

57. Pal S, Gupta R, Kim H, Wickramasinghe P, Baubet V, Showe LC, et al.. Alternative

transcription exceeds alternative splicing in generating the transcriptome diversity of cerebellar

development. *Genome Res*. 2011; doi: 10.1101/gr.120535.111.

58. Hoffman M, Wurtzel O. PLANAtools—An interactive gene expression repository for the

planarian Schmidtea mediterranea. *Front Cell Dev Biol*. 2023; doi: 10.3389/fcell.2023.1149537.

59. Sternberg PW, Van Auken K, Wang Q, Wright A, Yook K, Zarowiecki M, et al.. WormBase

2024:   status   and   transitioning   to   Alliance   infrastructure.   *Genetics*.   2024;   doi:

10.1093/genetics/iyae050.

60. Nowotarski SH, Davies EL, Robb SMC, Ross EJ, Matentzoglu N, Doddihal V, et al.. Planarian

Anatomy Ontology: a resource to connect data within and across experimental platforms.

Development. 2021; doi: 10.1242/dev.196097.

61. Castillo-Lara S, Abril JF. PlanNET: homology-based predicted interactome for multiple

planarian transcriptomes. *Bioinformatics*. 2018; doi: 10.1093/bioinformatics/btx738.

62. Cui G, Dong K, Zhou J-Y, Li S, Wu Y, Han Q, et al.. Spatiotemporal transcriptomic atlas

reveals the dynamic characteristics and key regulators of planarian regeneration. *Nat Commun*.

2023; doi: 10.1038/s41467-023-39016-0.

63. King HO, Owusu-Boaitey KE, Fincher CT, Reddien PW. A transcription factor atlas of stem

cell fate in planarians. *Cell Rep*. 2024; doi: 10.1016/j.celrep.2024.113843.

64. Fincher CT, Wurtzel O, de Hoog T, Kravarik KM, Reddien PW. Cell type transcriptome atlas

for the planarian *Schmidtea mediterranea*. *Science (1979)*. 2018; doi: 10.1126/science.aaq1736.

65. Plass M, Solana J, Wolf FA, Ayoub S, Misios A, Glažar P, et al.. Cell type atlas and lineage

tree of a whole complex animal by single-cell transcriptomics. *Science (1979)*. 2018; doi:

10.1126/science.aaq1723.

Figure Legends

Figure 1. **Identification rates and accuracy metrics for *in-silico* and SIRVs datasets.**

(A) - Transcript identification rates using *in-silico* data. Circles depict the four approaches tested

for short read (SR) data, using *de novo* (DN) or reference-based (RB) assembly and StringTie2

version 2.1.1 (211) or version 2.2.1 (221). (B) - Accuracy metrics of transcript isoform

identification using SIRVs on SR, mixed reads (MR) and merged reads (MeR) approaches.

Whiskers on the bar plots represent the variability in accuracy across five different StringTie2

parameter configurations tested.

Figure 2. **Workflow diagram for RNA-Seq pre-processing, transcript assembly, and**

functional annotation.

Figure 3. **User-driven inspection and refinement.**

Figure 4. **Annotation artifacts due to errors.**

(A) - Potentially fragmented gene scenario with SMESG000004200.1, SMESG000004201.1, and

MSTRG.967. (B) - The chimeric transcript prediction in SMESG000065897.1 due to an

excessively large intron in MSTRG.16892.1. (C) - The functionally chimeric gene

SMESG000077853.1 with divergent transcript annotations.

Figure 5. **Venn diagram and a word cloud of GO terms.**

(A) - Venn diagram of GO term overlaps between MSTRG (novel genes) and SMESG (reference

genes). (B) - Word cloud of unique GO terms assigned to SMESG (reference) genes. (C) - Word

cloud of common GO terms assigned to both SMESG and MSTRG genes. (D) - Word cloud of

unique GO terms assigned to MSTRG (novel) genes.

Figure 6. **Performance metrics and annotation features of transcriptome assemblies.**

(A) - Cumulative distribution of transcript lengths in selected annotations. The y-axis (logarithmic

scale) represents the cumulative count of transcripts/isoforms, indicating how many transcripts

exceed each length threshold on the x-axis (linear scale). (B) - BUSCO evaluation of unigene

assembly completeness.

Additional Files

Supplementary Table S1. Sequencing samples summary.

Supplementary Table S2. StringTie2 parameters tested on SIRVs.

Step 14: Analyse promoter & terminator sequences

Promoter 2.0

Extract intergenic sequences → Analyze promoter regions to verify regulatory coherence

ARNold

Analyze terminator regions to ensure consistency in gene models

Inspect promoter and terminator analysis results to determine if gene models should be merged

- Multiple alignments at identical genomic positions
- Inconsistent naming conventions
- Clustering of similar ncRNAs

Adress issues

Step 15: Annotate non-coding RNA (ncRNA)

seqkit rmdup

BLASTn

Download the RNAcentral database → Linearize → Deduplicate → Identify homologous regions in the reference genome → Classify ncRNA types → Convert alignment results to GTF format → Filter and refine ncRNA annotations

Step 16: Integrate collected information into a unified genome annotation file

Merge protein-coding gene annotations with ncRNA annotations

Step 17: Incorporate gene ontology (GO)

Download the GOA Uniprot database → Integrate GO annotations based on the functional annotations

Step 18: Perform final review and quality assurance

Validate the annotation using external databases or experimental data

A

SMESG

13403
(71.6%)

5180
(27.7%)

143
(0.8%)

MSTRG

B

Unique GO terms in SMESG genes

mitotic S phase
modulation by virus of host process
anchoring junction   axon development
fatty acid metabolic process   integral component of plasma membrane
RNA metabolic process   intracellular anatomical structure
cellular protein metabolic process   integral component of membrane   hemopoiesis
multicellular organism development   transport   viral process
kinesin binding   response to drug
microtubule associated complex
Golgi to plasma membrane protein transport
paranode region of axon

C

Common GO terms

endoplasmic reticulum
cytoskeleton   signal transduction
axon   mitochondrion   protein binding
metal ion binding   nucleus
nucleolus   plasma membrane   RNA binding   ATPase
centrosome   nucleoplasm   cytoplasm   cytosol   ATP binding
DNA binding   membrane   dendrite
synapse   zinc ion binding
identical protein binding   extracellular exosome
nucleic acid binding

D

Unique GO terms in MSTRG genes

8-hydroxy-dADP phosphatase activity
rRNA(pseudouridine-N3-)-methyltransferaseactivity
alginic acid biosynthetic process   dUMP biosynthetic process
ether hydrolase activity   archaealorbacterial-typeflagellum-dependentcellmotility
dUTP catabolic process   [protein-PII] uridylyltransferase activity
hyphal cell wall   viral DNA genome packaging   diterpenoid biosynthetic process
BRISC complex   terpene synthase activity
chondroitin 4-sulfotransferase activity
pons development   [heparansulfate]-glucosamine3-sulfotransferase1activity