A modular pipeline for evidence-integrated genome annotation across species: a case study

on *Schmidtea mediterranea*

Anastasiia Zaremba, Małgorzata Marszałek-Zeńczak, Annasha Dutta, Anna Samelak-Czajka,

Paulina Jackowiak#

Laboratory of Single Cell Analyses; Institute of Bioorganic Chemistry Polish Academy of

Sciences; Zygmunta Noskowskiego str. 12/14, 61-704 Poznań; Poland

azaremba@ibch.poznan.pl,   marszalekgosia@gmail.com,   adutta@ibch.poznan.pl,

asczajka@ibch.poznan.pl,

#Corresponding author: paulinaj@ibch.poznan.pl

Abstract

Despite advancements in genome annotation tools, challenges persist for non-classical

model organisms with limited genomic resources, such as *Schmidtea mediterranea*. To address

these challenges, we developed a flexible and scalable genome annotation pipeline that integrates

short-read (Illumina) and long-read (PacBio) sequencing technologies. The pipeline combines

reference-based and *de novo* assembly methods, effectively handling genomic variability and

alternative splicing events. To improve splice site detection accuracy, DeepSplice deep learning

predictions are used. Functional annotation is conducted to filter out low-confidence transcripts

and ensure biological relevance. Applying this pipeline to the asexual strain of *S. mediterranea*

revealed thousands of previously undescribed putative genes and transcripts, and improved the

existing gene models, highlighting its utility in annotating complex, underexplored genomes. The

modularity and comprehensiveness of our pipeline ensure its adaptability for genome annotation

across diverse species, making it a valuable tool for annotating genomes of non-model organisms

and supporting broader genomic research. The source code and implementation details are

available at https://github.com/Norreanea/SmedAnno.

Keywords

Genome annotation; Transcriptome assembly; *Schmidtea mediterranea*; Non-model organisms;

Functional annotation

Background

Understanding the genetic makeup of organisms is crucial for unraveling the

complexities of their biology, making genome annotation a foundational aspect of molecular and

genomic research. By precisely identifying coding regions, non-coding elements, and regulatory

sequences, genome annotation transforms raw genomic data into a practical blueprint that opens

up possibilities for studying living organisms at the molecular level.

Existing genome annotation methods can be broadly categorized into *ab initio* [1],

homology-based [2], evidence-based (reference-based) [3], and hybrid approaches [4–6], each

leveraging different strategies and tools to predict and validate genomic features. Each category

has its unique advantages and challenges. *Ab initio* methods, which predict genes solely from

intrinsic sequence signals without external evidence, are capable of identifying novel genes;

however, they tend to produce a higher rate of false positives and inaccurate gene models—such

as overpredicted exons, missed genes, and incorrect exon–intron boundaries—especially when

applied to complex or poorly characterized genomes. Homology☐based methods use

evolutionary conservation to achieve high accuracy for well☐conserved genes but often miss

novel or rapidly evolving genes that lack close homologs. Evidence☐based approaches utilize

transcriptomic or proteomic data to support gene models, thereby enhancing annotation quality

for expressed genes, yet they are inherently limited to regions where experimental data exist and

are as robust as the underlying evidence. Hybrid approaches combine intrinsic signals with

external evidence in an effort to balance sensitivity and specificity.

To streamline the annotation process, several genome annotation platforms and pipelines

have been developed. For instance, the RefSeq eukaryotic annotation pipeline at NCBI [7]

employs highly curated alignment-based evidence and gene prediction models, while another

recently reported pipeline TAGADA [8] aims to integrate evidence from multiple data sources

for eukaryotic genome annotation. Web-based platforms such as Galaxy Genome Annotation

(GGA) [9] provide an accessible interface for constructing annotation workflows, and tools like

Prokka [10] offer rapid prokaryotic annotation that automates the integration of gene prediction

and functional annotation, generating standardized output formats compatible with GenBank.

Additionally, PAGIT (Post Assembly Genome Improvement Toolkit) [11], the community-

driven Nextflow-based approach nf-core GenomeAnnotator [12], and other pipelines [13–15]

expand the available toolkit.

Despite these advancements, applying genome annotation methods to non-model or non-

classical model organisms presents significant challenges. Due to limited reference data, tools

must rely more heavily on *ab initio* methods, which—as noted above— are prone to errors,

particularly for complex genomes, and often lack sufficient training data. Repetitive sequences

and the presence of multiple isoforms complicate the annotation process. Genetic diversity and

evolutionary distance from well-studied species can impede the identification of orthologous

genes and conserved pathways, while limited experimental data often constrain the ability to

validate or refine gene models. Such constraints are further compounded by annotation

artifacts—including fragmented and chimeric transcripts—that can obscure true biological

functions and introduce errors into downstream analyses.

These challenges are particularly relevant to research on the sexual (S2F2 clonal line) and

asexual (CIW4) strains of *S. mediterranea*, a highly regenerative flatworm [16,17]. Currently,

reference genome sequences and comprehensive annotations are available primarily for the

sexual strain, including contig-level genome assemblies such as SmedGD_c1.3 [18,19],

dd_Smes_g4 [20,21], chromosome-scale genome assemblies schMedS2 [22] and recent

schMedS3 [23], as well as corresponding annotations like SmedGD_c1.3, high-confidence

SMESG, schMedS1, schMedS2, schMedS3_1 and schMedS3_2. Chromosome□scale assemblies

of the sexual strain indicate that sexual and asexual *S. mediterranea* share high genomic

similarity—they are both diploid with four pairs of chromosomes. Although the asexual strain

possesses a chromosomal translocation between chromosomes 1 and 3, these reference

sequences and annotations offer a strong starting point for transcriptome analysis. Nonetheless,

given potential strain□specific transcripts and structural variations, further refinement of the

asexual strain's transcript set is essential [24]. Additionally, the asexual strain exhibits a

simplified reproductive strategy [25–28] and it has been widely used in numerous studies, with

extensive data deposited in public repositories. As of January 2, 2025, there are 122 DNA and

1,694 RNA datasets on NCBI [29] for this strain, the majority of which are short-read data. In

comparison, for the sexual strain, there are only 18 DNA and 87 RNA datasets. The

predominance of RNA-seq data highlights the extensive transcriptomic resources available for

the asexual strain, which offers a strong foundation for gene expression and functional studies

and underscores the potential for leveraging RNA sequencing to enhance genome annotation

efforts. However, limited detailed reference resources for asexual strain remain a significant

challenge. Currently, the scientific community must rely mostly on reference transcriptomic

data—such as dd_Smed_v6.pcf.contigs.fasta [21] (a transcriptome assembly generated from

RNA☐Seq data) and a contig☐based genome assembly and annotation SmedAsxl_genome_v1.1

[19]. Beyond that, existing transcriptome assemblies are fragmented and incomplete, lacking the

high-resolution insights offered by newer sequencing technologies. To bridge this gap, our study

aims to implement a sophisticated bioinformatics pipeline to improve annotated genes and

transcripts in asexual *S. mediterranea* using the existing chromosome-scale genome of the sexual

strain (schMedS2) as a reference and in-house generated comprehensive collection of

transcriptomic data for asexual worms. Our dataset includes both short-read and long-read data,

providing a significant advantage in resolving complex isoforms. Addressing these challenges

requires the integration of various sequencing technologies and advanced bioinformatics tools.

Short-read sequencing platforms, such as Illumina, offer high accuracy and depth of coverage, a

low error rate, which is essential for identifying single-nucleotide polymorphisms, small indels,

and importantly, splice sites—ensuring accurate distinction of exon-intron boundaries [30].

Conversely, long-read technologies like PacBio provide valuable information on structural

variants and complex transcript isoforms, enhancing the resolution of genome assemblies

[31,32]. Combining these technologies benefits from their complementary strengths, enabling a

more comprehensive and accurate genome annotation [4].

Here we present a universal, modular, and adaptable pipeline for genome annotation to

process and interpret the vast amounts of sequencing data generated, that integrates both short-

and long-read sequencing data, employs state-of-the-art bioinformatics tools for transcriptome

assembly and genome characterization, and incorporates robust methods for functional and

homology annotation. By doing so, we aim to advance genome annotation efforts within *S.*

mediterranea and to provide a scalable framework applicable to other non-model organisms,

fostering broader genomic research and discovery.

Methods

Planarian culturing and sample collection

121   Asexual *S. mediterranea* were cultured as described in [33] and fed chicken liver twice

122   weekly, and were used either in wild-type form or after RNAi treatment, as detailed in Table S1.

123   Animals were collected, immediately flash☐frozen in liquid nitrogen to preserve RNA integrity,

124   and stored at −80☐°C until RNA isolation.

Dataset details

Full *S. mediterranea* dataset☐specific details including experimental treatments, RNA

extraction, quality control, are provided in Table☐S1 and in the associated GEO submissions.

Table S1 lists transcriptomic cDNA libraries from the asexual CIW4 strain of *S. mediterranea*: 3

Illumina short-read runs (2 on NextSeq 550, 1 on NovaSeq 6000) and 3 PacBio long-read runs (1

on Sequel II, 2 on Sequel IIe). Publicly available short-read Illumina RNA-seq data for

Arabidopsis thaliana (PRJNA477336) and *Caenorhabditis elegans* (CRA008476) were retrieved

from https://www.ncbi.nlm.nih.gov/ and https://ngdc.cncb.ac.cn/gsa, respectively.

Validation with simulated data

A synthetic genome was created using a custom Python script, which constructed a

single chromosome spanning 2 mega base pairs (bp). This genome included 1,000 genes

randomly distributed along the chromosome, each comprising between 2 and 20 exons with

lengths ranging from 100 to 1,000 nucleotides (nt) and introns of 100 to 1,000 nt. Paired-end

short-read RNA-seq data were simulated using ART Illumina tool, generating 100 bp reads at

10× coverage across six samples. Additionally, long-read sequencing data were produced using

PBSIM tool with 150 bp   reads at 10× coverage for one sample. To evaluate the pipeline's

ability to handle incomplete annotations, a random subset of genes was removed from the

annotation file using the custom bash script, resulting in a GTF file with missing gene

information.

Reference annotation polishing

The reference genome schMedS2 [21] was utilized for this study. High-confidence

annotations for this genome were obtained from two sources: SmedSxl smed_chr_ref_v1 UCLA

(SIMRBase (https://simrbase.stowers.org)) and schMedS2_PlanMine_high-confidence_liftover

(PlanMine [21]). Both annotations are compatible with the genome sequence; however, due to

minor differences between them, the genes were first unified, and the annotation was carefully

reviewed and corrected. Through detailed analysis and comparison, 27,676 SMEST transcripts

and 21,332 SMESG genes were identified as the most confident and consistent across both

sources. Next, 169 transcripts with identical exon structures using the AGAT tool [34] were

removed to eliminate redundancy and enhance annotation quality.

Benchmarking against BRAKER3

Input data

Illumina RNA-seq reads trimmed with TrimGalore were aligned to reference genome with

STAR v2.7.11a [3]. For *A. thaliana* and *C. elegans*, the following reference genomes and

annotations were used: GCF_000001735.4_TAIR10.1 [35,36] and PRJNA13758/ WBcel235

(WBPS19) [37,38] , respectively.

BRAKER3 test

BRAKER3 v2.1.8 [39]was launched in "RNA-seq only" mode: braker.pl --genome genome.fa --

bam star.bam --softmasking --threads 8.

SmedAnno test

Four runs were produced: no guide and no TransDecoder [40] (-TD) annotation added, no guide

but +TD, DeepSplice [41] guidance with -TD, DeepSplice guidance with +TD. All used

StringTie2 v2.1.1 and omitted long-read input.

Structural accuracy

Predictions were compared to references with gffcompare v0.12.6 [42]. Sensitivity, precision and

F1 score were extracted at exon, intron and transcript level.

Completeness

BUSCO v5.7.1 [43] was run in transcriptome mode (-m tran) with the following lineages:

Embryophyta_odb10 (*A. thaliana*), Nematoda_odb10 (*C. elegans*) and Metazoa_odb10 (*S.

mediterranea*).

StringTie2 parameter optimization

The transcriptome assembly was evaluated using the SIRV-set 4 RNA spike-in controls

to assess its accuracy, with specific StringTie2 parameters (-c, the minimum reads per bp

coverage to consider for multi-exon transcript, and -f, the minimum isoform fraction) being

tested (Table S2). Five combinations of StringTie2 parameters were used: default settings (-c 1; -

f 0.01), a stricter isoform fraction filter alone (-f 0.02), two higher coverage thresholds (-c 2 and -

c 5), and a mixed setting (-c 1.5, -f 0.02).

Assembly evaluation and comparative analysis

182   Completeness was assessed with BUSCO [43] (v5.7.1) with a Metazoa benchmark of 954

183 conserved genes, while assembly quality metrics were obtained using rnaQUAST [44] (v2.3.0)

184 and Transrate [45] (v1.0.3). Benchmarking procedures were conducted by comparing the

185 performance of the pipeline against existing annotations, with metrics such as the number of

186 correctly identified transcripts, assembly scores, and completeness indicators being analyzed.

ncRNA identification

All ncRNA sequences were downloaded from RNAcentral [46] and supplemented with

457 tRNA sequences from an external publication [47]. The sequences were linearized and

deduplicated using seqkit rmdup. The initial dataset contained 27,390,440 sequences, which was

reduced to 27,317,511 after deduplication. A BLASTn[2] search was conducted to identify

homologous sequences within the *S. mediterranea* nuclear and mitochondrial genomes. The

following parameters were used: -query: 27,317,511 ncRNA sequences; -db: *S. mediterranea*

genome (nuclear + mtDNA), -evalue: 1e-20, -word_size: 15, -qcov_hsp_perc: 90, -dust: no. The

output was then processed to extract the original sequence length of hits for further filtering. A

custom Python script (Table S3) was used to classify ncRNA types based on query ID patterns.

Hits were retained if alignment coverage met predefined thresholds (95% for tRNA, 90% for

miRNA, 70% for lncRNA, 90% for rRNA, 80% for other ncRNA). Then another custom R script

(Table S3) was used for converting to gtf format, refining and merging overlapping regions to

ensure only non-redundant ncRNA annotations were retained.

Analyses

Selection and validation of transcriptome assembly approach

204   One of the key steps in genome annotation is the assembly of the transcriptome. In our

efforts to enhance genome characterization for the asexual strain of *S. mediterranea*, we

employed StringTie2 [4], a widely recognized tool for transcriptome assembly. StringTie2 is

versatile, supporting both short-read and long-read sequencing datasets, which makes it well-

suited for comprehensive transcriptome analysis. However, StringTie2 has multiple versions,

each offering distinct features and performance enhancements. Therefore, before integrating

StringTie2 into our pipeline, we thoroughly optimized its parameters to ensure the highest

accuracy and reliability in transcript predictions. To this end, we employed two complementary

strategies.

213   In the first strategy, we generated an *in-silico* dataset to simulate genomic sequences with

corresponding full and incomplete annotations and short-read RNA-seq data (see Methods

section). This synthetic data allowed us to create scenarios with known simple transcript

structures, enabling a comprehensive evaluation of StringTie2 performance. Reads were aligned

to the *in-silico* reference genome using STAR. We then compared two commonly available

versions of StringTie2, 2.1.1 and 2.2.1, which allowed us to assess version-specific performance

improvements. We evaluated the results of the assembly step, by analyzing transcript

identification rates for both StringTie2 versions at base, exon, locus and transcript levels using

gffcompare. Our results indicated that StringTie2 version 2.1.1 provided a higher identification

rate of short-read data in reference-based mode than its descendant (Figure 1A), while there were

no differences in *de novo* mode. Although the StringTie2 documentation does not explicitly state

that version 2.1.1 is more sensitive for short-read data, the combination of release note details

and our experimental observation suggested that less stringent filtering of version 2.1.1 results in

a higher count of assembled isoforms. Meanwhile, version 2.2.1 has been optimized for hybrid-

read assembly (with the --mix option) and incorporates more conservative measures to reduce

false positives—which can lower the total transcript count when only short-read data are used.

This trade-off between sensitivity and precision is common in transcript assemblers. When

maximizing the number of detected isoforms from short reads is a priority, 2.1.1 may be

advantageous, whereas 2.2.1's improvements (including its ability to process mixed-read data)

make it preferable for datasets that include long reads. It should be noted that these version

choices are based on their prevalence at the time of our study, and although our benchmarking

indicates reliable performance, further testing in diverse datasets may be beneficial.

In the second strategy, we used Spike-In RNA Variants (SIRVs), a commercially

available collection of RNA spike-in controls, which were added to the RNA isolated from

planarians (see Table S1) prior to the preparation of sequencing libraries. These SIRVs included

three distinct modules — ERCC Mix 1 (a set of 92 synthetic RNA molecules ranging between

250 and 2,000 nt in length to resemble natural eukaryotic transcripts), Isoform Mix E0 (referred

here as multi-exon SIRVs), and long SIRVs (referred here as mono-exon SIRVs) — each

designed to mimic specific aspects of transcriptome complexity, including transcript abundance,

isoform diversity, and transcript length. RNA-seq reads were aligned to the reference SIRVs

sequences using STAR for short reads and Minimap2 for long reads. We then employed

StringTie2 using the "insufficient annotation" as the reference provided by the manufacturer of

SIRVs (SIRV_I), where some SIRVs that are actually present in the mixes are not annotated, and

compared the prediction to the correct annotation of all SIRVs (SIRV_C). Such a comparative

analysis demonstrated a proper reconstruction of missing spike-in transcripts. Then, focusing on

the mixture of synthetic RNA isoforms Iso Mix E0 category, we systematically assessed the

performances of seven assembly approaches for transcript variant detection, implementing

different combinations of read types—short and mixed—and assembly modes—reference-based

and *de novo*. These approaches included short reads in reference-based mode (SR_RB), short

reads in *de novo* mode (SR_DN), mixed reads in reference-based mode (MR_RB), mixed reads

in *de novo* mode (MR_DN), merged reads in *de novo* mode (MeR_DN), merged reads in

reference-based mode (MeR_RB), and merged reads in both modes (MeR_RB_DN). Mixed

reads refer to the combined use of short and long reads within a single assembly process,

whereas merged reads indicate separate assemblies of short reads and mixed reads, followed by

merging the resulting assemblies. For each approach, StringTie2 performance (Table S2) was

quantified in terms of accuracy metrics, reflecting the ability to correctly identify transcript

isoforms (Figure 1B). SR_DN approach achieved the lowest accuracy, ranging from 35.1% to

45.1% with its maximum at -c 1.5 and -f 0.02. Surprisingly, generally, the highest accuracy rate

of 81.1% was obtained when predictions from both short and mixed reads in both modes were

merged (MeR_RB_DN). This outcome underscores the synergistic benefits of integrating

multiple read types and assembly strategies.

Consequently, our pipeline employs a dual-version strategy—by default, StringTie2

version 2.1.1 is used for assembling short-read data and version 2.2.1 for mixed-read data,

ensuring optimal performance across a range of sequencing data types. Users should note that

while these versions provided robust performance in our tests, alternative versions may be

considered depending on specific experimental contexts and can be adjusted via command-line.

We set -c 1.5 and -f 0.02 based on pilot experiments with SIRVs, which balanced sensitivity and

precision in isoform detection (see Table S2). After completing individual assemblies, we

incorporated the MeR_RB_DN approach to merge reference-based and de novo transcripts,

resulting in a comprehensive and unified transcriptome annotation that captures a diverse array

of transcript isoforms.

Identification of artifactual transcripts

In our tests with *in-silico* data, we identified several types of artifactual transcripts,

including those with unusually long exons exceeding 100,000 nt, reversed duplicates (transcripts

overlapping the same genomic regions as the reference ones but on the opposite strand), as well

as fragmented and chimeric RNA. Fragmented transcripts are individual RNAs that have been

split into multiple predicted gene models due to incomplete transcript reconstruction or assembly

errors. Chimeric transcripts, on the other hand, erroneously combine sequences from multiple

distinct genes. In eukaryotic genomes, exons generally range in size from approximately 50 to

5,000 nt, with the majority falling between 100 and 1,000 nt [48]. Instances of exons exceeding

10,000 nt are exceptionally rare and are usually associated with specific organismal features,

such as immune-related genes or tandem duplications. Even in these rare cases, such lengths

remain an order of magnitude below 100,000 nt [49]. While highly effective in many contexts,

StringTie2 may incorrectly model transcripts, especially in regions with overlapping genes or

incomplete stop codons. Such inaccuracies can extend exon boundaries far beyond biologically

realistic lengths [50].

To address the identified challenges and eliminate artifacts, we added a custom function

to the pipeline that ensured that transcripts with exons exceeding 10,000 nt and/or transcripts that

surpassed 100,000 nt of the genomic region were removed, thus refining the annotation and

preventing the propagation of erroneous data. Additionally, we included a function to identify

and filter out reversed duplicates without inadvertently removing legitimate transcripts. We also

implemented criteria to flag potential fragmented and chimeric RNA based on exon length,

genomic proximity, and functional annotations. In our pipeline, indicators for fragmentation

include multiple predicted genes in close genomic proximity (within 1,000 nt) on the same strand

and adjacent or overlapping genomic coordinates. The exon length and genomic proximity

thresholds are based on general eukaryotic genome characteristics [49]. These parameters should

be adjusted as needed based on the specific organism or dataset characteristics. Transcripts and

genes are flagged as chimeric if they meet at least one of the following criteria: (i) they display

structural inconsistencies, such as widely spaced or non-contiguous exons; (ii) they span multiple

distinct genomic regions; or, in the case of genes, (iii) they possess multiple, unrelated functional

annotations. As part of our automated pipeline, we only added information indicating whether a

gene or transcript is possibly fragmented or chimeric. The decision to remove corresponding

genes or transcripts should be made by the user based on specialized evaluation. Incorporating

functional annotations into the pipeline as a part of manual curation can significantly improve the

accuracy of detecting artifacts.

Pipeline overview and workflow

Having resolved the challenges outlined above, we developed a versatile genome

annotation pipeline and demonstrated its performance by applying it to the asexual strain of *S.*

mediterranea. This pipeline integrates multiple sequencing technologies and bioinformatics tools

to achieve comprehensive and accurate gene and transcript annotations across different

organisms. The pipeline is divided into two main parts, each structured into distinct sequential

steps.

Part I: Pre-processing and assembly (SmedAnno)

The first part of the pipeline is container-orchestrated, fully automated, and encompasses

pre-processing, alignment, assembly, and functional annotation. The input data for the pipeline

can be either short-read or mixed, the latter including both short and long reads. While even a

single sample can be processed when only short reads are available, incorporating multiple

samples is recommended to achieve more reliable and comprehensive results. Mixed-read

datasets require at least one sample with short reads and one with long reads. Additionally, the

automated pipeline is optimized for paired-end reads in short-read mode. Mapping and transcript

assembly are performed independently for each sample. The workflow begins by pre-processing

the RNA-seq data, performing adapter and quality trimming with TrimGalore and Filtlong for

short- and long reads, respectively, followed by optional length and quality filtering. Next, the

pipeline focuses in particular on the removal of reads corresponding to ribosomal RNA (rRNA),

based on their alignment to user-provided rRNA reference sequences (Figure 2). This step was

found to be very important in the case of organisms with unusually high A/T content, such as *S.*

mediterranea. Library preparation methods typically involve rRNA depletion or polyA

enrichment to facilitate the study of protein-coding gene expression. However, commercially

available rRNA depletion kits are mostly optimized for model species, and polyA enrichment is

not effective in organisms with a high A/T content. Consequently, reads corresponding to rRNA

can be abundant in sequencing data for non-model organisms. In scenarios where rRNA

reference sequences are unavailable (not provided by the user), the pipeline bypasses the rRNA

removal step, continuing with the whole RNA-seq dataset. The pipeline then aligns the processed

RNA-seq reads to the reference genome using STAR for short reads and Minimap2 for long

reads. After completing this step, splice junctions can be optionally identified using a deep-

learning-based predictor DeepSplice [41]. It uses convolutional neural network architectures

(OpenSpliceAI-like) and supports CUDA acceleration for efficient computation. DeepSplice

infers splicing donor and acceptor motifs directly from the reference genome, based on an

integrated model. Currently the user can select a model, according to phylogenetic proximity to

the target organism, from five available ones (human, mouse, zebrafish, honeybee, and

thalecress). Next, the transcript assembly phase uses StringTie2 – by default, version 2.1.1 is

applied to short-read data, while version 2.2.1 is used for mixed-read data (which integrates both

short and long reads). Users can specify alternative versions if desired. After assembly, the

pipeline merges transcripts across all samples using StringTie2 to create unified sets of

transcripts. This merging process is performed separately for reference-based (if reference

annotation was provided) and *de novo* assemblies before combining them into the final complete

annotation. This dual approach by performing both reference-based and *de novo* annotation,

provides comprehensive assembly, thus capturing a wide range of transcript isoforms.

DeepSplice evidence is then merged with StringTie2 predictions. Subsequently, transcripts with

excessively long exons or that cover unusually large genomic regions are filtered out. Next, the

pipeline uses gffcompare to compare the predicted annotations against the reference annotations

(if provided), assessing the completeness and accuracy of the assembled transcripts. This

comparison generates a report that includes data summaries and accuracy estimations.

Additionally, it enriches the predicted annotations with detailed information about their overlaps

with known reference transcripts and genes. However, the generated annotations may contain

issues that can compromise the quality and usability of the annotation files. To address these

issues, the pipeline employs AGAT (Another GTF/GFF Tool), which systematically identifies

and corrects common syntax errors, ensuring that essential attributes such as gene_id and

transcript_id are present for each feature, preventing incomplete annotations, correcting improper

use of delimiters in attribute fields (such as replacing spaces with a semicolon), detecting and

removing duplicate entries, and ensuring correct key-value pair syntax. This correction phase is

crucial for maintaining high-quality genome annotation. The obtained set of transcripts is then

subjected to functional annotation in a process that involves the sequential application of several

tools, and which includes homology analysis and the identification of protein domains. To this

end, first, the open reading frames (ORFs) are predicted using TransDecoder. Following that,

BLASTp and BLASTx are used for homology search through the SwissProt database. Finally,

protein domains are identified using hmmscan through the PFAM database and InterProScan,

which aggregates multiple domain resources (SMART, Panther, TIGRFAM, etc.) and enriches

predictions with Gene Ontology (GO). This multi-faceted approach filters out low-confidence

annotations, retaining only the transcripts supported by multiple lines of evidence. We used the

SwissProt, PFAM and InterProScan databases as a part of the automated pipeline specifically for

their low to medium memory requirements. However, we recommend to use additional

resources, for example eggNOG, NCBI, UniRef, KEGG, RefSeq, COG, Reactome, and BioCyc

for conducting in-depth homology analyses, as these databases provide complementing

information, evolutionary relationships, and diverse protein characteristics. Next, the functional

annotations are integrated into the GTF file. This integration applies default thresholds: for

PFAM predictions, a bit score greater than 20, a conditional e-value below 1e-5, and coverage

exceeding 50%; for SwissProt predictions, a BLAST e-value threshold of 1e-5 and an identity

threshold of 25%; InterPro hits are retained if they meet the e-value threshold of < 1e-5 in at least

one of the member databases. These criteria ensure that low-confidence functional annotations

are discarded. However, all filtering thresholds can be configured by the user. Then, the pipeline

adds new columns to the GTF annotation file, including has_ORF (TRUE/FALSE),

best_SwissProt_blastp_hit, best_SwissProt_blastx_hit, best_Pfam_hit, and best_InterPro_hit

domain information. The next step is the identification of overlapping genes and transcripts,

which detects genes that share genomic regions and checks if each transcript in a gene has at

least one overlapping exonic region with any other transcript in the same gene. Lastly, the

pipeline identifies the aforementioned reversed duplicates and addresses fragmented and

chimeric genes and transcripts. At the completion of functional annotation, the pipeline emits an

interactive summary-and-advice block. Six boolean quality control flags

(*overlapped_predicted_gene*, *is_potentially_fragmented_reference,*

is_potentially_chimeric_reference, *is_potentially_fragmented_de_novo,*

is_potentially_chimeric_structure_de_novo, and is_potentially_chimeric_function_de_novo) are

printed together with plain-language recommendations (e.g. "merge fragmented loci", "split

structural chimeras"). Users are expected to review each flagged locus individually in IGV,

confirm read support, and decide case-by-case whether to merge, split, or retain the model

unchanged. All these steps (Figure 2) are implemented in the SmedAnno tool, forming Part I of

the entire pipeline. The source code can be accessed on GitHub

(https://github.com/Norreanea/SmedAnno).

Part II: Manual curation and refinement

The second part of the pipeline involves user-driven inspection and refinement of the

annotated data. Despite the implementation of several automated steps aimed at optimizing

transcript assembly and obtaining biologically relevant transcripts that encode functional protein

domains, resolving overlapping or fragmented transcripts remains a significant challenge that

requires specialized approaches. While the primary focus is on transcript assembly, gene features

are subsequently derived from the assembled transcripts. Fragmented and chimeric transcripts,

which primarily affect RNA-level annotations, are addressed by prioritizing the longest isoform

to define each gene, thereby reducing redundancy and improving annotation accuracy. To

accurately determine whether genes should be merged after being marked as potentially

fragmented and/or overlapped, promoter and terminator sequences should be analyzed using

Promoter 2.0 and ARNold tools (Figure 3). This analysis verifies the coherence of regulatory

regions, ensuring that merged gene models maintain consistent promoter and terminator

elements. These tools are accessible via their respective web pages and are not included in the

automated pipeline. Users must extract sequences between inspected genes to check for the

presence of promoters and terminators. After potential merges, manual adjustments are advised

to finalize gene models, ensuring accurate gene length prediction and overall annotation

reliability.

While genome annotation typically focuses on protein-coding genes, our pipeline also

includes a step for annotating non-coding RNAs (ncRNAs), which have significant functional

relevance. This step is optional, as it requires downloading the extensive (>20 GB) RNAcentral

database. The ncRNA sequences are first linearized and deduplicated using seqkit rmdup to

remove redundant entries. Homologous regions are then identified in the reference genome using

BLASTn. After the identification step, ncRNA types are classified based on the sequence

identifier. The matching results are then converted to GTF format. A multi-step filtering and

refinement process is used to improve the accuracy of the annotations. This process addresses

several common issues, including multiple alignments at identical genomic positions,

inconsistent naming conventions, and clustering.

The final step integrates all gathered information into a unified genome annotation file,

ready for downstream applications, including both coding and non-coding elements with

comprehensive functional insights.

The bioinformatics tools used in each pipeline step, along with their respective versions are

summarized in Table 1.

Table 1. Bioinformatics tools and pipeline steps

Step	Tool	Version
Step 0: Trim adapters & filter reads	TrimGalore Filtlong	0.7.1 0.2.1
Step 1: Remove rRNA	STAR Minimap2 samtools	2.7.9a
Step 2: Align reads to the reference genome	STAR Minimap2	2.7.9a
Step 3: Assemble genes & transcripts	StringTie2 DeepSplice	2.1.1 (short reads) 2.2.1 (mixed reads) 0.10
Step 4: Merge reference-based assemblies	StringTie2	2.2.1
Step 5: Merge *de novo* assemblies & create pre-final annotation	StringTie2	2.2.1

Step 6: Filter out transcripts artifacts	bash	
Step 7: Compare isoforms	gffcompare	0.12.6
Step 8: Correct & enhance annotation	AGAT	1.4.1
Step 9: Determine functional attributes	TransDecoder blastp blastx hmmscan InterProScan	5.7.1 2.15.0+ 2.15.0+ 3.3.2 5.67
Step 10: Integrate functional attributes	R	4.4.1
Step 11: Identify overlapping genes & transcripts	R	4.4.1
Step 12: Identify reversed duplicates	R	4.4.1
Step 13: Identify fragmented & chimeric genes	R	4.4.1
Step 14: Analyse promoter & terminator sequences	Promoter 2.0 ARNold	No official version number No official version number
Step 15: Annotate non-coding RNA (ncRNA)	blastn seqkit	2.15.1+ 2.8.2
Step 16: Integrate collected information into a unified genome annotation file	R	4.4.1
Step 17: Incorporate gene ontology (GO)	R	4.4.1
Step 18: Perform final review and quality assurance	R	4.4.1

SmedAnno benchmarking

To benchmark the automated Part I of our pipeline (SmedAnno), we compared its

performance against BRAKER3, a widely used genome annotation tool that integrates

GeneMark-ET and AUGUSTUS gene prediction, guided by short-read RNA-seq data [39]. The

comparison was carried out using three genomes: two classic models – *A. thaliana* (AT) and *C.*

elegans (CE) – and *S. mediterranea* (SM). Since BRAKER3 utilizes *ab initio* approach and

short-read data only, we ran a scaled-down version of SmedAnno without reference-based

guidance, exclusively on Illumina reads, to ensure a valid comparison. For *A. thaliana* and *C.*

elegans, publicly available data were used, while for *S. mediterranea* we employed in-house

generated datasets (Table S1). For DeepSplice, which is integrated into our pipeline, we

employed thalecress model for *A. thaliana*, and honebee model for *C. elegans* and *S.*

mediterranea. SmedAnno was run in four parameter configurations (with/without DeepSplice

and TransDecoder). After assembling transcripts for each genome, we evaluated the resulting

transcriptomes using BUSCO, which assesses the presence of conserved orthologs, by

comparing a transcriptome of interest to those within the Metazoa lineage, and gffcompare.

BRAKER3 reached slightly higher BUSCO completeness in AT (98.9 %) and CE (98.6

%) than the most comparable SmedAnno runs (~95 % and 88.8 %, respectively), but fell behind

in SM (76.9 % vs 78.6 % for both DeepSplice –TD and noGuide –TD) (Figure 4A). Notably,

SmedAnno yielded increased fraction of duplicated BUSCOs for all tested genomes (Figure 4A).

We found that SmedAnno configurations without TransDecoder generated more transcripts

being identified as single-copy BUSCOs. Next, the annotations obtained with BRAKER3 and

SmedAnno were compared with corresponding reference genome annotations using gffcompare.

The proportion of true positive predictions was assessed at transcript, exon and intron levels, and

depicted as F1 scores, where the higher value better reflects the reference. As expected,

BRAKER3 performed better in CE. BRAKER3's advantage in *C. elegans* is consistent with its

AUGUSTUS species-specific parameter set, which was trained on that very genome.

SmedAnno+DeepSplice (-TD) outperformed BRAKER3 in tests on AT and SM, displaying

higher F1 score at the exon and transcript levels in AT, and at all levels in SM (Figure 4B).

Genome annotation of the asexual strain of *S. mediterranea*

After demonstrating the efficiency of SmedAnno, we applied the whole pipeline to

generate a genome annotation of the asexual strain of *S. mediterranea*. To this end, we used the

available reference genome annotation and both Illumina and PacBio sequencing datasets (Table

S1). The SmedAnno output was subjected to manual curation and refinement, which included

identification of artifactual transcripts and detection of missing genes. The annotation,

schMed_Pn, is available in GEO repository under accession GSE294569.

Identifying fragmented and chimeric artifacts

In the context of genomic annotation, artifacts are erroneous gene or transcript models

that arise from limitations in the annotation process rather than representing true biological

entities. Specifically, we refer here to fragmented artifacts as those that occur when a single gene

is incorrectly split into multiple separate gene models. This fragmentation can result from

incomplete transcript assembly or sequencing errors, leading to discontinuous or truncated

representations of what is biologically a single gene. On the other hand, chimerism typically

arises from misassembly during transcriptome reconstruction, especially in regions with high

gene density or overlapping gene structures.

SmedAnno   employs   flags   such   as: *is_potentially_fragmented_de_novo*,

is_potentially_chimeric_structure_de_novo, and *is_potentially_chimeric_function_de_novo* to

identify these artifacts based on functional and structural criteria. Below, we discuss three

illustrative examples from schMed_Pn (depicted in Figure 5) that highlight the complexities and

implications of these annotation challenges. SMESGs here are reference genes and MSTRGs are

putative genes.

Prediction of gene fragments

In one genomic region, three genes—SMESG000004200.1, SMESG000004201.1, and

MSTRG.967—were located in close proximity on chromosome 1 (smed_chr1) (Figure 5A).

Specifically, gene SMESG000004200.1 spanned position 14,973,263 to 15,003,452 on the

positive strand, while SMESG000004201.1 extended from 15,003,453 to 15,011,292 on the

same strand. StringTie2, predicted an additional transcript, MSTRG.967, which encompassed the

regions of both genes (14,973,256 to 15,011,292). Both SMESG000004200.1 and

SMESG000004201.1 were independently annotated as "Epidermal growth factor receptors."

However, the prediction by StringTie2 that merged these two genes into a single transcript was

more biologically plausible. This is because the close genomic proximity and shared functional

annotations suggested a possible complex gene structure that is better represented by a single,

contiguous   transcript.   Flagging   these   genes   as   potentially   fragmented

(*is_potentially_fragmented_de_novo*) highlighted the necessity to consider merged annotations

in densely packed genomic regions to avoid misinterpretation of gene boundaries.

Chimeric transcript prediction due to excessive intron length

The   gene   SMESG000065897.1   initially   comprised   a   single   transcript,

SMEST065897001.1, spanning positions 325,241,809 to 325,242,781 on chromosome 1 (Figure

5B). However, StringTie2 predicted an additional transcript, MSTRG.16892.1, which extended

dramatically from 325,241,809 to 325,347,749, introducing an intron exceeding 100,000 nt. This

unusually   large   intron   length   is   atypical   and   suggested   a   misassembly,   classifying

MSTRG.16892.1 as potentially chimeric (*is_potentially_chimeric_structure_de_novo*).

The presence of such a large intron likely indicated that MSTRG.16892.1 erroneously

combined sequences from multiple genomic loci or failed to accurately represent the true

transcript structure of SMESG000065897.1. Consequently, the chimeric transcript should be

disregarded, and the gene localization should remain unchanged, basing solely on the original

transcript SMEST065897001.1.

Functionally chimeric gene due to divergent transcript annotations

The gene SMESG000077853.1 exhibited a complex transcriptional landscape (Figure

5C). While most transcripts, such as MSTRG.31576.1, MSTRG.31576.5, MSTRG.31576.6, and

all reference SMEST transcripts were annotated as "ankyrin repeat proteins," one transcript,

MSTRG.31576.11, was annotated as "Squamous cell carcinoma antigen recognized by T-cells 3

(SART-3)". This divergence in functional annotation across transcripts resulted in the gene being

flagged as chimeric (*is_potentially_chimeric_function_de_novo*). These inconsistent functional

annotations suggested that MSTRG.31576.11 may have been incorrectly assembled, possibly

incorporating sequences from a different gene or representing a mispredicted splice variant.

Given that "SART-3" functions are unrelated to "ankyrin repeat proteins," it is evident that

MSTRG.31576.11 does not belong to SMESG000077853.1. Consequently, MSTRG.31576.11

was filtered out.

In total, the automated run flagged 5.1% of all gene models, of which 1.2% were

confirmed as artifacts upon manual inspection.

Detecting missing genes through functional annotation and homology assignments

Functional annotations and homology assignments were carried out to explain the

biological function of the assembled transcripts and identify orthologous genes by using

established databases and computational tools. This approach facilitated the detection of missing

genes, thus highlighting potential gaps and guiding further refinement of the transcriptome

assembly.

To incorporate functional data into the newly developed *S. mediterranea* genome

annotation (schMed_Pn), we used the following databases: Pfam, SwissProt, and non-redundant

(nr) protein sequences from NCBI. Our annotation comprised 25,265 genes, of which 4,011 were

novel and 21,254 were identical to the reference genes. Each gene possessed at least one

function-related entry, either through a database hit or by being classified as having an ORF. For

18,552 genes, we successfully identified Gene Ontology (GO) terms using homology-based

strategies, which included the pfam2go package, UniProt GOA conversion, and a custom R

script (Table S3) to retrieve GO annotations based on NCBI IDs. Notably, 2,427 of these GO-

annotated genes were putative. To evaluate the similarity of GO terms between novel and

reference genes, Venn diagrams were constructed, highlighting shared and unique functional

categories across the gene sets. We assigned a total of 23,909 GO terms to genes within our

annotation. Specifically, 18,586 GO terms were assigned to SMESG (reference) genes, and

5,323 GO terms were assigned to MSTRG (putative) genes. Of these, 5,180 GO terms were

shared between both gene categories, while 143 GO terms were uniquely assigned to MSTRG

genes (Figure 6A). To visualize the functional distribution of the gene sets, we generated word

clouds representing the most frequent unique GO terms for each group: MSTRG (Figure 6D) and

SMESG (Figure 6B), as well as those that were common for two groups (Figure 6C).

Insights into planarian ncRNA diversity

Beyond refining the annotations of protein-coding genes, we also performed a

preliminary assessment of non-coding RNAs (ncRNAs) in *S. mediterranea*. Although not an

exhaustive characterization, our analysis revealed a diverse repertoire of regulatory RNAs,

highlighting significant differences between publicly available annotations and newly generated

data. Publicly available datasets from RNAcentral assigned a total of 154 rRNAs, 578 miRNAs

(including 269 precursors), 61 tRNAs, 2 snoRNAs, 12 snRNAs, and 2 SRP RNA hits to *S.*

mediterranea. After deduplication, these numbers were refined to 129 rRNA, 571 miRNAs (with

156 precursors), 60 tRNAs, 2 snoRNAs, 12 snRNAs, and 2 SRP RNA hits. Additionally,

according to recent literature [47], *Schmidtea* genome contains 457 tRNA hits, which were

reduced to 434 after deduplication. In contrast, using our annotation pipeline, we identified 154

rRNA hits across 2,287 genomic positions, 179 miRNA hits across 1,042 genomic positions,

488 tRNA hits distributed over 4,298 genomic positions, 2 snoRNA hits across 69 genomic

positions, 18 snRNA hits across 141 genomic positions, 6 miscellaneous RNAs spread across

4,752 genomic positions, 18 RNase P ncRNA components across 408 genomic positions, 21

lncRNA hits across 185 genomic positions and 2 sRNA hits across 187 genomic positions. The

discrepancy between published records assigned to *S. mediterranea* and our annotated RNAs

highlights the ongoing need for careful re-annotation and experimental validation in

underexplored organisms like planarians. In addition, we uncovered 40 copies of regions

encoding signal recognition particle RNAs (SRP_RNAs), localized on chromosome 3 and

flanked by the genes SMESG000020110.1 (BLAST-predicted as a lysine-specific demethylase)

and MSTRG.33721 (BLAST-predicted as containing a thrombospondin type 3 repeat).

Moreover, we identified 10 copies of TERC_RNA located on smed_chr4, while *TERT* gene

(SMESG000061912.1) was located on smed_chr3.

Comparative analysis of selected *S. mediterranea* genome annotations

To ensure the reliability of schMed_Pn genome annotation, we compared it with

commonly used genome annotations of *S. mediterranea*. First, we analyzed transcript length

distributions. It revealed that all annotations predominantly consisted of relatively short

transcripts, as reflected by a sharp decline in cumulative transcript counts with increasing

transcript length (Figure 7A). Specifically, dd_Smed_v6 showed the steepest decline in

cumulative transcript counts with increasing length, suggesting a higher fragmentation rate;

however, caution is advised in interpreting these distributions, as there is no "standard" shape for

planarian transcript length. In contrast, schMed_Pn demonstrated a more gradual decline, which

indicated a broader distribution of transcript lengths. Assemblies such as schMedS1, schMedS2,

schMedS3_1, and schMedS3_2 exhibited moderate slopes with slight differences, indicating

varying transcript length distributions among them. The SmedAsxl genome_v1.1 displayed a

narrower range of longer transcripts. In contrast, SmedGD_c1 and SMESG.1, similar to

schMed_Pn, demonstrated relatively smooth distributions, signifying balanced transcript

coverage with a significant proportion of longer transcripts.

The completeness of the transcriptome assemblies was evaluated using BUSCO (Figure

7B). The dd_Smed_v6 and SMESG.1 annotations achieved the highest single-copy ortholog

recovery—783 and 756 complete BUSCOs out of 954, respectively—with low duplication rates

(107 and 196 duplicated BUSCOs), indicative of very accurate core gene models but at the cost

of fewer total transcript isoforms. In contrast, schMed_Pn attained a balanced profile of

completeness and isoform richness, with 767 complete BUSCOs (426 single-copy, 341

duplicated), coupled with minimal fragmentation (34) and moderate missing counts (153). This

reflected our pipeline's emphasis on maximizing isoform discovery without substantially

sacrificing core□gene completeness. Other assemblies (schMedS1–3, SmedAsxl genome_v1.1)

showed lower complete counts and higher fragmentation or missing rates, consistent with greater

redundancy or gaps. Based on BUSCO analysis, both SMESG.1 and dd_Smed_v6 demonstrated

robust performance (albeit with different trade-offs in duplication and missing orthologs).

Notably, schMed_Pn also ranked among the stronger annotations, demonstrating moderate

duplication and missing ortholog levels, while retaining high completeness.

In addition, we employed rnaQUAST and Transrate to generate detailed assembly quality

metrics (Table 2). Notably, schMed_Pn exhibited a high number of transcripts over 1,000 nt and

maintained a competitive N50 value (the length of the shortest contig or scaffold covering 50%

of the total assembly), reflecting effective transcript coverage and solid assembly quality.

Table 2. Comparative metrics for selected transcriptome assemblies of *S. mediterranea*

Metric	dd_Smed_v6	schMed_Pn	schMedS1	schMedS2	schMedS3_1	schMedS3_2	SmedAsxl_v1.1	SmedGD_c1.3	SMESG.1
No. of Transcripts	41,745	41,396	22,348	27,790	41,358	41,036	22,855	29,850	31,102
Min transcript length (nt)	201	200	267	267	145	131	75	87	267
Max transcript length (nt)	19,524	48,466	86,373	1,022,601	76,733	94,201	43,226	12,11	48,121

Total Bases (M)	53.23	97.09	40.12	57.93	72.60	71.81	25.55	23.15	65.55
Mean transcript length (nt)	1,275	2,345	1,795	2,084	1,755	1,750	1,114	770	2,108
No. of transcripts >1k	18,487	31,199	16,082	20,727	29,402	28,932	8,610	7,218	23,648
No. of transcripts >10k	94	702	67	293	116	119	47	2	346
No. of transcripts with ORF	20,461	32,798	18,694	23,854	31,503	31,147	12,937	17,567	26,665
N50 (nt)	1,931	3,070	2,159	2,535	2,133	2,141	1,653	1,011	2,570
GC (%)	0.34	0.33	0.33	0.33	0.33	0.33	0.34	0.36	0.33

Performance benchmarking enabled the identification of improvements and areas needing

enhancement, while also positioning schMed_Pn within the context of current resources. These

analyses collectively demonstrated that our approach offers a balanced and complete annotation,

effectively capturing a wide range of transcript lengths and maintaining high completeness.

Discussion

Numerous pipelines and tools have been developed to tackle genome annotation across

diverse organisms. For well-studied species—such as *Drosophila melanogaster* or *Arabidopsis*

thaliana—community-maintained platforms (FlyBase [51] and TAIR [52,53] respectively)

provide robust resources that streamline annotation. In prokaryotes, automated tools like Prokka

[10] efficiently handle smaller, less complex genomes, while eukaryotic model organisms often

benefit from RefSeq [7] or large consortia efforts, which incorporate extensive experimental

validation. However, non-model eukaryotes, including many invertebrates and plants, typically

face an incomplete reference, novel repetitive elements, and limited training data for *ab initio*

predictions. Pipelines such as MAKER [5,54], BRAKER [55,56], GGA [9], nf-core

GenomeAnnotator [12], and others can automate workflow portions—especially when reference

data or orthologous gene sets are available. However, in certain species with high genomic

complexity or less functional annotation data, these tools may still underperform or require

extensive manual curation.

In this study, we developed and validated a comprehensive genome annotation

pipeline designed to address the challenges of working with non-classical model organisms,

while generating high-quality annotations applicable to broader genomic research. Our pipeline

uniquely combines: (i) the integration of multiple data types, (ii) automated artifact detection,

(iii) dual approach assembly —reference-based plus *de novo*, and (iv) broad functional

integration.

The BUSCO completeness of SmedAnno (the automated part of the pipeline) was within

4% of BRAKER3 in *A. thaliana*, surpassed it by approximately 2% in *S. mediterranea*, and

lagged behind by approximately 10% in *C. elegans*. Notably, SmedAnno runs consistently

doubled the proportion of duplicated BUSCO genes. Because BUSCO counts every alternatively

spliced isoform as a duplicate, such inflation is widely regarded as a proxy for improved isoform

recovery rather than assembly noise [57]. In practical terms, this means that the apparent

redundancy in SmedAnno is biologically informative when transcript diversity, rather than the

most parsimonious gene count, is the primary research goal.

Leveraging this enhanced isoform recovery, our analysis identified 4,011 novel genes and

multiple previously uncharacterized isoforms and ncRNAs in the asexual strain, enriching the

genomic landscape of *S. mediterranea*. Notably, the GO term analysis of putative MSTRG

genes revealed enrichment in diverse biological functions. The occurrence of these genes

necessitates further investigation to better understand their implications and potential

significance. While our transcriptomic data offer strain-specific insights, it should be considered

that the annotation relies on the genome of the sexual strain as the reference. Consequently,

differences between strains, such as chromosomal translocation and other genetic variations, may

compromise the accuracy of gene prediction and transcript assembly, leading to potential

inconsistencies in annotation quality.

The existing, planarian-specific repositories—such as PlanMine [21], PLANAtools [58],

WormBase [59], PlanoSphere [60], PlanNET [61], and various planaria single-cell datasets [62–

65]—provide essential resources for gene model comparisons and functional information in *S.*

mediterranea. Despite their value, discrepancies in naming conventions, annotation updates, and

differences between sexual and asexual strains can lead to inconsistencies. These challenges

highlight the importance of robust, well-documented pipelines that can reconcile or supplement

these external databases, thus ensuring more accurate gene models.

A detailed inspection of the curated annotation uncovered two intriguing features of the

genomic architecture: genes located in close proximity and non-contiguous transcripts. Although

these observations could potentially represent annotation artifacts, it is crucial to carefully assess

them in the context of established genomic knowledge. Genes located in close proximity can

reflect various biological scenarios. One possibility is the presence of functionally independent

genes that happen to be closely spaced, as observed in other genomes where gene density is high

[66,67]. For example, two planarian genes—SMESG000043996.1 and SMESG000043998.1—

that encode kynureninase and synaptotagmin XVII, respectively, are unrelated but positioned

close to each other. Alternatively, such arrangements may arise from evolutionary events such as

gene duplication followed by functional divergence, where both copies remain in close physical

proximity. Another scenario includes operon-like gene structures commonly seen in prokaryotes

but also identified in some eukaryotic contexts [68]. These structures allow the co-regulation of

adjacent genes through shared promoters or other regulatory elements. Bidirectional promoters,

enabling two genes to be transcribed in opposite directions from the same regulatory region,

represent another mechanism driving close gene arrangements [69]. On the other hand non-

contiguous transcripts, where exons are unusually spaced or appear fragmented, often arise from

structural variations, alternative splicing, or sequencing artifacts. Biologically, large introns or

dispersed exonic regions can be linked to long-range regulatory elements or structural genome

features such as transposon insertions. Moreover, such patterns may indicate complex gene

models, as seen in genes encoding for proteins with repetitive domains [70,71] or in cases where

alternative promoters and polyadenylation sites lead to diverse transcript forms   [72,73].

Consequently, accurately capturing these intricate gene architectures often requires more

comprehensive strategies than those employed for simpler genomes, underscoring the need for

user scrutiny and flexibility in annotation pipelines. The primary advantage of our pipeline lies in

its capacity to address these complexities. By combining reference-based and *de novo* methods

and incorporating both coding and non-coding RNA annotations it addresses the challenges

posed by scarcely studied genomes. However, the pipeline's reliance on user-defined thresholds

and manual curation introduces subjectivity, potentially impacting reproducibility.

While the automated steps of SmedAnno effectively identify fragmented and chimeric

transcripts, resolving these artifacts requires expertise and thus is not fully automated. Another

limitation concerns functional annotations, which can lead to false positives in identifying

chimeric genes. For instance, the same protein from different species or proteins representing

different subunits of a complex, may be annotated as distinct entries in functional databases. This

can lead to genes being incorrectly flagged as chimeric if the functional entries do not exactly

match. Accordingly, development of refined algorithms that account for near-identical

annotations and minor terminological differences is vital.

Future enhancements could include the integration of machine learning algorithms to

automate artifact resolution and improve annotation consistency. The present container

infrastructure already simplifies the implementation of graphics processing units (GPUs) and

central processing units (CPUs). Expanding the pipeline to support multi-omics data integration,

such as proteomics and epigenomics, would further enhance its utility. Community-driven efforts

to standardize annotation protocols and databases would ensure broader applicability and

reproducibility. Finally, experimental validation of novel genes and isoforms identified through

this pipeline will be critical to confirm their biological relevance.

Compared to pipelines that excel in either prokaryotic or model vertebrate annotation –

where reference data is abundant and well-curated – our approach fills the gap for non-classical

model systems with more limited resources. In doing so, it advances genome annotation efforts

for *S. mediterranea* and provides a robust framework for addressing similar challenges in other

species, fostering broader genomic discoveries.

Availability of Source Code and Requirements

Project name: SmedAnno

Project homepage: https://github.com/Norreanea/SmedAnno

Operating system: Linux

Container image: Docker ≥20.10, docker-compose ≥2.5

Programming languages: shell, R, Python, awk

License: MIT License

Data Availability

The *in-silico* datasets supporting the results and all other code used in this article are available in

the GitHub repository archived in Zenodo https://doi.org/10.5281/zenodo.15130394. RNA-seq

data can be accessed from the following GEO repositories: GSE293999 and GSE293997 from

NCBI BioProject PRJNA1243063; and GSE294569 from NCBI BioProject PRJNA1246370.

The final genome annotation file is available in GSE294569.

List of Abbreviations

BP: base pairs; NT: nucleotide; RNA: ribonucleic acid; rRNA: ribosomal RNA; ncRNA: non-

coding RNA; SIRV: Spike-In RNA Variants; ORF: open reading frame; GTF: Gene Transfer

Format; BLAST: Basic Local Alignment Search Tool; STAR: Spliced Transcripts Alignment to

a Reference; BUSCO: Benchmarking Universal Single-Copy Orthologs; GO: Gene Ontology;

cDNA: complementary DNA; DNA: deoxyribonucleic acid; RIN: RNA Integrity Number; GGA:

Galaxy Genome Annotation.

Competing Interests

Nothing to declare

Declaration of generative AI and AI-assisted technologies in the writing process

During the preparation of this work, the authors used the free version of OpenAI-ChatGPT

c2022 (v.GPT-o3-mini, February 2025) in order to improve manuscript readability and language.

After using this tool, the authors thoroughly reviewed and modified the manuscript and took full

responsibility for the content of the publication. The literature survey was done only by the

authors, with no input from the AI tools.

Funding

This work was financed by the National Science Centre, Poland, through a grant no.

2019/35/B/NZ2/02658 to PJ.

Authors' Contributions

AZ: Conceptualization, Methodology, Software, Visualization, Writing – Original Draft, Writing

– Review & Editing; MMZ: Conceptualization, Methodology, Writing – Review & Editing; AD:

Investigation, Writing – Review & Editing; ASCz: Investigation, Writing – Review & Editing;

PJ: Conceptualization, Supervision, Project administration, Funding Acquisition, Writing –

Review & Editing.

References

1. Stanke M, Keller O, Gunduz I, Hayes A, Waack S, Morgenstern B. AUGUSTUS: A b initio

prediction of alternative transcripts. *Nucleic Acids Res*. 2006; doi: 10.1093/nar/gkl200.

2. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al.. BLAST+:

Architecture and applications. *BMC Bioinformatics*. 2009; doi: 10.1186/1471-2105-10-421.

3. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al.. STAR: Ultrafast

universal RNA-seq aligner. *Bioinformatics*. 2013; doi: 10.1093/bioinformatics/bts635.

4. Shumate A, Wong B, Pertea G, Pertea M. Improved transcriptome assembly using a hybrid of

long and short reads with StringTie. *PLoS Comput Biol*. Public Library of Science; 2022; doi:

10.1371/journal.pcbi.1009730.

5. Cantarel BL, Korf I, Robb SMC, Parra G, Ross E, Moore B, et al.. MAKER: An easy-to-use

annotation pipeline designed for emerging model organism genomes. *Genome Res*. 2008; doi:

10.1101/gr.6743907.

6. Haas BJ, Salzberg SL, Zhu W, Pertea M, Allen JE, Orvis J, et al.. Automated eukaryotic gene

structure annotation using EVidenceModeler and the Program to Assemble Spliced Alignments.

Genome Biol. 2008; doi: 10.1186/gb-2008-9-1-r7.

7. O'Leary NA, Wright MW, Brister JR, Ciufo S, Haddad D, McVeigh R, et al.. Reference

sequence (RefSeq) database at NCBI: Current status, taxonomic expansion, and functional

annotation. *Nucleic Acids Res*. Oxford University Press; 2016; doi: 10.1093/nar/gkv1189.

8. Kurylo C, Guyomar C, Foissac S, Djebali S. TAGADA: a scalable pipeline to improve

genome annotations with RNA-seq data. *NAR Genom Bioinform*. Oxford University Press; 2023;

doi: 10.1093/nargab/lqad089.

9. Hiltemann S, Rasche H, Gladman S, Hotz HR, Larivière D, Blankenberg D, et al.. Galaxy

Training: A powerful framework for teaching! *PLoS Comput Biol*. Public Library of Science;

2023; doi: 10.1371/journal.pcbi.1010752.

10. Seemann T. Prokka: Rapid prokaryotic genome annotation. *Bioinformatics*. Oxford

University Press; 2014; doi: 10.1093/bioinformatics/btu153.

11. Swain MT, Tsai IJ, Assefa SA, Newbold C, Berriman M, Otto TD. A post-assembly genome-

improvement toolkit (PAGIT) to obtain annotated genomes from contigs. *Nat Protoc*. 2012; doi:

10.1038/nprot.2012.068.

12. Ewels PA, Peltzer A, Fillinger S, Patel H, Alneberg J, Wilm A, et al.. The nf-core framework

for community-curated bioinformatics pipelines. *Nat Biotechnol*. 2020; doi: 10.1038/s41587-

020-0439-x.

13. Wu S, Yuan S, Su Z. User-friendly genome assembly and gene annotation pipelines for

vertebrates.

14. Maia GA, Filho VB, Kawagoe EK, Teixeira Soratto TA, Moreira RS, Grisard EC, et al..

AnnotaPipeline: An integrated tool to annotate eukaryotic proteins using multi-omics data. *Front*

Genet. 2022; doi: 10.3389/fgene.2022.1020100.

15. Humann JL, Lee T, Ficklin S, Main D. Structural and Functional Annotation of Eukaryotic

Genomes with GenSAS.

16. Sánchez Alvarado A. The freshwater planarian Schmidtea mediterranea: Embryogenesis,

stem cells and regeneration. Curr Opin Genet Dev. Elsevier Ltd;

17. Baguñà J, Carranza S, Pala M, Ribera C, Giribet G, Arnedo MA, et al.. From morphology

and karyology to molecules. New methods for taxonomical identification of asexual populations

of freshwater planarians. A tribute to Professor Mario Benazzi. *Italian Journal of Zoology*.

Societa Paleontologica Italiana; 1999; doi: 10.1080/11250009909356258.

18. Robb SMC, Ross E, Alvarado AS. SmedGD: The Schmidtea mediterranea genome database.

Nucleic Acids Res. 2008; doi: 10.1093/nar/gkm684.

19. Robb SMC, Gotting K, Ross E, Sánchez Alvarado A. SmedGD 2.0: The Schmidtea

mediterranea genome database. *Genesis*. John Wiley and Sons Inc.; 2015; doi:

10.1002/dvg.22872.

20. Grohme MA, Schloissnig S, Rozanski A, Pippel M, Young GR, Winkler S, et al.. The

genome of Schmidtea mediterranea and the evolution of core cellular mechanisms. *Nature*.

Nature Publishing Group; 2018; doi: 10.1038/nature25473.

21. Rozanski A, Moon HK, Brandl H, Martín-Durán JM, Grohme MA, Hüttner K, et al..

PlanMine 3.0 - improvements to a mineable resource of flatworm biology and biodiversity.

Nucleic Acids Res. Oxford University Press; 2019; doi: 10.1093/nar/gky1070.

22. Guo L, Bloom JS, Dols-Serrate D, Boocock J, Ben-David E, Schubert OT, et al.. Island-

specific evolution of a sex-primed autosome in a sexual planarian. *Nature*. Nature Research;

2022; doi: 10.1038/s41586-022-04757-3.

23. Ivanković M, Brand JN, Pandolfini L, Brown T, Pippel M, Rozanski A, et al.. A comparative

analysis of planarian genomes reveals regulatory conservation in the face of rapid structural

divergence. *Nat Commun*. 2024; doi: 10.1038/s41467-024-52380-9.

24. Rabiasz A, Ziętkiewicz E. Schmidtea mediterranea as a Model Organism to Study the

Molecular Background of Human Motile Ciliopathies. Int J Mol Sci. Multidisciplinary Digital

Publishing Institute (MDPI);

25. Quinodoz S, Thomas MA, Dunkel J, Schötz EM. The More the Merrier?: Entropy and

Statistics of Asexual Reproduction in Freshwater Planarians. *J Stat Phys*. Springer New York

LLC; 2011; doi: 10.1007/s10955-011-0157-3.

26. Reddien PW. The Cellular and Molecular Basis for Planarian Regeneration. Cell. Cell Press;

27. Reddien PW, Sánchez Alvarado A. Fundamentals of planarian regeneration. Annu Rev Cell

Dev Biol.

28. Benham-Pyle BW, Mann FG, Brewster CE, Dewars ER, Vuu DM, Nowotarski SH, et al..

Planarians employ diverse and dynamic stem cell microenvironments to support whole-body

regeneration.

29. Sayers EW, Bolton EE, Brister JR, Canese K, Chan J, Comeau DC, et al.. Database resources

of the national center for biotechnology information. *Nucleic Acids Res*. Oxford University

Press; 2022; doi: 10.1093/nar/gkab1112.

30. Palmieri N, Schlötterer C. Mapping accuracy of short reads from massively parallel

sequencing and the implications for quantitative expression profiling. *PLoS One*. 2009; doi:

10.1371/journal.pone.0006323.

31. Sedlazeck FJ, Lee H, Darby CA, Schatz MC. Piercing the dark matter: bioinformatics of

long-range sequencing and mapping. *Nat Rev Genet*. 2018; doi: 10.1038/s41576-018-0003-4.

32. Chen Z, Ain NU, Zhao Q, Zhang X. From tradition to innovation: conventional and deep

learning frameworks in genome annotation. Brief Bioinform. Oxford University Press;

33. Newmark PA, Sánchez Alvarado A. Bromodeoxyuridine specifically labels the regenerative

stem cells of planarians. *Dev Biol*. Academic Press Inc.; 2000; doi: 10.1006/dbio.2000.9645.

34. Dainat J, Hereñú D, Dr. Murray KD, Davis E, Ugrin I, Crouch K, et al.. NBISweden/AGAT:

AGAT-v1.4.1 (v1.4.1). Zenodo;

35. Lamesch P, Berardini TZ, Li D, Swarbreck D, Wilks C, Sasidharan R, et al.. The Arabidopsis

Information Resource (TAIR): improved gene annotation and new tools. *Nucleic Acids Res*.

2012; doi: 10.1093/nar/gkr1090.

36. . Analysis of the genome sequence of the flowering plant Arabidopsis thaliana. *Nature*. 2000;

doi: 10.1038/35048692.

37. Howe KL, Bolt BJ, Shafie M, Kersey P, Berriman M. WormBase ParaSite – a

comprehensive resource for helminth genomics. *Mol Biochem Parasitol*. 2017; doi:

10.1016/j.molbiopara.2016.11.005.

38. . Genome Sequence of the Nematode *C. elegans*□: A Platform for Investigating Biology.

Science (1979). 1998; doi: 10.1126/science.282.5396.2012.

39. Gabriel L, Brůna T, Hoff KJ, Ebel M, Lomsadze A, Borodovsky M, et al.. BRAKER3: Fully

automated genome annotation using RNA-seq and protein evidence with GeneMark-ETP,

AUGUSTUS and TSEBRA.

40. Haas BJ, Papanicolaou A, Yassour M, Grabherr M, Blood PD, Bowden J, et al.. De novo

transcript sequence reconstruction from RNA-seq using the Trinity platform for reference

generation and analysis. *Nat Protoc*. 2013; doi: 10.1038/nprot.2013.084.

41. Abrar M, Hussain D, Khan IA, Ullah F, Haq MA, Aleisa MA, et al.. DeepSplice: a deep

learning approach for accurate prediction of alternative splicing events in the human genome.

Front Genet. 2024; doi: 10.3389/fgene.2024.1349546.

42. Pertea M, Pertea G. GFF Utilities: GffRead and GffCompare. *F1000Res*. F1000 Research

Ltd; 2020; doi: 10.12688/f1000research.23297.1.

43. Simão FA, Waterhouse RM, Ioannidis P, Kriventseva E V., Zdobnov EM. BUSCO:

Assessing genome assembly and annotation completeness with single-copy orthologs.

Bioinformatics. Oxford University Press; 2015; doi: 10.1093/bioinformatics/btv351.

44. Bushmanova E, Antipov D, Lapidus A, Suvorov V, Prjibelski AD. RnaQUAST: A quality

assessment tool for de novo transcriptome assemblies. *Bioinformatics*. Oxford University Press;

2016; doi: 10.1093/bioinformatics/btw218.

45. Smith-Unna R, Boursnell C, Patro R, Hibberd JM, Kelly S. TransRate: Reference-free

quality assessment of de novo transcriptome assemblies. *Genome Res*. Cold Spring Harbor

Laboratory Press; 2016; doi: 10.1101/gr.196469.115.

46. Sweeney BA, Petrov AI, Ribas CE, Finn RD, Bateman A, Szymanski M, et al.. RNAcentral

2021: secondary structure integration, improved sequence search and new member databases.

Nucleic Acids Res. 2021; doi: 10.1093/nar/gkaa921.

47. Lakshmanan V, Sujith TN, Bansal D, Shivaprasad P V., Palakodeti D, Krishna S.

Comprehensive annotation and characterization of planarian tRNA and tRNA-derived fragments

(tRFs). *RNA*. 2021; doi: 10.1261/rna.077701.120.

48. Francis WR, Wörheide G. Similar ratios of introns to intergenic sequence across animal

genomes. *Genome Biol Evol*. Oxford University Press; 2017; doi: 10.1093/gbe/evx103.

49. Gudlaugsdottir S, Boswell DR, Wood GR, Ma J. Exon size distribution and the origin of

introns. *Genetica*. 2007; doi: 10.1007/s10709-007-9139-4.

50. Bhasi A, Philip P, Manikandan V, Senapathy P. ExDom: An integrated database for

comparative analysis of the exon-intron structures of protein domains in eukaryotes. *Nucleic

Acids Res*. 2009; doi: 10.1093/nar/gkn746.

51. Öztürk-Çolak A, Marygold SJ, Antonazzo G, Attrill H, Goutte-Gattat D, Jenkins VK, et al..

FlyBase: updates to the *Drosophila* genes and genomes database. *Genetics*. 2024; doi:

10.1093/genetics/iyad211.

52. Berardini TZ, Reiser L, Li D, Mezheritsky Y, Muller R, Strait E, et al.. The arabidopsis

information resource: Making and mining the "gold standard" annotated reference plant genome.

genesis. 2015; doi: 10.1002/dvg.22877.

53. Cheng C, Krishnakumar V, Chan AP, Thibaud☐Nissen F, Schobel S, Town CD. Araport11:

a complete reannotation of the *Arabidopsis thaliana* reference genome. *The Plant Journal*. 2017;

doi: 10.1111/tpj.13415.

54. Holt C, Yandell M. MAKER2: an annotation pipeline and genome-database management

tool for second-generation genome projects. *BMC Bioinformatics*. 2011; doi: 10.1186/1471-

2105-12-491.

55. Hoff KJ, Lomsadze A, Borodovsky M, Stanke M. Whole-Genome Annotation with

BRAKER.

56. Brůna T, Hoff KJ, Lomsadze A, Stanke M, Borodovsky M. BRAKER2: automatic

eukaryotic genome annotation with GeneMark-EP+ and AUGUSTUS supported by a protein

database. *NAR Genom Bioinform*. 2021; doi: 10.1093/nargab/lqaa108.

57. Manni M, Berkeley MR, Seppey M, Zdobnov EM. BUSCO: Assessing Genomic Data

Quality and Beyond. *Curr Protoc*. 2021; doi: 10.1002/cpz1.323.

58. Hoffman M, Wurtzel O. PLANAtools—An interactive gene expression repository for the

planarian Schmidtea mediterranea. *Front Cell Dev Biol*. 2023; doi: 10.3389/fcell.2023.1149537.

59. Sternberg PW, Van Auken K, Wang Q, Wright A, Yook K, Zarowiecki M, et al.. WormBase

2024: status and transitioning to Alliance infrastructure. *Genetics*. 2024; doi:

10.1093/genetics/iyae050.

60. Nowotarski SH, Davies EL, Robb SMC, Ross EJ, Matentzoglu N, Doddihal V, et al..

Planarian Anatomy Ontology: a resource to connect data within and across experimental

platforms. *Development*. 2021; doi: 10.1242/dev.196097.

61. Castillo-Lara S, Abril JF. PlanNET: homology-based predicted interactome for multiple

planarian transcriptomes. *Bioinformatics*. 2018; doi: 10.1093/bioinformatics/btx738.

62. Cui G, Dong K, Zhou J-Y, Li S, Wu Y, Han Q, et al.. Spatiotemporal transcriptomic atlas

reveals the dynamic characteristics and key regulators of planarian regeneration. *Nat Commun*.

2023; doi: 10.1038/s41467-023-39016-0.

63. King HO, Owusu-Boaitey KE, Fincher CT, Reddien PW. A transcription factor atlas of stem

cell fate in planarians. *Cell Rep*. 2024; doi: 10.1016/j.celrep.2024.113843.

64. Fincher CT, Wurtzel O, de Hoog T, Kravarik KM, Reddien PW. Cell type transcriptome

atlas for the planarian *Schmidtea mediterranea*. *Science (1979)*. 2018; doi:

10.1126/science.aaq1736.

65. Plass M, Solana J, Wolf FA, Ayoub S, Misios A, Glažar P, et al.. Cell type atlas and lineage

tree of a whole complex animal by single-cell transcriptomics. *Science (1979)*. 2018; doi:

10.1126/science.aaq1723.

66. Arnone JT, Robbins-Pianka A, Arace JR, Kass-Gergi S, McAlear MA. The adjacent

positioning of co-regulated gene pairs is widely conserved across eukaryotes. *BMC Genomics*.

2012; doi: 10.1186/1471-2164-13-546.

67. Gherman A, Wang R, Avramopoulos D. Orientation, distance, regulation and function of

neighbouring genes. *Hum Genomics*. 2009; doi: 10.1186/1479-7364-3-2-143.

68. Kominek J, Doering DT, Opulente DA, Shen XX, Zhou X, DeVirgilio J, et al.. Eukaryotic

Acquisition of a Bacterial Operon. *Cell*. Cell Press; 2019; doi: 10.1016/j.cell.2019.01.034.

69. Ahmad SS, Samia NSN, Khan AS, Turjya RR, Khan MAAK. Bidirectional promoters: an

enigmatic genome architecture and their roles in cancers. Mol Biol Rep. Springer Science and

Business Media B.V.;

70. Hoock TC, Peters LL, Lux SE. Isoforms of Ankyrin-3 That Lack the NH2-terminal Repeats

Associate with Mouse Macrophage Lysosomes. *J Cell Biol*. 1997; doi: 10.1083/jcb.136.5.1059.

71. Yamankurt G, Wu HC, McCarthy M, Cunha SR. Exon Organization and Novel Alternative

Splicing of Ank3 in Mouse Heart. *PLoS One*. 2015; doi: 10.1371/journal.pone.0128177.

72. Aísa-Marín I, García-Arroyo R, Mirra S, Marfany G. The Alter Retina: Alternative Splicing

of Retinal Genes in Health and Disease. *Int J Mol Sci*. 2021; doi: 10.3390/ijms22041855.

73. Pal S, Gupta R, Kim H, Wickramasinghe P, Baubet V, Showe LC, et al.. Alternative

transcription exceeds alternative splicing in generating the transcriptome diversity of cerebellar

development. *Genome Res*. 2011; doi: 10.1101/gr.120535.111.

Figure Legends

Figure 1. **Identification rates and accuracy metrics for *in-silico* and SIRVs datasets.**

(A) - Transcript identification rates using *in-silico* data. Circles depict the four approaches tested

for short read (SR) data, using *de novo* (DN) or reference-based (RB) assembly and StringTie2

version 2.1.1 (211) or version 2.2.1 (221).   (B) - Accuracy metrics of transcript isoform

identification using SIRVs on SR, mixed reads (MR) and merged reads (MeR) approaches.

Whiskers on the bar plots represent the variability in accuracy across five different StringTie2

parameter configurations tested.

Figure 2. **Workflow diagram for RNA-seq pre-processing, transcript assembly, and**

functional annotation.

Figure 3. **Workflow diagram for user-driven inspection and refinement.**

Figure 4. **Benchmark of SmedAnno versus BRAKER3 on short-read data.**

(A) - BUSCO completeness for annotations of *A. thaliana* (AT), *C. elegans* (CE) and *S.*

mediterranea (SM) transcriptomes. Stacked bars depict the proportion of: complete single-copy,

complete duplicated, fragmented and missing BUSCOs for each pipeline: BRAKER3;

SmedAnno with DeepSplice junction hints merged but without TransDecoder

(SmedAnno_DeepSplice_-TD); the same plus TransDecoder (+TD); and SmedAnno runs

without DeepSplice guidance (noGuide_±TD). Numbers inside the bars correspond to raw

BUSCO counts. (B) - Structural accuracy. Bars show F1 scores (%, harmonic mean of sensitivity

and precision) at exon, intron and transcript level for the same pipelines. Colours denote

pipelines, panels group the three species.

Figure 5. **Representative annotation artifacts generated by SmedAnno pipeline.**

(A) - Fragmented gene. (B) Chimeric transcript. (C) - Functionally chimeric gene with divergent

transcript annotations. Genes are depicted with cyan arrows, while transcripts are shown as

combinations of gray lines and orange rectangles, denoting introns and exons, respectively.

Figure 6. **Venn diagram and a word cloud of GO terms assigned for novel and reference**

genes for the *S. mediterranea* **asexual strain.**

(A) - Venn diagram of GO term overlaps between MSTRG (novel genes) and SMESG (reference

genes). (B) - Word cloud of unique GO terms assigned to SMESG (reference) genes. (C) - Word

cloud of common GO terms assigned to both SMESG and MSTRG genes. (D) - Word cloud of

unique GO terms assigned to MSTRG (novel) genes.

Figure 7. **Comparative analysis of selected** *S. mediterranea* **genome annotations**

(A) - Cumulative distribution of transcript lengths in selected annotations. The y-axis

(logarithmic scale) represents the cumulative count of transcripts/isoforms, indicating how many

transcripts exceed each length threshold on the x-axis (linear scale). (B) - BUSCO completeness

of selected *S. mediterranea* annotations.

Additional Files

Supplementary Table S1. Description of sequencing datasets.

Supplementary Table S2. StringTie2 parameters tested on SIRVs.

Supplementary Table S3. Custom helper scripts used in the SmedAnno workflow

Run a containerized SmedAnno application using

Step 0: Trim adapters & filter reads
TrimGalore *Filtlong*

tool for short-read sequencing data
tool for long-read sequencing data

Step 1: Remove rRNA

Align to rRNA → Keep non-rRNA reads

STAR *Minimap2* *samtools*

Step 3: Assemble genes & transcripts

Generate DeepSplice guides → Assemle in reference-based mode → Assemble in *de novo* mode

StringTie *StringTie*

Step 4: Merge reference-based (RB) assemblies

Merge short-only assemblies Merge mixed assemblies → Merge RB-combo → Final merge with DeepSplice guides

Step 5: Merge *de novo* (DN) assemblies & create pre-final annotation

Merge short-only assemblies Merge mixed assemblies → Merge DN-combo

Step 6: Filter out transcripts artifacts

Identify transcripts with exons > X bp → Identify transcripts with genomic spans > X bp → Remove flagged

Step 7: Isoform comparison

gffcompare Compare assembled transcripts with reference

Step 8: Annotation correction & enhancement *AGAT* Add intron and gene features to annotation → Remove duplicates

Step 9: Determine functional attributes

blastp
blastx
hmmscan
interproscan

TransDecoder Predict ORFs for nuclear and mitochondrial transcripts → Functional prediction

Step 10: Integrate functional attributes

Filter the functional data → Add relevant functional data columns

Step 11: Identify overlapping genes & transcripts

Identify position overlaps → Detect overlapping exons among isoforms

Step 12: Identify reversed duplicates

Identify opposite strand overlaps with reference genes

Remove reversed duplicates ← yes — **Is there a predicted gene matching the reference strand?** — no → Retain reversed duplicates

Step 13: Identify fragmented & chimeric genes

Identify potentially fragmented and chimeric genes (vs. reference) → Identify potentially fragmented genes (de novo) → Identify potentially chimeric genes (functional conflict) → Identify potentially chimeric genes (structural anomaly)

Step 14: Analyse promoter & terminator sequences

Promoter 2.0

Extract intergenic sequences

Analyze promoter regions to verify regulatory coherence

Analyze terminator regions to ensure consistency in gene models

ARNold

Inspect promoter and terminator analysis results to determine if gene models should be merged

- Multiple alignments at identical genomic positions
- Clustering of similar ncRNAs
- Inconsistent naming conventions

Adress issues

Step 15: Annotate non-coding RNA (ncRNA)

seqkit rmdup

BLASTn

Download the RNAcentral database → Linearize → Deduplicate → Identify homologous regions in the reference genome → Classify ncRNA types → Convert alignment results to GTF format → Filter and refine ncRNA annotations

Step 16: Integrate collected information into a unified genome annotation file

Merge protein-coding gene annotations with ncRNA annotations

Step 17: Incorporate gene ontology (GO)

Download the GOA Uniprot database → Integrate GO annotations based on the functional annotations

Step 18: Perform final review and quality assurance

Validate the annotation using external databases or experimental data

A

SMESG

13403
(71.6%)

5180
(27.7%)

143
(0.8%)

MSTRG

B   **Unique GO terms in SMESG genes**

mitotic S phase

modulation by virus of host process

anchoring junction   axon development

fatty acid metabolic process

integral component of plasma membrane

RNA metabolic process

intracellular anatomical structure

cellular protein metabolic process   integral component of membrane   hemopoiesis

multicellular organism development

kinesin binding   transport   viral process   response to drug

microtubule associated complex

Golgi to plasma membrane protein transport

paranode region of axon

C   **Common GO terms**

endoplasmic reticulum

cytoskeleton   signal transduction

axon   mitochondrion   protein binding

metal ion binding   nucleus

nucleolus   plasma membrane   RNA binding

centrosome   nucleoplasm   cytoplasm   ATPase

DNA binding   membrane   cytosol   ATP binding

synapse   dendrite

identical protein binding   zinc ion binding

extracellular exosome

nucleic acid binding

D   **Unique GO terms in MSTRG genes**

8-hydroxy-dADP phosphatase activity

rRNA(pseudouridine-N3-)-methyltransferaseactivity

alginic acid biosynthetic process   dUMP biosynthetic process

ether hydrolase activity   archaeal or bacterial-type flagellum-dependent cell motility

dUTP catabolic process   [protein-PII] uridylyltransferase activity

hyphal cell wall   viral DNA genome packaging   diterpenoid biosynthetic process

BRISC complex   chondroitin 4-sulfotransferase activity   terpene synthase activity

pons development   [heparansulfate]-glucosamine3-sulfotransferase1activity