Title:

A co-speciation dilemma and a lifestyle transition with genomic consequences in *Wolbachia* of Neotropical *Drosophila*

Authors: Konstantinos Papachristos[1], Wolfgang J. Miller[2#], Lisa Klasson[1#]

Author affiliation: [1]Molecular evolution, Department of Cell and Molecular Biology, Science for Life Laboratory, Uppsala University, Uppsala, Sweden.

[2]Lab Genome Dynamics, Department Cell & Developmental Biology, Center for Anatomy and Cell Biology, Medical University of Vienna, Vienna, Austria.

[#]Corresponding authors: wolfgang.miller@meduniwien.ac.at and lisa.klasson@icm.uu.se

Emails:
konstantinos.papachristos@icm.uu.se
wolfgang.miller@meduniwien.ac.at

lisa.klasson@icm.uu.se

Keywords: *Wolbachia*, *Drosophila*, co-speciation, genome reduction, comparative genomics, symbiosis

Abstract

Background: Long-term persistent symbiotic associations may result in co-speciation and can be inferred if species trees of hosts and symbionts are congruent in topology and divergence times. Co-speciation has been seen to occur relatively frequently in obligate associations, but is less common in parasitic or facultative ones, mainly due to the difference in horizontal transmission rates. The long-term vertical inheritance and close host association of obligate endosymbionts also generally result in smaller genomes than in facultative endosymbionts. Here, we investigate co-speciation and genome reduction using highly similar strains of the endosymbiont *Wolbachia* infecting *Drosophila* species from the willistoni and saltans groups, where only one strain, *w*Pau, infecting *D. paulistorum*, is obligate.

Results: We sequenced the *Wolbachia* genomes from five species of the willistoni and saltans groups and constructed phylogenies. Topological congruence was found between these *Wolbachia* strains and the nuclear DNA of their hosts, except for *w*Pau and *D. paulistorum*, but full topological congruence was observed between *Wolbachia* and the host mitochondrial DNA. However, assuming temporal congruence, we estimated extremely low evolutionary rates in *Wolbachia* of 10^{-10}-10^{-11} changes/site/year. Additionally, the obligate *w*Pau strain was found to have a larger genome than closely related facultative strains, mainly due to an ongoing expansion of an IS4 element. Furthermore, *w*Pau has lost a large proportion of its prophage WO genes, but the *cif* genes, known to be involved in the CI phenotype, are intact. Finally, nine of the eleven genes from the prophage WO-associated Undecim cluster are uniquely duplicated.

Conclusions: The congruent topologies between *Wolbachia* and their willistoni and saltans group hosts indicate co-speciation. However, the high similarity between *Wolbachia* strains, which results in low mutation rate estimates, challenges this interpretation. Contrary to the expectations of the genome reduction theory, we observed an increase in genome size in the obligate *w*Pau strain, potentially driven by a decreased population size. Finally, the

duplication of the Undecim cluster despite a major loss of other prophage-associated genes

suggests that the genes in the Undecim cluster are under strong selection and potentially

play a role in the obligate association between wPau and their *D. paulistorum* hosts.

Background

Bacterial symbionts are ubiquitous among animals but vary widely in phenotypic effects,

level of dependency, and evolutionary persistence within host lineages. Mutualistic

symbioses often involve tight associations with the host that may become obligate and

thereby evolutionary persistent in a host lineage over time. In contrast, parasitic or

pathogenic symbionts are usually facultative and evolutionarily short-lived unless they can

manipulate their host to ensure their transmission across generations, by so-called

reproductive parasitism (Zeng and Wiens 2021). Hence, the evolutionary persistence of a

symbiont typically suggests that it is either an obligate mutualist or a reproductive parasite.

The phenotypic effect, dependency, and persistence of a symbiotic association are thus

interconnected and have been shown to influence how symbiont genomes evolve

(McCutcheon and Moran 2012; Fisher et al. 2017). Numerous studies (Wernegreen 2015;

McCutcheon and Moran 2012; Bobay and Ochman 2017) have demonstrated that obligate

symbionts typically have smaller genomes, fewer pseudogenes, repeats, and mobile

elements than facultative symbionts. The reduced genome size in obligate symbionts

reflects their long-term reliance on the host, with the loss of non-essential gene functions

and ongoing deletions facilitated by genetic drift. In contrast, facultative symbionts, which

maintain a more transient association with their hosts, retain larger genomes that provide

greater metabolic versatility and independence as well as more repeats and mobile

elements.

Wolbachia is an intracellular symbiont that infects many different arthropod species

as well as some nematodes (Kaur et al., 2021) and forms both facultative and obligate host

associations. It is mainly vertically transmitted via the maternal line to the next generation

(Russell et al. 2019) but is also frequently horizontally transmitted between host species

(Vancaester and Blaxter 2023). *Wolbachia* can affect many aspects of its host's biology,

including fecundity, immunity, and behaviour (Kaur et al. 2021). However, it is best known in

arthropods for its ability to alter host reproduction. One such reproductive phenotype is

cytoplasmic incompatibility (CI), where infected males are unable to produce offspring with

uninfected females. The reproductive advantage thereby gained by infected females over

uninfected females, can result in the spread of *Wolbachia* into a host population (Shropshire

et al. 2020) as well as persistence in a population once the infection has reached a high

frequency (Turelli and Hoffmann 1995). Even so, only a handful of examples of evolutionary

persistent *Wolbachia* are currently known. Phylogenetic congruence, which is a hallmark of

evolutionary persistence and co-speciation, has, for example, been observed between

obligate mutualistic *Wolbachia* in filarial nematodes (Lefoulon et al. 2016) and bed bugs

(Balvín et al. 2018), but also in non-obligate associations between *Wolbachia* and *Nasonia*

wasps where *Wolbachia* induces CI (Raychoudhury et al. 2008) and *Nomada* solitary bees

(Gerth and Bleidorn 2016) where the phenotypic effect of *Wolbachia* is unknown. However,

phylogenetic congruence between host mitochondrial DNA and *Wolbachia* is regularly

observed for closely related host species, suggesting that the cytoplasmic co-inheritance of

mitochondria and *Wolbachia* is frequently maintained at least over shorter times (Charlat et

al. 2009; Ilinsky 2013; Zhang et al. 2013; Cariou et al. 2017; Bakovic et al. 2020).

Wolbachia genomes generally conform to the genome reduction theory. Facultative

Wolbachia strains infecting various arthropods have genomes ranging between 1.1-1.8 Mb

in size (Werren et al. 2008; Vancaester and Blaxter 2023) with a substantial fraction of

mobile elements and other repeats and relatively many pseudogenes. Obligate mutualistic

Wolbachia of nematodes have smaller genomes, ranging from 0.5-1.0 Mb, with few

pseudogenes and a low fraction of mobile elements or repeats (Lefoulon et al. 2020; Dudzic

et al. 2022). Additionally, prophage WO regions are present in nearly all *Wolbachia* genomes

except in *Wolbachia* of nematodes, where they are absent or highly degraded (Gavotte et al.

2007; Bordenstein and Bordenstein 2022). This is important since prophage WO contains

genes responsible for reproductive parasitism, such as *cifA* and *cifB* responsible for CI

(Beckmann et al. 2017; LePage et al. 2017) and *wmk* involved in male-killing (Perlmutter et

al. 2019), as well as the Octomom region affecting *Wolbachia* titer and virulence (Chrostek

and Teixeira 2015; Duarte et al. 2021). Furthermore, the Eukaryotic Association Module

(EAM) of prophage WO encodes several proteins suggested to play a role in *Wolbachia*-host

interactions (Bordenstein and Bordenstein 2016).

In the Neotropical *Drosophila* of the willistoni and saltans groups, most tested

species carry *Wolbachia* with high similarity to each other and the well-known *w*Au strain

from *Drosophila simulans* (hereafter called *w*Au-like) (Miller and Riegler 2006). The high

frequency of infections and similarity between the *Wolbachia* strains found in the two

Drosophila groups indicates that their ancestor may have been infected with a *w*Au-like

Wolbachia and that the two partners may have been co-speciating. However, this

observation is puzzling, as co-speciation suggests evolutionary persistence and almost all

the *Wolbachia* strains in willistoni and saltans group flies are believed to be facultative

(Müller et al. 2012) and, in some cases, unable to induce CI and/or lack the *cif* genes

(Martinez et al. 2015, Martinez et al. 2021). Only the *Wolbachia* strain *w*Pau infecting *D.*

paulistorum from the willistoni group, a superspecies consisting of multiple reproductively

isolated semispecies, is considered obligate mutualistic, as previous results have shown that

all collected *D. paulistorum* flies are infected. Additionally, complete removal of *w*Pau by

antibiotic treatment causes lethality and reduces the *w*Pau titer, resulting in distorted ovary

development (Miller et al. 2010), thus affecting fecundity. Additionally, *w*Pau is likely involved

in the ongoing host speciation of *D. paulistorum* as it contributes to assortative mating

between different semispecies of *D. paulistorum* (Miller et al. 2010), affects pheromone

profiles (Schneider et al. 2019) and contributes to the distinct gene expression pattern in

heads and abdomens of three semispecies (Baião et al. 2019). *D. paulistorum* is also known

to carry two distinct mitotypes, α and β, and multiple mitochondrial introgressions with

related Neotropical *Drosophila* species likely occurred during divergence of its semispecies,

as indicated by discordant mitochondrial and nuclear phylogenies (Baião et al. 2023). Thus,

if a potential mitochondrial donor species carried *Wolbachia*, *w*Pau could have entered *D.*

paulistorum via introgression.

The Neotropical *Drosophila* system thus offers the opportunity to investigate

evolutionary persistence and co-speciation in non-obligate *Wolbachia* associations and

study genomic differences between very closely related facultative and obligate *Wolbachia*

strains. To do so, we sequenced and assembled *Wolbachia* genomes from five *Drosophila*

species: *D. paulistorum, D. willistoni* and *D. tropicalis* from the willistoni group*, and D.*

prosaltans and *D. septentriosaltans* from the saltans group and inferred whole genome

phylogenies. We found congruence between four of five *Wolbachia* strains and host nuclear

genes and full congruence between *Wolbachia* and host mitochondrial genomes.

Additionally, we discovered that the genome of the obligate *Wolbachia* strain *w*Pau is larger

than closely related facultative strains due to the expansion of IS4 elements. However, the

*w*Pau genome also had features commonly associated with ongoing genome reduction, like

a lower coding percentage and more pseudogenes. Finally, despite a reduced prophage WO

content, we found the WO-associated *cif* genes intact and a unique duplication of the

Undecim cluster, suggesting they are under selection and thus potentially important for the

symbiosis between *w*Pau and its host.

Materials and Methods

DNA Extraction and Sequencing

Drosophila lines used in this study and the sequencing data produced from each of them are

summarized in Tables S1 and S2. For *D. paulistorum* O11, total DNA for Oxford Nanopore

(ONT) sequencing was extracted from many adult flies of both sexes using the MagAttract

HMW DNA Kit (Qiagen) to obtain high molecular weight DNA. For Illumina sequencing, total

DNA was obtained as described in Baiaõ et al 2023. For *Wolbachia* strains *w*WilP98,

*w*ProPE2, *w*PauFG111, *w*PauPOA1 and *w*PauTP37, DNA samples used for PacBio

sequencing were enriched for *Wolbachia* using a differential centrifugation and filtration

procedure to obtain a *Wolbachia* cell pellet, which was then subjected to whole genome

amplification using the Repli-g midi kit (Qiagen) as described in (Ellegaard et al., 2013). For

*w*WilP98 and *w*ProPE2, Illumina sequences enriched for *Wolbachia* were obtained using the

same protocol.

The ONT library was produced and sequenced using an R9.4 PromethION flow cell

at the Uppsala Genome Center, Uppsala, Sweden. Basecalling was performed with Guppy

4.3.4 and the HAC model. Illumina TruSeq libraries were produced and sequenced at the

Uppsala SNP and Seq platform on an Illumina 2500 HiSeq machine generating 2x125 bp

reads. The Illumina reads were quality filtered and trimmed using Trimmomatic-0.30 with

parameters –phred33 ILLUMINACLIP:illumina_adapter.fasta:2:40:15 LEADING:3

TRAILING:3 SLIDINGWINDOW:4:15 MINLEN:95 (Bolger et al., 2014) and error-corrected

using SPAdes-3.5.0 --sc --only-errorcorrection (Bankevich et al., 2012). For PacBio, 5 kb

SMRTbell libraries were created. Each library was run with P6-C4 chemistry in one SMRT

cell on an RSII PacBio instrument at Uppsala Genome Center.

Genome Assemblies

Wolbachia genomes were assembled using several types of data and pipelines. For

*w*PauO11, reads from total DNA sequencing (i.e. including the *Drosophila* host) were filtered

using Filt-long v0.2.1 with quality priority and the parameters "--target_bases 12500000000 -

-min_length 1000 --mean_q_weight 10" and assembled with Nextdenovo v2.5.0 using

correction options: read_cutoff = 1k, genome_size = 250m, sort_options = -m 7g -t 2,

minimap2_options_raw = -t 3, pa_correction = 17, correction_options = -p 2 and assemble

options: minimap2_options_cns = -t 3, nextgraph_options = -a 1. The produced assembly

contained only one *Wolbachia* contig that was subsequently polished with Illumina reads

Pilon v1.24 (Walker et al. 2014) with the --fix indels parameter and circularized. The

genomes of the two strains *w*WilP98 and *w*ProPE2 were assembled using Illumina and

PacBio reads from amplified DNA enriched for *Wolbachia.* The Illumina reads were

assembled using Spades v3.15.5 (Bankevich et al. 2012) with kmer sizes 97,103,107,113

and the PacBio reads were used for scaffolding with the --pacbio parameter. Illumina and

PacBio reads were mapped against the assemblies using BWA-mem (Li and Durbin 2009)

with default parameters for Illumina reads and using PacBio settings for PacBio reads to

check the correctness of the genome assembly. Overlapping contigs were merged in

Consed (Gordon et al. 1998). See Table 1 for assembly statistics of complete genomes. For

all other assemblies, BWA-mem v0.7.17-r1198-dirty with default parameters was used to

map Illumina reads on the *w*PauO11, *w*ProPE and *w*WilP98 genomes and a set of

concatenated *Wolbachia* genomes representing the *Wolbachia* diversity (81 genomes from

supergroups A, B, C, D, E, F, L and M, Table S3). Mapped reads were filtered using

samtools v.1.13 (Danecek et al. 2021) view with the -f 1 -F 12 parameters to include only

pairs where both were mapped. Filtered reads were then subsampled to approximately 100X

with seqtk v1.3-r117-dirty sample -s 100 before assembly. The reads were then assembled

using Spades v3.15.5 with kmer sizes 97,103,107,113, and the *w*PauFG111, *w*PauPOA1

and *w*PauTP37 assemblies were scaffolded with PacBio reads using the flag -pacbio.

The Illumina assemblies were then filtered by removing contigs that deviated from

the mean sample distribution in kmer coverage and GC content. The kmer coverage of the

contigs was determined from the fasta headers in the output from Spades. In cases where

kmer coverage had a bimodal distribution, the minimum between the two peaks was

identified using the quantmod package in R (Ryan and Ulrich 2022) and used as threshold.

Contigs with kmer coverage below this threshold were blasted against the assembled

contigs and were removed if the sequence existed. This procedure ensured that redundant

low coverage contigs were removed. Remaining low coverage contigs were blasted against

additional *Wolbachia* genomes and proteins. Contigs with a blast alignment length longer

than 100 aa and percent identity over 60%, or blast alignment length longer than 200

nucleotides and percent identity over 70%, were kept. Any contig with an extreme GC value

(outside the 99% interval of a normal distribution of GC) was removed. Finally, all remaining

contigs were searched with Kraken against a custom database (containing a selection of

archaea, bacteria, plasmids, viruses, human, *D. melanogaster* and *D. willistoni* sequences)

and the contigs with a match to *Drosophila* were removed. All assemblies were polished

once using Pilon with the Illumina reads used for assembly.

Genome Annotation

For the complete genomes, *w*PauO11, *w*ProPE2 and *w*WilP98, Prokka v1.14.6 (Seemann

2014) with the parameters --genus Wolbachia --compliant --rfam --addgenes --usegenus

was used for annotation. Next, Pseudofinder v1.1.0 (Syberg-Olsen et al. 2022) with

parameters -ce and -db with a database containing all *Wolbachia* proteins from NCBI (2022-

03-02), was used to mark potential pseudogenes. Additionally, hmmsearch as implemented

in pfam_scan.pl was used for domain prediction with the PFAM database (Bateman et al.

2002). The Illumina reads used for the assemblies were mapped back to each genome with

BWA-mem to verify frameshifts and nonsense mutations. All marked pseudogenes were

inspected manually and annotated as pseudogenes if the start or stop codons were missing,

they contained nonsense or frameshift mutations or were truncated by, for example, an IS

element insertions.

The draft assemblies were annotated the same way, except that 1) Pseudofinder was

run with length_pseudo set at 0.8 and the protein database only included *Wolbachia*

genomes *w*Au, *w*Ha, *w*Ma, *w*No, *w*San, *w*Tei, *w*Yak, *w*PauO11, *w*ProPE2 and *w*WilP98 that

were manually annotated either in this or previous studies (Ellegaard et al. 2013; Baiaõ et al.

2021), 2) only pseudogene calls in CDS sequences were inspected manually, 3)

pseudogenes that we identified in the complete genomes (*w*PauO11, *w*ProPE2 and

*w*WilP98), but were not found by pseudofinder in the draft genomes, were transferred to the

draft genome annotation if they matched with high similarity and extensive coverage.

For identifying repeated sequences, nucmer from the MUMmer 4.0 package (Marçais et al.

2018) was used with flags --maxmatch --nosimplify to align each genome to itself. The

positions of non-overlapping repeat bases were then extracted using a perl script. ISEScan v

1.7.2.3 (Xie and Tang 2017) was used to identify IS elements. Bar plots of the percentage of

repeats and IS elements in each genome were made in R (R Core Team 2018; Wickham et

al. 2019). Phage WO regions were annotated by blastp searches to *Wolbachia* genomes

(*w*Mel, *w*Au, *w*Ha, *w*Ma, *w*No, *w*San, *w*Tei, *w*Yak) with previously identified phage WO

regions. All genomic regions that contained similarities to known phage WO genes were

then manually inspected, and borders were determined based on Bordenstein and

Bordenstein 2022.

MLST phylogeny of *Wolbachia* genomes

Genomes that contained the keyword "Wolbachia" in the taxon name, were downloaded

from NCBI using the command line tool "datasets" (https://github.com/ncbi/datasets) on May

the 6th 2022. tblastn of the Blast v2.7.1+ suite (Camacho et al. 2009) with the parameters --

max_hsps 1 --evalue 1e-10 was used to search for the five *w*Mel MLST proteins and

Wolbachia surface protein (Wsp) (GenBank accession numbers AAS13898.1, AAS14036.1,

AAS14201.1, AAS14415.1, AAS14719.1, AAS14884.1) in 1504 downloaded genomes.

Genomes with blast hits for all six genes were kept for phylogenetic inference. Nucleotide

sequences of the identified genes were extracted from each genome using samtools faidx,

and translated, aligned with MAFFT v6.850b using the linsi algorithm (Katoh and Toh 2008),

pruned to remove gap sites present in more than 50% of aligned sequences and then

backtranslated to nucleotides again. A phylogeny of the six genes from 1262 genomes was

then constructed using RaxML v8.2.9 (Stamatakis 2014) with parameters –f a –m

GTRGAMMA –x 12345 –p 12345 -# 100. The resulting phylogenetic tree as visualized in

Figure 1 and all other figures presenting phylogenetic trees (Fig. 2, Fig. 4B, Fig. S1, Fig. S2,

Fig S3, Fig. S5 and Fig. S6) were made using FigTree v1.4.4 (Rambaut 2009).

Protein clustering and strain phylogeny

At the time of downloading *Wolbachia* genomes, the ones from Vancaester and Blaxter 2023 were not yet available. Hence, to include them in our analysis, we downloaded the genomes classified as supergroup A by the authors. We then annotated the genomes from all strains classified as supergroup A *Wolbachia* in the MLST phylogeny and by Vancaester and Blaxter 2023 as described above for our draft assemblies, used their proteomes for clustering and inferred a phylogeny from single-copy genes (Table S4). Clustering was done with Orthofinder v2.5.4 (Emms and Kelly 2019) with -I 3 and filtered all-against-all blastp results as input. Blastp hits between two matching proteins were filtered so that the size of one should not be smaller than 60% of the size of the other, the e-value should be lower than 10^{-5} and the alignment length should be at least 80% of the size of the smaller protein. The nucleotide sequences of genes from orthogroups with single-copy proteins were extracted, translated, aligned with mafft-linsi, pruned to remove gap sites present in more than 50% of the aligned sequences and backtranslated to obtain codon-based nucleotide alignments.

To ensure that the genes used in the phylogeny did not show signs of recombination, we ran Phipack (Bruen et al. 2006) with the -p 100000, -o, -w 50 on all alignments. Phipack runs three tests to detect recombination, the Phi test, the max χ2 test and the NSS test. For a gene to be considered as recombined, at least two of the three recombination tests needed to have a p-value lower than 0.05. Single-copy gene alignments without recombination were then concatenated with geneStitcher.py from Utensils (https://github.com/ballesterus/Utensils/blob/master/geneStitcher.py).

IQ-TREE v2.2.0 (Minh et al. 2020) was then used for phylogenetic inference of the species tree using the concatenated alignments of the single-copy genes, a partition per gene, and the parameters -m MFP, -B 1000 -T 20 --seed 12345. Individual gene trees were inferred using the same parameters. The phylogenetic trees were visualized using FigTree v1.4.4. To test if the data fit better with the placement of *w*Pau according to the host nuclear

phylogeny rather than our strain phylogeny, we followed Shen et al. 2017 and used IQ-TREE

v1.6.12 with the same parameters as in the strain phylogeny together with the -z flag to add

the two topologies (shown in Figure S3) in Newick format and the -wsl and -wpl flags to get

the likelihood for each site and gene. The differences in the gene likelihoods between the

two topologies were calculated and plotted in R (R Core Team 2018; Wickham et al. 2019).

To measure the gene (gCFs) and site (sCFs) concordance factors, we used the same

pruned concatenated phylogeny as in the log-likelihood differences and we produced pruned

gene trees including the same strains with preserved branch lengths using a Python script

and the ete3 library (Huerta-Cepas et al. 2016). To calculate the gene and site concordance

factors, the pruned species tree was then provided to IQ-TREE with the flag -t, the fasta

alignments with -s, the partition file with -p, the gene trees were provided with --gcf and --scf

100 was used.

Substitution rates

Pairwise synonymous substitution rates were calculated with codeml from the PAML v4.8

package (Yang 2007) using alignments of the 620 single-copy *Wolbachia* genes from the

ortholog clustering (Table 2), as well as the alignments of 692 BUSCO genes and 13

mitochondrial genes from *Drosophila* (the same as in Baião et al 2023) generated the same

way as the *Wolbachia* alignments (see above). For substitution rate estimates from

MCMCTree, we calibrated our phylogeny based on estimates from Suvorov et al. 2022 on

the willistoni-saltans group divergence at around 14.2-20.6 million years and the *D. willistoni-*

D. tropicalis divergence at around 4.2-6.1 million years. A rough estimate of 19-21 million

years for the age of the tree and a time unit of 100 million years gives a beta of 156 for the

gamma distribution of rates. We used alpha parameters of 1, 5, and 10 for the shape of the

distribution to describe scenarios of low, medium, and high mutation rates. The whole

alignment of single-copy genes as one partition was given as input (ndata=0, seqtype==0),

and the GTR substitution model (model=8) was used with independent rates (clock=2). Each

of the three rate scenarios was run two times to verify convergence of estimates for the

chains, for 20,100,000 generations (burnin=100,000, sampfreq=1,000, nsample=20,000).

Gene order phylogeny

Syntenic blocks between the complete genomes listed in Table 1 were obtained using

SynChro_linux (January 2015) (Drillon et al. 2014) with a delta value of 2. The output from

SynChro was then used for phylogenetic reconstruction with PhyChro (downloaded July

2023 from http://www.lcqb.upmc.fr/phychro2/) (Drillon et al. 2020).

IS4 element and Undecim cluster copy numbers in draft *w*Pau genomes

To estimate the number of IS4 elements and Undecim cluster copies in the draft *w*Pau

genomes, we first masked all copies of the IS4 element and one copy of the Undecim cluster

in the *w*PauO11 genome. To do that, we created an IS4 element consensus sequence by

blasting (blastn) one IS4 element from *w*PauO11 against the *w*PauO11 genome, extracting

the sequences for all matches, aligning them with MAFFT v7.490 with flags "--maxiterate

1000" and "--localpair" and building a consensus with Jalview v2.11.4.0 (Waterhouse et al.

2009). The consensus IS4 element sequence was then blasted against the *w*PauO11

genome using blastn and all hits were masked. The Undecim cluster copies were first

identified by homology to the Undecim cluster genes in the *w*Mel genome. One of the

Undecim cluster copies, at positions 898.960-913.092 in *w*PauO11, was then manually

masked. The Illumina reads used to assemble the draft *w*Pau genomes were then mapped

to the masked *w*PauO11 genome and the IS4 element consensus sequence using BWA-

mem v0.7.17-r1198-dirty. Repeated areas were identified from the annotation (see above)

and non-repeated regions of the *w*PauO11 genome were identified using "bedtools

complement" and the repeated regions as input. Coverage over the IS4 element consensus

sequence and the unmasked Undecim cluster copy was calculated with "samtools coverage"

and the coverage of the non-repeated regions of the genome was calculated by getting the

average per-base coverage in these regions by using "samtools depth". The copy numbers

of the IS4 element and the Undecim cluster were calculated by dividing the average

coverage over these regions by the average coverage over the non-repeated regions of the

genome.

SNP phylogeny of *w*Pau genomes

To get a more resolved relationship between the *w*Pau strain variants, we called SNPs for all

*w*Pau strain variants from the Illumina reads mapped to the masked *w*PauO11 genome (see

above). Reads with MQ<1 were filtered out and bcftools was used to call SNPs using the

command "bcftools mpileup --threads 6 -Ou -d 1000 -f $genome -b bam.file.list.txt | bcftools

call --threads 6 -cv --ploidy 1 -Ob -o variants_cflag.bcf". To filter low-quality variants, the

command "bcftools view --types snps -i 'QUAL>30 & INFO/DP>10 & MQ>30'" was used.

The variants located in non-repeated areas of the *w*PauO11 genome were extracted using

"bcftools view" with the coordinates identified as described above and "bcftools consensus"

was used to create pseudo-chromosomes for each strain. The sequence for each pseudo-

chromosome was put into a multi-fasta file and used as input for IQ-TREE v2.2.0 using the

same parameters as in the *Wolbachia* strain phylogeny. The phylogeny was visualized using

FigTree v1.4.4.

IS4 element insertion dynamics

To detect differences in IS4 insertions between the *w*Pau genomes, we identified improperly

mapped read pairs in the Illumina reads that were mapped to the masked *w*PauO11 genome

(see above). If an IS4 element is present in the genomes with mapped reads, the insert size

of paired reads in this region is 0, as one read of the pair would map on the masked

Wolbachia genome and the other read would map on the consensus IS4 sequence. To get

such improperly paired reads, we filtered with samtools view using parameters -F2, MQ>0

and -e "tlen==0". Reads that mapped within 750 bases upstream of the start or 750 bases

downstream of the end of an IS4 element in wPauO11 (as specified in a provided bed file)

are from an IS4 element shared with *w*PauO11. Reads mapping at other places are

associated with IS4 insertions not present in *w*PauO11, and they were filtered by using

samtools view with the -U -L flag and the same bed file as above. If an IS4 element is

present in *w*PauO11 but absent in the genome with mapped reads, the reads from a pair are

further apart than expected, specifically the insert size would appear to be larger than the ca.

1500 bases of the IS4 element, and both reads of the pair should map on the masked

Wolbachia genome. To get such improperly paired reads, reads mapping 750 bases

upstream and 750 bases downstream of IS4 elements (as above) and that had an insert size

larger than 1500 bases but shorter than 5000 bases were filtered using samtools view -e

"(tlen>1500 && tlen<5000) || (tlen>-5000 && tlen < -1500)". The filtered read alignments of

improper pairs were visualized in IGV (v2.16.0) to identify unique IS4 element insertions

across the different *w*Pau strain variants. The *w*Pau whole genome SNP phylogeny was

then used to infer where and when IS4 copies were inserted in the *w*Pau strain variants.

Because complete deletions of IS4 elements occur rarely, differences were explained by

insertions, rather than deletions.

Additionally, we inferred the number of IS4 element copies present at each node of

the tree using complete *Wolbachia* genomes. We identified the shared IS4 elements across

the *w*Au-like strains, *w*Mel and *w*Cin2USA1 by blasting (blastn v2.7.1+) genomes to each

other and visualizing the regions flanking IS4 elements, identified by ISEscan, with Artemis

comparison tool (ACT) (Carver et al. 2005). An IS4 element was considered absent from a

genome if a region had the same gene order as an IS4 element-containing region of another

genome but didn't contain an IS4 element or if an IS4 element-containing region in other

genomes shared no synteny with the target genome.

Gene content variation between genomes *w*Au-like *Wolbachia*

To compare gene content across the different *Wolbachia* genomes, we used the gene count

output of Orthofinder to produce a hierarchical clustering of the genomes based on gene

presence/absence. First, we excluded all single-copy genes and then we assigned the count

of multi-copy genes to 1 so that we cluster based on presence and not copy-number. Then,

we used the "agnes" function for agglomerative clustering with the "average" method from

the cluster v2.1.4 package in R (Maechler et al. 2021).

Additionally, the variability in gene content between the wAu-like$_{ws}$ genomes was

investigated by identifying orthogroups where only one or several of the lineages, wPau,

wTro/wWil and wPro/wSpt, were present and one or several lineages were absent (Table 3,

Table S5 and Table S6). Unless otherwise stated, presence was counted if a gene was

present in the complete genome of the lineage (wPauO11, wWilP98 and wProPE2) and at

least one draft genome of the same lineage.

All figures presenting genome and gene comparisons (Fig. 4A, Fig. 7, Fig. S7 and

Fig. S8) were made with genoPlotR (Guy et al. 2010), after using blastn to search each

genome against every other genome.

Prophage WO classification and gene module assignment

To investigate the prophage WO content in detail, we identified prophage WO regions in the

complete wAu-like genomes, wPauO11, wProPE2, wWilP98, wInn, wAu, and in wMel,

wCin2USA1. All proteins of prophage WO origin were re-clustered to orthogroups with

OrthoFinder v2.5.4 with -I=1 and custom blastp (as above), which put most phage WO

paralogues into single orthogroups. We manually validated the orthogroups based on the

functional annotation and gene order in complete genomes (Table S7). The orthogroups

containing proteins with the same functional annotation that were syntenic were manually

merged. For orthogroup, we inferred a phylogeny using IQ-TREE v2.2.0 as described for

single-copy genes. Genes were assigned to WOa and WOb based on whether they formed

a clade with either WOMelA or WOMelB genes and/or were syntenic with WOMelA or

WOMelB. For orthogroups where a WOMelB gene was present, we followed the

classification of Bordenstein and Bordenstein 2022. Otherwise, we looked at what module

the neighboring genes belonged to and assigned the orthogroup to that module. If an

orthogroup had no functional annotation and could not be assigned to a module using the

above method, it was assigned to the EAM module if flanked by genes in a structural module

on one side and an EAM gene on the other side.

Results

We assembled three complete (*w*PauO11, *w*ProPE2 and *w*WilP98) and eleven draft

genomes (Table 1, Table S1) from *Wolbachia* strains infecting three different species of the

Drosophila willistoni group: seven *w*Pau strain variants from the Orinocan (OR) semispecies

of *D. paulistorum* (as determined by the host phylogeny), three *w*Wil strain variants from *D.*

willistoni, one *w*Tro strain variant from *D. tropicalis,* and two species of the *Drosophila*

saltans group: two *w*Pro strains variants from *D. prosaltans* and one *w*Spt strain variant from

D. septentriosaltans (Table S2)*.* We refer to each strain variant by the *Wolbachia* strain

name followed by the *Drosophila* line it was sequenced from, i.e., *w*PauO11 is the *w*Pau

strain variant from the *D. paulistorum* line O11*.*

Table 1. Genomic features in complete *w*Au-like and outgroup genomes.

	*w*PauO11[1]	*w*WilP98[1]	*w*ProPE2[1]	*w*Au[2]	*w*Inn[3]	*w*Mel[4]	*w*Cin2[5]
Genome size(bp)	1,365,198	1,268,527	1,339,466	1,268,455	1,290,587	1,267,782	1,538,351
GC (%)	34.8	35.2	35.2	35.2	35.3	35.2	35.1
CDSs	1000	1016	1033	996	1073	1011	1473
% Coding	73.8	77.5	76.3	75.7	70.1	76.0	86.2
Pseudogenes	177	123	130	122	227	111	216
Repeats (%)	22.9	11.4	15.8	11.0	16.8	9.8	13.6
Phage WO (%)	8.8	10.8	12.6	10.6	9.0	9.5	24.3
IS elements (%)	15.3	7.5	9.4	7.6	9.9	7.0	7.6

[1]*Wolbachia* genomes from Neotropical *Drosophila* sequenced in this study, [2] Baião GC et al., 2021, [3] Hill T et al., 2022, [4] Wu M et al., 2004, [5] Wolfe TM et al., 2021.

Co-speciation between *Wolbachia* and their Neotropical *Drosophila* hosts

To determine if *Wolbachia* co-speciated with their willistoni and saltans group hosts, we

started by producing a phylogeny of our new genomes and other closely related *Wolbachia*

genomes*.* To achieve that, we downloaded all publicly available *Wolbachia* genomes and

inferred a phylogeny using the five Multi-Locus Strain-Typing (MLST) genes (Baldo et al.

2006) and the *wsp* gene. We determined 23 of the 1613 downloaded genomes to be closely

related to ours (Table S4) and clustered their proteomes with the proteomes of our 14

genomes. The resulting 620 clusters with single-copy orthologous genes, all without

detectable recombination, were used to produce a phylogeny.

Figure 1. Phylogeny of *Wolbachia* genomes with high similarity to *w*Au-like_{ws} strains. The
branches with *Wolbachia* strains associated with *Drosophila* species from the willistoni and saltans
groups, i.e., the *w*Au-like_{ws} *Wolbachia* strains, are coloured. The clade referred to as the *w*Au-like is
indicated with an arrow. The tree was inferred from an alignment of 620 single-copy orthologs with IQ-
TREE. Support values from 1000 ultrafast bootstrap replicates are displayed on the nodes of the tree.
Only bootstrap values greater than 60 are shown.

The concatenated gene alignment contained only 0.55% parsimoniously informative

sites (3,088 of 562,755 sites), showing the very close relationship between all 37 strains.

Despite this, the inferred phylogeny had high support for a clade including the five *Wolbachia*

strains from the willistoni and saltans *Drosophila* groups, *w*Tro, *w*Wil, *w*Pau, *w*Pro and *w*Spt

(Figure 1). This clade, which also contained *Wolbachia* strain *w*Au from the old-world host *D.*

simulans, further included *Wolbachia* strains *w*Inn, *w*Bor and *w*Inc, which infect *Drosophila*

species of various other new-world species groups. Thus, from here on, these nine highly

similar *Wolbachia* strains are what we refer to as *w*Au-like, and the strains infecting willistoni

and saltans group flies are referred to as *w*Au-like_{ws}. Finally, as eight of the nine *Drosophila*

hosts are new-world species, we assume that *w*Au-like infections evolved in Americas and

have just recently infected old-world hosts like *D. simulans*.

The concatenated gene alignment of these nine *w*Au-like genomes contained only

0.3% informative sites and 13.2% of the genes had no informative sites at all. Even so,

inside the *w*Au-like clade, all strain and most strain variant relationships, except the position

of *w*Pau (bootstrap 62), were highly supported.

Thus, to evaluate the position of *w*Pau, we calculated the gene (gCF) and site

concordance factors (sCF) for the internal branches of the subtree that only included one

representative genome (*w*PauO11, *w*WilP98, *w*Tro, *w*ProPE2 and *w*SptPLR) of each *w*Au-

like$_{ws}$ strain, which are the focus of the co-speciation analysis. The concordance factors

represent the percentage of gene trees, gCF, or informative sites, sCF, that support a split in

the tree over all other possible splits. After removing single-copy genes without any

informative sites, we found that almost all informative sites (99.6-100%) place *w*PauO11

between the *w*Wil/*w*Tro and *w*Pro/*w*Spt clades and most of the genes (76.1-81.2%) also

agree with this placement (Figure S1). Thus, the position of *w*PauO11 in our inferred

phylogeny had strong support from informative sites and a majority of the individual genes.

Furthermore, the low bootstrap value for the split between *w*Pau and other strains only

occurred when the *w*Inc strain of *D. incompta* was included in our phylogenetic analysis.

When *w*Inc was excluded, the *w*Pau strain variants were in the same position in the tree but

with high bootstrap support (100) (Figure S2). Since our focus is on the topology of the *w*Au-

like$_{ws}$ *Wolbachia* strains, we also inferred phylogenies using only those genomes and an

outgroup. The resulting phylogeny confirms that the position of *w*Pau is supported (bootstrap

support 99) with respect to the other *w*Au-like$_{ws}$ *Wolbachia* strains (Figure 2, left part).

When comparing our inferred *w*Au-like$_{ws}$ strain phylogeny to the nuclear phylogeny of

the willistoni and saltans group hosts (Baião et al. 2023), we saw that the trees were

congruent except for the position of *w*Pau and *D. paulistorum* (Figure 2A). Rather than being

positioned inside the clade of the other two willistoni group strains, *w*Wil and *w*Tro, *w*Pau is

positioned between the willistoni and saltans group (*w*Pro and *w*Spt) strains. However, full

congruence was observed when comparing the *Wolbachia* tree with the phylogeny of

mitochondrial genomes from the same *Drosophila* species (Figure 2B) (Baiaõ et al. 2023).

Figure 2. Phylogenetic comparison between *w*Au-like *Wolbachia* and *Drosophila* of the
willistoni and saltans groups. A) Maximum Likelihood Phylogenies of *w*Au-like *Wolbachia* on the
left and 692 nuclear-encoded proteins of the host on the right (from Baião et al. 2023). The *w*Au-like
Wolbachia phylogeny is a subset of the tree in Figure 1. B) Maximum Likelihood Phylogenies of *w*Au-
like *Wolbachia* on the left (same as in A) and the whole mitochondrial genome of the neotropical
Drosophila host species on the right (from Baiaõ et al. 2023).
489
To test if our *Wolbachia* data fit better with the host nuclear or mitochondrial

topologies, we calculated the log-likelihood difference per gene between topologies

congruent with either the mitochondrial or the nuclear phylogeny. We found that most

Wolbachia genes strongly supported the mitochondrial-like topology, and a few genes

weakly supported the nuclear-like topology (Figure S3). Overall, our results indicate that the

*w*Au-like$_{ws}$ *Wolbachia* strains are highly congruent with the host mitochondrial phylogeny,

and partly congruent with the nuclear phylogeny of the hosts.

For co-speciation to be true, the divergence times of the hosts and *Wolbachia* strains

should also match (referred to as temporal congruence). To get an idea about temporal

congruence, we calculated relative pairwise synonymous substitution rates of *w*Au-like$_{ws}$

Wolbachia and host genes (Table S8) and compared them to previously described cases of

Wolbachia co-speciation in *Nasonia* and *Nomada* (Raychoudhury et al. 2008, Gerth and

Bleidorn 2016) and horizontal transmission in the *D. suzukii* subgroup (Conner et al. 2017).

The relative substitution rates ranged from 0.00025 for *Wolbachia* in the *D. suzukii* subgroup

to 0.31 for *Wolbachia* of *Nasonia* (Table 2), with *w*Au-like$_{ws}$ *Wolbachia* in between (0.0025 -

0.019).

Table 2. Evolutionary rates of *Wolbachia*

Compared *Wolbachia* strains	dS ratio *Wolbachia*/ Host nuclear	Host divergence time (MY)[1]	Mutation rate (substitutions/synonymous site/year)
*w*Wil - *w*Pau	0.019	4.3	5.7×10^{-10}
*w*Tro - *w*Wil	0.0036	5.1	1.2×10^{-10}
*w*Pro - *w*Spt	0.0069	-	-
*w*Pau - *w*Pro	0.0028	17.4	1.3×10^{-10}
*w*Wil - *w*Pro	0.0025	17.4	1.2×10^{-10}
*w*Suz - *w*Spc[2]	0.00025	4.0	7.6×10^{-12}
*w*NlonB1 - *w*NgirB[3]	0.31	0.4	9.0×10^{-9}
*w*Nleu - *w*Nfla[4]	0.015	0.3	4.0×10^{-10}
*w*Nleu - *w*Npa[4]	0.052	0.4	7.0×10^{-10}
*w*Nfla - *w*Npa[4]	0.055	0.4	7.0×10^{-10}
*w*Nfe - *w*Nleu[4]	0.13	2.4	1.0×10^{-9}
*w*Nfe - *w*Nfla[4]	0.13	2.4	1.0×10^{-9}
*w*Nfe - *w*Npa[4]	0.15	2.4	1.1×10^{-9}

[1]Host divergence times of all *Drosophila* species are from schemeA in Suvorov et al. 2022. dS ratios from *Wolbachia* of the *D.*
suzukii subgroup [2]Conner et al. 2017, dS values and host divergence times of *Nasonia* from [3]Raychoudhury et al. 2008, and
Nomada from [4]Gerth and Bleidorn 2016.

We also calculated mutation rates for these *Wolbachia* strains by dividing the

pairwise synonymous substitution rates by the divergence times of their hosts (if available).

The estimated mutation rates for *w*Au-like$_{ws}$ *Wolbachia* were between 1×10^{-10} to 5×10^{-10}

synonymous substitutions per synonymous site per year, which is in the same range as

some of the *Wolbachia* in *Nomada,* but again higher than in *Wolbachia* from the *D. suzukii*

subgroup, where *Wolbachia* was suggested to be horizontally transmitted and lower than the

ones in *Nasonia,* where *Wolbachia* is suggested to co-speciate with the hosts (Table 2,

Table S8). Hence, the evolutionary rates of *w*Au-like$_{ws}$ did not match those from previously

suggested examples of *Wolbachia* co-speciation or horizontal transmission.

Additionally, we estimated the mutation rate of *w*Au-like$_{ws}$ *Wolbachia* in a Bayesian

framework using host divergence times for the willistoni-saltans groups and *D. willistoni – D.*

tropicalis (Suvorov et al. 2022) as priors. This resulted in a mutation rate for *w*Au-like$_{ws}$

Wolbachia between 2.6 - 6.8x10^{-11} substitutions per site per year (Table S9), which is one

order of magnitude smaller than the rate based on synonymous substitutions.

Repeat expansion and rearrangements in the genome of obligate strain *w*Pau

Contrary to the expectations for an obligate strain, the *w*PauO11 genome was larger than all

other *w*Au-like *Wolbachia* genomes (Table 1), had a lower coding content and 25-50% more

repeats than both the genomes of *w*Au-like *Wolbachia* and two closely related facultative

outgroup strains, *w*Mel and *w*Cin2USA1 (Table 1, Figure 3A). Specifically, we discovered a

unique and extensive expansion of an IS4 element, which accounted for 9.9% of the total

genome size of *w*PauO11 (Figure 3B).

Figure 3. Repeats and IS-elements in complete genomes of *w*Au-like strains. A) Percentage of
repeats in each genome. B) Percentage of each IS family in the genomes, as identified by ISEscan.

Since repeats act as templates for genomic rearrangements via intragenomic

homologous recombination, we examined gene order differences between our closely

related genomes. From a whole genome alignment, we saw that the *w*PauO11 genome

exhibited many changes in gene order when compared to other *w*Au-like genomes (Figure

4A). To quantify this, we identified pairwise syntenic blocks between all genomes and used

this information to construct a phylogeny (Figure 4B). In the resulting tree, *w*PauO11 was on

the longest branch, indicating that it has experienced more genomic rearrangements than

the other *Wolbachia* genomes.

Figure 4. Genomic rearrangements in *w*Au-like strains. A) Whole genome alignments of complete
*Wolbachia w*Au-like genomes and outgroup. For each genome, genes are visualized with blue and
pseudogenes with green rectangles on either the forward or reverse strand (above and below the
midline respectively). Similarities between genomes are visualized as red and blue ribbons, with red
representing forward and blue reverse hits. B) Gene order phylogeny of the same genomes as in A.
Branch lengths written on each branch represent the estimated number of genomic rearrangements.

These findings highlight that the *w*PauO11 genome differs significantly from the

genomes of other *w*Au-like$_{ws}$ strains in several ways, such as having a high proportion of

non-coding DNA and repeats and frequent genomic rearrangements. Interestingly, a similar

pattern, except for the large repeat expansion, is also seen in the genome of the *w*Au-like

strain *w*Inn from *D. innubila.*

IS4 expansion is still ongoing in the *w*Pau genomes

Given the significant expansion of IS4 elements in *w*PauO11, we investigated whether these

elements are still actively transposing in *w*Pau genomes. By identifying all complete and

incomplete copies in the *w*PauO11 genome, we estimated that it originally had at least 96

complete IS4 elements (Figure S4). To quantify IS4 elements in all other *w*Pau genomes, we

masked all but one copy in the *w*PauO11 genome, mapped the Illumina reads of each *w*Pau

strain to it and calculated the coverage over the IS4 element relative to non-repetitive

regions. Applying this approach to Illumina reads from *w*PauO11, we estimated that this

genome harboured 98 IS4 copies, which closely matched our observation of 96 copies.

When extended to other *w*Pau variants, the analysis revealed variability in copy numbers of

the IS4 element across genomes (95-124 copies/genome) (Table S10). Additionally, by

examining improper read pairs, we identified the positions of IS4 element insertions that

occurred after the divergence of the *w*Pau strain variants (Table S11). To get a better

resolution of the *w*Pau strain variant relationships and thus be able to calculate the number

of insertions on each branch more correctly, we inferred a phylogeny based on whole

genome SNPs between the *w*Pau variants (Figure S5). Using this tree topology, we

identified seven IS4 element insertion events that occurred after the divergence of the *w*Pau

strain variants (Table S11, Figure S6). Hence, both read mapping approaches indicated that

the IS4 element is still actively transposing in these *w*Pau genomes.

Next, we approximated the rate and timing of IS4 element insertions in all complete

genomes of the *w*Au-like clade. Using copy number counts, synteny comparisons and our

phylogeny, we estimated that the genome of the ancestor of all *w*Au-like strains had four IS4

elements (Table S12) and the number of IS4 element insertions in all non-*w*Pau genomes to

be 22 (Figure S6). Using branch lengths as an approximation of time, we calculated a rate of

6,571 insertions/(substitution/site) for the non-*w*Pau genomes. We inferred 88 IS4 element

insertions on the branch leading up to the ancestor of the *w*Pau variants by estimating a

minimum of 93 IS4 element copies in the genome of the last common ancestor of *w*Pau.

This gave us a rate of 123,422 insertions/(substitution/site). The seven IS4 element

insertions in the *w*Pau strain variants gave us an estimated rate of 83,333

insertions/(substitution/site). The insertion rate of IS4 elements is thus 18.8 times greater in

the branch leading to the *w*Pau ancestor and 12.7 times greater in the current *w*Pau than in

the other analysed *Wolbachia* strains (Table S13). Our analyses thus showed that the IS4

element insertion rate was slightly higher in the *w*Pau genome in the past than it is today, but

expansion is still ongoing with a relatively fast rate.

Major loss of prophage WO genes, but retention of CI genes and duplication

of the Undecim cluster in *w*Pau

To explore which genes might be linked to phenotypic differences among *w*Au-like$_{ws}$ strains,

we clustered the genomes based on the presence and absence of genes.

Figure 5. Gene content clustering of *Wolbachia w*Au-like genomes. The cladogram on the left
shows how the *Wolbachia* strains cluster based on the presence and absence of orthogroups in each
genome, which is depicted in the plot to the right.

Our analysis revealed that the genomes of *w*Pro and *w*Spt clustered together, as did those

of *w*Wil, *w*Tro, and *w*Au, consistent with their phylogenetic relationships (Figure 1). However,

the gene content of *w*Pau was notably distinct from other the *w*Au-like$_{ws}$ genomes (Figure 5).

We analysed the variation in gene content between the *w*Au-like$_{ws}$ strains further, by

identifying which genes were uniquely present or absent in the three lineages, *w*Pau,

*w*Wil/*w*Tro and *w*Pro/*w*Spt. Our results showed that *w*Pau (defined as *w*PauO11 and at least

one *w*Pau draft genome) lacked many genes compared to the other *w*Au-like$_{ws}$ strains

(Figure 5, Table S5), rather than having unique ones (Table 3, Table S6). Most missing

genes were from prophage WO regions (Table 3), as suggested by the lower proportion of

prophage WO sequences in the *w*PauO11 genome. Of the 39 missing prophage WO-

associated genes, most encoded structural phage WO proteins, such as tail, head, or

baseplate proteins, whereas only six belonged to the Eukaryotic Association Module (EAM).

Table 3. Gene content variation between *w*Au-like$_{ws}$ *Wolbachia*. The numbers represent the presence or absence of an orthogroup in one lineage compared to the other two lineages in the table.

	*w*Pau	*w*Wil/*w*Tro	*w*Pro/*w*Spt
Absent (total)	**66**	**18**	**12**
Phage WO	39	6	-
Ankyrin	2	1	2
Hypothetical	9	3	3
Dozen island	-	4	-
Present (total)	**6**	**11**	**16**
Phage WO	-	5	6
Ankyrin	1	-	1
Hypothetical	4	6	2
Dozen island	-	-	6

Given the large number of missing prophage WO genes in *w*PauO11, we wanted to

determine if they all came from the same prophage WO copy. The prophage WO regions

identified in the complete *w*Au-like genomes were highly similar to either the WO-A or WO-B

regions in *w*Mel (Wu et al. 2004) and are referred to here as WOa and WOb, respectively.

By classifying the prophage WO proteins into WOa- and WOb-like, we found that WOa-like

prophages are generally more degraded than WOb-like prophages in all genomes, with the

Lysis, Tail, and Tail fiber modules of the former entirely missing in all *w*Au-like genomes as

well as in *w*Mel and *w*Cin2USA1 (Figure 6).

However, in *w*PauO11, the WOa-like prophage was almost completely absent,

except for a few genes within the Replication and Repair module. The WOb-like prophage in

*w*Pau also displayed a reduced gene content, particularly in the Connector/Baseplate

module. Conversely, the *w*Inn genome, which also has a low proportion of prophage WO,

retained all Baseplate/Connector genes in WOa but instead lost them in WOb. The presence

of genes associated with both WOa and WOb in *w*Pau and other complete *w*Au-like

genomes suggests that intact copies were likely present in their common ancestor. Thus,

*w*Pau has predominantly lost genes from the WOa copy, but both prophage WO copies have

experienced gene loss.

Figure 6. Prophage WO genes in complete *w*Au-like genomes and outgroup. Number of genes
per module for the two prophage WO variants identified in *w*Au-like and related genomes. WOa
genes were found in the same clade as genes from WO-A in *w*Mel and WOb genes were found in the
same clade as genes from WO-B in *w*Mel when inferring single gene phylogenies.

Apart from prophage WO genes, nine ankyrin repeat domain-containing genes

(ANKs) and nine hypothetical proteins were absent from all *w*Pau strain variants but present

in the *w*Wil/*w*Tro and *w*Pro/*w*Spt lineages. Of the remaining 11 protein clusters absent in all

*w*Pau, functions varied and included two non-WO phage proteins, a Fic protein, Type IV

secretion system protein VirB6, Malonyl-CoA decarboxylase, Cytochrome d ubiquinol

oxidase, subunit I and Exopolysaccharide synthesis ExoD-related protein. All missing protein

clusters in *w*Pau were found in *w*Au-like genomes from one or several of the non-*w*Au-like$_{ws}$

strains, indicating that they have likely been lost in *w*Pau rather than gained in the other

*w*Au-like$_{ws}$ lineages.

629   Much fewer protein clusters were uniquely absent in the other two *w*Au-like$_{ws}$

lineages. Twelve clusters were missing from the *w*Pro and *w*Spt genomes, and 20 were

absent in *w*Wil and *w*Tro (Table 3). Noteworthy, the *w*Wil/*w*Tro lineage lacks the prophage

WO-associated genes responsible for CI, *cifA* and *cifB*, as previously observed by Martinez

et al. 2021. Additionally, we found that *cifB* contains a frameshift mutation in the *w*Spt

genome, likely rendering it non-functional (Figure S7, Table S14). Furthermore, the *w*Bor

and *w*Inn strains also have frameshift mutations in *cifB* as well as non-sense mutations in

cifA, while *w*Inc has a non-sense mutation in *cifB* but an intact *cifA*. Thus, the only *w*Au-like

strains that potentially can induce CI are *w*Pro and *w*Pau. Since highly similar Type I *cif*

genes are found in all *w*Au-like genomes except in one clade (*w*Tro/*w*Wil/*w*Au), the common

ancestor of *w*Au-likes likely carried Type I *cif* genes. Hence, our results indicate that the *cif*

genes have been lost or pseudogenized independently multiple times in the *w*Au-like clade.

641   Only six protein-coding genes, encoding four hypothetical proteins, one ANK and a

ComF family protein, were present in *w*PauO11 (and at least one other draft *w*Pau genome)

but absent in the other *w*Au-like$_{ws}$ strains (Table 3). Potentially, one of the hypothetical

proteins and the ANK gene were acquired by *w*Pau, since no homologs were found in any

other *w*Au-like genome. Additionally, one of the hypothetical proteins is homologous to

WD0462 in *w*Mel, which correlated with CI induction in another set of *Wolbachia* strains

(Baião et al. 2021). However, since the gene is pseudogenized in *w*Pro, a strain known to

induce CI, the correlation was not seen among our strains. Finally, the ComF family protein,

with a potential role in DNA uptake, is pseudogenized in other *w*Au-like$_{ws}$ genomes.

More genes are uniquely present in the *w*Wil/*w*Tro and the *w*Pro/*w*Spt lineages. For

example, homologs of WD0463, the neighbouring gene of WD0462, are only intact in

*w*Pro/*w*Spt. This genomic region showed additional variability, with *w*Pau and *w*Pro carrying

an extra gene between WD0462 and WD0463 (Figure S8). Even though the function of

either of these genes is unknown, the region is highly variable between *Wolbachia* genomes

(Baião et al. 2021), even among these very close relatives. Additionally, we noted that six

genes in *w*Pro and five in *w*Spt from the Dozen Island, a genomic region likely stemming

from an integrated plasmid (Baião et al. 2021, Martinez et al. 2022), were not present in

*w*Pau or *w*Wil/*w*Tro. An additional four genes of the Dozen Island were found in *w*Pro/*w*Spt

and *w*Pau but lost from the wWil/wTro lineage.

Finally, although the loss of prophage WO genes appeared to be frequent in the

*w*Pau genome, we identified a large and unique duplication of a prophage WO-associated

region in *w*PauO11 containing ten of the eleven genes in the Undecim cluster (Bordenstein

and Bordenstein 2022) (Figure 7).

Figure 7. Duplication of the phage WO-associated genes in the Undecim cluster in *w*PauO11.
Similarities between genomes are visualized with grey bands and the duplication is marked in yellow.
Genes in orange are part of the Undecim cluster. Mobile elements are dark grey and pseudogenes
green, with the border colour indicating Undecimcluster or mobile element.

To determine if the duplication was present in the other *w*Pau strain variants, we estimated

the copy number by mapping reads from those genomes to the *w*PauO11 genome with one

copy masked. Coverage over the Undecim cluster was twice as large as coverage of non-

repeated sequences for *w*PauFG111, *w*PauPOA1 and *w*PauTP37 (Table S10), indicating

duplication of Undecim in those genomes. For *w*PauFG103, *w*PauFG295 and *w*PauRP,

coverage over the Undecim cluster was more like in non-repeated sequences. However, for

those lines, the standard deviation of coverage for non-repeats was as much as 30% of the

mean, compared to 10% for the other lines. This suggested that non-repeat coverage may

be noisy for some samples, and we thus plotted the coverage of reads from all strains

around and over the Undecim cluster of wPauO11 (Figure S9). Since the coverage was

approximately doubled compared to the surrounding single-copy regions for all *w*Pau strain

variants, the same genomic region is likely duplicated in all sequenced *w*Pau genomes.

680 Discussion

681 Neotropical *Drosophila* species of the willistoni and saltans groups were previously shown to

682 carry closely related *Wolbachia* strains with high similarity to the *w*Au strain from *D. simulans*

683 (Miller and Riegler 2006). Using 14 new *Wolbachia* genomes from five Neotropical

684 *Drosophila* species, we confirm that these strains are highly similar and belong to a

685 monophyletic clade including strain *w*Au – i.e., the *w*Au-likes.

686 The *w*Au-like *Wolbachia* show patterns of co-speciation and horizontal

687 transmission

To investigate the possibility of co-speciation between the *w*Au-like *Wolbachia* strains

infecting willistoni and saltans group flies, *w*Au-like$_{ws}$ strains, and their hosts, we inferred a

phylogeny and compared it to the nuclear and mitochondrial phylogenies of the hosts (Baião

et al. 2023). We found that the phylogeny of *w*Au-like$_{ws}$ strains was congruent with the

nuclear phylogeny of the host species, except for the *Wolbachia* strain *w*Pau and its host *D.*

paulistorum. Complete congruence was, however, found between the topologies of the two

maternally transmitted genetic entities, *Wolbachia* and mitochondria. Hence, based on

phylogenetic congruence, it seems plausible that a *w*Au-like strain was present in the

ancestor of the willistoni and saltans groups at their divergence ca. 20 million years ago

(Powell et al. 2003; Suvorov et al. 2022) and that this strain has co-diverged with the current

hosts ever since. However, phylogenetic congruence can be achieved in the absence of co-

speciation, a phenomenon referred to as pseudo-co-speciation and is especially common if

host switching preferentially occurs between closely related species (De Vienne et al. 2007).

Thus, to infer true co-speciation, temporal congruence between the two compared entities

should be observed. However, since no reliable estimate of the mutation rate exists for any

Wolbachia strain, dating the nodes in our tree will be highly unreliable. Nevertheless, to

attempt to investigate temporal congruence, we calculated the synonymous substitution rate

of *w*Au-like$_{ws}$ strains relative to that of their hosts and compared that to previously reported

cases of *Wolbachia*-host co-speciation. We found the relative rates in our dataset to be quite

constant between different *Wolbachia*-host pairs, with *D. paulistorum- D. willistoni* and their

Wolbachia having the highest ratio, around 3-7 times higher than the others. Even so, all

rates were between one and two orders of magnitude lower than in previous examples of co-

speciation (Raychoudhury et al. 2008; Gerth and Bleidorn 2016; Conner et al. 2017). Thus, if

all suggested co-speciation examples were true, the *w*Au-like$_{ws}$ strains either have evolved

slower than other co-speciating *Wolbachia* strains or the willistoni and saltans group flies

have evolved faster than estimated in previous studies. However, we don't know if the earlier

reported cases represent true examples of co-speciation, since neither has proven temporal

congruence. Additionally, mutation and substitution rates as well as generation times can

vary widely between different organisms, even between very closely related bacterial and

animal species. Hence, we also calculated mutation rates of *Wolbachia,* based on the

synonymous substitution rates and the substitution rate in single-copy genes, assuming that

Wolbachia divergence times are congruent with host divergence times. Using synonymous

substitutions, we got mutation rates between 10^{-10}–10^{-8} substitutions per site per year and

using all substitutions in single-copy genes, we got even lower estimates of the mutation rate

of $3\text{-}7\text{x}10^{-11}$ substitutions per site per year. Both rates are lower than the estimated mutation

rate in other bacteria (Gibson and Eyre-Walker 2019) as well as in *Wolbachia* strain *w*Mel,

where the mutation rate was estimated from population data to be between $2.88\text{x}10^{-9}$ –

1.29×10^{-8} substitutions per site per year (Richardson et al. 2012). However, Richardson et

al., note that the wMel rate is biased, as it includes slightly deleterious mutations only

observed at the population level, and not at the species level. Consequently, for our

calculated rates to match the generational mutation rates of bacteria (ranging between 10^{-10}–

10^{-8} synonymous substitutions per synonymous site per generation (Lynch et al. 2023), all

Wolbachia strains analysed here, except the ones in *Nasonia,* must have a generation time

of almost a year. Since this is even longer than the generation times of the hosts, our

estimated mutation rates are likely incorrect. There are indeed many potential sources of

error for our mutation rate estimates, for example, synonymous substitutions could be under

strong selection, or the recombination rate could be extremely high and eliminate diversity

between strains, both of which could yield underestimated mutation rates. Additionally, the

host divergence times could be wrong, which, depending on whether they are over- or

under-estimated, would result in either lower or higher mutation rates. At this point, we can't

completely rule out any of these factors, although selection seems unlikely given that the

effective population size of *Wolbachia* is probably low, and it is hard to understand how

recombination between the wAu-like$_{ws}$ strains could be so rampant if our *Wolbachia* strains

are only vertically transmitted. However, the divergence times of the hosts are likely a

smaller problem, as they would have to be off by orders of magnitude for our estimated rates

to be comparable to the rates of wMel and other bacteria. Alternatively, for co-speciation to

be believable, most of the *Wolbachia* strains analysed here must have a very low mutation

rate or long generation time, or a combination thereof. If not, neither the wAu-like$_{ws}$ strains

nor the *Wolbachia* strains from *Nomada* and *Nasonia* have co-speciated with their hosts but

may instead have been horizontally transmitted into their hosts via introgression or

alternative mechanisms, as suggested in other similar systems (Turelli et al. 2018; Cooper et

al. 2019).

It is unlikely that *Wolbachia* could have been recently transmitted between the

relatively distantly related species of the willistoni and saltans group via introgression, but

the topological congruence between mitochondria and *Wolbachia* could suggest that

introgressions have facilitated at least some horizontal transmission events. As previously

reported, *D. paulistorum* has two mitotypes, α and β, which were probably both introgressed

from an unknown species within the willistoni group (Baião et al. 2023). Hence, it is possible

that *w*Pau was introgressed into *D. paulistorum* from a willistoni group species together with

one of the mitotypes. Potentially, the β mitotype, since it is most common among *D.*

paulistorum semispecies and all *D. paulistorum* lines in this study carry it. As *w*Pau has

complete and thus likely functional *cif* genes, the spread of *w*Pau and the replacement of the

ancestral *D. paulistorum* mitochondrion could potentially have been driven by CI. However,

to fully answer this question, data from additional *D. paulistorum* semispecies, especially the

ones carrying the α-mitotype, are needed.

Additionally, at least four horizontal transmission events of *w*Au-like *Wolbachia*, i.e.,

*w*Inn, *w*Bor, *w*Inc, and *w*Au, between distantly related *Drosophila* species have likely

occurred relatively recently, as previously suggested (Ballard 2004; Miller and Riegler 2006;

Sheeley and McAllister 2009; Wallau et al. 2016), through mechanisms other than

introgression. We can only speculate about how these *w*Au-like *Wolbachia* strains infected

and spread in these hosts, but CI is not a likely mechanism for the spread since none of

them carry what appear to be functional *cif* genes. The well-studied strain *w*Au has long

been known to not induce CI in its native host, *D. simulans* (Hoffmann et al. 1996). However,

it does provide high antiviral protection in its host (Martinez et al., 2015), a potentially

beneficial phenotype that might have facilitated its spread. Nothing is currently known about

the phenotypic effects *w*Inc has on its host, but both *w*Inn and *w*Bor cause male-killing,

which could have contributed to their spread. Furthermore, *w*Inn was also seen to protect the

host against RNA viruses (Unckless and Jaenike 2012). It is also possible that *w*Au-like

strains provide yet unknown beneficial host effects.

In conclusion, more research is required to determine whether *Wolbachia* switches

hosts so frequently that pseudo-co-speciation is observed independently in multiple clades

and to identify mechanisms that facilitate the transfer and spread of *Wolbachia* infections.

The genome of the obligate *Wolbachia* strain *w*Pau is expanded and has

some genomic features similar to recently host-associated symbionts

When free-living bacteria become host-associated, many of their genes become superfluous

in the new nutrient-rich, stable environment of the host and are hence lost or pseudogenized

due to lack of selection (McCutcheon and Moran 2012). In addition, when symbionts

transition from free-living to host-associated, they experience a reduction in effective

population size that increases the effect of genetic drift (Moran and Plague 2004,

Wernegreen 2015). Consequently, their genomes tend to accumulate slightly deleterious

mutations, including many IS element insertions (Bobay and Ochman 2017). However, over

time, often after becoming obligate for their hosts, symbiont genomes generally shrink

further in size and lose repeated sequences and pseudogenes (McCutcheon and Moran

2012, Bobay and Ochman 2017), leading to a conserved gene order. This stage of the

genome reduction process can, for example, be observed in obligate mutualistic *Wolbachia*

of nematodes that have small genomes (Werren et al. 2008, Dudzic et al. 2022) with few

repeats and mobile elements (Lefoulon et al. 2020) compared to most facultative *Wolbachia*

strains.

To our knowledge, *w*Pau is the only *Wolbachia* strain found in the willistoni and

saltans group of flies that is obligate for the host (Miller et al. 2010). Hence, *w*Pau is

expected to have a smaller genome than its close facultative relatives. Contrary to the

expectation, we found that *w*PauO11 has the largest genome among the *w*Au-like

Wolbachia strains with complete genomes. Genome size increase in *Wolbachia* is generally

associated with a higher proportion of prophage WO (Kampfraath et al. 2019; Lefoulon et al.

2020; Wolfe et al. 2021; Vancaester and Blaxter 2023), but in *w*Pau, we rather found a

reduction of prophage WO content. Instead, we attribute the increase in genome size in

*w*Pau to a still ongoing expansion of an IS4 element that makes up almost 10% of the

genome and has greatly affected the gene order. The extensive IS4 element proliferation,

high rate of genomic rearrangements and reduced coding percentage in the *w*Pau genome

are genomic features that are more common in recently host-associated facultative

symbionts than evolutionary persistent obligate symbionts. All of them, however, suggest a

large influence of genetic drift, potentially caused by a reduction in effective population size.

We can think of at least three possible reasons why the effective population size of *w*Pau

could have decreased compared to its close relatives.

Firstly, under the assumption of horizontal transmission, *w*Pau might have reduced

the effective population size since only a few *Wolbachia* cells likely established the infection

in *D. paulistorum* initially. While we don't see an expansion of IS elements in other *w*Au-like

strains with complete genomes that have likely shifted host recently, i.e., *w*Inn and *w*Au, the

*w*Inn genome shares other features with the *w*Pau genome, including many rearrangements

and low coding density. Possibly, *w*Inn had more repeats but lost them over time due to

deletions. Although these similarities with *w*Inn might indicate that horizontal transmission

could initially cause a higher effect of genetic drift, we do not believe this to be the main

reason for the sustained high levels of genetic drift in *w*Pau.

Secondly, the transition from facultative to obligate host association could have

yielded a reduction in the effective population size of *w*Pau. Several similar genome features

can be found in the genome of the mutualistic *Wolbachia* strain *w*Cle infecting bedbugs

(Hosokawa et al. 2010), including an expansion of IS elements (with over 100 copies of IS5)

and reduced prophage WO content (Lefoulon et al. 2020). Potential IS element expansions

are also recorded in other obligate symbiont genomes, for example, the co-obligate *Serratia*

symbiotica strain Ct from the aphid *Cinara tujafilina* (Manzano-Marín and Latorre 2014),

where several IS element families have expanded (Renoz et al. 2021) and some species of

the genus *Sodalis* (Oakeson et al. 2014; Koga and Moran 2014; Ankrah et al. 2018;

Manzano-Marín et al. 2018; Garber et al. 2021). However, since no genome sequences of

closely related facultative strains are available in any of these cases, the IS element

expansion can't be linked to a transition to obligate host association.

Thirdly, a reduction in the effective population size of *w*Pau could be a consequence

of a reduction in the effective population size of its host, which might be linked to the fact that

D. paulistorum is a superspecies consisting of multiple fully or partly reproductively isolated

semispecies. Currently, no data about the effective population sizes of any species in the

willistoni or saltans groups exist; however, indirect evidence indicates a lower effective

population size in *D. paulistorum* compared to other willistoni and saltans group species.

Large eukaryotic genomes with a high proportion of transposable elements often occur in

species with low effective population sizes (Lynch and Conery 2003), due to the high

influence of genetic drift that allows mobile elements to accumulate. Since *D. paulistorum*

has the largest genome with the highest repetitive content of the sequenced willistoni group

species (Kim et al. 2021, Gebert et al. 2024), we might speculate that *D. paulistorum* has a

lower effective population size than other willistoni and saltans group species. If so, this

could explain not only the high rate of IS4 element insertions before the divergence of the

*w*Pau strain variants but also the continued high rate that seems to be ongoing in *w*Pau even

after its likely introgression and transition from facultative to obligate.

Prophage WO genes may be important for the obligate symbiosis of *w*Pau

The WO prophage is known to play an important role in facultative *Wolbachia* host

associations, given that it carries genes directly responsible for reproductive manipulation

(Beckmann et al. 2017; LePage et al. 2017; Perlmutter et al. 2019; Fricke and Lindsey

2024), infection titer and virulence (Chrostek and Teixeira 2015; Duarte et al. 2021). For the

obligate *w*Pau, we demonstrate loss of prophage WO genes compared to closely related

facultative strains. Specifically, *w*Pau has lost almost an entire copy (WOa) and some parts

of the other (WOb), suggesting a general lack of selection on WO-associated genes rather

than any specific WO prophage region. In contrast, nine of the eleven genes of the prophage

WO-associated Undecim cluster are instead duplicated in *w*PauO11 as well as the other

*w*Pau strain variants. Although the functions of the Undecim cluster genes are currently

unknown, they are thought to play a role in *Wolbachia*-host interactions, as they, among

other functions, encode proteins potentially involved in exopolysaccharide and/or

lipopolysaccharide biosynthesis (Bordenstein and Bordenstein 2022). Previous work has

also shown that the Undecim cluster has been horizontally transferred between several

unrelated insect endosymbionts and that the genes are constitutively highly expressed in

Wolbachia strain *w*Mel during the whole life cycle of *D. melanogaster* (Gutzwiller et al. 2015).

The fact that this prophage WO-associated set of genes is duplicated in *w*Pau, while many

other prophage WO genes are deleted, suggests that the region could encode proteins

potentially important for the obligate symbiosis between *w*Pau and its host. If true, prophage

WO might thus contribute genes for the persistence of a *Wolbachia* infection, even in the

absence of CI or other reproductive manipulations.

Conclusions

We conclude that the ancestor of Neotropical *Drosophila* in the willistoni and saltans groups

may have been infected with a *w*Au-like *Wolbachia* and co-speciated with their hosts, except

for the *w*Pau strain in *D. paulistorum*, which was likely introgression from another species

within the willistoni group together with one of the two mitotypes. However, such an

interpretation suggests that *Wolbachia* must have a very low mutation rate. Since the real

mutation rate of any strain of *Wolbachia i*s still unknown*,* further research is needed to

determine if co-speciation is a possible interpretation.

We also found that contrary to expectations, the genome of the obligate *w*Pau strain

was larger than its facultative relatives due to the ongoing proliferation of an IS4 element,

typically associated with genomes of recently host-associated symbionts. A reduced coding

percentage and more pseudogenes are also hallmarks of the early stages of genome

reduction, all of which are likely due to a reduction in the effective population size.

In general, the effective population size can be reduced multiple times after a

bacterium becomes host associated. Thus, the genome reduction process might take place

in a punctuated manner, as episodes of effective population size reduction can temporarily

lead to an increased genome size via repeat expansion, and a decrease in the proportion of

coding content, only to be reversed by homologous recombination at such repeats over time.

Consequently, only when mobile elements and repeats are lost from a genome altogether

can this cycle be broken.

Data availability

All *Wolbachia* genomes and annotations have been submitted to a public sequence

database and will be made available upon publication. The mitochondrial genomes and

nuclear data are available in NCBI Bioproject PRJNA643793 (Baiaõ et al. 2023). The

accession numbers of other *Wolbachia* genomes used in our analyses can be found in

Supplementary Tables S3 and S4.

Abbreviations

ACT: Artemis Comparison Tool

ANKs: ankyrin repeat domain-containing genes

CI: Cytoplasmic Incompatibility

EAM: Eukaryotic Association Module

ONT: Oxford Nanopore Technologies

SNP: Single Nucleotide Polymorphism

MLST: Multi-Locus Strain-Typing

gCF: Gene Concordance Factor

sCF: Site Concordance Factor

*w*Au-like_{ws}: *Wolbachia* strains infecting fruit flies of the willistoni and saltans groups

WOa/b: WO Phage sequences orthologous to WOMelA and WOMelB, respectively

908   References

Baião GC, Janice J, Galinou M, Klasson L. Comparative Genomics Reveals Factors
Associated with Phenotypic Expression of Wolbachia. Genome Biology and Evolution. 2021
July 1;13(7):evab111.

Baião GC, Schneider DI, Miller WJ, Klasson L. The effect of Wolbachia on gene expression
in Drosophila paulistorum and its implications for symbiont-induced host speciation. BMC
Genomics. 2019 Dec;20(1):465.

Baião GC, Schneider DI, Miller WJ, Klasson L. Multiple introgressions shape mitochondrial
evolutionary history in Drosophila paulistorum and the Drosophila willistoni group. Molecular
Phylogenetics and Evolution. 2023 Mar;180:107683.

Bakovic V, Schebeck M, Stauffer C, Schuler H. Wolbachia-Mitochondrial DNA Associations
in Transitional Populations of Rhagoletis cerasi. Insects. Multidisciplinary Digital Publishing
Institute; 2020 Oct;11(10):675.

Baldo L, Dunning Hotopp JC, Jolley KA, Bordenstein SR, Biber SA, Choudhury RR, et al.
Multilocus Sequence Typing System for the Endosymbiont Wolbachia pipientis. Applied and
Environmental Microbiology. American Society for Microbiology; 2006 Nov;72(11):7098–110.

Ballard JWO. Sequential Evolution of a Symbiont Inferred From the Host: Wolbachia and
Drosophila simulans. Molecular Biology and Evolution. 2004 Mar 1;21(3):428–42.

Balvín O, Roth S, Talbot B, Reinhardt K. Co-speciation in bedbug Wolbachia parallel the
pattern in nematode hosts. Sci Rep. 2018 June 11;8(1):8797.

Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: a new
genome assembly algorithm and its applications to single-cell sequencing. J Comput Biol.
2012 May;19(5):455–77.

Bateman A, Birney E, Cerruti L, Durbin R, Etwiller L, Eddy SR, et al. The Pfam protein
families database. Nucleic Acids Res. 2002 Jan 1;30(1):276–80.

Beckmann JF, Ronau JA, Hochstrasser M. A Wolbachia Deubiquitylating Enzyme Induces
Cytoplasmic Incompatibility. Nat Microbiol. 2017 Mar 1;2:17007.

Blasco-Costa I, Hayward A, Poulin R, Balbuena JA. Next-generation cophylogeny:
unravelling eco-evolutionary processes. Trends in Ecology & Evolution. 2021 Oct
1;36(10):907–18.

Bobay L-M, Ochman H. The Evolution of Bacterial Genome Architecture. Front. Genet.
[Internet]. 2017 [cited 2023 Aug 9];8. Available from:
https://www.frontiersin.org/articles/10.3389/fgene.2017.00072

Bordenstein SR, Bordenstein SR. Eukaryotic association module in phage WO genomes
from Wolbachia. Nat Commun. 2016 Oct 11;7(1):13155.

Bordenstein SR, Bordenstein SR. Widespread phages of endosymbionts: Phage WO
genomics and the proposed taxonomic classification of Symbioviridae. Matic I, editor. PLoS
Genet. 2022 June 6;18(6):e1010227.

Bourtzis K, Nirgianaki A, Markakis G, Savakis C. Wolbachia infection and cytoplasmic
incompatibility in Drosophila species. Genetics. 1996 Nov;144(3):1063–73.

Bruen TC, Philippe H, Bryant D. A simple and robust statistical test for detecting the
presence of recombination. Genetics. 2006 Apr;172(4):2665–81.

Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al. BLAST+:
architecture and applications. BMC Bioinformatics. 2009 Dec 15;10(1):421.

Cariou M, Duret L, Charlat S. The global impact of Wolbachia on mitochondrial diversity and
evolution. Journal of Evolutionary Biology. 2017;30(12):2204–10.

Carver TJ, Rutherford KM, Berriman M, Rajandream M-A, Barrell BG, Parkhill J. ACT: the
Artemis Comparison Tool. Bioinformatics. 2005 Aug 15;21(16):3422–3.

Charlat S, Duplouy A, Hornett EA, Dyson EA, Davies N, Roderick GK, et al. The joint
evolutionary histories of Wolbachia and mitochondria in Hypolimnas bolina. BMC
Evolutionary Biology. 2009 Mar 24;9(1):64.

Charleston MA. Cospeciation. Encyclopedia of Evolutionary Biology [Internet]. Elsevier; 2016
[cited 2023 Sept 25]. p. 381–6. Available from:
https://linkinghub.elsevier.com/retrieve/pii/B9780128000496002006

Chrostek E, Marialva MSP, Yamada R, O'Neill SL, Teixeira L. High Anti-Viral Protection
without Immune Upregulation after Interspecies Wolbachia Transfer. PLOS ONE. Public
Library of Science; 2014 June 9;9(6):e99025.

Chrostek E, Teixeira L. Mutualism Breakdown by Amplification of Wolbachia Genes. Malik
HS, editor. PLoS Biol. 2015 Feb 10;13(2):e1002065.

Conner WR, Blaxter ML, Anfora G, Ometto L, Rota-Stabelli O, Turelli M. Genome
comparisons indicate recent transfer of Ri-like Wolbachia between sister species Drosophila
suzukii and D. subpulchrella. Ecology and Evolution. John Wiley & Sons, Ltd; 2017
Nov;7(22):9391–404.

Cooper BS, Vanderpool D, Conner WR, Matute DR, Turelli M. Wolbachia Acquisition by
Drosophila yakuba-Clade Hosts and Transfer of Incompatibility Loci Between Distantly
Related Wolbachia. Genetics. 2019 Aug;212(4):1399–419.

Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, et al. Twelve years of
SAMtools and BCFtools. GigaScience. 2021 Jan 29;10(2):giab008.

De Vienne DM, Giraud T, Shykoff JA. When can host shifts produce congruent host and
parasite phylogenies? A simulation approach. Journal of Evolutionary Biology.
2007;20(4):1428–38.

Drillon G, Carbone A, Fischer G. SynChro: A Fast and Easy Tool to Reconstruct and
Visualize Synteny Blocks along Eukaryotic Chromosomes. PLOS ONE. Public Library of
Science; 2014 Mar 20;9(3):e92621.

Drillon G, Champeimont R, Oteri F, Fischer G, Carbone A. Phylogenetic Reconstruction
Based on Synteny Block and Gene Adjacencies. Mol Biol Evol. 2020 Sept 1;37(9):2747–62.

Duarte EH, Carvalho A, López-Madrigal S, Costa J, Teixeira L. Forward genetics in
Wolbachia: Regulation of Wolbachia proliferation by the amplification and deletion of an
addictive genomic island. PLOS Genetics. Public Library of Science; 2021 June
18;17(6):e1009612.

Dudzic JP, Curtis CI, Gowen BE, Perlman SJ. A highly divergent *Wolbachia* with a tiny
genome in an insect-parasitic tylenchid nematode. Proc. R. Soc. B. 2022 Sept
28;289(1983):20221518.

Ellegaard KM, Klasson L, Näslund K, Bourtzis K, Andersson SGE. Comparative Genomics
of Wolbachia and the Bacterial Species Concept. Matic I, editor. PLoS Genet. 2013 Apr
4;9(4):e1003381.

Emms DM, Kelly S. OrthoFinder: phylogenetic orthology inference for comparative
genomics. Genome Biology. 2019 Nov 14;20(1):238.

Fisher RM, Henry LM, Cornwallis CK, Kiers ET, West SA. The evolution of host-symbiont
dependence. Nat Commun. 2017 July 4;8:15973.

Frank AC, Amiri H, Andersson SGE. Genome deterioration: loss of repeated sequences and
accumulation of junk DNA. Genetica. 2002;

Fricke LC, Lindsey ARI. Identification of Parthenogenesis-Inducing Effector Proteins in
Wolbachia. Genome Biology and Evolution. 2024 Apr 1;16(4):evae036.

Gavotte L, Henri H, Stouthamer R, Charif D, Charlat S, Boulétreau M, et al. A Survey of the
Bacteriophage WO in the Endosymbiotic Bacteria Wolbachia. Molecular Biology and
Evolution. 2007 Feb 1;24(2):427–35.

Gerth M, Bleidorn C. Comparative genomics provides a timeframe for Wolbachia evolution
and exposes a recent biotin synthesis operon transfer. Nat Microbiol. 2016 Dec
22;2(3):16241.

Gibson B, Eyre-Walker A. Investigating Evolutionary Rate Variation in Bacteria. J Mol Evol.
2019 Dec 1;87(9):317–26.

Gordon D, Abajian C, Green P. Consed: a graphical tool for sequence finishing. Genome
Res. 1998 Mar;8(3):195–202.

Gutzwiller F, Carmo CR, Miller DE, Rice DW, Newton ILG, Hawley RS, et al. Dynamics of
Wolbachia pipientis Gene Expression Across the Drosophila melanogaster Life Cycle. G3
(Bethesda). 2015 Oct 23;5(12):2843–56.

Guy L, Roat Kultima J, Andersson SGE. genoPlotR: comparative gene and genome
visualization in R. Bioinformatics. 2010 Sept 15;26(18):2334–5.

Hill T, Unckless RL, Perlmutter JI. Positive Selection and Horizontal Gene Transfer in the
Genome of a Male-Killing Wolbachia. Molecular Biology and Evolution. 2022 Jan
1;39(1):msab303.

Hosokawa T, Koga R, Kikuchi Y, Meng X-Y, Fukatsu T. Wolbachia as a bacteriocyte-

associated nutritional mutualist. Proceedings of the National Academy of Sciences.

Proceedings of the National Academy of Sciences; 2010 Jan 12;107(2):769–74.

Huerta-Cepas J, Serra F, Bork P. ETE 3: Reconstruction, Analysis, and Visualization of

Phylogenomic Data. Molecular Biology and Evolution. 2016 June 1;33(6):1635–8.

Ilinsky Y. Coevolution of Drosophila melanogaster mtDNA and Wolbachia Genotypes. PLOS

ONE. Public Library of Science; 2013 Jan 17;8(1):e54373.

Kampfraath AA, Klasson L, Anvar SY, Vossen RHAM, Roelofs D, Kraaijeveld K, et al.

Genome expansion of an obligate parthenogenesis-associated Wolbachia poses an

exception to the symbiont reduction model. BMC Genomics. 2019 Dec;20(1):106.

Katoh K, Toh H. Recent developments in the MAFFT multiple sequence alignment program.

Briefings in Bioinformatics. 2008 July 1;9(4):286–98.

Kaur R, Shropshire JD, Cross KL, Leigh B, Mansueto AJ, Stewart V, et al. Living in the

endosymbiotic world of Wolbachia: A centennial review. Cell Host & Microbe. 2021

June;29(6):879–93.

Kim BY, Wang JR, Miller DE, Barmina O, Delaney E, Thompson A, et al. Highly contiguous

assemblies of 101 drosophilid genomes. Coop G, Wittkopp PJ, Sackton TB, editors. eLife.

eLife Sciences Publications, Ltd; 2021 July 19;10:e66405.

Lefoulon E, Bain O, Makepeace BL, d'Haese C, Uni S, Martin C, et al. Breakdown of

coevolution between symbiotic bacteria Wolbachia and their filarial hosts. PeerJ. PeerJ Inc.;

2016 Mar 28;4:e1840.

Lefoulon E, Clark T, Guerrero R, Cañizales I, Cardenas-Calligros JM, Junker K, et al.

Diminutive, degraded but dissimilar: Wolbachia genomes from filarial nematodes do not

conform to a single paradigm. Microb Genom. 2020 Dec;6(12):mgen000487.

LePage DP, Metcalf JA, Bordenstein SR, On J, Perlmutter JI, Shropshire JD, et al. Prophage

WO genes recapitulate and enhance Wolbachia-induced cytoplasmic incompatibility. Nature.

Nature Publishing Group; 2017 Mar;543(7644):243–7.

Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform.

Bioinformatics. 2009 July 15;25(14):1754–60.

Lynch M, Ali F, Lin T, Wang Y, Ni J, Long H. The divergence of mutation rates and spectra
across the Tree of Life. EMBO reports. John Wiley & Sons, Ltd; 2023 Oct 9;24(10):e57561.

Lynch M, Conery JS. The origins of genome complexity. Science. 2003 Nov
21;302(5649):1401–4.

Maechler M, Rousseeuw P, Struyf A, Hubert M, Hornik K. cluster: Cluster Analysis Basics
and Extensions [Internet]. 2021. Available from: https://CRAN.R-project.org/package=cluster

Manzano-Marín A, D'acier AC, Clamens A-L, Orvain C, Cruaud C, Barbe V, et al. A
Freeloader? The Highly Eroded Yet Large Genome of the Serratia symbiotica Symbiont of
Cinara strobi. Genome Biology and Evolution. Oxford University Press; 2018
Sept;10(9):2178.

Manzano-Marín A, Latorre A. Settling Down: The Genome of Serratia symbiotica from the
Aphid Cinara tujafilina Zooms in on the Process of Accommodation to a Cooperative
Intracellular Life. Genome Biology and Evolution. 2014 July 1;6(7):1683–98.

Marçais G, Delcher AL, Phillippy AM, Coston R, Salzberg SL, Zimin A. MUMmer4: A fast
and versatile genome alignment system. PLOS Computational Biology. Public Library of
Science; 2018 Jan 26;14(1):e1005944.

Martinez J, Ant TH, Murdochy SM, Tong L, Filipe A da S, Sinkins SP. Genome sequencing
and comparative analysis of Wolbachia strain wAlbA reveals Wolbachia-associated plasmids
are common. PLOS Genetics. Public Library of Science; 2022 Sept 19;18(9):e1010406.

Martinez J, Klasson L, Welch JJ, Jiggins FM. Life and Death of Selfish Genes: Comparative
Genomics Reveals the Dynamic Evolution of Cytoplasmic Incompatibility. Molecular Biology
and Evolution. 2021 Jan 4;38(1):2–15.

Martinez J, Ok S, Smith S, Snoeck K, Day JP, Jiggins FM. Should Symbionts Be Nice or
Selfish? Antiviral Effects of Wolbachia Are Costly but Reproductive Parasitism Is Not.
McGraw EA, editor. PLoS Pathog. 2015 July 1;11(7):e1005021.

McCutcheon JP, Moran NA. Extreme genome reduction in symbiotic bacteria. Nat Rev
Microbiol. 2012 Jan;10(1):13–26.

Miller WJ, Ehrman L, Schneider D. Infectious Speciation Revisited: Impact of Symbiont-
Depletion on Female Fitness and Mating Behavior of Drosophila paulistorum. Parrish C,
editor. PLoS Pathog. 2010 Dec 2;6(12):e1001214.

Miller WJ, Riegler M. Evolutionary Dynamics of *w* Au-Like *Wolbachia* Variants in Neotropical
Drosophila spp. Appl Environ Microbiol. 2006 Jan;72(1):826–35.

Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, Von Haeseler A, et al.
IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic
Era. Teeling E, editor. Molecular Biology and Evolution. 2020 May 1;37(5):1530–4.

Moran NA, Plague GR. Genomic changes following host restriction in bacteria. Current
Opinion in Genetics & Development. 2004 Dec;14(6):627–33.

Müller MJ, von Mühlen C, Valiati VH, Valente VL da S. Wolbachia pipientis is associated
with different mitochondrial haplotypes in natural populations of Drosophila willistoni. Journal
of Invertebrate Pathology. 2012 Jan 1;109(1):152–5.

Nikoh N, Hosokawa T, Moriyama M, Oshima K, Hattori M, Fukatsu T. Evolutionary origin of
insect–Wolbachia nutritional mutualism. Proceedings of the National Academy of Sciences.
Proceedings of the National Academy of Sciences; 2014 July 15;111(28):10257–62.

Perlmutter JI, Bordenstein SR, Unckless RL, LePage DP, Metcalf JA, Hill T, et al. The phage
gene wmk is a candidate for male killing by a bacterial endosymbiont. PLOS Pathogens.
Public Library of Science; 2019 Sept 10;15(9):e1007936.

Powell JR, Sezzi E, Moriyama EN, Gleason JM, Caccone A. Analysis of a Shift in Codon
Usage in Drosophila. J Mol Evol. 2003 Aug 1;57(1):S214–25.

R Core Team. R: A Language and Environment for Statistical Computing [Internet]. Vienna,
Austria: R Foundation for Statistical Computing; 2018. Available from: https://www.R-
project.org/

Rambaut A. FigTree v1.4: Tree Figure Drawing Tool. [Internet]. 2009. Available from:
http://tree.bio. ed.ac.uk/software/figtree/

Raychoudhury R, Baldo L, Oliveira DCSG, Werren JH. Modes of Acquisition of *Wolbachia*:
Horizontal Transfer, Hybrid Introgression, and Codivergence in the *Nasonia* Species
Complex. Evolution. 2008 Sept;63(1):165–83.

Renoz F, Arai H, Pons I. The genus Sodalis as a resource for understanding the
multifaceted evolution of bacterial symbiosis in insects. Symbiosis. 2024 Mar 1;92(2):187–
208.

Renoz F, Foray V, Ambroise J, Baa-Puyoulet P, Bearzatto B, Mendez GL, et al. At the Gate
of Mutualism: Identification of Genomic Traits Predisposing to Insect-Bacterial Symbiosis in
Pathogenic Strains of the Aphid Symbiont Serratia symbiotica. Front. Cell. Infect. Microbiol.
[Internet]. Frontiers; 2021 June 29 [cited 2024 Aug 8];11. Available from:
https://www.frontiersin.org/journals/cellular-and-infection-
microbiology/articles/10.3389/fcimb.2021.660007/full

Richardson MF, Weinert LA, Welch JJ, Linheiro RS, Magwire MM, Jiggins FM, et al.
Population Genomics of the Wolbachia Endosymbiont in Drosophila melanogaster. PLOS
Genetics. Public Library of Science; 2012 Dec 20;8(12):e1003129.

Russell SL, Chappell L, Sullivan W. A symbiont's guide to the germline. Current Topics in
Developmental Biology [Internet]. Elsevier; 2019 [cited 2023 July 18]. p. 315–51. Available
from: https://linkinghub.elsevier.com/retrieve/pii/S0070215319300559

Ryan JA, Ulrich JM. quantmod: Quantitative Financial Modelling Framework [Internet]. 2022.
Available from: https://CRAN.R-project.org/package=quantmod

Schneider DI, Ehrman L, Engl T, Kaltenpoth M, Hua-Van A, Le Rouzic A, et al. Symbiont-
Driven Male Mating Success in the Neotropical Drosophila paulistorum Superspecies. Behav
Genet. 2019 Jan;49(1):83–98.

Seemann T. Prokka: rapid prokaryotic genome annotation. Bioinformatics. 2014 July
15;30(14):2068–9.

Sheeley SL, McAllister BF. Mobile male-killer: similar Wolbachia strains kill males of
divergent Drosophila hosts. Heredity. Nature Publishing Group; 2009 Mar;102(3):286–92.

Shen X-X, Hittinger CT, Rokas A. Contentious relationships in phylogenomic studies can be
driven by a handful of genes. Nat Ecol Evol. Nature Publishing Group; 2017 Apr 10;1(5):1–
10.

Shropshire JD, Leigh B, Bordenstein SR. Symbiont-mediated cytoplasmic incompatibility:
What have we learned in 50 years? eLife. 2020;9:e61989.

Small ST, Tisch DJ, Zimmerman PA. Molecular epidemiology, phylogeny and evolution of
the filarial nematode Wuchereria bancrofti. Infect Genet Evol. 2014 Dec;0:33–43.

Suvorov A, Kim BY, Wang J, Armstrong EE, Peede D, D'Agostino ERR, et al. Widespread
introgression across a phylogeny of 155 *Drosophila* genomes. Current Biology. 2022 Jan
10;32(1):111-123.e5.

Syberg-Olsen MJ, Garber AI, Keeling PJ, McCutcheon JP, Husnik F. Pseudofinder:
Detection of Pseudogenes in Prokaryotic Genomes. Molecular Biology and Evolution. 2022
July 1;39(7):msac153.

Turelli M, Cooper BS, Richardson KM, Ginsberg PS, Peckenpaugh B, Antelope CX, et al.
Rapid Global Spread of *w*Ri-like *Wolbachia* across Multiple *Drosophila*. Current Biology.
2018 Mar 19;28(6):963-971.e8.

Turelli M, Hoffmann AA. Cytoplasmic incompatibility in Drosophila simulans: dynamics and
parameter estimates from natural populations. Genetics. 1995 Aug 1;140(4):1319–38.

Unckless RL, Boelio LM, Herren JK, Jaenike J. Wolbachia as populations within individual
insects: causes and consequences of density variation in natural populations. Proceedings
of the Royal Society B: Biological Sciences. Royal Society; 2009 May 6;276(1668):2805–11.

Unckless RL, Jaenike J. Maintenance of a Male-Killing Wolbachia in Drosophila Innubila by
Male-Killing Dependent and Male-Killing Independent Mechanisms. Evolution. 2012 Mar
1;66(3):678–89.

Vancaester E, Blaxter M. Phylogenomic analysis of Wolbachia genomes from the Darwin
Tree of Life biodiversity genomics project. Teixeira L, editor. PLoS Biol. 2023 Jan
23;21(1):e3001972.

Wallau GL, da Rosa MT, De Ré FC, Loreto ELS. Wolbachia from Drosophila incompta: just
a hitchhiker shared by Drosophila in the New and Old World? Insect Molecular Biology.
2016;25(4):487–99.

Waterhouse AM, Procter JB, Martin DMA, Clamp M, Barton GJ. Jalview Version 2—a
multiple sequence alignment editor and analysis workbench. Bioinformatics. 2009 May
1;25(9):1189–91.

Wernegreen JJ. Endosymbiont evolution: predictions from theory and surprises from
genomes: Endosymbiont genome evolution. Ann. N.Y. Acad. Sci. 2015 Dec;1360(1):16–35.

Werren JH, Baldo L, Clark ME. Wolbachia: master manipulators of invertebrate biology. Nat
Rev Microbiol. 2008 Oct;6(10):741–51.

Wickham H, Averick M, Bryan J, Chang W, McGowan LD, François R, et al. Welcome to the
tidyverse. Journal of Open Source Software. 2019;4(43):1686.

Wolfe TM, Bruzzese DJ, Klasson L, Corretto E, Lečić S, Stauffer C, et al. Comparative
genome sequencing reveals insights into the dynamics of Wolbachia in native and invasive
cherry fruit flies. Molecular Ecology. 2021;30(23):6259–72.

Wu M, Sun LV, Vamathevan J, Riegler M, Deboy R, Brownlie JC, et al. Phylogenomics of
the Reproductive Parasite Wolbachia pipientis wMel: A Streamlined Genome Overrun by
Mobile Genetic Elements. PLOS Biology. Public Library of Science; 2004 Mar 16;2(3):e69.

Xie Z, Tang H. ISEScan: automated identification of insertion sequence elements in
prokaryotic genomes. Bioinformatics. 2017 Nov 1;33(21):3340–7.

Yang Z. PAML 4: Phylogenetic Analysis by Maximum Likelihood. Molecular Biology and
Evolution. 2007 Apr 18;24(8):1586–91.

Zeng Y, Wiens JJ. Do mutualistic interactions last longer than antagonistic interactions?
Proceedings of the Royal Society B: Biological Sciences. Royal Society; 2021 Sept
8;288(1958):20211457.

Zhang Y-K, Ding X-L, Zhang K-J, Hong X-Y. Wolbachia Play an Important Role in Affecting
mtDna Variation of Tetranychus truncatus (Trombidiformes: Tetranychidae). Environmental
Entomology. 2013 Dec 1;42(6):1240–5.

Declarations

Ethics approval and consent to participate

Not applicable for this study.

Consent for publication

Not applicable for this study.

Competing Interests

The authors declare that they have no competing interests.

Funding

1192 This work was supported by the Swedish research council VR grant 2014-4353 to LK and by

1193 the Austrian Science Fund FWF grant P28255B22 to WJM.

1194 Acknowledgements

1195 We thank Mercè Montoliu Nerín for assembling the *Drosophila paulistorum* O11 genome

1196 from which the *w*PauO11 genome contig was extracted and for helpful discussions

1197 throughout the project, Tom Martin for performing the DNA extraction for Nanopore

1198 sequencing of *D. paulistorum* O11, Lina Juzokaite for *Wolbachia*-enriched DNA extractions

1199 for PacBio sequencing, and Illumina sequencing. We also thank Athanasios Alexiou for

1200 assembling the first version of the *w*WilP98 genome. Sequencing was performed by the

1201 SNP&SEQ Technology Platform and Uppsala Genome Center (UGC) in Uppsala, Sweden.

1202 The facilities are part of the National Genomics Infrastructure (NGI) Sweden and Science for

1203 Life Laboratory. Some of the data handling was enabled by resources provided by the

1204 Swedish National Infrastructure for Computing (SNIC) at UPPMAX. SNP&SEQ, UGC and

1205 Uppmax are supported by the Swedish Research Council and SNP&SEQ are supported by

1206 the Knut and Alice Wallenberg Foundation.

1207 Author contribution

1208 LK and WJM conceived and designed the study. KP and LK performed assemblies. KP and

1209 LK performed annotations. KP and LK performed comparative analyses. KP performed phy-

1210 logenetic analyses. LK and KP wrote the paper with contributions from WJM. All authors

1211 read and approved the manuscript.

49