Title:

A co-speciation dilemma and a lifestyle transition with genomics consequences in *Wolbachia* of Neotropical *Drosophila*

Authors: Konstantinos Papachristos[1], Wolfgang J. Miller[2#], Lisa Klasson[1#]

Author affiliation: [1]Molecular evolution, Department of Cell and Molecular Biology, Science for Life Laboratory, Uppsala University, Uppsala, Sweden.

[2]Lab Genome Dynamics, Department Cell & Developmental Biology, Center for Anatomy and Cell Biology, Medical University of Vienna, Vienna, Austria.

[#]Corresponding authors: wolfgang.miller@meduniwien.ac.at and lisa.klasson@icm.uu.se

Emails:

konstantinos.papachristos@icm.uu.se

wolfgang.miller@meduniwien.ac.at

lisa.klasson@icm.uu.se

Keywords: *Wolbachia*, *Drosophila*, co-speciation, genome reduction, comparative genomics, symbiosis

1 Abstract

Long-term relationships between hosts and their bacterial endosymbionts can result in co-
speciation. For obligate mutualistic associations, this occurs relatively frequently, but it is
less common in parasitic or facultative relationships. Here, we have sequenced genomes of
several very closely related *Wolbachia* strains, infecting fruit flies of the willistoni and saltans
groups of Neotropical *Drosophila*, of which only one, the *w*Pau strain infecting *D.*
paulistorum, is obligate for its host. Using these genomes, we constructed phylogenies and
found congruence between *Wolbachia* and their hosts, except for *w*Pau and *D. paulistorum*,
indicating mainly co-speciation between *Wolbachia* and willistoni and saltans group flies.
However, the high similarity between *Wolbachia* strains implies low evolutionary rates and
challenges the interpretation of our results as co-speciation. Additionally, contrary to the
expectations of the genome reduction theory, we observed a genome size increase in the
obligate *w*Pau strain, which is due to an ongoing expansion of an IS4 element. Finally,
although *w*Pau has lost a large proportion of prophage WO genes, the *cif* genes are intact
and nine genes from the WO-associated Undecim cluster are uniquely duplicated,
suggesting they are under strong selection and potentially play a role in the obligate
association.

Introduction

Bacterial symbionts are ubiquitous among animals but vary widely in phenotypic effects, level of dependency, and evolutionary persistence within host lineages. Mutualistic symbioses often involve tight associations with the host that may become obligate and thereby evolutionary persistent in a host lineage over time. In contrast, parasitic or pathogenic symbionts are usually facultative and evolutionarily short-lived unless they can manipulate their host to ensure their transmission across generations, by so-called reproductive parasitism (Zeng and Wiens 2021). Hence, the evolutionary persistence of a symbiont typically suggests that it is either an obligate mutualist or a reproductive parasite. The phenotypic effect, dependency, and persistence of a symbiotic association are thus interconnected and have been shown to influence how symbiont genomes evolve (McCutcheon and Moran 2012; Fisher et al. 2017). Numerous studies (Wernegreen 2015; McCutcheon and Moran 2012; Bobay and Ochman 2017) have demonstrated that obligate symbionts typically have smaller genomes, fewer pseudogenes, repeats, and mobile elements than facultative symbionts. The reduced genome size in obligate symbionts reflects their long-term reliance on the host, with the loss of non-essential gene functions and ongoing deletions facilitated by genetic drift. In contrast, facultative symbionts, which maintain a more transient association with their hosts, retain larger genomes that provide greater metabolic versatility and independence as well as more repeats and mobile elements.

   Wolbachia is an intracellular symbiont that infects many different arthropod species as well as some nematodes (Kaur et al., 2021) and forms both facultative and obligate host associations. It is mainly vertically transmitted via the maternal line to the next generation (Russell et al. 2019) but is also frequently horizontally transmitted between host species (Vancaester and Blaxter 2023). *Wolbachia* can affect many aspects of its host's biology, including fecundity, immunity, and behaviour (Kaur et al. 2021). However, it is best known in arthropods for its ability to alter host reproduction. One such reproductive phenotype is cytoplasmic incompatibility (CI), where infected males are unable to produce offspring with uninfected females. The reproductive advantage thereby gained by infected females over uninfected females, can result in the spread of *Wolbachia* into a host population (Shropshire et al. 2020) as well as persistence in a population once the infection has reached a high frequency (Turelli and Hoffmann 1995). Even so, only a handful of examples of evolutionary persistent *Wolbachia* are currently known. Phylogenetic congruence, which is a hallmark of evolutionary persistence and co-speciation, has, for example, been observed between obligate mutualistic *Wolbachia* in filarial nematodes (Lefoulon et al. 2016) and bed bugs

(Balvín et al. 2018), but also in non-obligate associations between *Wolbachia* and *Nasonia*
wasps where *Wolbachia* induces CI (Raychoudhury et al. 2008) and *Nomada* solitary bees
(Gerth and Bleidorn 2016) where the phenotypic effect of *Wolbachia* is unknown. However,
phylogenetic congruence between host mitochondrial DNA and *Wolbachia* is regularly
observed for closely related host species, suggesting that the cytoplasmic co-inheritance of
mitochondria and *Wolbachia* is frequently maintained at least over shorter times (Charlat et
al. 2009; Ilinsky 2013; Zhang et al. 2013; Cariou et al. 2017; Bakovic et al. 2020).

        Wolbachia genomes generally conform to the genome reduction theory. Facultative
Wolbachia strains infecting various arthropods have genomes ranging between 1.1-1.8 Mb
in size (Werren et al. 2008; Vancaester and Blaxter 2023) with a substantial fraction of
mobile elements and other repeats and relatively many pseudogenes. Obligate mutualistic
Wolbachia of nematodes have smaller genomes, ranging from 0.5-1.0 Mb, with few
pseudogenes and a low fraction of mobile elements or repeats (Lefoulon et al. 2020; Dudzic
et al. 2022). Additionally, prophage WO regions are present in nearly all *Wolbachia* genomes
except in *Wolbachia* of nematodes, where they are absent or highly degraded (Gavotte et al.
2007; Bordenstein and Bordenstein 2022). This is important since prophage WO contains
genes responsible for reproductive parasitism, such as *cifA* and *cifB* responsible for CI
(Beckmann et al. 2017; LePage et al. 2017) and *wmk* involved in male-killing (Perlmutter et
al. 2019), as well as the Octomom region affecting *Wolbachia* titer and virulence (Chrostek
and Teixeira 2015; Duarte et al. 2021). Furthermore, the Eukaryotic Association Module
(EAM) of prophage WO encodes several proteins suggested to play a role in *Wolbachia*-host
interactions (Bordenstein and Bordenstein 2016).

        In the Neotropical *Drosophila* of the willistoni and saltans groups, most tested
species carry *Wolbachia* with high similarity to each other and the well-known *w*Au strain
from *Drosophila simulans* (hereafter called *w*Au-like) (Miller and Riegler 2006). The high
frequency of infections and similarity between the *Wolbachia* strains found in the two
Drosophila groups indicates that their ancestor may have been infected with a *w*Au-like
Wolbachia and that the two partners may have been co-speciating. However, this
observation is puzzling, as co-speciation suggests evolutionary persistence and almost all
the *Wolbachia* strains in willistoni and saltans group flies are believed to be facultative
(Müller et al. 2012) and, in some cases, unable to induce CI and/or lack the *cif* genes
(Martinez et al. 2015, Martinez et al. 2021)*.* Only the *Wolbachia* strain *w*Pau infecting *D.
paulistorum* from the willistoni group, a superspecies consisting of multiple reproductively
isolated semispecies, is considered obligate mutualistic, as previous results have shown that
all collected *D. paulistorum* flies are infected. Additionally, complete removal of *w*Pau by
antibiotic treatment causes lethality and reducing the *w*Pau titer results in distorted ovary

development (Miller et al. 2010), thus affecting fecundity. Additionally, *w*Pau is likely involved

in the ongoing host speciation of *D. paulistorum* as it contributes to assortative mating

between different semispecies of *D. paulistorum* (Miller et al. 2010), affects pheromone

profiles (Schneider et al. 2019) and contributes to the distinct gene expression pattern in

heads and abdomens of three semispecies (Baião et al. 2019). *D. paulistorum* is also known

to carry two distinct mitotypes, α and β, and multiple mitochondrial introgressions with

related Neotropical *Drosophila* species likely occurred during divergence of its semispecies,

as indicated by discordant mitochondrial and nuclear phylogenies (Baião et al. 2023). Thus,

if a potential mitochondrial donor species carried *Wolbachia*, *w*Pau could have entered *D.*

paulistorum via introgression.

The Neotropical *Drosophila* system thus offers the opportunity to investigate

evolutionary persistence and co-speciation in non-obligate *Wolbachia* associations and

study genomic differences between very closely related facultative and obligate *Wolbachia*

strains. To do so, we sequenced and assembled *Wolbachia* genomes from five *Drosophila*

species: *D. paulistorum, D. willistoni* and *D. tropicalis* from the willistoni group*,* and *D.*

prosaltans and *D. septentriosaltans* from the saltans group and inferred whole genome

phylogenies. We found congruence between four of five *Wolbachia* strains and host nuclear

genes and full congruence between *Wolbachia* and host mitochondrial genomes.

Additionally, we discovered that the genome of obligate *Wolbachia* strain *w*Pau was larger

than closely related facultative strains due to the expansion of IS4 elements. However, the

*w*Pau genome also had features commonly associated with ongoing genome reduction, like

a lower coding percentage and more pseudogenes. Finally, despite a reduced prophage WO

content, we found the WO-associated *cif* genes intact and a unique duplication of the

Undecim cluster, suggesting they are under selection and thus potentially important for the

symbiosis between *w*Pau and its host.

Results

We assembled three complete (*w*PauO11, *w*ProPE2 and *w*WilP98) and eleven draft
genomes (Table 1, Table S1) from *Wolbachia* strains infecting three different species of the
Drosophila willistoni group: seven *w*Pau strain variants from the Orinocan (OR) semispecies
of *D. paulistorum* (as determined by the host phylogeny)*,* three *w*Wil strain variants from *D.
willistoni,* one *w*Tro strain variant from *D. tropicalis,* and two species of the *Drosophila*
saltans group: two *w*Pro strains variants from *D. prosaltans* and one *w*Spt strain variant from
D. septentriosaltans (Table S2)*.* We refer to each strain variant by the *Wolbachia* strain
name followed by the *Drosophila* line it was sequenced from, i.e., *w*PauO11 is the *w*Pau
strain variant from the *D. paulistorum* line O11*.*

The three complete genomes sequenced in this study, *w*PauO11, *w*WilP98 and
*w*ProPE2, were all similar in genome size and content to each other and to previously
sequenced genomes of facultative *Wolbachia* strains (Table 1).

Co-speciation between *Wolbachia* and their Neotropical *Drosophila* hosts

To determine if *Wolbachia* co-speciated with their willistoni and saltans group hosts, we
started by producing a phylogeny of our new genomes and other closely related *Wolbachia*
genomes. To achieve that, we downloaded all publicly available *Wolbachia* genomes and
inferred a phylogeny using the five Multi-Locus Strain-Typing (MLST) genes (Baldo et al.
2006) and the *wsp* gene. We determined 23 of the 1613 downloaded genomes to be closely
related to ours (Table S3) and clustered their proteomes with the proteomes of our 14
genomes. The resulting 620 clusters with single-copy orthologous genes, all without
detectable recombination, were used to produce a phylogeny.

The concatenated gene alignment contained only 0.55% parsimoniously informative
sites (3,088 of 562,755 sites), showing the very close relationship between all 37 strains.
Despite this, the inferred phylogeny had high support for a clade including the five *Wolbachia*
strains from the willistoni and saltans *Drosophila* groups*, w*Tro, *w*Wil, *w*Pau, *w*Pro and *w*Spt
(Figure 1). This clade, which also contained *Wolbachia* strain *w*Au from the old-world host *D.
simulans*, further included *Wolbachia* strains *w*Inn, *w*Bor and *w*Inc, which infect *Drosophila*
species of various other new-world species groups. Thus, from here on, these nine highly
similar *Wolbachia* strains are what we refer to as *w*Au-like, and the strains infecting willistoni
and saltans group flies are referred to as *w*Au-like$_{ws}$.

The concatenated gene alignment of these nine *w*Au-like genomes contained only
0.3% informative sites and 13.2% of the genes had no informative sites at all. Even so,
inside the *w*Au-like clade, all strain and most strain variant relationships, except the position
of *w*Pau (bootstrap 62), were highly supported. Thus, to evaluate the position of *w*Pau, we

calculated the gene (gCF) and site concordance factors (sCF) for the internal branches of

the subtree that only included one representative genome (wPauO11, wWilP98, wTro,

wProPE2 and wSptPLR) of each wAu-like$_{ws}$ strain, which are the focus of the co-speciation

analysis. The concordance factors represent the percentage of gene trees, gCF, or

informative sites, sCF, that support a split in the tree over all other possible splits. After

removing single-copy genes without any informative sites, we found that almost all

informative sites (99.6-100%) place wPauO11 between the wWil/wTro and wPro/wSpt

clades and most of the genes (76.1-81.2%) also agree with this placement (Figure S1).

Thus, the position of wPauO11 in our inferred phylogeny had strong support from informative

sites but a bit less from individual genes. Furthermore, the low bootstrap value for the split

between wPau and other strains only occurred when the wInc strain of *D. incompta* was

included in our phylogenetic analysis. When wInc was excluded, the wPau strain variants

were in the same position in the tree but with high bootstrap support (100) (Figure S2).

Although we were unable to find the underlying reason for the conflicting signal, our

analyses still indicate that the position of wPau is reliable.

When comparing our inferred wAu-like strain phylogeny to the nuclear phylogeny of

the willistoni and saltans group hosts (Baiaõ et al. 2023), we saw that the trees were

congruent except for the position of wPau and *D. paulistorum* (Figure 2A). Rather than being

positioned inside the clade of the other two willistoni group strains, wWil and wTro, wPau is

positioned between the willistoni and saltans group (wPro and wSpt) strains. However, full

congruence was observed when comparing the *Wolbachia* tree with the phylogeny of

mitochondrial genomes from the same *Drosophila* species (Figure 2B) (Baiaõ et al. 2023).

To test if our *Wolbachia* data fit better with the host nuclear or mitochondrial topologies, we

calculated the log-likelihood difference per gene between topologies congruent with either

the mitochondrial or the nuclear phylogeny. We found that most *Wolbachia* genes strongly

supported the mitochondrial-like topology, and a few genes weakly supported the nuclear-

like topology (Figure S3). Overall, our results indicate that the wAu-like$_{ws}$ *Wolbachia* strains

are highly congruent with the host mitochondrial phylogeny, and partly congruent with the

nuclear phylogeny of the hosts.

For co-speciation to be true, the divergence times of the hosts and *Wolbachia* strains

should match (referred to as temporal congruence). To get an idea about temporal

congruence, we calculated relative pairwise synonymous substitution rates of wAu-like$_{ws}$

Wolbachia and host genes (Table S4) and compared them to previously described cases of

co-speciation (*Nomada* and *Nasonia*) (Raychoudhury et al. 2008, Gerth and Bleidorn 2016)

and horizontal transmission (*D. suzukii* subgroup) (Conner et al. 2017) of *Wolbachia*.

Furthermore, we calculated mutation rates for wAu-like$_{ws}$ *Wolbachia* as well as for *Wolbachia*

strains in *Nomada, Nasonia* and the *D. suzukii* subgroup, by dividing the pairwise

synonymous substitutions rate by the divergence times of their hosts. The relative

substitution rates had a broad range from 0.0003 for *Wolbachia* in the *D. suzukii* subgroup,

where *Wolbachia* was suggested to be horizontally transmitted, to 0.31 for *Wolbachia* of

Nasonia, which was suggested to be co-speciating. *Wolbachia* of *Nomada* had ratios

between 0.015-0.15 and *w*Au-like$_{ws}$ *Wolbachia* between 0.0025 to 0.019, i.e., one to two

orders of magnitude lower than in *Nasonia* and some *Nomada* pairs where *Wolbachia* is

suggested to co-speciate with the hosts, but one to two orders of magnitude higher than in

the *D. suzukii* subgroup where *Wolbachia* was suggested to be horizontally transmitted. The

estimated mutation rates were 9×10^{-9} for *Wolbachia* in *Nasonia,* 4×10^{-10} to 1×10^{-9} in *Nomada*

Wolbachia and 7×10^{-12} in *Wolbachia* from the *D. suzukii* subgroup, while the mutation rate

for *w*Au-like$_{ws}$ *Wolbachia* was 1×10^{-10} to 5×10^{-10} synonymous substitutions per synonymous

site per year (Table S4). Hence, the evolutionary rates of *w*Au-like$_{ws}$ did not match those

from earlier suggested *Wolbachia* co-speciation or horizontal transmission examples.

Repeat expansion and rearrangements in the genome of obligate strain *w*Pau

Contrary to the expectations for an obligate strain, the *w*PauO11 genome was larger than all

other *w*Au-like *Wolbachia* genomes (Table 1), had a lower coding content and 25-50% more

repeats than both the genomes of *w*Au-like *Wolbachia* and two closely related facultative

outgroup strains, *w*Mel and *w*Cin2USA1 (Table 1, Figure 3A). Specifically, we discovered a

unique and extensive expansion of an IS4 element, which accounted for 9.9% of the total

genome size of *w*PauO11 (Figure 3B).

Since repeats act as templates for genomic rearrangements via intragenomic

homologous recombination, we examined gene order differences between our closely

related genomes.From a whole genome alignment, we saw that the *w*PauO11 genome

exhibited many changes in gene order when compared to other *w*Au-like genomes (Figure

4A). To quantify this, we identified pairwise syntenic blocks between all genomes and used

this information to construct a phylogeny (Figure 4B). In the resulting tree, *w*PauO11 was on

the longest branch, indicating that it has experienced more genomic rearrangements than

the other *Wolbachia* genomes.

These findings highlight that the *w*PauO11 genome differs significantly from the

genomes of other *w*Au-like$_{ws}$ strains in several ways, such as having a high proportion of

non-coding DNA and repeats and frequent genomic rearrangements. Interestingly, a similar

pattern, except for the large repeat expansion, is also seen in the genome of the *w*Au-like

strain *w*Inn from *D. innubila.*

IS4 expansion is still ongoing in the *w*Pau genomes

Given the significant expansion of IS4 elements in *w*PauO11, we investigated whether these elements are still actively transposing in *w*Pau genomes. By identifying all complete and incomplete copies in the *w*PauO11 genome, we estimated that it originally had at least 96 complete IS4 elements (Figure S4). To quantify IS4 elements in all other *w*Pau genomes, we masked all but one copy in the *w*PauO11 genome, mapped the Illumina reads of each *w*Pau strain to it and calculated the coverage over the IS4 element relative to non-repetitive regions. Applying this approach to Illumina reads from *w*PauO11, we estimated that this genome harboured 98 IS4 copies, which closely matched our observation of 96 copies. When extended to other *w*Pau variants, the analysis revealed variability in copy numbers of the IS4 element across genomes (95-124 copies/genome) (Table S5). Additionally, by examining improper read pairs, we identified the positions of IS4 element insertions that occurred after the divergence of the *w*Pau strain variants (Table S6). To get a better resolution of the *w*Pau strain variant relationships and thus be able to calculate the number of insertions on each branch more correctly, we inferred a phylogeny based on whole genome SNPs between the *w*Pau variants (Figure S5). Using this tree topology, we identified seven IS4 element insertion events that occurred after the divergence of the *w*Pau strain variants (Table S7, Figure S6). Hence, both read mapping approaches indicated that the IS4 element is still actively transposing in these *w*Pau genomes.

Next, we approximated the rate and timing of IS4 element insertions in all complete genomes of the *w*Au-like clade. Using copy number counts, synteny comparisons and our phylogeny, we estimated that the genome of the ancestor of all *w*Au-like strains had four IS4 elements and the number of IS4 element insertions in all non-*w*Pau genomes to be 22 (Figure S6). Using branch lengths as an approximation of time, we calculated a rate of 6,571 insertions/(substitution/site) for the non-*w*Pau genomes. We inferred 88 IS4 element insertions on the branch leading up to the ancestor of the *w*Pau variants, by estimating a minimum of 93 IS4 element copies in the genome of the last common ancestor of *w*Pau. This gave us a rate of 123,422 insertions/(substitution/site). The seven IS4 element insertions in the *w*Pau strain variants gave us an estimated rate of 83,333 insertions/(substitution/site). The insertion rate of IS4 elements is thus 18.8 times greater in the branch leading to the *w*Pau ancestor and 12.7 times greater in the current *w*Pau than in the other analysed *Wolbachia* strains (Table S8). Our analyses thus showed that the IS4 element insertion rate was slightly higher in the *w*Pau genome in the past than it is today, but expansion is still ongoing with a relatively fast rate.

Major loss of prophage WO genes, but retention of CI genes and duplication of the Undecim cluster in *w*Pau

To explore which genes might be linked to phenotypic differences among *w*Au-like$_{ws}$ strains,

we clustered the genomes based on the presence and absence of genes. Our analysis

revealed that the genomes of *w*Pro and *w*Spt clustered together, as did those of *w*Wil, *w*Tro,

and *w*Au, consistent with their phylogenetic relationships (Figure 1). However, the gene

content of *w*Pau was notably distinct from other the *w*Au-like$_{ws}$ genomes (Figure 5).

We analysed the variation in gene content between the *w*Au-like$_{ws}$ strains further, by

identifying which genes were uniquely present or absent in the three lineages, *w*Pau,

*w*Wil/*w*Tro and *w*Pro/*w*Spt. Our results showed that *w*Pau (defined as *w*PauO11 and at least

one *w*Pau draft genome) lacked many genes compared to the other *w*Au-like$_{ws}$ strains

(Figure 5, Table S9), rather than having unique ones (Table 2, Table S10). Most missing

genes were from prophage WO regions (Table 2), as suggested by the lower proportion of

prophage WO sequences in the *w*PauO11 genome. Of the 39 missing prophage WO-

associated genes, most encoded structural phage WO proteins, such as tail, head, or

baseplate proteins, whereas only six belonged to the Eukaryotic Association Module (EAM).

Given the large number of missing prophage WO genes in *w*PauO11, we wanted to

determine if they all came from the same prophage WO copy. The prophage WO regions

identified in the complete *w*Au-like genomes were highly similar to either the WO-A or WO-B

regions in *w*Mel (Wu et al. 2004) and are referred to here as WOa and WOb, respectively.

By classifying the prophage WO proteins into WOa- and WOb-like, we found that WOa-like

prophages are generally more degraded than WOb-like prophages in all genomes, with the

Lysis, Tail, and Tail fiber modules of the former entirely missing in all *w*Au-like genomes as

well as in *w*Mel and *w*Cin2USA1 (Figure 6). However, in *w*PauO11, the WOa-like prophage

was almost completely absent, except for a few genes within the Replication and Repair

module. The WOb-like prophage in *w*Pau also displayed a reduced gene content,

particularly in the Connector/Baseplate module. Conversely, the *w*Inn genome, which also

has a low proportion of prophage WO, retained all Baseplate/Connector genes in WOa but

instead lost them in WOb. The presence of genes associated with both WOa and WOb in

*w*Pau and other complete *w*Au-like genomes suggests that intact copies were likely present

in their common ancestor. Thus, *w*Pau has predominantly lost genes from the WOa copy,

but both prophage WO copies have experienced gene loss.

Apart from prophage WO genes, nine ankyrin repeat domain-containing genes

(ANKs) and nine hypothetical proteins were absent from all *w*Pau strain variants but present

in the *w*Wil/*w*Tro and *w*Pro/*w*Spt lineages. Of the remaining 11 protein clusters absent in all

*w*Pau, functions varied and included two non-WO phage proteins, a Fic protein, Type IV

secretion system protein VirB6, Malonyl-CoA decarboxylase, Cytochrome d ubiquinol

oxidase, subunit I and Exopolysaccharide synthesis ExoD-related protein. All missing protein

clusters in *w*Pau were found in *w*Au-like genomes from one or several of the non-*w*Au-like$_{ws}$

strains, indicating that they have likely been lost in *w*Pau rather than gained in the other

*w*Au-like$_{ws}$ lineages.

Much fewer protein clusters were uniquely absent in the other two *w*Au-like$_{ws}$

lineages. Twelve clusters were missing from the *w*Pro and *w*Spt genomes, and 20 were

absent in *w*Wil and *w*Tro (Table 2). Noteworthy, the *w*Wil/*w*Tro lineage lacks the prophage

WO-associated genes responsible for CI, *cifA* and *cifB*, as previously observed by Martinez

et al. 2021. Additionally, we found that *cifB* contains a frameshift mutation in the *w*Spt

genome, likely rendering it non-functional (Figure S7, Table S11). Furthermore, the *w*Bor

and *w*Inn strains also have frameshift mutations in *cifB* as well as non-sense mutations in

cifA, while *w*Inc has a non-sense mutation in *cifB* but an intact *cifA*. Thus, the only *w*Au-like

strains that potentially can induce CI are *w*Pro and *w*Pau. Since highly similar Type I *cif*

genes are found in all *w*Au-like genomes except in one clade (*w*Tro/*w*Wil/*w*Au), the common

ancestor of *w*Au-likes likely carried Type I *cif* genes. Hence, our results indicate that the *cif*

genes have been lost or pseudogenized independently multiple times in the *w*Au-like clade.

Only six protein-coding genes, encoding four hypothetical proteins, one ANK and a

ComF family protein, were present in *w*PauO11 (and at least one other draft *w*Pau genome)

but absent in the other *w*Au-like$_{ws}$ strains (Table 2). Potentially, one of the hypothetical

proteins and the ANK gene were acquired by *w*Pau, since no homologs were found in any

other *w*Au-like genome. Additionally, one of the hypothetical proteins is homologous to

WD0462 in *w*Mel, which correlated with CI induction in another set of *Wolbachia* strains

(Baião et al. 2021). However, since the gene is pseudogenized in *w*Pro, a strain known to

induce CI, the correlation was not seen among our strains. Finally, the ComF family protein,

with a potential role in DNA uptake, is pseudogenized in other *w*Au-like$_{ws}$ genomes.

More genes are uniquely present in the *w*Wil/*w*Tro and the *w*Pro/*w*Spt lineages. For

example, homologs of WD0463, the neighbouring gene of WD0462, are only intact in

*w*Pro/*w*Spt. This genomic region showed additional variability, with *w*Pau and *w*Pro carrying

an extra gene between WD0462 and WD0463 (Figure S8). Even though the function of

either of these genes is unknown, the region is highly variable between *Wolbachia* genomes

(Baião et al. 2021), even among these very close relatives. Additionally*,* we noted that six

genes in *w*Pro and five in *w*Spt from the Dozen Island, a genomic region likely stemming

from an integrated plasmid (Baião et al. 2021, Martinez et al. 2022), were not present in

*w*Pau or *w*Wil/*w*Tro. An additional four genes of the Dozen Island were found in *w*Pro/*w*Spt

and *w*Pau but lost from the wWil/wTro lineage.

Finally, although the loss of prophage WO genes appeared to be frequent in the

*w*Pau genome, we identified a large and unique duplication of a prophage WO-associated

region in *w*PauO11 containing ten of the eleven genes in the Undecim cluster (Bordenstein

and Bordenstein 2022) (Figure 7). To determine if the duplication was present in the other

*w*Pau strain variants, we estimated the copy number by mapping reads from those genomes

to the *w*PauO11 genome with one copy masked. Coverage over the Undecim cluster was

twice as large as coverage of non-repeated sequences for *w*PauFG111, *w*PauPOA1 and

*w*PauTP37 (Table S5), indicating duplication of Undecim in those genomes. For

*w*PauFG103, *w*PauFG295 and *w*PauRP, coverage over the Undecim cluster was more like

in non-repeated sequences. However, for those lines, the standard deviation of coverage for

non-repeats was as much as 30% of the mean, compared to 10% for the other lines. This

suggested that non-repeat coverage may be noisy for some samples, and we thus plotted

the coverage of reads from all strains around and over the Undecim cluster of wPauO11

(Figure S9). Since the coverage was approximately doubled compared to the surrounding

single-copy regions for all *w*Pau strain variants, the same genomic region is likely duplicated

in all sequenced *w*Pau genomes.

Discussion

Neotropical *Drosophila* species of the willistoni and saltans groups were previously shown to carry closely related *Wolbachia* strains with high similarity to the *w*Au strain from *D. simulans* (Miller and Riegler 2006). Using 14 new *Wolbachia* genomes from five Neotropical *Drosophila* species, we confirm that these strains are highly similar and belong to a monophyletic clade including strain *w*Au – i.e. the *w*Au-likes.

The *w*Au-like *Wolbachia* show patterns of co-speciation and horizontal transmission

To investigate the possibility of co-speciation between the *w*Au-like *Wolbachia* strains infecting willistoni and saltans group flies, *w*Au-like$_{ws}$ strains, and their hosts, we inferred a phylogeny and compared it to the nuclear and mitochondrial phylogenies of the hosts (Baião et al. 2023). We found that the phylogeny of *w*Au-like$_{ws}$ strains was congruent with the nuclear phylogeny of the host species, except for the *Wolbachia* strain *w*Pau and its host *D. paulistorum*. Complete congruence was, however, found between the topologies of the two maternally transmitted genetic entities, *Wolbachia* and mitochondria. Hence, based on phylogenetic congruence, it seems plausible that a *w*Au-like strain was present in the ancestor of the willistoni and saltans groups at their divergence ca. 20 million years ago (Powell et al. 2003) and that this strain has co-diverged with the current hosts ever since.

However, phylogenetic congruence can be achieved in the absence of co-speciation, a phenomenon referred to as pseudo-co-speciation, especially common if host switching preferentially occurs between closely related species (De Vienne et al. 2007). Thus, to infer true co-speciation, temporal congruence between the two compared entities should ideally be observed. However, since no reliable estimate of the mutation rate exists for any *Wolbachia* strain, and the inferred divergence dates of host species are not confirmed by fossil data, dating the nodes in either of our trees will be highly unreliable. Nevertheless, to attempt to investigate temporal congruence, we calculated the synonymous substitution rate of *w*Au-like$_{ws}$ strains relative to that of their hosts and compared that to previously reported cases of *Wolbachia*-host co-speciation. We found the relative rates in our dataset to be quite constant between different *Wolbachia*-host pairs, with *D. paulistorum- D. willistoni* and their *Wolbachia* having the highest ratio around 3-7 times higher than the others. Even so, all rates were in general between one and two orders of magnitude lower than in previous co-speciation cases (Raychoudhury et al. 2008; Gerth and Bleidorn 2016; Conner et al. 2017). Thus, if all co-speciation cases were true, the *w*Au-like$_{ws}$ strains must either have evolved much slower than the other co-speciating *Wolbachia* strains or the willistoni and saltans group flies must have evolved much faster than the hosts in previous studies. However, this

comparison is not conclusive as we don't know if the earlier reported cases represent true

examples of co-speciation, since neither has proven temporal congruence. Additionally,

mutation and substitution rates as well as generation times can vary widely between different

organisms, even between very closely related bacterial and animal species.

Therefore, we also calculated mutation rates of *Wolbachia* based on the synonymous

substitution rates and the host divergence times. Using this method, we got mutation rates

between 10^{-10}–10^{-8} synonymous substitutions per synonymous site per year, which is lower

than in other bacteria (Gibson and Eyre-Walker 2019). Consequently, for our calculated

rates to match the generational mutation rates of bacteria (ranging between 10^{-10}–10^{-8}

synonymous substitutions per synonymous site per generation (Lynch et al. 2023)), all

Wolbachia strains*,* except the ones in *Nasonia,* must have a generation time of almost a

year. Since such a generation time is even longer than the hosts', our estimated mutation

rates are likely incorrect. There are indeed many potential sources of error for our mutation

rate estimates, for example, synonymous substitutions could be under strong selection, or

the recombination rate could be extremely high and eliminate any diversity between strains,

both of which could yield underestimated mutation rates. Additionally, the host divergence

times could be wrong, which, depending on if they are over- or under-estimated, could result

in either lower or higher mutation rates. At this point, we can't completely rule out any of

these factors, although selection seems unlikely given that the effective population size of

Wolbachia is probably low, and it is hard to understand how recombination between the

*w*Au-like$_{ws}$ strains could be so rampant in the absence of horizontal transmission. Even

though there is a good chance that the divergence times of the hosts are wrong, they would

have to be off by orders of magnitude, which is less likely. Alternatively, for co-speciation to

be believable, *Wolbachia* must have a very low mutation rate or a long generation time, or a

combination thereof. If not, neither the *w*Au-like$_{ws}$ strains nor the *Wolbachia* strains from

Nomada and *Nasonia* have co-speciated with their hosts, but may instead have been

horizontally transmitted into their hosts via introgression or alternative mechanisms, as

suggested in other similar systems (Turelli et al. 2018; Cooper et al. 2019).

It is unlikely that *Wolbachia* could have been recently transmitted between the

relatively distantly related species of the willistoni and saltans group via introgression, but full

congruence between mitochondria and *Wolbachia* suggests that introgressions have

facilitated at least some horizontal transmission. As previously reported, *D. paulistorum* has

two mitotypes, α and β, which were probably both introgressed from an unknown species

within the willistoni group (Baião et al. 2023). Hence, it is likely that *w*Pau was introgressed

into *D. paulistorum* from a willistoni group species together with one of the mitotypes,

potentially β since it is most common among *D. paulistorum* semispecies and all *w*Pau

strains in this study are from *D. paulistorum* lines carrying the β mitotype. As *w*Pau carries
complete and thus likely functional *cif* genes, the spread of *w*Pau and the replacement of the
ancestral *D. paulistorum* mitochondrion could potentially have been driven by CI. However,
to fully answer this question, data from additional *D. paulistorum* semispecies, especially the
ones carrying the α-mitotype, are needed.

Additionally, at least four horizontal transmission events of *w*Au-like *Wolbachia, w*Inn,
*w*Bor, *w*Inc, and *w*Au, into distantly related *Drosophila* species have likely occurred relatively
recently, as previously suggested (Ballard 2004; Miller and Riegler 2006; Sheeley and
McAllister 2009; Wallau et al. 2016), but through mechanisms other than introgression. We
can only speculate about how these *w*Au-like *Wolbachia* strains infected and spread in their
new hosts, but CI is not a likely mechanism for the spread since none of them carry what
appear to be functional *cif* genes. The well-studied strain *w*Au has long been known to not
induce CI in its native host, *D. simulans* (Hoffmann et al. 1996). However, it does provide
high antiviral protection in its host (Martinez et al., 2015), a potentially beneficial phenotype
that might have facilitated its spread. Nothing is currently known about the phenotypic effects
*w*Inc has on its host, but both *w*Inn and *w*Bor cause male-killing, which could have
contributed to their spread. Furthermore, *w*Inn was also seen to protect the host against
RNA viruses (Unckless and Jaenike 2012). It is also possible that *w*Au-like strains provide
yet unknown beneficial host effects.

In conclusion, more research is required to determine whether *Wolbachia* switches
hosts so frequently that pseudo-co-speciation is observed independently in multiple clades
and to identify mechanisms that facilitate the transfer and spread of *Wolbachia* infections.

434 The genome of obligate *Wolbachia* strain *w*Pau is expanded and has some
435 genomic features similar to recently host-associated symbionts

When free-living bacteria become host-associated, many of their genes become superfluous
in the new nutrient-rich, stable environment of the host and are hence lost or pseudogenized
due to lack of selection (McCutcheon and Moran 2012). In addition, when symbionts
transition from free-living to host-associated, they experience a reduction in effective
population size that increases the effect of genetic drift (Moran and Plague 2004,
Wernegreen 2015). Consequently, their genomes tend to accumulate slightly deleterious
mutations, including many IS element insertions (Bobay and Ochman 2017). However, over
time, often after becoming obligate for their hosts, symbiont genomes generally shrink
further in size and lose repeated sequences and pseudogenes (McCutcheon and Moran
2012, Bobay and Ochman 2017), leading to a conserved gene order. This stage of the
genome reduction process can, for example, be observed in obligate mutualistic *Wolbachia*

of nematodes that have small genomes (Werren et al. 2008, Dudzic et al. 2022) with few

repeats and mobile elements (Lefoulon et al. 2020) compared to most facultative *Wolbachia*

strains.

To our knowledge, *w*Pau is the only *Wolbachia* strain found in the willistoni and

saltans group of flies which is obligate for the host (Miller et al. 2010). Hence, *w*Pau is

expected to have a smaller genome than its close facultative relatives. Contrary to the

expectation, we found that *w*PauO11 has the largest genome among the *w*Au-like

Wolbachia strains with complete genomes. Genome size increase in *Wolbachia* is generally

associated with a higher proportion of prophage WO (Kampfraath et al. 2019; Lefoulon et al.

2020; Wolfe et al. 2021; Vancaester and Blaxter 2023), but in *w*Pau, we rather found a

reduction of prophage WO content. Instead, we attribute the increase in genome size in

*w*Pau to a still ongoing expansion of an IS4 element that makes up almost 10% of the

genome and has greatly affected the gene order. The extensive IS4 element proliferation,

high rate of genomic rearrangements and reduced coding percentage in the *w*Pau genome

are genomic features that are more common in recently host-associated facultative

symbionts than evolutionary persistent obligate symbionts. All of them, however, suggest a

large influence of genetic drift, potentially caused by a reduction in effective population size.

We can think of at least three possible reasons why the effective population size of *w*Pau

could have decreased compared to its close relatives.

Firstly, the horizontal transmission of *w*Pau might have reduced the effective

population size since only a few *Wolbachia* cells likely established the infection in *D.*

paulistorum initially. While we don't see an expansion of IS elements in other *w*Au-like

strains with complete genomes that have likely shifted host recently, i.e. *w*Inn and *w*Au, the

*w*Inn genome shares other features with the *w*Pau genome, including many rearrangements

and low coding density. Possibly, *w*Inn had more repeats but lost them over time due to

deletions. Although these similarities with *w*Inn might indicate that horizontal transmission

could initially cause a higher effect of genetic drift, we do not believe this to be the main

reason for the sustained high levels of genetic drift in *w*Pau.

Secondly, the transition from facultative to obligate host association could have

yielded a reduction in the effective population size of *w*Pau. Several similar genome features

can be found in the genome of the mutualistic *Wolbachia* strain *w*Cle infecting bedbugs

(Hosokawa et al. 2010), including an expansion of IS elements (with over 100 copies of IS5)

and reduced prophage WO content (Lefoulon et al. 2020). Potential IS element expansions

are also recorded in other obligate symbiont genomes, for example, the co-obligate *Serratia*

symbiotica strain Ct from the aphid *Cinara tujafilina* (Manzano-Marín and Latorre 2014),

where several IS element families have expanded (Renoz et al. 2021) and some species of

the genus *Sodalis* (Oakeson et al. 2014; Koga and Moran 2014; Ankrah et al. 2018;

Manzano-Marín et al. 2018; Garber et al. 2021). However, since no genome sequences of

closely related facultative strains are available in any of these cases, the IS element

expansion can't be linked to a transition to obligate host association.

Thirdly, a reduction in the effective population size of *w*Pau could be a consequence

of a reduction in the effective population size of its host, which might be linked to the fact that

D. paulistorum is a superspecies consisting of multiple fully or partly reproductively isolated

semispecies. Currently, no data about the effective population sizes of any species in the

willistoni or saltans groups exist, however, indirect evidence indicates a lower effective

population size in *D. paulistorum* compared to other willistoni and saltans group species.

Large eukaryotic genomes with a high proportion of transposable elements often occur in

species with low effective population sizes (Lynch and Conery 2003), due to the high

influence of genetic drift that allows mobile elements to accumulate. Since *D. paulistorum*

has the largest genome with the highest repetitive content of the sequenced willistoni group

species (Kim et al. 2021, Gebert et al. 2024), we might speculate that *D. paulistorum* has a

lower effective population size than other willistoni and saltans group species. If so, this

could explain not only the high rate of IS4 element insertions before the divergence of the

*w*Pau strain variants but also the continued high rate that seems to be ongoing in *w*Pau even

after its likely introgression and transition from facultative to obligate.

Prophage WO genes may be important for the obligate symbiosis of *w*Pau

The WO prophage is known to play an important role in facultative *Wolbachia* host

associations, given that it carries genes directly responsible for reproductive manipulation

(Beckmann et al. 2017; LePage et al. 2017; Perlmutter et al. 2019; Fricke and Lindsey

2024), infection titer and virulence (Chrostek and Teixeira 2015; Duarte et al. 2021). For the

obligate *w*Pau, we demonstrate loss of prophage WO genes compared to closely related

facultative strains. Specifically, *w*Pau has lost almost an entire copy (WOa) and some parts

of the other (WOb), suggesting a general lack of selection on WO-associated genes rather

than any specific WO prophage region. In contrast, nine of the eleven genes of the prophage

WO-associated Undecim cluster are instead duplicated in *w*PauO11 as well as the other

*w*Pau strain variants. Although the functions of the Undecim cluster genes are currently

unknown, they are thought to play a role in *Wolbachia*-host interactions, as they, among

other functions, encode proteins potentially involved in exopolysaccharide and/or

lipopolysaccharide biosynthesis (Bordenstein and Bordenstein 2022). Previous work has

also shown that the Undecim cluster has been horizontally transferred between several

unrelated insect endosymbionts and that the genes are constitutively highly expressed in

Wolbachia strain *w*Mel during the whole life cycle of *D. melanogaster* (Gutzwiller et al. 2015).

The fact that this prophage WO-associated set of genes is duplicated in *w*Pau, while many

other prophage WO genes are deleted, suggests that the region could encode proteins

potentially important for the obligate symbiosis between *w*Pau and its host. If true, prophage

WO might thus contribute genes for the persistence of a *Wolbachia* infection, even in the

absence of CI or other reproductive manipulations.

524 Conclusions

We conclude that the ancestor of Neotropical *Drosophila* in the willistoni and saltans groups

may have been infected with a *w*Au-like *Wolbachia* and co-speciated with their hosts, except

for the *w*Pau strain in *D. paulistorum*, which was likely introgression from another species

within the willistoni group together with one of the two mitotypes. However, such an

interpretation suggests that *Wolbachia* must have a very low mutation rate and/or a long

generation time. Further research is needed to determine if this is the case.

We also found that contrary to expectations, the genome of the obligate *w*Pau strain

was larger than its facultative relatives due to the ongoing proliferation of an IS4 element,

typically associated with genomes of recently host-associated symbionts. A reduced coding

percentage and more pseudogenes are also hallmarks of the early stages of genome

reduction, all of which are likely due to a reduction in the effective population size.

In general, the effective population size can be reduced multiple times after a

bacterium becomes host-associated. Thus, the genome reduction process might take place

in a punctuated manner, as episodes of effective population size reduction can temporarily

lead to an increased genome size via repeat expansion, and a decrease in the proportion of

coding content, only to be reversed by homologous recombination over such repeats over

time. Consequently, only when mobile elements and repeats are lost from a genome

altogether, can this cycle be broken.

Materials and Methods

DNA Extraction and Sequencing

Drosophila lines used in this study and the sequencing data produced from each of them are summarized in (Table S2). For *D. paulistorum* O11, total DNA for Oxford Nanopore (ONT) sequencing was extracted from many adult flies of both sexes using the MagAttract HMW DNA Kit (Qiagen) to obtain high molecular weight DNA. For Illumina sequencing, total DNA was obtained as described in Baiaõ et al 2023. For *Wolbachia* strains *w*WilP98, *w*ProPE2, *w*PauFG111, *w*PauPOA1 and *w*PauTP37, DNA samples used for PacBio sequencing were enriched for *Wolbachia* using a differential centrifugation and filtration procedure to obtain a *Wolbachia* cell pellet, which was then subjected to whole genome amplification using the Repli-g midi kit (Qiagen) as described in (Ellegaard et al., 2013). For *w*WilP98 and *w*ProPE2, Illumina sequences enriched for *Wolbachia* were obtained using the same protocol.

The ONT library was produced and sequenced using an R9.4 PromethION flow cell at the Uppsala Genome Center, Uppsala, Sweden. Basecalling was performed with Guppy 4.3.4 and the HAC model. Illumina TruSeq libraries were produced and sequenced at the Uppsala SNP and Seq platform on an Illumina 2500 HiSeq machine generating 2x125 bp reads. The Illumina reads were quality filtered and trimmed using Trimmomatic-0.30 with parameters –phred33 ILLUMINACLIP:illumina_adapter.fasta:2:40:15 LEADING:3 TRAILING:3 SLIDINGWINDOW:4:15 MINLEN:95 (Bolger et al., 2014) and error-corrected using SPAdes-3.5.0 –sc –only-errorcorrection (Bankevich et al., 2012). For PacBio, 5 kb SMRTbell libraries were created. Each library was run with P6-C4 chemistry in one SMRT cell on an RSII PacBio instrument at Uppsala Genome Center.

Genome Assemblies

Wolbachia genomes were assembled using several types of data and pipelines. For *w*PauO11, reads from total DNA sequencing (i.e. including the *Drosophila* host) were filtered using Filt-long v0.2.1 with quality priority and the parameters "--target_bases 12500000000 --min_length 1000 --mean_q_weight 10" and assembled with Nextdenovo v2.5.0 using correction options: read_cutoff = 1k, genome_size = 250m, sort_options = -m 7g -t 2, minimap2_options_raw = -t 3, pa_correction = 17, correction_options = -p 2 and assemble options: minimap2_options_cns = -t 3, nextgraph_options = -a 1. The produced assembly contained only one *Wolbachia* contig that was subsequently polished with Illumina reads Pilon v1.24 (Walker et al. 2014) with the --fix indels parameter and circularized. The genomes of the two strains *w*WilP98 and *w*ProPE2 were assembled using Illumina and PacBio reads from amplified DNA enriched for *Wolbachia*. The Illumina reads were assembled using Spades v3.15.5 (Bankevich et al. 2012) with kmer sizes 97,103,107,113

and the PacBio reads were used for scaffolding with the --pacbio parameter. Illumina and

PacBio reads were mapped against the assemblies using BWA-mem (Li and Durbin 2009)

with default parameters for Illumina reads and using PacBio settings for PacBio reads to

check the correctness of the genome assembly. Overlapping contigs were merged in

Consed (Gordon et al. 1998). For all other assemblies, BWA-mem v0.7.17-r1198-dirty with

default parameters was used to map Illumina reads on the *w*PauO11, *w*ProPE and *w*WilP98

genomes and a set of concatenated *Wolbachia* genomes representing the *Wolbachia*

diversity (81 genomes from supergroups A, B, C, D, E, F, L and M, Table S12). Mapped

reads were filtered using samtools v.1.13 (Danecek et al. 2021) view with the -f 1 -F 12

parameters to include only pairs where both were mapped. Filtered reads were then

subsampled to approximately 100X with seqtk v1.3-r117-dirty sample -s 100 before

assembly. The reads were then assembled using Spades v3.15.5 with kmer sizes

97,103,107,113, and the *w*PauFG111, *w*PauPOA1 and *w*PauTP37 assemblies were

scaffolded with PacBio reads using the flag -pacbio.

The Illumina assemblies were then filtered by removing contigs that deviated from

the mean sample distribution in kmer coverage and GC content. The kmer coverage of the

contigs was determined from the fasta headers in the output from Spades. In cases where

kmer coverage had a bimodal distribution, the minimum between the two peaks was

identified using the quantmod package in R (Ryan and Ulrich 2022) and used as threshold.

Contigs with kmer coverage below this threshold were blasted against the assembled

contigs and were removed if the sequence existed. This procedure ensured that redundant

low coverage contigs were removed. Remaining low coverage contigs were blasted against

additional *Wolbachia* genomes and proteins. Contigs with a blast alignment length longer

than 100 aa and percent identity over 60%, or blast alignment length longer than 200

nucleotides and percent identity over 70%, were kept. Any contig with an extreme GC value

(outside the 99% interval of a normal distribution of GC) was removed. Finally, all remaining

contigs were searched with Kraken against a custom database (containing a selection of

archaea, bacteria, plasmids, viruses, human, *D. melanogaster* and *D. willistoni* sequences)

and the contigs with a match to *Drosophila* were removed. All assemblies were polished

once using Pilon with the Illumina reads used for assembly.

Genome Annotation

For the complete genomes, *w*PauO11, *w*ProPE2 and *w*WilP98, Prokka v1.14.6 (Seemann

2014) with the parameters –genus Wolbachia –compliant –rfam --addgenes --usegenus was

used for annotation. Next, Pseudofinder v1.1.0 (Syberg-Olsen et al. 2022) with parameters -

ce and -db with a database containing all *Wolbachia* proteins from NCBI (DATE), was used

to mark potential pseudogenes. Additionally, hmmsearch as implemented in pfam_scan.pl was used for domain prediction with the PFAM database (Bateman et al. 2002). The Illumina reads used for the assemblies were mapped back to each genome with BWA-mem to verify frameshifts and nonsense mutations. All marked pseudogenes were inspected manually and annotated as pseudogenes if the start or stop codons were missing, they contained nonsense or frameshift mutations or were truncated by, for example, an IS element insertions.

The draft assemblies were annotated the same way, except that 1) Pseudofinder was run with length_pseudo set at 0.8 and the protein database only included *Wolbachia* genomes *w*Au, *w*Ha, *w*Ma, *w*No, *w*San, *w*Tei, *w*Yak, *w*PauO11, *w*ProPE2 and *w*WilP98 that were manually annotated either in this or previous studies (Ellegaard et al. 2013; Baião et al. 2021), 2) only pseudogene calls in CDS sequences were inspected manually, 3) pseudogenes that we identified in the complete genomes (*w*PauO11, *w*ProPE2 and *w*WilP98), but were not found by pseudofinder in the draft genomes, were transferred to the draft genome annotation if they matched with high similarity and extensive coverage. For identifying repeated sequences, nucmer from the MUMmer 4.0 package (Marçais et al. 2018) was used with flags --maxmatch --nosimplify to align each genome to itself. The positions of non-overlapping repeat bases were then extracted using a perl script. ISEScan v 1.7.2.3 (Xie and Tang 2017) was used to identify IS elements. Bar plots of the percentage of repeats and IS elements in each genome were made in R (R Core Team 2018; Wickham et al. 2019). Phage WO regions were annotated by blastp searches to *Wolbachia* genomes (*w*Mel, *w*Au, *w*Ha, *w*Ma, *w*No, *w*San, *w*Tei, *w*Yak) with previously identified phage WO regions. All genomic regions that contained similarities to known phage WO genes were then manually inspected, and borders were determined based on Bordenstein and Bordenstein 2022.

MLST phylogeny of *Wolbachia* genomes

Genomes that contained the keyword "Wolbachia" in the taxon name, were downloaded from NCBI using the command line tool "datasets" (https://github.com/ncbi/datasets) on May the 6[th] 2022. tblastn of the Blast v2.7.1+ suite (Camacho et al. 2009) with the parameters --max_hsps 1 --evalue 1e-10 was used to search for the five *w*Mel MLST proteins and Wolbachia surface protein (Wsp) (GenBank accession numbers AAS13898.1, AAS14036.1, AAS14201.1, AAS14415.1, AAS14719.1, AAS14884.1) in 1504 downloaded genomes. Genomes with blast hits for all six genes were kept for phylogenetic inference. Nucleotide sequences of the identified genes were extracted from each genome using samtools faidx, and translated, aligned with MAFFT v6.850b using the linsi algorithm (Katoh and Toh 2008),

pruned to remove gap sites present in more than 50% of aligned sequences and then

backtranslated to nucleotides again. A phylogeny of the six genes from 1262 genomes was

then constructed using RaxML v8.2.9 (Stamatakis 2014) with parameters –f a –m

GTRGAMMA –x 12345 –p 12345 -# 100. The resulting phylogenetic tree as visualized in

Figure 1 and all other figures presenting phylogenetic trees (Fig. 2, Fig. 4B, Fig. S1, Fig. S2,

Fig S3, Fig. S5 and Fig. S6) were made using FigTree v1.4.4 (Rambaut 2009).

Protein clustering and strain phylogeny

At the time of downloading *Wolbachia* genomes, the ones from Vancaester and Blaxter 2023

were not yet available. Hence, to include them in our analysis, we downloaded the genomes

classified as supergroup A by the authors. We then annotated the genomes from all strains

classified as supergroup A *Wolbachia* in the MLST phylogeny and by Vancaester and

Blaxter 2023 as described above for our draft assemblies, used their proteomes for

clustering and inferred a phylogeny from single-copy genes. Clustering was done with

Orthofinder v2.5.4 (Emms and Kelly 2019) with –I 3 and filtered all-against-all blastp results

as input. Blastp hits between two matching proteins were filtered so that the size of one

should not be smaller than 60% of the size of the other, the e-value should be lower than 10^{-5}

and the alignment length should be at least 80% of the size of the smaller protein. The

nucleotide sequences of genes from orthogroups with single-copy proteins were extracted,

translated, aligned with mafft-linsi, pruned to remove gap sites present in more than 50% of

the aligned sequences and backtranslated to obtain codon-based nucleotide alignments.

To ensure that the genes used in the phylogeny did not show signs of recombination,

we ran Phipack (Bruen et al. 2006) with the -p 100000, -o, -w 50 on all alignments. Phipack

runs three tests to detect recombination, the Phi test, the max χ2 test and the NSS test and

at least two of the three recombination tests needed to have a p-value lower than 0.05, for a

gene to be considered as recombined. Single-copy gene alignments without recombination

were then concatenated with geneStitcher.py from Utensils

(https://github.com/ballesterus/Utensils/blob/master/geneStitcher.py).

IQ-TREE v2.2.0 (Minh et al. 2020) was then used for phylogenetic inference of the

species tree using the concatenated alignments of the single-copy genes, a partition per

gene, and the parameters -m MFP, -B 1000 -T 20 --seed 12345. Individual gene trees were

inferred using the same parameters. The phylogenetic trees were visualized using FigTree

v1.4.4. To test if the data fit better with the placement of *w*Pau according to the host nuclear

phylogeny rather than our strain phylogeny, we followed Shen et al. 2017 and used IQ-TREE

v1.6.12 with the same parameters as in the strain phylogeny together with the -z flag to add

the two topologies (shown in Figure S3) in Newick format and the -wsl and -wpl flags to get

the likelihood for each site and gene. The differences in the gene likelihoods between the

two topologies were calculated and plotted in R (R Core Team 2018; Wickham et al. 2019).

To measure the gene (gCFs) and site (sCFs) concordance factors, we used the same

pruned concatenated phylogeny as in the log-likelihood differences and we produced pruned

gene trees including the same strains with preserved branch lengths using a Python script

and the ete3 library (Huerta-Cepas et al. 2016). To calculate the gene and site concordance

factors, the pruned species tree was then provided to IQ-TREE with the flag -t, the fasta

alignments with -s, the partition file with -p, the gene trees were provided with --gcf and --scf

100 was used.

Substitution rates

693 Pairwise synonymous substitution rates were calculated with codeml from the PAML v4.8

694 package (Yang 2007) using alignments of the 620 single-copy *Wolbachia* genes from the

695 ortholog clustering, as well as the alignments of 692 BUSCO genes and 13 mitochondrial

696 genes from *Drosophila* (the same as in Baião et al 2013) generated the same way as the

697 *Wolbachia* alignments (see above).

Gene order phylogeny

699 Syntenic blocks between the complete genomes listed in Table 1 were obtained using

700 SynChro_linux (January 2015) (Drillon et al. 2014) with a delta value of 2. The output from

701 SynChro was then used for phylogenetic reconstruction with PhyChro (downloaded July

702 2023 from http://www.lcqb.upmc.fr/phychro2/) (Drillon et al. 2020).

IS4 element and Undecim cluster copy numbers in draft *w*Pau genomes

To estimate the number of IS4 elements and Undecim cluster copies in the draft *w*Pau

genomes, we first masked all copies of the IS4 element and one copy of the Undecim cluster

in the *w*PauO11 genome. To do that, we created an IS4 element consensus sequence by

blasting (blastn) one IS4 element from *w*PauO11 against the *w*PauO11 genome, extracting

the sequences for all matches, aligning them with MAFFT v7.490 with flags "--maxiterate

1000" and "--localpair" and building a consensus with Jalview v2.11.4.0 (Waterhouse et al.

2009). The consensus IS4 element sequence was then blasted against the *w*PauO11

genome using blastn and all hits were masked. The Undecim cluster copies were first

identified by homology to the Undecim cluster genes in the *w*Mel genome. One of the

Undecim cluster copies, at positions 898.960-913.092 in *w*PauO11, was then manually

masked. The Illumina reads used to assemble the draft *w*Pau genomes were then mapped

to the masked *w*PauO11 genome and the IS4 element consensus sequence using BWA-

mem v0.7.17-r1198-dirty. Repeated areas were identified from the annotation (see above)

and non-repeated regions of the *w*PauO11 genome were identified using "bedtools

complement" and the repeated regions as input. Coverage over the IS4 element consensus

sequence and the unmasked Undecim cluster copy was calculated with "samtools coverage"

and the coverage of the non-repeated regions of the genome was calculated by getting the

average per-base coverage in these regions by using "samtools depth". The copy numbers

of the IS4 element and the Undecim cluster were calculated by dividing the average

coverage over these regions by the average coverage over the non-repeated regions of the

genome.

SNP phylogeny of *w*Pau genomes

To get a more resolved relationship between the *w*Pau strain variants, we called SNPs for all

*w*Pau strain variants from the Illumina reads mapped to the masked *w*PauO11 genome (see

above). Reads with a MQ<1 were filtered out and bcftools was used to call SNPs using the

command "bcftools mpileup --threads 6 -Ou -d 1000 -f $genome -b bam.file.list.txt | bcftools

call --threads 6 -cv --ploidy 1 -Ob -o variants_cflag.bcf". To filter low-quality variants, the

command "bcftools view --types snps -i 'QUAL>30 & INFO/DP>10 & MQ>30'" was used.

The variants located in non-repeated areas of the *w*PauO11 genome were extracted using

"bcftools view" with the coordinates identified as described above and "bcftools consensus"

was used to create pseudo-chromosomes for each strain. The sequence for each pseudo-

chromosome was put into a multi-fasta file and used as input for IQ-TREE v2.2.0 using the

same parameters as in the *Wolbachia* strain phylogeny. The phylogeny was visualized using

FigTree v1.4.4.

IS4 element insertion dynamics

To detect differences in IS4 insertions between the *w*Pau genomes, we identified improperly

mapped read pairs in the Illumina reads that were mapped to the masked *w*PauO11 genome

(see above). If an IS4 element is present in the genomes with mapped reads, the insert size

of paired reads in this region is 0, as one read of the pair would map on the masked

Wolbachia genome and the other read would map on the consensus IS4 sequence. To get

such improperly paired reads, we filtered with samtools view using parameters -F2, MQ>0

and -e "tlen==0". Reads that mapped within 750 bases upstream of the start or 750 bases

downstream of the end of an IS4 element in wPauO11 (as specified in a provided bed file)

are from an IS4 element shared with *w*PauO11. Reads mapping at other places are

associated with IS4 insertions not present in *w*PauO11, and they were filtered by using

samtools view with the -U -L flag and the same bed file as above. If an IS4 element is

present in *w*PauO11 but absent in the genome with mapped reads, the reads from a pair are

further apart than expected, specifically the insert size would appear to be larger than the ca.

1500 bases of the IS4 element, and both reads of the pair should map on the masked

Wolbachia genome. To get such improperly paired reads, reads mapping 750 bases

upstream and 750 bases downstream of IS4 elements (as above) and that had an insert size

larger than 1500 bases but shorter than 5000 bases were filtered using samtools view -e

"(tlen>1500 && tlen<5000) || (tlen>-5000 && tlen < -1500)". The filtered read alignments of

improper pairs were visualized in IGV (v2.16.0) to identify unique IS4 element insertions

across the different *w*Pau strain variants. The *w*Pau whole genome SNP phylogeny was

then used to infer where and when IS4 copies were inserted in the *w*Pau strain variants.

Because complete deletions of IS4 elements occur rarely, differences were explained by

insertions, rather than deletions.

Additionally, we inferred the number of IS4 element copies present at each node of

the tree using complete *Wolbachia* genomes. We identified the shared IS4 elements across

the *w*Au-like strains, *w*Mel and *w*Cin2USA1 by blasting (blastn v2.7.1+) genomes to each

other and visualizing the regions flanking IS4 elements, identified by ISEscan, with Artemis

comparison tool (ACT) (Carver et al. 2005). An IS4 element was considered absent from a

genome if a region had the same gene order as an IS4 element-containing region of another

genome but didn't contain an IS4 element or if an IS4 element-containing region in other

genomes shared no synteny with the target genome.

Gene content variation between genomes *w*Au-like *Wolbachia*

To compare gene content across the different *Wolbachia* genomes, we used the gene count

output of Orthofinder to produce a hierarchical clustering of the genomes based on gene

presence/absence. First, we excluded all single-copy genes and then we assigned the count

of multi-copy genes to 1 so that we cluster based on presence and not copy-number. Then,

we used the "agnes" function for agglomerative clustering with the "average" method from

the cluster v2.1.4 package in R (Maechler et al. 2021).

Additionally, the variability in gene content between the wAu-like$_{ws}$ genomes was

investigated by identifying orthogroups where only one or several of the lineages, *w*Pau,

*w*Tro/*w*Wil and *w*Pro/*w*Spt, were present and one or several lineages were absent (Table 2,

Table S9 and Table S10). Unless otherwise stated, presence was counted if a gene was

present in the complete genome of the lineage (*w*PauO11, *w*WilP98 and *w*ProPE2) and at

least one draft genome of the same lineage.

All figures presenting genome and gene comparisons (Fig. 4A, Fig. 7, Fig. S7 and

Fig. S8) were made with genoPlotR (Guy et al. 2010), after using blastn to search each

genome against every other genome.

Prophage WO classification and gene module assignment

To investigate the prophage WO content in detail, we identified prophage WO regions in the
complete *w*Au-like genomes, *w*PauO11, *w*ProPE2, *w*WilP98, *w*Inn, *w*Au, and in *w*Mel,
*w*Cin2USA1. All proteins of prophage WO origin were re-clustered to orthogroups with
OrthoFinder v2.5.4 with -I=1 and custom blastp (as above), which put most phage WO
paralogues into single orthogroups. We manually validated the orthogroups based on the
functional annotation and gene order in complete genomes (Table S13). The orthogroups
containing proteins with the same functional annotation that were syntenic were manually
merged. For orthogroup, we inferred a phylogeny using IQ-TREE v2.2.0 as described for
single-copy genes. Genes were assigned to WOa and WOb based on whether they formed
a clade with either WOMelA or WOMelB genes and/or were syntenic with WOMelA or
WOMelB. For orthogroups where a WOMelB gene was present, we followed the
classification of Bordenstein and Bordenstein 2022. Otherwise, we looked at what module
the neighboring genes belonged to and assigned the orthogroup to that module. If an
orthogroup had no functional annotation and could not be assigned to a module using the
above method, it was assigned to the EAM module if it was flanked by genes in a structural
module on one side and an EAM gene on the other side.

Acknowledgements

We thank Mercè Montoliu Nerín for assembling the *Drosophila paulistorum* O11 genome
from which the *w*PauO11 genome contig was extracted and for helpful discussions
throughout the project, Tom Martin for performing the DNA extraction for Nanopore
sequencing of *D. paulistorum* O11, Lina Juzokaite for *Wolbachia*-enriched DNA extractions
for PacBio sequencing, and Illumina sequencing. We also thank Athanasios Alexiou for
assembling the first version of the *w*WilP98 genome. Sequencing was performed by the
SNP&SEQ Technology Platform and Uppsala Genome Center (UGC) in Uppsala, Sweden.
The facilities are part of the National Genomics Infrastructure (NGI) Sweden and Science for
Life Laboratory. Some of the data handling was enabled by resources provided by the
Swedish National Infrastructure for Computing (SNIC) at UPPMAX. SNP&SEQ, UGC and
Uppmax are supported by the Swedish Research Council and SNP&SEQ are supported by
the Knut and Alice Wallenberg Foundation.

Author contribution

817 LK and WJM conceived and designed the study. KP and LK performed assemblies. KP and
818 LK performed annotations. KP and LK performed comparative analyses. KP performed phy-
819 logenetic analyses. LK and KP wrote the paper with contributions from WJM. All authors
820 read and approved the manuscript.

Data availability

The data underlying this study will be made available at publication.

Figure legends

Figure 1. Phylogeny of *Wolbachia* genomes with high similarity to *w*Au-like$_{ws}$ strains. The branches with *Wolbachia* strains associated with *Drosophila* species from the willistoni and saltans groups, i.e., the *w*Au-like$_{ws}$ *Wolbachia* strains, are coloured. The clade referred to as the *w*Au-like is indicated with an arrow. The tree was inferred from an alignment of 620 single-copy orthologs with IQ-TREE. Support values from 1000 ultrafast bootstrap replicates are displayed on the nodes of the tree. Only bootstrap values greater than 60 are shown.

Figure 2. Phylogenetic comparison between *w*Au-like *Wolbachia* and *Drosophila* of the willistoni and saltans groups. A) Maximum Likelihood Phylogenies of *w*Au-like *Wolbachia* on the left and 692 nuclear-encoded proteins of the host on the right (from Baião et al. 2023). The *w*Au-like *Wolbachia* phylogeny is a subset of the tree in Figure 1. B) Maximum Likelihood Phylogenies of *w*Au-like *Wolbachia* on the left (same as in A) and the whole mitochondrial genome of the neotropical *Drosophila* host species on the right (from Baiaõ et al. 2023).

Figure 3. Repeats and IS-elements in complete genomes of *w*Au-like strains. A) Percentage of repeats in each genome. B) Percentage of each IS family in the genomes, as identified by ISEscan.

Figure 4. Genomic rearrangements in *w*Au-like strains. A) Whole genome alignments of complete *Wolbachia w*Au-like genomes and outgroup. For each genome, genes are visualized with blue and pseudogenes with green rectangles on either the forward or reverse strand (above and below the midline respectively). Similarities between genomes are visualized as red and blue ribbons, with red representing forward and blue reverse hits. B) Gene order phylogeny of the same genomes as in A. Branch lengths written on each branch represent the estimated number of genomic rearrangements.

Figure 5. Gene content clustering of *Wolbachia w*Au-like genomes. The cladogram on the left shows how the *Wolbachia* strains cluster based on the presence and absence of orthogroups in each genome, which is depicted in the plot to the right.

Figure 6. Prophage WO genes in complete *w*Au-like genomes and outgroup. Number of genes per module for the two prophage WO variants identified in *w*Au-like and related genomes. WOa genes were found in the same clade as genes from WO-A in *w*Mel and WOb genes were found in the same clade as genes from WO-B in *w*Mel when inferring single gene phylogenies.

855 **Figure 7. Duplication of the phage WO-associated genes in the Undecim cluster in**

856 ***w*PauO11.** Similarities between genomes are visualized with grey bands and the duplication

857 is marked in yellow. Genes in orange are part of the Undecim cluster. Mobile elements are

858 dark grey and pseudogenes green, with the border colour indicating Undecim cluster or

859 mobile element.

Tables

Table 1. Genomic features in complete *w*Au-like and outgroup genomes.

	*w*PauO11[1]	*w*WilP98[1]	*w*ProPE2[1]	*w*Au[2]	*w*Inn[3]	*w*Mel[4]	*w*Cin2[5]
Genome size (bp)	1,365,198	1,268,527	1,339,466	1,268,455	1,290,587	1,267,782	1,538,351
GC (%)	34.8	35.2	35.2	35.2	35.3	35.2	35.1
CDSs	1000	1016	1033	996	1073	1011	1473
% Coding	73.8	77.5	76.3	75.7	70.1	76.0	86.2
Pseudogenes	177	123	130	122	227	111	216
Repeats (%)	22.9	11.4	15.8	11.0	16.8	9.8	13.6
Phage WO (%)	8.8	10.8	12.6	10.6	9.0	9.5	24.3
IS elements (%)	15.3	7.5	9.4	7.6	9.9	7.0	7.6

[1]*Wolbachia* genomes from Neotropical *Drosophila* sequenced in this study, [2] Baião GC et al., 2021, [3] Hill T et al., 2022, [4] Wu M et al., 2004, [5] Wolfe TM et al., 2021.

Table 2. Gene content variation between *w*Au-like$_{ws}$ *Wolbachia*. The numbers represent the presence or absence of an orthogroup in one lineage compared to the other two lineages in the table.

	*w*Pau	*w*Wil/*w*Tro	*w*Pro/*w*Spt
Absent (total)	**66**	**18**	**12**
Phage WO	39	6	-
Ankyrin	2	1	2
Hypothetical	9	3	3
Dozen island	-	4	-
Present (total)	**6**	**11**	**16**
Phage WO	-	5	6
Ankyrin	1	-	1
Hypothetical	4	6	2
Dozen island	-	-	6

References

Baião GC, Janice J, Galinou M, Klasson L. 2021. Comparative Genomics Reveals Factors Associated with Phenotypic Expression of Wolbachia. *Genome Biology and Evolution* 13:evab111.

Baião GC, Schneider DI, Miller WJ, Klasson L. 2019. The effect of Wolbachia on gene expression in Drosophila paulistorum and its implications for symbiont-induced host speciation. *BMC Genomics* 20:465.

Baião GC, Schneider DI, Miller WJ, Klasson L. 2023. Multiple introgressions shape mitochondrial evolutionary history in Drosophila paulistorum and the Drosophila willistoni group. *Molecular Phylogenetics and Evolution* 180:107683.

Bakovic V, Schebeck M, Stauffer C, Schuler H. 2020. Wolbachia-Mitochondrial DNA Associations in Transitional Populations of Rhagoletis cerasi. *Insects* 11:675.

Baldo L, Dunning Hotopp JC, Jolley KA, Bordenstein SR, Biber SA, Choudhury RR, Hayashi C, Maiden MCJ, Tettelin H, Werren JH. 2006. Multilocus Sequence Typing System for the Endosymbiont Wolbachia pipientis. *Applied and Environmental Microbiology* 72:7098–7110.

Ballard JWO. 2004. Sequential Evolution of a Symbiont Inferred From the Host: Wolbachia and Drosophila simulans. *Molecular Biology and Evolution* 21:428–442.

Balvín O, Roth S, Talbot B, Reinhardt K. 2018. Co-speciation in bedbug Wolbachia parallel the pattern in nematode hosts. *Sci Rep* 8:8797.

Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, et al. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477.

Bateman A, Birney E, Cerruti L, Durbin R, Etwiller L, Eddy SR, Griffiths-Jones S, Howe KL, Marshall M, Sonnhammer ELL. 2002. The Pfam protein families database. *Nucleic Acids Res* 30:276–280.

Beckmann JF, Ronau JA, Hochstrasser M. 2017. A Wolbachia Deubiquitylating Enzyme Induces Cytoplasmic Incompatibility. *Nat Microbiol* 2:17007.

Blasco-Costa I, Hayward A, Poulin R, Balbuena JA. 2021. Next-generation cophylogeny: unravelling eco-evolutionary processes. Trends in Ecology & Evolution 36:907–918.

Bobay L-M and Ochman H (2017) The Evolution of Bacterial Genome Architecture. Front. Genet. 8:72. doi: 10.3389/fgene.2017.00072

Bordenstein Sarah R., Bordenstein Seth R. 2016. Eukaryotic association module in phage WO genomes from Wolbachia. *Nat Commun* 7:13155.

Bordenstein Sarah R., Bordenstein Seth R. 2022. Widespread phages of
endosymbionts: Phage WO genomics and the proposed taxonomic classification of
Symbioviridae.Matic I, editor. *PLoS Genet* 18:e1010227.

Bourtzis K, Nirgianaki A, Markakis G, Savakis C. 1996. Wolbachia infection and
cytoplasmic incompatibility in Drosophila species. *Genetics* 144:1063–1073.

Bruen TC, Philippe H, Bryant D. 2006. A Simple and Robust Statistical Test for
Detecting the Presence of Recombination. *Genetics* 172:2665–2681.

Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL.
2009. BLAST+: architecture and applications. BMC Bioinformatics 10:421.

Cariou M, Duret L, Charlat S. 2017. The global impact of Wolbachia on mitochondrial
diversity and evolution. *Journal of Evolutionary Biology* 30:2204–2210.

Charlat S, Duplouy A, Hornett EA, Dyson EA, Davies N, Roderick GK, Wedell N,
Hurst GD. 2009. The joint evolutionary histories of Wolbachia and mitochondria in
Hypolimnas bolina. *BMC Evolutionary Biology* 9:64.

Carver TJ, Rutherford KM, Berriman M, Rajandream M-A, Barrell BG, Parkhill J.
2005. ACT: the Artemis Comparison Tool. Bioinformatics 21:3422–3423.

Charleston MA. 2016. Cospeciation. In: Encyclopedia of Evolutionary Biology.
Elsevier. p. 381–386.

Chrostek E, Marialva MSP, Yamada R, O'Neill SL, Teixeira L. 2014. High Anti-Viral
Protection without Immune Upregulation after Interspecies Wolbachia Transfer.
PLOS ONE 9:e99025.

Chrostek E, Teixeira L. 2015. Mutualism Breakdown by Amplification of Wolbachia
Genes.Malik HS, editor. *PLoS Biol* 13:e1002065.

Conner WR, Blaxter ML, Anfora G, Ometto L, Rota-Stabelli O, Turelli M. 2017.
Genome comparisons indicate recent transfer of Ri-like Wolbachia between sister
species Drosophila suzukii and D. subpulchrella. Ecology and Evolution 7:9391–
9404.

Cooper BS, Vanderpool D, Conner WR, Matute DR, Turelli M. 2019. Wolbachia
Acquisition by Drosophila yakuba-Clade Hosts and Transfer of Incompatibility Loci
Between Distantly Related Wolbachia. Genetics 212:1399–1419.

Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A,
Keane T, McCarthy SA, Davies RM, et al. 2021. Twelve years of SAMtools and
BCFtools. *GigaScience* 10:giab008.

De Vienne DM, Giraud T, Shykoff JA. 2007. When can host shifts produce congruent
host and parasite phylogenies? A simulation approach. Journal of Evolutionary
Biology 20:1428–1438.

Drillon G, Carbone A, Fischer G. 2014. SynChro: A Fast and Easy Tool to
Reconstruct and Visualize Synteny Blocks along Eukaryotic Chromosomes. *PLOS*
ONE 9:e92621.

Drillon G, Champeimont R, Oteri F, Fischer G, Carbone A. 2020. Phylogenetic
Reconstruction Based on Synteny Block and Gene Adjacencies. *Mol Biol Evol*
37:2747–2762.

Duarte EH, Carvalho A, López-Madrigal S, Costa J, Teixeira L. 2021. Forward
genetics in Wolbachia: Regulation of Wolbachia proliferation by the amplification and
deletion of an addictive genomic island. *PLOS Genetics* 17:e1009612.

Dudzic JP, Curtis CI, Gowen BE, Perlman SJ. 2022. A highly divergent *Wolbachia*
with a tiny genome in an insect-parasitic tylenchid nematode. *Proc. R. Soc. B.*
289:20221518.

Ellegaard KM, Klasson L, Näslund K, Bourtzis K, Andersson SGE. 2013.
Comparative Genomics of Wolbachia and the Bacterial Species Concept.Matic I,
editor. *PLoS Genet* 9:e1003381.

Emms DM, Kelly S. 2019. OrthoFinder: phylogenetic orthology inference for
comparative genomics. *Genome Biology* 20:238.

Fisher RM, Henry LM, Cornwallis CK, Kiers ET, West SA. 2017. The evolution of
host-symbiont dependence. *Nat Commun* 8:15973.

Frank AC, Amiri H, Andersson SGE. 2002. Genome deterioration: loss of repeated
sequences and accumulation of junk DNA. *Genetica*.

Fricke LC, Lindsey ARI. 2024. Identification of Parthenogenesis-Inducing Effector
Proteins in Wolbachia. Genome Biology and Evolution 16:evae036.

Gavotte L, Henri H, Stouthamer R, Charif D, Charlat S, Boulétreau M, Vavre F. 2007.
A Survey of the Bacteriophage WO in the Endosymbiotic Bacteria Wolbachia.
Molecular Biology and Evolution 24:427–435.

Gerth M, Bleidorn C. 2016. Comparative genomics provides a timeframe for
Wolbachia evolution and exposes a recent biotin synthesis operon transfer. *Nat*
Microbiol 2:16241.

Gordon D, Abajian C, Green P. 1998. Consed: a graphical tool for sequence
finishing. *Genome Res* 8:195–202.

Gutzwiller F, Carmo CR, Miller DE, Rice DW, Newton ILG, Hawley RS, Teixeira L,
Bergman CM. 2015. Dynamics of Wolbachia pipientis Gene Expression Across the
Drosophila melanogaster Life Cycle. G3 (Bethesda) 5:2843–2856.

Guy L, Roat Kultima J, Andersson SGE. 2010. genoPlotR: comparative gene and
genome visualization in R. Bioinformatics 26:2334–2335.

Hill T, Unckless RL, Perlmutter JI. 2022. Positive Selection and Horizontal Gene
Transfer in the Genome of a Male-Killing Wolbachia. Molecular Biology and
Evolution 39:msab303.

Hosokawa T, Koga R, Kikuchi Y, Meng X-Y, Fukatsu T. 2010. Wolbachia as a
bacteriocyte-associated nutritional mutualist. Proceedings of the National Academy
of Sciences 107:769–774.

Huerta-Cepas J, Serra F, Bork P. 2016. ETE 3: Reconstruction, Analysis, and
Visualization of Phylogenomic Data. Molecular Biology and Evolution 33:1635–1638.

Ilinsky Y. 2013. Coevolution of Drosophila melanogaster mtDNA and Wolbachia
Genotypes. *PLOS ONE* 8:e54373.

Kampfraath AA, Klasson L, Anvar SY, Vossen RHAM, Roelofs D, Kraaijeveld K,
Ellers J. 2019. Genome expansion of an obligate parthenogenesis-associated
Wolbachia poses an exception to the symbiont reduction model. *BMC Genomics*
20:106.

Katoh K, Toh H. 2008. Recent developments in the MAFFT multiple sequence
alignment program. *Briefings in Bioinformatics* 9:286–298.

Kaur R, Shropshire JD, Cross KL, Leigh B, Mansueto AJ, Stewart V, Bordenstein
Sarah R., Bordenstein Seth R. 2021. Living in the endosymbiotic world of Wolbachia:
A centennial review. *Cell Host & Microbe* 29:879–893.

Kim BY, Wang JR, Miller DE, Barmina O, Delaney E, Thompson A, Comeault AA,
Peede D, D'Agostino ER, Pelaez J, et al. 2021. Highly contiguous assemblies of 101
drosophilid genomes.Coop G, Wittkopp PJ, Sackton TB, editors. eLife 10:e66405.

Lefoulon E, Bain O, Makepeace BL, d'Haese C, Uni S, Martin C, Gavotte L. 2016.
Breakdown of coevolution between symbiotic bacteria Wolbachia and their filarial
hosts. PeerJ 4:e1840.

Lefoulon E, Clark T, Guerrero R, Cañizales I, Cardenas-Calligros JM, Junker K,
Vallarino-Lhermitte N, Makepeace BL, Darby AC, Foster JM, et al. 2020. Diminutive,
degraded but dissimilar: Wolbachia genomes from filarial nematodes do not conform
to a single paradigm. *Microb Genom* 6:mgen000487.

LePage DP, Metcalf JA, Bordenstein Sarah R., On J, Perlmutter JI, Shropshire JD,
Layton EM, Funkhouser-Jones LJ, Beckmann JF, Bordenstein Seth R. 2017.

Prophage WO genes recapitulate and enhance Wolbachia-induced cytoplasmic incompatibility. *Nature* 543:243–247.

Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* 25:1754–1760.

Lynch M, Ali F, Lin T, Wang Y, Ni J, Long H. 2023. The divergence of mutation rates and spectra across the Tree of Life. EMBO reports 24:e57561.

Lynch M, Conery JS. 2003. The origins of genome complexity. Science 302:1401–1404.

Maechler M, Rousseeuw P, Struyf A, Hubert M, Hornik K. 2021. cluster: Cluster Analysis Basics and Extensions. Available from: https://CRAN.R-project.org/package=cluster

Manzano-Marín A, D'acier AC, Clamens A-L, Orvain C, Cruaud C, Barbe V, Jousselin E. 2018. A Freeloader? The Highly Eroded Yet Large Genome of the Serratia symbiotica Symbiont of Cinara strobi. Genome Biology and Evolution 10:2178.

Manzano-Marín A, Latorre A. 2014. Settling Down: The Genome of Serratia symbiotica from the Aphid Cinara tujafilina Zooms in on the Process of Accommodation to a Cooperative Intracellular Life. Genome Biology and Evolution 6:1683–1698.

Marçais G, Delcher AL, Phillippy AM, Coston R, Salzberg SL, Zimin A. 2018. MUMmer4: A fast and versatile genome alignment system. *PLOS Computational Biology* 14:e1005944.

Martinez J, Ant TH, Murdochy SM, Tong L, Filipe A da S, Sinkins SP. 2022. Genome sequencing and comparative analysis of Wolbachia strain wAlbA reveals Wolbachia-associated plasmids are common. *PLOS Genetics* 18:e1010406.

Martinez J, Klasson L, Welch JJ, Jiggins FM. 2021. Life and Death of Selfish Genes: Comparative Genomics Reveals the Dynamic Evolution of Cytoplasmic Incompatibility. *Molecular Biology and Evolution* 38:2–15.

Martinez J, Ok S, Smith S, Snoeck K, Day JP, Jiggins FM. 2015. Should Symbionts Be Nice or Selfish? Antiviral Effects of Wolbachia Are Costly but Reproductive Parasitism Is Not.McGraw EA, editor. *PLoS Pathog* 11:e1005021.

McCutcheon JP, Moran NA. 2012. Extreme genome reduction in symbiotic bacteria. *Nat Rev Microbiol* 10:13–26.

Miller WJ, Ehrman L, Schneider D. 2010. Infectious Speciation Revisited: Impact of Symbiont-Depletion on Female Fitness and Mating Behavior of Drosophila paulistorum.Parrish C, editor. *PLoS Pathog* 6:e1001214.

Miller WJ, Riegler M. 2006. Evolutionary Dynamics of *w* Au-Like *Wolbachia* Variants
in Neotropical *Drosophila* spp. *Appl Environ Microbiol* 72:826–835.

Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, Von Haeseler A,
Lanfear R. 2020. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic
Inference in the Genomic Era.Teeling E, editor. *Molecular Biology and Evolution*
37:1530–1534.

Moran NA, Plague GR. 2004. Genomic changes following host restriction in bacteria.
Current Opinion in Genetics & Development 14:627–633.

Müller MJ, von Mühlen C, Valiati VH, Valente VL da S. 2012. Wolbachia pipientis is
associated with different mitochondrial haplotypes in natural populations of
Drosophila willistoni. *Journal of Invertebrate Pathology* 109:152–155.

Nikoh N, Hosokawa T, Moriyama M, Oshima K, Hattori M, Fukatsu T. 2014.
Evolutionary origin of insect–Wolbachia nutritional mutualism. Proceedings of the
National Academy of Sciences 111:10257–10262.

Perlmutter JI, Bordenstein Sarah R., Unckless RL, LePage DP, Metcalf JA, Hill T,
Martinez J, Jiggins FM, Bordenstein Seth R. 2019. The phage gene wmk is a
candidate for male killing by a bacterial endosymbiont. *PLOS Pathogens*
15:e1007936.

Powell JR, Sezzi E, Moriyama EN, Gleason JM, Caccone A. 2003. Analysis of a shift
in codon usage in Drosophila. J Mol Evol 57 Suppl 1:S214-225.

R Core Team. 2018. R: A Language and Environment for Statistical Computing.
Vienna, Austria: R Foundation for Statistical Computing Available from:
https://www.R-project.org/

Rambaut, A., 2009. FigTree v1.4: Tree Figure Drawing Tool. Available at:
http://tree.bio. ed.ac.uk/software/figtree/

Raychoudhury R, Baldo L, Oliveira DCSG, Werren JH. 2008. Modes of Acquisition of
Wolbachia: Horizontal Transfer, Hybrid Introgression, and Codivergence in the
Nasonia Species Complex. *Evolution* 63:165–183.

Renoz F, Arai H, Pons I. 2024. The genus Sodalis as a resource for understanding
the multifaceted evolution of bacterial symbiosis in insects. Symbiosis 92:187–208.

Renoz F, Foray V, Ambroise J, Baa-Puyoulet P, Bearzatto B, Mendez GL,
Grigorescu AS, Mahillon J, Mardulyn P, Gala J-L, et al. 2021. At the Gate of
Mutualism: Identification of Genomic Traits Predisposing to Insect-Bacterial
Symbiosis in Pathogenic Strains of the Aphid Symbiont Serratia symbiotica. Front.
Cell. Infect. Microbiol. 11.

Russell SL, Chappell L, Sullivan W. 2019. A symbiont's guide to the germline. In:
Current Topics in Developmental Biology. Vol. 135. Elsevier. p. 315–351.

Ryan JA, Ulrich JM. 2022. quantmod: Quantitative Financial Modelling Framework.
Available from: https://CRAN.R-project.org/package=quantmod

Schneider DI, Ehrman L, Engl T, Kaltenpoth M, Hua-Van A, Le Rouzic A, Miller WJ.
2019. Symbiont-Driven Male Mating Success in the Neotropical Drosophila
paulistorum Superspecies. *Behav Genet* 49:83–98.

Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics*
30:2068–2069.

Sheeley SL, McAllister BF. 2009. Mobile male-killer: similar Wolbachia strains kill
males of divergent Drosophila hosts. Heredity 102:286–292.

Shen X-X, Hittinger CT, Rokas A. 2017. Contentious relationships in phylogenomic
studies can be driven by a handful of genes. Nat Ecol Evol 1:1–10.

Shropshire JD, Leigh B, Bordenstein SR. 2020. Symbiont-mediated cytoplasmic
incompatibility: What have we learned in 50 years? *eLife* 9:e61989.

Small ST, Tisch DJ, Zimmerman PA. 2014. Molecular epidemiology, phylogeny and
evolution of the filarial nematode Wuchereria bancrofti. *Infect Genet Evol* 0:33–43.

Syberg-Olsen MJ, Garber AI, Keeling PJ, McCutcheon JP, Husnik F. 2022.
Pseudofinder: Detection of Pseudogenes in Prokaryotic Genomes. *Molecular Biology*
and Evolution 39:msac153.

Turelli M, Hoffmann AA. 1995. Cytoplasmic incompatibility in Drosophila simulans:
dynamics and parameter estimates from natural populations. Genetics 140:1319–
1338.

Turelli M, Cooper BS, Richardson KM, Ginsberg PS, Peckenpaugh B, Antelope CX,
Kim KJ, May MR, Abrieux A, Wilson DA, et al. 2018. Rapid Global Spread of wRi-like
Wolbachia across Multiple Drosophila. Current Biology 28:963-971.e8.

Unckless RL, Boelio LM, Herren JK, Jaenike J. 2009. Wolbachia as populations
within individual insects: causes and consequences of density variation in natural
populations. *Proceedings of the Royal Society B: Biological Sciences* 276:2805–
2811.

Unckless RL, Jaenike J. 2012. Maintenance of a Male-Killing Wolbachia in
Drosophila Innubila by Male-Killing Dependent and Male-Killing Independent
Mechanisms. Evolution 66:678–689.

Vancaester E, Blaxter M. 2023. Phylogenomic analysis of Wolbachia genomes from
the Darwin Tree of Life biodiversity genomics project.Teixeira L, editor. *PLoS Biol*
21:e3001972.

Wallau GL, da Rosa MT, De Ré FC, Loreto ELS. 2016. Wolbachia from Drosophila
incompta: just a hitchhiker shared by Drosophila in the New and Old World? Insect
Molecular Biology 25:487–499.

Waterhouse AM, Procter JB, Martin DMA, Clamp M, Barton GJ. 2009. Jalview
Version 2—a multiple sequence alignment editor and analysis workbench.
Bioinformatics 25:1189–1191.

Wernegreen JJ. 2015. Endosymbiont evolution: predictions from theory and
surprises from genomes: Endosymbiont genome evolution. *Ann. N.Y. Acad. Sci.*
1360:16–35.

Werren JH, Baldo L, Clark ME. 2008. Wolbachia: master manipulators of
invertebrate biology. *Nat Rev Microbiol* 6:741–751.

Wickham H, Averick M, Bryan J, Chang W, McGowan LD, François R, Grolemund G,
Hayes A, Henry L, Hester J, et al. 2019. Welcome to the tidyverse. *Journal of Open*
Source Software 4:1686.

Wolfe TM, Bruzzese DJ, Klasson L, Corretto E, Lečić S, Stauffer C, Feder JL,
Schuler H. 2021. Comparative genome sequencing reveals insights into the
dynamics of Wolbachia in native and invasive cherry fruit flies. *Molecular Ecology*
30:6259–6272.

Wu M, Sun LV, Vamathevan J, Riegler M, Deboy R, Brownlie JC, McGraw EA,
Martin W, Esser C, Ahmadinejad N, et al. 2004. Phylogenomics of the Reproductive
Parasite Wolbachia pipientis wMel: A Streamlined Genome Overrun by Mobile
Genetic Elements. PLOS Biology 2:e69.

Xie Z, Tang H. 2017. ISEScan: automated identification of insertion sequence
elements in prokaryotic genomes. *Bioinformatics* 33:3340–3347.

Yang Z. 2007. PAML 4: Phylogenetic Analysis by Maximum Likelihood. Molecular
Biology and Evolution 24:1586–1591.

Zeng Y, Wiens JJ. 2021. Do mutualistic interactions last longer than antagonistic
interactions? Proceedings of the Royal Society B: Biological Sciences
288:20211457.

Zhang Y-K, Ding X-L, Zhang K-J, Hong X-Y. 2013. Wolbachia Play an Important
Role in Affecting mtDna Variation of Tetranychus truncatus (Trombidiformes:
Tetranychidae). *Environmental Entomology* 42:1240–1245.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7