Intestinal Tissue Mechanics Regulate Angiogenesis and Stem Cell Proliferation via Vascular Piezo

Jade A. Phillips[1‡], Jessica Perochon[2‡*], Cai T. Johnson[2,3], Matthew Walker[3,5], Colin Nixon[1], Mark Hughes[1], André Barros-Carvalho[4], Yachuan Yu[1], Louise Mitchell[1], Karen Blyth[1,2], Massimo Vassalli[3,5], Julia B. Cordero[1,2*]

[1] CRUK Scotland Institute, Garscube Estate, Switchback Road, Glasgow, G61 1BD, U.K.

[2] School of Cancer Sciences, Wolfson Wohl Cancer Research Centre, University of Glasgow, Garscube Estate, Switchback Road, Glasgow, G61 1QH, U.K.

[3] James Watt School of Engineering, University of Glasgow, University Avenue, Glasgow, G12 8QQ, U.K.

[4] IBMC - Instituto de Biologia Molecular e Celular; i3S - Instituto de Investigação e Inovação em Saúde, Universidade do Porto, 4200-135 Porto, Portugal.

[5] School of Molecular Biosciences, Centre for the Cellular Microenvironment, Joseph Black Building, University of Glasgow, University Avenue, Glasgow, G12 8QQ, U.K

[‡] Equally contributing authors

*Corresponding authors: julia.cordero@glasgow.ac.uk and jessica.perochon@glasgow.ac.uk

Keywords: intestinal biomechanics, vascular intestinal stem cell niche, intestinal stem cells, intestinal regeneration, intestinal tumourigenesis, interorgan signaling.

Abstract

The vasculature is a prominent component of developmental and adult tissue
microenvironments. How, tissue specific characteristics and environmental states
influence vascular biology and function, remains largely understudied.

Previously, we discovered crosstalk between the adult intestinal epithelium
and the vasculature-like tracheal system of the fruit fly *Drosophila melanogaster*,
which is driven by reactive oxygen species (ROS) during pathogen induced-intestinal
regeneration. However, chemical stress signals alone are insufficient to explain the
rich diversity of vasculature/tissue interactions in living systems and justify the widely
observed adaptation of the vascular network in physiology and disease. Here, we
uncover reciprocal, mechanochemical interorgan communication between the adult
intestine and its vascular niche, which shapes vascular and epithelial tissue
adaptations and drives stem cell proliferation during intestinal regeneration and
tumour growth. Mechanistically, apoptotic epithelial cells within the regenerating
intestine induce local and global mechanical changes in the gut, which results in
activation and upregulation of the mechanosensitive ion channel Piezo in a subset of
gut-associated trachea. Piezo drives a specific molecular program within the trachea
through activation of the mechanosensitive transcription factor Yorkie/YAP, leading to
tracheal remodelling and intestinal stem cell proliferation. Furthermore, we identify a
non-redundant role of vascular Piezo1 driving remodelling of the intestinal crypt
vasculature and inducing crypt growth, WNT signalling activity, and stem cell
proliferation in the regenerating mouse small intestine.

Our cross-species *in vivo* study reveals previously unrecognised
mechanosensory regulation of intestinal regeneration and tumourigenesis
through the vascular-stem cell niche and highlights the importance of studying tissue
and context specific vascular cell biology to understand intestinal plasticity and the
complexity of tissue/vasculature interactions within a living organ.

Introduction

The adult intestine is a remarkably plastic organ, capable of responding to a myriad of environmental stimuli through intrinsic cellular adaptations and complex interactions with a highly heterogeneous microenvironment (1-3). Within the intestinal epithelium, intestinal stem cells (ISCs) have a vast regenerative capacity, required to enable a high turnover of the epithelial lining and maintain tissue homeostasis (4-6), while diverse enteroendocrine cell populations endow the intestine with rich sensory and signalling functions (7, 8) and enterocytes contribute to absorptive and metabolic functions of the intestinal epithelium (9, 10). How, specialised interactions between the intestinal epithelium and it's diverse microenvironment contribute to tissue plasticity and the functional versatility of the adult intestine, is poorly understood.

The vasculature interacts with target tissues through the exchange of oxygen, metabolites and angiocrine factors (11-13). Molecular analysis of vascular endothelial cells across tissues, reveals tissue specific characteristics of the vascular microenvironment, likely tailored to individual needs of each target tissue. Beyond supporting organ specific functions, the vascular system of highly self-renewing and plastic tissues, such as the intestine and the skin, must adjust to frequent and often dramatic tissue remodelling processes, which range from discrete cellular adaptations to whole organ resizing, in response to a diverse range of environmental stimuli and organismal states (14-20). The biology underpinning context dependent cellular interactions and adaptations of the vascular microenvironment within an individual tissue remains significantly understudied.

The diversity of interactions between the adult intestine and its vascular microenvironment are reflected in the crosstalk between the midgut and its rich tracheal network in the fruit fly *Drosophila melanogaster* (14-17). The *Drosophila* tracheal system is a mesh-like structure composed of oxygen-delivering tubes, functionally and molecularly alike the mammalian blood vasculature and respiratory system (21, 22). *Drosophila* terminal tracheal cells (TTCs), which are related to vertebrate vascular tip cells (23), extend prominent cytoplasmic projections to supply oxygen to their target tissues (21, 24, 25). Previous work from us and others showed a crosstalk between the intestinal epithelium and the vasculature like

tracheal tissue in *Drosophila* during pathogen-induced intestinal regeneration and

tumour growth triggered by ROS dependent signalling (14, 15). However, chemical

stress signals alone are insufficient to explain the broad tracheal/midgut interactions

observed in the adult fly midgut (14, 15, 17) or adaptation of the mammalian vascular

network in broad physiological settings (18-20).

Through *in vivo* genetic studies in flies and mice, combined with live imaging,

quantitative biology and biophysical approaches, we reveal previously unrecognised

mechanosensory regulation of intestinal regeneration and tumorigenesis through the

vascular- intestinal stem cell niche and present the first report on the role of the

blood vasculature and vascular expressed Piezo1 as functional contributors to crypt

regeneration in the adult mouse small intestine.

Results

Cellular and tissue-scale re-sizing of the regenerating intestine correlate with remodelling of associated tracheal/vascular tissue.

The adult *Drosophila melanogaster* gut is densely surrounded by TTCs, which we labelled with *GFP* driven under the control of the Serum Response Factor promoter (*dSRF>GFP*) (14, 15). Remodelling of the adult gut tracheal network in *Drosophila* is regulated through cross-tissue chemical signalling downstream of ROS, which are a prominent feature of midguts subjected to damage by oral pathogenic infection or oxidative stress (14, 15) (Fig. 1a, a' and Extended data Fig.1a, b). However, the presence of ROS is not significant in midguts exposed to the basal membrane disruptor dextran sodium sulphate (DSS)(Extended data Fig. 1c, d) (26), which equally depend on active tracheal remodelling for their regeneration (14) (Fig. 1b, b'). Furthermore, treating flies with the antioxidant N-acetyl cysteine (NAC) alongside DSS has no significant impact on DSS induced tracheal remodelling or ISC proliferation (Extended data Fig. 1e-g). Such observations suggested that, remodelling of the adult gut associated trachea, is a response to diverse cues working cooperatively or independently from oxidative stress signals. Understanding such alternative processes of gut/tracheal interactions might explain the aetiology of organ associated vascular remodelling observed in broad physiological and pathological settings (18-20, 27-30).

Organ resizing are poorly understood phenomenology associated with adult midgut regeneration in *Drosophila* (14, 31). Image segmentation and quantitative morphometric analysis of control and damaged midguts revealed increased thickness of the dorsal-ventral (D/V) axis and/or shortening of the anterior-posterior (A/P) length in posterior midguts during *Pe* or DSS treatment (Fig. 1c-h). Next, we used available tools for image segmentation and morphometric analysis (32) to quantify cellular morphology parameters within the midgut epithelium from masks of segmented enterocytes (EC), which represent approximately 80% of the midgut epithelium. We observed a significant increase in eccentricity— a deviation from cellular circularity associated with mechanical changes (33, 34)— in ECs from *Pe* (Fig. 1i-j) or DSS treated midguts (Fig. 1k-l).

Next, we asked whether only introducing a mechanical stimulus to the intestinal epithelium would be sufficient to induce the tissue and cellular resizing as well as the tracheal remodelling observed in regenerating midguts. To address this, we fed flies with an indigestible fibre, methylcellulose, previously shown to induce an 'overfull' phenotype in the gut (35). We observed that methylcellulose feeding in the absence of damage recapitulated all morphological changes of regenerating midguts, including shortening of the A/P length of the gut tube, thickening of the D/V axis, and increased EC eccentricity (Fig. 1m-q). Importantly, gut morphological changes in methylcellulose fed animals were accompanied by a significant increase in remodelling of midgut associate terminal tracheal cells (TTCs) (Fig. 1r, s).

Finally, we measured cell morphology in control and regenerating crypts of the mouse small intestine, following 72 hrs post-injury by whole body irradiation (IR), which represents the point of maximal regenerative proliferation in the small intestinal crypts (36). We used immunostaining for E-cadherin to segment crypt cells for morphological analysis (Fig. 1t), which revealed a significant increase in cellular eccentricity of crypts from irradiated versus untreated mice (Fig. 1t', u). Observations reported in our previous study indicated an increase in CD31+ endothelial cells surrounding regenerative crypts of the mouse small intestine (14). A refined assessment of the blood vasculature through whole mount tissue preparations and immune-detection of Pecam-1, a marker of blood vascular endothelial cells, confirmed increased branching of crypt blood vasculature in the regenerating mouse small intestine (Fig. 1v, w). Therefore, as in the *Drosophila* midgut, changes in intestinal epithelial cell morphology correlate with increased remodelling of the blood vasculature surrounding regenerating crypts of the mouse small intestine.

Altogether these results suggest that mechanical changes of the intestinal epithelium are prominent and consistent features across regenerating intestinal models, which co-exist with ROS in a pathogen induced model of midgut regeneration. Furthermore, mechanical forces in the absence of intestinal damage are sufficient to drive remodelling of the midgut associated tracheal network.

The midgut epithelium experiences regionalised mechanical forces during intestinal regeneration.

While cell morphological changes are indicative of mechanical stress, further
information is necessary to quantify mechanical forces within a tissue. Prior studies
have compartmentalised the adult fly gut into histologically, molecularly and
metabolically defined sub-regions (37, 38) (Fig. 2a). Despite the global nature of the
gut epithelial damage used in our models of intestinal regeneration, phenotypes of
tracheal remodelling and EC cell changes were regionalised across the posterior
midgut. We observed that, while tracheal remodelling and EC aspect ratio were
significantly increased in the R4 of the posterior midgut from *Pe* fed animals, both
parameters remained largely stable in the R5 segment (Fig. 2b-e). Therefore, we
took advantage of this phenomenology to more directly correlate changes in local
intestinal mechanics with tracheal remodelling within individual midguts. First, we
used an available Bayesian Force Inference model in MATLAB to predict relative
forces in the midgut tissue across the R4 and R5 regions from tile scan images of *Pe*
infected and control guts, using the R5 gut region as internal control. We observed a
significant increase in cell pressure in the R4 region relative to R5 upon infection,
which was not observed in homeostasis (Fig. 2f-g'). To directly quantify regional
mechanical properties in the midgut, we next used nanoindentation to measure
stiffness of the gut tube across the R4 and R5 regions of freshly dissected control
and orally infected midguts (Fig. 2h). We detected a significant increase in stiffness
specific to the R4 region of the gut upon infection (Fig. 2i), which correlates with the
increased tensional state of this midgut region as evaluated from the force inference
analysis. Altogether, these results provide novel quantitative and qualitative data
correlating biomechanics of the intestinal epithelium and whole midgut with
remodelling of the tracheal gut microenvironment.

Midgut enterocytes and gut associated tracheal cells establish physical
contact during intestinal regeneration

Cellular projections of TTCs appear located between the visceral muscle layer
of the gut and the midgut epithelium (14). We reasoned that one way for trachea to
sense morphological changes within the midgut epithelium would be through direct
physical contact with ECs. To test this, we used the GRASP (GFP reconstitution
across synaptic partners) system (39) to visualise points of close proximity between
tracheal cells and ECs by expressing the 11[th] strand of a split GFP in the main
tracheal branches under a *Btl-lexA* driver and strands 1-10 in ECs under the *Mex-*

Gal4 driver (Fig. 2j). While a constitutive red signal will allow visualisation of gut
associated trachea (Fig. 2j), a GFP signal would only be detected upon close
interaction between tracheal cells and ECs. While in homeostatic conditions we saw
RFP but little GFP reconstitution, *Pe* infected midguts revealed the expected
increased tracheal branching and a significant increase in the number of GFP
positive enterocytes within the epithelium (Fig. 2k, l). We observed no GFP signal in
midguts independently expressing individual GRASP components (Fig. 2m, n).
These results suggest the presence of physical interaction between ECs and gut
associated trachea, which are induced during midgut regeneration. As such, gut-
associated tracheal cells may be equipped to directly sense mechanical changes
within the intestinal epithelium.

Enterocyte apoptosis is necessary and sufficient to influence the
biomechanical properties of the intestine and gut associated tracheal
remodelling.

Imbalance between cell proliferation and cell loss within the *Drosophila* midgut
epithelium has been associated with major organ resizing (31, 40). Our previous light
microscopy analysis of adult midgut associated trachea suggested that EC cell death
rather than ISC proliferation determines tracheal cell remodelling (14). Here, we
used a dual driver genetic system to induce adult EC cell death by overexpressing
the pro-apoptotic gene *bax* (*Mex^{ts}>bax*) while fluorescently labelling the tracheal
system, with a previously characterised tracheal specific *QF/QUAS* line (*QF6>mtd-*
Tomato) (15). Using this approach, we confirmed that EC death in the absence of
external damaging cues is sufficient to induce tracheal remodelling (Fig. 3a, b). We
therefore hypothesised that cell death could be driving morphological and
mechanical changes in the midgut epithelium, leading to tracheal remodelling.

Firstly, we observed a significant increase in cellular eccentricity in the
posterior midgut epithelium, following a short pulse of global Bax overexpression
within ECs from the undamaged midgut (Fig. 3c-d). To refine our analysis of EC
apoptosis into a localised rather than global setting, we next performed a time course
of oral *Pe* infection to monitor the earliest appearance of discrete cell death using a
live apoptotic sensor (41). We observe areas with individual apoptotic ECs 2hs after
initiation of *Pe* feeding (Fig. 3e; GFP^{+ve}; arrow). Segmentation and morphometric

analysis revealed a localized pocket of epithelia with increased cellular eccentricity
restricted to the area immediately surrounding an individual apoptotic cell (Fig. 3e'-f).
Lastly, blocking EC apoptosis by overexpressing the apoptotic inhibitor, p35
(*Mex^{ts}>p35*) significantly reduced cellular eccentricity (Fig 3g-h) and decreased gut
stiffness upon *Pe* infection (Fig. 3i). Importantly, while EC p35 overexpression
inhibited damage induced tracheal remodelling (Fig. 3j, k), it did not affect ROS
levels in *Pe* infected midguts (Fig. 3l, m). This set of data suggests that EC cell death
is a determinant factor in the morphological and biomechanical changes quantified in
regenerating intestines.

Mechanical changes within the intestine are sensed by tracheal
mechanosensory channel Piezo

The mammalian vasculature is a highly mechanosensitive tissue. Stimulation
of vascular endothelial cells by intrinsic shear stress and stiffness are capable of
driving angiogenesis and vascular remodelling (42-44). Mechanical signals within the
vascular endothelium are largely transduced through TRP and Piezo family ion
channels (45, 46). While the role of Piezo1 in vascular development is well
documented (47), little is known about mechanisms of regulation and/or function of
vascular Piezo1 in adult self-renewing epithelia.

Recent studies in adult *Drosophila* and mouse intestine reported a role of
stem/progenitor cell expressed Piezo in homeostatic stem cell maintenance and
differentiation of the secretory lineage of intestinal stem cells (35, 48). While a single
piezo gene exists in *Drosophila* (49), described functions of *Piezo* within the
mammalian intestinal epithelium, are redundant between *Piezo1* and *Piezo2* (48).

Given the physical interaction observed between midgut ECs and tracheal
cells (Fig. 2j-l), we hypothesised that, changes in the mechanical properties of the
regenerating midgut epithelium, could be detected through mechanosensory
channels expressed in the gut associated tracheal tissue. Using an available *GAL4*
reporter of endogenous *piezo* gene expression *(piezo-GAL4)* (35) we identified
subpopulations of tracheal cells expressing *piezo* in the adult posterior midgut
(Fig.4a). Strikingly, we observed a significant increase in tracheal *piezo* expression
upon intestinal damage with orally administered *Pe* or DSS (Fig. 4a, b). Tracheal

piezo upregulation was confirmed with a reporter of Piezo protein expression
Piezo[mimic] (Extended data Fig. 2a, b) (50, 51). Consistent with previous reports (14,
15), general tracheal remodelling of *Pe* treated midguts was significantly reduced
upon neutralization of ROS with NAC treatment (Extended data Fig. 2c). However,
piezo upregulation within trachea of *Pe* treated midguts was not dependent on
oxidative stress (Extended data Fig. 2d, e). However, we noted that *piezo* expressing
TTCs showed partially reduced remodelling following *Pe* treatment (Extended data
Fig. 2d), suggesting that some of the mechanisms leading to tracheal remodelling
downstream of *piezo* may be shared with those activated by oxidative stress (14, 15).
Methylcellulose induced mechanical stretch of the intestine in the absence of
damage, recapitulated *piezo* upregulation in TTCs (Extended data Fig. 2f, g).
Selective chemical activation of Piezo by agonist Yoda1 was sufficient to induce
upregulation of *piezo* expression within the trachea, tracheal remodelling and ISC
proliferation (Fig. 4c-f and Extended data Fig. 2h). Overall, these results indicate that
mechanical changes within the regenerating *Drosophila* midgut epithelium lead to
non- autonomous upregulation of *piezo* in a subpopulation of gut associated trachea.
Remarkably, Piezo activation upregulates *piezo* gene expression within adult
Drosophila midgut trachea (Fig. 4c, d).

Next, we used RNA interference gene knockdown to assess the functional
role of tracheal *piezo* in TTC remodelling and ISC proliferation in regenerating
midguts. Our results showed that, knocking down *piezo* in adult TTCs
(*dSRF^{ts}>piezo-IR*), impaired tracheal remodelling and ISC proliferation upon midgut
epithelium injury through oral administration of *Pe* or DSS (Fig. 4g-i). However,
epithelial eccentricity was not affected in *dSRF^{ts}>piezo-IR* midguts (Extended data
Fig. 2i, j), suggesting that mechanical changes of the midgut epithelium precede
tracheal *piezo* activation and function in midgut regeneration. Additionally,
overexpression of tracheal *piezo* upon infection significantly potentiated tracheal
remodelling and ISC proliferation during midgut regeneration and impaired the
recovery of tracheal and ISC homeostasis following removal of damage (Fig. 4j-l).
Therefore, tracheal *piezo* is necessary and sufficient for remodelling of gut
associated TTCs and it is an instrumental regulator of intestinal regeneration
dynamics during and post-damage.

Dynamic *piezo* regulation and activity during damage induced midgut
regeneration.

To further characterize *piezo* expression in the midgut, we performed a time
course experiment, which revealed a gradual increase in gut-associated tracheal
piezo expression during a 16h period of *Pe* feeding (Fig. 4m, n). The observed time-
dependent pattern of *piezo* expression correlated with gut tracheal remodelling
dynamics and nanoindentation measures of intestinal tube stiffness (Fig. 4o, p).

Signalling downstream of Piezo channel activity involves the increase of
intracellular Calcium (Ca^{2+}) influx (46, 47). We therefore used a genetic Ca^{2+} sensor
(*GCamp7s*) expressed in tracheal cells to measure Piezo activity during a time
course of intestinal damage (Fig. 4q). Intracellular Ca^{2+} levels in TTCs depicted a
temporal pattern of increase consistent with the one established for tracheal
remodelling and *piezo* expression during the monitored stages of midgut damage
(Fig. 4q, r). Importantly, we confirmed that *GCamp7s* activation in gut associated
trachea was dependent on Piezo (Extended data Fig. 2k, Supplementary Movies 1-
3). Furthermore, longitudinal studies through in vivo imaging allowed us to visualise
discrete Ca^{2+} pulses in extending and fusing cytoplasmic projections of TTCs as
early as 2hs after the start of *Pe* feeding (Fig. 4s). This timeframe precedes a
detectable increase in global tracheal remodelling (Fig. 4m).

In summary, this set of results suggests that temporal patterns of *piezo* gene
upregulation and Piezo channel activation in gut associated trachea overlap with the
dynamic of gut trachea remodelling and gut biomechanical changes observed during
midgut regeneration. Importantly, our *in vivo* tracking of Ca^{2+} influx in tracheal cells
suggest that Piezo activation in gut associated trachea precedes a global increase in
tracheal remodelling.

Piezo activates a Yorkie/YAP-TAZ dependent mechanosensory program
for tracheal remodelling in the regenerating midgut.

We next looked into the downstream mediators of the role of Piezo in midgut
tracheal remodelling, by first assessing its impact on an oxidative stress responsive
tracheal programme, including regulation of Fibroblast growth factor-like *branchless*

(*bnl*) and its receptor breathless (*btl*) in TTCs (14, 15). Interestingly, while *piezo*
knockdown from trachea impaired Bnl upregulation following oral *Pe* infection (Fig.
5a, b), it did not affect upregulation of the gene encoding for the Bnl receptor, *btl*
(Extended data Fig. 3a, b). On the other hand, *piezo* knockdown led to significant
decrease in Myc expression in TTCs (Fig. 5c, d), which is also a target of oxidative
stress signals (14, 15). These results suggest that mechanical and oxidative stress
signals partially overlap in the molecular programs they regulate in midgut
associated trachea during midgut regeneration.

How does Piezo activation lead to transcriptional changes in TTCs? We
hypothesise that the mechanosensitive transcription factors YAP/TAZ, which have
been functionally linked to Piezo 1 in mammalian endothelial cells (52-54), may
mediate the transcriptional effects of tracheal Piezo during midgut regeneration.
Drosophila Yorkie (Yki) is the orthologue of mammalian YAP/TAZ. As with their
mammalian counterparts, Yki activation following cell intrinsic or extrinsic mechanical
cues leads to its translocation from the cytoplasm to the nucleus to drive
transcriptional target regulation. Using an endogenous Yki protein reporter line, we
monitored Yki localisation within gut associated trachea in homeostatic and
regenerating midguts. In homeostasis, Yki expression was largely restricted to the
cytoplasm of TTCs, however, upon pathogenic infection or DSS feeding we saw
significant translocation of Yki to TTCs nuclei (Fig. 5e-g). Importantly, *piezo*
knockdown inhibited translocation of Yki in TTCs form regenerating midguts (Fig. 5e-
g). Therefore, Yki is activated in remodelling trachea from the regenerating midgut in
a Piezo dependent manner.

We next assessed the functional role of tracheal Yki in TTC remodelling and
regenerative ISC proliferation in the adult midgut. Knocking down *yki* from the adult
trachea using two independent RNAi lines resulted in a significant decrease in TTC
remodelling and ISC proliferation in *Pe* infected midguts (Fig. 6a-c). Conversely,
overexpression of a constitutively active Yki in adult TTCs potentiated tracheal
remodelling during the damage phase of intestinal regeneration and delayed
homeostasis recovery of both TTC remodelling and ISC proliferation even 48hs after
removing flies from pathogen containing food (Fig. 6d-f). Finally, to further interrogate
the functional link between Piezo and Yki in midgut associated trachea, we assessed
a potential dependency on Yki for Bnl and Myc upregulation in TTCs from

regenerating midguts. Consistently with the interrelationship observed between
Piezo and Yki (Fig. 5e, g), tracheal *yki* knockdown lead to significant impairment in
both, Bnl and Myc upregulation in TTCs from regenerating midguts (Fig. 6g-j).

Our results reveal a mechanosensitive model of gut/tracheal interaction during
adult midgut regeneration where Piezo activation and upregulation within gut
associated trachea is translated into a cell intrinsic molecular program through the
activation of Yki in terminal tracheal cells, leading to regenerative tracheal
remodelling and ISC proliferation (Fig. 6k)

The blood vasculature and vascular Piezo 1 are functional components of the regenerative stem cell niche in the mammalian small intestine.

The rich microenvironment of the mammalian gastrointestinal tract is
composed of diverse cellular subtypes, including multiple mesenchymal cell
populations, enteric neurons, and endothelial cells that together provide physical
support, contractility, and multiple stem cell niche factors to the adult intestinal
epithelium (55-61). While lymphatic vessels are known as functional components of
the mammalian stem cell niche, supporting intestinal stem cell function during tissue
homeostasis and regeneration (55, 62), the role of the intestinal blood vasculature
remains unknown.

Using single molecule RNA in situ hybridisation, we detected co-expression of
Piezo1 and vascular *Pecam-1* in the mouse small intestinal crypt (Fig. 7a). We used
a tamoxifen inducible *Pdgf-β-Cre-GFP* (63) for visualisation of the intestinal blood
vasculature from whole mount tissue preparations (Fig. 7b) (64) and to investigate
the role of the blood vessels and vascular Piezo1 in the mouse intestine. We used
control *Pdgf-β-CreER-GFP* mice (*Piezo 1$^{wt/wt}$*) or *Pdgf-β-CreER-GFP* mice combined
with an inducible Piezo1 knockout allele (*Piezo 1$^{fl/fl}$*) to knock out *Piezo1* from the
adult blood vasculature. Adult mice were treated with tamoxifen for Cre activation
and allowed to age for 5 days prior to induction of intestinal damage by 10 Gy whole
body irradiation (IR), followed by sampling at 72 hours post IR (Extended data Fig.
4a).

Strikingly, knocking out *Piezo1* from the adult vasculature had no impact on
homeostatic intestinal vascular remodelling (Extended data Fig 4b-d) or self-renewal
of the intestinal epithelium (Fig. 7c-e and Extended data Fig. 4 e-h). However,

vascular *Piezo1* knock out animals showed a significant impairment in crypt vascular
remodelling (Fig. 7f, g), and a decrease in the total number of regenerating crypts
and proliferating cell within intestinal crypts following irradiation-induced damage (Fig.
7h-j). Furthermore, Wnt signalling activity as per levels of Myc and Sox9 (65, 66) in
post-damage regenerating crypts was impaired in *Piezo^{fl/fl}* mice compared to
Piezo^{wt/wt} counterparts (Fig7k-m). Therefore, while dispensable to adult intestinal and
intestinal vasculature homeostasis, endothelial Piezo1 is necessary for vascular
remodelling and intestinal cell proliferation within regenerating crypts of the mouse
small intestine, following irradiation induced damage.

Piezo drives tracheal remodelling and intestinal hyperplasia in a
***Drosophila* model of intestinal tumourigenesis.**

Previous work reported active remodelling and a functional role of gut
associated trachea in fly models of oncogenic *ras* and *Notch* driven tumourigenesis
(15). Hyperplasia of the adult intestinal midgut epithelium induced through stem cell
overexpression of the *Src* oncogene (*esg^{ts}>Src*), leads to adenoma like structures,
which are likely to cause mechanical stress within the intestinal epithelium (67).
Consistently, *esg^{ts}>Src* midguts showed whole tissue and cellular morphology
changes akin to those observed in damaged or methylcellulose treated midguts (Fig.
7n-r). We generated a new transgenic system for Src overexpression (*lexAop-Src*) to
allow independent genetic labelling and manipulation of tracheal cells combined with
intestinal stem cell induction of *Src* overexpression. We confirmed the presence of
intestinal stem cell hyperproliferation and increased tracheogenesis in *esg^{ts}-*
LexA/LexAop-Src midguts (Extended data Fig. 4i-k). Furthermore, a Piezo protein
reporter fly line, revealed significant increase in tracheal Piezo expression following
Src overexpression in the intestinal epithelium (Fig. 7s, t). Importantly, knocking
down *piezo* in TTCs, inhibited tracheal remodelling and intestinal hyperplasia in Src
overexpressing midguts (Fig. 7u-w). Therefore, we show that tracheal Piezo is also
necessary for tumour induced tracheal remodelling and the reciprocal action of the
trachea in intestinal tumorigenesis in *Drosophila*.

Altogether, our cross-species study reports a novel pro-angiogenic
mechanical crosstalk between the intestinal epithelium and the tracheal/vascular
tissue. Mechanical forces generated by apoptotic intestinal epithelial cells during

intestinal regeneration contribute to cellular and tissue resizing and mechanical

changes within the intestinal epithelium, which induce activation of a subset of

trachea/vascular cells expressing the mechanosensory channel Piezo. Piezo

activation in the vascular microenvironment is necessary and sufficient to drive

remodelling of the vascular niche and intestinal stem cell proliferation during

regeneration and tumourigenesis.

Discussion

The vasculature is as an intrinsically mechanosensitive tissue and an

important constituent of the microenvironment of developing and adult organs.

Whether and how those vascular characteristics converge during the support of

target tissues is not well understood.

A few of the many diverse contexts of vascular remodelling in physiology and

disease include pregnancy and foetal growth, wound healing, bone fracture repair,

tumour growth and metastasis (20, 30, 68-72). A seemingly common theme amongst

the mentioned examples is the presence of vascular adaptation to environmental

and/or host organ changes. In contrast to the vast number of studies on intrinsic

forces driving vascularization, such as fluid shear stress (73), the impact and identity

of extrinsic physical cues on angiogenesis and vascular remodelling remains largely

understudied. Here, we report a novel cross-species pro-angiogenic mechanical

crosstalk between the intestinal epithelium and the tracheal/vascular system, which

drives reciprocal adaptation of the intestine and vascular microenvironment during

intestinal regeneration and tumorigenesis.

A diversity of extrinsic signals leading to common phenomenology of

post-developmental vascular remodelling

Hypoxia and oxidative stress modulate vascular/tracheal remodelling across

species (74-77). Nevertheless, adaptive vascularization of adult organs is observed

in the context of a wide range of chemical, physiological and environmental cues

(14-20, 78).

Our previous observations showed a conserved phenomenology of
tracheal/vasculature changes upon diverse intestinal insults (14). Here, we
demonstrate that mechanical forces from the intestinal epithelium are a common
denominator in gut/tracheal interactions during intestinal regeneration and
tumorigenesis. As such, our new findings may explain mechanisms of vascular
remodelling observed in board physiological and pathological contexts.

What underpins the diversity of vascular/tracheal adaptations across and
within tissues? Our results revealed Piezo expression in a subpopulation of gut
associated trachea (Fig. 4), which do not appear to depend on FGFR/Btl for their
function (Extended data Fig. 3). Furthermore, while *btl* expression is dependent on
ROS (14), *piezo* expression is largely refractory to the action of the antioxidant NAC
(Extended data Fig. 2). These results suggest that TTCs in the adult intestine may
be highly heterogenous. For instance, Piezo and Btl expressing trachea could
represent two distinct sub-populations. Tracheal cell diversity may explain the ability
of the tracheal/vascular microenvironment to respond to different cues and support
intestinal function accordingly.

Despite the heterogeneous nature of *piezo* expression in midgut trachea,
gene knock down from TTCs results in rather global effect on tracheal remodelling
(Fig. 4). A similar phenomenon is observed upon *btl* knockdown (14, 15). How do
context specific cues sensed by a sub-set of tracheal cells translate into global
vascular/tracheal responses? One possible answer to this open question may lie in
the existence of paracrine interaction amongst distinct tracheal cell populations.
While definite proof of such a phenomenon remains to be obtained, our live imaging
visualization of intracellular Ca^{2+} signals in gut-associated trachea revealed the
presence of branch fusion events between distinct tracheal cells and transfer of
intracellular calcium, during TTC remodelling (Fig. 4).

Cellular apoptosis as a regulator of intestinal biomechanics and gut
associated tracheal remodelling during intestinal regeneration.

Here, we show enterocyte morphology and mechanical changes within the
intestine are conserved across different challenges, including chemical and
pathogenic damage, physical forces and tumour formation (Figs.1 and 7). Whilst the
intestinal visceral muscle and the extracellular matrix likely contribute to the

biomechanical properties of the gut, intestinal epithelial changes appear to be a
predominant determinant of cellular and tissue mechanical properties measured in
regenerating midguts (Figs. 1-3).

We identified physical interaction between trachea and intestinal EC and EC
apoptosis as an important cue driving mechanical changes in regenerating midguts
as well as tracheal remodelling. Previous studies have characterized a force
generating role of cell death in developmental and homeostatic contexts. Apoptosis
influences shape and size of developing tissues (79, 80). Furthermore, in a
phenomenology reminiscent to our observations on individual apoptotic cells (Fig.3),
cells closest to apoptotic cells experience tensile stretching in a model of
homeostatic compensatory proliferation *in vitro* (81). Our findings suggest that
proposed roles of apoptosis in tissue biomechanics may also apply to a complex,
multilayer epithelial tissue such as intestine. However, apoptosis is likely a context
dependent rather than universal driver of biomechanical organ/vascular interactions.

Tumorigenesis is associated with significant mechanical changes such as
increased tissue stiffness (44). The non-receptor tyrosine kinase Src is activated
within the intestinal epithelium in response to mechanical cues from the
mesenchymal microenvironment (82). Consistently, intestinal tumours
overexpressing Src induce midgut tracheal remodelling in a Piezo dependent
manner (Fig. 7). The role of apoptosis in tumour-induced mechanical changes in the
intestine and gut associated trachea, remains to be addressed.

Multicellular role of Piezo mechanosensing in the adult intestine

Pioneering work in *Drosophila* discovered a cell autonomous role of Piezo
within stem/progenitor cells in the intestine, during differentiation of the secretory
lineage of ISCs (35). Similarly, subsequent work in the mammalian intestine reported
redundant roles of intestinal stem cell Piezo1 and Piezo2 in homeostatic stem cell
maintenance and differentiation of secretory cell lineages in the intestine (48).

Using *Drosophila* and mouse models, our results demonstrate a distinct and
non-redundant role of vascular/tracheal Piezo1 in intestinal regeneration,
contributing to angiogenesis and ISC proliferation during intestinal regeneration and
tumorigenesis. Whilst we show that vascular *Piezo1* is necessary for damage
induced stem cell proliferation and crypt regeneration in the mouse small intestine,

Its impairment does not impact homeostatic vascular remodelling or self-renewal of
the intestinal epithelium (Fig. 7). Furthermore, we observed no significant impact in
secretory lineage differentiation in either homeostatic or regenerating intestines
following *Piezo1* knockout from the adult vasculature.

As such, our *in vivo* work reports a novel function for Piezo1 in the intestine
and suggests an exciting therapeutic opportunity to target Piezo1 for regenerative
and anti-tumour purposes with low impact on basal tissue homeostasis.

Materials and Methods

Fly stocks and husbandry.

A list of the fly and mouse lines and full genotypes of animals used in this
study can be found in a dedicated section of the Materials and Methods. For
temperature sensitive experiments, using the *GAL4/UAS* or *LexA/LexAop; GAL80^{ts}*
system, flies were crossed, the F_1 progeny were reared, and experimental adults
were aged for 3-5 days after eclosion at 18°C. Then, animals were moved to 29°C
for 5 days to allow transgenes activation. Experiments that did not involve
temperature-sensitive transgenes were performed with animals kept at 25°C and
experimental adults aged for 5 days prior to any procedure. All experimental animals
were transferred to fresh food every 2 days. Only posterior midguts from adult mated
females were used in this study.

Generation of *lexAop-Src* transgenic line: Genomic DNA from *UAS-Src* flies
(67) was extracted and used as a template to amplify the inserted *Src* sequence by
PCR using the following vector specific primers:
AGATCTGCGGCCGCATCCTCTAGAGTCGATCGACAT (forward),
AGGTACCCTCGAGCCTGCAGGCATGCAAGCTT (reverse). The resulting PCR
product was purified and cloned into NotI-XhoI sites of *pJFRC19* (Cat No 26224;
Addgene) to generate *pJFRC19-Src*. Following *pJFRC19-Src* sequence confirmation,
the transgene-containing plasmid was injected for the generation of *13xLexAop-Src*
flies by BestGene Inc.

***Drosophila* intestinal damage models.**

Bacterial infection - *Pseudomonas entomophila* (*Pe*) – females were fed on filter

papers containing a 5% sucrose solution only or sucrose containing *Pe* OD_{600}=100

for 16h.

DSS treatment - (Sigma-Aldrich, cat no. 42867) – females were fed on filter papers

containing a 5% sucrose solution only or sucrose containing 3% DSS for 2 days with

fresh medium applied daily.

Methylcellulose feeding - (Sigma-Aldrich, cat no. 274429) – females were fed on

normal food as a control or on sucrose food containing 10% methylcellulose for 72h.

Flies were transferred onto new food every two days. Prior to dissection, control flies

were transferred over night to 5% sucrose on filter papers to clear the gut. Flies

exhibiting guts with over a 50% increase in diameter were selected as exhibiting the

'full gut' methylcellulose phenotype.

Drosophila chemical treatments.

NAC treatment - (Sigma-Aldrich, cat no. A7250) – Flies were placed on filter paper

discs soaked with a 5% sucrose solution containing 20mM NAC for 24h. Then, the

animals were fed with either 5% sucrose + NAC or 5% sucrose + NAC + *Pe*

(OD_{600}=100) for 16hs or 5% sucrose + NAC + 3% DSS for 48hs.

DHE treatment - (Invitrogen Molecular Probes, cat no. D23107) – Tissues were

dissected in Schneiders medium and then incubated in 30mM DHE solution for 5-

7mins at room temperature and in a dark chamber. Then, midguts were washed

three times with Schneiders medium for 5min, mounted in DAPI containing media

followed by immediate imaging and image capture.

Yoda1 treatment – Yoda1 (Tocris, cat no. 5586) was suspended in DMSO and

diluted to 1 µM in 5% sucrose. Vehicle containing an equivalent volume of DMSO in

5% sucrose was used as control. Flies were placed in vials containing filter paper

discs soaked with either solution for 16h at 29°C prior to dissection.

Drosophila immunohistochemistry.

Guts from adult female flies were dissected in PBS and immediately fixed in 4%

formaldehyde (FA, EM grade, Polysciences) diluted in PBS for 1h at room

temperature (RT) and then washed three times in PBS-T (PBS + 0.2% Triton X100).

Samples were incubated with the primary antibody diluted in PBT (PBS-T + 2 % BSA)

overnight at 4°C. Guts were washed three times in PBS-T and then incubated with
secondary antibody in PBT for 2h at RT. After incubation, guts were washed three
times for 20min in PBS-T and mounted on glass slides with 13mm x 0.12mm
spacers (sigma) in Vectashield mounting medium with DAPI (Vector Laboratories).
For anti-Myc, anti-β-gal and anti-Bnl staining, fixation was performed in 4%
formaldehyde for 1h, followed by 5min methanol fixation added to the sample
gradually and then 10min in 100% methanol. Samples were blocked in 7% goat
serum for 1h before primary antibody incubation. The following primary antibodies
were used: chicken anti-GFP (1:200, Abcam, cat. no. ab13970), rabbit anti-DsRed
(1:1000, Clontech, cat no. 632496), mouse anti-Dlg (1:50, DSHB 4F3E2, AB 528203),
mouse anti-PH3 (1:100; Cell Signaling, cat. No. 9706), rabbit anti-β-gal (1:1,000; MP
Biochemicals, cat no. 559761), guinea pig anti-Myc (1:100, gift from G. Morata), rat
anti-Bnl (1:50, M. Krasnow. Secondary antibodies: Alexa Fluor anti-chicken 488
(1;200), anti-rabbit 594 (1:100) and anti-mouse 647 (1:100) (Invitrogen) were used.
Guts were mounted in Vectashield anti-fade mounting medium for fluorescence with
DAPI (Vector Laboratories, Inc.) to visualize all nuclei.

Image acquisition.

Confocal Microscopy (Zeiss LSM 780/880) – Each image represents half-volume of
the posterior midgut (area between hindgut and the copper cell region) and was
acquired with a ×20, ×40 or ×63 lens on a Zeiss LSM 780 or 880 confocal
microscope using identical acquisition conditions for all samples from a given
experiment. The images represent the maximal intensity projection of Z-stacks and
were processed using ImageJ (Version 1.54f) and Carl Zeiss Zen 3.0 to adjust the
brightness and contrast.

Nanoindentation – Flies were anesthetised on ice and midguts were freshly
dissected on the day of measurement in Schneider's *Drosophila* medium (1x Sigma
Aldrich 50146) and then transferred onto a 0.28 mg/ml Cell-Tak (Corning, 354240)
pretreated glass bottom petri dish (thickness 1,5) with Shneider's *Drosophila* medium
for tissue hydration. Petri dishes were coated with 10μl of Cell-Tak overnight followed
by 2 washes with PBS and air drying. A Chiaro nanoindentation device (Optics11,
The Netherlands) integrated above an inverted Zeiss Axiovert 200M microscope was
used to probe gut mechanics using a procedure previously described (83). A

cantilever (stiffness (K)= 0.52 Nm^{-1}) with a spherical tip (radius= 3.0-3.5 µm) was

used to probe the midguts *ex vivo*. After probe calibration, an initial indentation was

performed to tune the experimental parameters. The threshold, amount of cantilever

bending required to recognise a substrate, and find surface values, distance in µm

the probe moves after indentation for subsequent indentation, were tuned according

to the initial force displacement curve produced. A matrix scan was configured at 25

points per scan at intervals of 5µm along the specified midgut region, scans were

repeated twice per region of interest for every midgut. The midguts were approached

at 5 µm/s and indented with displacement as defined in the displacement control

mode.

Ex vivo live imaging – Flies were placed in vials containing filter paper discs with

either 400 µl *Pe* or sucrose for 6 hours prior to imaging. Coverslips were glued into

imaging mounts and 10 µl Imaging Agarose (Low gelling temperature agarose 6%

(Sigma, A4018), Schneider Insect medium (Thermo, 21720024)) was placed on the

surface of the coverslip. Flies were dissected in cold imaging medium (Schneider

Insect medium (Sigma, 50146), 5% Fly Extract (VDRC), 1:1000 penicillin-

streptomycin (Invitrogen, 15140), 10% FBS (Sigma, F4135) and placed on the

agarose in modified Schneider's medium as previously described (84). Plates were

sealed with lids containing damp paper to maintain humidity. Z stacks of the gut were

acquired at a ZEISS 780 in a time series at a rate of 1 stack every 5 minutes for 10

hours unless otherwise indicated.

In vivo live imaging – Flies were placed on Pe or sucrose solution for two hours prior

to imaging. Flies were glued on their dorsal side and the A1 segment of abdominal

cuticle removed. A small portion of the R4 region was gently exposed with forceps

and covered with 2 percent low gelling temperature agarose (Sigma, A4018) in

Schneider's Insect Medium (Sigma 50146). The dorsal side of the fly was then

covered with Imaging medium (Schneider Insect Medium, 5 percent Fly Extract

(VDRC), 0.1 percent Penicillin-Streptomycin (Invitrogen, 15140), 10 percent FBS

(Sigma F4135)). A coverslip was gently placed over the dorsal side of the fly before

imaging on a Zeiss 880 confocal microscope. Acquisitions were taken at 5-minute

intervals. Protocol adapted from (85).

Quantification Methods.

<u>A/P and D/V length</u> – Gut dorsal-ventral (DV) and anterior-posterior length (AP) were
quantified using ImageJ (Version 1.54f) software. Length was measured using the
freehand tool in the A/P axis of the posterior midgut. The D/V axis was measured
from the widest point of the R4 region.

<u>ISC proliferation -</u> Proliferating cells were detected by immunostaining against PH3.
Olympus BX51fluorescence microscope was used to manually, on visual inspection,
count the number of cells positive for PH3 staining per posterior midgut.

<u>Tracheal coverage</u> – detailed protocol described in (14). In brief, tracheal tile scans
were Z projected in ImageJ (Version 1.54f) and binarised from thresholding. A
skeleton was then produced of the R4 and R5 regions of the gut and mean grey
value was used to measure coverage of tracheal branches.

<u>ROS fluorescence</u> – For quantification of total ROS levels, pixel intensities of Z
stacks (sum slices) were used. The mean of the summed DHE intensities averaged
from each tissue was used for statistical analysis.

<u>Eccentricity</u> - Images of the intestinal epithelium were projected using the local z
projector plugin in ImageJ (Version 1.54f) to produce a curvature corrected Z
projection and heightmap. Z projected images were then segmented in cellpose.
Cellpose segmentations were converted to 4-bit connectivity images using an
ImageJ plugin morpholibJ to create a cell mask (32). Masks and heightmaps from
each sample were then analysed in Matlab using Deproj to analyse cellular
eccentricity (86).

<u>Bayesian Force Inference of epithelium</u> - Z projections and segmentation masks of
R4/R5 tile scans were generated from Z-stacks as described above using
localzprojector and cellpose respectively. Segmentation masks were then run
through TissueAnalyser to produce coordinate data files of cell edges and vertices
within the segmentation mask (87). Coordinate files were then processed in
BayesianForceInference programme in MATLAB to generate tension and pressure
data(19).

<u>Stiffness (Nanoindentation)</u> - Acquired force-displacement curves (*F-z*) were loaded
in the graphical user interface (GUI) Nanoprepare (83), which was used to clean the
data by deselecting any curves with failed contact or inappropriate noise. Collating

this cleaned data set, formal data analysis was carried out on the approach segment

of the curve in the GUI SofTMech (83). The curves were then filtered to smooth any

background noise generated by movement through the solution before contact. The

segment denoting force applied (forward segment) to the midgut was then isolated.

The contact point was defined by a threshold algorithm, transforming the curves from

F-z to force-indentation (*F–δ*). To these *F–δ* curves, the Hertz model was applied to

the max indentation of $\delta = 0.1R$ to quantify the elastic property or stiffness (Pa) of the

midguts.

<u>GRASP signal</u> – GRASP positive cells were detected by immunostaining for GFP.

Zeiss 780 confocal microscope was used to manually count the number of GRASP+

cells per posterior midgut.

<u>*lacZ* reporters</u> – Antibodies against β-galactosidase were used to detect btl-lacZ

reporters. Images were taken using confocal microscopy and staining was quantified

using ImageJ. For each gut quantified, the background staining signal was

subtracted from the total signal of β-galactosidase detected in the TTCs, ISCs/EBs

or ECs. This value was then divided by the background signal to normalise the data.

<u>Myc and Bnl staining</u> – Midguts stained with antibodies to detect these proteins

included a methanol fixation step between the paraformaldehyde fixation and PBST

wash steps of the standard protocol, as described previously (36). Images were

acquired using confocal microscopy and staining was quantified using ImageJ

(Version 1.54f). For each quantified gut, the background staining signal was

subtracted from the total antibody signal in DAPI+ cells. This value was then divided

by the background signal to normalize the data.

<u>Tracheal expression of *piezo* gene and Piezo protein trap</u> - Quantifications were

generated from a single z-stack of the R4 region for each midgut by confocal

microscope and analysis was performed using ImageJ (Version 1.54f). A maximum

intensity projection was produced for all images and channels were split. A skeleton

was created in the channel corresponding to the gut trachea and threshold adjusted

accordingly to ensure good selection of all tracheal branches. A region of interest

was created from the trachea and applied to the *piezo* gene or Piezo protein trap

expressing channel and fluorescence intensity quantified.

<u>Tracheal GCamp signal</u> - Quantifications were generated from a single z-stack of the
R4 region for each midgut by confocal microscopy and analysis was performed using
ImageJ (Version 1.54f). A maximum intensity projection was produced for all images
and channels were split. A skeleton was created in the channel corresponding to the
gut trachea (red) and threshold adjusted accordingly to ensure good selection of
tracheal branches. A region of interest was created from the active GCamp
expressing trachea (green) and applied to the general trachea expressing channel
(red) and fluorescence intensity quantified.

<u>Yki::GFP reporter</u> – Quantifications were generated from a single Z stack of the R4
region of each midgut by confocal microscope and analysis was performed using
ImageJ (Version 1.54f). A single Z plane was used to quantify each body cell where
the nucleus was visible. Intensity of GFP was measured in the cytoplasm and the
nucleus of each body cell and a ratio of nuclear to cytoplasmic intensity was
generated to quantify nuclear translocation.

<u>Mouse models</u>

All mouse experiments were performed in accordance with the UK Home Office and
ARRIVE guidelines and were reviewed by the Animal Welfare and Ethical Review
Board of the University of Glasgow. Experiments were performed under PPL
PP7226959 and PP6345023. Transgenic mouse strains were backcrossed with
C57Bl6J for 10 generations prior to experimental work. *Pfgfb-CreER* were bred to
Piezo fl/fl to generate cohort (*CreER;Piezo$^{fl/fl}$*) and control (*CreER;Piezo$^{wt/wt}$*) animals.
Both male and female mice between 8-16 weeks of age were used for experiments.
Mice were induced by 200µl of 15mg/ml Tamoxifen (Sigma T5648) in corn oil by oral
gavage on two consecutive days. For regeneration experiments, mice were exposed
to 10Gy whole body irradiation using the Xtrahl irradiator 5 days after Tamoxifen
administration and then sampled 72 hours post radiation treatment. Non-irradiated,
tamoxifen treated mice were harvested 8 days post induction for assessment of
intestinal homeostasis. Equal numbers of males and females, which were matched
in the control and cohort groups were used. Intestines were flushed with cold PBS
and sections taken for wholemount immunostaining and cryosections which were
snap frozen in OCT. Ten x 1cm sections from the duodenum and jejenum were

wrapped in surgical tape to make parcels and fixed in 4 percent paraformaldehyde
overnight..

<u>Immunohistochemistry of mouse tissue samples</u>

All haematoxylin and eosin (H&E), immunohistochemistry (IHC), Alcian Blue (AB)
and dual *in-situ* hybridization immunofluorescence (ISH-IF) staining was
performed on 4µm formalin fixed paraffin embedded sections (FFPE) which had
previously been ovened at 60□C for 2 hours. FFPE sections for c-Myc (ab32072,
Abcam), chromogranin A (ab15160, Abcam) and Sox 9 (AB5535, Millipore)
staining were stained using an Agilent autostainer Link48. H&E staining was
performed on a Leica autostainer (ST5020). Staining for AB was performed
manually on FFPE sections that were dewaxed and rehydrated through xylene
and a graded ethanol series before washing in water. Rehydrated slides were
stained for 10 minutes in Alcian Blue solution before rinsing in tap water and then
counterstained with Eosin for 30 seconds.

<u>RNAscope of mouse tissue samples</u>

Dual *ISH-IF* detection for Pecam1 (316728-C2) and Piezo (500518-C1), 3-plex
negative probe (320878) and 3-plex positive probe (320885) (all Bio-Techne)
mRNA was performed using Multiplex fluorescent kit (322800; Bio-Techne) strictly
according to the manufacturer's instructions. The staining was visualised staining
with Opal dyes (FP1487001KT, FP1488001KT, Akoya Biosciences). To complete
staining, sections were placed in a 60ºC oven for 30 minutes till dry, then sections
were cover slipped using Ecomount (320409, Bio-Techne).

<u>Whole mount Immunostaining of mouse small intestine</u>

Samples were prepared using a protocol adapted from (64). Sections of small
intestine were flushed with cold PBS, cut open and pinned longitudinally on silicone
plates. Sections were then fixed overnight in 4 percent PFA at 4 degrees on a shaker.
Samples were then washed in PBS before washing in 10 percent sucrose in PBS
and then 20 percent sucrose with 10 percent glycerol in PBS. Samples were then
dipped in 0.1 percent sodium azide for sample preservation. Some sections were
stored in sodium azide for up to 3 months at 4 degrees for further immunostaining.
Samples were then washed in PBS before being placed in blocking buffer (PBS 5

percent donkey serum, 0.5 percent BSA, 0.3 percent Triton, 0.1 percent sodium

azide) for 2 hours before incubation with primary antibody for 48 hours at 4 degrees.

Primary antibodies were diluted to the following concentrations in blocking buffer:

chicken anti-GFP (1:400, Abcam), rat anti-Pecam (1:400, Cell Signalling), rabbit anti-

Ecadherin (1:500, Cell Signalling). Samples were then washed in blocking buffer

before incubation in secondary antibody for 16 hours at 4 degrees. All secondary

antibodies were diluted in blocking buffer to a concentration of 1:500. Samples were

washed in wash buffer (PBS 0.3 percent Triton) before post fixation in 1 percent PFA

at 4 degrees overnight. Samples were washed in PBS before mounting. Small strips

1-2 villi thick were cut from intestine sections and mounted in Vectashield with DAPI.

Images were acquired on a Zeiss 880. Vascular coverage was quantified using the

Mexican Hat filter and Skeletonise plugins on Z projections of villi in ImageJ (Version

1.54f). Epithelial quantifications were performed the same way as in the *Drosophila*

model using LocalZprojector, cellpose and Deproj. Five crypts per mouse were

chosen at random.

Statistics

Data were analysed using two-way ANOVA with Sidak's multiple comparisons or T-

test as indicated where $P>0.05$ was statistically significant. Results were analysed

using Prism Graphpad. Individual sample data points are shown with group means

presented as mean + standard error, where $*P=0.05$, $**P = 0.01$, $***P=0.001$ and

$****P= 0.0001$.

Key Resources Table

REAGENT OR RESOURCE	SOURCE	IDENTIFIER
Antibodies		
rat anti-Bnl	M. Krasnow	(88)
Chicken anti- βeta-Galactosidase, 1:400	Abcam	ab-9361
Rabbit anti-Chromogranin A, 1:700	Abcam	ab15160
Chicken anti-GFP, 1:4000	Abcam	ab-13970
Rabbit anti-DsRed	Clontech	632496
Guinea pig Anti-dMyc	Gift from G.Morata	N/A

Rabbit Anti-c-Myc, 1:200	Abcam	ab32072
Mouse anti-Dlg 1:100	DSHB	528203
Rabbit Anti-Ki67	Cell Signalling Technology	12202
Rabbit Anti-Sox-9	Millipore	AB5535
Rabbit anti pH3 S10, 1:100	Cell Signaling	NEB-9701
Mouse anti pH3 S10, 1:100	Cell Signaling	9706S
Rabbit anti-p-Src 1:100	Invitrogen	44660G
Rat anti-Pecam1 1:500	BD Pharmingen	557355/MEC13.3 1
Rabbit anti-Ecadherin 1:500	Cell Signalling	24E10
RNAscope *Mm-Piezo1-O1*	ACDBio	500518
RNAscope *Mm-Pecam1-C2*	ACDBo	316728-C2
Goat Alexa Fluor anti-chicken IgY 488, 1:100	Invitrogen	A11039
Goat Alexa Fluor anti-chicken IgY 594, 1:100	Invitrogen	A11042
Goat Alexa Fluor anti-mouse IgG 488, 1:100	Invitrogen	A11029
Goat Alexa Fluor anti-mouse IgG 594, 1:100	Invitrogen	A11032
Goat Alexa Fluor anti-mouse IgG 647, 1:100	Invitrogen	A21235
Goat Alexa Fluor anti-Rabbit IgG 488, 1:100	Invitrogen	A11008
Goat Alexa Fluor anti-Rabbit IgG 594, 1:100	Invitrogen	A11034
Donkey Alexa Fluor anti-Rat IgG 594, 1:100	Invitrogen	A21209
Opal 520	Akoya Biosciences	FP1487001KT
Opal 570	Akoya Biosciences	FP1488001KT
Chemicals		
NAC	Sigma	A7250
Sucrose	Fisher	S/8600/60

	Chemical	
Yoda1	Tocris	5586
Methylcellulose	Sigma	274429
DSS	Sigma	42867-5G
Formaldehyde	FA, EM grade,	30525-89-4
Tamoxifen	Sigma	T5648
Triton X100	Sigma	X-100-1L
Normal Goat serum	Abcam	ab7481
Vectashield with DAPI	Vector Laboratories	H-1200
Bovine Serum Albumin	Sigma	A3294-50G
Schnieder's Medium	Sigma	50146
Fly Extract	VDRC	
Pen-Strep	Invitrogen	15140
Whatman GF/B Glass microfiber filters 21mm	Cytiva	1821-021
Corn Oil	Sigma	C8267
QuikSil	World Precision Instruments	
DHE	Invitrogen	D23107
DMSO	ThermoFisher	J66650.AK
Cell-Tak	Sigma	CLS354240
Low gelling temperature agarose	Sigma	A4018
FBS	Sigma	F4135
Ecomount	Bio-Techne	320409
Sodium Azide	Sigma	26628-22-8
Plasmids and primer sequences		
pJFRC19	Addgene	Cat No 26224
lexAop-Src (forward) *AGATCTGCGGCCGCATCCTCTAGAGTCGATCGACAT*	Thermo Fisher Scientific	N/A
lexAop-Src (reverse) *AGGTACCCTCGAGCCTGCAGGCATGCAAGCTT*	Thermo Fisher Scientific	N/A

Experimental models: **Drosophila melanogaster** strains:		
Wild type strain (w^{1118})	Gift from Ross Cagan	N/A
dSRF-gal4, UAS-CD8GFP	Gift from Irene Miguel-Aliaga	(89)
esg-gal4, UAS-CD8GFP; tub-gal80^{ts}	S. Hayashi	(90)
esg-lexA	Bloomington Drosophila Stock Center	66632
lexAop-GFP	Bloomington Drosophila Stock Center	32210
esg-lexA, lexAop-GFP	Generated in the Cordero lab	N/A
piezo-gal4	Gift from Norbert Perrimon	(35)
QF6, QUAS mtdTomato	Gift from Chrysoula Pitsouli	(15)
Mex-gal4, tub-gal80^{ts}	Gift from Irene Miguel-Aliaga	N/A
UAS-piezo-IR1	Bloomington Drosophila Stock Center	2796
UAS-piezo-IR2	Bloomington Drosophila Stock Center	105132
btl-lacZ	Bloomington Drosophila Stock Center	12086
bnl-lacZ	Bloomington Drosophila Stock Center	11704
Yki::GFP	Gift from Barry Thompson	(91)
UAS-GC3Ai	Gift from Andrew Davidson	N/A
UAS-Yki-IR1	Bloomington *Drosophila* Stock Center	31965
UAS-Yki-IR2	Bloomington *Drosophila* Stock Center	34067
UAS-Yki^{CA}	Bloomington *Drosophila* Stock Center	28817
lexAop-Src	Generated in the Cordero lab for this study	N/A

lgl::mcherry	Gift from Eurico Morais-de-Sá	N/A
btl-LexA	Gift from Thomas Kornberg	(92)
UAS-GFP1-10, lexAop-GFP11	Gift from Thomas Kornberg	N/A
lexAop-Cherry-CAAX	Gift from Thomas Kornberg	(93)
QUAS-GCamp6m	Bloomington *Drosophila* Stock Center	91988
Piezo::GFP mimic	Bloomington *Drosophila* Stock Center	60209
UAS-Piezo-EGFP	Bloomington *Drosophila* Stock Center	58772
UAS-bax, tubgal80^{ts}	Gift from Cedric Polesello	N/A
Mouse Strains		
Pdgfb-CreER	Marcus Fruttiger	(63)
Piezo1$^{fl/fl}$	Jackson Laboratories	029213

Detailed Genotypes:

Figure 1

1. dSRF-Gal4, UAS-CD8GFP/+; tub-Gal80^{ts}/+
2. dSRF-Gal4, UAS-CD8GFP/lgl::mcherry; MKRS/+
3. dSRF-Gal4, UAS-CD8GFP/+; tub-Gal80^{ts}/+
4. dSRF-Gal4, UAS-CD8GFP/lgl::mcherry; tub-Gal80^{ts}/+
5. Pdgf-beta Cre-GFP Piezo$^{wt/wt}$ (C57BL/6)

Extended data Figure 1

1. dSRF-Gal4, UAS-CD8GFP/+; tub-Gal80^{ts}/+

Figure 2

1. dSRF-Gal4, UAS-CD8GFP/lgl::mcherry; tub-Gal80^{ts}/+
2. dSRF-Gal4, UAS-CD8GFP/+; tub-Gal80^{ts}/+
3. Mex-Gal4/Btl-LexAHG, lexAop-Cherry-CAAX; UAS-GFP1-10, lexAop-
GFP11/tub-Gal80^{ts}
4. Mex-Gal4/+; UAS-GFP1-10, lexAop-GFP11/tub-Gal80^{ts}
5. Btl-LexAHG, lexAop-Cherry-CAAX/+; UAS-GFP1-10, lexAop-GFP11/tub-
Gal80^{ts}

Figure 3

1. Mex-Gal4/UAS-Bax, tub-Gal80^{ts}; QF6::QUAS-TdTomato/tub-Gal80^{ts}
2. Mex-Gal4/UAS-Bax, tub-Gal80ts; tub-Gal80^{ts}/MKRS
3. Mex-Gal4/lgl::mcherry; UAS-GC3Ai/MKRS

4. Mex-Gal4/+; tub-Gal80^{ts}/UAS-p35

5. Mex-Gal4,tub-Gal80^{ts}/+; QF6::QUAS-TdTomato/UAS-p35

6. Mex-Gal4/+; tub-Gal80^{ts}/UAS-p35

Figure 4

1. Piezo-Gal4/+; UAS-mCD8GFP/QF6::QUAS-TdTomato

2. dSRF-gal4, UAS-CD8GFP/+; tub-gal80^{ts}/+

3. dSRF-gal4, UAS-CD8GFP/UAS-piezo-IR2; tub-gal80^{ts}/+

4. dSRF-gal4, UAS-CD8GFP/+; tub-gal80^{ts}/UAS-piezo-IR1

5. dSRF-Gal4, UAS-CD8GFP/+; tub-Gal80^{ts}/+

6. dSRF-Gal4, UAS-CD8GFP/UAS-Piezo-EGFP; tub-Gal80^{ts}/+

7. Piezo-Gal4/+; UAS-mCD8GFP/QF6::QUAS-TdTomato

8. QF6::QUAS-TdTomato/QUAS-GCamp6m

Extended data Figure 2

1. Piezo::GFP mimic/ QF6::QUAS-TdTomato

2. Piezo-Gal4/+; UAS-mCD8GFP/QF6::QUAS-TdTomato

3. dSRF-Gal4, UAS-CD8GFP/+; tub-Gal80^{ts}/+

4. dSRF-Gal4, UAS-CD8GFP/+; tub-Gal80^{ts}/UAS-PiezoIR1

5. QF6::QUAS-TdTomato/QUAS-GCamp6m

Figure 5

1. dSRF-gal4, UAS-CD8GFP/+; tub-gal80^{ts}/+

2. dSRF-gal4, UAS-CD8GFP/UAS-piezo-IR2; tub-gal80^{ts}/bnl-lacZ

3. dSRF-gal4, UAS-CD8GFP/+; tub-gal80^{ts}/+

4. dSRF-gal4, UAS-CD8GFP/+; tub-gal80^{ts}/UAS-piezo-IR1

5. Yki::GFP/ Yki::GFP; QF6, QUAS mtdTomato/+

Figure 6

1. dSRF-gal4, UAS-CD8GFP/+; tub-gal80^{ts}/+

2. dSRF-gal4, UAS-CD8GFP/+; tub-gal80^{ts}/UAS-Yki-IR1

3. dSRF-gal4, UAS-CD8GFP/UAS-Yki-IR2; tub-gal80^{ts}/+

4. dSRF-gal4, UAS-CD8GFP/+; tub-gal80^{ts}/ UAS-Yki^{CA}

Extended data Figure 3

1. dSRF-gal4, UAS-CD8GFP/+; tub-gal80^{ts}/btl-lacZ

2. dSRF-gal4, UAS-CD8GFP/UAS-piezo-IR2; tub-gal80^{ts}/btl-lacZ

Figure 7

1. Pdgf-beta CreER-GFP Piezo$^{wt/wt}$ (C57BL/6)

2. Pdgf-beta CreER-GFP Piezo$^{fl/fl}$ (C57BL/6)

3. esg-lexA, LexAop-GFP/+; QF6::QUAS-TdTomato/ tub-Gal80^{ts}

4. esg-lexA, lexOp-GFP/lexAopSrc; QF6::QUAS-TdTomato/ tub-Gal80^{ts}

5. esg-lexA/Piezo::GFP mimic; QF6::QUAS-TdTomato/ tub-Gal80^{ts}

6. esg-lexA/dSRF-Gal4, UAS-CD8GFP; UAS-PiezoIR1/ tub-Gal80^{ts}

7. esg-lexA/dSRF-Gal4, UAS-CD8GFP; +/lexAop-Src, tub-Gal80^{ts}

8. esg-lexA/dSRF-Gal4, UAS-CD8GFP; +/tub-Gal80^{ts}

9. esg-lexA/dSRF-Gal4, UAS-CD8GFP; UAS-PiezoIR1/lexAop-Src, tub-Gal80^{ts}

Extended data Figure 4

1. Pdgf-beta CreER-GFP Piezo$^{wt/wt}$ (C57BL/6)

2. Pdgf-beta CreER-GFP Piezo$^{fl/fl}$ (C57BL/6)

3. esg-lexA, lexAop-GFP/+; QF6::QUAS-TdTomato/ tub-Gal80^{ts}

4. esg-lexA, lexAop-GFP/lexAop-Src; QF6::QUAS-TdTomato/ tub-Gal80^{ts}

Figure Legends

Fig 1. Morphological changes in the regenerating adult intestine occur in
parallel of tracheal/vascular remodelling. a, b. Representative confocal images of
the adult *Drosophila* midgut associated trachea (green) in the context of sucrose and
Pe (**a**) or DSS (**b**) feeding. Scale bars, 20 microns. **a', b'**. Quantification of tracheal
remodelling as in a (n=21,21) and b (n=21,23), respectively. **c.** Overview of adult
Drosophila midgut in the context of sucrose and *Pe* feeding. Posterior midgut
regions are shaded in red. Anterior-posterior (A.P.) and dorsal-central (D.V) lengths
are marked with arrows. Scale bar, 100 microns. **d, e.** Quantification of anterior-
posterior (A/P) (**d**) and dorsal-ventral (D/V) (**e**) lengths from midguts as in c
(n=22,21,22,21). **f.** Overview of adult *Drosophila* midgut in the context of sucrose
and DSS feeding. Posterior midgut regions are shaded in red. Anterior-posterior
(A.P.) and dorsal-central (D.V) lengths are marked with arrows. Scale bar, 100
microns. **g, h.** Quantification of gut A/P (**g**) and D/V (**h**) length from midguts as in f
(n=12,21,12,17). **i**. Confocal images of the intestinal epithelium in the context of
sucrose feeding and *Pe* infection. *lgl*::mcherry labels cell junctions (red) and DAPI
labels all cell nuclei (white). Scale bar, 20 microns. **i'**. Representative images of
epithelial eccentricity in midguts as in i. More eccentric cells have a value close to 1
indicated by the yellow colouring and more circular cells have a value close to 0
indicated by the blue colouring. Scale bar, 10 microns. **j**. Quantification of epithelial
eccentricity as in i'. Smaller circles in the background indicate individual cell values
and large outlined circles indicate the average cell eccentricity value per gut
analysed. Each colour indicates a different biological replicate. (n=13, 12). **k**.
Confocal images of the intestinal epithelium in the context of sucrose feeding and
DSS damage. *lgl*::mcherry labels cell junctions (red) and DAPI labels all cell nuclei
(white). Scale bar, 20 microns. **k'**. Representative images of cellular eccentricity in

midguts as in k. Scale bar, 10 microns. **l**. Quantification of epithelial eccentricity as in

k'. Smaller circles in the background indicate individual cell values and large outlined

circles indicate the average cell eccentricity value per gut analysed. Each colour

indicates a different biological replicate (n=5,5). **m**. Confocal images of posterior

midgut of flies fed normal food (NF) or methylcellulose (MC). Posterior midgut region

is shaded in red, and A/P and D/V lengths are marked with arrows. Scale bar, 100

microns. **n, o.** Quantification of A/P length (**n**) and D/V axis (**o**) from midguts as in m

(n=12,22,12,22) **p**. Confocal images of epithelial cells in the midgut of flies fed

normal food (NF) or methylcellulose (MC). *lgl*::mcherry labels cell junctions (red) and

DAPI labels all cell nuclei (white). Scale bar, 20 microns. **p'**. representative images

of eccentricity in midguts as in p. **q**. Quantification of eccentricity as in p'. Smaller

circles in the background indicate individual cell values and large outlined circles

indicate the average cell eccentricity value per gut analysed. Each colour indicates a

different biological replicate (n=11,13) **r**. Representative confocal images of midgut

associated trachea (green) in the context of normal food (NF) or methylcellulose (MC)

feeding. Scale bar, 20 microns. **s**. Quantification of tracheal coverage in midguts as

in r (n=19,34). **t.** Confocal images of mouse intestinal crypts from control (non-

irradiated) and irradiated mice 72 hours post treatment. Images are produced using

wholemount immunostaining and cell boundaries are detected using E-cadherin

staining (red); DAPI (blue). Scale bar, 20 microns. **t'**. Representative image of

epithelial eccentricity in mouse crypts as in t. **u**. Quantification of epithelial

eccentricity as in t'. Smaller circles in the background indicate individual cell values

and large outlined circles indicate the average cell eccentricity value per gut

analysed (n=6,5). **v**. Wholemount immunostaining of crypt vasculature using Pecam-

1 (green) in the context of homeostasis (non-irradiated) and 72 hours post whole-

body irradiation (irradiated). Cell boundaries are detected using E-cadherin staining

(red). Scale bar, 20 microns. **w**. Quantification of crypt vascular branches in

intestines as in v (n=3,6). Data represent the mean ± s.e.m. n numbers indicated

for each graph are listed from left to right and correspond to the numbers of guts

analysed. <u>Statistics</u>: Two-tailed unpaired Student's t-test. *P=0.05, ** P = 0.01, ***

P=0.001 and **** P= 0.0001. DAPI, 4,6-diamidino-2-phenylindole.

Fig 2. **Enterocytes experience significant mechanical changes upon damage. a.** Overview of an adult *Drosophila* gut with R1-R5 regions labelled. Scale bar, 100 microns. **b, c** Confocal images of trachea (**b**) and intestinal epithelium (**c**) from R4 and R5 regions of the posterior midgut in the context of Sucrose or *Pe* infection. Scale bar, 20 microns. **d**. Quantification of tracheal coverage in the R4 and R5 regions of the midgut upon *Pe* infection. (n=6,6,6,6). **e**. Quantification of cell aspect ratio in R4 and R5 regions of the gut upon infection. Each data point represents an individual enterocyte. (n =6,6,6,6). **f**. Tilescans across the R4 and R5 regions of the posterior midgut where enterocytes cell junctions have been detected with lgl (green). Scale bar, 50 microns. **f'**. Representative image of inferred relative cell pressure calculated using Bayesian Force Inference. Cells of higher pressure are red/yellow and lower pressure cells blue/green. **g**. Quantification of relative pressure across the R4 (grey) and R5 (green) regions of the midgut upon sucrose feeding or *Pe* infection, measured by Bayesian Force Inference in MATLAB. Smaller circles represent individual cells and larger circles in foreground represent average of cells per gut (n=4,4,5,5). **h**. Representative image of nanoindentation setup. *Drosophila* gut is placed on cell-Tak adhesive and cantilever is used to detect stiffness by touching the outside of the gut. **i**. Quantification of stiffness across midgut R4 and R5 regions using nanoindentation in the context of sucrose feeding or *Pe* infection. Smaller circles represent individual measurements within a gut (n=4,5,5,3). **j**. Overview of GFP reconstitution across synaptic partners (GRASP) system. Components of a split GFP barrel are expressed in two different cell types where close contact results in GFP reconstitution and fluorescence. When expressed in the trachea (*btl-lexA*; red) and enterocytes (*Mex-gal4)* GFP (green) reconstitution is observed upon *Pe* damage as a result of close contact. **k**. Confocal images of GRASP expression in the intestinal epithelium and surrounding trachea following sucrose or *Pe* infection. Scale bar, 20 microns. **l**. Quantification of the number of GRASP positive cells per posterior midgut (n=20,23). **m, n**. Controls without *btl-lexA* driver system (**m**) and without *Mex-Gal4* driver system (**n**). Scale bar, 20 microns. Data represent the mean □±□s.e.m. n numbers indicated for each graph are listed from left to right and they correspond to the numbers of guts analysed. <u>Statistics</u>: **l**. Two-tailed unpaired Student's t-test. **d, e, g, i.** Two-way ANOVA followed by Tukey's multiple comparisons test *P=0.05, ** P = 0.01, *** P=0.001 and **** P= 0.0001. DAPI, 4,6-diamidino-2-phenylindole.

Fig 3. Cell death is necessary and sufficient for mechanical changes and tracheal remodelling in the regenerating intestine. a. Representative confocal images of midgut associated trachea (*QF>mtdTomato*; red) in wildtype midgut or following *bax* overexpression in enterocytes (*Mex^{ts}>bax*). Scale bar, 20 microns. **b**. Quantification of tracheal coverage in midguts as in a (n=6,11). **c.** Confocal images of enterocytes detected by Dlg staining (red) in wildtype midgut or following *bax* overexpression in enterocytes (*Mex^{ts}>bax*). Scale bar, 20 microns. **c'**. Representative image of eccentricity quantification in midguts as in c. More eccentric cells have a value closer to 1 in yellow and more circular cells have a value close to 0 in blue and green. **d**. Eccentricity quantification as in c'. Smaller circles in the background indicate individual cell values and large outlined circles indicate the average cell eccentricity value per gut analysed. Each colour indicates a different biological replicate (n=11,14). **e**. Confocal images of midgut expressing an apoptotic sensor (GC3Ai; green) in enterocytes (*Mex-Gal4*) and cell boundary marker (*lgl::mCherry*; red) at 2 hours post infection. The righ panel shows a zoom of area indicated by the dotted box in left panel. Arrow points to an individual apoptotic cell. Scale bar, 20 microns. **e'.** Segmentation of cell boundaries in midgut as in e. Apoptotic cell (orange; arrow) and Row1 (red) and Row3 (blue) from apoptotic cell are highlighted. **e''**. Representative image of epithelial eccentricity as in e. Row1 and Row3 are labelled as in e'. Arrow points to an individual apoptotic cell.   **f**. Quantification of epithelial eccentricity in cells witin Row1 and Row3 from apoptotic cell after 2 hours of *Pe* infection (n=3,3). **g**. Confocal images of the epithelium (Dlg; red) in wildtype midguts or following *p35* overexpression in enterocytes (*Mex^{ts}>p35*) upon sucrose or *Pe* feeding. Scale bar, 20 microns. **g'** Representative images of eccentricity in midguts as in g. **h**. Quantification of eccentricity as in g' (n=9,8,9,8). **i**. Quantification of stiffness (young's modulus) in measured using nanoindentation in midguts as in g. Smaller dots in the background represent individual cells and larger circles in the foreground represent average per gut (n=3,3,3,3). **j**. Confocal images of the trachea (*QF>mtdTomato*; red) in wildtype midguts or following *p35* overexpression in enterocytes (*Mex^{ts}>p35*) upon sucrose or *Pe* feeding. Scale bar, 20 microns. **K**. Quantification of tracheal remodelling from midguts as in j (n=9,12,3,7). **l**. Confocal images of DHE staining (red) in midguts as in j. Scale bar,

20 microns. **m**. Quantification of DHE staining from midguts as in l (n=11,10,11,13).

Data represent the mean ± s.e.m. n numbers indicated for each graph are listed

from left to right and they correspond to the numbers of guts analysed. <u>Statistics</u>: **b,**

d, f. Two-tailed unpaired Student's t-test. **h, j, k, m.** Two-way ANOVA followed by

Tukey's multiple comparisons test ** P = 0.01, *** P=0.001 and **** P= 0.0001. DAPI;

4,6-diamidino-2-phenylindole. DHE; Dihydroethidium.

Fig 4. Piezo activity dictates tracheal remodelling and ISC proliferation in the

***Drosophila* intestine. a.** Confocal images of midgut associated trachea

(*QF>mtdTomato*; green pseudocoloured) and *piezo* expression (magenta) in flies

treated with Sucrose, Pe or DSS. Arrows point to a *piezo* negative tracheal cell.

Scale bar, 20 microns. **b**. Quantification of *piezo* expression relative to total tracheal

coverage in midguts as in a (n=24,9,22). **c**. Confocal images of midgut associated

trachea (*QF>mtdTomato*; green pseudocoloured) and *piezo* expression (magenta)

following Yoda1 treatment. Scale bar, 20 microns. **d**. Quantification of *piezo*

expression relative to total tracheal coverage in midguts as in c (n=22,24). **e, f.**

Quantification of tracheal coverage (**e**) (n=23,30) and ISC proliferation (PH3+) (**f**)

(n=23,30) in midguts as in c. **g**. Representative confocal images of the midgut

associated trachea (*dSRF^{ts}>gfp*; green) in the context sucrose feeding, *Pe* infection

or DSS feeding in control animals or following RNAi *piezo* knockdown in trachea

(*piezo-IR1*). Scale bar, 20 microns. **h, i.** Quantification of tracheal coverage (**h**)

(n=23,19,8,21,23,14,13,14)   and   ISC   proliferation   (PH3+)   (**i**)

(n=14,19,9,30,23,11,13,14) in midguts as in g. **j**. Representative confocal images of

midgut associated tracheal tissue (*dSRF^{ts}>gfp*; green) in control or tracheal *piezo*

overexpressing animals upon 0 and 16hs of *Pe* infection or 48hs after *Pe* removal

(recovery). Scale bar, 20 microns. **k, l**. Quantification of tracheal coverage (**k**)

(n=29,16,36,23,23,19) and ISC proliferation (**l**) (n=25,22,20,23,22,17) in midguts as

in j. **m**. Confocal images of midgut associated trachea (*QF>mtdTomato*; green

pseudocoloured) and *piezo* expression (magenta) across a time course of 0-16

hours of *Pe* infection. Scale bar, 20 microns. **n**. Quantification of tracheal *piezo*

expression relative to total tracheal coverage in midguts as in m (n=14,19,11,25). **o**.

Correlation graph between tracheal coverage and *piezo* expression in each of the

time points represented in m. Pearson's correlation coefficient (r=0.9915). **p**.

Quantification of stiffness measured by nanoindentation (young's modulus) in
midguts from timepoints as in m. Each dot represents a single measurement within a
gut (n=3,3,3,3). **q**. Representative confocal images of trachea (*QF>mtdTomato*; red)
and GCamp6m calcium sensor (green) of a *Pe* infection time course from 0 to 16
hours of *Pe* feeding. Scale bar, 20 microns. **r**. Correlation graph between tracheal
coverage and GCamp6m signal in each of the time points represented in q.
Pearson's correlation coefficient (r=0.7345). **s**. Close up still frames of GCamp6m *in*
vivo live imaging in the context of *Pe* infection. Arrows indicate branches which show
a calcium impulse prior to a remodelling event. Scale bar, 20 microns. Data
represent the mean□±□s.e.m. n numbers indicated for each graph are listed from
left to right and they correspond to the numbers of guts analysed. Statistics: **b, d, e,**
f.Two-tailed unpaired Student's t-test. **h, i, k, l, n, p.** Two-way ANOVA followed by
Tukey's multiple comparisons test ** P = 0.01, *** P=0.001 and **** P= 0.0001. DAPI,
4,6-diamidino-2-phenylindole.

1029

Fig 5. Tracheal Piezo regulates Yki activity and remodelling pathways in gut
associated trachea. a. Representative confocal images of midgut associated
trachea (*dSRF^{ts}>gfp*; green) and *bnl-LacZ* expression (grey) upon sucrose or *Pe*
feeding in control midguts or following tracheal *piezo* knockdown (*piezo-IR1*). Scale
bar, 20 microns. **b**. Quantification of *bnl-LacZ* expression per TTC in midguts as in a
(n=18,16,22,22). **c**. Representative confocal images of midgut associated trachea
(*dSRF^{ts}>gfp*; green) and Myc expression (magenta) upon sucrose or *Pe* feeding in
control midguts or following tracheal *piezo* knockdown (*piezo-IR1*). Scale bar, 20
microns. **d**. Quantification of Myc intensity in TTCs as in c (n=18,23,13,26). **e**.
Representative confocal images of midgut associated tracheal tissue
(*QF>mtdTomato*; green pseudocoloured) and Yki::GFP (grey) in midguts from
sucrore, *Pe* or DSS fed control or *Pe* fed animals following tracheal *piezo* knockdown
(*piezo-IR1*). Scale bar, 20 microns. **f, g**. Quantification of Yki nuclear to cytoplasmic
ratio in midguts as in e. Each data point represents a single tracheal body cell; **f**
(n=10,10); **g** (n=8,7,8,10). Data represent the mean□±□s.e.m. n numbers indicated
for each graph are listed from left to right and they correspond to the numbers of
guts analysed. Statistics: **f.** Two-tailed unpaired Student's t-test. **b, d, g.**Two-way

ANOVA followed by Tukey's multiple comparisons test *** P=0.001 and **** P=
0.0001. DAPI, 4,6-diamidino-2-phenylindole.

1049

Fig 6. Yki translates mechanical signals into tracheal remodelling in the
***Drosophila* midgut. a.** Confocal images of trachea (*dSRF^{ts}>gfp*; green) upon
sucrose or *Pe* feeding in control midguts or following tracheal *yki* knockdown (*yki-*
IR1). Scale bar, 20 microns. **b, c.** Quantification of tracheal coverage (**b**)
(n=9,9,9,11,11,11) and ISC proliferation (PH3+) (**c**) (n=10,9,10,11,12,12) in midguts
as in a. **d**. Representative confocal images of midgut associated trachea
(*dSRF^{ts}>gfp*; green) without (control) or with overexpression of constitutively active
Yki (*Yki-CA*) upon 16hs of *Pe* infection or 48hs after *Pe* removal (recovery). Scale
bar, 20 microns. **e, f.** Quantification of tracheal coverage (**e**) (n=16,17,20,14) and ISC
proliferation (PH3+) (**f**) (n=16,17,20,14) in midguts as in d. **g**. Representation
confocal images of midgut associated trachea (*dSRF^{ts}>gfp*; green) and anti-Bnl
expression (grey) upon sucrose or *Pe* feeding in control midguts or following tracheal
yki knockdown (*yki-IR1*). Scale bar, 20 microns. **h**. Quantification of Bnl expression
per TTC in midguts as in g (n=15,18,18,16). **i**. Representative confocal images of
midgut associated trachea (*dSRF^{ts}>gfp*; green) and Myc expression (magenta) upon
sucrose or *Pe* feeding in control midguts or following tracheal *yki* knockdown (*yki-*
IR1). Scale bar, 20 microns. **j**. Quantification of Myc expression per TTC in midguts
as in i (n=14,16,13,18). Data represent the mean ± s.e.m. n numbers indicated for
each graph are listed from left to right and they correspond to the numbers of guts
analysed. **k**. Proposed working model of gut/tracheal mechanosensory crosstalk
during intestinal regeneration in *Drosophila*. Statistics: Two-way ANOVA followed by
Tukey's multiple comparisons test *P=0.05, ** P = 0.01, *** P=0.001 and **** P=
0.0001. DAPI, 4,6-diamidino-2-phenylindole.

1073

Fig 7. The role of vascular Piezo is conserved in mammalian intestinal
regeneration and *Drosophila* intestinal hyperplasia. a. Confocal images of single
molecule RNA in situ detection of *Pecam-1* (red) and *Piezo-1* (green; grey) in mouse
small intestinal crypts. DAPI (blue). Centre and right panels are magnified views of
the area highlighted by the dotted box in the left panel. Arrows point to *Piezo-1* signal

within a crypt. Scale bar, 60 microns. **b**. Confocal images of wholemount preparation

of intestinal tissue from *Pdgf-beta-CreER-GFP* mice. Cre expression (green); anti-

Pecam-1 (grey). Scale bar, 20 microns. **c**. H and E and Ki67 staining of small

intestinal samples from non-irradiated *Pdgf-beta-CreER Piezo1$^{wt/wt}$* and *Piezo1$^{fl/fl}$*

mice. Scale bar, 20 microns. **d, e.** Quantification of the number of crypts per

circumference (**d**) (n=6,6) and number of Ki67+ cells per intestinal crypt (**e**) (n=3,3) in

mice as in c. **f**. Wholemount immunostaining of the crypt intestinal vasculature of

irradiated *Pdgf-beta-CreER Piezo1$^{wt/wt}$* and *Piezo1$^{fl/fl}$* mice using anti-Pecam-1

antibody (green). Cell boundaries are detected using E-cadherin staining (red);

DAPI (blue). Scale bar, 20 microns. **g**. Quantification of crypt vascular branch

number in intestines as in f (n=6,6). **h**. H and E and Ki67 staining of small intestinal

samples from irradiated *Pdgf-beta-CreER Piezo1$^{wt/wt}$* and *Piezo1$^{fl/fl}$* mice. Scale bar,

20 microns. **i, j**. Quantification of number of crypts per circumference (**i**) (n=6,6) and

number of Ki67+ cells per intestinal crypt (**j**) (n=3,3) in mice as in h. **k.**

Immunostaining of Myc and Sox9 in intestinal crypts of *Pdgf-beta-CreER Piezo1$^{wt/wt}$*

and *Piezo1$^{fl/fl}$* mice following irradiation. Scale bars, 20 microns. **l, m.** Quantification

of Myc (**l**) (n=6,6) and Sox9 (**m**) (n=6,6) as a percentage of staining positive cells per

crypt in intestines as in k. **n**. Representative confocal images of adult *Drosophila*

midgut from control flies or upon overexpression of *Src* in stem/progenitor cells (*esg-*

lexA^{ts}). Posterior midgut regions are shaded in red. Anterior-posterior (A.P.) and

dorsal-central (D.V) lengths are marked with arrows. Scale bar, 100 microns. **o, p.**

Quantification of A/P length (**o**) (n=14,16) and D/V axis (**p**) (n=14,16) in midguts as in

n. **q**. Confocal images of the intestinal epithelium of control flies or upon

overexpression of *Src* in stem/progenitor cells (*esg-lexA^{ts}*). Enterocyte junctions are

labelled with Dlg (grey) and progenitor cells express GFP (green). Scale bar, 20

microns. **q'**. Eccentricity representative images of epithelia from midguts as in q. **r**.

Quantification of epithelial eccentricity as in q' (n=13,16). **s**. Confocal images of

control flies or upon overexpression of *Src* in stem/progenitor cells (*esg-lexA^{ts}*) and

stained with anti-phosphorylated Src (pSrc; green); Piezo expression (magenta); gut

associated trachea (*Q6>mtdTomato*; red). Scale bar, 20 microns. **t**. Quantification of

tracheal Piezo expression in midguts as in s (n=17,13). **u**. Confocal images of gut

associated trachea (*dSRF^{ts}>gfp*; green) from control flies or upon overexpression of

Src in stem/progenitor cells (*esg-lexA^{ts}*); with and without piezo knockdown (*piezo-*

IR1) in tracheal cells. Scale bar, 20 microns. **v, w.** Quantification of ISC proliferation

(PH3+) (**v**) (n=26,20,22,19) and tracheal coverage (**w**) (n=22,26,25,29) in midguts as in u. Data represent the mean□±□s.e.m. n numbers indicated for each graph are listed from left to right and they correspond to the numbers of guts analysed. <u>Statistics</u>: **d, e, g, i, j, l, m, o, p, r, t**. Two-tailed unpaired Student's t-test. **v, w.** Two-way ANOVA followed by Tukey's multiple comparisons test *P=0.05, ** P = 0.01, *** P=0.001 and **** P= 0.0001. DAPI, 4,6-diamidino-2-phenylindole.

Extended Data Fig 1. Mechanical changes in the intestine are sufficient to drive tracheal remodelling. a. Representative confocal images of intestinal DHE staining (red) for reactive oxigene species detection in sucrose and *Pe* fed flies. Scale bar, 20 microns. **b**. Quantification of DHE intensity in midguts as in a (n=5,9). **c**. Confocal images of intestinal DHE staining (red) in sucrose and DSS fed flies. Scale bars, 20 microns. **d**. Quantification of gut DHE intensity in midguts as in c (n=13,14). **e**. Confocal images of midgut associated trachea (*dSRF^{ts}>gfp*; green) from sucrose and DSS fed flies with and without treatment with the anti-oxidant NAC. Scale bar, 20 microns. **f, g.** Quantification of tracheal coverage (**f**) (n=17,17,16,17) and ISC proliferation (PH3+) (**g**) (n=17,17,16,17) in midguts as in e. Data represent the mean□±□s.e.m. n numbers indicated for each graph are listed from left to right and they correspond to the numbers of guts analysed. <u>Statistics</u>: **b, d**. Two-tailed unpaired Student's t-test. **f, g.** Two-way ANOVA followed by Tukey's multiple comparisons test ** P = 0.01 and **** P= 0.0001. DHE; Dihydroethidium. NAC; N-acetylcysteine.

Extended Data Fig 2. Mechanical changes in the intestine are sensed by tracheal Piezo. a. Representative confocal images of trachea (*QF>mtdTomato*; green pseudocoloured) and Piezo (magenta) in in sucrose and *Pe* fed flies. Scale bar, 20 microns. **b**. Quantification of tracheal Piezo expression relative to tracheal coverage in midguts as in a (n=27,29). **c**. Quantification of tracheal coverage in midguts from sucrose and *Pe* fed flies with and without treatment with the anti-oxidant NAC (n=12,23,9,21). **d.** Representative confocal images of *piezo* expression (magenta) in midguts from sucrose and *Pe* fed flies with and without treatment with the anti-oxidant NAC. Scale bar, 20 microns. **e.** Quantification of *piezo* expression relative to tracheal coverage in midguts as in d (n=12,23,9,21). **f**. Confocal images of

trachea (*QF>mtdTomato*; green pseudocoloured) and *piezo* expression (magenta) in the posterior midgut after feeding with normal food (NF) or methylcellulose (MC). Scale bar, 20 microns. **g**. Quantification of tracheal *piezo* expression relative to tracheal coverage in midguts as in f (n= 6, 8). **h**. Confocal images of midgut associated trachea (*dSRF^{ts}>gfp*; green) following sucrose or Yoda1 feeding. Scale bar, 20 microns. **i**. Confocal images of the intestinal epithelium (anti-Dlg; red) of control flies or flies overexpressing Piezo RNAi (*piezo-IR1*) in tracheal cells, upon sucrose or *Pe* feeding. DAPI (blue). Scale bar, 20 microns. **i'**. Representative images of eccentricity in midguts as in i. **j**. Quantification of eccentricity in midguts as in i' (n=7,8,9,7). **k**. Still frames of *in vivo* live imaging of trachea (*QF>mtdTomato*; red) and GCamp6s calcium reporter (green) in sucrose or *Pe* fed control animals or upon tracheal knockdown of *piezo* (*piezo-IR1*) and Pe feeding. Scale bar, 20 microns. Data represent the mean □±□ s.e.m. n numbers indicated for each graph are listed from left to right and they correspond to the numbers of guts analysed. <u>Statistics</u>: **b,g.** Two-tailed unpaired Student's t-test. **c, e, j.** Two-way ANOVA followed by Tukey's multiple comparisons test *P=0.05, ** P = 0.01, *** P=0.001 and **** P= 0.0001. DAPI, 4,6-diamidino-2-phenylindole. NAC; N-acetylcysteine.

Extended Data Fig 3. Tracheal Btl is not regulated by Piezo. a. Representative confocal images of gut trachea (*dSRF^{ts}>gfp*; green) and *btl-lacZ* (grey) upon sucrose or *Pe* feeding in control midguts or following tracheal *piezo* knockdown (*piezo-IR1*). Scale bar, 20 microns. **b**. Quantification of average *btl-lacZ* expression per TTC in midguts as in a (n=15,10,14,18). Boxed areas indicate regions selected for magnified view, which are presented in insets witin the grey scale panels. Data represent the mean □±□ s.e.m. n numbers indicated for each graph are listed from left to right and they correspond to the numbers of guts analysed. <u>Statistics</u>: Two-tailed unpaired Student's t-test. *** P=0.001 and **** P= 0.0001. DAPI, 4,6-diamidino-2-phenylindole.

Extended Data Fig 4. The role of vascular Piezo is conserved in mammalian intestinal regeneration and *Drosophila* intestinal hyperplasia. a. Experimental design of mouse experiments. Mice were induced with 3mg Tamoxifen on two consecutive days by oral gavage and sacrificed at day 8 after induction. For the damage-induced regeneration model, mice were given 10 Gy whole body irradiation at day 5 after transgene induction and intestinal tissue sampled 72 hours later. **b**. Method used to segment and quantify vascular branching in whole mount

immunostaining of the intestinal vasculature using anti-Pecam-1 antibody. Confocal
images are projected and a Mexican hat filter is applied before skeletonization. Scale
bar, 20 microns. **c**. Whole mount immunostaining of anti-Pecam-1 (green) in crypt
villus images of *Pdgf-beta-CreER Piezo1*$^{wt/wt}$ and *Piezo1*$^{fl/fl}$ mice without or with
irradiation. **d**. Quantification of the number of vascular branches in non-irradiated
(n=6,5) and irradiated (n=6,6) mice. **e**. Immunostaining of Chromogranin A for
enteroendocrine cell detection in small intestinal samples from mice as in c. Scale
bar, 20 microns. **f**. Quantification of Chromogranin A cells per crypt villus axis in non-
irradiated (n=3,3) and irradiated (n=3,3) mice. **g**. Staining with Alcian blue for goblet
cell detection in intestinal samples from animals as in c. Scale bar, 20 microns. **h**.
Quantification of Alcian Blue cells per crypt villus axis in non-irradiated (n=3,3) and
irradiated mice (n=3,3). **i**. Confocal images of control or *Src* overexpressing midgut
stem progenitor cells (*esg-lexA*^{ts}; green) and surrounding tracheal tissue
(*QF>mtdTomato*; red). DAPi (blue). Scale bar, 20 microns. **j, k.** Quantification of ISC
proliferation (**j**) (n=14,13) and tracheal coverage (**K**) (n=19,23) in midguts as in i.
Data represent the mean ± s.e.m. n numbers indicated for each graph are listed
from left to right and they correspond to the numbers of guts analysed. <u>Statistics</u>:
Two-tailed unpaired Student's t-test. **** P= 0.0001. DAPI, 4,6-diamidino-2-
phenylindole.

1197

Supplementary Movie 1

Time-lapse *in vivo* live imaging of a control *Drosophila* posterior midgut from Sucrose
fed mated female fly. The general tracheal network was labelled with *Q6>mtdTomato*
(red) and the calcium reporter *GCamp6s* (green) is expressed under the control of
the tracheal driver *dSRF-gal4*. Fly was starved and then fed sucrose for 2 hours prior
to imaging for 3 hours.

Supplementary Movie 2

Time-lapse *in vivo* live imaging of a control *Drosophila* posterior midgut from *Pe* fed
mated female fly. The general tracheal network was labelled with *Q6>mtdTomato*
(red) and the calcium reporter *GCamp6s* (green) is expressed under the control of
the tracheal driver *dSRF-gal4*. Fly was starved and then fed *Pe* for 2 hours prior to
imaging for 3 hours.

Supplementary Movie 3

Time-lapse *in vivo* live imaging of a *Drosophila* posterior midgut from *Pe* fed mated

female fly. The general tracheal network was labelled with *Q6>mtdTomato* (red) and

the calcium reporter *GCamp6s* (green) and a *piezo-RNAi* (*piezo-IR1*) are expressed

under the control of the tracheal driver *dSRF-gal4*. Fly was starved and then fed *Pe*

for 2 hours prior to imaging for 3 hours.

1216

References

1.   Hageman JH, Heinz MC, Kretzschmar K, Van Der Vaart J, Clevers H,
Snippert HJG. Intestinal Regeneration: Regulation by the Microenvironment.
Developmental Cell. 2020;54(4):435-46.
2.   Sato T, Van Es JH, Snippert HJ, Stange DE, Vries RG, Van Den Born M, et al.
Paneth cells constitute the niche for Lgr5 stem cells in intestinal crypts. Nature.
2011;469(7330):415-8.
3.   Li H, Jasper H. Gastrointestinal stem cells in health and disease: from flies to
humans. Disease Models & Mechanisms. 2016;9(5):487-99.
4.   Barker N, Van Es JH, Kuipers J, Kujala P, Van Den Born M, Cozijnsen M, et al.
Identification of stem cells in small intestine and colon by marker gene Lgr5. Nature.
2007;449(7165):1003-7.
5.   Micchelli CA, Perrimon N. Evidence that stem cells reside in the adult
Drosophila midgut epithelium. Nature. 2006;439(7075):475-9.
6.   Ohlstein B, Spradling A. The adult Drosophila posterior midgut is maintained
by pluripotent stem cells. Nature. 2006;439(7075):470-4.
7.   Medina AB, Perochon J, Johnson C, Polcowñuk S, Tian Y, Yu Y, et al.
Neuroendocrine Control of Intestinal Regeneration Through the Vascular Niche
in*Drosophila*. 2024.
8.   Touhara KK, Rossen ND, Deng F, Castro J, Harrington AM, Chu T, et al.
Topological segregation of stress sensors along the gut crypt–villus axis. Nature.
2025.
9.   Miguel-Aliaga I, Jasper H, Lemaitre B. Anatomy and Physiology of the
Digestive Tract of *Drosophila melanogaster*. Genetics. 2018;210(2):357-96.
10.   Vereecke L, Beyaert R, van Loo G. Enterocyte death and intestinal barrier
maintenance in homeostasis and disease. Trends in Molecular Medicine.
2011;17(10):584-93.
11.   Ding B-S, Cao Z, Lis R, Nolan DJ, Guo P, Simons M, et al. Divergent
angiocrine signals from vascular niche balance liver regeneration and fibrosis.
Nature. 2014;505(7481):97-102.
12.   Rafii S, Butler JM, Ding B-S. Angiocrine functions of organ-specific endothelial
cells. Nature. 2016;529(7586):316-25.
13.   Rivas-Carrillo SD, Kanamune J, Iwanaga Y, Uemoto S, Daneri-Navarro A,
Rivas-Carrillo JD. Endothelial Cells Promote Pancreatic Stem Cell Activation During
Islet Regeneration in Mice. Transplantation Proceedings. 2011;43(9):3209-11.

14.   Perochon J, Yu Y, Aughey GN, Medina AB, Southall TD, Cordero JB. Dynamic
adult tracheal plasticity drives stem cell adaptation to changes in intestinal
homeostasis in Drosophila. Nature Cell Biology. 2021;23(5):485-96.
15.   Tamamouna V, Rahman MM, Petersson M, Charalambous I, Kux K, Mainor H,
et al. Remodelling of oxygen-transporting tracheoles drives intestinal regeneration
and tumorigenesis in Drosophila. Nature Cell Biology. 2021;23(5):497-510.
16.   Gerit, Jacobson J, Karl, Hudry B, Christo, Dormann D, et al. Neuronal Control
of Metabolism through Nutrient-Dependent Modulation of Tracheal Branching. Cell.
2014;156(1-2):69-83.
17.   Blackie L, Gaspar P, Mosleh S, Lushchak O, Kong L, Jin Y, et al. The sex of
organ geometry. Nature. 2024;630(8016):392-400.
18.   Kam CY, Singh ID, Gonzalez DG, Matte-Martone C, Solá P, Solanas G, et al.
Mechanisms of skin vascular maturation and maintenance captured by longitudinal
imaging of live mice. Cell. 2023;186(11):2345-60.e16.
19.   Kong W, Loison O, Chavadimane Shivakumar P, Chan EH, Saadaoui M,
Collinet C, et al. Experimental validation of force inference in epithelia from cell to
tissue scale. Scientific Reports. 2019;9(1).
20.   Ichijo R, Kabata M, Kidoya H, Muramatsu F, Ishibashi R, Abe K, et al.
Vasculature-driven stem cell population coordinates tissue scaling in dynamic organs.
Science Advances. 2021;7(7):eabd2575.
21.   Ghabrial A, Luschnig S, Metzstein MM, Krasnow MA. Branching
Morphogenesis of the Drosophila Tracheal System. Annual Review of Cell and
Developmental Biology. 2003;19(Volume 19, 2003):623-47.
22.   Centanin L, Gorr TA, Wappner P. Tracheal remodelling in response to hypoxia.
Journal of Insect Physiology. 2010;56(5):447-54.
23.   Eilken HM, Adams RH. Dynamics of endothelial cell behavior in sprouting
angiogenesis. Current Opinion in Cell Biology. 2010;22(5):617-25.
24.   Affolter M, Bellusci S, Itoh N, Shilo B, Thiery J-P, Werb Z. Tube or Not Tube.
Developmental Cell. 2003;4(1):11-8.
25.   Hayashi S, Kondo T. Development and Function of the *Drosophila* Tracheal
System. Genetics. 2018;209(2):367-80.
26.   Amcheslavsky A, Jiang J, Ip YT. Tissue Damage-Induced Intestinal Stem Cell
Division in Drosophila. Cell Stem Cell. 2009;4(1):49-61.
27.   Biteau B, Jasper H. EGF signaling regulates the proliferation of intestinal stem
cells in *Drosophila*. Development. 2011;138(6):1045-55.
28.   Farnsworth RH, Lackmann M, Achen MG, Stacker SA. Vascular remodeling in
cancer. Oncogene. 2014;33(27):3496-505.
29.   Osol G, Moore LG. Maternal Uterine Vascular Remodeling During Pregnancy.
Microcirculation. 2014;21(1):38-47.
30.   Spradley FT, Ge Y, Granger JP, Chade AR. Utero-placental vascular
remodeling during late gestation in Sprague-Dawley rats. Pregnancy Hypertension.
2020;20:36-43.
31.   Buchon N, Broderick NA, Kuraishi T, Lemaitre B. DrosophilaEGFR pathway
coordinates stem cell proliferation and gut remodeling following infection. BMC
Biology. 2010;8(1):152.
32.   Stringer C, Wang T, Michaelos M, Pachitariu M. Cellpose: a generalist
algorithm for cellular segmentation. Nature Methods. 2021;18(1):100-6.
33.   Bodenschatz JFE, Ajmail K, Skamrahl M, Vache M, Gottwald J, Nehls S, et al.
Epithelial cells sacrifice excess area to preserve fluidity in response to external
mechanical stress. Communications Biology. 2022;5(1):855.

34.   Rangamani P, Lipshtat A, Azeloglu Evren U, Calizo Rhodora C, Hu M,
Ghassemi S, et al. Decoding Information in Cell Shape. Cell. 2013;154(6):1356-69.
35.   He L, Si G, Huang J, Samuel ADT, Perrimon N. Mechanical regulation of
stem-cell differentiation by the stretch-activated Piezo channel. Nature.
2018;555(7694):103-6.
36.   Johansson J, Naszai M, Hodder MC, Pickering KA, Miller BW, Ridgway RA, et
al. RAL GTPases Drive Intestinal Stem Cell Function and Regeneration through
Internalization of WNT Signalosomes. Cell Stem Cell. 2019;24(4):592-607.e7.
37.   Buchon N, Osman D, David FPA, Fang HY, Boquete J-P, Deplancke B, et al.
Morphological and molecular characterization of adult midgut compartmentalization
in Drosophila. Cell Reports. 2013;3(5):1725-38.
38.   Marianes A, Spradling AC. Physiological and stem cell compartmentalization
within the Drosophila midgut. Elife. 2013;2:e00886.
39.   Feinberg EH, Vanhoven MK, Bendesky A, Wang G, Fetter RD, Shen K, et al.
GFP Reconstitution Across Synaptic Partners (GRASP) Defines Cell Contacts and
Synapses in Living Nervous Systems. Neuron. 2008;57(3):353-63.
40.   Lucy, Sarah, Li X, Bilder D. Altered Modes of Stem Cell Division Drive
Adaptive Intestinal Growth. Cell. 2011;147(3):603-14.
41.   Schott S, Ambrosini A, Barbaste A, Benassayag C, Gracia M, Proag A, et al. A
fluorescent toolkit for spatiotemporal tracking of apoptotic cells in living *Drosophila*
tissues. Development. 2017;144(20):3840-6.
42.   Fischer KS, Henn D, Zhao ET, Sivaraj D, Litmanovich B, Hahn WW, et al.
Elevated Shear Stress Modulates Heterogenous Cellular Subpopulations to Induce
Vascular Remodeling. Tissue Engineering Part A. 2024;30(23-24):752-65.
43.   Sun Y, Liu J, Xu Z, Lin X, Zhang X, Li L, et al. Matrix stiffness regulates
myocardial differentiation of human umbilical cord mesenchymal stem cells. Aging.
2021;13(2):2231-50.
44.   Zanotelli MR, Reinhart-King CA. Mechanical Forces in Tumor Angiogenesis.
Springer International Publishing; 2018. p. 91-112.
45.   Negri S, Faris P, Berra-Romani R, Guerra G, Moccia F. Endothelial Transient
Receptor Potential Channels and Vascular Remodeling: Extracellular Ca2 + Entry for
Angiogenesis, Arteriogenesis and Vasculogenesis. Frontiers in Physiology. 2020;10.
46.   Coste B, Mathur J, Schmidt M, Earley TJ, Ranade S, Petrus MJ, et al. Piezo1
and Piezo2 Are Essential Components of Distinct Mechanically Activated Cation
Channels. Science. 2010;330(6000):55-60.
47.   Ranade SS, Qiu Z, Woo S-H, Hur SS, Murthy SE, Cahalan SM, et al. Piezo1,
a mechanically activated ion channel, is required for vascular development in mice.
Proceedings of the National Academy of Sciences. 2014;111(28):10347-52.
48.   Baghdadi MB, Houtekamer RM, Perrin L, Rao-Bhatia A, Whelen M, Decker L,
et al. PIEZO-dependent mechanosensing is essential for intestinal stem cell fate
decision and maintenance. Science. 2024;386(6725):eadj7615.
49.   Kim SE, Coste B, Chadha A, Cook B, Patapoutian A. The role of Drosophila
Piezo in mechanical nociception. Nature. 2012;483(7388):209-12.
50.   Zechini L, Amato C, Scopelliti A, Wood W. Piezo acts as a molecular brake on
wound closure to ensure effective inflammation and maintenance of epithelial
integrity. Current Biology. 2022;32(16):3584-92.e4.
51.   Nagarkar-Jaiswal S, Lee P-T, Campbell ME, Chen K, Anguiano-Zarate S,
Cantu Gutierrez M, et al. A library of MiMICs allows tagging of genes and reversible,
spatial and temporal knockdown of proteins in Drosophila. eLife. 2015;4:e05338.

52. Yang Y, Wang D, Zhang C, Yang W, Li C, Gao Z, et al. Piezo1 mediates
endothelial atherogenic inflammatory responses via regulation of YAP/TAZ activation.
Human Cell. 2022;35(1):51-62.
53. Lan Y, Lu J, Zhang S, Jie C, Chen C, Xiao C, et al. Piezo1‐Mediated
Mechanotransduction Contributes to Disturbed Flow‐Induced Atherosclerotic
Endothelial Inflammation. Journal of the American Heart Association.
2024;13(21):e035558.
54. Mao J, Yang R, Yuan P, Wu F, Wei Y, Nie Y, et al. Different stimuli induce
endothelial dysfunction and promote atherosclerosis through the Piezo1/YAP
signaling axis. Archives of Biochemistry and Biophysics. 2023;747:109755.
55. Niec RE, Chu T, Schernthanner M, Gur-Cohen S, Hidalgo L, Pasolli HA, et al.
Lymphatics act as a signaling hub to regulate intestinal stem cell activity. Cell Stem
Cell. 2022;29(7):1067-82.e18.
56. Wang P, Kljavin N, Nguyen TTT, Storm EE, Marsh B, Jiang J, et al. Adrenergic
nerves regulate intestinal regeneration through IL-22 signaling from type 3 innate
lymphoid cells. Cell Stem Cell. 2023;30(9):1166-78.e8.
57. Bernier-Latmani J, Petrova TV. Intestinal lymphatic vasculature: structure,
mechanisms and functions. Nature Reviews Gastroenterology & Hepatology.
2017;14(9):510-26.
58. Bernier-Latmani J, González-Loyola A, Petrova TV. Mechanisms and
functions of intestinal vascular specialization. Journal of Experimental Medicine.
2024;221(1).
59. Stzepourginski I, Nigro G, Jacob J-M, Dulauroy S, Sansonetti PJ, Eberl G, et
al. CD34 $^+$ mesenchymal cells are a major component of the intestinal stem cells
niche at homeostasis and after injury. Proceedings of the National Academy of
Sciences. 2017;114(4):E506-E13.
60. McCarthy N, Manieri E, Storm EE, Saadatpour A, Luoma AM, Kapoor VN, et
al. Distinct Mesenchymal Cell Populations Generate the Essential Intestinal BMP
Signaling Gradient. Cell Stem Cell. 2020;26(3):391-402.e5.
61. Kim J-E, Fei L, Yin W-C, Coquenlorge S, Rao-Bhatia A, Zhang X, et al. Single
cell and genetic analyses reveal conserved populations and signaling mechanisms
of gastrointestinal stromal niches. Nature Communications. 2020;11(1).
62. Palikuqi B, Rispal J, Reyes EA, Vaka D, Boffelli D, Klein O. Lymphangiocrine
signals are required for proper intestinal repair after cytotoxic injury. Cell Stem Cell.
2022;29(8):1262-72.e5.
63. Claxton S, Kostourou V, Jadeja S, Chambon P, Hodivala-Dilke K, Fruttiger M.
Efficient, inducible Cre-recombinase activation in vascular endothelium. genesis.
2008;46(2):74-80.
64. Bernier-Latmani J, Petrova TV. High-resolution 3D analysis of mouse small-
intestinal stroma. Nature Protocols. 2016;11(9):1617-29.
65. Blache P, Van De Wetering M, Duluc I, Domon C, Berta P, Freund J-NL, et al.
SOX9 is an intestine crypt transcription factor, is regulated by the Wnt pathway, and
represses the *CDX2* and *MUC2* genes. The Journal of Cell Biology. 2004;166(1):37-
47.
66. Liu L, Rao JN, Zou T, Xiao L, Smith A, Zhuang R, et al. Activation of Wnt3a
signaling stimulates intestinal epithelial repair by promoting c-Myc-regulated gene
expression. American Journal of Physiology-Cell Physiology. 2012;302(1):C277-C85.
67. Cordero JB, Ridgway RA, Valeri N, Nixon C, Frame MC, Muller WJ, et al. c-
Src drives intestinal regeneration and transformation. The EMBO Journal. 2014.

68. Chen P, Zhang G, Jiang S, Ning Y, Deng B, Pan X, et al. Mechanosensitive Piezo1 in endothelial cells promotes angiogenesis to support bone fracture repair. Cell Calcium. 2021;97:102431.

69. Kang H, Hong Z, Zhong M, Klomp J, Bayless KJ, Mehta D, et al. Piezo1 mediates angiogenesis through activation of MT1-MMP signaling. American Journal of Physiology-Cell Physiology. 2019;316(1):C92-C103.

70. Peralta M, Dupas A, Larnicol A, Lefebvre O, Goswami R, Stemmelen T, et al. Endothelial calcium firing mediates the extravasation of metastatic tumor cells. iScience. 2025;28(2):111690.

71. Boeldt DS, Bird IM. Vascular adaptation in pregnancy and endothelial dysfunction in preeclampsia. Journal of Endocrinology. 2017;232(1):R27-R44.

72. Guelfi S, Hodivala-Dilke K, Bergers G. Targeting the tumour vasculature: from vessel destruction to promotion. Nature Reviews Cancer. 2024;24(10):655-75.

73. Roux E, Bougaran P, Dufourcq P, Couffinhal T. Fluid Shear Stress Sensing by the Endothelial Layer. Frontiers in Physiology. 2020;11.

74. Fraisl P, Mazzone M, Schmidt T, Carmeliet P. Regulation of Angiogenesis by Oxygen and Metabolism. Developmental Cell. 2009;16(2):167-79.

75. Gervais L, Bardin AJ. Tracheal remodelling supports stem cells. Nature Cell Biology. 2021;23(6):580-2.

76. Fortuño A, José GS, Moreno MU, Díez J, Zalba G. Oxidative stress and vascular remodelling. Experimental Physiology. 2005;90(4):457-62.

77. Centanin L, Dekanty A, Romero N, Irisarri M, Gorr TA, Wappner P. Cell Autonomy of HIF Effects in Drosophila: Tracheal Cells Sense Hypoxia and Induce Terminal Branch Sprouting. Developmental Cell. 2008;14(4):547-58.

78. Mukherjee P, El-Abbadi MM, Kasperzyk JL, Ranes MK, Seyfried TN. Dietary restriction reduces angiogenesis and growth in an orthotopic mouse brain tumour model. British Journal of Cancer. 2002;86(10):1615-21.

79. Matamoro-Vidal A, Cumming T, Davidović A, Levillayer F, Levayer R. Patterned apoptosis has an instructive role for local growth and tissue shape regulation in a fast-growing epithelium. Current Biology. 2024;34(2):376-88.e7.

80. Roellig D, Theis S, Proag A, Allio G, Bénazéraf B, Gros J, et al. Force-generating apoptotic cells orchestrate avian neural tube bending. Developmental Cell. 2022;57(6):707-18.e6.

81. Kawaue T, Yow I, Pan Y, Le AP, Lou Y, Loberas M, et al. Inhomogeneous mechanotransduction defines the spatial pattern of apoptosis-induced compensatory proliferation. Developmental Cell. 2023;58(4):267-77.e5.

82. Yui S, Azzolin L, Maimets M, Pedersen MT, Fordham RP, Hansen SL, et al. YAP/TAZ-Dependent Reprogramming of Colonic Epithelium Links ECM Remodeling to Tissue Regeneration. Cell Stem Cell. 2018;22(1):35-49.e7.

83. Ciccone G, Azevedo Gonzalez Oliva M, Antonovaite N, Lüchtefeld I, Salmeron-Sanchez M, Vassalli M. Experimental and Data Analysis Workflow for Soft Matter Nanoindentation. Journal of Visualized Experiments. 2022(179).

84. Marchetti M, Zhang C, Edgar BA. An improved organ explant culture method reveals stem cell lineage dynamics in the adult Drosophila intestine. eLife. 2022;11:e76010.

85. Martin JL, Sanders EN, Moreno-Roman P, Jaramillo Koyama LA, Balachandra S, Du X, et al. Long-term live imaging of the Drosophila adult midgut reveals real-time dynamics of division, differentiation and loss. eLife. 2018;7:e36248.

86.   Herbert S, Valon L, Mancini L, Dray N, Caldarelli P, Gros J, et al.
LocalZProjector and DeProj: a toolbox for local 2D projection and accurate
morphometrics of large 3D microscopy images. BMC Biology. 2021;19(1).
87.   Aigouy B, Umetsu D, Eaton S. Segmentation and Quantitative Analysis of
Epithelial Tissues. Springer New York; 2016. p. 227-39.
88.   Newton H, Wang YF, Camplese L, Mokochinski JB, Kramer HB, Brown AEX,
et al. Systemic muscle wasting and coordinated tumour response drive
tumourigenesis. Nat Commun. 2020;11(1):4653.
89.   Gervais L, Casanova J. The Drosophila homologue of SRF acts as a boosting
mechanism to sustain FGF-induced terminal branching in the tracheal system.
Development. 2011;138(7):1269-74.
90.   Goto S, Hayashi S. Proximal to distal cell communication in the Drosophila leg
provides a basis for an intercalary mechanism of limb patterning. Development.
1999;126(15):3407-13.
91.   Fletcher GC, Diaz-de-la-Loza MD, Borreguero-Muñoz N, Holder M, Aguilar-
Aragon M, Thompson BJ. Mechanical strain regulates the Hippo pathway in
Drosophila. Development. 2018;145(5).
92.   Roy S, Huang H, Liu S, Kornberg TB. Cytoneme-mediated contact-dependent
transport of the Drosophila decapentaplegic signaling protein. Science.
2014;343(6173):1244624.
93.   Yagi R, Mayer F, Basler K. Refined LexA transactivators and their use in
combination with the *Drosophila* Gal4 system. Proceedings of the National Academy
of Sciences. 2010;107(37):16166-71.

1471

1472

Acknowledgements:

We would like to thank Norbert Perrimon (Harvard), Irene Miguel-Aliaga (The Francis
Crick Institute), Shigeo Hayashi (Rinken Centre, Kobe, Japan), Chrysoula Pitsouli
(University of Cyprus), Thomas Kornberg and Wanpeng Wang (Kornberg Lab-UCSF),
Barry Denholm (University of Edinburgh), Barry Thompson (EMBL Australia), Gines
Morata (Autonomous University of Madrid), Lucy O'Brien (Stanford University),
Susumu Hirabayashi (Imperial College London), Magali Suzanne (CBI Toulouse),
Yanlan Mao (UCL) and Andrew Davidson (University of Glasgow) for generously
sharing flies and reagents, and Marcus Fruttiger (UCL) for sharing the *Pdgf-beta-*
Cre-GFP mice. We thank the Bloomington Drosophila Stock Centre; the Vienna
Drosophila Resource Center and the *Drosophila* Studies Hybridoma Bank for fly
stocks and antibodies. We would like to thank the Core Services and Advanced
Technologies at the CRUK Scotland Institute, which is core funded by Cancer
Research UK (A31287). With particular thanks to Leo Carlin, Claire Mitchell, and
Peter Thomason from the Beatson Advanced Imaging Resource for imaging support,
Derek Miller and Evarest Onwubiko from the Biological Service Unit for assistance
with animal whole body perfusions for whole mount intestinal sample preparations,
and Dimitris Athineos and Dale Watt for assistance with stock maintenance and back
crossing of the *Piezo1^{fl/fl}* mice. We are grateful to Jeremiah Bernier-Latmani and
Tatiana Petrova for their support and advice during intestinal vascular tissue
preparation and imaging. The manuscript was critically reviewed by Catherine
Winchester (CRUK Scotland Institute). We thank all members of the Cordero
laboratory for general advice on the project.

Data availability: All source data related to this study will be made freely available
through institutional public repositories and accessible via a DOI link at the time of
publication.

1499

1500 **Funding sources:** Work in the Cordero laboratory is funded through a Wellcome
1501 Trust and Royal Society Sir Henry Dale Fellowship (104103/Z/14/Z), Wellcome
1502 Senior Research Fellowship (223091/Z/21/Z) and core Institutional funds from CRUK
1503 to the CRUK Scotland Institute (A31287) to J.B.C., a Wellcome ISSF Returner
1504 Scheme (204820/Z/16/Z) and Tenovus (Tenovus/S22-28) Fellowships to J.P, and an

EPSRC/MVLS studentship to C.J. (A31GA-7101). The Blyth laboratory is supported by CRUK funding (A29799) to K.B. Work from the Vassalli laboratory is supported by Leverhulme Trust grant RPG-2022-324 and EPSRC grant Mechanomeds EP/X033554 to M.V.

Authors contributions:

J.A.P. Designed, performed and analysed experiments, including all the work on tissue segmentation and morphometric analysis, and mouse experiments.

J.P. Designed, performed and analysed experiments, including the study of signalling mechanisms downstream of Piezo. Contributed to study conceptualization, supervision and funding acquisition.

C.T.J. Designed, performed and analysed experiments involving the use of Chiaro nanoindenter, and the time course of Piezo expression and activation.

M.W. Provided training and assistance with nanoindentation data acquisition and analysis.

C. N. and M.H. Processed samples for mammalian histology.

A.B.C. Contributed to early efforts in live imaging.

Y.Y. Provided general technical assistance and generated the *lexAop-Src* line.

L.M and K.B. Provided support with the supervision and planning of the mouse work.

M.V. Planning, supervision and analysis of nanoindentation work and funding acquisition.

J.B.C. Study conceptualization, supervision, data analysis and funding acquisition.

J.A.P., J.P., C.T.J, and J.B.C., Wrote the manuscript with input from the rest of the authors.

Conflict of Interests: The authors declare no conflict of interests.

Phillips and Perochon et al. Figure 1

Phillips and Perochon et al. Figure 2

Phillips and Perochon et al. Figure 3

Phillips and Perochon et al. Figure 4

dSRF^{ts}> GFP

a Control *piezo-IR1*

GFP / Dapi   *bnl-LacZ*

Sucrose

Pe

b
- Sucrose
- *Pe*

TTC β-gal signal relative to background (AU)

****

****   ***

Control   *piezo-IR1*

c

GFP / Myc   Myc

Sucrose

Pe

d
- Sucrose
- *Pe*

TTC Myc intensity relative to background (AU)

****

****   ****

Control   *piezo-IR1*

dSRF>; QF6>mtdTomato

e
mtdTomato/   Dapi/Yki::GFP Yki::GFP
Dapi/Yki::GFP

Sucrose

DSS

Pe

Pe + piezo-IR1

f
TTC Yki::GFP (nuclear/cytoplasmic ratio)

***

Suc   DSS

g
- Sucrose
- *Pe*

TTC Yki::GFP (nuclear/cytoplasmic ratio)

****

****   ns

Control   *piezo-IR1*

Phillips and Perochon et al. Figure 5

Phillips and Perochon et al. Figure 6

Phillips and Perochon et al. Figure 7

Phillips and Perochon et al. Extended data Fig. 1

Phillips and Perochon et al. Extended data Fig. 2

Phillips and Perochon et al. Extended data Figure 3

a Tamoxifen 3mg O.G — Day 0, Day 1; 10 Gy Whole body Irradiation — Day 5; Sample — Day 8

b Subtract Background | Mexican Hat Filter | Skeletonize

Pdgf-B-Cre-GFP

c *Piezo1^{wt/wt}* | *Piezo1^{fl/fl}*
Non-irradiated
Irradiated 72hr 10Gy
Pecam-1

d Total vascular branches
Non-irradiated — ns — *Piezo1^{wt/wt}*, *Piezo1^{fl/fl}*
Irradiated — ns — *Piezo1^{wt/wt}*, *Piezo1^{fl/fl}*

e *Piezo1^{wt/wt}* | *Piezo1^{fl/fl}*
Non-irradiated
Irradiated 72hr 10Gy
Chromogranin A

f Average ChrA+ cells /crypt-villus
Non-irradiated — ns — *Piezo1^{wt/wt}*, *Piezo1^{fl/fl}*
Irradiated — ns — *Piezo1^{wt/wt}*, *Piezo1^{fl/fl}*

g *Piezo1^{wt/wt}* | *Piezo1^{fl/fl}*
Non-irradiated
Irradiated 72hr 10Gy
Alcian Blue

h Alcian Blue+ cells /crypt villus
Non-irradiated — ns — *Piezo1^{wt/wt}*, *Piezo1^{fl/fl}*
Irradiated — ns — *Piezo1^{wt/wt}*, *Piezo1^{fl/fl}*

esg-lexA^{ts}>lexAop-GFP; QF6>mtdTomato

i Control / *Src*
Esg/Dapi | Esg/Dapi/QF6

j PH3+ cells post. midgut — Control, *Src* — ****

k Tracheal coverage post. midgut (AU) — Control, *Src* — ****

Phillips and Perochon et al. Extended data Fig. 4