Strain-specific structural variant landscapes shape mutation retention following mutagenesis in

Caenorhabditis elegans

R. Kapila[1,2*], S. Saber[1,2], R.K. Verma[1,2], G. Blanco[1,2], V.K. Eggers[1,2,] J.L. Fierst[1,2*]

[1]Department of Biological Sciences and [2]Biomolecular Sciences Institute

Florida International University, 11200 8th Street, Miami, 33199, FL, USA

*Corresponding author(s). E-mail(s): rkapila@fiu.edu; jfierst@fiu.edu

Author contributions

J.L.F. secured funding. R.K. and J.L.F. designed the experiments. R.K., with assistance from S.S.,

performed the fitness assays, and, with assistance from G.B., conducted the male frequency and

outcrossing rate assays. V.K.E. carried out the transposable element analysis. R.K. performed the

genomic analysis with guidance from J.L.F. R.K.V. developed the Python code for worm counting.

J.L.F. developed the theoretical model and conducted the population simulations. R.K. and J.L.F. co-

wrote the first draft. R.K. and J.F.K. prepared the final manuscript with contributions from V.K.E.

Acknowledgements

The authors would like to thank the members of the Fierst lab and Jessica Gonzalez, Jason Pienaar

and Jesualdo Fuentes-Gonzalez for suggestions and careful feedback. We also thank four anonymous

reviewers who provided thorough review and greatly improved the work.

Competing Interests:

The authors declare no competing financial or non-financial interests.

Classification

Keywords: Mutations, Outcrossing, Structural variations, Genome, *Caenorhabditis elegans*

This PDF file includes

Main Text

Figures 1 to 7

Abstract

Classical mutational theories centered on single nucleotide polymorphisms suggest that outcrossing

enhances the purging of deleterious mutations by promoting recombination. However, larger

structural variants, such as insertions, deletions, and inversions, can suppress recombination and

create linkage blocks. Using experimental evolution and whole-genome long- and short-read

sequencing, we characterized structural and nucleotide mutation landscapes in three *Caenorhabditis*

elegans strains following repeated mutagen exposure and recovery. We found substantial strain-

specific differences in structural variant accumulation and mutation retention. The strain with the

highest outcrossing propensity exhibited the greatest structural variant burden and a higher fraction

of single nucleotide polymorphisms within structural variant intervals. Consistent with this pattern,

our population genetic simulations showed that structural variants can persist more readily under

higher outcrossing rates. Together, these results indicate that structural variant architecture may

influence mutation retention dynamics and highlight strain-specific constraints on purging following

mutagenesis in *C. elegans*.

Introduction

The accumulation of deleterious mutations increases genetic load and can reduce population fitness,

potentially leading to long-term decline or extinction [1,2]. Mutations are broadly categorized into two

types: single nucleotide polymorphisms (SNPs) and structural variants (SVs). SNPs are single base

pair changes, resulting in localized effects on amino acids or regulatory elements[3]. In contrast, SVs

including deletions, duplications, inversions, and translocations, affect larger genomic regions and

can alter multiple genes or regulatory networks[4].

While mutation purging operates on both SNPs and SVs, the dynamics may differ because

the majority of SNPs may impact simpler genomic interactions[5]. In contrast, SVs can span larger

genomic regions and may suppress recombination, potentially complicating removal[6]. Evolutionary

theory has traditionally viewed the spectrum of genomic changes under the broad category of

"mutation", and much of the research on mutation purging has disproportionately focused on SNPs

due to their easier detection [7]. However, the advent of large-scale genome sequencing has revealed

the ubiquity of SVs. Despite the foundational role of mutation purging in evolutionary theory[8–10],

how this process operates on SVs remains poorly understood. Addressing this gap is essential for

refining theoretical models and understanding how mutation size and severity influence

evolutionary dynamics.

Mating systems influence mutation purging through genetic diversity, recombination, and

expression patterns of deleterious mutations[11,12]. Population genetics theory predicts that

outcrossing populations, through recombination, create novel genetic combinations able to purge

deleterious variation[13]. However, outcrossing can also potentially dilute the effects of deleterious

mutations and slow purging[14]. In contrast, reduced recombination and genetic diversity in self-

fertilizing populations may allow mildly deleterious mutations to accumulate through genetic drift[15].

At the same time, increased homozygosity under selfing can expose harmful recessive mutations to

selection and enabling more effective purging[15]. These processes may differ for mutations of

different physical sizes asSVs can suppress recombination creating linkage blocks limiting the

removal of deleterious variation[16,17]. Consequently, the interaction between mating system,

recombination suppression by SVs, and mutation purging remains poorly understood. This

complexity is further influenced by transposable elements (TEs), a dynamic source of genomic

variation [18,19]. TEs can generate SVs through transposition or ectopic recombination and may

respond to mutagenic or environmental stress[20]. Mating systems may shape TE dynamics:

outcrossing can facilitate TE spread but also introduce recombination that helps remove them,

whereas selfing can promote TE accumulation through reduced recombination or enhance TE

elimination through increased homozygosity [21,22]. Despite their role in genome evolution, how TEs

contribute to mutational landscapes following acute mutagenic stress is also poorly understood.

In this study we use experimental evolution to examine whether strains differing in

outcrossing propensity differ in post-mutagenesis mutation retention, and whether these patterns

vary with mutation size and genomic context. Specifically, we ask whether strains differing in male

frequency and outcrossing rates exhibit distinct patterns of SV accumulation, SNP localization, and

mutation retention following repeated mutagen exposure and recovery. By comparing three

genetically distinct isolates (N2, AB1, and CB4856), we evaluate whether mutation size and genomic

context contribute to strain-specific differences in the persistence of induced mutations. C. elegans

are androdiecious with populations consisting of both males and self-fertile hermaphrodites. The

frequency of males and outcrossing varies drastically across different strains [23–25] making C. elegans

an excellent system to test these ideas. Previous experimental evolution studies in C. elegans show

that increased mutational input can transiently elevate male frequency and outcrossing, yet does

not consistently enhance purging of deleterious variation, with outcomes strongly dependent on

genetic background [26–29]. Comparative genomic analyses across Caenorhabditis species further

demonstrate that mating system differences shape genome-wide patterns of genetic variation, with

self-fertilizing species such as C. elegans typically exhibiting reduced nucleotide diversity and

stronger linkage among mutations than outcrossing relatives such as *C. remanei* [30,31]. Together, these
studies indicate that reproductive mode can strongly influence mutation dynamics across genomes.
However, how these processes affect the retention of different mutation classes—particularly
structural variants—following acute mutagenic stress remains largely unexplored.
100
To test whether strain-specific differences in mating behavior and genomic background
influence mutation retention following mutagenesis, we hypothesized that strains differing in
outcrossing propensity would exhibit distinct patterns of SNP and SV retention after recovery. We
subjected *C. elegans* strains to two mutagens, EMS to induce single nucleotide mutations and
formaldehyde to introduce complex lesions in the genome[32]. Populations were mutagenized for five
consecutive generations and allowed to recover for three generations. We then tracked relative
fitness, changes in male frequency, and outcrossing rates to quantify phenotypic response and
investigate patterns of mutation retention after mutagen exposure and recovery. We employed
both Pacific Biosystems HIFI long-read and Illumina short-read sequencing to quantify the size,
location, and types of mutations that persist after recovery.
Results
Strain-specific phenotypic and genomic responses to mutagenesis
To evaluate whether variation in outcrossing propensity is associated with differences in mutation
retention following mutagenesis, we subjected three strains of *C. elegans* (N2 (Bristol), AB1, and
CB4856 (Hawaiian)), to five generations of mutagenesis followed by three generations of recovery,
with four replicate populations for each experimental treatment (12 in total).
117
Male frequency was differentially affected by mutagenesis
Male frequency determines the opportunity for outcrossing in *C. elegans* [24,25] and changes in male
production indicate how mating system dynamics respond to mutagenic stress. Non-mutagenized
CB4856 populations had the highest male frequencies (Figure 1A), significantly greater than either
AB1 ($p < 0.0001$) or N2 ($p < 0.01$). CB4856 male frequency was reduced after mutagenesis with EMS
($p = 0.03$; Figure 1A) and formaldehyde, although this comparison was not statistically significant (p
$= 0.11$). N2 male frequency did not change in response to either mutagen ($p = 1.00$ for both).
Similarly, AB1 male frequencies did not differ across populations or treatments (all p-values were >
0.05; Supplementary Table 1; Figure 1A).
Outcrossing frequencies varied between C. elegans strains
To determine whether mutagenesis altered mating behavior among strains with different male
frequencies, we quantified outcrossing frequency using paired mating assays. Cross-fertilization in *C.*

elegans produces approximately 50% male progeny while, selfing hermaphrodites produce few or
no males. We used male frequency as a proxy for determinig whether outcrossing occurred[33]. In
each assay, a single male was paired with a single hermaphrodite, and lifetime male production
recorded. Each plate was classified "outcrossing" or "selfing-dominated" using strain-specific male-
frequency thresholds established a priori. N2 and AB1 plates with male frequencies >5% were
classified as outcrossing. CB4856 exhibited elevated baseline male frequencies and plates were
classified as "outcrossing" when male frequency ≥25%. Outcrossing frequency was calculated as the
proportion of plates meeting these criteria within each strain and treatment. Thus, outcrossing
frequency reflects the probability of successful mating rather than mating efficiency. CB4856
populations had the highest outcrossing frequencies under all conditions, (Figure 1B). In contrast, N2
populations displayed the lowest outcrossing frequencies. AB1 populations were intermediate to
these.

Relative fitness did not differ significantly from parental levels following recovery

Relative fitness, defined here as competitive performance against a GFP-marked reference strain,
quantified from relative abundances after ~8 days of co-culture. Across all strains, recovered
populations were statistically indistinguishable from their non-mutagenized ancestors indicating
convergence toward baseline fitness (Figure 1C; Supplementary Table 2). CB4856 relative fitness did
not differ significantly between non-mutagenized, mutagenized, or recovered populations under
either mutagen (all pairwise comparisons p ≥ 0.18). AB1 showed no significant fitnesss differences
following either formaldehyde (p = 0.94) or EMS exposure (p = 0.76), although recovered EMS-
mutagenized AB1 populations exhibited a modest fitness increase relative to during mutagenesis (p
= 0.022). N2 populations exhibited a transient response to formaldehyde exposure, with relative
fitness significantly higher immediately after mutagenesis compared to non-mutagenized controls (p
= 0.026), an effect not observed under EMS treatment (p = 0.46). After recovery, N2 fitness was
statistically indistinguishable from the parental baseline (Recovered vs. Non-mutagenized, p ≥ 0.99).

Numerous SVs and SNPs were retained after recovery from mutagenesis

Despite rapid fitness recovery, populations harboured numerous *de novo* mutations. CB4856
exhibited significantly higher susceptibility to mutagenesis compared to AB1. When mutagenized
with formaldehyde, CB4856 populations showed a 1.12-fold increase in *de novo* SNPs compared
with AB1 (t=5.55, *p*=0.0014) and comparable *de novo* SNP counts to N2 (t < 0.01, *p*= 0.996). In
contrast, with EMS as the mutagen, CB4856 showed a significantly higher *de novo* SNP count when
compared with AB1 (t = 6.79, p < 0.01) and a marginally higher *de novo* SNP count when compared
with N2 (t = 2.45, p = 0.05; Figure 2A).

Due to marked heteroscedasticity in SV counts we performed Mann–Whitney U tests to

statistically evaluate *de novo* SV counts. CB4856 populations had significantly higher *de novo* SV

counts (Figure 2A) when compared with AB1 (W=16, *p* = 0.03 for both mutagens) and N2

populations (W=16, *p* = 0.03 for both mutagens).

Across strains and treatments, most of the detected SVs were intrachromosomal insertions,

deletions, duplications and inversions. Interchromosomal translocations were just ~0.7–4.3% of all

identified SVs (Supplementary Table 3). Thus, most mutagen-induced SVs reflect within-

chromosome rearrangements rather than widespread between-chromosome exchanges.

CB4856 harbored high numbers of SNPs and SVs in exons

To evaluate whether mutation retention differed across functional genomic regions, we examined

the distribution of SNPs and SVs relative to annotated exons. CB4856 populations had the highest

number of mutations in coding regions (Figure 2B) and the highest number of SNPs located within

SVs (Figure 3A). CB4856 had 2.20 times more exonic SNPs than AB1 in formaldehyde-mutagenized

population (W=0, *p*= 0.03) and 1.12 times more than N2 (W=12, *p*=0.34). CB4856 had 2.26 times

more exonic SNPs compared to AB1 in EMS mutagenized population (W=0, *p*=0.03) and 1.38 times

more exonic SNPs compared to N2 (W=16, *p*=0.03) (Figure 2B).

Exons were similarly affected by SVs. AB1 populations had fewest exons containing SVs,

whereas CB4856 populations had the most number. AB1 formaldehyde populations had 16.25-fold

fewer exonic SVs than CB4856 (W = 0, p = 0.02) and 6.92-fold fewer than N2 (W = 16, p = 0.03).

CB4856 EMS populations had 22.31-fold more exonic SVs compared to AB1 (W = 0, p = 0.02), and N2

had 7.81-fold more when compared with AB1 (W = 16, p = 0.03; Figure 2B).

CB4856 had the highest number of SNPs located within SVs interval (77.13% for

formaldehyde and 82.46% for EMS treatment; Figure 3A). SVs in AB1 populations contained 6.36%

(formaldehyde) and 12.01% (EMS) of *de novo* SNPs. N2 populations had similarly low numbers of

SNPs contained in SVs (EMS 7.12%; formaldehyde 6.80%).

Non-random enrichment of SNPs within structural variant regions

SNP counts within SV intervals were significantly greater than random expectation (all p < 0.001;

Supplementary Table 4). Enrichment ranged from 2.6–3.0× in AB1, 2.6–4.9× in N2, and ~1.9× in

CB4856 (Supplementary Table 4). CB4856 exhibited substantially greater SV genome coverage (~40–

43%) (Figure 4) with SNPs significantly enriched within SV regions relative to genomic expectation.

Breakpoint microhomology differed among strains but did not explain SV burden

To test whether strain-specific differences in SV abundance reflect differences in DNA repair

signatures, we examined breakpoint microhomology across intrachromosomal SVs. Non-

Homologous End Joining (NHEJ) results in 0–2 bp mutational 'scars' while Microhomology-Mediated

End Joining (MMEJ) leaves larger 3–10 bp signatures. Homology-mediated repair such as Single-
Strand Annealing (SSA) results in ~11-50 bp mutations while mutations >50bp indicate different
mechanisms of Homology-Mediated Repair (HMR). Accordingly, we classified breakpoints into bins
of 0 bp (NHEJ) 1–2 bp (NHEJ/MMEJ), 3–10 bp (MMEJ), 11–50 bp (SSA), and >50 bp (HMR)
(Supplementary Table 5). NHEJ and NHEJ/MMEJ occurred at comparable frequencies across strains.
MMEJ and SSA tracts were enriched in AB1 while CB4856 and N2 were enriched for larger HMR
signatures. N2 had the highest enrichment of extended HMR mutations despite having an
intermediate overall SV burden, indicating that breakpoint architecture alone does not explain strain
differences in SV abundance.

Active transposable elements were responsible for a small portion of SV mutations

Because both mutagenesis and outcrossing can influence transposable element (TE) activity,
we evaluated whether TE mobilization contributed to SVs in mutagenized populations using the
TransposonUltimate pipeline [34].

Total repeat content comprised 23.21%, 25.74%, and 21.70% of the N2, AB1, and CB4856
reference genomes, respectively. Class II DNA transposons represented ~94% of annotated TEs in
each genome (Figure 3B). Zator elements were the most abundant, accounting for ~46% of DNA TEs.
AB1 contained a smaller proportion of hAT elements but a higher proportion of Sola and MITE
elements relative to N2 and CB4856. Tc1-Mariner (9%), CMC (13%), Helitrons (0.5%), and Novosib
(0.3%) elements occurred at similar frequencies across genomes. As expected for *C. elegans*, TE
density followed the characteristic chromosome arm–center pattern, with higher TE abundance on
chromosome arms (Figure 5).

To assess TE mobilization after mutagenesis, we intersected reference TE annotations with
de novo SV calls using the TEdeTEct pipeline implemented in TransposonUltimate. Only a small
fraction of annotated TEs showed evidence of transposition: 0.07–0.08% in N2, 0.04–0.05% in AB1,
and 0.24–0.27% in CB4856. Consequently, TE-associated events accounted for <15% of all de novo
SVs. Despite having the lowest genomic TE proportion, CB4856 exhibited ~4× more transposition
events than N2 or AB1 (Supplementary Table 6).

TE activity was broadly similar between EMS and formaldehyde treatments. In N2, 6 of 21
(29%) transposition events occurred under both mutagens. CB4856 and AB1 showed 12 of 67 (18%)
and 1 of 22 (4.5%) shared events between treatments, respectively, and no identical transposition
events were observed across strains. Zator elements accounted for ~55% of all detected
transposition events (Figure 3B). Together, these results indicate that TE activity contributes to the
formation of de novo SVs but, it accounts for a minority of the SVs observed following mutagenesis.

Mutagenesis induced large SV mutations

Although most SVs were small (<10⁴ bp), both formaldehyde and EMS generated large-scale

mutations affecting substantial portions of the genome (Supplementary Figure 1). Across all strains

and treatments, SVs larger than 10 kb accounted for only ~1.4% of detected events (Supplementary

Table 7), yet these included several megabase-scale inversions capable of reshaping large genomic

regions (Figure 4).

In the N2 formaldehyde-treated population, insertion added 471,238 nucleotides (~0.48% of

the 100.8 Mb *C. elegans* genome). In comparison, 74,044 nucleotides (0.06%) were deleted, 44,318

nucleotides (0.04%) duplicated, and 1,289,368 nucleotides (1.29%) affected by inversions. Mutations

in the EMS-mutagenized N2 populations were of similar scale (Supplementary Table 8). Large

inversion events were also detected in N2 populations, including a 1.09 Mb inversion on

chromosome IV in formaldehyde-treated populations and a 2.49 Mb inversion on chromosome X in

EMS-treated populations (Supplementary Table 9).

AB1 populations had the smallest number of nucleotides affected by SV mutations. In

formaldehyde-treated AB1, 35,468 nucleotides were inserted and 46,363 deleted. The notable

exception was inversion, which spanned 2,539,375 nucleotides in formaldehyde-treated and

4,181,603 nucleotides in EMS-treated populations. Some transposition events were shared between

mutagen treatments. For example, a 2.54 Mb inversion on the right arm of chromosome 5 occurred

in both AB1 Formaldehyde and EMS-mutagenized populations. AB1 EMS-mutagenized populations

also experienced a 1.56 Mb inversion on the left arm of chromosome 4 (Supplementary Table 9).

CB4856 populations harbored substantially more DNA affected by SVs (Figure 4;

Supplementary Table 3). Roughly 1 Mb of new DNA was inserted under both treatments

(Supplementary Table 3). Compared to AB1, CB4856 had 14.11-fold more insertions under

formaldehyde and 15.18-fold more under EMS. Relative to N2, it had 1.50-fold and 1.66-fold more

insertions under formaldehyde and EMS, respectively. CB4856 also contained large amounts of

duplicated DNA, totaling 1.7 Mb under formaldehyde and 21.7 Mb under EMS, largely driven by a

12.31 Mb duplication on chromosome IV and an 8.82 Mb duplication on chromosome I. Inversions

represented the largest class of structural change, spanning 49.7 Mb—nearly half the genome—in

formaldehyde-treated CB4856 populations, including 13.91 Mb and 7.1 Mb inversions on

chromosome V, an 11.24 Mb inversion on chromosome III, and 10.5 Mb and 6.29 Mb inversions on

chromosome II. In EMS-treated CB4856 populations, inversions spanned 32.8 Mb—nearly one-third

of the genome—including large inversions on chromosomes II, I, and V (Supplementary Table 9).

When SVs larger than 10 kb were excluded, the remaining mutations followed the

characteristic arms–center pattern of the *C. elegans* genome, with more nucleotides affected on

chromosome arms than in centers (Figure 5), consistent with the recombination landscape where

recombination is elevated on arms and suppressed in central regions. However, the presence of

multiple megabase-scale SVs indicates that mutagenesis generates structural changes capable of

influencing recombination and mutation linkage across large genomic regions.

SVs and transposon activity varied across chromosomes

The distribution of SVs varied across chromosomes, strains, and mutagens (Figure 6A). SV mutations

were more frequent on Chromosomes IV and V compared with Chromosome X. TE activity did not

explain these patterns; for example, TEs were most active on Chromosome IV in CB4856, whereas

Chromosome V harbored more SVs (Figure 6B). TE-associated duplications and inversions were less

frequent than deletions across all populations. These patterns indicate that TE activity alone does

not account for the chromosomal distribution of SVs.

Simulation support of experimental results

Although there is a body of work modelling the influence of outcrossing on mutation in populations

these models typically assume that mutations are occurring at the level of an individual SNP or

broader, abstract 'allele' level[35–37]. To explore whether our experimental results could emerge under

different evolutionary scenarios, we developed a population model describing evolution of SVs in

the genome. We implemented this model in stochastic simulations using the SLiM 4.3 software[38] and

measured the impact of outcrossing rates on SVs in the population.

Consistent with our experimental observations, the simulations showed that outcrossing

populations retained SVs, particularly when outcrossing rates consistently exceeded 30%. In

contrast, fully selfing populations rapidly purged SVs. SVs were retained in outcrossing populations

when each mutation carried a small deleterious effect on fitness, drawn from a gamma

distribution[39–41] (Supplementary Figure 2). The fitness landscape had a negative curvature or

synergistic epistasis (Supplementary Figure 3). Under these conditions, SVs persisted in simulated

populations. These results suggest that outcrossing can, under specific conditions, promote the

persistence of structural mutations (Figure 7).

Discussion

In this study we compared genomic responses to repeated mutagen exposure and recovery in three

genetically distinct *C. elegans* isolates. Integrating experimental evolution, whole-genome

sequencing, and population genetic simulations revealed strain-specific differences in mutation

retention associated with mating behavior and genomic background. CB4856, which exhibited the

highest male frequency and outcrossing rate, retained large numbers of mutations despite

recovering relative fitness after mutagenesis. Notably, many SNPs occurred within SV intervals and

frequently overlapped exonic regions. Although SNP enrichment within SV regions occurred in all

strains, the absolute number was far greater in CB4856, consistent with genome wide SVs coverage
suggesting that genome architecture influences mutation retention following mutagenesis.

C. elegans has been widely used to study mutation accumulation and genomic responses to
mutational stress through both mutation-accumulation experiments and analyses of naturally
occurring variation [42,43]. Experimental evolution studies show that increased mutational input can
elevate male frequency and outcrossing rates, likely as short-term responses to mutational load [26,29].
However, elevated outcrossing does not consistently result in sustained purging of deleterious
mutations, and natural variation among isolates highlights strong genetic background effects [26,28,29].
Our results extend these findings by showing that strain-specific genomic architecture can influence
genome-wide patterns of mutation retention following mutagenesis.

We explored these factors in our population simulations and found that under specific
assumptions increased levels of outcrossing would retain SVs. Importantly, the phenotypic effect
size of the segregating mutations was very small. In a fully outcrossing simulated population, there
was an average of 10 mutations per individual after 100,000 generations, each with an average
phenotypic effect size of 0.0035. Larger mutation effect sizes were rapidly purged from outcrossing
populations. However, deleterious mutations with small effects on the phenotype and fitness were
stably maintained in the simulated outcrossing populations for long periods. Because mating system
was explicitly manipulated in the simulations but not in the experimental populations, we interpret
the simulations as providing mechanistic support for (rather than direct validation of) the empirical
associations observed among strains.

Genomic responses to mutational stress depend in part on the type of DNA lesion and the
repair mechanisms that resolve it. Point mutations are typically repaired through efficient pathways
like base excision repair (BER) or nucleotide excision repair (NER), while complex DNA lesions
generating double-strand breaks must be resolved through pathways such as homologous
recombination (HR), classical non-homologous end joining (NHEJ), and microhomology-mediated
end joining (MMEJ), which can produce SVs during repair[44]. We expected that EMS should create
more SNPs while formaldehyde would create more SVs in the genome. However, we found that 5
consecutive generations of treatment with either chemical had similar mutational effects on the *C.*
elegans genome. Breakpoint microhomology analysis revealed that homology patterns differed
among strains but did not correspond directly with SV abundance (Supplementary Material 7). In
particular, CB4856 exhibited the highest SV burden despite displaying breakpoint signatures broadly
similar to those observed in N2. These results suggest that differences in SV abundance are unlikely
to arise solely from variation in repair pathway usage and may instead reflect differences in genome
architecture.

SV in *Caenorhabditis* is dominated by intrachromosomal rearrangements, whereas

interchromosomal translocations are rare[46]. Recent mutation-accumulation studies provide

benchmarks for spontaneous SVs in *C. elegans*[47], which occur at low frequency and are dominated

by insertions and deletions. Consistent with this pattern, the SVs detected following mutagenesis in

our study were overwhelmingly intrachromosomal across all strains and treatments (Supplementary

Table 4). Comparisons between the CB4856 and N2 reference genomes have used long-read DNA

sequencing to reveal extensive structural divergence at the same time as chromosome-scale synteny

is preserved[48,49], analysis of the non-mutagenized populations mirrored this pattern, with CB4856

harboring substantially greater standing SV variation than N2. The mutagen-induced SVs we

observed exhibited similar qualitative properties: across strains and treatments, SVs were primarily

insertions and deletions, intrachromosomal rearrangements predominated, and exon overlap

occurred at frequencies comparable to those reported for spontaneous structural mutations[47]

(Supplementary Tables 3–10). These similarities suggest that mutagenesis increases the frequency

and genomic impact of SVs without generating qualitatively different classes of rearrangements.

These strains also differ extensively at the nucleotide level, with hundreds of thousands of SNPs and

numerous short insertions and deletions distributed across the genome [49,50]. Because long-read

sequencing was performed at a single post-recovery time point, our data capture the net genomic

outcome of recovery from mutagenesis rather than directly measuring purging rates. Nevertheless,

all strains recovered relative fitness within three generations, indicating no irreversible fitness

decline. Differences in mutational load across strains could, in principle, reflect variation in initial

susceptibility to mutagenesis or DNA repair efficiency, in addition to differences in purging

dynamics. However, all strains recovered relative fitness within three generations following

mutagenesis, indicating that none suffered irreversible fitness impairment under these conditions.

Moreover, population genetic simulations that hold repair efficacy constant recapitulated the

empirical observation that higher outcrossing rates are sufficient to retain SV that would otherwise

be purged in predominantly selfing populations, even in the absence of strain-specific differences in

DNA repair.

AB1 exhibited relatively low numbers of SNPs and SVs, limited exonic overlap, and moderate

SNP enrichment within SV regions (~2.5–3× above expectation). Because SVs span a relatively small

fraction of the genome and contain comparatively few coding mutations, this configuration may

reduce functional disruption. In this context, SNP enrichment likely reflects localized clustering of

mutations within a comparatively constrained mutational landscape, potentially buffering AB1

against large-scale functional perturbation following mutagenesis.

In CB4856 populations, SVs spanned approximately 40–43% of the genome and contained
the highest absolute number of SNPs embedded within SV regions. Although enrichment relative to
genomic expectation (~1.9×) was lower than in N2, the overall burden of coupled structural and
nucleotide variation was substantially greater. Many of these variants intersected coding regions,
increasing the likelihood of functional disruption. SVs in CB4856 also spanned a broad size range
(10^3–10^7 bp; Supplementary Figure 1), generating a genome-wide mutational landscape that may
promote genomic plasticity but also increase susceptibility to instability under chemical
mutagenesis. Despite this elevated mutational burden, CB4856 populations recovered relative
fitness to levels indistinguishable from the ancestral baseline within three generations, indicating
that short-term fitness does not fully capture underlying genomic load.

N2 presents a contrasting genomic response. Although SVs span only a small proportion of
the genome (~1–3%), SNP enrichment within SV regions was among the highest observed (~2.6–4.9×
above expectations), indicating strong local clustering of nucleotide variation within structurally
altered segments despite limited overall SV coverage. Because the total number and genomic span
of SVs remain relatively constrained, the absolute burden of linked structural and nucleotide
variation is potentially lower than in CB4856. N2 populations also showed relatively stable male
frequencies and rapid fitness recovery, suggesting that SVs were spatially restricted. Interestingly,
relative fitness in N2 increased transiently following formaldehyde exposure before returning to
ancestral levels after recovery (Fig. 1B). This short-term increase may reflect consequences of long-
term laboratory domestication. Unlike the wild isolates AB1 and CB4856, N2 has experienced
decades of selection in stable laboratory environments[51], whereas CB4856 represents a genetically
diverged natural isolate[52]. The transient increase in relative fitness observed in N2 following
formaldehyde exposure may reflect a compensatory physiological response analogous to hormesis,
in which short-term stress induces beneficial physiological effects[53,54].

Our observation that SNPs are consistently enriched within (SV) regions across strains
suggests that nucleotide and structural mutations frequently occur within the same genomic
intervals. Such clustering can reduce the effective separation of linked mutations during
recombination, particularly when SVs span large genomic regions. These patterns provide a genomic
context for previous experimental evolution studies in *C. elegans*, which show that increased
mutational input can elevate male frequency and outcrossing but does not consistently enhance
purging of deleterious mutations [26,28,29]. Together, these results suggest that the physical
organization of mutations within genomes may influence how efficiently recombination can
separate and remove deleterious variants following mutagenic stress.

Transposable elements (TEs) can play a critical role in shaping genomic resilience by
introducing genetic variation and driving structural changes in response to stress[55]. Increased
outcrossing has been associated with elevated TE activity, likely due to the enhanced recombination
rates and reduced efficiency of silencing mechanisms in outcrossing populations[56,57]. Our results
show strain-specific differences in TE activity following mutagenesis. CB4856, which had the highest
outcrossing frequency, also exhibited the highest transposition rates, consistent with increased
genomic instability under mutagenic stress. In contrast, although AB1 had the highest number of
annotated transposons, its transposition rates were lower, suggesting fewer active TEs. N2 showed
the highest overlap in TE events between mutagens (~30%), suggesting common TE activations
under stress.

Unlike earlier studies focusing on the Tc1-Mariner family of TEs [58,59], we found that Zator
was the most active and abundant transposon across all three strains and both mutagens. This result
is consistent with recent work showing that few TEs remain active in *C. elegans*, most of which
belong to the Zator family[49,60]. The transposon landscape also showed the expected arm–center
pattern reported in previous studies, consistent with the known recombination landscape of *C.*
elegans, in which recombination is elevated on chromosome arms and reduced in chromosome
centers[61]. Together, these observations suggest that elevated TE activity in CB4856 may contribute
to its increased genomic instability under mutagenic stress. However, because TEs account for only a
small fraction of the total SVs in our dataset, TE mobilization is unlikely to explain the full extent of
strain differences in mutation retention.

In conclusion, we found strain specific fitness and mutation responses which may reflect
larger patterns of evolutionary response. Although the N2 "Bristol" strain, originally collected by
Sydney Brenner[51] has been developed as a model system, its fitness, behavior and genomics differ
substantially from other strains of *C. elegans*. N2 is a highly lab adapted strain with little genetic
variation, few males and low outcrossing frequencies [62]. In contrast, CB4856 and AB1 are more
recently collected "wild worms" with substantial genomic differences [50] and high natural variation in
male frequency and outcrossing [24,28]. Additionally, our analyses indicate that SNPs are significantly
enriched within SV regions relative to genomic expectations, suggesting that strain specific genomic
landscapes determine the region in which mutations accumulate and remain linked. Such clustering
may influence how mutational load is distributed across the genome and how efficiently outcrossing
and recombination can separate deleterious variants following mutagenic stress. However, because
these comparisons involve distinct genetic backgrounds, the present results do not establish a causal
relationship between outcrossing and SV persistence. Forward-time simulations were therefore used
to explore how variation in mating system parameters could influence SV persistence under

controlled conditions; these are presented as a mechanistic framework consistent with, but not
demonstrating, causality. Future studies manipulating mating system within a common genetic
background, for example using *him* mutants to increase male frequency or *mab* mutants to prevent
mating, together with strain-specific analyses using *fog* or *xol* backgrounds and additional mutagenic
treatments, will help clarify how mating dynamics influence mutation retention. Overall, our results
suggest that genome architecture and mating system jointly shape how populations respond to
mutational stress.
440
441
442
443 **Funding**
444 This work was supported by NSF award 2225796 and NIGMS award R35GM147245 to J.L.F.
445
446 **Data Availability**
447 Scripts, bioinformatic workflows and analyses are available at github.com/RohitKapila. Python code
448 used for image analysis is available at github.com/FierstLab/WormImage. Fitness data are available
449 as Supplemental Materials associated with the manuscript. Sequence reads are available at the NCBI
450 SRA under BioProject PRJNA1201181.
451
Methods
Strains and mutagens
Three *C. elegans* strains with varying male frequencies were chosen: N2 (Bristol), CB4856 (Hawaiian),
and AB1. ST2 served as a common competitor in relative fitness assays. All strains were obtained
from the Caenorhabditis Genetics Center and maintained frozen until the experiment. Frozen strains
were thawed on 75 mm × 13 mm NGM plates seeded with *E. coli* OP50. After one generation (~3.5
days at 20 °C) for recovery, each strain was divided into nine replicate populations. Prior to the
experiment, replicates were synchronized by bleaching. Over five generations, four replicates per
strain were mutagenized with formaldehyde, four with EMS, and one remained untreated as a non-
mutagenized control. All replicates for a given strain were handled in parallel on the same day using
the same batch of plates to minimize intra-strain variability due to plate effects. A single non-
mutagenized replicate was included per strain as a reference baseline rather than as a fully
replicated treatment. This design choice reflects the extremely low rates of spontaneous SNP and
structural variant (SV) accumulation in *C. elegans* (approximately 0.003–0.03 SVs per genome per
generation)[42,47]. Over the five-generation duration of this experiment, the expected number of

spontaneous de novo SVs is therefore near zero. Moreover, because genomic analyses were
performed using pooled long-read sequencing, any spontaneous variants arising in untreated
populations would be present at very low frequencies and fall below reliable detection thresholds.
Under these conditions, additional untreated replicates would be expected to accumulate negligible
mutational differences and provide little additional biological information. Replication was instead
prioritized within mutagenized treatments to maximize power and sequencing depth for detecting
treatment-specific genomic and fitness consequences. Following the mutagenesis phase, all
populations were frozen at −80 °C for downstream analyses.

Mutagenesis
Four replicates per strain were treated with 1 mM ethyl methanesulfonate (EMS) in M9 solution,
while the other four replicates were treated with 0.1% (w/v) formaldehyde solution in M9. These
concentrations were chosen because they are non-lethal yet sufficient to induce detectable genomic
mutations. At each transfer, entire populations were moved in suspension (300 µL per replicate),
preserving large census sizes and minimizing bottlenecks. Based on subsampling counts of the
transfer suspension, populations were estimated to contain approximately 3,000–4,000 individuals
per replicate Mutagenesis was carried out for five consecutive generations. During this phase,
populations were exposed to the mutagen for 4 hours every fourth day before being transferred
back to standard NGM plates seeded with *E. coli* OP50. Following the mutagenesis phase, all
populations were allowed to recover for three additional generations without mutagen exposure,
during which worms were maintained under standard laboratory conditions. We followed the worm
mutagenesis protocol described by Brenner et al. (1974) with minor modifications. Every generation
(~4 days), worms were washed from the plates with M9, collected in tubes, and resuspended in their
respective mutagen solutions for 4 hours at 20 °C, during which they were gently rocked to ensure
uniform exposure. After treatment, worms were transferred to fresh plates with NGM and *E. coli*
OP50, ensuring maintenance of replicate identity and avoidance of mixing. Population transfers
were performed without developmental synchronization. Generations were overlapping, i.e.,
individuals at different developmental stages contributed to subsequent generations.Population sizes
were not altered at any point during the experiment. During both transfers, from the plates to the
tubes and from the tubes back to the plates, all worms were transferred using 300 microliters of
liquid. To approximate census size, we counted worms in replicate 10-µL subsamples of the transfer
suspension and extrapolated to 300 µL (average ≈ 3,300 worms per transfer). Specifically, M9 buffer
was used for the transfer from plates to tubes, and the mutagen was used for the transfer from
tubes back to plates. After the mutagen treatment, the liquid was carefully transferred onto the

plate, causing parts of the *E. coli* colony to be dislodged. As a result, the worms would initially swim

on the plate. We then left the plate open, allowing the liquid to be absorbed within 30-40 minutes,

after which the worms resumed crawling on the plate.

<u>*Male frequency assay*</u>

To evaluate the impact of five generations of mutational exposure on male frequency, a male

frequency assay was conducted using both non-mutagenized (parent) worms and worms

immediately after five generations of mutagen treatment. This assay was performed as an initial

phenotypic screen prior to downstream fitness and genomic analyses. Worms from each strain

*treatment combination were bleached for age synchronization. The lifetime progeny production of

15 replicates per strain per treatment was monitored by placing a single L4 worm on a 75mm x

13mm Petri plate, transferring it to a fresh plate every 24 hours. After 96 hours, the worm was

discarded, and the progeny produced over the four days were sexed and counted. Any worms that

died during the transfer process were excluded from the analysis. Male frequency was calculated for

each assay plate individually, and treatment means ± SE were calculated across plates.

<u>*Outcrossing frequency assay*</u>

Outcrossing frequency was assessed using paired matings (one L4 hermaphrodite + one adult male).

For each strain and mutagen treatment, the lifetime progeny production of 15 independent mating

plates was monitored, while 30 independent mating plates were assayed for the non-mutagenized

parental control. The larger sample size for the non-mutagenized parental control was chosen to

provide a high-confidence estimate of baseline male and outcrossing frequencies for each strain,

which serve as reference points for classifying outcrossing events in the mutagenized treatments.

Each mating pair was placed on a 75 mm × 13 mm Petri plate and transferred to a fresh plate every

24 hours. After 96 hours, both the male and hermaphrodite were discarded, and progeny produced

over the four days were allowed to develop to adulthood. Mating plates were pooled across lines

sharing the same treatment history.

Because cross-fertilization in *C. elegans* produces approximately 50% male progeny, whereas

selfing hermaphrodites produce few or no males, male frequency provides a reliable indicator of

whether outcrossing occurred during the assay window [33]. The goal of this assay was to determine

whether successful mating occurred at least once on a plate, rather than to quantify mating

efficiency. Plates were therefore classified as either "outcrossing" or "selfing-dominated" using

strain-specific criteria informed by baseline male production.

Based on our own baseline measurements, spontaneous male production in the N2 and AB1
strains was consistently below 1% both before and after mutagenesis. For these strains, plates were
visually screened for males rather than exhaustively counted. Plates exhibiting male frequencies
clearly exceeding the expected spontaneous range (1–5%) were classified as outcrossing. In practice,
outcrossed plates were readily distinguishable by the presence of numerous males, whereas selfing-
dominated plates contained few or no males and required extended inspection to detect even a
single male. In contrast, CB4856 naturally exhibits elevated baseline male frequencies (~15–17%).
For this strain, progeny on each plate were counted and sexed, and male frequency was calculated
as the proportion of male progeny. To conservatively distinguish true outcrossing events from
baseline variation, CB4856 plates were classified as outcrossing only when male frequency reached
or exceeded 25%, a threshold approximately one standard deviation above baseline male frequency
observed in this strain. To verify that our classification was not sensitive to the chosen cutoff, we
repeated the analysis using alternative thresholds (20% and 30% male frequency) for CB4856. These
sensitivity analyses produced qualitatively identical estimates of outcrossing frequency across
treatments. Outcrossing frequency was calculated as the proportion of plates meeting these criteria
within each strain and treatment.

Relative fitness assay
After five generations of mutagenesis, worms were frozen at -80°C and later recovered for relative
fitness assay at sixth generation- representing the fitness immediately after five generations of
mutagenesis and again after three generations of recovery from mutations using the standard _C._
elegans maintenance protocol at 20 °C. During recovery, worms were maintained using the standard
worm maintenance protocol. For the relative fitness assay, one early-stage L1 worm from each focal
strain (N2, CB4856, AB1) was paired with one L1 worm from the non-mutagenized common
competitor strain ST2 expressing green fluorescent protein (GFP). Both strains were co-cultured on
NGM plates with _E. coli_ OP50 for eight days (~2 generations) at 20 °C. Washed worms from replicate
plates (11 per generation/mutant/strain combination) were collected in M9 on a slide, imaged with
and without a GFP filter on a fluorescent microscope, and counted using a Python-based code
specific for worm morphology (available on the Github). Plates with zero worms of either type were
excluded from analysis, assuming that the worm did not survive to lay eggs. Images captured
without the GFP filter counted all worms present on the plate- both the focal strain and the common
competitor strain. Conversely, images captured with the GFP filter only revealed the green,
fluorescent competitor worms, allowing for estimation of their abundance relative to the total worm
population. Relative fitness was quantified from focal and competitor counts using a ratio-based
metric as described in the Statistical Analysis section.

Fitness was compared across non-mutagenized, post-mutagenized, and recovered lineages
using automated image analysis (Supplementary Material Table 2). To validate the accuracy of the
image analysis pipeline, we randomly selected a subset of samples and manually counted
individuals. This confirmed that the automated method produced reliable counts, with no observed
false positives or false negatives. We found that three generations of recovery from mutagenesis
were sufficient for each strain to recover pre-mutagenesis fitness levels for both the mutagens
(Figure 1C).

DNA extraction and Sequencing

To assess the genomic impact of mutagenesis, high molecular weight DNA was extracted from
worms (after 3 generations of recovery) using the Cytiva kit (detailed procedure available in the
Supplementary Materials). Before starting out the DNA extraction we washed a minimum of ~10,000
worms with M9 buffer from the plate into a tube and kept the worms rocking overnight to remove
associated microbiome. Pooled PacBio long-read sequencing was performed for parental worms of
all three strains (N2, CB4856, AB1) and one population per strain per mutagen. Additionally, Illumina
short-read sequencing was conducted for each replicate of each treatment (control, EMS,
formaldehyde) for all strains. This strategy enabled comprehensive analysis of genomic variants
induced by mutagenesis and their phenotypic consequences in subsequent generations.

For the comprehensive genomic analysis, we sequenced a total of 36 samples. Using PacBio
long-read sequencing, with an average depth of coverage of 94.52x, we sequenced nine samples:
the three parental strains (N2, CB4856, AB1) as controls, and six mutagenized replicates i.e., one
replicate per strain treated with each mutagen. Additionally, we performed Illumina short-read
sequencing on 27 treatment replicates, achieving an average depth of coverage of 72.11x. These 27
samples included the three parental strains, 12 EMS-treated replicates, and 12 formaldehyde-
treated replicates, with four replicates per treatment for each strain. Short-read data were used to
identify SNPs, whereas both long- and short-read data were integrated to generate a high-
confidence SV dataset.

The male frequency and outcrossing rate assays were conducted on worms derived from the
same batch that underwent five generations of mutagenesis. In contrast, the worms used for the
relative fitness assay and DNA extraction came from a different set, although they were
mutagenized in the same manner as the first batch.

Population simulations

To quantitatively examine the influence of outcrossing on SV purging and retention we implemented

our theoretical model in population simulations using the software SLiM 4.3[64]. Our model described

a genomic system where mutations occur as SVs, either insertions, deletions or inversions. Each

individual in a population of size K carried n structural mutations. We constructed dioecious

populations containing equal numbers of males and females and self-fertile populations with a

proportion of individuals reproducing through outcrossing and a proportion reproducing through

selfing. Each population started at a specified size K and evolved through overlapping generations

with reproduction and fitness differences creating variable population sizes. Each individual had a

diploid chromosome 10,000 nucleotides in length. Recombination occurred along the chromosomes

with uniform probability at a specified rate r. After reproduction SV mutations occurred at a rate μ

and were removed at a rate ν, for example by back mutation via gene conversion [65].

We did not explicitly model the physical size of mutations but assumed that mutations had

discrete phenotypic effect sizes that cumulatively influenced fitness. The probability of a SV

mutation occurring at any location was uniform across the chromosome and each SV mutation

conferred a small deleterious phenotypic effect drawn from $\Gamma(k, \theta)$ [39–41](Supplementary Figure 1).

Individual SVs interacted to decrease fitness faster-than-linearly and the fitness landscape had a

negative curvature or synergistic epistasis (Supplementary Figure 3). The fitness of an individual

carrying n SVs was $w(n) = exp^{-(an+(bn^2)/2)}$ where a and b were parameters determining the

curvature of the fitness landscape [36]. Our simulated populations evolved for 100,000 generations

with the parameters $k = 1, \theta = 2, \mu = 0.01, \nu = 0.001, a = 0.001$ and $b = 0.0006$. We evolved

populations under a range of evolutionary scenarios including recombination rates r=0.5-1x10^{-5},

K=100 to 5,000 and levels of self-fertility ranging from 0 to 100%. (Supplementary Figure 2). We also

experimented with the curvature of the landscape, modeling a linear fitness landscape as $w(n) =$

exp^{-an} with $a = 0.00635$ and a positive or antagonistic curvature where fitness impacts attenuate

as they accumulate as $w(n) = exp^{-(an+(bn^2)/2)}$ with $a = 0.001$ and $b = 0.045$ (Supplementary

Figure 3).

Data analysis

Male frequency assay

Male frequency was calculated as the ratio of male progeny to the total progeny laid by each worm,

and this proportion was used as the unit of analysis. We assessed the normality of the data using the

Shapiro-Wilk test and by visualizing its distribution with a histogram. As expected, the distribution

was right-skewed, with a high proportion of zero values, indicating zero-inflation. Given the bounded

nature (0 to 1) of the data and the excess zero values, we used a zero-inflated Beta regression for

analysis. Since this model requires proportion values strictly between 0 and 1, we applied an epsilon

adjustment by adding a small constant to all zero values to make the data more appropriate for the

model. We performed the analysis using the glmmTMB package in R [66], implementing the following

model:

Male Frequency~Strain× Mutagen

In this model Strain (N2, AB1, CB4856) and Mutagen treatment (EMS, Formaldehyde and

non-mutagenized parents were included as fixed factors and an interaction term was added to check

if the effect of strain on male frequency varied by mutagen identity. This model accounts for both (1)

the continuous variation in male frequency and (2) the excess zero values separately, making it a

better fit than standard regression models. Further we did Tukey's HSD comparison (Abdi &

Williams, 2010) on estimated marginal means using emmeans package in R [68]. For visualization

purposes, figures display arithmetic means ± 1 standard error calculated directly from individual

assay plates, rather than model-estimated marginal means.

Relative fitness assay

To quantify the relative fitness of the _C. elegans_ strains N2, CB4856, and AB1, each plate was washed

with M9 buffer solution to prepare the worms for imaging. Two aliquots from the washed plate

were then placed onto a microscope slide and covered with a coverslip. We performed imaging to

count the GFP-expressing competitor strain and the non-GFP-expressing focal strain worms within

these aliquots. The allelic frequency of the non-fluorescent allele was calculated as the square root

of proportion of non-GFP-expressing worms to the total worm population. For statistical analysis,

two-way ANOVAs were conducted separately for each mutagen to assess the effects of strain and

treatment duration on relative fitness. Data normality was examined using the Shapiro-Wilk test and

the residual distribution showed slight variation from normality (Shapiro-Wilk test: W = 0.9868, $p <$

0.01), indicating variation from normality. However, visual inspection of the residuals with histogram

suggested an approximately normal distribution. Since ANOVA is robust to minor deviations, we

proceeded with ANOVA followed by pairwise comparisons using Tukey's HSD test. The following

linear model was applied:

anova(lm(RF ~ Strain*Generation, data = a)).

Following ANOVA, we performed pairwise comparisons using Tukey's Honest Significant

Difference (HSD) test to assess differences in relative fitness across strains and generations. We

conducted separate Tukey HSD tests within each strain to compare relative fitness before

mutagenesis, after five generations of mutagenesis, and after three generations of recovery.

Bioinformatic analysis

De novo mutations were defined as variants detected in recovered populations that were absent

from the corresponding non-mutagenized parental populations when compared with the reference

genome. The bioinformatic analysis for identifying *de novo* mutations following PacBio sequencing

involved a comprehensive workflow to process and analyse long-read data, distinguishing mutations

that arose in the worms after mutagenesis from those existing in parental strains. Initially, raw

PacBio reads were aligned and indexed to reference genomes using minimap2 [69] and bcftools [70]; for

the N2 and CB4856 strains, NCBI reference genomes were used, while the AB1 strain employed the

recently annotated genome of its descendant strain, SX3368[71], to account for genomic divergences

within the lineage. Subsequent variant calling was performed using Sniffles [72] for SVs, such as

insertions, deletions, and inversions, capitalizing on the accuracy offered by long-read data.

For the specific identification of de novo mutations induced by mutagenesis, a comparative

analysis using bcftools contrasted the variant profiles of generation 8 worms (post three generations

of recovery) with their control parents, filtering out pre-existing genetic variations. All parental and

mutagenized samples were processed using identical library preparation, sequencing, and

bioinformatics pipelines to ensure comparability in variant detection. Further quality control and

data filtering ensured only high-quality SNPs and SVs (quality score $\geq$ 20, depth $\geq$ 10) were retained.

We then used bedtools to provide insights into their potential impacts on exonic regions and

regulatory elements.

Additionally, similar bioinformatic steps were conducted with Illumina sequencing data.

Alignment and indexing were again performed on reference genomes, followed by SNP and SV

calling using Freebayes [73] and Manta [74], respectively. We initially called variants in pooled-population

mode (FreeBayes 'pooled') to account for minor alleles in the combined worm samples. However,

when intersecting these SNP calls with structural variants, we observed inflated counts due to

multi-allelic sites being represented multiple times. To avoid this over-counting artifact, we

generated a separate VCF using non-pooled calling parameters and used these calls for all

subsequent intersection and annotation analyses. Enhanced SV detection was achieved with

VGtoolkit [75] through graph-based variant calling, which was then integrated with SVs identified by

Sniffles 2.0 using Survivor 1.0.3[76], creating a comprehensive view of the genomic alterations induced

by the mutagens.

Long-read sequencing was performed on the three non-mutagenized parent strains and
one replicate per strain for each mutagen treatment. Short-read sequencing covered all replicates
across all strains. For structural variation analysis, we merged variants identified by both long-read
and short-read callers using Survivor 1.0.3. For SNP analysis, we relied solely on variants identified by
short-read callers. To isolate mutagen-induced effects, the ancestral (pre-treatment) genome was
used as the reference baseline for all de novo variant calling. Parallel non-mutagenized controls
were not sequenced because spontaneous SVs (SVs) in *C. elegans* occur at extremely low
frequencies (approximately 0.003–0.03 events per genome per generation;[42,47]). Given the three-
generation duration of our experiment and the detection limits of pooled long-read sequencing, the
expected contribution of spontaneous SVs is negligible. Accordingly, the SVs identified here
predominantly reflect high-confidence, mutagen-induced genomic changes

To test whether SNPs were enriched within SV regions, the expected number of SNPs within
SV intervals was calculated as: Expected = Total SNPs × (SV base pairs / Genome base pairs).
Observed SNP counts within SV intervals were then compared with expectation using one-sided
binomial tests for each strain–treatment combination.

<u>Annotation of Transposable Elements and Intersection with Structural Variants</u>
Transposon annotation is facilitated in complete and contiguous sequences, thus annotation of TEs
was done in the N2, AB, and CB reference genomes rather than directly from the raw sequence data.
We used the TransposonUltimate reasonaTE pipeline for TE annotation in each of the reference
genomes. Briefly, TEUltimate reasonaTE wraps 13 previously published TE annotation tools, which
include: RepeatMasker v4.1.1 , RepeatModeler v2.0.1, SINE-Scan , SINE-Finder, LTRPred ,
LTRHarvest, TIRvish , MUSTv2 , MITE-Tracker, MITEFinderII, HelitronScanner, TransposonPSI for
transposon discovery. While some functions are optional, all subprograms were run with default
parameters in this study. The resulting annotations are filtered, merged, and clustered using CD-HIT
v4.8.1 [77] and BLASTN v2.10.1 [78]. The final annotations from FinalAnnotations_Transposons.gff3 in the
final Results folder were used for downstream analysis for each of the three focal strains.
Evidence of transposon movement was assumed from structural variants which overlapped
with transposon annotations. The TEUltimatedeTEct pipeline[34] Click or tap here to enter
text.was used for intersection of transposon and structural variant annotations. De novo VCF
structural variant files, as generated above, were intersected with transposon annotations and
reported as an event if overlapping by at least 10% and lengths were similar by at least 50%. Finally,

any annotation smaller than 50 base pairs, or larger than 1% of the genome was discarded. Plots

were generated from the transpositionEvents.gff3 file in RStudio with R v4.3.2 (R Core Team, 2023).

<u>Literature Cited</u>

1.   Lynch, M. *et al. The Mutational Meltdown in Asexual Populations. We Are Very*
Grateful To. Journal of Heredity vol. 84 (1993).

2.   Lynch, M. *et al.* Perspective: Spontaneous deleterious mutation. *Evolution (N. Y).* **53**,
645–663 (1999).

3.   Robert, F. & Pelletier, J. Exploring the Impact of Single-Nucleotide Polymorphisms on
Translation. *Frontiers in Genetics* vol. 9 Preprint at
https://doi.org/10.3389/fgene.2018.00507 (2018).

4.   Feuk, L., Carson, A. R. & Scherer, S. W. Structural variation in the human genome.
Nature Reviews Genetics vol. 7 85–97 Preprint at https://doi.org/10.1038/nrg1767
(2006).

5.   Collins, R. L. *et al.* A structural variation reference for medical and population
genetics. *Nature* **581**, 444–451 (2020).

6.   Mahmoud, M. *et al.* Structural variant calling: The long and the short of it. *Genome*
Biology vol. 20 Preprint at https://doi.org/10.1186/s13059-019-1828-7 (2019).

7.   Ho, S. S., Urban, A. E. & Mills, R. E. Structural variation in the sequencing era. *Nature*
Reviews Genetics vol. 21 171–189 Preprint at https://doi.org/10.1038/s41576-019-
0180-9 (2020).

8.   Glémin, S. How are deleterious mutations purged? Drift versus nonrandom mating.
Evolution (N. Y). **57**, 2678–2687 (2003).

9.   Whitlock, M. C. & Agrawal, A. F. Purging the genome with sexual selection: Reducing
mutation load through selection on males. *Evolution* vol. 63 569–582 Preprint at
https://doi.org/10.1111/j.1558-5646.2008.00558.x (2009).

10.   García-Dorado, A. Understanding and predicting the fitness decline of shrunk
populations: Inbreeding, purging, mutation, and standard selection. *Genetics* **190**,
1461–1476 (2012).

11.   Byers, D. L. & Waller, D. M. *Do Plant Populations Purge Their Genetic Load? Effects of*
Population Size and Mating History on Inbreeding Depression. Source: Annual Review
of Ecology and Systematics vol. 30 https://about.jstor.org/terms (1999).

12. Szövényi, P. *et al.* Efficient purging of deleteriousmutations in plants with haploid
selfing. *Genome Biol. Evol.* **6**, 1238–1252 (2014).

13. MacPherson, B., Scott, R. & Gras, R. Sex and recombination purge the genome of
deleterious alleles: An Individual Based Modeling Approach. *Ecological Complexity* **45**,
(2021).

14. Sianta, S. A., Peischl, S., Moeller, D. A. & Brandvain, Y. The efficacy of selection may
increase or decrease with selfing depending upon the recombination environment.
Evolution (N. Y). **77**, 394–408 (2023).

15. Hartfield, M. Evolutionary genetic consequences of facultative sex and outcrossing. *J.*
Evol. Biol. **29**, 5–22 (2016).

16. Hoffmann, A. A. & Rieseberg, L. H. Revisiting the impact of inversions in evolution:
From population genetic markers to drivers of adaptive shifts and speciation? *Annual*
Review of Ecology, Evolution, and Systematics vol. 39 21–42 Preprint at
https://doi.org/10.1146/annurev.ecolsys.39.110707.173532 (2008).

17. Kirkpatrick, M. & Barton, N. Chromosome inversions, local adaptation and speciation.
Genetics **173**, 419–434 (2006).

18. Kazazian, H. H. *Mobile Elements: Drivers of Genome Evolution*.
https://www.science.org.

19. Feschotte, C. & Pritham, E. J. *DNA Transposons and the Evolution of Eukaryotic*
Genomes.

20. Belyayev, A. Bursts of transposable elements as an evolutionary driving force. *Journal*
of Evolutionary Biology vol. 27 2573–2584 Preprint at
https://doi.org/10.1111/jeb.12513 (2014).

21. Wright, S. & Finnegan, D. *Genome Evolution: Sex and the Transposable Element*.
Current Biology vol. 11 (2001).

22. Boutin, T. S., Le Rouzic, A. & Capy, P. How does selfing affect the dynamics of selfish
transposable elements? *Mob. DNA* **3**, (2012).

23. Anderson, J. L., Morran, L. T. & Phillips, P. C. Outcrossing and the maintenance of
males within C. elegans populations. in *Journal of Heredity* vol. 101 (2010).

24. Wegewitz, V., Schulenburg, H. & Streit, A. Experimental insight into the proximate
causes of male persistence variation among two strains of the androdioecious
Caenorhabditis elegans (Nematoda). *BMC Ecol.* **8**, (2008).

25. Manoel, D., Carvalho, S., Phillips, P. C. & Teotónio, H. Selection against males in
Caenorhabditis elegans under two mutational treatments. *Proceedings of the Royal*
Society B: Biological Sciences **274**, 417–424 (2007).

26.   Stewart, A. D. & Phillips, P. C. *Selection and Maintenance of Androdioecy in*
Caenorhabditis Elegans.

27.   Gray, J. C. & Cutter, A. D. Mainstreaming Caenorhabditis elegans in experimental
evolution. *Proceedings of the Royal Society B: Biological Sciences* vol. 281 Preprint at
https://doi.org/10.1098/rspb.2013.3055 (2014).

28.   Manoel, D., Carvalho, S., Phillips, P. C. & Teotónio, H. Selection against males in
Caenorhabditis elegans under two mutational treatments. *Proceedings of the Royal*
Society B: Biological Sciences **274**, 417–424 (2007).

29.   Cutter, A. D. Mutation and the experimental evolution of outcrossing in
Caenorhabditis elegans. *J. Evol. Biol.* **18**, 27–34 (2005).

30.   Chelo, I. M. *et al.* Partial selfing can reduce genetic loads while maintaining diversity
during experimental evolution. *G3: Genes, Genomes, Genetics* **9**, 2811–2821 (2019).

31.   Teterina, A. A. *et al.* Genomic diversity landscapes in outcrossing and selfing
Caenorhabditis nematodes. *PLoS Genet.* **19**, (2023).

32.   Kutscher, L. M. & Shaham, S. Forward and reverse mutagenesis in C. elegans.
WormBook: the online review of C. elegans biology 1–26 Preprint at
https://doi.org/10.1895/wormbook.1.167.1 (2014).

33.   Yin, D. & Haag, E. S. Evolution of sex ratio through gene loss. *Proc. Natl. Acad. Sci. U.*
S. A. **116**, 12919–12924 (2019).

34.   Riehl, K., Riccio, C., Miska, E. A. & Hemberg, M. TransposonUltimate: software for
transposon classification, annotation and detection. *Nucleic Acids Res.* **50**, E64 (2022).

35.   Kondrashov, A. S. Selection against harmful mutations in large sexual and asexual
populations. *Genet. Res.* **40**, 325–332 (1982).

36.   Charlesworth, B. OPTIMIZATION MODELS, QUANTITATIVE GENETICS, AND
MUTATION. *Evolution (N. Y).* **44**, 520–538 (1990).

37.   Charlesworth, B. Mutation-selection balance and the evolutionary advantage of sex
and recombination. *Genet. Res.* **55**, 199–221 (1990).

38.   Haller, B. C. & Messer, P. W. SLiM 4: Multispecies Eco-Evolutionary Modeling.
American Naturalist **201**, E127–E139 (2023).

39.   Eyre-Walker, A. & Keightley, P. D. The distribution of fitness effects of new mutations.
Nature Reviews Genetics vol. 8 610–618 Preprint at https://doi.org/10.1038/nrg2146
(2007).

40. Keightley, P. D. & Eyre-Walker, A. Joint inference of the distribution of fitness effects
of deleterious mutations and population demography based on nucleotide
polymorphism frequencies. *Genetics* **177**, 2251–2261 (2007).

41. Racimo, F. & Schraiber, J. G. Approximation to the Distribution of Fitness Effects
across Functional Categories in Human Segregating Polymorphisms. *PLoS Genet.* **10**,
(2014).

42. Konrad, A. *et al.* Mitochondrial mutation rate, spectrum and heteroplasmy in
Caenorhabditis elegans spontaneous mutation accumulation lines of differing
population size. *Mol. Biol. Evol.* **34**, 1319–1334 (2017).

43. Estes, S., Phillips, P. C., Denver, D. R., Thomas, W. K. & Lynch, M. *Mutation*
Accumulation in Populations of Varying Size: The Distribution of Mutational Effects for
Fitness Correlates in Caenorhabditis Elegans.

44. Ciccia, A. & Elledge, S. J. The DNA Damage Response: Making It Safe to Play with
Knives. *Molecular Cell* vol. 40 179–204 Preprint at
https://doi.org/10.1016/j.molcel.2010.09.019 (2010).

45. Limpose, K. L., Corbett, A. H. & Doetsch, P. W. BERing the burden of damage: Pathway
crosstalk and posttranslational modification of base excision repair proteins regulate
DNA damage management. *DNA Repair* vol. 56 51–64 Preprint at
https://doi.org/10.1016/j.dnarep.2017.06.007 (2017).

46. Hillier, L. D. W. *et al.* Comparison of C. elegans and C. briggsae genome sequences
reveals extensive conservation of chromosome organization and synteny. *PLoS Biol.*
5, 1603–1616 (2007).

47. Saxena, A. S. & Baer, C. F. High rate of mutation and efficient removal by selection of
structural variants from natural populations of Caenorhabditis elegans. Preprint at
https://doi.org/10.1101/2025.03.22.644739 (2025).

48. Kim, C. *et al.* Long-read sequencing reveals intra-species tolerance of substantial
structural variations and new subtelomere formation in C. elegans. *Genome Res.* **29**,
1023–1035 (2019).

49. Bush, Z. D. *et al.* Transposable elements and heterochromatic regions are enriched
for structural variation and sequence divergence in the genome of wild-type
Caenorhabditis elegans. *G3: Genes, Genomes, Genetics* **15**, (2025).

50. Thompson, O. A. *et al.* Remarkably divergent regions punctuate the genome assembly
of the Caenorhabditis elegans hawaiian strain CB4856. *Genetics* **200**, 975–989 (2015).

51. Brenner, S. *THE GENETICS OF CAENORHABDZTZS ELEGANS.*

52. Andersen, E. C. *et al.* Chromosome-scale selective sweeps shape Caenorhabditis
elegans genomic diversity. *Nat. Genet.* **44**, 285–290 (2012).

53. Ristow, M. & Schmeisser, K. Mitohormesis: Promoting health and lifespan by
increased levels of reactive oxygen species (ROS). *Dose-Response* **12**, 288–341 (2014).

54. Cypser, J. R. & Johnson, T. E. *Multiple Stressors in Caenorhabditis Elegans Induce*
Stress Hormesis and Extended Longevity. Journal of Gerontology vol. 57
https://academic.oup.com/biomedgerontology/article/57/3/B109/550368 (2002).

55. Negi, P., Rai, A. N. & Suprasanna, P. Moving through the stressed genome: Emerging
regulatory roles for transposons in plant stress response. *Frontiers in Plant Science*
vol. 7 Preprint at https://doi.org/10.3389/fpls.2016.01448 (2016).

56. Wright, S. I. & Schoen, D. J. *Transposon Dynamics and the Breeding System. Genetica*
vol. 107 (1999).

57. Dolgin, E. S. & Charlesworth, B. The fate of transposable elements in asexual
populations. *Genetics* **174**, 817–827 (2006).

58. Emmons, S. W. & Yesner, L. *High-Frequency Excision of Transposable Element Tel in*
the Nematode Caenorhabditis Elegans Is Limited to Somatic Cells. Cell vol. 36 (1964).

59. Mori, I., Benian, G. M., Moermant, D. G. & Waterston, R. H. *Transposable Element Tcl*
of Caenorhabditis Elegans Recognizes Specific Target Sequences for Integration (Site
Specificity/Transposon/Unc-22/Nematode). Proc. Natl. Acad. Sci. USA vol. 85 (1988).

60. Bao, W., Jurka, M. G., Kapitonov, V. V. & Jurka, J. New superfamilies of eukaryotic
DNA tyransposons and their internal divisions. *Mol. Biol. Evol.* **26**, 983–993 (2009).

61. Laricchia, K. M., Zdraljevic, S., Cook, D. E. & Andersen, E. C. Natural Variation in the
Distribution and Abundance of Transposable Elements Across the Caenorhabditis
elegans Species. *Mol. Biol. Evol.* **34**, 2187–2202 (2017).

62. Sterken, M. G., Snoek, L. B., Kammenga, J. E. & Andersen, E. C. The laboratory
domestication of Caenorhabditis elegans. *Trends in Genetics* vol. 31 224–231 Preprint
at https://doi.org/10.1016/j.tig.2015.02.009 (2015).

63. Anderson, J. L., Morran, L. T. & Phillips, P. C. Outcrossing and the maintenance of
males within C. elegans populations. in *Journal of Heredity* vol. 101 (2010).

64. Haller, B. C. & Messer, P. W. SLiM 4: Multispecies Eco-Evolutionary Modeling.
American Naturalist **201**, E127–E139 (2023).

65. Katju, V., LaBeau, E. M., Lipinski, K. J. & Bergthorsson, U. Sex change by gene
conversion in a Caenorhabditis elegans fog-2 mutant. *Genetics* **180**, 669–672 (2008).

66. Bolker, B. *Getting Started with the GlmmTMB Package.* (2022).

67. Abdi, H. & Williams, L. J. *Encyclopedia of Research Design*.
http://www.utd.edu/~herve.

68. Searle, S. R., Speed, F. M. & Milliken, G. A. Population marginal means in the linear
model: An alternative to least squares means. *American Statistician* **34**, 216–221
(1980).

69. Li, H. Minimap2: Pairwise alignment for nucleotide sequences. *Bioinformatics* **34**,
3094–3100 (2018).

70. Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *Gigascience* **10**, (2021).

71. Riccio C. Riccio, C., 2022. Duplication is a prominent mechanism of recent gene birth
in Caenorhabditis elegans (Doctoral dissertation). (2022).

72. Smolka, M. *et al.* Detection of mosaic and population-level structural variants with
Sniffles2. *Nat. Biotechnol.* https://doi.org/10.1038/s41587-023-02024-y (2024)
doi:10.1038/s41587-023-02024-y.

73. Garrison, E. & Marth, G. Haplotype-based variant detection from short-read
sequencing. http://arxiv.org/abs/1207.3907 (2012).

74. Chen, X. *et al.* Manta: Rapid detection of structural variants and indels for germline
and cancer sequencing applications. *Bioinformatics* **32**, 1220–1222 (2016).

75. Hickey, G. *et al.* Genotyping structural variants in pangenome graphs using the vg
toolkit. *Genome Biol.* **21**, (2020).

76. Jeffares, D. C. *et al.* Transient structural variations have strong effects on quantitative
traits and reproductive isolation in fission yeast. *Nat. Commun.* **8**, (2017).

77. Fu, L., Niu, B., Zhu, Z., Wu, S. & Li, W. CD-HIT: Accelerated for clustering the next-
generation sequencing data. *Bioinformatics* **28**, 3150–3152 (2012).

78. Camacho, C. *et al.* BLAST+: Architecture and applications. *BMC Bioinformatics* **10**,
(2009).

79. Riehl, K., Riccio, C., Miska, E. A. & Hemberg, M. TransposonUltimate: software for
transposon classification, annotation and detection. *Nucleic Acids Res.* **50**, E64 (2022).

934 **Figures**

Figure 1. **Male frequency, outcrossing behaviour, and relative fitness reveal strain-specific responses to mutagenic stress:** (A) Male frequency across strains and treatments. Colored points represent individual assay plates, while open diamonds represent the mean male frequency for each strain–treatment combination. Error bars indicate ±1 standard error (SE) of the mean across plates. Non-mutagenized CB4856 populations exhibited significantly higher male frequencies than AB1 and N2. Mutagenesis had strain-specific effects, reducing male frequency in CB4856 but producing no significant change in AB1 or N2. (B) Outcrossing frequency across strains and treatments. Triangles represent the proportion of assay plates classified as outcrossing for each strain–treatment combination. CB4856 exhibited the highest outcrossing frequencies across conditions, N2 the lowest, and AB1 intermediate levels.

948 (C) Relative fitness distributions across strains and generations. Violin plots show the distribution of
949 plate-level relative fitness values, with boxplots indicating median and interquartile ranges. Colors
950 represent strains and shading indicates generation: non-mutagenized (light), mutagenized
951 (medium), and recovered (dark). Across all strains, relative fitness returned to levels statistically
952 indistinguishable from non-mutagenized baselines following recovery.

Figure 2: Comparison of de novo SNPs and SVs mutations and their localization in exons across strains (A) Mean counts of de novo SNPs (Circles) and SVs (Triangles) induced by EMS and Formaldehyde across AB1 (Green), CB4856 (Red), and N2 (Blue). De novo SNPs are significantly more abundant than de novo SVs under both mutagen treatments. CB4856 has the maximum number of de novo SNPs and de novo SVs. (B) Mean number of de novo SNPs and de novo SVs localized in exons for the same strains and mutagen treatments, with de novo SNPs again being more prevalent than de novo SVs and CB4856 having more SVs in exons than the other two strains and more SNPs in exons than AB1. Compared to N2, CB4856 had more SNPs in exon than N2 when treated with EMS and comparable SNPs in exon when treated with Formaldehyde.

Figure 3: Mutational and TE landscape across strains and mutagens **(A)**: Number of SNPs, SVs and
SNPs in SVs for AB1, CB4856 and N2; **(B)**: Classification and count of overall TE families contributing
to transposition events across strain-treatment combinations. DNA transposons such as Zator and
Tc1-Mariner were among the most active elements, with CB4856 showing the broadest TE activity
profile.

Figure 4. Structural variation patterns across chromosome II following EMS mutagenesis. Comparison of structural variants detected in reference and mutagenized populations of CB4856, AB1, and N2. Black tick marks indicate individual structural variant breakpoints, while colored blocks denote larger structural rearrangements classified by mutation type (deletion, duplication, insertion, inversion, or other). Shaded connectors indicate large inversion or rearrangement events between reference and mutagenized genomes. CB4856 populations exhibit a greater density and span of structural variants relative to AB1 and N2.

Figure 5. Genomic distribution of SNPs, structural variants (SVs), and transposable elements (TEs) across the genomes of three Caenorhabditis elegans strains following EMS mutagenesis. Columns represent the six chromosomes, and rows correspond to the three strains (AB1, CB4856, and N2). Points show the percentage of 10-kb genomic windows containing each feature across genomic position. Smoothed lines represent locally estimated scatterplot smoothing (LOESS) fits for transposable elements (blue), structural variants (red), and SNPs (green). Black downward-facing triangles indicate positions of predicted active transposable elements. Features exceeding 10 kb in length were excluded from the visualization; details of these larger structural variants are provided in Supplementary Table 9.

1011

1012
1013 Figure 6: Chromosomal distribution of SVs and TE-associated SVs across strains and mutagens. (A)
1014 Stacked bar plots showing the total count and types of SVs including Deletions, Insertions,
1015 Duplications, Inversions, Translocations, and Unclassified (NA) across chromosomes 1–5 and X, for
1016 each strain (AB1, CB4856, N2) and mutagen (EMS and formaldehyde). (B) TE associated SVs were
1017 primarily Deletions.

1018

1019

1020

1021

1022

1023

1024

1025

1026

1027

1028 Figure 7. A) Simulated populations evolving with less than 10% of individuals reproducing through

1029 outcrossing do not retain SVs. B) The mean number of structural variant mutations is lower in

1030 populations with lower quantitative levels of reproduction through outcrossing. Here K=1000,

1031   =0.001,   =0.0006,   =0.01,   =0.001 and recombination rate   =0.0001.

1032