<u>**Outcrossing Complicates Mutation Purging by Trapping Single Nucleotide Polymorphisms in**</u>

<u>**Structural Variant Mutations**</u>

R. Kapila[1,2*], S. Saber[1,2], R.K. Verma[1,2], G. Blanco[1,2], V.K. Eggers[1,2,] J.L. Fierst[1,2*]

[1]Department of Biological Sciences and [2]Biomolecular Sciences Institute

Florida International University, 11200 8th Street, Miami, 33199, FL, USA

*Corresponding author(s). E-mail(s): rkapila@fiu.edu; jfierst@fiu.edu

<u>**Author contributions**</u>

J.L.F. secured funding. R.K. and J.L.F. designed the experiments. R.K., with assistance from S.S.,

performed the fitness assays, and, with assistance from G.B., conducted the male frequency and

outcrossing rate assays. V.K.E. carried out the transposable element analysis. R.K. performed the

genomic analysis with guidance from J.L.F. R.K.V. developed the Python code for worm counting.

J.L.F. developed the theoretical model and conducted the population simulations. R.K. and J.F.K. co-

wrote the first draft. R.K. and J.F.K. prepared the final manuscript with contributions from V.K.E.

<u>**Acknowledgements**</u>

The authors would like to thank the members of the Fierst lab and Jessica Gonzalez, Jason Pienaar

and Jesualdo Fuentes-Gonzalez for suggestions and careful feedback.

<u>**Competing Interests:**</u>

The authors declare no competing financial or non-financial interests.

<u>**Classification**</u>

<u>**Keywords:**</u> Mutations, Outcrossing, Structural variations, Genome, *Caenorhabditis elegans*

<u>**This PDF file includes**</u>

Main Text

Figures 1 to 6

Abstract

Classical mutational theories centered on single nucleotide polymorphisms suggest that outcrossing enhances the purging of deleterious mutations by promoting recombination. However, larger structural variants, such as insertions, deletions, and inversions, can suppress recombination and create linkage blocks. In this study, using experimental evolution in *C. elegans* lines and whole-genome sequencing, we show that outcrossing populations struggle to purge structural mutations and instead retain mutations small and large as single nucleotide polymorphisms become trapped within larger structural variants. Supporting this, our population genetic simulations showed that structural variants accumulate more readily under higher outcrossing rates. These findings contradict classical population genetic expectations, showing that outcrossing can hinder the removal of structural mutations and complicate mutation purging by trapping small variants within larger mutations. Together, our results challenge the prevailing view that sex uniformly promotes genomic resilience and reveals a previously underappreciated constraint on mutation purging.

Introduction

The accumulation of deleterious mutations can have severe consequences for populations, leading to long-term fitness decline or, in extreme cases, extinction (Lynch et al., 1993). The ability to effectively eliminate or purge harmful mutations is critical for reducing genetic load and ensuring long-term evolutionary fitness (Lynch et al., 1999). Based on their size and scale, mutations are broadly categorized into two types: single nucleotide polymorphisms (SNPs) and structural variants (SVs). SNPs are small-scale changes involving a single base pair, often resulting in localized effects on amino acids or regulatory elements (Robert & Pelletier, 2018). In contrast, SVs are large-scale genomic changes including deletions, duplications, inversions, and translocations. These can impact larger portions of the genome, frequently altering multiple genes or regulatory networks (Feuk et al., 2006).

While mutation purging operates on both SNPs and SVs, the process may be more straightforward for SNPs due to their smaller scale, localized effects, and simpler genomic dynamics (Collins et al., 2020). In contrast, purging SVs may be more complex due to their larger genomic footprint and ability to suppress recombination (Mahmoud et al., 2019). Evolutionary theory has traditionally viewed the spectrum of genomic changes under the broad category of "mutation". Much of the research on mutation purging has disproportionately focused on SNPs as they are more easily and accurately detected (Ho et al., 2020). However, the advent of large-scale genome sequencing has revealed the ubiquity of SVs. Despite the foundational role of mutation purging in evolutionary theory (García-Dorado, 2012; Glémin, 2003; Whitlock & Agrawal, 2009), our

understanding of how it operates on SVs remains unexplored. Addressing this is essential to refining

theoretical models, validating predictions with empirical evidence and understanding how mutation

size and severity interact with established evolutionary predictions.

Mating systems influence mutation purging by affecting genetic diversity, recombination,

and the expression patterns of deleterious mutations (Byers & Waller, 1999; Szövényi et al.,

2014). Current consensus in the field is that outcrossing populations, through recombination, create

novel genetic combinations which facilitates the purging of deleterious variation (MacPherson et al.,

2021). However, outcrossing can also potentially dilute the effects of deleterious mutations and slow

their purging (Sianta et al., 2023). Reduced recombination and genetic diversity in populations that

reproduce through self-fertility may lead to the accumulation of mildly deleterious mutations

through genetic drift (Hartfield, 2016). Selfing however, can also increase homozygosity, exposing

harmful recessive mutations to selection and enabling more effective purging (Hartfield, 2016).

These processes may differ for mutations of different physical sizes. SNPs, with largely localized

effects, may be more directly exposed to selection. Purging deleterious SVs may be complicated by

their larger size and diverse impacts, both of which can vary significantly with mating system

(Hoffmann & Rieseberg, 2008; Kirkpatrick & Barton, 2006). Outcrossing can facilitate purging

through recombination but this process could be physically disrupted by SVs, which are capable of

suppressing recombination and creating linkage blocks. The complex interplay between mating

system, recombination suppression by SVs, and mutation purging is not well understood.

In this study we use experimental evolution to address two key questions: First, how do

levels of outcrossing influence the genome's ability to purge deleterious mutations? Second, how

does this ability vary with mutation size and genomic location? To investigate these questions, we

experimentally evaluated how three strains of *Caenorhabditis elegans*, which differ in natural male

frequency (N2, AB1, and CB4856) respond to mutagenesis and recovery. As an androdiecious species

C. elegans populations consist of both males and hermaphrodites, allowing for reproduction through

either self-fertilization or outcrossing with males. While self-fertilization predominates in *C. elegans*,

the frequency of males and outcrossing varies drastically across different strains (Gaertner &

Phillips, 2010) making *C. elegans* a prime system to test these ideas.

We subjected *C. elegans* strains to two contrasting mutagens, EMS to induce single

nucleotide mutations and formaldehyde to introduce complex lesions in the genome (Kutscher &

Shaham, 2014). We mutagenized worms for five consecutive generations and allowed them to

recover from induced mutations for three generations. We then tracked the population's relative

fitness, changes in male frequency, and outcrossing rates to investigate their ability to cope with

mutational input and purge deleterious mutations. We employed both long-read and short-read

sequencing to quantify the size, location, and types of mutations that persist after recovery.

Results

Experimental test of the outcrossing prediction

To test the impact of outcrossing on mutations we subjected three strains of *C. elegans* (N2 (Bristol),

AB1, and CB4856 (Hawaiian)), obtained from the *Caenorhabditis* Genetics Center, University of

Minnesota) to five generations of mutagenesis followed by three generations of recovery in large,

high-density populations. AB1 and CB4856 are wild isolates, whereas N2 is a long-term laboratory

adapted strain, allowing us to test the impact of mutagenesis on diverse genetic backgrounds

(Brenner 1974; Sterken et al. 2015). Each strain was maintained in two distinct lineages: one

mutagenized with EMS and the other with formaldehyde, with four replicate populations per

lineage.

Male frequency was differentially affected by mutagenesis

Non-mutagenized CB4856 populations had the highest male frequencies (Figure 1A), significantly

greater than either AB1 ($p < 0.0001$) or N2 ($p < 0.01$). Mutagenesis had a strain-specific effect on

male frequency. In CB4856 populations male frequency was reduced after mutagenesis with EMS (p

= 0.03; Figure 1A) and formaldehyde, although this comparison was not statistically significant (p =

0.11). In N2 populations, male frequency did not change in response to either mutagen (p = 1.00 for

both). Similarly, AB1 male frequencies did not differ across populations or treatments (all p-values

were > 0.05; Supplementary Material Table 1; Figure 1A).

Outcrossing frequencies varied between C. elegans strains

We found that CB4856 populations had the highest outcrossing frequencies under all conditions,

with peaks following mutagenesis (Figure 1A). In contrast, N2 populations displayed the lowest

outcrossing frequencies. AB1 populations were intermediate to these with notably higher values in

EMS mutagenized populations compared with both non-mutagenized strains and the AB1

formaldehyde mutagenized population.

C. elegans populations quickly recovered fitness

Three generations of recovery from mutagenesis were sufficient to restore fitness to pre-

mutagenesis levels across all strains (Figure 1B). For CB4856 relative fitness was not affected by

formaldehyde (p=1.00) or EMS (p=0.18). Similarly, for AB1 pre-mutagenesis and post-recovery

fitness did not change with treatment by either formaldehyde (p=0.94) or EMS (p=0.76). The initial
response to mutagenesis was different for N2 populations which showed an increase in relative
fitness immediately after exposure to formaldehyde (p<0.03) but not EMS (p<0.46). However, N2
relative fitness decreased to pre-mutagenesis levels after recovery (p=1.00).

Genomic analyses showed numerous SVs and SNPs were retained after recovery
Despite the rapid fitness recovery, sequence analyses showed that populations harbored numerous
de novo mutations. To comprehensively identify SVs and SNPs we sequenced whole genome DNA
libraries with both Pacific Biosystems long-read and Illumina short-read DNA sequencing. Short-read
data was used to characterize SNPs and both short- and long-read data were used to generate a
high-confidence SV dataset.

We found that CB4856 exhibited significantly higher susceptibility to mutagenesis compared
to AB1. When mutagenized with formaldehyde, CB4856 populations showed a 1.12-fold increase in
de novo SNPs compared to AB1 (t=5.55, _p_=0.0014) and comparable SNP counts to N2 (t < 0.01, _p_=
0.996). In contrast, with EMS as the mutagen, CB4856 showed a significantly higher _de novo_ SNP
count when compared with AB1 (t = 6.79, p < 0.01) and a marginally higher _de novo_ SNP count when
compared with N2 populations (t = 2.45, p = 0.05; Figure 2A).

Due to marked heteroscedasticity in SV counts we performed Mann–Whitney U tests to
compare _de novo_ SV counts among strains. Under both EMS and formaldehyde treatments, CB4856
populations had significantly higher _de novo_ SV counts (Figure 2A) when compared with AB1 (W=16,
_p_ = 0.03 for both treatments) and N2 populations (W=16, _p_ = 0.03 for both treatments).

CB4856 harbored high numbers of SNPs and SVs in exons
CB4856 populations had the highest number of mutations in coding regions (Figure 2B) and the
highest number of SNPs trapped within SVs (Figure 3A). When formaldehyde was used as the
mutagen, CB4856 had 2.20 times more exonic SNPs than AB1 (W=0, _p_= 0.03) and 1.12 times more
than N2 (W=12, _p_=0.34). Similarly, when mutagenized with EMS, CB4856 had 2.26 times more
exonic SNPs compared to AB1 (W=0, _p_=0.03) and 1.38 times more exonic SNPs compared to N2
(W=16, _p_=0.03) (Figure 2B).

Exons were similarly affected by SVs. AB1 populations showed the least number of exons
containing SVs, whereas CB4856 populations had the highest number. AB1 formaldehyde
populations had 16.25-fold fewer exonic SVs than CB4856 (W = 0, p = 0.02) and 6.92-fold fewer than
N2 (W = 16, p = 0.03). CB4856 EMS populations had 22.31-fold more exonic SVs compared to AB1 (W
= 0, p = 0.02), and N2 had 7.81-fold more when compared with AB1 (W = 16, p = 0.03) (Figure 2B).

We found that CB4856 had the highest number of SNPs trapped within SVs (77.13% for

formaldehyde and 82.46% for EMS treatment; Figure 3A). In contrast, SVs in AB1 populations

contained 6.36% (formaldehyde) and 12.01% (EMS) of *de novo* SNPs. N2 populations had similarly

low numbers of SNPs contained in SVs (EMS 7.12%; formaldehyde 6.80%).

Active transposable elements were responsible for a small portion of SV mutations

Both mutagenesis and outcrossing can activate TEs. To determine if TE activity was responsible for

the SVs in our mutagenized populations we used TransposonUltimate (Riehl et al., 2022) to annotate

repeats and TEs. We compared populations pre- and post- mutagenesis to both analyze *de novo* TE-

associated SV mutations and identify potentially active TEs (Riehl et al., 2022).

We found that *de novo* TE-associated SVs constituted a small proportion (<15% in all

populations) of the total *de novo* SVs. The parental AB1 population had the highest genomic repeat

content (25.71%) but the fewest TE-associated SVs (Figure 3B). In contrast, CB4856 had the lowest

genomic repeat content (21.68%) and the most TE-associated SVs. The N2 parental population had

23.19% genomic repeat content. Some TEs, including Copia, Gypsy, and Sola in CB4856, were strain-

specific or uniquely active in one population. Others like the DNA transposon Zator (Figure 3B), were

consistently active in all populations and across both mutagens. Zator was responsible for over half

of the TE-associated SVs in CB4856, N2 and formaldehyde-mutagenized AB1 populations.

Mutagenesis induced large SV mutations

We found that while most SVs were small (<10^4 bp), both formaldehyde and EMS induced numerous

large mutations and affected substantial portions of the genome (Figure 4). In the N2 formaldehyde-

treated population, insertion mutations created 471,238 nucleotides of new genetic material,

approximately 0.48% of the 100.8 Mb *C. elegans* genome (Supplementary Material Table 3). In

comparison, 74,044 nucleotides (0.06%) were deleted, 44,038 nucleotides (0.04%) were duplicated,

and 1,289,368 nucleotides (1.29%) were affected by inversions. Mutations in the EMS-mutagenized

N2 populations were of similar scale.

AB1 populations showed the smallest number of nucleotides affected by SV mutations. In

formaldehyde-treated AB1, 35,468 nucleotides were inserted and 46,363 deleted. The notable

exception was inversion mutations, which spanned 2,539,375 nucleotides in formaldehyde-treated

and 4,181,603 nucleotides in EMS-treated populations. CB4856 populations harbored substantially

more DNA affected by structural variation. Roughly 1 Mb of new DNA was inserted under both

treatments (Supplementary Material Table 3). Compared to AB1, CB4856 had 14.11-fold more

insertions under formaldehyde and 15.18-fold more under EMS. Relative to N2, it had 1.50-fold and

1.66-fold more insertions under formaldehyde and EMS, respectively. CB4856 also had large
amounts of duplicated DNA: 1.2 Mb under formaldehyde and 2.1 Mb under EMS. Strikingly,
inversion mutations accounted for 49.7 Mb—nearly half the genome—in formaldehyde-treated
CB4856, including large 13 Mb and 8 Mb inversions, with additional megabase-scale and smaller
rearrangements. In EMS-treated CB4856, inversions spanned 32.8 Mb—almost a third of the
genome (Figure 4).
203
SVs and transposon activity varied across chromosomes
The distribution of SVs varied across chromosomes, regardless of strain or mutagen (Figure 5A). SV
mutations occurred in higher numbers on Chromosome IV and V and lower numbers on
Chromosome X. TE activity did not explain these patterns as, for example, TEs were most active on
Chromosome IV in CB4856 but Chromosome V harbored the highest number of SVs (Figure 5B). TE-
associated duplications and inversions were less frequent than deletions across all populations.
Additionally, we found shared transposition events between the two mutagens. The highest overlap
between formaldehyde and EMS TE-associated mutations was observed in N2 where 33.3% of the
TE-associated mutations were shared. In CB4856 17.9% of the TE-associated mutations were shared
between EMS and formaldehyde treated populations while in AB1 it was just 4.5%. These findings
indicate that while TEs contribute to structural variation, they do not account for the full spectrum
of observed SVs.
216
Simulation support of experimental results
Although there is a body of work modelling the influence of outcrossing on mutation in populations
these models typically assume that mutations are occurring at the level of an individual SNP or
broader, abstract 'allele' level (Kondrashov, 1982; Charlesworth, 1990a). To explore whether our
experimental results could emerge under different evolutionary scenarios, we developed a
population model describing evolution of SVs in the genome. We implemented this model in
stochastic simulations using the SLiM 4.3 software (Haller & Messer, 2023) and measured the
impact of outcrossing rates on SVs in the population.
Consistent with our experimental observations, the simulations showed that outcrossing
populations retained SVs, particularly when outcrossing rates consistently exceeded 30%. In
contrast, fully selfing populations rapidly purged SVs. SVs were retained in outcrossing populations
when each mutation carried a small deleterious effect on fitness, drawn from a gamma distribution
(Supplementary Figure 1; Eyre-Walker & Keightley, 2007; Keightley & Eyre-Walker, 2007;
Racimo & Schraiber, 2014). The fitness landscape had a negative curvature or synergistic epistasis
(Supplementary Figure 2). Under these conditions, SVs persisted in simulated populations. These

results suggest that outcrossing can, under specific conditions, promote the persistence of structural

mutations (Figure 6).

<u>Discussion</u>

In this study we investigated how quantitative levels of outcrossing influence the genome's ability to

purge deleterious mutations and how this ability varies with mutation size and genomic location.

Leveraging the unique mating system of *C. elegans*, we combined experimental evolution with

genomic analyses to assess how outcrossing influences the purging of mutations of different sizes

and locations. We found that CB4856, characterized by highest male frequency, outcrossing rate and

elevated TE activity, regained fitness after mutagenesis and recovery. Despite this, CB4856

populations retained large numbers of *de novo* SNPs and SVs. Importantly, we found many SNPs

were trapped within SVs and overlapped with exonic regions, making it difficult to purge these

variants through recombination and outcrossing.

C. elegans is one of the most tractable model systems and has emerged as a powerful model

organism for unraveling the intricate dynamics of mutation accumulation. Traditionally, researchers

have employed two primary approaches to investigate this phenomenon. The first approach involves

manipulating DNA repair pathways, creating "mutation accumulation" lines with compromised

repair mechanisms (Estes et al., 2024). This allows for controlled observation of how increased

mutation rates impact population fitness and adaptation over generations. The second approach

takes a more observational stance, tracking the accumulation and consequences of naturally

occurring mutations over extended periods (Konrad et al., 2017). Both approaches have yielded

valuable insights into the remarkable ability of *C. elegans* populations to tolerate and potentially

recover from mutational stress. However, despite these advancements, a crucial aspect remains

underexplored: how mating system variation and reproductive strategies interact with mutation size

and the resulting impacts on genomic variation.

We explored these factors in our population simulations and found that under specific

assumptions increased levels of outcrossing would retain SVs. Importantly, the phenotypic effect

size of the segregating mutations was very small. In a fully outcrossing simulated population, there

was an average of 10 mutations per individual after 100,000 generations, each with an average

phenotypic effect size of 0.0035. Larger mutation effect sizes were rapidly purged from outcrossing

populations. However, deleterious mutations with small effects on the phenotype and fitness were

stably maintained in the simulated outcrossing populations for long periods.

The mechanisms by which genomes cope with mutational threats are multifaceted and

depend heavily on the type of lesion incurred. Point mutations like SNPs predominantly affect gene

expression and protein structure and are typically repaired through efficient pathways like base
excision repair (BER) or nucleotide excision repair (NER). In contrast, complex lesions like SVs
(structural variants) pose a more significant challenge to genome stability, often leading to
chromosomal aberrations (Ciccia & Elledge, 2010). Repairing these intricate injuries often relies
on more intricate mechanisms like homologous recombination (HR) and non-homologous end
joining (NHEJ) (Limpose et al., 2017). In our study we used two types of mutagens that are known to
affect the genome in different ways. EMS is known to cause more point mutations and
formaldehyde is known to cause complex lesions. We expected that EMS should be creating more
SNPs and formaldehyde should be causing more structural variations in the genome. However, we
found that long term mutagenic abuse (5 consecutive generations in our case) with either of the
mutagen had similar mutational effects on the *C. elegans* genome and resulted in SNPs and SVs
ranging from 10s of nucleotides to multiple megabases in length (Supplementary Material Table 3).

The potential impact of male frequency and outcrossing on mutational burden remains
highly debated. Studies in dioecious species suggest that intense male competition for mates drives
stronger selection against deleterious mutations, potentially benefiting populations facing genomic
stress (Whitlock & Agrawal, 2009). However, bottlenecks arising from such strong selection can
ironically counteract this advantage by allowing the accumulation of detrimental mutations.
Additionally, the relevance of these findings to androdiecious species like *C. elegans* is unclear. Due
to their self-fertilizing nature, *C. elegans* males might not experience the same level of selective
pressure as those in dioecious species (Artieri et al., 2008). Consequently, increased male frequency
in *C. elegans* might not directly translate to mutational resilience. Nevertheless, higher male
proportions could offer another advantage: increased outcrossing, which can introduce new genetic
variation. This influx of novel alleles could enhance the population's adaptability and resilience in the
face of unpredictable challenges like ongoing mutagenic stress. Moreover, mutation purging
processes differ significantly between self-fertilizing and outcrossing mating systems, particularly for
SNPs and SVs. In self-fertilizing populations, increased homozygosity facilitates rapid purging of
highly deleterious mutations from the genome. However, reduced genetic variation weakens
selection against slightly deleterious mutations, potentially leading to the accumulation of slightly
deleterious SNPs and SVs. In contrast, outcrossing populations can maintain both highly deleterious
and slightly deleterious mutations if they are recessive.

Our results suggest that there are strain- and mutagen-specific effects, with CB4856 showing
the highest male frequency among the three strains. However, it exhibited the worst relative fitness
after three generations of recovery from mutagenic treatment and carried the highest mutational
load for both mutagens. AB1, in contrast, had second largest male frequency but was least affected

by mutagenic abuse and had the minimum mutagenic load in the genome when compared with the
other two strains. N2 had the lowest male frequency but the highest relative fitness during
mutagenesis; however, its fitness declined during recovery, suggesting that the elevated fitness
under mutagenic stress may have come with costs that became apparent in a stable environment.

Strain-specific differences in coping with mutagenic stress may reflect distinct genomic
architectures. For example, AB1 had few SNPs within SVs and the lowest exonic overlap of all strains,
potentially minimizing disruption to coding regions. This pattern could act as a genomic "buffer,"
reducing the functional impact of mutations and contributing to resilience after mutagenic stress.

CB4856 had the highest number of de novo SNPs and SVs. A high proportion of SNPs were
trapped within SVs and a significant number of these mutations intersected with exonic regions,
potentially disrupting essential genes. This genomic configuration likely explains CB4856's reduced
resilience under mutagenic stress, as the interplay between structural and single nucleotide variants
can amplify the deleterious effects of mutations, complicating the purging process. The broad and
evenly distributed mutational landscape especially for SV ranging from 1,000 base pairs to 10^7 base
pairs (Fig. 4) may favor greater genomic plasticity, improving a species generation of genetic
variation and potential adaptation in uncertain environments. However, CB4856 experienced the
most pronounced decline in fitness following mutagenesis. The delicate balance between
evolutionary plasticity and mutational resilience may tip toward instability under the stress of
chemical mutagenesis.

N2 populations showed relatively stable relative fitness responses without significant
increases in male frequencies or outcrossing frequencies and relatively low genomic responses to
mutagenesis. This suggests that N2 relied heavily on DNA repair mechanisms, rather than increased
genetic variation through outcrossing, to mitigate the negative effects of induced mutations. The
combination of self-fertilization and the retention of small SVs may minimize genomic disruptions,
ensuring genomic integrity over generations. This may balance minimizing risk through self-
fertilization and small SV accumulation, while still allowing for some level of genomic flexibility. This
type of response may allow an organism to persist in environments where maintaining genomic
integrity is essential for survival. Interestingly, the relative fitness of N2 with both the mutagens
increased which came back to the pre-mutagenizes levels after recovery phase. This suggest that the
initial increase in fitness might come with a long-term cost, potentially limiting its adaptive potential
over time.

Transposable elements (TEs) can play a critical role in shaping genomic resilience by
introducing genetic variation and driving structural changes in response to stress (Negi et al., 2016).
Increased outcrossing has been associated with elevated TE activity, likely due to the enhanced

recombination rates and reduced efficiency of silencing mechanisms in outcrossing populations
(Dolgin & Charlesworth, 2006; Wright & Schoen, 1999). Our results highlight significant strain-
specific differences in TE activity post-mutagenesis. TEs like Tc1, known for their cut-and-paste
transposition mechanism and historical activity in *C. elegans* (Plasterk, R. H. A, 1996), were not the
dominant contributors to SVs in our findings. In fact, in N2 and AB1 under formaldehyde treatment,
Tc1 was entirely absent, yet these strains still retained substantial structural variation, indicating
that SV retention cannot be attributed solely to Tc1 or any other single transposable element's
activity. CB4856, with the highest outcrossing frequency, exhibited the highest transposition rates,
reflecting its propensity for increased genomic instability under mutagenic stress. Interestingly,
although AB1 had the highest number of annotated transposons, its transposition rates were lower,
suggesting fewer active TEs. Elevated TE activity in CB4856, coupled with unique transposons such as
Copia, Gypsy, and Sola, may indicate a potential for adaptability through increased genetic variation
and a capacity to respond to environmental challenges. Interestingly, N2 exhibited the highest
overlap in TE events between mutagens (33.3%), suggesting common TE activations under stress.
Across all strains, Chromosome IV emerged as a TE hotspot, likely reflecting shared genomic
characteristics that tolerate or facilitate transposition in this region.

Increased outcrossing enhances recombination rates, creating genomic instability that can
activate TE mobilization (Arkhipova & Meselson, 2000). Additionally, the introduction of genetic
variation through outcrossing may temporarily disrupt TE silencing mechanisms, further contributing
to higher TE activity in CB4856 (Hollister & Gaut, 2009). These results indicate that while higher
outcrossing may promote the generation of genetic diversity, it can also trigger processes that
increase susceptibility to mutagenic stress. This susceptibility may stem from strain-specific genomic
architectures, the accumulation of deleterious mutations, or the difficulty of purging large-scale
genomic changes under conditions of frequent outcrossing.

In conclusion, we found strain specific fitness and mutation responses which may reflect
larger patterns of evolutionary response. Although the N2 "Bristol" strain, originally collected by
Sydney Brenner (Brenner 1974) has been developed as a model system, its fitness, behavior and
genomics differ substantially from other strains of *C. elegans*. N2 is a highly lab adapted strain with
little genetic variation, few males and low outcrossing frequencies (Sterken et al., 2015). In contrast,
CB4856 and AB1 are more recently collected "wild worms" with substantial genomic differences
(Thompson et al., 2015) and high natural variation in male frequency (Anderson et al., 2010) and
outcrossing (Teotonio et al., 2012). Additionally, our study reveals that SVs can act as genomic traps
for SNPs thereby shielding these mutations from purging through recombination and outcrossing.
This mechanism was most prominent in outcrossing-prone strains, suggesting that higher rates of

recombination may not always facilitate genome cleanup, particularly for large or complex
mutations. The natural genetic variation in *Caenorhabditis* remains a powerful resource for
dissecting how genome architecture and mating systems interact to shape evolutionary outcomes.
371

Funding
This work was supported by NSF award 2225796 and NIGMS award R35GM147245 to J.L.F.

Data Availability
Scripts, bioinformatic workflows and analyses are available at github.com/RohitKapila. Python code
used for image analysis is available at github.com/FierstLab/WormImage. Fitness data are available
as Supplemental Materials associated with the manuscript. Sequence reads are available at the NCBI
SRA under BioProject PRJNA1201181.
380

Methods
Strains and mutagens
Three *C. elegans* strains with varying male frequencies were chosen: N2 (Bristol), CB4856 (Hawaiian),
and AB1. ST2 served as a common competitor in relative fitness assays. All strains were obtained
from the Caenorhabditis Genetics Center and maintained frozen until the experiment. Frozen strains
were thawed on 75 mm × 13 mm NGM plates seeded with *E. coli* OP50. After one generation (~3.5
days at 20°C) for recovery, each strain was divided into nine replicate populations. Prior to the
experiment, replicates were synchronized by bleaching. Over five generations, four replicates per
strain were mutagenized with formaldehyde, four with EMS, and one remained untreated as a non-
mutagenized control. All replicates for a given strain were handled in parallel on the same day using
the same batch of plates to minimize intra-strain variability due to plate effects. Following the
mutagenesis phase, all populations were frozen and stored at −80°C for downstream analyses.
393

Mutagenesis
Four replicates per strain were treated with 1 mM ethyl methanesulfonate (EMS) in M9 solution,
while the other four replicates were treated with 0.1% (w/v) formaldehyde solution in M9. These
concentrations were chosen because they are non-lethal yet sufficient to induce detectable genomic
mutations. During both the mutagenesis and recovery phases, worms were maintained on standard
laboratory plates. However, during the mutagenesis phase, worms were suspended in mutagen
solutions for 4 hours every fourth day before being transferred back to standard plates. In contrast,
during the recovery phase, worms were not exposed to the mutagen and were allowed to grow

normally on standard laboratory plates. We followed the worm mutagenesis protocol described by

Brenner et al. (1974) with minor modifications. Every generation (~4 days), worms were washed

from the plates with M9, collected in tubes, and resuspended in their respective mutagen solutions

for 4 hours at 20 °C, during which they were gently rocked to ensure uniform exposure. After

treatment, worms were transferred to fresh plates with NGM and *E. coli* OP50, ensuring

maintenance of replicate identity and avoidance of mixing.

Population sizes were not altered at any point during the experiment. During both transfers,

from the plates to the tubes and from the tubes back to the plates, all worms were transferred using

300 microliters of liquid. Specifically, M9 buffer was used for the transfer from plates to tubes, and

the mutagen was used for the transfer from tubes back to plates. After the mutagen treatment, the

liquid was carefully transferred onto the plate, causing parts of the *E. coli* colony to be dislodged. As

a result, the worms would initially swim on the plate. We then left the plate open, allowing the

liquid to be absorbed within 30-40 minutes, after which the worms resumed crawling on the plate.

Male frequency assay

To evaluate the impact of five generations of mutational exposure on male frequency, a male

frequency assay was conducted using both non-mutagenized (parent) worms and worms

immediately after five generations of mutagen treatment. Worms from each replicate population

were bleached for age synchronization. The lifetime progeny production of worms was monitored by

placing a single L4 worm on a 75mm x 13mm Petri plate, transferring it to a fresh plate every 24

hours. After 96 hours, the worm was discarded, and the progeny produced over the four days were

sexed and counted. Any worms that died during the transfer process were excluded from the

analysis.

Outcrossing frequency assay

As with the male frequency assay, outcrossing frequency was assessed using the same set of

replicate plates. The lifetime progeny production of each worm was monitored by placing a single L4

worm on a 75mm x 13mm Petri plate, transferring both male and hermaphrodite to a fresh plate

every 24 hours. After 96 hours, both the male and hermaphrodite were discarded, and the progeny

produced over the four days were allowed to grow to adulthood.

Outcrossing in *C. elegans* is known to produce approximately 50% male progeny(Yin &

Haag, 2019). Based on male frequency assays, we determined that spontaneous male production

in the N2 and AB1 strains, both before and after mutagenesis, was relatively low. Thus, plates for

these strains were visually scanned for males, and plates with male frequencies exceeding the

expected spontaneous range (1%–5%) and approaching 50% were categorized as outcrossing plates.

For the CB4856 strain, which naturally exhibits higher baseline male frequencies (15–17%), male
frequency was calculated for each plate. A threshold of 25% males was set, and plates with male
frequencies meeting or exceeding this value were marked as outcrossing plates.

Relative fitness assay
After five generations of mutagenesis, worms were frozen at -80°C and later recovered for relative
fitness assay at sixth generation- representing the fitness immediately after five generations of
mutagenesis and again after three generations of recovery from mutations using the standard _C._
elegans maintenance protocol at 20 °C. During recovery, worms were maintained using the standard
worm maintenance protocol. For the relative fitness assay, one early-stage L1 worm from each focal
strain (N2, CB4856, AB1) was paired with one L1 worm from the non-mutagenized common
competitor strain ST2 expressing green fluorescent protein (GFP). Both strains were co-cultured on
NGM plates with _E. coli_ OP50 for eight days (~2 generations) at 20 °C. Washed worms from replicate
plates (11 per generation/mutant/strain combination) were collected in M9 on a slide, imaged with
and without a GFP filter on a fluorescent microscope, and counted using a Python-based code
specific for worm morphology (available on the Github). Plates with zero worms of either type were
excluded from analysis, assuming that the worm did not survive to lay eggs. Images captured
without the GFP filter counted all worms present on the plate- both the focal strain and the common
competitor strain. Conversely, images captured with the GFP filter only revealed the green,
fluorescent competitor worms, allowing for estimation of their abundance relative to the total worm
population.

Fitness was compared across non-mutagenized, post-mutagenized, and recovered lineages using
automated image analysis (Supplementary Material Table 2). To validate the accuracy of the image
analysis pipeline, we randomly selected a subset of samples and manually counted individuals. This
confirmed that the automated method produced reliable counts, with no observed false positives or
false negatives. We found that three generations of recovery from mutagenesis were sufficient for
each strain to recover pre-mutagenesis fitness levels for both the mutagens (Figure 1B).

DNA extraction and Sequencing
To assess the genomic impact of mutagenesis, high molecular weight DNA was extracted from
worms (after 3 generations of recovery) using the Cytiva kit (detailed procedure available in the
Supplementary Materials). Before starting out the DNA extraction we washed a minimum of
~10,0000 worms with M9 buffer from the plate into a tube and kept the worms rocking overnight to

remove associated microbiome. Pooled PacBio long-read sequencing was performed for parental
worms of all three strains (N2, CB4856, AB1) and one population per strain per mutagen.
Additionally, Illumina short-read sequencing was conducted for each replicate of each treatment
(control, EMS, formaldehyde) for all strains. This strategy enabled comprehensive analysis of
genomic variants induced by mutagenesis and their phenotypic consequences in subsequent
generations.

For the comprehensive genomic analysis, we sequenced a total of 36 samples. Using PacBio
long-read sequencing, with an average depth of coverage of 94.52x, we sequenced nine samples:
the three parental strains (N2, CB4856, AB1) as controls, and six mutagenized replicates i.e., one
replicate per strain treated with each mutagen. Additionally, we performed Illumina short-read
sequencing on 27 treatment replicates, achieving an average depth of coverage of 72.11x. These 27
samples included the three parental strains, 12 EMS-treated replicates, and 12 formaldehyde-
treated replicates, with four replicates per treatment for each strain.

The male frequency and outcrossing rate assays were conducted on worms derived from the
same batch that underwent five generations of mutagenesis. In contrast, the worms used for the
relative fitness assay and DNA extraction came from a different set, although they were
mutagenized in the same manner as the first batch.

Population simulations
To quantitatively examine the influence of outcrossing on SV purging and retention we implemented
our theoretical model in population simulations using the software SLiM 4.3 (Haller & Messer,
2023). Our model described a genomic system where mutations occur as SVs, either insertions,
deletions or inversions. Each individual in a population of size K carried n structural mutations. We
constructed dioecious populations containing equal numbers of males and females and self-fertile
populations with a proportion of individuals reproducing through outcrossing and a proportion
reproducing through selfing. Each population started at a specified size K and evolved through
overlapping generations with reproduction and fitness differences creating variable population sizes.
Each individual had a diploid chromosome 10,000 nucleotides in length. Recombination occurred
along the chromosomes with uniform probability at a specified rate r. After reproduction SV
mutations occurred at a rate μ and were removed at a rate ν, for example by back mutation via gene
conversion (Katju et al., 2008).

We did not explicitly model the physical size of mutations but assumed that mutations had
discrete phenotypic effect sizes that cumulatively influenced fitness. The probability of a SV
mutation occurring at any location was uniform across the chromosome and each SV mutation

conferred a small deleterious phenotypic effect drawn from $\Gamma(k,\theta)$ (Supplementary Figure 1;

Eyre-Walker & Keightley, 2007; Keightley & Eyre-Walker, 2007; Racimo & Schraiber, 2014).

Individual SVs interacted to decrease fitness faster-than-linearly and the fitness landscape had a

negative curvature or synergistic epistasis (Supplementary Figure 2). The fitness of an individual

carrying n SVs was $w(n) = exp^{-(an+(bn^2)/2)}$ where a and b were parameters determining the

curvature of the fitness landscape (Charlesworth 1990). Our simulated populations evolved for

100,000 generations with the parameters $k = 1, \theta = 2, \mu = 0.01, v = 0.001, a = 0.001$ and

$b = 0.0006$. We evolved populations under a range of evolutionary scenarios including

recombination rates r=0.5-1x10^{-5}, K=100 to 5,000 and levels of self-fertility ranging from 0 to 100%.

Data analysis

<u>Male frequency assay</u>

Male frequency was calculated as the ratio of male progeny to the total progeny laid by each worm,

and this proportion was used as the unit of analysis. We assessed the normality of the data using the

Shapiro-Wilk test and by visualizing its distribution with a histogram. As expected, the distribution

was right-skewed, with a high proportion of zero values, indicating zero-inflation. Given the bounded

nature (0 to 1) of the data and the excess zero values, we used a zero-inflated Beta regression for

analysis. Since this model requires proportion values strictly between 0 and 1, we applied an epsilon

adjustment by adding a small constant to all zero values to make the data more appropriate for the

model. We performed the analysis using the glmmTMB package in R (R version 4.1.0, released 2021-

05-18)(Bolker, 2022), implementing the following model:

Male Frequency~Strain× Mutagen

In this model Strain (N2, AB1, CB4856) and Mutagen treatment (EMS, Formaldehyde and

non-mutagenized parents were included as fixed factors and an interaction term was added to check

if the effect of strain on male frequency varied by mutagen identity. This model accounts for both (1)

the continuous variation in male frequency and (2) the excess zero values separately, making it a

better fit than standard regression models. Further we did Tukey's HSD comparison (Abdi &

Williams, 2010) using emmeans package in R (Searle et al., 1980).

<u>*Relative fitness assay*</u>

To quantify the relative fitness of the *C. elegans* strains N2, CB4856, and AB1, each plate was washed

with M9 buffer solution to prepare the worms for imaging. Two aliquots from the washed plate

were then placed onto a microscope slide and covered with a coverslip. We performed imaging to

count the GFP-expressing competitor strain and the non-GFP-expressing focal strain worms within

these aliquots. The allelic frequency of the non-fluorescent allele was calculated as the square root

of proportion of non-GFP-expressing worms to the total worm population. For statistical analysis,

two-way ANOVAs were conducted separately for each mutagen to assess the effects of strain and

treatment duration on relative fitness. Data normality was examined using the Shapiro-Wilk test and

the residual distribution showed slight variation from normality (Shapiro-Wilk test: W = 0.9868, $p <$

0.01), indicating variation from normality. However, visual inspection of the residuals with histogram

suggested an approximately normal distribution. Since ANOVA is robust to minor deviations, we

proceeded with ANOVA followed by pairwise comparisons using Tukey's HSD test. The following

linear model was applied:

anova(lm(RF ~ Strain*Generation, data = a)).

Following ANOVA, we performed pairwise comparisons using Tukey's Honest Significant

Difference (HSD) test to assess differences in relative fitness across strains and generations. We

conducted separate Tukey HSD tests within each strain to compare relative fitness before

mutagenesis, after five generations of mutagenesis, and after three generations of recovery.

Bioinformatic analysis

The bioinformatic analysis for identifying *de novo* mutations following PacBio sequencing involved a

comprehensive workflow to process and analyse long-read data, distinguishing mutations that arose

in the worms after mutagenesis from those existing in parental strains. Initially, raw PacBio reads

were aligned and indexed to reference genomes using minimap2 (Li, 2018) and bcftools (Danecek et

al., 2021); for the N2 and CB4856 strains, NCBI reference genomes were used, while the AB1 strain

employed the recently annotated genome of its descendant strain, SX3368 (Riccio C, 2022), to

account for genomic divergences within the lineage. Subsequent variant calling was performed using

Sniffles (Smolka et al., 2024) for structural variants (SVs), such as insertions, deletions, and

inversions, capitalizing on the accuracy offered by long-read data.

For the specific identification of de novo mutations induced by mutagenesis, a comparative

analysis using bcftools contrasted the variant profiles of generation 8 worms (post three generations

of recovery) with their control parents, filtering out pre-existing genetic variations. All parental and

mutagenized samples were processed using identical library preparation, sequencing, and

bioinformatics pipelines to ensure comparability in variant detection. Further quality control and

data filtering ensured only high-quality SNPs and SVs (quality score ≥ 20, depth ≥ 10) were retained.

We then used bedtools to provide insights into their potential impacts on exonic regions and

regulatory elements.

Additionally, similar bioinformatic steps were conducted with Illumina sequencing data.
Alignment and indexing were again performed on reference genomes, followed by SNP and SV
calling using Freebayes (Ypsilanti et al.,2021)and Manta (Chen et al., 2016), respectively. We initially
called variants in pooled-population mode (FreeBayes 'pooled') to account for minor alleles in the
combined worm samples. However, when intersecting these SNP calls with structural variants, we
observed inflated counts due to multi-allelic sites being represented multiple times. To avoid this
over-counting artifact, we generated a separate VCF using non-pooled calling parameters and used
these calls for all subsequent intersection and annotation analyses. Enhanced SV detection was
achieved with VGtoolkit (Hickey et al., 2020) through graph-based variant calling, which was then
integrated with SVs identified by Sniffles 2.0 using Survivor 1.0.3 (Jeffares et al., 2017), creating a
comprehensive view of the genomic alterations induced by the mutagens.

Long-read sequencing was performed on the three non-mutagenized parent strains and
one replicate per strain for each mutagen treatment. Short-read sequencing covered all replicates
across all strains. For structural variation analysis, we merged variants identified by both long-read
and short-read callers using Survivor 1.0.3. For SNP analysis, we relied solely on variants identified by
short-read callers.

<u>Annotation of Transposable Elements and Intersection with Structural Variants</u>
Transposon annotation is facilitated in complete and contiguous sequences, thus annotation of TEs
was done in the N2, AB, and CB reference genomes rather than directly from the raw sequence data.
We used the TransposonUltimate reasonaTE (Riehl et al., 2022) pipeline for TE annotation in each of
the reference genomes. Briefly, TEUltimate reasonaTE wraps 13 previously published TE annotation
tools, which include: RepeatMasker v4.1.1 **,** RepeatModeler v2.0.1, SINE-Scan , SINE-Finder, LTRPred
, LTRHarvest, TIRvish , MUSTv2 , MITE-Tracker, MITEFinderII**,** HelitronScanner, TransposonPSI for
transposon discovery. While some functions are optional, all subprograms were run with default
parameters in this study. The resulting annotations are filtered, merged, and clustered using CD-HIT
v4.8.1 (Fu et al., 2012) and BLASTN v2.10.1 (Camacho et al., 2009). The final annotations from
FinalAnnotations_Transposons.gff3 in the final Results folder were used for downstream analysis for
each of the three focal strains.

Evidence of transposon movement was assumed from structural variants which overlapped
with transposon annotations. The TEUltimatedeTEct pipeline (Riehl et al., 2022) was used for
intersection of transposon and structural variant annotations. De novo VCF structural variant files, as
generated above, were intersected with transposon annotations and reported as an event if
overlapping by at least 10% and lengths were similar by at least 50%. Finally, any annotation smaller

than 50 base pairs, or larger than 1% of the genome was discarded. Plots were generated from the

transpositionEvents.gff3 file in RStudio with R v4.3.2 (R Core Team, 2023).

<u>Literature Cited</u>

Abdi, H., & Williams, L. J. (2010). Principal component analysis. In *Wiley*
Interdisciplinary Reviews: Computational Statistics (Vol. 2, Issue 4, pp. 433–
459). https://doi.org/10.1002/wics.101

Anderson, J. L., Morran, L. T., & Phillips, P. C. (2010). Outcrossing and the
maintenance of males within C. elegans populations. *Journal of Heredity*,
101(SUPPL. 1). https://doi.org/10.1093/jhered/esq003

Arkhipova, I., & Meselson, M. (2000). *Transposable elements in sexual and ancient*
asexual taxa. Harvard School of Public Health. www.pnas.org

Artieri, C. G., Haerty, W., Gupta, B. P., & Singh, R. S. (2008). Sexual selection and
maintenance of sex: Evidence from comparisons of rates of genomic
accumulation of mutations and divergence of sex-related genes in sexual and
hermaphroditic species of Caenorhabditis. *Molecular Biology and Evolution*,
25(5), 972–979. https://doi.org/10.1093/molbev/msn046

Bolker, B. (2022). *Getting started with the glmmTMB package*.

Brenner S. (1974). The genetics of *Caenorhabditis elegans*. Genetics 77: 71-94.

Byers, D. L., & Waller, D. M. (1999). Do Plant Populations Purge Their Genetic
Load? Effects of Population Size and Mating History on Inbreeding Depression.
In *Source: Annual Review of Ecology and Systematics* (Vol. 30).
https://about.jstor.org/terms

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., &
Madden, T. L. (2009). BLAST+: Architecture and applications. *BMC*
Bioinformatics, *10*. https://doi.org/10.1186/1471-2105-10-421

Charlesworth, B. (1990a). Mutation-selection balance and the evolutionary
advantage of sex and recombination. *Genetical Research*, *55*(3), 199–221.
https://doi.org/10.1017/S0016672300025532

Charlesworth, B. (1990b). OPTIMIZATION MODELS, QUANTITATIVE GENETICS,
AND MUTATION. *Evolution*, *44*(3), 520–538. https://doi.org/10.1111/j.1558-
5646.1990.tb05936.x

Charlesworth, B., & Charlesworth, D. (1983). The population dynamics of
transposable elements. *Genetical Research*, *42*(1), 1–27.
https://doi.org/10.1017/S0016672300021455

Chen, X., Schulz-Trieglaff, O., Shaw, R., Barnes, B., Schlesinger, F., Källberg, M.,
Cox, A. J., Kruglyak, S., & Saunders, C. T. (2016). Manta: Rapid detection of
structural variants and indels for germline and cancer sequencing applications.
Bioinformatics, *32*(8), 1220–1222. https://doi.org/10.1093/bioinformatics/btv710

Ciccia, A., & Elledge, S. J. (2010). The DNA Damage Response: Making It Safe to
Play with Knives. In *Molecular Cell* (Vol. 40, Issue 2, pp. 179–204).
https://doi.org/10.1016/j.molcel.2010.09.019

Collins, R. L., Brand, H., Karczewski, K. J., Zhao, X., Alföldi, J., Francioli, L. C.,
Khera, A. V., Lowther, C., Gauthier, L. D., Wang, H., Watts, N. A., Solomonson,
M., O'Donnell-Luria, A., Baumann, A., Munshi, R., Walker, M., Whelan, C. W.,
Huang, Y., Brookings, T., … Talkowski, M. E. (2020). A structural variation
reference for medical and population genetics. *Nature*, *581*(7809), 444–451.
https://doi.org/10.1038/s41586-020-2287-8

Danecek, P., Bonfield, J. K., Liddle, J., Marshall, J., Ohan, V., Pollard, M. O.,
Whitwham, A., Keane, T., McCarthy, S. A., & Davies, R. M. (2021). Twelve
years of SAMtools and BCFtools. *GigaScience*, *10*(2).
https://doi.org/10.1093/gigascience/giab008

Dolgin, E. S., & Charlesworth, B. (2006). The fate of transposable elements in
asexual populations. *Genetics*, *174*(2), 817–827.
https://doi.org/10.1534/genetics.106.060434

Estes, S., Phillips, P. C., Denver, D. R., Thomas, W. K., & Lynch, M. (2004).
Mutation Accumulation in Populations of Varying Size: The Distribution of
Mutational Effects for Fitness Correlates in Caenorhabditis elegans.

Eyre-Walker, A., & Keightley, P. D. (2007). The distribution of fitness effects of new
mutations. In *Nature Reviews Genetics* (Vol. 8, Issue 8, pp. 610–618).
https://doi.org/10.1038/nrg2146

Feuk, L., Carson, A. R., & Scherer, S. W. (2006). Structural variation in the human
genome. In *Nature Reviews Genetics* (Vol. 7, Issue 2, pp. 85–97).
https://doi.org/10.1038/nrg1767

Fu, L., Niu, B., Zhu, Z., Wu, S., & Li, W. (2012). CD-HIT: Accelerated for clustering
the next-generation sequencing data. *Bioinformatics*, *28*(23), 3150–3152.
https://doi.org/10.1093/bioinformatics/bts565

Gaertner, B. E., & Phillips, P. C. (2010). Caenorhabditis elegans as a platform for
molecular quantitative genetics and the systems biology of natural variation.
Genetics Research, *92*(5–6), 331–348.
https://doi.org/10.1017/S0016672310000601

García-Dorado, A. (2012). Understanding and predicting the fitness decline of shrunk
populations: Inbreeding, purging, mutation, and standard selection. *Genetics,*
190(4), 1461–1476. https://doi.org/10.1534/genetics.111.135541

Glémin, S. (2003). How are deleterious mutations purged? Drift versus nonrandom
mating. *Evolution, 57*(12), 2678–2687. https://doi.org/10.1111/j.0014-
3820.2003.tb01512.x

Haller, B. C., & Messer, P. W. (2023a). SLiM 4: Multispecies Eco-Evolutionary
Modeling. *American Naturalist, 201*(5), E127–E139.
https://doi.org/10.1086/723601

Hartfield, M. (2016). Evolutionary genetic consequences of facultative sex and
outcrossing. *Journal of Evolutionary Biology, 29*(1), 5–22.
https://doi.org/10.1111/jeb.12770

Hickey, G., Heller, D., Monlong, J., Sibbesen, J. A., Sirén, J., Eizenga, J., Dawson,
E. T., Garrison, E., Novak, A. M., & Paten, B. (2020). Genotyping structural
variants in pangenome graphs using the vg toolkit. *Genome Biology, 21*(1).
https://doi.org/10.1186/s13059-020-1941-7

Ho, S. S., Urban, A. E., & Mills, R. E. (2020). Structural variation in the sequencing
era. In *Nature Reviews Genetics* (Vol. 21, Issue 3, pp. 171–189). Nature
Research. https://doi.org/10.1038/s41576-019-0180-9

Hoffmann, A. A., & Rieseberg, L. H. (2008). Revisiting the impact of inversions in
evolution: From population genetic markers to drivers of adaptive shifts and
speciation? In *Annual Review of Ecology, Evolution, and Systematics* (Vol. 39,
pp. 21–42). https://doi.org/10.1146/annurev.ecolsys.39.110707.173532

Hollister, J. D., & Gaut, B. S. (2009). Epigenetic silencing of transposable elements:
A trade-off between reduced transposition and deleterious effects on
neighboring gene expression. *Genome Research, 19*(8), 1419–1428.
https://doi.org/10.1101/gr.091678.109

Jeffares, D. C., Jolly, C., Hoti, M., Speed, D., Shaw, L., Rallis, C., Balloux, F.,
Dessimoz, C., Bähler, J., & Sedlazeck, F. J. (2017). Transient structural
variations have strong effects on quantitative traits and reproductive isolation in
fission yeast. *Nature Communications, 8.* https://doi.org/10.1038/ncomms14061

Katju, V., LaBeau, E. M., Lipinski, K. J., & Bergthorsson, U. (2008). Sex change by
gene conversion in a Caenorhabditis elegans fog-2 mutant. *Genetics, 180*(1),
669–672. https://doi.org/10.1534/genetics.108.090035

Keightley, P. D., & Eyre-Walker, A. (2007). Joint inference of the distribution of
fitness effects of deleterious mutations and population demography based on
nucleotide polymorphism frequencies. *Genetics, 177*(4), 2251–2261.
https://doi.org/10.1534/genetics.107.080663

Kirkpatrick, M., & Barton, N. (2006). Chromosome inversions, local adaptation and
speciation. *Genetics*, *173*(1), 419–434.
https://doi.org/10.1534/genetics.105.047985

Kondrashov, A. S. (1982). Selection against harmful mutations in large sexual and
asexual populations. *Genetical Research*, *40*(3), 325–332.
https://doi.org/10.1017/S0016672300019194

Konrad, A., Thompson, O., Waterston, R. H., Moerman, D. G., Keightley, P. D.,
Bergthorsson, U., & Katju, V. (2017). Mitochondrial mutation rate, spectrum and
heteroplasmy in Caenorhabditis elegans spontaneous mutation accumulation
lines of differing population size. *Molecular Biology and Evolution*, *34*(6), 1319–
1334. https://doi.org/10.1093/molbev/msx051

Kutscher, L. M., & Shaham, S. (2014). Forward and reverse mutagenesis in C.
elegans. In *WormBook : the online review of C. elegans biology* (pp. 1–26).
https://doi.org/10.1895/wormbook.1.167.1

Li, H. (2018). Minimap2: Pairwise alignment for nucleotide sequences.
Bioinformatics, *34*(18), 3094–3100. https://doi.org/10.1093/bioinformatics/bty191

Limpose, K. L., Corbett, A. H., & Doetsch, P. W. (2017). BERing the burden of
damage: Pathway crosstalk and posttranslational modification of base excision
repair proteins regulate DNA damage management. In *DNA Repair* (Vol. 56, pp.
51–64). Elsevier B.V. https://doi.org/10.1016/j.dnarep.2017.06.007

Lynch, M., Blanchard, J., Houle, D., Kibota, T., Schultz, S., Vassilieva, L., & Willis, J.
(1999). Perspective: Spontaneous deleterious mutation. *Evolution*, *53*(3), 645–
663. https://doi.org/10.1111/j.1558-5646.1999.tb05361.x

Lynch, M., Burger, R., Butcher, D., Gabriel, W., Charlesworth, B., Hansen, T., Houle,
D., Kondrashov, A., Lande, R., & Willis, J. (1993). The Mutational Meltdown in
Asexual Populations. We are very grateful to. In *Journal of Heredity* (Vol. 84).

MacPherson, B., Scott, R., & Gras, R. (2021). Sex and recombination purge the
genome of deleterious alleles: An Individual Based Modeling Approach.
Ecological Complexity, *45*. https://doi.org/10.1016/j.ecocom.2021.100910

Mahmoud, M., Gobet, N., Cruz-Dávalos, D. I., Mounier, N., Dessimoz, C., &
Sedlazeck, F. J. (2019). Structural variant calling: The long and the short of it. In
Genome Biology (Vol. 20, Issue 1). BioMed Central Ltd.
https://doi.org/10.1186/s13059-019-1828-7

Negi, P., Rai, A. N., & Suprasanna, P. (2016). Moving through the stressed genome:
Emerging regulatory roles for transposons in plant stress response. In *Frontiers
in Plant Science* (Vol. 7, Issue OCTOBER2016). Frontiers Media S.A.
https://doi.org/10.3389/fpls.2016.01448

Ospina, R., & Ferrari, S. L. P. (2012). On Bias Correction in a Class of Inflated Beta
Regression Models. *International Journal of Statistics and Probability*, *1*(2).
https://doi.org/10.5539/ijsp.v1n2p269

Otto, S. P. (2008). Unravelling the evolutionary advantage of sex: A commentary on
'Mutation-selection balance and the evolutionary advantage of sex and
recombination' by Brian Charlesworth. In *Genetics Research* (Vol. 89, Issues 5–
6, pp. 447–449). https://doi.org/10.1017/S001667230800966X

Racimo, F., & Schraiber, J. G. (2014). Approximation to the Distribution of Fitness
Effects across Functional Categories in Human Segregating Polymorphisms.
PLoS Genetics, *10*(11). https://doi.org/10.1371/journal.pgen.1004697

Riccio C. (2022). *Riccio, C., 2022. Duplication is a prominent mechanism of recent
gene birth in Caenorhabditis elegans (Doctoral dissertation).*

Riddle DL, B. T. M. B. et al. ,. (1997). Mechanism of Tc1 Transposition. In *Cold
Spring Harbor Laboratory Press; : Vol. Section VI*. Cold Spring Harbor
Laboratory Press.

Riehl, K., Riccio, C., Miska, E. A., & Hemberg, M. (2022a). TransposonUltimate:
software for transposon classification, annotation and detection. *Nucleic Acids
Research*, *50*(11), E64. https://doi.org/10.1093/nar/gkac136

Robert, F., & Pelletier, J. (2018). Exploring the Impact of Single-Nucleotide
Polymorphisms on Translation. In *Frontiers in Genetics* (Vol. 9). Frontiers Media
S.A. https://doi.org/10.3389/fgene.2018.00507

Searle, S. R., Speed, F. M., & Milliken, G. A. (1980). Population marginal means in
the linear model: An alternative to least squares means. *American Statistician*,
34(4), 216–221. https://doi.org/10.1080/00031305.1980.10483031

Sianta, S. A., Peischl, S., Moeller, D. A., & Brandvain, Y. (2023). The efficacy of
selection may increase or decrease with selfing depending upon the
recombination environment. *Evolution*, *77*(2), 394–408.
https://doi.org/10.1093/evolut/qpac013

Smolka, M., Paulin, L. F., Grochowski, C. M., Horner, D. W., Mahmoud, M., Behera,
S., Kalef-Ezra, E., Gandhi, M., Hong, K., Pehlivan, D., Scholz, S. W., Carvalho,
C. M. B., Proukakis, C., & Sedlazeck, F. J. (2024). Detection of mosaic and
population-level structural variants with Sniffles2. *Nature Biotechnology*.
https://doi.org/10.1038/s41587-023-02024-y

Sterken, M. G., Snoek, L. B., Kammenga, J. E., & Andersen, E. C. (2015). The
laboratory domestication of Caenorhabditis elegans. In *Trends in Genetics* (Vol.
31, Issue 5, pp. 224–231). Elsevier Ltd. https://doi.org/10.1016/j.tig.2015.02.009

Szövényi, P., Devos, N., Weston, D. J., Yang, X., Hock, Z., Shaw, J. A., Shimizu, K.
K., McDaniel, S. F., & Wagner, A. (2014). Efficient purging of deleterious

mutations in plants with haploid selfing. *Genome Biology and Evolution*, *6*(5),
1238–1252. https://doi.org/10.1093/gbe/evu099

Teotonio, H., Carvalho, S., Manoel, D., Roque, M., & Chelo, I. M. (2012). Evolution
of outcrossing in experimental populations of caenorhabditis elegans. *PLoS*
ONE, *7*(4). https://doi.org/10.1371/journal.pone.0035811

Thompson, O. A., Snoek, L. B., Nijveen, H., Sterken, M. G., Volkers, R. J. M.,
Brenchley, R., van't Hof, A., Bevers, R. P. J., Cossins, A. R., Yanai, I., Hajnal,
A., Schmid, T., Perkins, J. D., Spencer, D., Kruglyak, L., Andersen, E. C.,
Moerman, D. G., Hillier, L. D. W., Kammenga, J. E., & Waterston, R. H. (2015).
Remarkably divergent regions punctuate the genome assembly of the
Caenorhabditis elegans hawaiian strain CB4856. *Genetics*, *200*(3), 975–989.
https://doi.org/10.1534/genetics.115.175950

Whitlock, M. C., & Agrawal, A. F. (2009a). Purging the genome with sexual selection:
Reducing mutation load through selection on males. In *Evolution* (Vol. 63, Issue
3, pp. 569–582). https://doi.org/10.1111/j.1558-5646.2008.00558.x

Wright, S. I., & Schoen, D. J. (1999). Transposon dynamics and the breeding
system. In *Genetica* (Vol. 107).

Ypsilanti, A. R., Pattabiraman, K., Catta-Preta, R., Golonzhka, O., Lindtner, S., Tang,
K., Jones, I. R., Abnousi, A., Juric, I., Hu, M., Shen, Y., Dickel, D. E., Visel, A.,
Pennacchio, L. A., Hawrylycz, M., Thompson, C. L., Zeng, H., Barozzi, I., Nord,
A. S., & Rubenstein, J. L. (2021). *Transcriptional network orchestrating regional*
patterning of cortical progenitors. https://doi.org/10.1073/pnas.2024795118/-
/DCSupplemental

Yin, D., & Haag, E. S. (2019). Evolution of sex ratio through gene loss. *Proceedings*
of the National Academy of Sciences of the United States of America, *116*(26),
12919–12924. https://doi.org/10.1073/pnas.1903925116

<u>**Figures**</u>

Figure 1. Male frequency, outcrossing behaviour, and relative fitness reveal strain-specific responses
to mutagenic stress in *C. elegans*. (A) Mean male frequency across strains, represented by solid
circles, before mutagenesis (non-mutagenized) and following EMS or formaldehyde exposure.
CB4856 exhibited the highest male frequency across all treatments, followed by AB1 and N2.
Outcrossing frequency, shown as solid triangles, was consistently higher in CB4856 across
treatments, suggesting strain-specific differences in mating behaviour. (B) Relative fitness across
strains and generations, comparing non-mutagenized (lightest shade), mutagenized (medium
shade), and recovered (darkest shade) populations. AB1 (green) demonstrated the highest post-
mutagenesis resilience, while CB4856 (red) showed the lowest recovery in fitness.

Figure 2: Comparison of de novo SNPs and SVs mutations and their localization in exons across strains (A) Mean counts of de novo SNPs (Circles) and SVs (Triangles) induced by EMS and Formaldehyde across AB1 (Green), CB4856 (Red), and N2 (Blue). De novo SNPs are significantly more abundant than de novo SVs under both mutagen treatments. CB4856 has the maximum number of de novo SNPs and de novo SVs. (B) Mean number of de novo SNPs and de novo SVs localized in exons for the same strains and mutagen treatments, with de novo SNPs again being more prevalent than de novo SVs and CB4856 having more SVs in exons than the other two strains and more SNPs in exons than AB1. Compared to N2, CB4856 had more SNPs in exon than N2 when treated with EMS and comparable SNPs in exon when treated with Formaldehyde.

Figure 3: Mutational and TE landscape across strains and mutagens **(A)**: Number of SNPs, SVs and SNPs in SVs for AB1, CB4856 and N2; **(B)**: Classification and count of active TE families contributing to transposition events across strain-treatment combinations. DNA transposons such as Zator and Tc1-Mariner were among the most active elements, with CB4856 showing the broadest TE activity profile.

859 Figure 4: Size distribution of structural variants (SVs) across *C. elegans* strains and mutagens: Density
860 distribution of SVs sizes for all the three strains for both the mutagens. x- axis Size of SVs, Y-axis:
861 density of SVs.

Figure 5: Chromosomal distribution of SVs and TE-associated SVs across strains and mutagens. (A)
Stacked bar plots showing the total count and types of SVs including Deletions, Insertions,
Duplications, Inversions, Translocations, and Unclassified (NA) across chromosomes 1–5 and X, for
each strain (AB1, CB4856, N2) and mutagen (EMS and formaldehyde). (B) TE associated SVs were
primarily Deletions.

Figure 6. A) Simulated populations evolving with less than 10% of individuals reproducing through
outcrossing do not retain SVs. B) The mean number of structural variant mutations is lower in
populations with lower quantitative levels of reproduction through outcrossing. Here K=1000,
=0.001, =0.0006, =0.01, =0.001 and recombination rate =0.0001.

A Mean and Standard Error of Mutations by Strain and Mutagen

B Mean and Standard Error of Mutations in Exon by Strain and Mutagen

SV Density Distribution by Strain and Mutagen