1 Research Article

Screening analysis of
light-responsive phenotype of leaf bacteria

Hiroyuki Iguchi*, Michihiro Kanagasaki, Saya Ishii, Kazuma Taguchi

Department of Agriculture and Food Technology, Faculty of Bioenvironmental Sciences,

Kyoto University of Advanced Science.

* Corresponding author:

Hiroyuki Iguchi, PhD

Associate Professor

Department of Agriculture and Food Technology, Faculty of Bioenvironmental Sciences,

Kyoto University of Advanced Science, 1-1 Nanjo-Ohtani, Sogabe, Kameoka, Kyoto

621-8555, Japan.

Phone: +81 771 29 3452.   Fax: +81 771 29 3429.

E-mail: iguchi.hiroyuki@kuas.ac.jp

Keywords

light response, pigment, stress resistance, epiphyte, plant colonization

Abstract

The phyllosphere is a harsh environment in which microorganisms experience various stresses, many of which are attributed to sunlight. To understand the adaptation strategy of bacteria to the phyllospheric environment under sunlight, we examined light-responsive phenotypes by cultivating leaf bacterial isolates under light. Seven bacterial strains belonging to the genera *Massilia*, *Kocuria*, *Gordonia*, *Agromyces*, *Microbacterium*, and *Arthrobacter* were identified to enhance the production of yellow or red carotenoid pigments under light exposure. Light-responsive phenotypes were further analyzed in *Massilia* sp. ST22. This bacterium was confirmed to have the ability to colonize as well as produce pigments on plants. Analysis of several forms of stress resistance revealed that light positively affected UV resistance and negatively affected resistance to osmotic pressure and oxidative stress. These results uncover the light-responsive phenotypes of the leaf bacteria and indicate that light regulates several bacterial abilities necessary for plant colonization.

Introduction

In natural environments, microorganisms encounter various stresses, and their successful growth or survival requires adaptation to these stresses. In particular, the aerial part of plants (phyllosphere) is considered a hostile environment for microbial colonization (1, 2). The leaf surface conditions are intensely affected by climate changes (i.e., sun, rainfall, and wind), which can alter temperature, humidity, osmotic pressure, and light (UV) intensity. Furthermore, plants release antimicrobial compounds and limited nutrients. Despite the presence of these severe stresses, many microorganisms including bacteria, and fungi colonize the phyllosphere, and some play important roles in plant growth, plant health, and the carbon and nitrogen cycle (1-5). Thus, to better understand microbial ecology or implement phyllospheric microorganisms in plant cultivation, their ability to effectively colonize plant leaves has been studied (6-8). *Methylobacterium* species can utilize methanol, a niche carbon compound released from leaves, that is believed to allow this bacterial species to become dominant on leaves (9). Other analyses showed that leaf colonization requires flagella for localization and the production of trehalose, catalase, and pigment for protection from stresses (10-13).

A major influencer of the leaf surface environment is sunlight, which affects temperature, light energy, humidity, and plant metabolism. During daytime, the intensity of sunlight is altered irregularly by shading and regularly in a circadian manner. The leaf surface area is directly irradiated by sunlight, while some light energy and wavelengths are absorbed by the surface layer in soil and water environments (14, 15). The high temperature, desiccation, and DNA injury due to sunlight are harmful to microbial cells.

If microorganisms perceive changes in light intensity as a signal, they may effectively combat cognate stresses before the stresses reach a damaging level. Some bacteria employ light receptor proteins, which include LOV, BLUF, PYP, and cryptochrome for blue wavelength, and phytochrome for red wavelengths (16). Light and its cognate receptor were reported to regulate

the functions necessary for plant colonization, i.e., motility, attachment, and exopolysaccharide

production (17-20). However, these studies on light response used a limited panel of bacteria

such as plant pathogens. The bacterial population on leaves varies widely, mainly consisting of

Proteobacteria, *Bacteroidetes*, *Firmicutes*, and *Actinobacteria* (2, 21). Thus, knowledge of the

light response of the entire leaf bacterial population is lacking, and whether leaf bacteria use

light signals to facilitate growth on plants remains unknown.

In this study, we collected leaf bacterial isolates from natural environments and analyzed the

phenotypes under light culturing conditions. We identified light-induced pigment production in

leaf bacteria belonging to six genera. We further revealed that several forms of stress resistance

of one bacterium, *Massilia*, altered under the light condition. The findings of the current study

provide new information on the strategies used by leaf bacteria to grow under sunlight.

Materials and Methods

Sampling of plant leaves. Plant leaves were sampled in Kyoto, Hyogo and Osaka

prefectures in Japan. The sampling period occurred from spring to summer in 2015, 2016, and

2017.

Screening of light-responsive bacteria. Each leaf sample and 1 mL of sterilized water

were placed in a polypropylene tube. The tube was shaken for 1 hour, and then a portion of the

suspension was spread on an agar plate containing 0.5% (w/v) cycloheximide. We employed

two media, commonly used for plant bacterial cultivation (22, 23): R2A broth (Nihon

Pharmaceutical, Tokyo, Japan) was used in 2015 and 2016, and 2-fold-diluted tryptic soy broth

(TSB; Becton, Dickinson and Company, NJ, USA) was used in 2017. The dilution of TSB was

performed for the modulation of the bacterial growth rate. Colonies were collected from the

agar plates to make a master plate, which was incubated at 30ºC. The collected colonies were

streaked on a fresh plate containing the same agar media (1.5% agar) and cultivated at 30ºC

under both dark and light (PPFD, 3 µmol m^{-2} s^{-1}) conditions, the latter of which was generated

with a fluorescent light (FL20SW lamp; Toshiba, Tokyo, Japan). After 2 days of cultivation, the

two plates (dark and light conditions) were visually compared.

16S rRNA gene sequencing and accession numbers. The 16S rRNA gene was amplified

by PCR with genomic DNA, the 27f-1492r primer set (24), and Blend Taq (Toyobo, Osaka,

Japan), and then sequenced. These sequences were submitted to GenBank and assigned the

following accession numbers: LC416392–LC416398.

Spectral analysis. The agar plates on which the bacteria were streaked were cultivated in

the dark and under fluorescent light (PPFD, 3 µmol m^{-2} s^{-1}). To extract pigments, 0.5 mg (wet

weight) of the colonies collected from the plates was mixed with 1 mL of methanol or acetone

(for strain S136). The suspensions were incubated for 1 hour at room temperature or 50ºC (for

strain S138), and then the supernatant generated by centrifugation was analyzed by a

spectrophotometer (UV-2550; Shimadzu, Kyoto, Japan).

HPLC analysis. Strain ST22 was grown on *Arabidopsis* plants by seed inoculation

describes below, or on R2A agars. Cells of strain ST22 were harvested from the R2A liquid

culture and washed and resuspended with water. *Arabidopsis thaliana* Col-0 seeds were

sterilized with 70% ethanol. The seeds were incubated in 1 mL cell suspension (each

OD_{600}=0.1) with shaking on a rotator for 2 hours. The seeds were sown on Murashige and

Skoog medium including vitamins (Duchefa, Haarlem, Netherlands) and supplemented with

0.25% sucrose and 0.8% agar. The plants enclosed in petri dishes were gnotobiotically

incubated in a plant growth chamber (Nippon Medical & Chemical Instruments, Osaka, Japan)

at 25°C (14 hours light/10 hours darkness) for 3 weeks. The light condition was prepared with

two fluorescent lights (FHF32EX-N-HX-S lamp, NEC, Tokyo, Japan; FHF32N-EDL lamp,

Toshiba), and the light intensity (PPFD) at the petri dishes was 110 μmol m^{-2} s^{-1}.

The bacterial cells on the plant were collected from the aerial parts of six plants by

suspension with water followed by centrifugation. The collected bacterial cells from plants or

agars were mixed with methanol to extract the pigments, and the supernatant was subjected to

HPLC analysis.

HPLC analysis was performed on a UFLC system LC-20 series (Shimadzu). Each sample

was separated on an ODS column (column size 2 × 150 mm, Tosoh, Tokyo, Japan), and the

column flow was maintained at 0.2 mL/min with a linear gradient of 20 to 100% acetonitrile for

15 minutes. The UV/Vis detector was set at 450 nm.

ROS (reactive oxygen species) assay. ROS production was evaluated by a

dihydroethidium-based assay (25). Strain ST22 cells, grown on R2A agar in the dark or under

fluorescent light (PPFD, 3 or 10 μmol m^{-2} s^{-1}) for 2 days, were suspended in Dulbecco's

phosphate-buffered saline (D-PBS; Nacalai Tesque, Kyoto, Japan). The supernatant, generated

by undisturbed settling for a few minutes, was collected and adjusted to OD_{600} 0.5 with D-PBS.

One hundred microliters of the cell suspension was mixed with 50 μL of 15 μM

dihydroethidium (final concentration 5 μM) and incubated at 30°C for 30 minutes. The cells

were collected by centrifugation and suspended in D-PBS. The fluorescence was measured

using a fluorescence microplate reader (Gemini EM; Molecular Devices, CA, USA) with an

excitation wavelength of 485 nm and emission wavelength of 590 nm.

Liquid culture growth.   Strain ST22 cells were grown on R2A medium at 30°C and 160

rpm in the dark or under fluorescent light (PPFD, 6 μmol m^{-2} s^{-1}).

Stress resistance assay.   Strain ST22 cells, grown on R2A agar in the dark or under

fluorescent light (PPFD, 3 μmol m^{-2} s^{-1}) for 2 days, were suspended in water. The supernatant,

generated by undisturbed settling for a few minutes, was diluted to OD_{600}=1 and used for the

following analysis. Results of the analysis were expressed as viability, which was calculated by

comparing the CFU of stress-treated cells with that of untreated cells. Colony formation for

CFU evaluation was performed in the dark. (i) UV resistance. The cell suspension was further

diluted and spread on eight plates of R2A agar. Five of eight plates were exposed to UV-C light

(GL10 lamp, NEC; peak spectrum, 254 nm; full width at half maximum, 7 nm) for 10 seconds,

and the three remaining plates were untreated. (ii) Heat resistance. The cell suspension was

divided into four tubes (each 500 μL, OD_{600}=0.1), incubated at 50°C for 15 minutes, and then

immediately cooled at 4°C. The appropriately diluted suspensions were spread on R2A agar.

(iii) NaCl resistance. The cell suspension was divided into four tubes containing 2 M NaCl

(final OD_{600}=0.1), incubated for 30 minutes, and then immediately diluted with water. (iv) H_2O_2

resistance. The cell suspension was divided into four tubes containing 0.05 % hydrogen

peroxide (each 500 μL, final OD_{600}=0.1), incubated for 15 minutes, and then immediately mixed

with 5 μL of 0.2 M sodium thiosulfate for neutralization.

Results

Screening of light-responsive bacteria from plant leaves

We investigated the light-response behavior of the bacterial isolates from plant leaves. We used R2A agar, an oligotrophic medium, to cultivate bacteria from leaf samples. We isolated 382 bacterial strains, and determined whether they have light-responsive phenotypes. Visual comparisons between agar cultures grown under dark and fluorescent light conditions revealed two strains (Fig. 1 and Table 1): strain Kn16 (identified in the first year) produced yellow pigments under the light condition, and strain ST22 (identified in the second year) exhibited enhanced yellow pigment production under the light condition. Other light-responsive phenotypes such as motility, colony morphology, and growth rate were not detected under the tested condition. Sequencing of the 16S rRNA gene (ca. 1,500 bp) revealed that strains Kn16 and ST22 both belonged to the genus *Massilia*, with a sequence difference of 3.7%. These two strains were isolated from different plant species (Table 1). Then, isolation and cultivation of leaf bacteria were carried out using TSB agar, which is a nutritious medium. Among the 224 isolated strains, we identified five strains with enhanced pigment production under the light condition (Fig. 1 and Table 1). Strains S71, S138, S821, and S1029 produced yellow pigments, while strain 136 produced orange pigment. The 16S rRNA gene sequences revealed that all five strains belonged to the phylum Actinobacteria. Based on the closest relatives, the strains were classified as *Kocuria* sp. S71, *Gordonia* sp. S136, *Agromyces* sp. S138, *Microbacterium* sp. S821, and *Arthrobacter* sp. S1029.

The pigment production induced under the light condition was analyzed with a spectrophotometer. The pigments were extracted from cells grown on agar under dark and light conditions. The spectra of the pigment extracts of seven strains displayed a peak representative of carotenoid, with the maximum absorption wavelength at 450 nm (Fig. 2). The pigment production was induced 2.6- to 17-fold by light exposure, based on the absorbance value at 450

nm.

Plant colonization ability of *Massilia* sp. ST22

We determined if the pigments are produced during plant colonization. We used the isolate

Massilia sp. ST22, as knowledge of the light-response and plant colonization of this genus is

lacking. The strain ST22 was inoculated on sterilized *Arabidopsis* seeds, which were cultivated

gnotobiotically under a dark/light cycle. The aqueous suspension containing the phyllospheric

portions of the plants formed colonies on R2A agar, demonstrating that strain ST22 have the

ability to grow on plants. Then, the pigment production on plants was analyzed by HPLC to

detect the relatively small number of cells colonizing the plants. The pigments in bacterial cells

collected from the plant phyllosphere were extracted with methanol. The extract of the strain

showed a peak at 13.5 min (Fig. 3a), which corresponds to the pigment produced on agar (Fig.

3b), and the eluted fraction displayed a yellow color. These results showed that strain ST22

produces pigments during colonization on *Arabidopsis* plants.

Analysis of light-responsive, stress-resistant phenotypes of *Massilia* sp. ST22

Next, we further analyzed light-responsive phenotypes with respect to stress resistance to

determine whether light regulates the plant colonization ability. First, the light conditions were

set to minimize cellular damage. The ROS level was evaluated based on the

dihydroethidium-based fluorescence assay for the cells grown on agar. The ROS level was

comparable between light and dark conditions at a light intensity of 3 μmol m^{-2} s^{-1}, whereas

ROS were accumulated at a light intensity of 10 μmol m^{-2} s^{-1} (Fig 4a). This light intensity (3

μmol m^{-2} s^{-1}) also induced pigment production as shown in Fig. 1 and was thus employed in the

following tests with agar cultures.

The resistance of *Massilia* sp. ST22 wild-type to several stresses was compared under dark

and light conditions. It was found that UV resistance was elevated under the light condition,

while the resistance to NaCl and hydrogen peroxide was decreased under the light condition

(Fig. 4b). Light did not affect the heat resistance. Growth was compared in liquid culture using

the flask shaker equipped with a fluorescent light (6 μmol m^{-2} s^{-1}). A similar growth curve was

generated for the dark and light conditions (Fig. 4c). These results indicated that light positively

regulates UV resistance, and negatively regulates on resistance to osmotic pressure (NaCl) and

oxidative stress (H_2O_2) in *Massilia* sp. ST22.

Discussion

The phyllosphere is known to be a stressful environment, and microbial colonizers are expected to be able to effectively respond to the stresses for growth and survival. Our focus in this study was to determine whether leaf bacteria use sunlight as a signal to address the stresses present in the phyllosphere. To elucidate the light-responsive phenotypes, we cultivated a total of 606 bacterial isolates from leaves under light and dark conditions and identified seven strains that produce more pigments under light conditions (Fig. 1 and Table 1). These strains belong to the two major phyla in the phyllospheric bacterial community (Proteobacteria and Actinobacteria) (2, 21). This is the first report on the light-induced pigmentation of the two genera *Massilia* and *Kocuria*. Pigment production under light conditions has been reported in four other genera (*Gordonia*, *Agromyces*, *Microbacterium*, and *Arthrobacter*) (26-30). These six genera are widely distributed in environments such as soil, water, animals, and plants (phyllosphere and rhizosphere) (31-36). Some species of these genera were reported to not only colonize plants but also have beneficial interaction with plants (i.e., plant growth promotion) (35-40). The air pollutant biodegradation capacity of bacteria on leaf surfaces was also reported (41). Thus, these bacteria are likely to play an important ecological role in plant-associated environments.

Some isolated strains produced pigments at a low level in the dark, whereas strains S71, S138, and S1029 showed almost no pigment production in the dark, indicating strict repression of pigment synthesis (Fig. 1 and 2). Once pigments are produced intracellularly, they are usually maintained in the cells during their lifetime. Accordingly, light induced-pigmentation is likely to occur when the bacteria first encounter light. The advantages of the light-induced pigmentation system in bacteria remain unknown. One explanation is that bacteria can conserve the biological energy for pigment production in dark environments such as soils where pigment function is possibly not required. The plant pathogen *P. syringae* was reported to reversibly change its

behavior for plant infection via the blue light receptor between the soil (dark) and phyllospheric (light) environments as well as day and night. Under blue light conditions, which partly represent the phyllospheric environment, motility and virulence are repressed, while attachment to plant leaves is promoted (17). At nightfall, these traits can be reversed (i.e., activation of motility and virulence) by altering gene expression levels. The defense response of plants is also regulated in a circadian manner, with higher susceptibility to infection at night (42).

The induced pigments were considered carotenoids (Fig. 2), and carotenoid pigments are known to function in protection from oxidative stresses, especially ROS generated by UV radiation (43, 44). Previous studies demonstrated that pigment-deficient bacterial mutants became more sensitive to oxidative stresses (12, 45, 46) with reduced ability to colonize plants (12, 44). The plant colonization ability encompasses many traits, including acquisition and metabolism of nutrients, response to environmental stresses (temperature, light, humidity, osmotic pressure, ROS, and antimicrobial compounds), motility, and adhesion (2). Several light-responsive phenotypes possibly relative to plant colonization were further revealed in *Massilia* sp. ST22. Light induced UV resistance (Fig. 4b), representing the close connection between light and resistance to damage caused by light. Thus, this bacterium is suspected to generate a defense system against sunlight damage by enhancing its UV resistance during daytime. Light-induced UV resistance has been observed in some microorganisms (47, 48) and probably occurs via controlling the functions of DNA repair, protein folding, and ROS scavenging at the gene expression level (43, 49-51). In the context of UV resistance, the resistance to oxidative stress, which is caused by light, was expected to be induced, but the resistance to H_2O_2 was decreased (Fig. 4b). The resistance to osmotic pressure (NaCl), which is unlikely to relate to light, was also found to be decreased. Thus, not all light-induced phenotypes are useful for plant colonization, and we suspect that light triggers multiple regulatory systems (i.e., positive and negative regulation).

In summary, cultivation of bacteria under light conditions has clarified the light-responsive

phenotypes of seven phyllospheric bacteria. Our test conditions using agar culture clearly

revealed the pigmentation phenotype. The extensive analysis of different bacterial abilities

revealed that several stress-resistance phenotypes of *Massilia* sp. ST22 affected by light. This

finding suggests the possibility that light-responsive phenotypes are widely distributed in

phyllospheric bacteria, including those not assessed in this study. The light-responsive

phenotypes such as stress resistance also should play a role in growth and survival in the

phyllosphere. It is still unclear whether changes in stress resistance under the light condition

occurred via sensing of light signals with photoreceptors or were triggered by light-induced cell

injury. Further work is required to determine the light-responsive regulatory mechanism,

including photoreceptor proteins, transcriptional factors and regulated genes. In addition,

techniques controlling bacterial behavior using specific wavelengths and photoreceptors are

promising research topics for future investigation (52-54). The combined knowledge of ecology

and molecular biology will facilitate the application of light to control the abilities of

phyllospheric bacteria, including growth property, pathogenicity, and bioactive compound

production.

Funding

This study was supported in part by a Grant-in-Aid for Young Scientists (18K14381, to H.I.), and a Grant-in-Aid for Young Scientists (B) (15K18671, to H.I.) from the Japan Society for the Promotion of Science.

References

References

1. **Lindow SE, Brandl MT.** 2003. Microbiology of the phyllosphere. Appl Environ Microbiol
69:1875-1883.

2. **Vorholt JA.** 2012. Microbial life in the phyllosphere. Nat Rev Microbiol **10:**828-840.

3. **Sandhu A, Halverson LJ, Beattie GA.** 2007. Bacterial degradation of airborne phenol in the
phyllosphere. Environ Microbiol **9:**383-392.

4. **Galbally IE, Kirstine W.** 2002. The production of methanol by flowering plants and the global
cycle of methanol. Journal of Atmospheric Chemistry **43:**195-229.

5. **Penuelas J, Terradas J.** 2014. The foliar microbiome. Trends Plant Sci **19:**278-280.

6. **Tani A, Takai Y, Suzukawa I, Akita M, Murase H, Kimbara K.** 2012. Practical application of
methanol-mediated mutualistic symbiosis between *Methylobacterium* species and a roof
greening moss, *Racomitrium japonicum*. PLoS ONE **7:**e33800.

7. **Arthurs S, Dara SK.** 2018. Microbial biopesticides for invertebrate pests and their markets in
the United States. J Invertebr Pathol doi:10.1016/j.jip.2018.01.008.

8. **J P Blakeman a, Fokkema NJ.** 1982. Potential for biological control of plant diseases on the
phylloplane. Annu Rev Phytopathol **20:**167-190.

9. **Sy A, Timmers AC, Knief C, Vorholt JA.** 2005. Methylotrophic metabolism is advantageous
for *Methylobacterium extorquens* during colonization of *Medicago truncatula* under competitive
conditions. Appl Environ Microbiol **71:**7245-7252.

10. **Freeman BC, Chen C, Beattie GA.** 2010. Identification of the trehalose biosynthetic loci of
Pseudomonas syringae and their contribution to fitness in the phyllosphere. Environ Microbiol
12:1486-1497.

11. **Guo M, Block A, Bryan CD, Becker DF, Alfano JR.** 2012. *Pseudomonas syringae* catalases
are collectively required for plant pathogenesis. J Bacteriol **194:**5054-5064.

12. **Mohammadi M, Burbank L, Roper MC.** 2012. Biological role of pigment production for the

309   bacterial phytopathogen *Pantoea stewartii* subsp. stewartii. Appl Environ Microbiol

310   **78:**6859-6865.

311  13. **Rossez Y, Holmes A, Wolfson EB, Gally DL, Mahajan A, Pedersen HL, Willats WG, Toth**

312   **IK, Holden NJ.** 2014. Flagella interact with ionic plant lipids to mediate adherence of

313   pathogenic *Escherichia coli* to fresh produce plants. Environ Microbiol **16:**2181-2195.

314  14. **Levine JS, MacNichol EF.** 1982. Color vision in fishes. Sci Am **246:**140-149.

315  15. **Mandoli DF, Ford GA, Waldron LJ, Nemson JA, Briggs WR.** 1990. Some spectral properties

316   of several soil types: implications for photomorphogenesis. Plant Cell Environ **13:**287-294.

317  16. **Gomelsky M, Hoff WD.** 2011. Light helps bacteria make important lifestyle decisions. Trends

318   Microbiol **19:**441-448.

319  17. **Rio-Alvarez I, Rodriguez-Herva JJ, Martinez PM, Gonzalez-Melendi P, Garcia-Casado G,**

320   **Rodriguez-Palenzuela P, Lopez-Solanilla E.** 2014. Light regulates motility, attachment and

321   virulence in the plant pathogen *Pseudomonas syringae* pv tomato DC3000. Environ Microbiol

322   **16:**2072-2085.

323  18. **Bonomi HR, Posadas DM, Paris G, Carrica Mdel C, Frederickson M, Pietrasanta LI,**

324   **Bogomolni RA, Zorreguieta A, Goldbaum FA.** 2012. Light regulates attachment,

325   exopolysaccharide production, and nodulation in *Rhizobium leguminosarum* through a

326   LOV-histidine kinase photoreceptor. Proc Natl Acad Sci U S A **109:**12135-12140.

327  19. **Kraiselburd I, Alet AI, Tondo ML, Petrocelli S, Daurelio LD, Monzon J, Ruiz OA, Losi A,**

328   **Orellano EG.** 2012. A LOV protein modulates the physiological attributes of *Xanthomonas*

329   *axonopodis* pv. citri relevant for host plant colonization. PLoS ONE **7:**e38226.

330  20. **Moriconi V, Sellaro R, Ayub N, Soto G, Rugnone M, Shah R, Pathak GP, Gartner W, Casal**

331   **JJ.** 2013. LOV-domain photoreceptor, encoded in a genomic island, attenuates the virulence of

332   *Pseudomonas syringae* in light-exposed *Arabidopsis* leaves. Plant J **76:**322-331.

333  21. **Bulgarelli D, Schlaeppi K, Spaepen S, Ver Loren van Themaat E, Schulze-Lefert P.** 2013.

Structure and functions of the bacterial microbiota of plants. Annu Rev Plant Biol **64:**807-838.

22.   **Bai Y, Muller DB, Srinivas G, Garrido-Oter R, Potthoff E, Rott M, Dombrowski N, Munch**

PC, Spaepen S, Remus-Emsermann M, Huttel B, McHardy AC, Vorholt JA, Schulze-Lefert

P. 2015. Functional overlap of the *Arabidopsis* leaf and root microbiota. Nature **528:**364-369.

23.   **Lebeis SL.** 2017. Plant microbiome identification and characterization. Curr Protc in Plant Biol

2:135-146.

24.   **Weisburg WG, Barns SM, Pelletier DA, Lane DJ.** 1991. 16S ribosomal DNA amplification for

phylogenetic study. J Bacteriol **173:**697-703.

25.   **Owusu-Ansah E, Yavari A, Banerjee U.** 2008. A protocol for *in vivo* detection of reactive

oxygen species. Protocol Exchange doi:10.1038/nprot.2008.23.

26.   **Jeon BY, Kim BY, Jung IL, Park DH.** 2012. Metabolic roles of carotenoid produced by

non-photosynthetic bacterium *Gordonia alkanivorans* SKF120101. J Microbiol Biotechnol

22:1471-1477.

27.   **Meddeb-Mouelhi F, Moisan JK, Bergeron J, Daoust B, Beauregard M.** 2016. Structural

characterization of a novel antioxidant pigment produced by a photochromogenic

Microbacterium oxydans strain. Appl Biochem Biotechnol **180:**1286-1300.

28.   **Trutko SM, Dorofeeva LV, Evtushenko LI, Ostrovskii DN, Hintz M, Wiesner J, Jomaa H,**

Baskunov BP, Akimenko VK. 2005. Isoprenoid pigments in representatives of the family

Microbacteriaceae. Mikrobiologiia **74:**335-341.

29.   **Linos A, Steinbuchel A, Sproer C, Kroppenstedt RM.** 1999. *Gordonia polyisoprenivorans* sp.

nov., a rubber-degrading actinomycete isolated from an automobile tyre. Int J Syst Bacteriol

49:1785-1791.

30.   **Sutthiwong N, Dufosse L.** 2014. Production of carotenoids by *Arthrobacter arilaitensis* strains

isolated from smear-ripened cheeses. FEMS Microbiol Lett **360:**174-181.

31.   **Baldani JI, Rouws L, Cruz LM, Olivares FL, Schmid M, Hartmann A.** 2014. The family

Oxalobacteraceae, p 919-974. *In* Rosenberg E, DeLong EF, Lory S, Stackebrandt E, Thompson

F (ed), The Prokaryotes: Alphaproteobacteria and Betaproteobacteria. Springer, Berlin,

Heidelberg.

32.   **Stackebrandt E, Schumann P.** 2015. *Kocuria*. Bergey's Manual of Systematics of Archaea and

Bacteria, Online doi:10.1002/9781118960608.gbm00120.

33.   **Busse HJ, Wieser M, Buczolits S.** 2015. *Arthrobacter*. Bergey's Manual of Systematics of

Archaea and Bacteria, Online doi: 10.1016/j.jip.2018.01.008.

34.   **Goodfellow M, Kumar Y, Maldonado LA.** 2015. *Gordonia*. Bergey's Manual of Systematics of

Archaea and Bacteria, Online doi:10.1002/9781118960608.gbm00030.

35.   **Evtushenko LI, Ariskina EV, Prisyazhnaya NV, Starodumova IP.** 2017. *Agromyces*. Bergey's

Manual of Systematics of Archaea and Bacteria, Online

doi:10.1002/9781118960608.gbm00093.pub2.

36.   **Suzuki K-I, Hamada M.** 2015. *Microbacterium*. Bergey's Manual of Systematics of Archaea

and Bacteria, Online doi:10.1002/9781118960608.gbm00104.

37.   **Haidar B, Ferdous M, Fatema B, Ferdous AS, Islam MR, Khan H.** 2018. Population

diversity of bacterial endophytes from jute (*Corchorus olitorius*) and evaluation of their potential

role as bioinoculants. Microbiol Res **208:**43-53.

38.   **Kayasth M, Kumar V, Gera R.** 2014. *Gordonia* sp.: a salt tolerant bacterial inoculant for

growth promotion of pearl millet under saline soil conditions. 3 Biotech **4:**553-557.

39.   **Xu X, Xu M, Zhao Q, Xia Y, Chen C, Shen Z.** 2018. Complete genome sequence of

Cd(II)-resistant *Arthrobacter* sp. PGP41, a plant growth-promoting bacterium with potential in

microbe-assisted phytoremediation. Curr Microbiol **75:**1231-1239.

40.   **Chimwamurombe PM, Grönemeyer JL, Reinhold-Hurek B.** 2016. Isolation and

characterization of culturable seed-associated bacterial endophytes from gnotobiotically grown

Marama bean seedlings. FEMS Microbiol Ecol **92:**fiw083.

41.   **Wei X, Lyu S, Yu Y, Wang Z, Liu H, Pan D, Chen J.** 2017. Phylloremediation of air

385   pollutants: exploiting the potential of plant leaves and leaf-associated microbes. Frontiers in

386   Plant Science **8:**1318.

387   42.   **Bhardwaj V, Meier S, Petersen LN, Ingle RA, Roden LC.** 2011. Defence responses of

388   *Arabidopsis thaliana* to infection by *Pseudomonas syringae* are regulated by the circadian cock.

389   PLoS ONE **6:**e26968.

390   43.   **Ziegelhoffer EC, Donohue TJ.** 2009. Bacterial responses to photo-oxidative stress. Nat Rev

391   Microbiol **7:**856-863.

392   44.   **Jacobs JL, Carroll TL, Sundin GW.** 2005. The role of pigmentation, ultraviolet radiation

393   tolerance, and leaf colonization strategies in the epiphytic survival of phyllosphere bacteria.

394   Microb Ecol **49:**104-113.

395   45.   **Orlandi VT, Bolognese F, Chiodaroli L, Tolker-Nielsen T, Barbieri P.** 2015. Pigments

396   influence the tolerance of *Pseudomonas aeruginosa* PAO1 to photodynamically induced

397   oxidative stress. Microbiology **161:**2298-2309.

398   46.   **Liu GY, Essex A, Buchanan JT, Datta V, Hoffman HM, Bastian JF, Fierer J, Nizet V.** 2005.

399   *Staphylococcus aureus* golden pigment impairs neutrophil killing and promotes virulence

400   through its antioxidant activity. J Exp Med **202:**209-215.

401   47.   **Massier S, Bouffartigues E, Rincé A, Maillot O, Feuilloley MGJ, Orange N, Chevalier S.**

402   2013. Effects of a pulsed light-induced stress on *Enterococcus faecalis*. J Appl Microbiol

403   **114:**186-195.

404   48.   **Wood ER, Ghané F, Grogan DW.** 1997. Genetic responses of the thermophilic archaeon

405   *Sulfolobus acidocaldarius* to short-wavelength UV light. J Bacteriol **179:**5693-5698.

406   49.   **Goosen N, Moolenaar GF.** 2008. Repair of UV damage in bacteria. DNA Repair **7:**353-379.

407   50.   **Fröls S, White Malcolm F, Schleper C.** 2009. Reactions to UV damage in the model archaeon

408   *Sulfolobus solfataricus*. Biochem Soc Trans **37:**36-41.

51.   **Massier S, Rincé A, Maillot O, Feuilloley MGJ, Orange N, Chevalier S.** 2012. Adaptation of

Pseudomonas aeruginosa to a pulsed light-induced stress. J Appl Microbiol **112:**502-511.

52.   **Idnurm A, Crosson S.** 2009. The photobiology of microbial pathogenesis. PLoS Pathog

5:e1000470.

53.   **Trivedi S, Choudhary OP, Gharu J.** 2011. Different proposed applications of

bacteriorhodopsin. Recent patents on DNA & gene sequences **5:**35-40.

54.   **Ramírez MS, Müller GL, Pérez JF, Golic AE, Mussi MA.** 2015. More than just light: clinical

relevance of light perception in the nosocomial pathogen *Acinetobacter baumannii* and other

members of the genus *Acinetobacter*. Photochem Photobiol **91:**1291-1301.

Table 1. Bacterial isolates from leaves with light-induced pigmentation

Strain	Isolation origin	Closest relative
Kn16	*Salix koriyanagi*	*Massilia timonae* UR/MT95[T] (98%)
ST22	*Toxicodendron sylvestre*	*Massilia putida* 6NM-7[T] (98%)
S71	*Osmanthus fragrans var. aurantiacus*	*Kocuria palustris* TAGA27 [T] (100%)
S136	Not identified	*Gordonia hongkongensis* HKU50 [T] (98%)
S138	*Rhododendron pulchrum* cv. *Oomurasaki*	*Agromyces aureus* AR33 [T] (99%)
S821	*Cayratia japonica*	*Microbacterium ginsengisoli* Gsoil 259 [T] (99%)
S1029	*Conyza canadensis*	*Arthrobacter histidinolovorans* DSM 20115 [T] (99%)

420   Closest relative is with the type strains. The sequence identity is shown in parentheses.

Fig. 1. Colony of leaf bacterial isolates grown under dark or light conditions. Bacteria from leaf samples were grown on agar, and isolates were stored on a master plate. The stored bacteria were inoculated on two agar plates, and the colony phenotypes were compared between the dark and light culture conditions.

Fig. 2. Spectrum of cell extracts of leaf bacterial isolates grown under dark (dotted line) or light (solid line) conditions. Pigments were solvent-extracted from cells grown on agar. The value indicates the fold change of the absorbance at 450 nm, which represents the maximum absorption wavelength of carotenoid pigment, between the cells grown under dark and light conditions, respectively.

Fig. 3. HPLC analysis of pigment production in *Massilia* sp. ST22. Pigments were extracted from bacterial cells collected from the phyllosphere of *Arabidopsis* plants (a) and agar cultures (b). An arrow indicates the peak of the pigment.

Fig. 4. Analysis of light responsive functions in *Massilia* sp. ST22. (a) ROS generation in cells grown under dark and light conditions. Values indicate the relative fluorescent intensity based on the dihydroethidium assay. Data are provided as the means ± standard deviations (N=3). (b) Stress resistance of cells grown under dark and light conditions. The viability is expressed as the relative CFU ratio of stress-treated cells to untreated cells. Data are provided as the means ± standard deviations (N=4). D, dark condition; L, light condition at 3 μmol m^{-2} s^{-1} intensity; LL, light condition at 10 μmol m^{-2} s^{-1}. Statistical significance was assessed using Student's t-test. **, *P*<0.01. (c) Liquid culture growth under dark (circle) and light (triangle) conditions.