Social Context Restructures Behavioral Syntax in Mice

Marti Ritter[1,2]*, Hope Shipley[2,3], Serena Deiana[2], Bastian Hengerer[2], Carsten T. Wotjak[2],
Michael Brecht[1], Amarender R. Bogadhi[2]

[1]Department of Biology, Faculty of Life Sciences, Humboldt University, Berlin, Germany

[2]Neuroscience and Mental Health Discovery Research, Boehringer Ingelheim Pharma GmbH &
Co. KG, Biberach, Germany

[3]University of Manchester, Manchester, United Kingdom

* Correspondence:

Marti Ritter (Research Group of Michael Brecht)

Philippstraße 13, House 6
10115 Berlin
Germany
Tel: + 49-(0)30-20936770

E-mail: ritermaq@hu-berlin.de

Keywords: mouse, social behavior, social withdrawal, context, behavioral syntax,
behavioral decomposition, behavioral repertoire

Abstract

Investigation of social behavior in mice is increasingly relevant to understand associated brain circuits and thereby treating psychiatric symptoms involving social withdrawal or social anxiety. A data-driven description of behavior and its modulation by solitary and social context is lacking. In this study, we asked if and how the behavioral repertoire of mice differs in solitary and social ("dyadic") context through behavioral decomposition ("Syllables") using motion sequencing ("MoSeq"). Our results reveal a significant modulation in a minority (25%) of syllables, containing mostly stationary and undirected behaviors, by social context. However, the modulation is associated with spatial proximity to the conspecific rather than active contact. Interestingly, a network analysis of syllable transitions shows that modulated syllables play a regulatory role with significant changes in network influence, independent of the number of connected syllables. Furthermore, we found significant changes to syllable composition at the time of mice contact behavior and identified two families of syllable sequences that dominate mice approach or leave behaviors. However, we did not identify unique sequences of syllables that map onto specific mice social behaviors. Overall, our findings suggest that a subset of syllables underlie the changes in the behavioral composition of mice under social context and may mediate transitions within the behavioral repertoire.

1   Introduction

Social interactions are some of the most complex processes across all basic functions required for survival, leading to a high level of specialization of the involved structures and a high cost of dysfunction (Cacioppo et al. 2014; Porcelli et al. 2019). At the same time this high level of complexity leads to an increased vulnerability, e.g. to pathogens or disorders, an example being the occurrence of social withdrawal as a shared feature across major depressive disorder (MDD), Alzheimer's disease (AD), and schizophrenia (Porcelli et al. 2019). These conditions have a large impact on quality of life and carry a high cost to both affected individual (Chevance et al. 2020) and society, leading to an urgent need for the discovery of treatments (Greenberg et al. 2015; Jia et al. 2018). After previous approaches showed limited success, perspective on translational neuroscience has shifted towards a focus on psychopathology rooted in observable behavior and neurobiological measures (Research Domain Criteria, RDoC), suggesting that focusing on social impairments in complex disorders might lead to an earlier detection and better treatment outcomes (Garner 2014; Casey et al. 2013; Cuthbert 2018; Insel et al. 2010).

To facilitate early and effective identification of promising candidate treatments, pre-clinical research makes extensive use of rodent models of disorders found in humans (Bolivar, Walters, and Phoenix 2007). This approach can also be used in the search for treatments of social impairments (Luo et al. 2023). Mice possess a complex social behavior, influenced by age, kinship, and sex, among other factors. They utilize ultrasonic vocalizations, pheromones, and body contact as modalities, to cite some (Sangiamo, Warren, and Neunuebel 2020; Ricceri, Moles, and Crawley 2007). Recent developments in translational research led to refocusing from constrained tasks at shorter timescales, such as the three-chamber social approach task (Nadler et al. 2004) or tube test to determine the social hierarchy between mice (Fan et al. 2019), towards an open arena layout to enable the mice to show more complex behavioral structures and social interactions (Peleh, Ike, et al. 2019; Shemesh et al. 2013; Forkosh et al. 2019). But the complexity of social behavior also required agreed-upon definitions for specific interaction motifs to achieve quantifiable and reproducible descriptions.

At its inception, traditional ethological research was limited to qualitative, higher-level descriptions to characterize complex behaviors (Anderson and Perona 2014). This required a reduced set of dimensions to make behaviors observable (Gomez-Marin et al. 2014), such as focusing on vocalizations or Infra-red beam breaks. Even though some of the earliest descriptions of mouse behavior contained "ethograms" matching complex and specific behaviors (Abeelen 1964), it was not possible at that time to capture the full complexity of behavior into an ethogram in a data-driven, quantifiable manner. As this limited the possibilities of the neuroscience community to investigate unrestrained, naturalistic behavior, an urgent goal was the development of appropriate tools (Datta et al. 2019). The advent of convolutional neural networks and marker-less pose tracking tools, such as DeepLabCut (Lauer et al. 2022) or SLEAP (Pereira et al. 2022), allowed for the automatic extraction of key body points capturing much of the pose information available in video recordings collected by one or more cameras (Ebbesen and Froemke 2022), for single (Pereira et al. 2019; Mathis et al. 2018) or multiple animals (Romero-Ferrero et al. 2019; Bewley et al. 2016). In parallel, other approaches to behavior tracking and decomposition arose (Wiltschko et al. 2015), which utilized the recent advances in machine learning and computation (Dennis et al. 2021). The behavioral decomposition approach, which allows identification of recurring short segments of behavior ('syllables') that form the

basis of complex behavior, was effective in characterizing behavioral fingerprints of drugs in
disorders, such as epilepsy (Gschwind et al. 2023), using both three dimensional recordings of
single mice (Wiltschko et al. 2020) and utilizing body point representations of single or multiple
mice (Hsu and Yttri 2021; Luxem et al. 2022).

A multitude of tools that focus on automated tracking and annotation of social interactions in
mice can be found in recent literature (Peleh, Bai, et al. 2019; Chaumont et al. 2012; Giancardo
et al. 2013), including open-source recording setups (Chaumont et al. 2018). Despite advances
towards an integration of ethological and comparative psychological approaches (Bordes et al.
2023) most of these approaches are based on parametric definitions of specific behaviors, as in
supervised tools (Goodwin et al. 2024), and do not leverage potential advantages of unsupervised
behavior classification or standardized benchmarking (Choi and Kumar 2024). For example, use
of motion sequencing in pain quantification (Jhumka and Abdus-Saboor 2022) and evaluation of
behavioral effects of analgesics (Bohic et al. 2023) revealed unique insights into the difference
between baseline and analgesia induced behaviors, as well as common global markers of
ongoing pain, such as specific pause and grooming modules.

  99   To investigate if an unsupervised behavioral decomposition approach yields insights into the
structure of social behavior in mice, we applied Keypoint-MoSeq to videos of mice in dyadic and
solitary contexts. By comparing syllable frequency and structure in dyadic and solitary contexts,
we show a significant modulation in a subset of stationary, undirected syllables that are not fully
dependent on contact with a conspecific. Instead, these modulated syllables play a key role in
transitions between syllables as the behavior becomes more diverse in a dyadic context.
Furthermore, social behaviors (e.g. social approach or leave) defined through parametric
thresholds can be represented in syllable space. However, we did not find syllables or syllable
sequences that distinctly map onto social behaviors. Our results suggest that syllables or syllable
sequences are sensitive markers for the changes in behavior in the presence of a conspecific but
may not correspond to specific "social" behaviors occurring in this context.

2   Materials and Methods

2.1   Animals

Twenty female C57BL/6J mice (~6 weeks old at time of first experiment) were obtained from Charles River Germany. They were housed in groups of two to three, with the other cage-mates not used in this study and kept in a light- and humidity-controlled facility. All animals were orally injected with a vehicle treatment intraperitoneally 20 minutes before their recordings started to account for adverse effects caused by the oral gavage procedure. All procedures followed the regulations for animal experimentation enforced by the local district administration's animal welfare commissioner of the state of Baden-Württemberg.

2.2   Open Field Arena Setup

The experiments were run in a 453mm x 453mm x 400mm (Width x Depth x Height) square arena made from clear plexiglass. The arena was equipped with dimmable LED strips and an analog camera (768px x 576px, 25Hz). All sides except for the front were enclosed by a wooden frame, with the front being covered with a red plexiglass door, and the floor being removable to facilitate cleaning. During the tests a stable illumination of around 220lux was maintained.

2.3   Experimental procedure

Half of the mice (n=10) were kept on a regular 12h light-dark cycle, with the other 10 animals being kept in a reversed 12h light-dark cycle. This applied to both the solitary as well as the dyadic experiments, with the dyadic experiments occurring between mice with the same light cycle. All experiments began between 8 and 10 am. The timeline of experiments and other procedures is shown below.

2.3.1 Solitary open field test

Mice were orally injected with Hydroxypropyl-β-cyclodextrin 20% McIlVaine and kept in their home cages for 20 minutes. The oral gavage was performed to consider the impact of treatment stress for future studies on pharmacological treatment. After 20 minutes had passed, mice were transferred into the recording setups and video-recorded for one hour. The recordings were started automatically through Stoelting Any-Maze software. Of the one hour of solitary recordings, only the first 20 minutes were used for behavioral analysis in this study, to maintain data parity to the dyadic context. The timeline for a single solitary recording is shown below.

2.3.2 Dyadic social interation test

Five weeks after the solitary test recordings, mice were tested in a dyadic social interaction test. Two mice from separate cages were orally injected Hydroxypropyl-β-cyclodextrin 20% McIIVaine at the same time and housed in their home cage for 20 minutes before starting the dyadic social interaction test. Mice were allowed to freely interact and explore the open field arena for 20 minutes. To allow for discrimination between the two mice in each recording, one mouse was shaved eight to nine days before the recording date at a constant location at the upper back behind the neck. The timeline for a single dyadic recording is shown below.

2.4   Data collection and analysis

2.4.1 Stoelting Any-Maze

Stoelting Any-Maze version 7.4 was used to record the videos used in this analysis and to record metadata. No further tracking parameters were extracted from Anymaze for this study. We utilized the built-in scoring utility to collect experimenter-scored active and passive contact epochs in the video data.

2.4.2 SLEAP

SLEAP (Pereira et al. 2022) version 1.3.3 was used to track 12 key points on the body of each mouse in the solitary and dyadic video recordings. 4 of the key points were located on the head of the mouse, 4 along the spine, another 2 were located one each at the hips, and 2 further points were labelled along the tail. For the model, we selected a bottom-up design with 800 frames taken from dyadic context and 600 frames taken from solitary context for the training set.

2.4.3 Keypoint-MoSeq

Keypoint-MoSeq (Weinreb et al. 2024) version 0.4.1 was used to extract the behavioral syllables from the keypoints extracted in the previous step. We used 10 of the 12 body points tracked in SLEAP (excluding the tail) to fit the model. The model parameter Kappa was selected iteratively such that the resulting syllable length distribution had its median at 10 frames, representing roughly 400ms of behavior. The default threshold proposed in the analysis tools provided by the Keypoint-MoSeq Python package was set to 0.5% of total onset proportion and we followed that threshold here.

Using the tools provided in the Keypoint-MoSeq package, we named all 32 syllables with a bout proportion above 0.5% of the global distribution. Names and descriptions were chosen before further analysis according to the behaviors shown in the majority of the 24 extracted example videos for each syllable. For a full list and descriptions, see Table 1.

Table 1. *Table of syllables.* Labels and descriptions were determined by using the tools provided with the Keypoint-MoSeq package to create a 24-video overview of the occurrences of each syllable. Behaviors were described according to the behavior shown in most of the videos (see fractions in description column). Labelling was performed prior to the analysis, and minor edits were done for readability.

Syllable	Label	Short description
0	dart_forward	20/24 videos show relatively quick movement from stationary position forward, generally, if possible straight ahead no turns
1	turn_left	17/24 show left turn, 8/17 related to down from rearing… due to 2D? Usually stationary
2	turn_right_move	16/24 show right turn, 7/16 related to (down from, sometimes aborted) rearing, same as 1, maybe due to 2D? Compared to 1 this one has after the turn some movement
3	turn_right_stationary	17/24 show right turn, this time mainly stationary, 11/17 related to moving up into rearing.
4	unclear_contraction	8/24 show down from rearing, generally seems to have some rightward bias, but not specific
5	movement_forward	22/24 show general, constant movement forward, usually average velocity

6	sniffing	23/24 show sniffing, 20/23 related to the corners and edges of the box
7	rearing	19/24 show movement related to rearing, or "waving" during rearing, mounting… seems to detect more "last" phase of rearing shortly before or during down
8	turn_right_move2	19/24 usually first light right turn then moving forwards, generally long
9	turn_left_short	21/24 short left turn, sometimes movement afterwards
10	sniffing2	19/24 sniffing again
11	turn_right_stationary2	20/24 right turn on spot
12	head_raise	22/24 showing some form of raising head, either on floor from grooming/sniffing up or often related to rearing (14/22)
13	unclear_sniffing	19/24 some stationary sniffing, but not specific
14	head_raise_left	19/24 raising head, leftwards bias, sometimes weak (17/19)
15	turn_left_with_retract	19/24 show left turn, most often after short retraction, happens often during down from rearing (leftwards) (14/19), but not specific
16	turn_right	23/24 show right turn, basically the rightwards version of 15, also with down from rearing often (16/23)
17	rearing_right	18/24 show some form of rearing/mounting, mainly rightwards (14/18)
18	unclear_sniffing2	18/24 show some form of sniffing on floor or in the air… maybe righward bias, but unclear head raise/down
19	unclear_stop	17/24 show some form of stopping from movement

20	rearing_start	17/24 show start of rearing some of the rest shows rearing related (down or currently rearing)
21	rearing_left_unclear	16/24 show some movement turning leftwards during a rear
22	sniffing_left	19/24 show some sniffing at walls while turning their head slightly left and moving sometimes forwards
23	rear_down	17/24 show down from rearing
24	unclear	No clear behavior
25	sniffing_right	21/24 show some sniffing, usually against walls, while turning head slightly right… rightwards version of 22
26	sniffing_left_turn	16/24 appear to be sniffing while turning slightly left
27	turn_right_stationary3	20/24 show some turning right in various contexts
28	turn_right_stationary4	19/24 show right turn without much other movement
29	movement_unclear	Appears to be a general movement syllable, maybe slight right turn, but not very specific
30	rearing_sniffing	17/24 show sniffing while on hindlegs, sometimes with movement forward
31	stop_with_head_down	16/24 not very clear, but appears to be rather abrupt stop with a short head down movement

2.4.4 Body center tracking

To explore any differences in mouse behavior, body center tracking was used to calculate the
distance moved by a single mouse (Figure 1B), its position (Figure 4A), or inter-mice distance
(Figure 4B-E, G). For tracking of body center, we used the mouse centroid calculated in
Keypoint-MoSeq by taking the median of all used body points (10 points on the mouse body,
excluding the tail). We then converted this pixel-based measure to a distance-based measure
using the pixel per millimeter resolution in our arenas, calculated based on the known distance
between the corners of the arena and the same distance in pixel measured in the video (measure

extracted through the Python package OpenCV2, version 4.11.0; using the `selectROI`
function).

2.4.5 Trajectory analysis

We extracted the syllable trajectories from the tracking information collected across all mice. To
evaluate similarity between a pair of syllable trajectories, we calculated the cosine distance
between position vectors of the pair of syllables, using the implementation in SciPy (version
1.14.0). Cosine distance is a measure of distance between two vectors (u and v) calculated as
$\mathbf{d_{cos}} = \mathbf{1} - \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{u}\|_2 \|\mathbf{v}\|_2}$. The cosine distance between all pairs is used for hierarchical clustering of
syllables in a dendrogram (Figure 2A, Suppl. Figure 3A). The data shown in Figure 2 is recorded
from the vehicle treated animals used in this study. As this study was part of a larger experiment,
40 additional mice of the same characteristics were recorded in parallel. When we used the larger
dataset and looked at the hierarchical clustering of syllables in a dendrogram (Suppl. Figure 3A),
we found 6 out of 8 dyadic-modulated syllables being separated into a distinct arm of the
dendrogram.

The dendrograms shown here were created based on the output of the functions `linkage` and
`dendrogram` contained in the SciPy package (Version 1.14.0) for Python. The exemplar
trajectories for each syllable (suppl. Figure 1A) were selected by the same algorithm as
implemented in `get_typical_trajectories` in the Keypoint-MoSeq package for Python.

2.4.6 Eigenvector centrality

To identify syllable nodes that may play transitory role in a directed network of syllable
transitions, we calculated eigenvector centrality measures for each syllable in the transition
network of syllables in solitary and dyadic context (Figure 2B). Eigenvector centrality measure
allows for the detection of nodes that are influential in the network, but not necessarily connected
to many nodes themselves. The eigenvector centrality is defined by the formula: $\mathbf{x_v} =$
$\frac{1}{\lambda} \sum_{\mathbf{t} \in \mathbf{M(v)}} \mathbf{x_t}$, with $\mathbf{M(v)}$ being the set of neighbors of node $\mathbf{v}$. To calculate eigenvector centrality,
we used the `eigenvector_centrality` function implemented in the NetworkX Python
package (version 3.3) on the normalized transition probabilities (outgoing transitions sum to 1
for a given source syllable) for each animal in solitary and dyadic contexts. Overall, for a given
syllable, we obtained 20 eigenvector centrality measures in each of solitary and dyadic contexts
(Figure 2B).

2.4.7 Social contact scoring

The experimenter was trained in the recognition of social behaviors and the usage of the built-in
video scoring functionality of Any-Maze and was blind to the light cycle and treatment of mice.
Since this study was part of a larger series, the scorer also scored further videos whose data is not
shown here. The scorer classified two main types of behaviors: *active social interaction*, that is
mice engaged in directed interactions, such as touching of the conspecific's body with the nose
and sniffing behavior; *passive social interaction* was identified as body contacts without active
sniffing. If there was uncertainty about how to score a specific segment, the scorer was instructed
to mark the segment as both states. The uncertain phases can occur, for example, at the transition
between active and passive contact, as the exact moment of transition can be unclear. Around

35% of active contacts and 10% of passive contacts were overlapping in this way. This overlap
can also be explained by the input delay on the keyboard used for scoring, and the fact that
continuous active contacts do not occur often and last only a few moments, while passive
interactions can last multiple seconds. Since we aimed to evaluate the time course of change in
syllable and syntax composition during active and passive social contact (see Figure 3B, C), we
pooled bouts with less than 6 seconds duration of the same contact type. This allowed for the
application of a 5 second (~125 frames) pre-contact control window, as shown in Figure 3B and
C. We used these pre-contact control windows to ensure that there was a clean onset at the
observed bouts, without other bouts of the evaluated behavior occurring in the control window.

2.4.8 Kullback-Leibler Divergence

To measure differences between distributions of syllables (Figure 3B) and syntax (Figure 4F),
we applied the Kullback-Leibler divergence (D_{KL}), or relative entropy. The D_{KL} measures the
distance between an observed probability distribution and an expected probability distribution. It
is defined as $\mathbf{D_{KL}(P||Q)} = \sum_{\mathbf{x \in X}} \mathbf{P(x)} \, \mathbf{log}\left(\frac{\mathbf{P(x)}}{\mathbf{Q(x)}}\right)$, with P being the observed and Q being the
expected distribution. In this study we used a base of 2 for the logarithm, leading to a relative
entropy in bits. To calculate the Kullback-Leibler divergence, we used the implementation
contained in the `entropy` function in SciPy (passing the distribution to be compared as `pk` and
the reference (expected) distribution as `qk`).

2.4.9 Parametric Behavior Classes

To gain a separation of behavioral syntax into parametrically defined behaviors, we used two
parameters - inter-mouse distance (IMD) and change in distance (CID) as shown in Figure 4C.
The peaks from kernel density plot in Figure 4C were separated into corresponding contact
related class (syntax starting and/or ending below an IMD < 100mm) and control class (marked
with hatched lines). Movement classes (approach and leave) were defined with an absolute CID
> 50mm and a contact threshold of 100mm based on the preferred IMD shown in Figure 4B. The
relative movement threshold (absolute CID) of 50mm was chosen as half of contact threshold.
These two thresholds along with limitations of the arena (maximum IMD and maximum CID ≤
diagonal of arena) resulted in the polygons defining approach and leave behavior classes.

Control classes were defined as those that did not lead to inter-mouse contact but matched the
same movement parameters (see dashed arrows in Figure 4C). For example, the control for
approach behavior was any movement that had an CID < -50mm and led to a final IMD between
100mm and 200mm. The leave control class was defined as CID > 50mm and a final IMD
between 100mm and 200mm. Finally, the contact control class used a larger distance criterion
and captured the largest peak in the IMD-CID distribution, at an IMD > 200mm and an absolute
CID < 50mm.

2.4.10   Hamming Distance

The hamming distance between two sequences is the number of positions that are different in the
two sequences. This only includes substitutions, but not rearrangements. In this study, we
defined the syntax family as a set of syntax with hamming distance < 1 relative to the name-
giving syntax – for the syntax family (9,0,5), the syntax (9,0,10) would be a member but not
(0,5,10). This was used to aggregate the diverse, but highly similar set of syntax influencing

inter-mouse approaches and leaves as shown in Figure 4D. Further analysis of these families can
be found in Figure 4E.

The formula for two sequences of identical length could be written as $d_{Hamming} =$
$\sum_{i=1}^{n_x} eq(x_i, y_i)$, with x and y representing the two sequences, and n_x being the length of
sequence x.

2.4.11 Principal Component Analysis (PCA)

We used dimensionality reduction with PCA to evaluate if DM syllables have a larger
contribution to the variability across videos (keypoint tracks) of mice in solitary and dyadic
contexts. We used syllable frame proportions for each individual mice, concatenated across
animals, in solitary and dyadic contexts as input to the PCA. Since syllable frequencies summed
up to 1, we did not apply further scaling to individual frequencies. The loadings for the first 15
PCs are split for DM or DU syllables (Figure 5A). Similarly, we also evaluated the contribution
of DM syntax to the variability across videos (keypoint tracks) of mice in solitary and dyadic
contexts (Figure 5B). We used the implementation of PCA in the scikit-learn package (version
1.5.1).

2.5 Statistical tests

286 All statistical tests were performed in Python (version 3.11.9). Unless stated otherwise we used a
287 significance threshold of α=0.05 and corrected for multiple testing by applying the Bonferroni
288 correction where applicable.

2.5.1 Mann-Whitney U-Test

We used the Mann Whitney U-test in the following analysis:

• In Figure 1B, we tested if the total distance moved differed between solitary and dyadic
recordings. Applying a one-sided test showed a significant difference between both
groups (p<0.0001).
• To find DM syllables, we applied a two-sided test in Figure 1C.
• As mentioned in the section on the eigenvector centrality measure, we tested if
eigenvector centrality differed between DM and DU syllables, as shown in Figure 2B.
We applied a two-sided test on the eigenvector centrality for DU and DM syllables,
extracted per recording and compared between solitary and dyadic recordings. 5 out of 8
DM syllables showed a significantly different eigenvector centrality (p≤0.05) between
solitary and dyadic recordings. No DU syllables showed a significant difference.
• Based on the results shown in Figure 2C, we tested for significantly modulated transitions
between solitary and dyadic contexts. We applied a two-sided test to the transition counts
observed in dyadic and solitary context for each transition observed in both contexts.
After correction for multiple testing, we found 9 significantly modulated transitions with
8 of those targeting DM syllables.
• We also applied a one-sided test on the absolute value of the PCA loadings shown in
Figure 5A, B to verify whether DM syllables or syntax provided a greater influence on
the first 5 PCs compared to DU syllables or syntax. The tests showed that DM syllables

309		had significantly higher loadings than DU syllables (p<0.002), but that the same effect
310		was not visible in DM syntax when compared to DU syntax.
311	•	The data shown in Suppl. Figure 1B was tested with a two-sided test and we did not find
312		significant differences between dyadic groups and the corresponding solitary onset
313		proportions (after correction for multiple comparisons).
314	•	For each parameter as shown in Suppl. Figure 1C, we performed a one-sided test between
315		DM syllable trajectories and DU syllable trajectories and found the two direction-based
316		parameters to be significantly reduced for the DM syllables (after correction for multiple
317		comparisons).
318	•	A two-sided test was applied for the pairwise comparisons of the data shown in Suppl.
319		Figure 2B. We found that all the comparisons were significant (after correction for
320		multiple comparisons).
321	•	The comparison between the count of unique syllable transitions observed in dyadic
322		context and solitary context yielded a significant result when we applied a one-sided test.

had significantly higher loadings than DU syllables (p<0.002), but that the same effect
was not visible in DM syntax when compared to DU syntax.
• The data shown in Suppl. Figure 1B was tested with a two-sided test and we did not find
significant differences between dyadic groups and the corresponding solitary onset
proportions (after correction for multiple comparisons).
• For each parameter as shown in Suppl. Figure 1C, we performed a one-sided test between
DM syllable trajectories and DU syllable trajectories and found the two direction-based
parameters to be significantly reduced for the DM syllables (after correction for multiple
comparisons).
• A two-sided test was applied for the pairwise comparisons of the data shown in Suppl.
Figure 2B. We found that all the comparisons were significant (after correction for
multiple comparisons).
• The comparison between the count of unique syllable transitions observed in dyadic
context and solitary context yielded a significant result when we applied a one-sided test.

We used SciPy's (version 1.14.0) implementation of the Mann-Whitney U-Test (function
`mannwhitneyu`) with significance levels set at $\alpha=0.05$.

2.5.2 χ^2 Contingency Test

The χ^2 contingency test is used to test for independence of frequencies of two groups
summarized in a contingency table. To verify which DM syllables showed a significant effect of
experimenter-scored contact in Figure 1D (left panel), we applied this test to the onset counts of
each syllable during and outside contact compared to the onset counts of any other syllable
during and outside contact. The test revealed two syllables (7 and 18) with significant effects
($p\leq0.05$, Bonferroni correction). In Suppl. Figure 6A, we applied the χ^2 contingency test to the
onset counts of each syllable family within or outside the behavior class, compared to the onset
counts of all length 3 syntax within or outside the same behavior class. We found all
comparisons to be significantly different, except for the comparison between syntax family
(11,2,8) to all other syntax in the leave control behavior class. We used the implementation
(`chi2_contingency` function) contained in the SciPy package for Python.

2.5.3 Two-way ANOVA

In Figure 1D, we compared the onset proportion of each DM syllable between solitary and
dyadic contexts. Within dyadic context, we tested for an interaction between onset proportions of
DM syllables and contact between conspecifics. Both contact (either experimenter-scored or
based on distance quartiles; for distance distribution, see Figure 4B) and DM syllable were
categorical variables. We performed a two-way ANOVA (formula `proportion ~`
`C(syllable) * C(contact)`) with the help of the `anova_lm` and `ols` functions contained in
the statsmodels package (version 0.14.2) for Python. Results showed significant main effects on
proportion and interactions between the categorical dependent variables (see Results).

2.5.4 Bonferroni correction

We corrected for multiple comparisons by using the Bonferroni correction. For example, we
used the following equation to adapt the significance threshold for the 32 syllables to maintain
our desired overall $\alpha=0.05$: $\alpha_{Bonferroni} = \alpha_{uncorrected}/n_{tests}$ with n_{tests} corresponding to the

number of syllables. This correction was applied to the results of the χ^2 contingency test used in
Figure 1D ($\mathbf{n_{tests}}$=8); Mann Whitney U-tests in Figure 2B ($\mathbf{n_{tests}}$ =32), Figure 2C ($\mathbf{n_{tests}}$ =99),
Suppl. Figure 1B ($\mathbf{n_{tests}}$ =6), Suppl. Figure 1C ($\mathbf{n_{tests}}$ =3), Suppl. Figure 2B ($\mathbf{n_{tests}}$ =6); the χ^2
contingency test used in Suppl. Figure 6A ($\mathbf{n_{tests}}$ =12).

2.5.5 Benjamini-Hochberg correction

To control for the false discovery rate in the Mann-Whitney U-test used to find the 8 DM
syllables (see Figure 1C), the implementation of the Benjamini-Hochberg procedure provided by
the SciPy package was applied. We used this in place of the Bonferroni correction as the
visualizations provided in the Keypoint-MoSeq package also used the Benjamini-Hochberg
correction.

2.5.6 Z-Score representation of syllable and syntax during contact

To identify whether the experimenter-scored contact behaviors had a significantly different
representation in syllable space when compared to the global average of behavior, we computed
z-score for each of the syllable or syntax occurring during contact behaviors using the
distribution of the same syllable or syntax in the predefined baseline window (Figure 3B, C). We
calculated the z-score for data shown in Figure 3B and C using the following formula: $\mathbf{z = \frac{x - \mu}{\sigma}}$.

In Figure 3B, a baseline period was defined between 5 seconds and 2 seconds prior to the scored
contact. The z-score of each trace in figure 3B was calculated based on the distribution of values
for the same trace within the baseline window. In addition, we also calculated z-score for the
active and passive contact traces based on the entire randomized trace (labeled "randomized" in
the legend). We evaluated both z-scores framewise for every trace and defined frames as
significant where both z-scores lay above 3. The frames found to be significant are marked for
each trace in the upper subplot.

In Figure 3C, we again defined a baseline period between 5 seconds and 2 seconds prior to the
scored contact. We computed z-score for each trace (syntax) based on the distribution of values
for the same trace within the baseline window. We again defined a z-score above 3 as significant.
All traces shown in black (active, n=25 and passive, n=24) showed significance across at least
50% of the test windows starting from 2 seconds prior to 2 seconds after the scored contact. The
remaining traces are shown in gray (n_{active}=79, $n_{passive}$=80). Four syntax traces with the highest z-
score were marked as the top 4 syntax associated with each contact type and are shown in color.

3   Results

We ran two open field tests (OFT) with video recordings, separated by 5 weeks, on 20 group housed female C57BL/6J mice. The first OFT recording contained an individual mouse in the open arena (referred to as solitary context), and the second OFT recording, performed after 5 weeks, contained two mice from separate cages in the same arena (referred to as dyadic context). In preparation of future pharmacological studies, all mice were orally injected with a vehicle treatment 20 minutes before the start of the recordings to account for potential adverse effects caused by the oral gavage procedure. Considering potential light cycle effects on behavior, one half of the mice were housed under normal, and the second half were housed under reversed light cycle conditions. However, we could not identify significant differences in syllable expression between the two light cycle groups and hence the data were pooled from the two light cycle groups.

We analyzed 20 minutes of video containing a single mouse in solitary and dyadic contexts using SLEAP (Pereira et al. 2022) to track a set of 12 key body points on each mouse. We fitted a Keypoint-MoSeq model (Weinreb et al. 2023) to a subset of 10 key body points to extract repeated elements of mice behavior, referred to as behavioral syllables (see Methods; Figure 1A). The output syllables were aggregated by either their onset (bout) proportion, representing the proportion of syllable bouts regardless of individual bout lengths in frames, or frame proportion, representing the total proportion of the recorded frames in one syllable. After filtering for total onset proportion, we extracted a total of 32 syllables which were used in subsequent analyses (see Methods; Suppl. Figure 1A). Syllables with a bout proportion below the threshold often occurred in clusters (Suppl. Figure 2A) and were observed to relate mostly to self-grooming (Suppl. Figure 2C, D). These clusters occurred in both solitary and dyadic contexts but did not meaningfully differ between the two contexts (Suppl. Figure 2B).

3.1   Dyadic context modulates a small subset of MoSeq identified syllables

To investigate the structure of social behavior through behavior decomposition, we first asked if behavioral 'syllables' identified through decomposition (Keypoint-MoSeq) are sensitive to modulation in solitary and dyadic contexts and whether this modulation is dependent on physical proximity between the two animals. Before comparing syllables in solitary and dyadic contexts, we verified whether parameters related to movement revealed significant changes in mouse behavior between solitary and dyadic contexts (L. M. von Ziegler et al. 2023). By approximating the output of traditional body center tracking (see Methods), we detected a significant reduction in distance moved of mice between solitary and dyadic contexts (Mann-Whitney U test, $p \approx 1.83e\text{-}5$; Figure 1B).

A comparison of onset proportion of all 32 syllables in solitary and dyadic contexts revealed a subset of 8 syllables that differed significantly between contexts (Figure 1C; see Methods; Mann-Whitney U-Test, Benjamini-Hochberg correction, $p \leq 0.05$). From here onwards, the 8 significantly modulated syllables will be referred to as dyadic-modulated (DM) syllables and the remaining 24 syllables will be referred to as dyadic-unmodulated (DU) syllables.

Figure 1. *Identification of behavioral syllables in solitary and dyadic contexts.* (A) Panels left to right: C57BL/6J mice were placed into an open field arena, 20 minutes after an oral vehicle injection, in solitary or dyadic context. After video recording in the open field arena, Key points on the body and tail were tracked using SLEAP; key points on the body were used to extract behavioral syllables using Keypoint-MoSeq (see Methods). (B) A comparison of the distances moved in solitary and dyadic context recordings. Asterisk indicates statistical significance (Mann-Whitney U-test, p<0.0001; see Results). (C) For each of the 32 syllables, a comparison of onset proportions between solitary and dyadic context (Mann-Whitney U-Test, Benjamini-Hochberg correction, p<0.05) reveals 8 significantly modulated (DM; orange colored syllable numbers) and 24 unmodulated syllables (DU). (D) Left panel: A comparison of DM syllables' onset proportion based on its occurrence during or outside of an active contact behavior as scored by an experimenter (see Methods). We performed a χ2 proportion test for each syllable's onset proportion. 2 syllables (7 and 18) show a significant (Bonferroni correction, p<0.05) difference during and outside of an active contact in the dyadic context. Right panel: A comparison of DM syllables' onset proportion based on the distance between the mice, divided into quartiles, during the syllable occurrence (see Methods). (E) Trajectories of the 8 DM syllables belonging to stationary (7, 12, 13, 18, 23, 30) or non-directional movement (all). (F) A selection of DU syllable trajectories with directed (0, 2, 9, 11) and traversal movement (0, 2, 8, 9, 22). In panels C & D, filled circles and error bars indicate mean and 95% CI respectively.

Because the mice move less in dyadic context (Figure 1B), one may deduce that DM syllables are likely associated with active contact but may not be influenced by physical separation between mice outside of an active contact. To evaluate the effect of physical proximity on DM syllables, we compared the onset proportions of syllables during experimenter-scored active contact (for definition see Figure 3A) to no contact. Results from a two-way ANOVA revealed a significant main effect for DM syllables and active contact ($F(7, 299)=49.61$, $p<1e-45$; $F(1, 299)=5.07$, $p=0.03$) with a significant interaction between DM syllables and active contact ($F(7, 299)=2.21$, $p=0.03$). However, a closer inspection for each syllable revealed few significant syllables (2 out of 8, χ^2 contingency test, Bonferroni correction, $p<0.05$). In addition, we checked if the distance between mice is associated with the modulation of the DM syllables by comparing onset frequency in different distance quartile. A two-way ANOVA revealed a significant main effect for DM syllables and distance quartiles ($F(7, 603)=93.62$, $p<1e-91$; $F(3, 603)=3.87$, $p=0.01$) with a significant interaction between DM syllables and distance quartile ($F(21, 603)=8.67$, $p<1e-22$). These findings reveal that majority of the DM syllables are not directly linked to an active contact but are associated with physical proximity to a conspecific (Figure 1D, left panel; but also Suppl. Figure 1B).

An inspection of the trajectories of all DM syllables reveals high proportion of stationary, and non-directional behaviors, for example rearing as in syllable 7 (Figure 1E). In contrast, most directional behavior (left or right oriented movement as in syllable 9 and 11) falls into the DU group (Figure 1F and Suppl. Figure 1C; one-sided Mann-Whitney U-test, Bonferroni correction, $p<0.05$). To quantify the differences within and across DM and DU syllables, we used cosine distance to quantify similarity (see Methods; Figure 2A). The dendrogram of similarity measure did not reveal any systematic clustering within and across DM and DU syllables. However, using an extended dataset (see Methods), we found that most of the DM syllables are similar when compared to DU syllables (Suppl. Figure 3A).

These results show that a subset of behavioral 'syllables' identified through decomposition are sensitive to modulation in solitary and dyadic contexts. Surprisingly, most of the modulated syllables, dominated by stationary undirected behaviors, are not linked to an active contact but are associated with physical proximity to a conspecific outside of an active contact.

3.2 Modulated syllables drive the syllable transitions in dyadic context

Because there was no association of an active contact with most behavioral syllables in the presence of a conspecific (dyadic context), we next investigated whether there is a distinct role for DM syllables in the transitions between behavioral syllables to form a complex social behavior.

We employed Eigenvector centrality to quantify the role of DM syllables in the transitions between syllables (see Methods). Briefly, Eigenvector centrality is a measure of how well a node is connected to nodes which themselves are well connected to individual nodes within a network (Negre et al. 2018). Here, the Eigenvector centrality of a syllable is calculated based on the directed graph of transition probabilities. We found a significant change in eigenvector centrality, during the dyadic context, for most of the DM syllables but not for any of the DU

463   syllables (Figure 2B; two-sided Mann-Whitney U-test, Bonferroni correction, p≤0.05). This
464   finding reveals a crucial transitory role for DM syllables in the network of syllables defining
465   complex social behavior in the presence of a conspecific (dyadic context).

Figure 2. *Role of dyadic modulated syllables in the network of syllable transitions during solitary and dyadic contexts.* (A) The dendrogram of inter-syllable cosine distances shows limited clustering of DM syllables (orange colored syllable numbers). (B) A comparison of eigenvector centrality measure between solitary and dyadic contexts (see Methods), based on the network of transition probabilities (outgoing probabilities were normalized to sum 1) between syllables, reveals significant differences for DM syllables (5 out of 8) but not for DU syllables. Asterisk indicates statistical significance (two-sided Mann-Whitney-U test, Bonferroni correction, p<0.05). Filled circles and error bars indicate mean and 95% CI. (C) A directed network of syllable transitions modulated by dyadic context with edges indicating transition probabilities between syllable nodes. Edge colors indicate negative (blue) or positive (red) modulation of the transition probability in dyadic context; edge thickness indicates amount of change in transition probability in dyadic context. Syllables are nodes in the network and node diameter corresponds to changes in eigenvector centrality of that node. Syllable nodes with a significant change in eigenvector centrality are indicated in bold syllable numbers (see panel B).

During the dyadic context, most of the significantly modulated transitions (8/9, two-sided Mann-
Whitney U-test, Bonferroni correction, p≤0.05) target DM syllables as reflected by eigenvector
centrality (see arrows pointing to the first column with DM syllables in Figure 2C). In addition,
we observed an overall increase in syllable transitions (Suppl. Figure 3B; one-sided Mann-
Whitney U Test, p≤0.05) and the strongest changes in transition probabilities (thick lines in
Figure 2C) were mostly a result of reduced transition probabilities (blue lines), originating in DU
syllables (see Suppl. Figure 3C).

Due to a weak association between contact and DM syllables (Figure 1D), the exact role of these
syllables in social interactions is not clear. However, we observed that behavior in general
becomes less stereotypical (or structured) when mice are in a dyadic context (see Suppl. Figure
3D). Regardless, these results reveal an important role for DM syllables as "transfer nodes" in
the behavioral program to construct a complex social behavior in the presence of a conspecific
(dyadic context).

481 3.3   Changes in syllable composition reflect active and passive contact behaviors

Given that DM syllables play an important role of 'transfer node' in the behavioral syllable
transitions to form complex behavior in dyadic context, we investigated whether complex
behaviors, such as active or passive contact behaviors, are captured through changes in the
composition of syllables or sequences of syllables, referred to as syntax in the following
observations.

For this purpose, we tasked a fellow researcher to score active and passive social contacts in the
dyadic recordings (see Methods). To quantify changes in syllable composition, in an individual
mouse during active and passive behaviors, we computed Kullback-Leibler divergence (D_{KL}), a
measure of statistical distance between two distributions, between framewise distribution of
syllable proportions in active or passive contacts to an overall distribution of syllable proportions
in dyadic context (see Methods). The time course of D_{KL} reflects changes in syllable composition
aligned to start of active or passive behaviors (Left panel in Figure 3B). The results show a
significant change in syllable composition (D_{KL}), in an individual mouse, at the time of active
and passive behaviors. We verified that this change is fully attributable to active and passive
behaviors since the same analysis on randomized sequence of syllables did not show any change
in syllable composition (Figure 3B; z-score see Methods).

We next asked if active and passive behaviors also reflect changes in joint syllable composition
of two mice by computing framewise Kullback-Leibler divergence (D_{KL}) between joint
distributions of syllable proportions between two mice during active or passive contacts to an
overall joint distribution of syllable proportions in dyadic context. The time course of D_{KL} shows
a weak but continuous and significant change in joint syllable composition aligned to start of
active behavior but less so for passive behavior (Right panel in Figure 3B).

To investigate if sequence of syllables (syntax) signal active and passive contact behaviors, we
extracted length 3 syntax (i.e., sequences comprising 3 syllables), as this corresponded to about a
second of behavior with a median syllable duration of 10 frames (see Methods). A multitude of
syntax are significantly associated with active (25 syntax) and passive (24 syntax) contact
behaviors (Figure 3C; see Methods for statistics). However, there was little overlap of syntax
between active and passive contact behaviors (Figure 3C, 8 significantly associated syntax
shared between active and passive). In addition, we observed few DM syllables in the top 4 most
significantly associated syntax for both contact behaviors. Furthermore, inspection of example
trajectories for the two syntax most associated with active contact suggests a behavior related to
circling around a conspecific (Suppl. Figure 4A, B). Similarly, the syntax most associated with

passive contact included mostly stationary behavior, while the second syntax was related to a
turning behavior followed by movement (Suppl. Figure 4C, D).

Figure 3. *Syllable and syntax composition during experimenter scored active and passive contact behaviors.* (A) Schematic of an experimenter scoring active and passive contact between mice in the dyadic video recordings. For details on scoring instructions, see Methods. (B) Left-bottom panel: Time course of difference in frame-wise syllable composition of an individual mice, quantified as DKL, between experimenter-scored behavior and all behaviors is plotted for active and passive behaviors. Left-Top panel: Time course of statistical significance for data in bottom panel compared to shaded baseline window. Right panel: Same as Left panel but using frame-wise syllable composition of two mice instead of a single mouse (see Methods). (C) The frequency of length 3 syllable syntax was aligned to the start of a scored behavior and transformed to z-score using the baseline window (from 100ms to 50ms of the start; shaded window). Time course of syntax aligned to the start of scored behavior is shown for active contact (left panel) and passive contact (right panel). The top 4 syntax are indicated with colored lines and their corresponding syllable composition is shown in legend. Orange colored syllable number indicates DM syllable.

These results show that experimenter defined active and passive contact behaviors are
represented in the syllable space and there are a diverse set of syllable sequences that are
modulated in contact behaviors. Furthermore, DM syllables are underrepresented in the most of
the contact-associated syllable sequences, in line with the previously observed modulation of
DM syllables regardless of contact (Figure 1D), suggesting their role in aiding transitions
between syntax associated with contact behaviors.

3.4 Individual syllable sequences do not map onto specific behaviors

The finding that experimenter defined contact behaviors are represented in syllable and syntax space prompted us to ask whether social behaviors, such as social approach and leave defined in a parametric space (Peleh, Bai, et al. 2019), can be mapped onto syllable and syntax space. First, we verified if there is a propensity for targeted behavior in our dataset by calculating distributions of mice positions (Figure 4A) and probability of inter-mouse distance (Figure 4B). Our data shows that mice preferred to position themselves in corners of arena (Figure 4A), and that the co-location of two mice in the same corner was a coordinated event, rather than a random occurrence (Figure 4B).

Before parametrically defining behaviors, we extracted inter-mouse distance during length 3 syntax and mapped them onto a space representing the inter-mouse distance at the start of syntax and change in distance by the end of the syntax (Figure 4C). From the density plot of all syntax in this space (blue contours in Figure 4C), we were able to identify four density zones that specify parametrically defined behaviors – targeted approach and leave, as well as stationary behaviors in low distance (contact) and stationary behaviors with high distance. We further defined control zones as those that showed the same relative movements but did not relate to contact between the mice (see dotted arrows and hatched areas in Figure 4C & Methods). We choose to use syntax rather than syllables since syntax timescales were more apt for the description of targeted long-distance movements and longer stationary behaviors.

For the two distinct behaviors of social approach and leave (Figure 4C), we investigated whether there is a dominant representation by specific syntax. We tracked the change in distance over a rolling 30 frame window, matching our definition of length 3 syntax. It is evident that nearly all the windows in which inter-mouse distance changed by more than 200mm were fully represented by either one of two syntax families (see Figure 4D). A syntax family is defined as a set of syntax with at most one changed syllable relative to a reference syntax (the family was named after the reference syntax; see Hamming Distance in Methods). We found that social approach and leave were fully represented by two syntax families (syntax 9,0,5 and syntax 11,2,8; see Figure 4D) and both contained a syllable with a long-distance traversal trajectory (syllables 9&0 and 2 respectively, see Figure 1F), suggesting both syntax families capture behavior either targeted towards or away from a conspecific. Inspection of sample trajectories included in these syntax families revealed that one syntax family, (9,0,5), was usually associated with a leftwards turn followed by a movement straight ahead while the other syntax family, (11,2,8), was associated with a rightwards turn followed by movement (Suppl. Figure 5A, B). We interpret these syntaxes as corresponding to a mouse reorienting towards or away from a conspecific, during an approach and leave respectively.

We asked if the above finding that social approach and leave were fully represented by two syntax families extended in the opposite direction i.e. whether the two syntax families specifically map onto the social approach and leave behaviors. When we mapped all occurrences of the two syntax families onto parameter space (Figure 4E), we did not identify a direct mapping between a syntax family and a specific behavior class as defined in our parametric space (Suppl. Figure 6A; χ^2 contingency test, Bonferroni correction, $p \leq 0.05$, but see Methods). Since we could not identify an association between a syntax family and a behavior, we checked if syllable composition differed between behavior classes. We calculated D_{KL} on the syllables

contained within each syntax, between a pair of behavior classes. We observed no separation
between approach and leave classes further confirming that there is not a specific syntax, at least
in our dataset, that directly maps onto approach and leave classes. Furthermore, we noted a
modest separation between all movement classes (approach and leave) and stationary (contact)
classes respectively (Figure 4F, for analysis of syntax see Suppl. Figure 6B). However, mapping
all occurrences of syntax associated with active contact (from Figure 3C) onto our parameter
space did not reveal any specific syntax associated with active contact behavior (Figure 4G). We
also confirmed that there was no specific syntax associated with passive contact behavior (Suppl.
Figure 6C).

Figure 4. *Syntax and parametric behavior classes.* (A) 2D kernel density estimate of mouse positions sampled from the recordings in dyadic context (2 million data points). Marginal plots show isolated kernel density estimate of each axis. (B) Histograms of inter-mouse distances in observed data, shuffled and random control data. Observed refers to inter-mouse distances in recorded data; shuffled refers to inter-mouse distances with the individual positions being shuffled; Random refers to inter-mouse distances simulated with randomly selected positions within the arena (10 million point-pairs). The difference of the observed and shuffled histogram is shown in black. (C) Parametric behavior classes were derived from observed length 3 syntax. Inter-mouse distance (x-axis) and change of that distance (y-axis) were plotted as a kernel density estimate shown as blue contours (KDE, see methods). Based on clusters in the KDE, we defined 3 behavior classes: contact (x<100mm, x+y<100mm), approach (x>100mm, x+y<100mm), and leave (x<100mm, x+y>100mm). Control classes were defined to match change in distance (y-axis) but excluding contact scenarios (see dotted arrows). (D) A rolling window (length 30 frames) was used to calculate change in inter-mouse distance (x-axis) and is plotted against the proportion of the window spent in one of the two syntax family (9,0,5) or (11,2,8) (see Methods). Line with shaded region represents median and 95% CI. (E) KDE with same parameters as in C but limited to one of the two syntax families referred to in panel D. The KDE from panel C was subtracted as baseline. Red areas represent higher density than in C, blue areas with reduced relative density. (F) Heatmap of DKL between syllable distributions found in syntax belonging to the behavior classes defined in panel C. Darker colors indicate lower DKL. (G) Top 4 syntax correlated with experimenter-scored active contact, taken from Figure 3C, plotted as scatter on top of the KDE taken from panel C.

These findings show that while mapping of a behavior to syllable space is possible with a one-to-
many mapping, we could not find evidence for distinct syllable sequences that map onto specific
behaviors.

3.5 Syllables allow discrimination of solitary and dyadic behavioral modes

Based on the lack of evidence for specific syntax or syllable descriptors of social behavior, we
aimed to find out if syllables or syntax provide discrimination of solitary and dyadic contexts.
First, we evaluated the variance captured by syllables and syntax using principal component
analysis on the proportion of frames containing each syllable for each mouse (see Methods). We
separated the contributions of dyadic modulated and unmodulated syllables to each of the
principal components (Figure 5A). Results reveal that DM syllables capture significantly more
variance compared to DU syllables – first five PCs explaining ~73% variance have significantly
higher DM syllable loadings (one-sided Mann Whitney U-test, p<0.005; see Figure 5A).
However, we did not find such a difference for syntax (same test, Figure 5B).

Furthermore, using the above identified principal components, we observed a clear separation
between solitary and dyadic contexts, while the same approach did not lead to a separation
between two light cycle conditions (Figure 5C). These results demonstrate that syllables and

syntax, particularly DM syllables, provide a sensitive description of behavior to discriminate if
behavior is in the presence or absence of a conspecific.

Figure 5. *Syllables and syntax discriminate solitary and dyadic context but not normal and reverse light-cycle.* (A) Results from principal component analysis (PCA) of frame-wise occurrences of modulated (DM) and unmodulated (DU) syllables in dyadic and solitary recordings are shown (see Methods). The box plots show (with y-axis on the left) the absolute of the loadings on each component for DM and DU syllables. The y-axis on the right shows the percentage variance explained by each of the component. The first 5 PCs explain 74% of the variance and DM syllables contribute significantly more to the loadings of these PCs than DU syllables (one-sided Mann-Whitney U-test, Bonferroni correction, p<0.05). (B) Results from PCA on syntax proportions are shown with same conventions as in panel A. DM syntax was defined as containing at least 2 DM syllables. (C) Scatter plot of the first 2 PCs from syllable-based PCA (top plots) or syntax-based PCA (bottom plots). Colors separate tracks of individual mice in either solitary/dyadic recordings (left plots) or normal/reversed light cycles (right plots).

4   Discussion and conclusion

A growing number of tools are available to accomplish the data-driven segmentation of the full behavioral repertoire of mice (See reviews: Choi & Kumar, 2024; Datta et al., 2019; L. von Ziegler, Sturman, & Bohacek, 2021). However, it is not known if such an approach, to segment or decompose the behavior into its recurring elements, yields insights into the structure of social behavior in mice. Here, we investigated the structure of social behavior using Keypoint-MoSeq (Weinreb et al. 2024) to videos of mice in a dyadic and solitary context. We found that in the presence of a conspecific, mice moved less and significantly modulated the usage of a small subset of mostly stationary and undirected syllables, referred to as dyadic modulated (DM) syllables (Figure 1B, C; Suppl. Figure 1A, C). Surprisingly, for most of the DM syllables, this modulation in usage was not associated with the occurrence of experimenter-scored contact between the mice (Figure 1D). Instead, DM syllables played a significant role as transitory nodes in the syllable transition network (Figure 2B, C). Importantly, we found significant changes in composition of syllables and syntax (length 3 syllable sequence) aligned to experimenter-defined contact behaviors (Figure 3B, C). In addition, we identified 2 syntax families that captured approach and leave behavior as defined through parametric thresholds (Figure 4C, D; Suppl. Figure 5A, B). However, we did not find evidence for distinct syntax that map onto specific complex behaviors. Overall, our results show that behavioral syllables can be used as descriptive and sensitive markers to detect changes in behavior in the presence of a conspecific, but specific social behaviors may not map onto individual behavioral syllables or a specific syntax thereof.

A possible caveat of our study could be that the experiments were performed with one set of mice across both contexts. We ran the two recordings (solitary, dyadic) in the same set of open field arenas 5 weeks apart. It is possible that during the second recording the arena may not have been a novel environment for mice. Based on available literature the long-term spatial memory of the arena will be negligible after a span of 5 weeks, as most assessments of spatial memory work with periods of up to 2 weeks after training session over multiple days (Sharma, Rakoczy, and Brown-Borg 2010). Similarly, movement patterns of our mice could potentially also evolve during inter-experimental phase (5 weeks) and might have contributed to our findings, but it can be expected that social contact with a conspecific outweighs these effects. The estrous cycle most likely varies across individuals and recordings and might contribute to social preference (Chari et al. 2020). However, as solitary and dyadic context behaviors were clearly distinguishable in our dataset, we conclude that the estrous cycle-related variance must be below the level of behavioral changes in a dyadic context.

To detect as much of the behavioral repertoire as possible the Keypoint-MoSeq model was fitted to the entire recorded dataset, including full one hour solitary recordings and 20-minute dyadic recordings. Potentially this 3:1 disbalanced dataset could introduce a bias towards behaviors shown in solitary context, but as we treated the solitary state as the baseline, effects of the dyadic context would still be visible in the following analysis. In addition, we limited the analysis of the extracted syllables to the first 20 minutes of all recordings to restore data parity and analyze segments of identical activity (see Methods). Distinct behaviors occurring only in a dyadic context would have been visible with this approach, like behavioral syllables related to self-grooming (see Suppl. Figure 2C, D). While some distinct behaviors (self-grooming and jumping) were captured as individual syllables as expected, these occurred too rarely to be included in our analysis (see Methods).

It was possible to detect contact-specific changes in the behavioral syllable composition for
individual mice (Figure 3B, left panel), but we could not observe the same prominent effect for
the joint syllable composition across two mice (Figure 3B, right panel). This difference in results
between individual mice and two mice is very likely because joint occurrences of syllables are
sparse, and a higher number of joint occurrences are necessary when analyzing the joint behavior
of two mice. Since the significant effect of contact occurrence on the syllable distribution in
individual mice is clear in our dataset, we decided to show the across-mice effect as a
comparison to encourage further studies. Perhaps, an approach to decompose multi-animal
behavior, from large datasets to overcome sparse social events, instead of decomposing
individual animal behavior could be beneficial in detecting 'social' syllables.

Social interactions capture rich and multi-modal behavioral repertoire (Porcelli et al. 2019) and
thus need a data-driven extraction of behaviors. Recent studies addressed this need by using
either parametric methods (Peleh, Bai, et al. 2019) or machine learning applications (Goodwin et
al. 2024; Lin et al. 2024). It is also necessary to aim for explainable readouts from data-driven
analysis of complex behaviors and a fruitful approach is to limit analysis to a small set of well-
defined parametric approximations. We applied this perspective in our analysis (Figure 4C) and
defined a set of four parametric categories in the context of mouse interaction (close contact,
approach, leave, other stationary behaviors). Our original expectation with the behavioral
decomposition approach was to find distinct syllables or sequence of syllables (syntax) that
represent the parametrically defined categories. While we did find syntax families that capture
most occurrences of social approach and leave (Figure 4C, D), a direct and distinct mapping of
behavioral syllables onto social behavior classes was not observable (Figure 4E, G). It is possible
that behavioral decomposition based on the key points may have contributed to the lack of
distinct mapping of behavioral syllables onto social behaviors in our dataset because extraction
of key points inherently limits the amount of pose information unlike pose extracted from depth
recordings (Lin et al. 2024). In addition, a fruitful approach could be to use the joint pose
information between mice, and this requires overcoming the limitation of sparseness in social
interaction datasets.

In summary, our results show that, in the presence of a conspecific, mice possess a subset of
behavioral syllables that were significantly modulated, and these modulated syllables change
their transitory hub-like role in the transition network of all syllables. While we did not find
bidirectional mapping between a syllable or syntax and specific social behavior, our results show
that syllables or syntax are sensitive descriptors to detect changes in behavior because of change
in social context (Figure 5C). This finding taken together with previous studies, showing the
sensitivity of behavioral syllables to detect different pharmacological interventions (Wiltschko et
al. 2015; 2020), suggests that manipulation of experimental context (social or non-social
enrichment) may offer to be a valuable dimension in phenotypic screening of drugs using
behavioral decomposition methods.

Table of abbreviations

Abbreviation	Full Term
DM	Dyadic Modulated
DU	Dyadic Unmodulated
PCA	Principal Component Analysis
IMD	Inter-Mouse Distance
CID	Change In Distance
RDoC	Research Domain Criteria
MoSeq	Motion Sequencing
LED	Light Emitting Diode

Acknowledgements

We want to thank the other members of the data science team for their feedback on this project.
We also thank Alina Ritter for her support during the recording of data, and Michael Winter, Zoi
Balla, Johannes Freudenreich, and Carmen Weiss for their help using software related to the
recording of data.

Data availability statement

The original contributions presented in the study are publicly available. This data can be found
here: https://doi.org/10.5281/zenodo.15237564

Code availability statement

The code used to generate the data shown in this study is publicly available. It can be found here:
https://github.com/Marti-Ritter/social-context-restructures-behavioral-syntax-in-mice

Ethics statement

All procedures followed the regulations for animal experimentation enforced by the local district
administration's animal welfare commissioner of the state of Baden-Württemberg.

Author Contributions

	Conceptualization	Data curation	Formal analysis	Funding acquisition	Investigation	Methodology	Project administration	Resources	Software	Supervision	Validation	Visualization	Writing – original draft	Writing – review & editing
MR	x	x	x		x	x			x		x	x	x	x
HS		x	x		x						x	x		
SD	x					x	x	x		x				x
BH	x									x				x
CW	x			x		x	x			x				x
MB	x									x				x
AB	x					x	x			x		x		x

Funding

The authors declare that financial support was received for the research, authorship, and/or publication of this article. This work was supported by Boehringer Ingelheim Pharma GmbH & Co. KG, Biberach, Germany.

Conflict of interest

Marti Ritter, Hope Shipley, Serena Deiana, Bastian Hengerer, Carsten T. Wotjak, and Amarender Bogadhi received salaries from Boehringer Ingelheim Pharma GmbH & Co. KG.

Generative AI statement

The authors declare that no Generative AI was used in the creation of this manuscript.

References

Abeelen, J. H. F. van. 1964. "Mouse Mutants Studied by Means of Ethological Methods." *Genetica* 34 (1): 79–94. https://doi.org/10.1007/bf01664181.

Anderson, David J., and Pietro Perona. 2014. "Toward a Science of Computational Ethology." *Neuron* 84 (1): 18–31. https://doi.org/10.1016/j.neuron.2014.09.005.

Bewley, Alex, Zongyuan Ge, Lionel Ott, Fabio Ramos, and Ben Upcroft. 2016. "Simple Online and Realtime Tracking." *2016 IEEE International Conference on Image Processing (ICIP)*, 3464–68. https://doi.org/10.1109/icip.2016.7533003.

Bohic, Manon, Luke A. Pattison, Z. Anissa Jhumka, Heather Rossi, Joshua K. Thackray, Matthew Ricci, Nahom Mossazghi, et al. 2023. "Mapping the Neuroethological Signatures of Pain, Analgesia, and Recovery in Mice." *Neuron* 111 (18): 2811-2830.e8. https://doi.org/10.1016/j.neuron.2023.06.008.

Bolivar, Valerie J., Samantha R. Walters, and Jennifer L. Phoenix. 2007. "Assessing Autism-like Behavior in Mice: Variations in Social Interactions among Inbred Strains." *Behavioural Brain Research* 176 (1): 21–26. https://doi.org/10.1016/j.bbr.2006.09.007.

Bordes, Joeri, Lucas Miranda, Bertram Müller-Myhsok, and Mathias V. Schmidt. 2023. "Advancing Social Behavioral Neuroscience by Integrating Ethology and Comparative Psychology Methods through Machine Learning." *Neuroscience & Biobehavioral Reviews* 151:105243. https://doi.org/10.1016/j.neubiorev.2023.105243.

Cacioppo, John T., Stephanie Cacioppo, Stephanie Dulawa, and Abraham A. Palmer. 2014. "Social Neuroscience and Its Potential Contribution to Psychiatry." *World Psychiatry* 13 (2): 131–39. https://doi.org/10.1002/wps.20118.

Casey, B. J., Nick Craddock, Bruce N. Cuthbert, Steven E. Hyman, Francis S. Lee, and Kerry J. Ressler. 2013. "DSM-5 and RDoC: Progress in Psychiatry Research?" *Nature Reviews Neuroscience* 14 (11): 810–14. https://doi.org/10.1038/nrn3621.

Chari, Trishala, Sophie Griswold, Nick A. Andrews, and Michela Fagiolini. 2020. "The Stage of the Estrus Cycle Is Critical for Interpretation of Female Mouse Social Interaction Behavior." *Frontiers in Behavioral Neuroscience* 14:113. https://doi.org/10.3389/fnbeh.2020.00113.

Chaumont, Fabrice de, Renata Dos-Santos Coura, Pierre Serreau, Arnaud Cressant, Jonathan Chabout, Sylvie Granon, and Jean-Christophe Olivo-Marin. 2012. "Computerized Video Analysis of Social Interactions in Mice." *Nature Methods* 9 (4): 410–17. https://doi.org/10.1038/nmeth.1924.

Chaumont, Fabrice de, Elodie Ey, Nicolas Torquet, Thibault Lagache, Stéphane Dallongeville,
Albane Imbert, Thierry Legou, et al. 2018. "Live Mouse Tracker: Real-Time Behavioral
Analysis of Groups of Mice." *BioRxiv*, 345132. https://doi.org/10.1101/345132.

Chevance, Astrid, Philippe Ravaud, Anneka Tomlinson, Catherine Le Berre, Birgit Teufer,
Suzanne Touboul, Eiko I Fried, Gerald Gartlehner, Andrea Cipriani, and Viet Thi Tran. 2020.
"Identifying Outcomes for Depression That Matter to Patients, Informal Caregivers, and
Health-Care Professionals: Qualitative Content Analysis of a Large International Online
Survey." *The Lancet Psychiatry* 7 (8): 692–702. https://doi.org/10.1016/s2215-
0366(20)30191-7.

Choi, Jessica D., and Vivek Kumar. 2024. "A New Era in Quantification of Animal Social
Behaviors." *Neuroscience & Biobehavioral Reviews* 157:105528.
https://doi.org/10.1016/j.neubiorev.2023.105528.

Cuthbert, Bruce N. 2018. "The PRISM Project: Social Withdrawal from an RDoC Perspective."
Neuroscience & Biobehavioral Reviews 97:34–37.
https://doi.org/10.1016/j.neubiorev.2018.08.006.

Datta, Sandeep Robert, David J. Anderson, Kristin Branson, Pietro Perona, and Andrew Leifer.
2019. "Computational Neuroethology: A Call to Action." *Neuron* 104 (1): 11–24.
https://doi.org/10.1016/j.neuron.2019.09.038.

Dennis, Emily Jane, Ahmed El Hady, Angie Michaiel, Ann Clemens, Dougal R. Gowan Tervo,
Jakob Voigts, and Sandeep Robert Datta. 2021. "Systems Neuroscience of Natural Behaviors
in Rodents." *The Journal of Neuroscience* 41 (5): 911–19.
https://doi.org/10.1523/jneurosci.1877-20.2020.

Ebbesen, Christian L., and Robert C. Froemke. 2022. "Automatic Mapping of Multiplexed
Social Receptive Fields by Deep Learning and GPU-Accelerated 3D Videography." *Nature
Communications* 13 (1): 593. https://doi.org/10.1038/s41467-022-28153-7.

Fan, Zhengxiao, Hong Zhu, Tingting Zhou, Sheng Wang, Yan Wu, and Hailan Hu. 2019. "Using
the Tube Test to Measure Social Hierarchy in Mice." *Nature Protocols* 14 (3): 819–31.
https://doi.org/10.1038/s41596-018-0116-4.

Forkosh, Oren, Stoyo Karamihalev, Simone Roeh, Uri Alon, Sergey Anpilov, Chadi Touma,
Markus Nussbaumer, et al. 2019. "Identity Domains Capture Individual Differences from
across the Behavioral Repertoire." *Nature Neuroscience* 22 (12): 2023–28.
https://doi.org/10.1038/s41593-019-0516-y.

Garner, Joseph P. 2014. "The Significance of Meaning: Why Do Over 90% of Behavioral
Neuroscience Results Fail to Translate to Humans, and What Can We Do to Fix It?" *ILAR
Journal* 55 (3): 438–56. https://doi.org/10.1093/ilar/ilu047.

Giancardo, Luca, Diego Sona, Huiping Huang, Sara Sannino, Francesca Managò, Diego
  Scheggia, Francesco Papaleo, and Vittorio Murino. 2013. "Automatic Visual Tracking and
  Social Behaviour Analysis with Multiple Mice." *PLoS ONE* 8 (9): e74557.
  https://doi.org/10.1371/journal.pone.0074557.

Gomez-Marin, Alex, Joseph J Paton, Adam R Kampff, Rui M Costa, and Zachary F Mainen.
  2014. "Big Behavioral Data: Psychology, Ethology and the Foundations of Neuroscience."
  Nature Neuroscience 17 (11): 1455–62. https://doi.org/10.1038/nn.3812.

Goodwin, Nastacia L., Jia J. Choong, Sophia Hwang, Kayla Pitts, Liana Bloom, Aasiya Islam,
  Yizhe Y. Zhang, et al. 2024. "Simple Behavioral Analysis (SimBA) as a Platform for
  Explainable Machine Learning in Behavioral Neuroscience." *Nature Neuroscience*, 1–14.
  https://doi.org/10.1038/s41593-024-01649-9.

Greenberg, Paul E, Andree-Anne Fournier, Tammy Sisitsky, Crystal T Pike, and Ronald C
  Kessler. 2015. "The Economic Burden of Adults With Major Depressive Disorder in the
  United States (2005 and 2010)." *The Journal of Clinical Psychiatry* 76 (02): 155–62.
  https://doi.org/10.4088/jcp.14m09298.

Gschwind, Tilo, Ayman Zeine, Ivan Raikov, Jeffrey E. Markowitz, Winthrop F. Gillis, Sylwia
  Felong, Lori L. Isom, Sandeep Robert Datta, and Ivan Soltesz. 2023. "Hidden Behavioral
  Fingerprints in Epilepsy." *Neuron* 111 (9): 1440-1452.e5.
  https://doi.org/10.1016/j.neuron.2023.02.003.

Hsu, Alexander I., and Eric A. Yttri. 2021. "B-SOiD, an Open-Source Unsupervised Algorithm
  for Identification and Fast Prediction of Behaviors." *Nature Communications* 12 (1): 5188.
  https://doi.org/10.1038/s41467-021-25420-x.

Insel, Thomas, Bruce Cuthbert, Marjorie Garvey, Robert Heinssen, Daniel S. Pine, Kevin Quinn,
  Charles Sanislow, and Philip Wang. 2010. "Research Domain Criteria (RDoC): Toward a
  New Classification Framework for Research on Mental Disorders." *American Journal of
  Psychiatry* 167 (7): 748–51. https://doi.org/10.1176/appi.ajp.2010.09091379.

Jhumka, Z. Anissa, and Ishmail J. Abdus-Saboor. 2022. "Next Generation Behavioral
  Sequencing for Advancing Pain Quantification." *Current Opinion in Neurobiology*
  76:102598. https://doi.org/10.1016/j.conb.2022.102598.

Jia, Jianping, Cuibai Wei, Shuoqi Chen, Fangyu Li, Yi Tang, Wei Qin, Lina Zhao, et al. 2018.
  "The Cost of Alzheimer's Disease in China and Re-Estimation of Costs Worldwide."
  Alzheimer's & Dementia 14 (4): 483–91. https://doi.org/10.1016/j.jalz.2017.12.006.

Lauer, Jessy, Mu Zhou, Shaokai Ye, William Menegas, Steffen Schneider, Tanmay Nath,
  Mohammed Mostafizur Rahman, et al. 2022. "Multi-Animal Pose Estimation, Identification
  and Tracking with DeepLabCut." *Nature Methods* 19 (4): 496–504.
  https://doi.org/10.1038/s41592-022-01443-0.

Lin, Sherry, Winthrop F. Gillis, Caleb Weinreb, Ayman Zeine, Samuel C. Jones, Emma M. Robinson, Jeffrey Markowitz, and Sandeep Robert Datta. 2024. "Characterizing the Structure of Mouse Behavior Using Motion Sequencing." *Nature Protocols* 19 (11): 3242–91. https://doi.org/10.1038/s41596-024-01015-w.

Luo, Fei, Jun-yang Deng, Xuan Sun, Jian Zhen, and Xiao-dan Luo. 2023. "Anterior Cingulate Cortex Orexin Signaling Mediates Early-Life Stress-Induced Social Impairment in Females." *Proceedings of the National Academy of Sciences* 120 (20): e2220353120. https://doi.org/10.1073/pnas.2220353120.

Luxem, Kevin, Petra Mocellin, Falko Fuhrmann, Johannes Kürsch, Stephanie R. Miller, Jorge J. Palop, Stefan Remy, and Pavol Bauer. 2022. "Identifying Behavioral Structure from Deep Variational Embeddings of Animal Motion." *Communications Biology* 5 (1): 1267. https://doi.org/10.1038/s42003-022-04080-7.

Mathis, Alexander, Pranav Mamidanna, Kevin M. Cury, Taiga Abe, Venkatesh N. Murthy, Mackenzie Weygandt Mathis, and Matthias Bethge. 2018. "DeepLabCut: Markerless Pose Estimation of User-Defined Body Parts with Deep Learning." *Nature Neuroscience* 21 (9): 1281–89. https://doi.org/10.1038/s41593-018-0209-y.

Nadler, J J, S S Moy, G Dold, D Trang, N Simmons, A Perez, N B Young, et al. 2004. "Automated Apparatus for Quantitation of Social Approach Behaviors in Mice." *Genes, Brain and Behavior* 3 (5): 303–14. https://doi.org/10.1111/j.1601-183x.2004.00071.x.

Negre, Christian F. A., Uriel N. Morzan, Heidi P. Hendrickson, Rhitankar Pal, George P. Lisi, J. Patrick Loria, Ivan Rivalta, Junming Ho, and Victor S. Batista. 2018. "Eigenvector Centrality for Characterization of Protein Allosteric Pathways." *Proceedings of the National Academy of Sciences* 115 (52): E12201–8. https://doi.org/10.1073/pnas.1810452115.

Peleh, Tatiana, Xuesheng Bai, Martien J.H. Kas, and Bastian Hengerer. 2019. "RFID-Supported Video Tracking for Automated Analysis of Social Behaviour in Groups of Mice." *Journal of Neuroscience Methods* 325:108323. https://doi.org/10.1016/j.jneumeth.2019.108323.

Peleh, Tatiana, Kevin G.O. Ike, Emma J. Wams, Evan P. Lebois, and Bastian Hengerer. 2019. "The Reverse Translation of a Quantitative Neuropsychiatric Framework into Preclinical Studies: Focus on Social Interaction and Behavior." *Neuroscience & Biobehavioral Reviews* 97:96–111. https://doi.org/10.1016/j.neubiorev.2018.07.018.

Pereira, Talmo D., Diego E. Aldarondo, Lindsay Willmore, Mikhail Kislin, Samuel S.-H. Wang, Mala Murthy, and Joshua W. Shaevitz. 2019. "Fast Animal Pose Estimation Using Deep Neural Networks." *Nature Methods* 16 (1): 117–25. https://doi.org/10.1038/s41592-018-0234-5.

Pereira, Talmo D., Nathaniel Tabris, Arie Matsliah, David M. Turner, Junyu Li, Shruthi Ravindranath, Eleni S. Papadoyannis, et al. 2022. "SLEAP: A Deep Learning System for

Multi-Animal Pose Tracking." *Nature Methods* 19 (4): 486–95. https://doi.org/10.1038/s41592-022-01426-1.

Porcelli, Stefano, Nic Van Der Wee, Steven van der Werff, Moji Aghajani, Jeffrey C. Glennon, Sabrina van Heukelum, Floriana Mogavero, et al. 2019. "Social Brain, Social Dysfunction and Social Withdrawal." *Neuroscience & Biobehavioral Reviews* 97:10–33. https://doi.org/10.1016/j.neubiorev.2018.09.012.

Ricceri, Laura, Anna Moles, and Jacqueline Crawley. 2007. "Behavioral Phenotyping of Mouse Models of Neurodevelopmental Disorders: Relevant Social Behavior Patterns across the Life Span." *Behavioural Brain Research* 176 (1): 40–52. https://doi.org/10.1016/j.bbr.2006.08.024.

Romero-Ferrero, Francisco, Mattia G. Bergomi, Robert C. Hinz, Francisco J. H. Heras, and Gonzalo G. de Polavieja. 2019. "Idtracker.Ai: Tracking All Individuals in Small or Large Collectives of Unmarked Animals." *Nature Methods* 16 (2): 179–82. https://doi.org/10.1038/s41592-018-0295-5.

Sangiamo, Daniel T., Megan R. Warren, and Joshua P. Neunuebel. 2020. "Ultrasonic Signals Associated with Different Types of Social Behavior of Mice." *Nature Neuroscience* 23 (3): 411–22. https://doi.org/10.1038/s41593-020-0584-z.

Sharma, Sunita, Sharlene Rakoczy, and Holly Brown-Borg. 2010. "Assessment of Spatial Memory in Mice." *Life Sciences* 87 (17–18): 521–36. https://doi.org/10.1016/j.lfs.2010.09.004.

Shemesh, Yair, Yehezkel Sztainberg, Oren Forkosh, Tamar Shlapobersky, Alon Chen, and Elad Schneidman. 2013. "High-Order Social Interactions in Groups of Mice." *ELife* 2:e00759. https://doi.org/10.7554/elife.00759.

Weinreb, Caleb, Mohammed Abdal Monium Osman, Libby Zhang, Sherry Lin, Jonah Pearl, Sidharth Annapragada, Eli Conlin, et al. 2023. "Keypoint-MoSeq: Parsing Behavior by Linking Point Tracking to Pose Dynamics." *BioRxiv*, 2023.03.16.532307. https://doi.org/10.1101/2023.03.16.532307.

Weinreb, Caleb, Jonah E. Pearl, Sherry Lin, Mohammed Abdal Monium Osman, Libby Zhang, Sidharth Annapragada, Eli Conlin, et al. 2024. "Keypoint-MoSeq: Parsing Behavior by Linking Point Tracking to Pose Dynamics." *Nature Methods* 21 (7): 1329–39. https://doi.org/10.1038/s41592-024-02318-2.

Wiltschko, Alexander B., Tatsuya Tsukahara, Ayman Zeine, Rockwell Anyoha, Winthrop F. Gillis, Jeffrey E. Markowitz, Ralph E. Peterson, Jesse Katon, Matthew J. Johnson, and Sandeep Robert Datta. 2020. "Revealing the Structure of Pharmacobehavioral Space through Motion Sequencing." *Nature Neuroscience* 23 (11): 1433–43. https://doi.org/10.1038/s41593-020-00706-3.

Wiltschko, Alexander B., Matthew J. Johnson, Giuliano Iurilli, Ralph E. Peterson, Jesse M. Katon, Stan L. Pashkovski, Victoria E. Abraira, Ryan P. Adams, and Sandeep Robert Datta. 2015. "Mapping Sub-Second Structure in Mouse Behavior." *Neuron* 88 (6): 1121–35. https://doi.org/10.1016/j.neuron.2015.11.031.

Ziegler, Lukas M von, Fabienne K Roessler, Oliver Sturman, Rebecca Waag, Mattia Privitera, Sian N Duss, Eoin C O'Connor, and Johannes Bohacek. 2023. "Analysis of Behavioral Flow Resolves Latent Phenotypes." https://doi.org/10.1101/2023.07.27.550778.

Ziegler, Lukas von, Oliver Sturman, and Johannes Bohacek. 2021. "Big Behavior: Challenges and Opportunities in a New Era of Deep Behavior Profiling." *Neuropsychopharmacology* 46 (1): 33–44. https://doi.org/10.1038/s41386-020-0751-7.