Social Context Restructures Behavioral Syntax in Mice

Marti Ritter[1,2]*, Hope Shipley[2,3], Serena Deiana[2], Bastian Hengerer[2], Carsten T. Wotjak[2], Michael
Brecht[1], Amarender R. Bogadhi[2]

[1]Department of Biology, Faculty of Life Sciences, Humboldt University, Berlin, Germany

[2]Neuroscience and Mental Health Discovery Research, Boehringer Ingelheim Pharma GmbH & Co. KG,
Biberach, Germany

[3]University of Manchester, Manchester, United Kingdom

* Correspondence:

Marti Ritter (Research Group of Michael Brecht)

Philippstraße 13, House 6
10115 Berlin
Germany
Tel: + 49-(0)30-20936770

E-mail: ritermaq@hu-berlin.de

Keywords: mouse, social behavior, social withdrawal, context, behavioral syntax, behavioral
decomposition, behavioral repertoire

Abstract

The study of social behavior in mice has grown increasingly relevant for unraveling associated brain
circuits and advancing the development of treatments for psychiatric symptoms involving social
withdrawal or social anxiety. However, a data-driven understanding of behavior and its modulation in
solitary and social contexts is lacking. In this study, we employed motion sequencing ("MoSeq") to
decompose mouse behaviors into discrete units ("Syllables") and investigate whether—and how—the
behavioral repertoire differs between solitary and dyadic (social) settings.

Our results reveal that social context significantly modulates a minority (25%) of syllables, containing
predominantly stationary and undirected behaviors. Notably, these changes are associated with spatial
proximity to another mouse rather than active social contact. Interestingly, a network analysis of syllable
transitions shows that context-sensitive syllables exhibit altered network influence, independent of the
number of connected syllables, suggesting a regulatory role. Furthermore, syllable composition changes
significantly during social contact events with two distinct sequence families governing approach and
withdrawal behaviors. However, no unique syllable sequences mapped to specific social interactions.

Overall, our findings suggest that a subset of syllables drives contextual behavioral adaptation in mice,
potentially facilitating transitions within the broader behavioral repertoire. This highlights the utility of
MoSeq in dissecting nuanced, context-dependent behavioral dynamics.

1   Introduction

Social interactions are some of the most complex processes across all basic functions required for survival, leading to a high level of specialization of the involved structures and a high cost of dysfunction (Cacioppo et al. 2014; Porcelli et al. 2019). At the same time this high level of complexity leads to an increased vulnerability, e.g. to pathogens or disorders, an example being the occurrence of social withdrawal as a shared feature across major depressive disorder (MDD), Alzheimer's disease (AD), and schizophrenia (Porcelli et al. 2019). These conditions have a large impact on quality of life and carry a high cost to both affected individual (Chevance et al. 2020) and society, leading to an urgent need for the discovery of treatments (Greenberg et al. 2015; Jia et al. 2018). After previous approaches showed limited success, perspective on translational neuroscience has shifted towards a focus on psychopathology rooted in observable behavior and neurobiological measures (Research Domain Criteria, RDoC), suggesting that focusing on social impairments in complex disorders might lead to an earlier detection and better treatment outcomes (Garner 2014; Casey et al. 2013; Cuthbert 2018; Insel et al. 2010).

To facilitate early and effective identification of promising candidate treatments, pre-clinical research makes extensive use of rodent models of disorders found in humans (Bolivar, Walters, and Phoenix 2007). This approach can also be used in the search for treatments of social impairments (Luo et al. 2023). Mice possess a complex social behavior, influenced by age, kinship, and sex, among other factors. They utilize ultrasonic vocalizations, pheromones, and body contact as modalities, to cite some (Sangiamo, Warren, and Neunuebel 2020; Ricceri, Moles, and Crawley 2007). Recent developments in translational research led to refocusing from constrained tasks at shorter timescales, such as the three-chamber social approach task (Nadler et al. 2004) or tube test to determine the social hierarchy between mice (Fan et al. 2019), towards an open arena layout to enable the mice to show more complex behavioral structures and social interactions (Peleh, Ike, et al. 2019; Shemesh et al. 2013; Forkosh et al. 2019). But the complexity of social behavior also required agreed-upon definitions for specific interaction motifs to achieve quantifiable and reproducible descriptions.

At its inception, traditional ethological research was limited to qualitative, higher-level descriptions to characterize complex behaviors (Anderson and Perona 2014). This required a reduced set of dimensions to make behaviors observable (Gomez-Marin et al. 2014), such as focusing on vocalizations or Infra-red beam breaks. Even though some of the earliest descriptions of mouse behavior contained "ethograms" matching complex and specific behaviors (Abeelen 1964), it was not possible at that time to capture the full complexity of behavior into an ethogram in a data-driven, quantifiable manner. As this limited the possibilities of the neuroscience community to investigate unrestrained, naturalistic behavior, an urgent goal was the development of appropriate tools (Datta et al. 2019). The advent of convolutional neural networks and marker-less pose tracking tools, such as DeepLabCut (Lauer et al. 2022) or SLEAP (Pereira et al. 2022), allowed for the automatic extraction of key body points capturing much of the pose information available in video recordings collected by one or more cameras (Ebbesen and Froemke 2022), for single (Pereira et al. 2019; Mathis et al. 2018) or multiple animals (Romero-Ferrero et al. 2019; Bewley et al. 2016). In parallel, other approaches to behavior tracking and decomposition arose (Wiltschko et al. 2015), which utilized the recent advances in machine learning and computation (Dennis et al. 2021). The behavioral decomposition approach, which allows identification of recurring short segments of behavior ('syllables') that form the basis of complex behavior, was effective in characterizing behavioral fingerprints of drugs in disorders, such as epilepsy (Gschwind et al. 2023), using both three

dimensional recordings of single mice (Wiltschko et al. 2020) and utilizing body point representations of
single or multiple mice (Hsu and Yttri 2021; Luxem et al. 2022).

A multitude of tools that focus on automated tracking and annotation of social interactions in mice can
be found in recent literature (Peleh, Bai, et al. 2019; Chaumont et al. 2012; Giancardo et al. 2013),
including open-source recording setups (Chaumont et al. 2018). Despite advances towards an integration
of ethological and comparative psychological approaches (Bordes et al. 2023) most of these approaches
are based on parametric definitions of specific behaviors, as in supervised tools (Goodwin et al. 2024),
and do not leverage potential advantages of unsupervised behavior classification or standardized
benchmarking (Choi and Kumar 2024). For example, use of motion sequencing in pain quantification
(Jhumka and Abdus-Saboor 2022) and evaluation of behavioral effects of analgesics (Bohic et al. 2023)
revealed unique insights into the difference between baseline and analgesia induced behaviors, as well
as common global markers of ongoing pain, such as specific pause and grooming modules.

To investigate if an unsupervised behavioral decomposition approach yields insights into the structure of
social behavior in mice, we applied Keypoint-MoSeq to videos of mice in dyadic and solitary contexts. By
comparing syllable frequency and structure in dyadic and solitary contexts, we show a significant
modulation in a subset of stationary, undirected syllables that are not fully dependent on contact with a
conspecific. Instead, these modulated syllables play a key role in transitions between syllables as the
behavior becomes more diverse in a dyadic context. Furthermore, social behaviors (e.g. social approach
or leave) defined through parametric thresholds can be represented in syllable space. However, we did
not find syllables or syllable sequences that distinctly map onto social behaviors. Our results suggest that
syllables or syllable sequences are sensitive markers for the changes in behavior in the presence of a
conspecific but may not correspond to specific "social" behaviors occurring in this context.

2   Materials and Methods

2.1   Animals

Twenty female C57BL/6J mice (~6 weeks old at time of first experiment) were obtained from Charles River Germany. They were housed in groups of two to three, with the other cage-mates not used in this study and kept in a light- and humidity-controlled facility. All animals were orally injected with a vehicle treatment intraperitoneally 20 minutes before their recordings started to account for adverse effects caused by the oral gavage procedure. All procedures followed the regulations for animal experimentation enforced by the local district administration's animal welfare commissioner of the state of Baden-Württemberg.

2.2   Open Field Arena Setup

The experiments were run in a 453mm x 453mm x 400mm (Width x Depth x Height) square arena made from clear plexiglass. The arena was equipped with dimmable LED strips and an analog camera (768px x 576px, 25Hz). All sides except for the front were enclosed by a wooden frame, with the front being covered with a red plexiglass door, and the floor being removable to facilitate cleaning. During the tests a stable illumination of around 220lux was maintained.

2.3   Experimental procedure

Half of the mice (n=10) were kept on a regular 12h light-dark cycle, with the other 10 animals being kept in a reversed 12h light-dark cycle. This applied to both the solitary as well as the dyadic experiments, with the dyadic experiments occurring between mice with the same light cycle. All experiments began between 8 and 10 am. The timeline of experiments and other procedures is shown below.

2.3.1 Solitary open field test

Mice were orally injected with Hydroxypropyl-β-cyclodextrin 20% McIIVaine and kept in their home cages for 20 minutes. The oral gavage was performed to consider the impact of treatment stress for future studies on pharmacological treatment. After 20 minutes had passed, mice were transferred into the recording setups and video-recorded for one hour. The recordings were started automatically through Stoelting Any-Maze software. Of the one hour of solitary recordings, only the first 20 minutes were used for behavioral analysis in this study, to maintain data parity to the dyadic context. The timeline for a single solitary recording is shown below.

2.3.2 Dyadic social interation test

Five weeks after the solitary test recordings, mice were tested in a dyadic social interaction test. Two mice from separate cages were orally injected Hydroxypropyl-β-cyclodextrin 20% McIIVaine at the same time and housed in their home cage for 20 minutes before starting the dyadic social interaction test. Mice were allowed to freely interact and explore the open field arena for 20 minutes. To allow for discrimination between the two mice in each recording, one mouse was shaved eight to nine days before the recording date at a constant location at the upper back behind the neck. The timeline for a single dyadic recording is shown below.

2.4   Data collection and analysis

2.4.1 Stoelting Any-Maze

Stoelting Any-Maze version 7.4 was used to record the videos used in this analysis and to record metadata. No further tracking parameters were extracted from Anymaze for this study. We utilized the built-in scoring utility to collect experimenter-scored active and passive contact epochs in the video data.

2.4.2 SLEAP

SLEAP (Pereira et al. 2022) version 1.3.3 was used to track 12 key points on the body of each mouse in the solitary and dyadic video recordings. 4 of the key points were located on the head of the mouse, 4 along the spine, another 2 were located one each at the hips, and 2 further points were labelled along the tail. For the model, we selected a bottom-up design with 800 frames taken from dyadic context and 600 frames taken from solitary context for the training set.

2.4.3 Keypoint-MoSeq

Keypoint-MoSeq (Weinreb et al. 2024) version 0.4.1 was used to extract the behavioral syllables from the
keypoints extracted in the previous step. We used 10 of the 12 body points tracked in SLEAP (excluding
the tail) to fit the model. The model parameter Kappa was selected iteratively such that the resulting
syllable length distribution had its median at 10 frames, representing roughly 400ms of behavior. The
default threshold proposed in the analysis tools provided by the Keypoint-MoSeq Python package was
set to 0.5% of total onset proportion and we followed that threshold here.

Using the tools provided in the Keypoint-MoSeq package, we named all 32 syllables with a bout
proportion above 0.5% of the global distribution. Names and descriptions were chosen before further
analysis according to the behaviors shown in the majority of the 24 extracted example videos for each
syllable. For a full list and descriptions, see Table 1.

Table 1. *Table of syllables.* Labels and descriptions were determined by using the tools provided with the
Keypoint-MoSeq package to create a 24-video overview of the occurrences of each syllable. Behaviors
were described according to the behavior shown in most of the videos (see fractions in description
column). Labelling was performed prior to the analysis, and minor edits were done for readability.

Syllable	Label	Short description
0	dart_forward	20/24 videos show relatively quick movement from stationary position forward, generally, if possible straight ahead no turns
1	turn_left	17/24 show left turn, 8/17 related to down from rearing… due to 2D? Usually stationary
2	turn_right_move	16/24 show right turn, 7/16 related to (down from, sometimes aborted) rearing, same as 1, maybe due to 2D? Compared to 1 this one has after the turn some movement
3	turn_right_stationary	17/24 show right turn, this time mainly stationary, 11/17 related to moving up into rearing.
4	unclear_contraction	8/24 show down from rearing, generally seems to have some rightward bias, but not specific
5	movement_forward	22/24 show general, constant movement forward, usually average velocity
6	sniffing	23/24 show sniffing, 20/23 related to the corners and edges of the box
7	rearing	19/24 show movement related to rearing, or "waving" during rearing, mounting… seems to detect more "last" phase of rearing shortly before or during down
8	turn_right_move2	19/24 usually first light right turn then moving forwards, generally long
9	turn_left_short	21/24 short left turn, sometimes movement afterwards
10	sniffing2	19/24 sniffing again
11	turn_right_stationary2	20/24 right turn on spot
12	head_raise	22/24 showing some form of raising head, either on floor from grooming/sniffing up or often related to rearing (14/22)
13	unclear_sniffing	19/24 some stationary sniffing, but not specific
14	head_raise_left	19/24 raising head, leftwards bias, sometimes weak (17/19)

15	turn_left_with_retract	19/24 show left turn, most often after short retraction, happens often during down from rearing (leftwards) (14/19), but not specific
16	turn_right	23/24 show right turn, basically the rightwards version of 15, also with down from rearing often (16/23)
17	rearing_right	18/24 show some form of rearing/mounting, mainly rightwards (14/18)
18	unclear_sniffing2	18/24 show some form of sniffing on floor or in the air… maybe righward bias, but unclear head raise/down
19	unclear_stop	17/24 show some form of stopping from movement
20	rearing_start	17/24 show start of rearing some of the rest shows rearing related (down or currently rearing)
21	rearing_left_unclear	16/24 show some movement turning leftwards during a rear
22	sniffing_left	19/24 show some sniffing at walls while turning their head slightly left and moving sometimes forwards
23	rear_down	17/24 show down from rearing
24	unclear	No clear behavior
25	sniffing_right	21/24 show some sniffing, usually against walls, while turning head slightly right… rightwards version of 22
26	sniffing_left_turn	16/24 appear to be sniffing while turning slightly left
27	turn_right_stationary3	20/24 show some turning right in various contexts
28	turn_right_stationary4	19/24 show right turn without much other movement
29	movement_unclear	Appears to be a general movement syllable, maybe slight right turn, but not very specific
30	rearing_sniffing	17/24 show sniffing while on hindlegs, sometimes with movement forward
31	stop_with_head_down	16/24 not very clear, but appears to be rather abrupt stop with a short head down movement

2.4.4 Body center tracking

To explore any differences in mouse behavior, body center tracking was used to calculate the distance moved by a single mouse (Figure 1B), its position (Figure 4A), or inter-mice distance (Figure 4B-E, G). For tracking of body center, we used the mouse centroid calculated in Keypoint-MoSeq by taking the median of all used body points (10 points on the mouse body, excluding the tail). We then converted this pixel-based measure to a distance-based measure using the pixel per millimeter resolution in our arenas, calculated based on the known distance between the corners of the arena and the same distance in pixel measured in the video (measure extracted through the Python package OpenCV2, version 4.11.0; using the selectROI function).

2.4.5 Trajectory analysis

We extracted the syllable trajectories from the tracking information collected across all mice. To evaluate similarity between a pair of syllable trajectories, we calculated the cosine distance between position vectors of the pair of syllables, using the implementation in SciPy (version 1.14.0). Cosine distance is a measure of distance between two vectors (u and v) calculated as $\mathbf{d_{cos}} = \mathbf{1} - \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{u}\|_2 \|\mathbf{v}\|_2}$. The cosine

distance between all pairs is used for hierarchical clustering of syllables in a dendrogram (Figure 2A, Suppl. Figure 3A). The data shown in Figure 2 is recorded from the vehicle treated animals used in this study. As this study was part of a larger experiment, 40 additional mice of the same characteristics were recorded in parallel. When we used the larger dataset and looked at the hierarchical clustering of syllables in a dendrogram (Suppl. Figure 3A), we found 6 out of 8 dyadic-modulated syllables being separated into a distinct arm of the dendrogram.

The dendrograms shown here were created based on the output of the functions `linkage` and `dendrogram` contained in the SciPy package (Version 1.14.0) for Python. The exemplar trajectories for each syllable (suppl. Figure 1A) were selected by the same algorithm as implemented in `get_typical_trajectories` in the Keypoint-MoSeq package for Python.

2.4.6 Eigenvector centrality

To identify syllable nodes that may play transitory role in a directed network of syllable transitions, we calculated eigenvector centrality measures for each syllable in the transition network of syllables in solitary and dyadic context (Figure 2B). Eigenvector centrality measure allows for the detection of nodes that are influential in the network, but not necessarily connected to many nodes themselves. The eigenvector centrality is defined by the formula: $\mathbf{x_v} = \frac{1}{\lambda}\sum_{\mathbf{t}\in\mathbf{M(v)}}\mathbf{x_t}$, with $\mathbf{M(v)}$ being the set of neighbors of node $\mathbf{v}$. To calculate eigenvector centrality, we used the `eigenvector_centrality` function implemented in the NetworkX Python package (version 3.3) on the normalized transition probabilities (outgoing transitions sum to 1 for a given source syllable) for each animal in solitary and dyadic contexts. Overall, for a given syllable, we obtained 20 eigenvector centrality measures in each of solitary and dyadic contexts (Figure 2B).

2.4.7 Social contact scoring

The experimenter was trained in the recognition of social behaviors and the usage of the built-in video scoring functionality of Any-Maze and was blind to the light cycle and treatment of mice. Since this study was part of a larger series, the scorer also scored further videos whose data is not shown here. The scorer classified two main types of behaviors: *active social interaction*, that is mice engaged in directed interactions, such as touching of the conspecific's body with the nose and sniffing behavior; *passive social interaction* was identified as body contacts without active sniffing. If there was uncertainty about how to score a specific segment, the scorer was instructed to mark the segment as both states. The uncertain phases can occur, for example, at the transition between active and passive contact, as the exact moment of transition can be unclear. Around 35% of active contacts and 10% of passive contacts were overlapping in this way. This overlap can also be explained by the input delay on the keyboard used for scoring, and the fact that continuous active contacts do not occur often and last only a few moments, while passive interactions can last multiple seconds. Since we aimed to evaluate the time course of change in syllable and syntax composition during active and passive social contact (see Figure 3B, C), we pooled bouts with less than 6 seconds duration of the same contact type. This allowed for the application of a 5 second (~125 frames) pre-contact control window, as shown in Figure 3B and C. We used these pre-contact control windows to ensure that there was a clean onset at the observed bouts, without other bouts of the evaluated behavior occurring in the control window.

2.4.8 Kullback-Leibler Divergence

To measure differences between distributions of syllables (Figure 3B) and syntax (Figure 4F), we applied the Kullback-Leibler divergence (D_{KL}), or relative entropy. The D_{KL} measures the distance between an observed probability distribution and an expected probability distribution. It is defined as $\mathbf{D_{KL}(P||Q)} = \sum_{\mathbf{x} \in \mathbf{X}} \mathbf{P(x)} \log\left(\frac{\mathbf{P(x)}}{\mathbf{Q(x)}}\right)$, with P being the observed and Q being the expected distribution. In this study we used a base of 2 for the logarithm, leading to a relative entropy in bits. To calculate the Kullback-Leibler divergence, we used the implementation contained in the `entropy` function in SciPy (passing the distribution to be compared as pk and the reference (expected) distribution as qk).

2.4.9 Parametric Behavior Classes

To gain a separation of behavioral syntax into parametrically defined behaviors, we used two parameters - inter-mouse distance (IMD) and change in distance (CID) as shown in Figure 4C. The peaks from kernel density plot in Figure 4C were separated into corresponding contact related class (syntax starting and/or ending below an IMD < 100mm) and control class (marked with hatched lines). Movement classes (approach and leave) were defined with an absolute CID > 50mm and a contact threshold of 100mm based on the preferred IMD shown in Figure 4B. The relative movement threshold (absolute CID) of 50mm was chosen as half of contact threshold. These two thresholds along with limitations of the arena (maximum IMD and maximum CID ≤ diagonal of arena) resulted in the polygons defining approach and leave behavior classes.

Control classes were defined as those that did not lead to inter-mouse contact but matched the same movement parameters (see dashed arrows in Figure 4C). For example, the control for approach behavior was any movement that had an CID < -50mm and led to a final IMD between 100mm and 200mm. The leave control class was defined as CID > 50mm and a final IMD between 100mm and 200mm. Finally, the contact control class used a larger distance criterion and captured the largest peak in the IMD-CID distribution, at an IMD > 200mm and an absolute CID < 50mm.

2.4.10 Hamming Distance

The hamming distance between two sequences is the number of positions that are different in the two sequences. This only includes substitutions, but not rearrangements. In this study, we defined the syntax family as a set of syntax with hamming distance < 1 relative to the name-giving syntax – for the syntax family (9,0,5), the syntax (9,0,10) would be a member but not (0,5,10). This was used to aggregate the diverse, but highly similar set of syntax influencing inter-mouse approaches and leaves as shown in Figure 4D. Further analysis of these families can be found in Figure 4E.

The formula for two sequences of identical length could be written as $d_{Hamming} = \sum_{i=1}^{n_x} eq(x_i, y_i)$, with x and y representing the two sequences, and n_x being the length of sequence x.

2.4.11 Principal Component Analysis (PCA)

We used dimensionality reduction with PCA to evaluate if DM syllables have a larger contribution to the variability across videos (keypoint tracks) of mice in solitary and dyadic contexts. We used syllable frame proportions for each individual mice, concatenated across animals, in solitary and dyadic contexts as input to the PCA. Since syllable frequencies summed up to 1, we did not apply further scaling to individual frequencies. The loadings for the first 15 PCs are split for DM or DU syllables (Figure 5A).

264 Similarly, we also evaluated the contribution of DM syntax to the variability across videos (keypoint
265 tracks) of mice in solitary and dyadic contexts (Figure 5B). We used the implementation of PCA in the
266 scikit-learn package (version 1.5.1).

2.5 Statistical tests

268 All statistical tests were performed in Python (version 3.11.9). Unless stated otherwise we used a
269 significance threshold of α=0.05 and corrected for multiple testing by applying the Bonferroni correction
270 where applicable.

2.5.1 Mann-Whitney U-Test

We used the Mann Whitney U-test in the following analysis:

• In Figure 1B, we tested if the total distance moved differed between solitary and dyadic
recordings. Applying a one-sided test showed a significant difference between both groups
($p<0.0001$).
• To find DM syllables, we applied a two-sided test in Figure 1C.
• As mentioned in the section on the eigenvector centrality measure, we tested if eigenvector
centrality differed between DM and DU syllables, as shown in Figure 2B. We applied a two-sided
test on the eigenvector centrality for DU and DM syllables, extracted per recording and
compared between solitary and dyadic recordings. 5 out of 8 DM syllables showed a significantly
different eigenvector centrality ($p≤0.05$) between solitary and dyadic recordings. No DU syllables
showed a significant difference.
• Based on the results shown in Figure 2C, we tested for significantly modulated transitions
between solitary and dyadic contexts. We applied a two-sided test to the transition counts
observed in dyadic and solitary context for each transition observed in both contexts. After
correction for multiple testing, we found 9 significantly modulated transitions with 8 of those
targeting DM syllables.
• We also applied a one-sided test on the absolute value of the PCA loadings shown in Figure 5A, B
to verify whether DM syllables or syntax provided a greater influence on the first 5 PCs
compared to DU syllables or syntax. The tests showed that DM syllables had significantly higher
loadings than DU syllables ($p<0.002$), but that the same effect was not visible in DM syntax when
compared to DU syntax.
• The data shown in Suppl. Figure 1B was tested with a two-sided test and we did not find
significant differences between dyadic groups and the corresponding solitary onset proportions
(after correction for multiple comparisons).
• For each parameter as shown in Suppl. Figure 1C, we performed a one-sided test between DM
syllable trajectories and DU syllable trajectories and found the two direction-based parameters
to be significantly reduced for the DM syllables (after correction for multiple comparisons).
• A two-sided test was applied for the pairwise comparisons of the data shown in Suppl. Figure 2B.
We found that all the comparisons were significant (after correction for multiple comparisons).
• The comparison between the count of unique syllable transitions observed in dyadic context and
solitary context yielded a significant result when we applied a one-sided test.

We used SciPy's (version 1.14.0) implementation of the Mann-Whitney U-Test (function
`mannwhitneyu`) with significance levels set at α=0.05.

2.5.2 χ² Contingency Test

The χ² contingency test is used to test for independence of frequencies of two groups summarized in a
contingency table. To verify which DM syllables showed a significant effect of experimenter-scored
contact in Figure 1D (left panel), we applied this test to the onset counts of each syllable during and
outside contact compared to the onset counts of any other syllable during and outside contact. The test
revealed two syllables (7 and 18) with significant effects (p≤0.05, Bonferroni correction). In Suppl. Figure
6A, we applied the χ² contingency test to the onset counts of each syllable family within or outside the
behavior class, compared to the onset counts of all length 3 syntax within or outside the same behavior
class. We found all comparisons to be significantly different, except for the comparison between syntax
family (11,2,8) to all other syntax in the leave control behavior class. We used the implementation
(`chi2_contingency` function) contained in the SciPy package for Python.

2.5.3 Two-way ANOVA

In Figure 1D, we compared the onset proportion of each DM syllable between solitary and dyadic
contexts. Within dyadic context, we tested for an interaction between onset proportions of DM syllables
and contact between conspecifics. Both contact (either experimenter-scored or based on distance
quartiles; for distance distribution, see Figure 4B) and DM syllable were categorical variables. We
performed a two-way ANOVA (formula `proportion ~ C(syllable) * C(contact)`) with the help
of the `anova_lm` and `ols` functions contained in the statsmodels package (version 0.14.2) for Python.
Results showed significant main effects on proportion and interactions between the categorical
dependent variables (see Results).

2.5.4 Bonferroni correction

We corrected for multiple comparisons by using the Bonferroni correction. For example, we used the
following equation to adapt the significance threshold for the 32 syllables to maintain our desired overall
α=0.05: $\alpha_{\mathbf{Bonferroni}} = \alpha_{\mathbf{uncorrected}}/\mathbf{n_{tests}}$ with $\mathbf{n_{tests}}$ corresponding to the number of syllables. This
correction was applied to the results of the χ² contingency test used in Figure 1D ($\mathbf{n_{tests}}$=8); Mann
Whitney U-tests in Figure 2B ($\mathbf{n_{tests}}$ =32), Figure 2C ($\mathbf{n_{tests}}$ =99), Suppl. Figure 1B ($\mathbf{n_{tests}}$ =6), Suppl. Figure 1C
($\mathbf{n_{tests}}$ =3), Suppl. Figure 2B ($\mathbf{n_{tests}}$ =6); the χ² contingency test used in Suppl. Figure 6A **($\mathbf{n_{tests}}$ =12)**.

2.5.5 Benjamini-Hochberg correction

333 To control for the false discovery rate in the Mann-Whitney U-test used to find the 8 DM syllables (see
334 Figure 1C), the implementation of the Benjamini-Hochberg procedure provided by the SciPy package was
335 applied. We used this in place of the Bonferroni correction as the visualizations provided in the Keypoint-
336 MoSeq package also used the Benjamini-Hochberg correction.

2.5.6 Z-Score representation of syllable and syntax during contact

To identify whether the experimenter-scored contact behaviors had a significantly different
representation in syllable space when compared to the global average of behavior, we computed z-score
for each of the syllable or syntax occurring during contact behaviors using the distribution of the same

syllable or syntax in the predefined baseline window (Figure 3B, C). We calculated the z-score for data
shown in Figure 3B and C using the following formula: $z = \frac{x - \mu}{\sigma}$.

In Figure 3B, a baseline period was defined between 5 seconds and 2 seconds prior to the scored
contact. The z-score of each trace in figure 3B was calculated based on the distribution of values for the
same trace within the baseline window. In addition, we also calculated z-score for the active and passive
contact traces based on the entire randomized trace (labeled "randomized" in the legend). We evaluated
both z-scores framewise for every trace and defined frames as significant where both z-scores lay above
3. The frames found to be significant are marked for each trace in the upper subplot.

In Figure 3C, we again defined a baseline period between 5 seconds and 2 seconds prior to the scored
contact. We computed z-score for each trace (syntax) based on the distribution of values for the same
trace within the baseline window. We again defined a z-score above 3 as significant. All traces shown in
black (active, n=25 and passive, n=24) showed significance across at least 50% of the test windows
starting from 2 seconds prior to 2 seconds after the scored contact. The remaining traces are shown in
gray (n_{active}=79, $n_{passive}$=80). Four syntax traces with the highest z-score were marked as the top 4 syntax
associated with each contact type and are shown in color.

3   Results

We ran two open field tests (OFT) with video recordings, separated by 5 weeks, on 20 group housed female C57BL/6J mice. The first OFT recording contained an individual mouse in the open arena (referred to as solitary context), and the second OFT recording, performed after 5 weeks, contained two mice from separate cages in the same arena (referred to as dyadic context). In preparation of future pharmacological studies, all mice were orally injected with a vehicle treatment 20 minutes before the start of the recordings to account for potential adverse effects caused by the oral gavage procedure. Considering potential light cycle effects on behavior, one half of the mice were housed under normal, and the second half were housed under reversed light cycle conditions. However, we could not identify significant differences in syllable expression between the two light cycle groups and hence the data were pooled from the two light cycle groups.

We analyzed 20 minutes of video containing a single mouse in solitary and dyadic contexts using SLEAP (Pereira et al. 2022) to track a set of 12 key body points on each mouse. We fitted a Keypoint-MoSeq model (Weinreb et al. 2023) to a subset of 10 key body points to extract repeated elements of mice behavior, referred to as behavioral syllables (see Methods; Figure 1A). The output syllables were aggregated by either their onset (bout) proportion, representing the proportion of syllable bouts regardless of individual bout lengths in frames, or frame proportion, representing the total proportion of the recorded frames in one syllable. After filtering for total onset proportion, we extracted a total of 32 syllables which were used in subsequent analyses (see Methods; Suppl. Figure 1A). Syllables with a bout proportion below the threshold often occurred in clusters (Suppl. Figure 2A) and were observed to relate mostly to self-grooming (Suppl. Figure 2C, D). These clusters occurred in both solitary and dyadic contexts but did not meaningfully differ between the two contexts (Suppl. Figure 2B).

3.1   Dyadic context modulates a small subset of MoSeq identified syllables

To investigate the structure of social behavior through behavior decomposition, we first asked if behavioral 'syllables' identified through decomposition (Keypoint-MoSeq) are sensitive to modulation in solitary and dyadic contexts and whether this modulation is dependent on physical proximity between the two animals. Before comparing syllables in solitary and dyadic contexts, we verified whether parameters related to movement revealed significant changes in mouse behavior between solitary and dyadic contexts (L. M. von Ziegler et al. 2023). By approximating the output of traditional body center tracking (see Methods), we detected a significant reduction in distance moved of mice between solitary and dyadic contexts (Mann-Whitney U test, p≈1.83e-5; Figure 1B).

A comparison of onset proportion of all 32 syllables in solitary and dyadic contexts revealed a subset of 8 syllables that differed significantly between contexts (Figure 1C; see Methods; Mann-Whitney U-Test, Benjamini-Hochberg correction, p≤0.05). From here onwards, the 8 significantly modulated syllables will be referred to as dyadic-modulated (DM) syllables and the remaining 24 syllables will be referred to as dyadic-unmodulated (DU) syllables.

Figure 1. *Identification of behavioral syllables in solitary and dyadic contexts.* (A) Panels left to right: C57BL/6J mice were placed into an open field arena, 20 minutes after an oral vehicle injection, in solitary or dyadic context. After video recording in the open field arena, Key points on the body and tail were tracked using SLEAP; key points on the body were used to extract behavioral syllables using Keypoint-MoSeq (see Methods). (B) A comparison of the distances moved in solitary and dyadic context recordings. Asterisk indicates statistical significance (Mann-Whitney U-test, p<0.0001; see Results). (C) For each of the 32 syllables, a comparison of onset proportions between solitary and dyadic context (Mann-Whitney U-Test, Benjamini-Hochberg correction, p<0.05) reveals 8 significantly modulated (DM; orange colored syllable numbers) and 24 unmodulated syllables (DU). (D) Left panel: A comparison of DM syllables' onset proportion based on its occurrence during or outside of an active contact behavior as scored by an experimenter (see Methods). We performed a χ2 proportion test for each syllable's onset proportion. 2 syllables (7 and 18) show a significant (Bonferroni correction, p<0.05) difference during and outside of an active contact in the dyadic context. Right panel: A comparison of DM syllables' onset proportion based on the distance between the mice, divided into quartiles, during the syllable occurrence (see Methods). (E) Trajectories of the 8 DM syllables belonging to stationary (7, 12, 13, 18, 23, 30) or non-directional movement (all). (F) A selection of DU syllable trajectories with directed (0, 2, 9, 11) and traversal movement (0, 2, 8, 9, 22). In panels C & D, filled circles and error bars indicate mean and 95% CI respectively. Mouse (Ethan and Kravitz 2020) and syringe (Bates 2021) vector graphics adapted from SciDraw.

Because the mice move less in dyadic context (Figure 1B), one may deduce that DM syllables are likely
associated with active contact but may not be influenced by physical separation between mice outside of
an active contact. To evaluate the effect of physical proximity on DM syllables, we compared the onset
proportions of syllables during experimenter-scored active contact (for definition see Figure 3A) to no
contact. Results from a two-way ANOVA revealed a significant main effect for DM syllables and active
contact ($F_{(7, 299)}=49.61$, $p<1e-45$; $F_{(1, 299)}=5.07$, $p=0.03$) with a significant interaction between DM
syllables and active contact ($F_{(7, 299)}=2.21$, $p=0.03$). However, a closer inspection for each syllable
revealed few significant syllables (2 out of 8, χ^2 contingency test, Bonferroni correction, $p<0.05$). In
addition, we checked if the distance between mice is associated with the modulation of the DM syllables
by comparing onset frequency in different distance quartile. A two-way ANOVA revealed a significant
main effect for DM syllables and distance quartiles ($F_{(7, 603)}=93.62$, $p<1e-91$; $F_{(3, 603)}=3.87$, $p=0.01$)
with a significant interaction between DM syllables and distance quartile ($F_{(21, 603)}=8.67$, $p<1e-22$).
These findings reveal that majority of the DM syllables are not directly linked to an active contact but are
associated with physical proximity to a conspecific (Figure 1D, left panel; but also Suppl. Figure 1B).

An inspection of the trajectories of all DM syllables reveals high proportion of stationary, and non-
directional behaviors, for example rearing as in syllable 7 (Figure 1E). In contrast, most directional
behavior (left or right oriented movement as in syllable 9 and 11) falls into the DU group (Figure 1F and
Suppl. Figure 1C; one-sided Mann-Whitney U-test, Bonferroni correction, $p<0.05$). To quantify the
differences within and across DM and DU syllables, we used cosine distance to quantify similarity (see
Methods; Figure 2A). The dendrogram of similarity measure did not reveal any systematic clustering
within and across DM and DU syllables. However, using an extended dataset (see Methods), we found
that most of the DM syllables are similar when compared to DU syllables (Suppl. Figure 3A).

These results show that a subset of behavioral 'syllables' identified through decomposition are sensitive
to modulation in solitary and dyadic contexts. Surprisingly, most of the modulated syllables, dominated
by stationary undirected behaviors, are not linked to an active contact but are associated with physical
proximity to a conspecific outside of an active contact.

3.2   Modulated syllables drive the syllable transitions in dyadic context

Because there was no association of an active contact with most behavioral syllables in the presence of a
conspecific (dyadic context), we next investigated whether there is a distinct role for DM syllables in the
transitions between behavioral syllables to form a complex social behavior.

We employed Eigenvector centrality to quantify the role of DM syllables in the transitions between
syllables (see Methods). Briefly, Eigenvector centrality is a measure of how well a node is connected to
nodes which themselves are well connected to individual nodes within a network (Negre et al. 2018).
Here, the Eigenvector centrality of a syllable is calculated based on the directed graph of transition
probabilities. We found a significant change in eigenvector centrality, during the dyadic context, for most
of the DM syllables but not for any of the DU syllables (Figure 2B; two-sided Mann-Whitney U-test,

Bonferroni correction, p≤0.05). This finding reveals a crucial transitory role for DM syllables in the
network of syllables defining complex social behavior in the presence of a conspecific (dyadic context).

Figure 2. ***Role of dyadic modulated syllables in the network of syllable transitions during solitary and dyadic contexts.*** (A) The dendrogram of inter-syllable cosine distances shows limited clustering of DM syllables (orange colored syllable numbers). (B) A comparison of eigenvector centrality measure between solitary and dyadic contexts (see Methods), based on the network of transition probabilities (outgoing probabilities were normalized to sum 1) between syllables, reveals significant differences for DM syllables (5 out of 8) but not for DU syllables. Asterisk indicates statistical significance (two-sided Mann-Whitney-U test, Bonferroni correction, p<0.05). Filled circles and error bars indicate mean and 95% CI. (C) A directed network of syllable transitions modulated by dyadic context with edges indicating transition probabilities between syllable nodes. Edge colors indicate negative (blue) or positive (red) modulation of the transition probability in dyadic context; edge thickness indicates amount of change in transition probability in dyadic context. Syllables are nodes in the network and node diameter corresponds to changes in eigenvector centrality of that node. Syllable nodes with a significant change in eigenvector centrality are indicated in bold syllable numbers (see panel B).

During the dyadic context, most of the significantly modulated transitions (8/9, two-sided Mann-
Whitney U-test, Bonferroni correction, p≤0.05) target DM syllables as reflected by eigenvector centrality
(see arrows pointing to the first column with DM syllables in Figure 2C). In addition, we observed an
overall increase in syllable transitions (Suppl. Figure 3B; one-sided Mann-Whitney U Test, p≤0.05) and
the strongest changes in transition probabilities (thick lines in Figure 2C) were mostly a result of reduced
transition probabilities (blue lines), originating in DU syllables (see Suppl. Figure 3C).

Due to a weak association between contact and DM syllables (Figure 1D), the exact role of these syllables
in social interactions is not clear. However, we observed that behavior in general becomes less
stereotypical (or structured) when mice are in a dyadic context (see Suppl. Figure 3D). Regardless, these

results reveal an important role for DM syllables as "transfer nodes" in the behavioral program to
construct a complex social behavior in the presence of a conspecific (dyadic context).

3.3   Changes in syllable composition reflect active and passive contact behaviors

Given that DM syllables play an important role of 'transfer node' in the behavioral syllable transitions to
form complex behavior in dyadic context, we investigated whether complex behaviors, such as active or
passive contact behaviors, are captured through changes in the composition of syllables or sequences of
syllables, referred to as syntax in the following observations.

For this purpose, we tasked a fellow researcher to score active and passive social contacts in the dyadic
recordings (see Methods). To quantify changes in syllable composition, in an individual mouse during
active and passive behaviors, we computed Kullback-Leibler divergence (D_{KL}), a measure of statistical
distance between two distributions, between framewise distribution of syllable proportions in active or
passive contacts to an overall distribution of syllable proportions in dyadic context (see Methods). The
time course of D_{KL} reflects changes in syllable composition aligned to start of active or passive behaviors
(Left panel in Figure 3B). The results show a significant change in syllable composition (D_{KL}), in an
individual mouse, at the time of active and passive behaviors. We verified that this change is fully
attributable to active and passive behaviors since the same analysis on randomized sequence of syllables
did not show any change in syllable composition (Figure 3B; z-score see Methods).

We next asked if active and passive behaviors also reflect changes in joint syllable composition of two
mice by computing framewise Kullback-Leibler divergence (D_{KL}) between joint distributions of syllable
proportions between two mice during active or passive contacts to an overall joint distribution of syllable
proportions in dyadic context. The time course of D_{KL} shows a weak but continuous and significant
change in joint syllable composition aligned to start of active behavior but less so for passive behavior
(Right panel in Figure 3B).

To investigate if sequence of syllables (syntax) signal active and passive contact behaviors, we extracted
length 3 syntax (i.e., sequences comprising 3 syllables), as this corresponded to about a second of
behavior with a median syllable duration of 10 frames (see Methods). A multitude of syntax are
significantly associated with active (25 syntax) and passive (24 syntax) contact behaviors (Figure 3C; see
Methods for statistics). However, there was little overlap of syntax between active and passive contact
behaviors (Figure 3C, 8 significantly associated syntax shared between active and passive). In addition,
we observed few DM syllables in the top 4 most significantly associated syntax for both contact
behaviors. Furthermore, inspection of example trajectories for the two syntax most associated with
active contact suggests a behavior related to circling around a conspecific (Suppl. Figure 4A, B). Similarly,

the syntax most associated with passive contact included mostly stationary behavior, while the second
syntax was related to a turning behavior followed by movement (Suppl. Figure 4C, D).

Figure 3. *Syllable and syntax composition during experimenter scored active and passive contact behaviors.* (A) Schematic of an experimenter scoring active and passive contact between mice in the dyadic video recordings. For details on scoring instructions, see Methods. (B) Left-bottom panel: Time course of difference in frame-wise syllable composition of an individual mice, quantified as DKL, between experimenter-scored behavior and all behaviors is plotted for active and passive behaviors. Left-Top panel: Time course of statistical significance for data in bottom panel compared to shaded baseline window. Right panel: Same as Left panel but using frame-wise syllable composition of two mice instead of a single mouse (see Methods). (C) The frequency of length 3 syllable syntax was aligned to the start of a scored behavior and transformed to z-score using the baseline window (from 100ms to 50ms of the start; shaded window). Time course of syntax aligned to the start of scored behavior is shown for active contact (left panel) and passive contact (right panel). The top 4 syntax are indicated with colored lines and their corresponding syllable composition is shown in legend. Orange colored syllable number indicates DM syllable.

481   These results show that experimenter defined active and passive contact behaviors are represented in
482   the syllable space and there are a diverse set of syllable sequences that are modulated in contact
483   behaviors. Furthermore, DM syllables are underrepresented in the most of the contact-associated
484   syllable sequences, in line with the previously observed modulation of DM syllables regardless of contact
485   (Figure 1D), suggesting their role in aiding transitions between syntax associated with contact behaviors.

3.4 Individual syllable sequences do not map onto specific behaviors

The finding that experimenter defined contact behaviors are represented in syllable and syntax space prompted us to ask whether social behaviors, such as social approach and leave defined in a parametric space (Peleh, Bai, et al. 2019), can be mapped onto syllable and syntax space. First, we verified if there is a propensity for targeted behavior in our dataset by calculating distributions of mice positions (Figure 4A) and probability of inter-mouse distance (Figure 4B). Our data shows that mice preferred to position themselves in corners of arena (Figure 4A), and that the co-location of two mice in the same corner was a coordinated event, rather than a random occurrence (Figure 4B).

Before parametrically defining behaviors, we extracted inter-mouse distance during length 3 syntax and mapped them onto a space representing the inter-mouse distance at the start of syntax and change in distance by the end of the syntax (Figure 4C). From the density plot of all syntax in this space (blue contours in Figure 4C), we were able to identify four density zones that specify parametrically defined behaviors – targeted approach and leave, as well as stationary behaviors in low distance (contact) and stationary behaviors with high distance. We further defined control zones as those that showed the same relative movements but did not relate to contact between the mice (see dotted arrows and hatched areas in Figure 4C & Methods). We choose to use syntax rather than syllables since syntax timescales were more apt for the description of targeted long-distance movements and longer stationary behaviors.

For the two distinct behaviors of social approach and leave (Figure 4C), we investigated whether there is a dominant representation by specific syntax. We tracked the change in distance over a rolling 30 frame window, matching our definition of length 3 syntax. It is evident that nearly all the windows in which inter-mouse distance changed by more than 200mm were fully represented by either one of two syntax families (see Figure 4D). A syntax family is defined as a set of syntax with at most one changed syllable relative to a reference syntax (the family was named after the reference syntax; see Hamming Distance in Methods). We found that social approach and leave were fully represented by two syntax families (syntax 9,0,5 and syntax 11,2,8; see Figure 4D) and both contained a syllable with a long-distance traversal trajectory (syllables 9&0 and 2 respectively, see Figure 1F), suggesting both syntax families capture behavior either targeted towards or away from a conspecific. Inspection of sample trajectories included in these syntax families revealed that one syntax family, (9,0,5), was usually associated with a leftwards turn followed by a movement straight ahead while the other syntax family, (11,2,8), was associated with a rightwards turn followed by movement (Suppl. Figure 5A, B). We interpret these syntaxes as corresponding to a mouse reorienting towards or away from a conspecific, during an approach and leave respectively.

We asked if the above finding that social approach and leave were fully represented by two syntax families extended in the opposite direction i.e. whether the two syntax families specifically map onto the social approach and leave behaviors. When we mapped all occurrences of the two syntax families onto parameter space (Figure 4E), we did not identify a direct mapping between a syntax family and a specific behavior class as defined in our parametric space (Suppl. Figure 6A; χ^2 contingency test, Bonferroni correction, $p \leq 0.05$, but see Methods). Since we could not identify an association between a syntax family and a behavior, we checked if syllable composition differed between behavior classes. We calculated D_{KL} on the syllables contained within each syntax, between a pair of behavior classes. We observed no separation between approach and leave classes further confirming that there is not a specific syntax, at

least in our dataset, that directly maps onto approach and leave classes. Furthermore, we noted a

modest separation between all movement classes (approach and leave) and stationary (contact) classes

respectively (Figure 4F, for analysis of syntax see Suppl. Figure 6B). However, mapping all occurrences of

syntax associated with active contact (from Figure 3C) onto our parameter space did not reveal any

specific syntax associated with active contact behavior (Figure 4G). We also confirmed that there was no

specific syntax associated with passive contact behavior (Suppl. Figure 6C).

Figure 4. *Syntax and parametric behavior classes.* (A) 2D kernel density estimate of mouse positions sampled from the recordings in dyadic context (2 million data points). Marginal plots show isolated kernel density estimate of each axis. (B) Histograms of inter-mouse distances in observed data, shuffled and random control data. Observed refers to inter-mouse distances in recorded data; shuffled refers to inter-mouse distances with the individual positions being shuffled; Random refers to inter-mouse distances simulated with randomly selected positions within the arena (10 million point-pairs). The difference of the observed and shuffled histogram is shown in black. (C) Parametric behavior classes were derived from observed length 3 syntax. Inter-mouse distance (x-axis) and change of that distance (y-axis) were plotted as a kernel density estimate shown as blue contours (KDE, see methods). Based on clusters in the KDE, we defined 3 behavior classes: contact (x<100mm, x+y<100mm), approach (x>100mm, x+y<100mm), and leave (x<100mm, x+y>100mm). Control classes were defined to match change in distance (y-axis) but excluding contact scenarios (see dotted arrows). (D) A rolling window (length 30 frames) was used to calculate change in inter-mouse distance (x-axis) and is plotted against the proportion of the window spent in one of the two syntax family (9,0,5) or (11,2,8) (see Methods). Line with shaded region represents median and 95% CI. (E) KDE with same parameters as in C but limited to one of the two syntax families referred to in panel D. The KDE from panel C was subtracted as baseline. Red areas represent higher density than in C, blue areas with reduced relative density. (F) Heatmap of DKL between syllable distributions found in syntax belonging to the behavior classes defined in panel C. Darker colors indicate lower DKL. (G) Top 4 syntax correlated with experimenter-scored active contact, taken from Figure 3C, plotted as scatter on top of the KDE taken from panel C.

These findings show that while mapping of a behavior to syllable space is possible with a one-to-many
mapping, we could not find evidence for distinct syllable sequences that map onto specific behaviors.

3.5   Syllables allow discrimination of solitary and dyadic behavioral modes

Based on the lack of evidence for specific syntax or syllable descriptors of social behavior, we aimed to
find out if syllables or syntax provide discrimination of solitary and dyadic contexts. First, we evaluated
the variance captured by syllables and syntax using principal component analysis on the proportion of
frames containing each syllable for each mouse (see Methods). We separated the contributions of dyadic
modulated and unmodulated syllables to each of the principal components (Figure 5A). Results reveal
that DM syllables capture significantly more variance compared to DU syllables – first five PCs explaining
~73% variance have significantly higher DM syllable loadings (one-sided Mann Whitney U-test, p<0.005;
see Figure 5A). However, we did not find such a difference for syntax (same test, Figure 5B).

Furthermore, using the above identified principal components, we observed a clear separation between
solitary and dyadic contexts, while the same approach did not lead to a separation between two light
cycle conditions (Figure 5C). These results demonstrate that syllables and syntax, particularly DM

syllables, provide a sensitive description of behavior to discriminate if behavior is in the presence or
absence of a conspecific.

Figure 5. *Syllables and syntax discriminate solitary and dyadic context but not normal and reverse light-cycle.* (A) Results from principal component analysis (PCA) of frame-wise occurrences of modulated (DM) and unmodulated (DU) syllables in dyadic and solitary recordings are shown (see Methods). The box plots show (with y-axis on the left) the absolute of the loadings on each component for DM and DU syllables. The y-axis on the right shows the percentage variance explained by each of the component. The first 5 PCs explain 74% of the variance and DM syllables contribute significantly more to the loadings of these PCs than DU syllables (one-sided Mann-Whitney U-test, Bonferroni correction, p<0.05). (B) Results from PCA on syntax proportions are shown with same conventions as in panel A. DM syntax was defined as containing at least 2 DM syllables. (C) Scatter plot of the first 2 PCs from syllable-based PCA (top plots) or syntax-based PCA (bottom plots). Colors separate tracks of individual mice in either solitary/dyadic recordings (left plots) or normal/reversed light cycles (right plots).

4   Discussion and conclusion

A growing number of tools are available to accomplish the data-driven segmentation of the full behavioral repertoire of mice (See reviews: Choi & Kumar, 2024; Datta et al., 2019; L. von Ziegler, Sturman, & Bohacek, 2021). However, it is not known if such an approach, to segment or decompose the behavior into its recurring elements, yields insights into the structure of social behavior in mice. Here, we investigated the structure of social behavior using Keypoint-MoSeq (Weinreb et al. 2024) to videos of mice in a dyadic and solitary context. We found that in the presence of a conspecific, mice moved less and significantly modulated the usage of a small subset of mostly stationary and undirected syllables, referred to as dyadic modulated (DM) syllables (Figure 1B, C; Suppl. Figure 1A, C). Surprisingly, for most of the DM syllables, this modulation in usage was not associated with the occurrence of experimenter-scored contact between the mice (Figure 1D). Instead, DM syllables played a significant role as transitory nodes in the syllable transition network (Figure 2B, C). Importantly, we found significant changes in composition of syllables and syntax (length 3 syllable sequence) aligned to experimenter-defined contact behaviors (Figure 3B, C). In addition, we identified 2 syntax families that captured approach and leave behavior as defined through parametric thresholds (Figure 4C, D; Suppl. Figure 5A, B). However, we did not find evidence for distinct syntax that map onto specific complex behaviors. Overall, our results show that behavioral syllables can be used as descriptive and sensitive markers to detect changes in behavior in the presence of a conspecific, but specific social behaviors may not map onto individual behavioral syllables or a specific syntax thereof.

A possible caveat of our study could be that the experiments were performed with one set of mice across both contexts. We ran the two recordings (solitary, dyadic) in the same set of open field arenas 5 weeks apart. It is possible that during the second recording the arena may not have been a novel environment for mice. Based on available literature the long-term spatial memory of the arena will be negligible after a span of 5 weeks, as most assessments of spatial memory work with periods of up to 2 weeks after training session over multiple days (Sharma, Rakoczy, and Brown-Borg 2010). Similarly, movement patterns of our mice could potentially also evolve during inter-experimental phase (5 weeks) and might have contributed to our findings, but it can be expected that social contact with a conspecific outweighs these effects. The estrous cycle most likely varies across individuals and recordings and might contribute to social preference (Chari et al. 2020). However, as solitary and dyadic context behaviors were clearly distinguishable in our dataset, we conclude that the estrous cycle-related variance must be below the level of behavioral changes in a dyadic context.

To detect as much of the behavioral repertoire as possible the Keypoint-MoSeq model was fitted to the entire recorded dataset, including full one hour solitary recordings and 20-minute dyadic recordings. Potentially this 3:1 disbalanced dataset could introduce a bias towards behaviors shown in solitary context, but as we treated the solitary state as the baseline, effects of the dyadic context would still be visible in the following analysis. In addition, we limited the analysis of the extracted syllables to the first 20 minutes of all recordings to restore data parity and analyze segments of identical activity (see Methods). Distinct behaviors occurring only in a dyadic context would have been visible with this approach, like behavioral syllables related to self-grooming (see Suppl. Figure 2C, D). While some distinct behaviors (self-grooming and jumping) were captured as individual syllables as expected, these occurred too rarely to be included in our analysis (see Methods).

It was possible to detect contact-specific changes in the behavioral syllable composition for individual
mice (Figure 3B, left panel), but we could not observe the same prominent effect for the joint syllable
composition across two mice (Figure 3B, right panel). This difference in results between individual mice
and two mice is very likely because joint occurrences of syllables are sparse, and a higher number of
joint occurrences are necessary when analyzing the joint behavior of two mice. Since the significant
effect of contact occurrence on the syllable distribution in individual mice is clear in our dataset, we
decided to show the across-mice effect as a comparison to encourage further studies. Perhaps, an
approach to decompose multi-animal behavior, from large datasets to overcome sparse social events,
instead of decomposing individual animal behavior could be beneficial in detecting 'social' syllables.

Social interactions capture rich and multi-modal behavioral repertoire (Porcelli et al. 2019) and thus
need a data-driven extraction of behaviors. Recent studies addressed this need by using either
parametric methods (Peleh, Bai, et al. 2019) or machine learning applications (Goodwin et al. 2024; Lin
et al. 2024). It is also necessary to aim for explainable readouts from data-driven analysis of complex
behaviors and a fruitful approach is to limit analysis to a small set of well-defined parametric
approximations. We applied this perspective in our analysis (Figure 4C) and defined a set of four
parametric categories in the context of mouse interaction (close contact, approach, leave, other
stationary behaviors). Our original expectation with the behavioral decomposition approach was to find
distinct syllables or sequence of syllables (syntax) that represent the parametrically defined categories.
While we did find syntax families that capture most occurrences of social approach and leave (Figure 4C,
D), a direct and distinct mapping of behavioral syllables onto social behavior classes was not observable
(Figure 4E, G). It is possible that behavioral decomposition based on the key points may have contributed
to the lack of distinct mapping of behavioral syllables onto social behaviors in our dataset because
extraction of key points inherently limits the amount of pose information unlike pose extracted from
depth recordings (Lin et al. 2024). In addition, a fruitful approach could be to use the joint pose
information between mice, and this requires overcoming the limitation of sparseness in social
interaction datasets.

In summary, our results show that, in the presence of a conspecific, mice possess a subset of behavioral
syllables that were significantly modulated, and these modulated syllables change their transitory hub-
like role in the transition network of all syllables. While we did not find bidirectional mapping between a
syllable or syntax and specific social behavior, our results show that syllables or syntax are sensitive
descriptors to detect changes in behavior because of change in social context (Figure 5C). This finding
taken together with previous studies, showing the sensitivity of behavioral syllables to detect different
pharmacological interventions (Wiltschko et al. 2015; 2020), suggests that manipulation of experimental
context (social or non-social enrichment) may offer to be a valuable dimension in phenotypic screening
of drugs using behavioral decomposition methods.

634 Table of abbreviations

Abbreviation	Full Term
DM	Dyadic Modulated
DU	Dyadic Unmodulated
PCA	Principal Component Analysis
IMD	Inter-Mouse Distance
CID	Change In Distance
RDoC	Research Domain Criteria
MoSeq	Motion Sequencing
LED	Light Emitting Diode

635 Acknowledgements

636 We want to thank the other members of the data science team for their feedback on this project. We
637 also thank Alina Ritter for her support during the recording of data, and Michael Winter, Zoi Balla,
638 Johannes Freudenreich, and Carmen Weiss for their help using software related to the recording of data.

639 Data availability statement

640 The original contributions presented in the study are publicly available. This data can be found here:
641 https://doi.org/10.5281/zenodo.15237564

642 Code availability statement

643 The code used to generate the data shown in this study is publicly available. It can be found here:
644 https://github.com/Marti-Ritter/social-context-restructures-behavioral-syntax-in-mice

645 Ethics statement

646 All procedures followed the regulations for animal experimentation enforced by the local district
647 administration's animal welfare commissioner of the state of Baden-Württemberg.

649 **Author Contributions**

	Conceptualization	Data curation	Formal analysis	Funding acquisition	Investigation	Methodology	Project administration	Resources	Software	Supervision	Validation	Visualization	Writing – original draft	Writing – review & editing
MR	x	x	x		x	x			x		x	x	x	x
HS		x	x		x						x	x		x
SD	x					x	x	x		x				x
BH	x									x				x
CW	x			x		x	x			x				x
MB	x									x				x
AB	x					x	x			x			x	x

Funding

The authors declare that financial support was received for the research, authorship, and/or publication
of this article. This work was supported by Boehringer Ingelheim Pharma GmbH & Co. KG, Biberach,
Germany.

Conflict of interest

Marti Ritter, Hope Shipley, Serena Deiana, Bastian Hengerer, Carsten T. Wotjak, and Amarender R.
Bogadhi received salaries from Boehringer Ingelheim Pharma GmbH & Co. KG.

Generative AI statement

The authors declare that no Generative AI was used in the creation of this manuscript.

References

Abeelen, J. H. F. van. 1964. "Mouse Mutants Studied by Means of Ethological Methods." *Genetica* 34 (1): 79–94. https://doi.org/10.1007/bf01664181.

Anderson, David J., and Pietro Perona. 2014. "Toward a Science of Computational Ethology." *Neuron* 84 (1): 18–31. https://doi.org/10.1016/j.neuron.2014.09.005.

Bates, Alexander. 2021. "Syringe." Zenodo. https://doi.org/10.5281/zenodo.4421220.

Bewley, Alex, Zongyuan Ge, Lionel Ott, Fabio Ramos, and Ben Upcroft. 2016. "Simple Online and Realtime Tracking." *2016 IEEE International Conference on Image Processing (ICIP)*, 3464–68. https://doi.org/10.1109/icip.2016.7533003.

Bohic, Manon, Luke A. Pattison, Z. Anissa Jhumka, Heather Rossi, Joshua K. Thackray, Matthew Ricci, Nahom Mossazghi, et al. 2023. "Mapping the Neuroethological Signatures of Pain, Analgesia, and Recovery in Mice." *Neuron* 111 (18): 2811-2830.e8. https://doi.org/10.1016/j.neuron.2023.06.008.

Bolivar, Valerie J., Samantha R. Walters, and Jennifer L. Phoenix. 2007. "Assessing Autism-like Behavior in Mice: Variations in Social Interactions among Inbred Strains." *Behavioural Brain Research* 176 (1): 21–26. https://doi.org/10.1016/j.bbr.2006.09.007.

Bordes, Joeri, Lucas Miranda, Bertram Müller-Myhsok, and Mathias V. Schmidt. 2023. "Advancing Social Behavioral Neuroscience by Integrating Ethology and Comparative Psychology Methods through Machine Learning." *Neuroscience & Biobehavioral Reviews* 151:105243. https://doi.org/10.1016/j.neubiorev.2023.105243.

Cacioppo, John T., Stephanie Cacioppo, Stephanie Dulawa, and Abraham A. Palmer. 2014. "Social Neuroscience and Its Potential Contribution to Psychiatry." *World Psychiatry* 13 (2): 131–39. https://doi.org/10.1002/wps.20118.

Casey, B. J., Nick Craddock, Bruce N. Cuthbert, Steven E. Hyman, Francis S. Lee, and Kerry J. Ressler. 2013. "DSM-5 and RDoC: Progress in Psychiatry Research?" *Nature Reviews Neuroscience* 14 (11): 810–14. https://doi.org/10.1038/nrn3621.

Chari, Trishala, Sophie Griswold, Nick A. Andrews, and Michela Fagiolini. 2020. "The Stage of the Estrus Cycle Is Critical for Interpretation of Female Mouse Social Interaction Behavior." *Frontiers in Behavioral Neuroscience* 14:113. https://doi.org/10.3389/fnbeh.2020.00113.

Chaumont, Fabrice de, Renata Dos-Santos Coura, Pierre Serreau, Arnaud Cressant, Jonathan Chabout, Sylvie Granon, and Jean-Christophe Olivo-Marin. 2012. "Computerized Video Analysis of Social Interactions in Mice." *Nature Methods* 9 (4): 410–17. https://doi.org/10.1038/nmeth.1924.

Chaumont, Fabrice de, Elodie Ey, Nicolas Torquet, Thibault Lagache, Stéphane Dallongeville, Albane
Imbert, Thierry Legou, et al. 2018. "Live Mouse Tracker: Real-Time Behavioral Analysis of Groups of
Mice." *BioRxiv*, 345132. https://doi.org/10.1101/345132.

Chevance, Astrid, Philippe Ravaud, Anneka Tomlinson, Catherine Le Berre, Birgit Teufer, Suzanne
Touboul, Eiko I Fried, Gerald Gartlehner, Andrea Cipriani, and Viet Thi Tran. 2020. "Identifying
Outcomes for Depression That Matter to Patients, Informal Caregivers, and Health-Care
Professionals: Qualitative Content Analysis of a Large International Online Survey." *The Lancet
Psychiatry* 7 (8): 692–702. https://doi.org/10.1016/s2215-0366(20)30191-7.

Choi, Jessica D., and Vivek Kumar. 2024. "A New Era in Quantification of Animal Social Behaviors."
Neuroscience & Biobehavioral Reviews 157:105528.
https://doi.org/10.1016/j.neubiorev.2023.105528.

Cuthbert, Bruce N. 2018. "The PRISM Project: Social Withdrawal from an RDoC Perspective."
Neuroscience & Biobehavioral Reviews 97:34–37. https://doi.org/10.1016/j.neubiorev.2018.08.006.

Datta, Sandeep Robert, David J. Anderson, Kristin Branson, Pietro Perona, and Andrew Leifer. 2019.
"Computational Neuroethology: A Call to Action." *Neuron* 104 (1): 11–24.
https://doi.org/10.1016/j.neuron.2019.09.038.

Dennis, Emily Jane, Ahmed El Hady, Angie Michaiel, Ann Clemens, Dougal R. Gowan Tervo, Jakob Voigts,
and Sandeep Robert Datta. 2021. "Systems Neuroscience of Natural Behaviors in Rodents." *The
Journal of Neuroscience* 41 (5): 911–19. https://doi.org/10.1523/jneurosci.1877-20.2020.

Ebbesen, Christian L., and Robert C. Froemke. 2022. "Automatic Mapping of Multiplexed Social
Receptive Fields by Deep Learning and GPU-Accelerated 3D Videography." *Nature Communications*
13 (1): 593. https://doi.org/10.1038/s41467-022-28153-7.

Ethan, Tyler, and Lex Kravitz. 2020. "Mouse." Zenodo. https://doi.org/10.5281/zenodo.3925901.

Fan, Zhengxiao, Hong Zhu, Tingting Zhou, Sheng Wang, Yan Wu, and Hailan Hu. 2019. "Using the Tube
Test to Measure Social Hierarchy in Mice." *Nature Protocols* 14 (3): 819–31.
https://doi.org/10.1038/s41596-018-0116-4.

Forkosh, Oren, Stoyo Karamihalev, Simone Roeh, Uri Alon, Sergey Anpilov, Chadi Touma, Markus
Nussbaumer, et al. 2019. "Identity Domains Capture Individual Differences from across the
Behavioral Repertoire." *Nature Neuroscience* 22 (12): 2023–28. https://doi.org/10.1038/s41593-019-
0516-y.

Garner, Joseph P. 2014. "The Significance of Meaning: Why Do Over 90% of Behavioral Neuroscience
Results Fail to Translate to Humans, and What Can We Do to Fix It?" *ILAR Journal* 55 (3): 438–56.
https://doi.org/10.1093/ilar/ilu047.

Giancardo, Luca, Diego Sona, Huiping Huang, Sara Sannino, Francesca Managò, Diego Scheggia,
Francesco Papaleo, and Vittorio Murino. 2013. "Automatic Visual Tracking and Social Behaviour

Analysis with Multiple Mice." *PLoS ONE* 8 (9): e74557.
https://doi.org/10.1371/journal.pone.0074557.

Gomez-Marin, Alex, Joseph J Paton, Adam R Kampff, Rui M Costa, and Zachary F Mainen. 2014. "Big
Behavioral Data: Psychology, Ethology and the Foundations of Neuroscience." *Nature Neuroscience*
17 (11): 1455–62. https://doi.org/10.1038/nn.3812.

Goodwin, Nastacia L., Jia J. Choong, Sophia Hwang, Kayla Pitts, Liana Bloom, Aasiya Islam, Yizhe Y. Zhang,
et al. 2024. "Simple Behavioral Analysis (SimBA) as a Platform for Explainable Machine Learning in
Behavioral Neuroscience." *Nature Neuroscience*, 1–14. https://doi.org/10.1038/s41593-024-01649-9.

Greenberg, Paul E, Andree-Anne Fournier, Tammy Sisitsky, Crystal T Pike, and Ronald C Kessler. 2015.
"The Economic Burden of Adults With Major Depressive Disorder in the United States (2005 and
2010)." *The Journal of Clinical Psychiatry* 76 (02): 155–62. https://doi.org/10.4088/jcp.14m09298.

Gschwind, Tilo, Ayman Zeine, Ivan Raikov, Jeffrey E. Markowitz, Winthrop F. Gillis, Sylwia Felong, Lori L.
Isom, Sandeep Robert Datta, and Ivan Soltesz. 2023. "Hidden Behavioral Fingerprints in Epilepsy."
Neuron 111 (9): 1440-1452.e5. https://doi.org/10.1016/j.neuron.2023.02.003.

Hsu, Alexander I., and Eric A. Yttri. 2021. "B-SOiD, an Open-Source Unsupervised Algorithm for
Identification and Fast Prediction of Behaviors." *Nature Communications* 12 (1): 5188.
https://doi.org/10.1038/s41467-021-25420-x.

Insel, Thomas, Bruce Cuthbert, Marjorie Garvey, Robert Heinssen, Daniel S. Pine, Kevin Quinn, Charles
Sanislow, and Philip Wang. 2010. "Research Domain Criteria (RDoC): Toward a New Classification
Framework for Research on Mental Disorders." *American Journal of Psychiatry* 167 (7): 748–51.
https://doi.org/10.1176/appi.ajp.2010.09091379.

Jhumka, Z. Anissa, and Ishmail J. Abdus-Saboor. 2022. "Next Generation Behavioral Sequencing for
Advancing Pain Quantification." *Current Opinion in Neurobiology* 76:102598.
https://doi.org/10.1016/j.conb.2022.102598.

Jia, Jianping, Cuibai Wei, Shuoqi Chen, Fangyu Li, Yi Tang, Wei Qin, Lina Zhao, et al. 2018. "The Cost of
Alzheimer's Disease in China and Re-Estimation of Costs Worldwide." *Alzheimer's & Dementia* 14 (4):
483–91. https://doi.org/10.1016/j.jalz.2017.12.006.

Lauer, Jessy, Mu Zhou, Shaokai Ye, William Menegas, Steffen Schneider, Tanmay Nath, Mohammed
Mostafizur Rahman, et al. 2022. "Multi-Animal Pose Estimation, Identification and Tracking with
DeepLabCut." *Nature Methods* 19 (4): 496–504. https://doi.org/10.1038/s41592-022-01443-0.

Lin, Sherry, Winthrop F. Gillis, Caleb Weinreb, Ayman Zeine, Samuel C. Jones, Emma M. Robinson,
Jeffrey Markowitz, and Sandeep Robert Datta. 2024. "Characterizing the Structure of Mouse
Behavior Using Motion Sequencing." *Nature Protocols* 19 (11): 3242–91.
https://doi.org/10.1038/s41596-024-01015-w.

Luo, Fei, Jun-yang Deng, Xuan Sun, Jian Zhen, and Xiao-dan Luo. 2023. "Anterior Cingulate Cortex Orexin
Signaling Mediates Early-Life Stress-Induced Social Impairment in Females." *Proceedings of the*
National Academy of Sciences 120 (20): e2220353120. https://doi.org/10.1073/pnas.2220353120.

Luxem, Kevin, Petra Mocellin, Falko Fuhrmann, Johannes Kürsch, Stephanie R. Miller, Jorge J. Palop,
Stefan Remy, and Pavol Bauer. 2022. "Identifying Behavioral Structure from Deep Variational
Embeddings of Animal Motion." *Communications Biology* 5 (1): 1267.
https://doi.org/10.1038/s42003-022-04080-7.

Mathis, Alexander, Pranav Mamidanna, Kevin M. Cury, Taiga Abe, Venkatesh N. Murthy, Mackenzie
Weygandt Mathis, and Matthias Bethge. 2018. "DeepLabCut: Markerless Pose Estimation of User-
Defined Body Parts with Deep Learning." *Nature Neuroscience* 21 (9): 1281–89.
https://doi.org/10.1038/s41593-018-0209-y.

Nadler, J J, S S Moy, G Dold, D Trang, N Simmons, A Perez, N B Young, et al. 2004. "Automated Apparatus
for Quantitation of Social Approach Behaviors in Mice." *Genes, Brain and Behavior* 3 (5): 303–14.
https://doi.org/10.1111/j.1601-183x.2004.00071.x.

Negre, Christian F. A., Uriel N. Morzan, Heidi P. Hendrickson, Rhitankar Pal, George P. Lisi, J. Patrick
Loria, Ivan Rivalta, Junming Ho, and Victor S. Batista. 2018. "Eigenvector Centrality for
Characterization of Protein Allosteric Pathways." *Proceedings of the National Academy of Sciences*
115 (52): E12201–8. https://doi.org/10.1073/pnas.1810452115.

Peleh, Tatiana, Xuesheng Bai, Martien J.H. Kas, and Bastian Hengerer. 2019. "RFID-Supported Video
Tracking for Automated Analysis of Social Behaviour in Groups of Mice." *Journal of Neuroscience*
Methods 325:108323. https://doi.org/10.1016/j.jneumeth.2019.108323.

Peleh, Tatiana, Kevin G.O. Ike, Emma J. Wams, Evan P. Lebois, and Bastian Hengerer. 2019. "The Reverse
Translation of a Quantitative Neuropsychiatric Framework into Preclinical Studies: Focus on Social
Interaction and Behavior." *Neuroscience & Biobehavioral Reviews* 97:96–111.
https://doi.org/10.1016/j.neubiorev.2018.07.018.

Pereira, Talmo D., Diego E. Aldarondo, Lindsay Willmore, Mikhail Kislin, Samuel S.-H. Wang, Mala
Murthy, and Joshua W. Shaevitz. 2019. "Fast Animal Pose Estimation Using Deep Neural Networks."
Nature Methods 16 (1): 117–25. https://doi.org/10.1038/s41592-018-0234-5.

Pereira, Talmo D., Nathaniel Tabris, Arie Matsliah, David M. Turner, Junyu Li, Shruthi Ravindranath, Eleni
S. Papadoyannis, et al. 2022. "SLEAP: A Deep Learning System for Multi-Animal Pose Tracking."
Nature Methods 19 (4): 486–95. https://doi.org/10.1038/s41592-022-01426-1.

Porcelli, Stefano, Nic Van Der Wee, Steven van der Werff, Moji Aghajani, Jeffrey C. Glennon, Sabrina van
Heukelum, Floriana Mogavero, et al. 2019. "Social Brain, Social Dysfunction and Social Withdrawal."
Neuroscience & Biobehavioral Reviews 97:10–33. https://doi.org/10.1016/j.neubiorev.2018.09.012.

Ricceri, Laura, Anna Moles, and Jacqueline Crawley. 2007. "Behavioral Phenotyping of Mouse Models of
Neurodevelopmental Disorders: Relevant Social Behavior Patterns across the Life Span." *Behavioural*
Brain Research 176 (1): 40–52. https://doi.org/10.1016/j.bbr.2006.08.024.

Romero-Ferrero, Francisco, Mattia G. Bergomi, Robert C. Hinz, Francisco J. H. Heras, and Gonzalo G. de
Polavieja. 2019. "Idtracker.Ai: Tracking All Individuals in Small or Large Collectives of Unmarked
Animals." *Nature Methods* 16 (2): 179–82. https://doi.org/10.1038/s41592-018-0295-5.

Sangiamo, Daniel T., Megan R. Warren, and Joshua P. Neunuebel. 2020. "Ultrasonic Signals Associated
with Different Types of Social Behavior of Mice." *Nature Neuroscience* 23 (3): 411–22.
https://doi.org/10.1038/s41593-020-0584-z.

Sharma, Sunita, Sharlene Rakoczy, and Holly Brown-Borg. 2010. "Assessment of Spatial Memory in
Mice." *Life Sciences* 87 (17–18): 521–36. https://doi.org/10.1016/j.lfs.2010.09.004.

Shemesh, Yair, Yehezkel Sztainberg, Oren Forkosh, Tamar Shlapobersky, Alon Chen, and Elad
Schneidman. 2013. "High-Order Social Interactions in Groups of Mice." *ELife* 2:e00759.
https://doi.org/10.7554/elife.00759.

Weinreb, Caleb, Mohammed Abdal Monium Osman, Libby Zhang, Sherry Lin, Jonah Pearl, Sidharth
Annapragada, Eli Conlin, et al. 2023. "Keypoint-MoSeq: Parsing Behavior by Linking Point Tracking to
Pose Dynamics." *BioRxiv*, 2023.03.16.532307. https://doi.org/10.1101/2023.03.16.532307.

Weinreb, Caleb, Jonah E. Pearl, Sherry Lin, Mohammed Abdal Monium Osman, Libby Zhang, Sidharth
Annapragada, Eli Conlin, et al. 2024. "Keypoint-MoSeq: Parsing Behavior by Linking Point Tracking to
Pose Dynamics." *Nature Methods* 21 (7): 1329–39. https://doi.org/10.1038/s41592-024-02318-2.

Wiltschko, Alexander B., Tatsuya Tsukahara, Ayman Zeine, Rockwell Anyoha, Winthrop F. Gillis, Jeffrey E.
Markowitz, Ralph E. Peterson, Jesse Katon, Matthew J. Johnson, and Sandeep Robert Datta. 2020.
"Revealing the Structure of Pharmacobehavioral Space through Motion Sequencing." *Nature
Neuroscience* 23 (11): 1433–43. https://doi.org/10.1038/s41593-020-00706-3.

Wiltschko, Alexander B., Matthew J. Johnson, Giuliano Iurilli, Ralph E. Peterson, Jesse M. Katon, Stan L.
Pashkovski, Victoria E. Abraira, Ryan P. Adams, and Sandeep Robert Datta. 2015. "Mapping Sub-
Second Structure in Mouse Behavior." *Neuron* 88 (6): 1121–35.
https://doi.org/10.1016/j.neuron.2015.11.031.

Ziegler, Lukas M von, Fabienne K Roessler, Oliver Sturman, Rebecca Waag, Mattia Privitera, Sian N Duss,
Eoin C O'Connor, and Johannes Bohacek. 2023. "Analysis of Behavioral Flow Resolves Latent
Phenotypes." https://doi.org/10.1101/2023.07.27.550778.

Ziegler, Lukas von, Oliver Sturman, and Johannes Bohacek. 2021. "Big Behavior: Challenges and
Opportunities in a New Era of Deep Behavior Profiling." *Neuropsychopharmacology* 46 (1): 33–44.
https://doi.org/10.1038/s41386-020-0751-7.