ZFT is the major iron and zinc transporter in *Toxoplasma gondii*

Dana Aghabi[1], Cecilia Gallego Rubio[1], Miguel Cortijo Martinez[1], Augustin Pouzache[1,] Erin J. Gibson[1,2], Lucas Pagura[1] and Clare R. Harding[1]

1. Centre for Parasitology, School of Infection and Immunity, University of Glasgow, Glasgow, UK
2. Wellcome-Wolfson Institute for Experimental Medicine, School of Medicine, Dentistry, and Biomedical Sciences, Queen's University Belfast, Belfast, UK

Abstract

Transition metals, such as iron and zinc, are indispensable trace elements for eukaryotic life, acting as co-factors in essential processes ranging from respiration and metabolism to DNA replication. These metals can be transported into cells by an evolutionary-conserved family of metal transporters; however, how the ubiquitous mammalian parasite *Toxoplasma gondii* acquired essential metals has been unknown. Here, we have identified and characterised the first iron and zinc importer in *T. gondii*. This transporter, named ZFT, localised to the parasite plasma membrane and is essential for the parasite's life cycle. We find ZFT is regulated by iron availability and overexpression sensitises cells to excess iron and zinc. Using a conditional knockdown system, we find that knockdown of ZFT leads to a reduction in mitochondrial respiration and a switch to a more quiescent lifecycle stage. To confirm transport activity, we find that knockdown of ZFT leads to a reduction is parasite-associated zinc and iron, and ZFT complements loss of zinc transporter activity in a yeast model. Overall, we have identified the first metal uptake transporter in *T. gondii* and demonstrated the importance iron and zinc transport to the cell. This finding reveals a critical piece of the puzzle in understanding how this obligate intracellular parasite thrives within its host.

Introduction

Iron, and other transition metals such as zinc, manganese and copper, are essential nutrients for almost all life, playing vital roles in biological processes including DNA replication, translation, cellular respiration and metabolism (Teh et al., 2024). As metals cannot diffuse across membranes, organisms have developed a range of strategies to take up metals, e.g. endocytosis of iron-bound transferrin by mammalian cells, or bacterial siderophore uptake systems. However, iron and other transition metals can catalyse reactive oxygen species, and so intracellular levels are tightly regulated (Galaris et al., 2019; Imlay et al., 1988).

Toxoplasma gondii is an obligate intracellular apicomplexan parasite which chronically infects around 30% of the world's population (Bisetegn et al., 2023). Although typically asymptomatic, Toxoplasmosis can be dangerous in immunocompromised individuals and to the developing foetus, resulting in severe congenital defects (Desmonts and Couvreur, 1974; Wong and Remington, 1994). Unusually among the Apicomplexa, *T. gondii* can infect all warm-blooded species, where it causes significant losses in ruminants (Stelzer et al., 2019) and it is

promiscuous, capable of infecting all nucleated cells (Tenter et al., 2000). Iron is essential for the parasite, and depletion using iron chelators restricts parasite replication *in vitro* and *in vivo* (Aghabi et al., 2023; Ferrer et al., 2012; Mahmoud, 1999; Raventos-Suarez et al., 1982; Sloan et al., 2025). *Toxoplasma* uses iron in essential cellular processes, including haem biosynthesis (Bergmann et al., 2020; Kloehn et al., 2021) and iron-sulphur (Fe-S) cluster biogenesis pathways (Aw et al., 2021). Beyond iron, the role of other transition metals in *Toxoplasma* biology is less well understood. It is predicted at least 800 *Toxoplasma* proteins require zinc, several of which have already been shown to be essential (Gissot et al., 2017; Semenovskaya et al., 2020). Although both iron and zinc are essential for the parasite, the mechanisms of their uptake in *Toxoplasma gondii* have not yet been established.

The ZIP (Zrt-/Irt-like protein)-domain containing proteins represent a large, highly conserved family of metal transporters (Guerinot, 2000). ZIP-domain proteins mediate iron and zinc uptake and intracellular trafficking in mammalian, plant and fungal cells. ZIP-domain containing proteins have also been utilised in eukaryotic parasites; a *Leishmania* ZIP-domain containing protein was identified as the major iron importer and was essential for intracellular replication (Huynh et al., 2006; Jacques et al., 2010). In *Plasmodium,* the plasma membrane-localised ZIP-domain containing protein ZIPCO was shown to be essential only in the liver stage of the parasite and was identified as a potential iron and zinc transporter based on partial rescue of a growth phenotype with iron and zinc (Sahu et al., 2014). Further, DMT1 was recently identified as an essential transporter in asexual *Plasmodium* replication. DMT1 was proposed to transport iron from the food vacuole to the cytosol, where it could be utilised by the apicoplast and mitochondrion (Loveridge and Sigala, 2024; Zhong and Zhou, 2023).

Toxoplasma gondii encodes four predicted ZIP-domain containing proteins: TGME49_261720, TGME49_254080, TGME49_500163 and TGME49_225530, none of which have been previously characterised. Our previous work identified that TGME49_261720 mRNA was stabilised under low iron by the presence of a stem-loop structure in the 3' UTR (Sloan et al., 2025). Loss of this regulation inhibited parasite survival under low iron, however the function of the protein was not addressed. Here, we characterise TGME49_261720, named Zn and Fe Transporter (ZFT), in detail. We find ZFT localisation and abundance is dynamic and dependent on parasite replication, vacuolar stage and iron availability. ZFT is essential for the parasite's lytic lifecycle and knockdown leads to inhibition of mitochondrial respiration and initiation of stage conversion and cannot be rescued by addition of exogenous metals. By quantifying parasite-associated metals, we show ZFT knockdown parasites have significantly less iron and zinc, and expression of ZFT rescues zinc uptake in mutant yeast. These data strongly support the role of ZFT as the major iron and zinc transporter in *Toxoplasma gondii.*

Results

Iron transport in *Toxoplasma gondii*

We examined the genome of *Toxoplasma* and identified four genes encoding predicted ZIP-domains (**Table 1**) (Amos et al., 2022). Of these, the ZIP domain of TGME49_254080 was relatively poorly conserved, and TGME49_500163 was predicted to contain only 3 transmembrane domains (rather than the 8-9 which characterise ZIP transporters) and so were removed from further analysis. Both TGME49_261720 and TGME49_225530 share limited homology with PfZIPCO (21% and 26% identity respectively) suggesting they are not orthologs of the *Plasmodium* gene. We examined the sequence conservation at the binuclear metal centres (BNC) M1 and M2 in transmembrane domains α4 and α5, which have been shown to be required for metal binding in a bacterial ZIP homolog (BpZIP) (Wiuf et al., 2022). Several residues are conserved in *Toxoplasma* ZIP proteins, including E181 and E211, which have been shown to be essential for transport activity (Wiuf et al., 2022) (**Figure 1a**). At M1, *Toxoplasma* TGME49_261720 contains an unusual HGxxEG motif, which we found only in coccidian ZIPs and in some marine organisms (e.g. *Branchiostoma floridae*), the impact of which is not clear (**Figure 1a**). In contrast, TGME49_225530 contains a HAxxEG motif, also found in prokaryotic ZIP9-like proteins (Hu, 2021). Interestingly, all parasite ZIP-domain proteins examined have a HK motif at the M2 metal binding site (**Figure 1a**), characteristic of the ZIPI family of ZIP-domain proteins, shared in HsZIP1 and HsZIP2 (Dufner-Beattie et al., 2003). This lysine likely prevents metal binding at this site, however has been shown to be required for transport activity in HsZIP2, possibly with a role in pH-responsivity of the transporter (Gyimesi et al., 2019). Based on the unusual conserved motif, the predicted plasma membrane localisation, essentiality in tachyzoites and our previous results demonstrating iron-mediated regulation (Sloan et al., 2025), we decided to functionally characterise TGME49_261720. We aligned the structural prediction of TGME49_261720 with the human ZIP-domain transporter ZIP1, a zinc importer (Gaither and Eide, 2001) and saw good alignment in the transmembrane domains, although TGME49_261720 has a large, unstructured loop, predicted to be on the outer lumen of the membrane (**Figure 1b**).

ID	Length (aa)	Predicted localisation (Barylyuk et al., 2020)	Phenotype score (Sidik et al., 2016)	TM domains	ZIP domain (E value)
TGME49_261720 (ZFT)	448	PM-integral	-4.31	8	5.9E-34
TGME49_254080	490	rhoptries 2	1.41	6	4.4E-16
TGME49_500163	1347	n/a	-1.4	3	5.9E-30
TGME49_225530	717	n/a	-2.94	8	7.2E-30

Table 1. Summarising known features of the four ZIP-domain containing proteins of *T. gondii.*

ZFT localisation is dynamic and vacuolar stage dependent

We have previously shown that the 3' UTR of ZFT plays an important role in iron-mediated regulation (Sloan et al., 2025) and so we tagged ZFT with a 3xHA epitope tag with the endogenous 3' UTR (ZFT-3HA$_{zft}$) as previously described (Sloan et al., 2025) (**Figure 1c**). Western blot demonstrated the protein ran at the expected size of around 50 kDa, although with multiple, non-specific lower bands (asterisks) (**Figure 1d**). Using immunofluorescence, we found that ZFT localised to the periphery of the parasite with a second foci at the basal pole of the cell (**Figure 1e**). To determine if changing the UTR affected ZFT localisation, we also tagged it with a 3xHA epitope tag followed by the *sag1* 3' UTR (ZFT-3HA$_{sag1}$) as previously described (Sloan et al., 2025) (**Figure S1a**) and saw a similar localisation (**Figure S1b**). Interestingly, localisation was highly dependent on the number of parasites per vacuole. In early stages (2-4 parasites/vacuole) ZFT localised peripherally, however this shifted to basal localisation in larger vacuoles (>8 parasites) (**Figures 1e, f**). Accompanying this change in localisation, we quantified the amount of ZFT-3HA$_{zft}$ and found that the mean fluorescence intensity (MFI) of ZFT/parasite was significantly higher in vacuoles with fewer parasites, compared to larger vacuoles with 8 parasites (< 0.001, one-way ANOVA, Tukey corrected for multiple comparisons) (**Figure 1g**). This suggests that ZFT is being degraded in a vacuolar stage-dependent manner. Previously, we showed that ZFT levels are regulated by low iron (Sloan et al., 2025). To determine if the localisation of ZFT was affected under iron starvation, we treated the parasites with the iron chelator deferoxamine (DFO) for 24 h. Under low iron conditions, ZFT-3HA$_{zft}$ localisation was maintained at the parasite periphery, although some intracellular vesicles could be observed (**Figure S1c**).

These data show ZFT expression and localisation is dynamic depending on vacuolar stage.

Figure 1. ZFT is a ZIP-domain containing protein with dynamic localisation. a. Alignment of ZIP proteins from *T. gondii* (TGME49_261720), *T. gondii* (TGME49_266800), *P. falciparum* (PF3D7_1022300), HsZIP1 and HsZIP2. Key conserved residues in the binuclear metal centres (BNC) M1 and M2 in transmembrane domains a4 and a5, which have been shown to be required for metal binding are highlighted. Conserved glutamate residues highlighted in purple. **b.** Alignment of the predicted structure of TGME49_261720 with the human transporter ZIP1. **c.** Schematic of F3ΔHX ZFT-3HA$_{zft}$ strain construction to endogenously tag ZFT with 3xHA epitope tags at the C-terminal under the endogenous promoter with the endogenous 3' UTR. **d.** Western blot analysis of ZFT-3HA$_{zft}$ confirming the expected band size (50 kDa). Non-specific bands indicated with asterisks. CDPK1 (TGME49_301440) used as a loading control. **e.** Immunofluorescence of ZFT-3HA$_{zft}$

ZFT expression is regulated by excess iron

The localisation and expression of transporters is often regulated by substrate availability (Conte and Walker, 2011; Gao and Dubos, 2021; Lee et al., 2020), and we have shown the iron transporter VIT is regulated by iron levels (Aghabi et al., 2023). We have previously demonstrated the *zft* 3' UTR is sufficient to stabilise the mRNA under low iron (Sloan et al., 2025), however expression under excess iron was not examined. To determine if ZFT-3HA$_{zft}$ is regulated by excess iron, we treated parasites with exogenous ferric ammonium chloride (FAC). At 18 h post treatment, ZFT-3HA$_{zft}$ levels were slightly (to around 87% of the untreated) but significantly ($p = 0.028$, two-tailed paired t test) reduced (**Figures 2a, b**). However, by 24 h post treatment we found that ZFT-3HA$_{zft}$ expression was reduced to around 40% of the untreated ($p = 0.0041$, two-tailed paired t test) (**Figures 2c, d**). We confirmed loss of ZFT-3HA$_{zft}$ expression under excess iron by immunofluorescence (**Figures 2e, f**), demonstrating ZFT expression is significantly reduced under excess iron.

Overexpression of ZFT leads to a change in iron sensitivity

Initial attempts to overexpress ZFT were unsuccessful, suggesting that overexpression was toxic. However, we were able to stably overexpress ZFT by coupling a second copy of ZFT-Ty expressed from a strong promotor, pTUB8 (Frénal et al., 2010) with DHFR using a P2A-slip peptide (Markus et al., 2019) and maintaining the line in pyrimethamine (**Figure 2g**). We validated the presence of the gene by PCR (**Figure S2a**), and the protein by western blotting (**Figure 2h**). To determine the effect of overexpression of ZFT on parasite fitness, we performed a plaque assay of both parental and ZFT-Ty parasites (**Figure 2i**). Parasites overexpressing ZFT form significantly ($p = 0.0027$, unpaired two-tailed t test) fewer plaques than parental parasites (**Figure 2j**), with significantly smaller area ($p < 0.0001$, two-tailed unpaired t test with Welch's correction) (**Figure 2k**), confirming that overexpression of ZFT inhibits parasite replication. Overexpression of metal transporters can alter the sensitivity to environmental metal conditions (Connolly et al., 2002; Li and Kaplan, 1998). To test this, we quantified the effects of ZFT overexpression on iron sensitivity. Our data show parasites overexpressing ZFT-Ty are significantly ($p = 0.0027$, two-tailed unpaired t test) more sensitive (EC$_{50}$ = 0.386 mM, 95% C.I. 0.315 to 0.479 mM)

Figure 2. ZFT-3HA$_{zft}$ is regulated by iron, and overexpressing ZFT leads to a change in iron sensitivity. a. Western blot analysis showing ZFT-3HA$_{zft}$ expression levels 18 h post infection in untreated and high iron (500 µM FAC) conditions. CDPK1 was used as a loading control. **b.** Quantification of ZFT-3HA$_{zft}$ from three independent experiments. Bars are at mean ± SD. *p* values from two-tailed paired *t* test. **c.** As above, at 24 h post infection in untreated and high iron (500 µM FAC) conditions. **d.** Quantification of ZFT-3HA$_{zft}$ from three independent experiments. Bars are at mean ± SD. *p* values from two-tailed paired *t* test. **e.** Immunofluorescence assay of ZFT-3HA$_{zft}$ at 24 h post infection in untreated and high iron (500 µM FAC) conditions. Scale bars 5 µm. **f.** Quantification of ZFT-3HA$_{zft}$ localisation (peripheral/basal or no signal) in untreated and high iron conditions. Bars are mean of three independent experiments, ± SD. **g.** Schematic of ZFT-Ty strain construction to overexpress ZFT from the TUB8 promoter in RHtdTomato parental line. **h.** Western blot of ZFT-Ty showing expected size. **i.** Plaque assay of parental and ZFT-Ty parasites in untreated and high iron (500 µM FAC) conditions. Quantification of plaque number (**j**) and area (**k**) for parental and ZFT-Ty parasites, *p* values are unpaired two-tailed *t* test. Points represent individual plaque areas from three independent experiments, bars at mean

to excess iron, compared to the parental line (EC_{50} = 1.29 mM, 95% C.I. 0.813 to 2.01 mM) (**Figures 2l, S2b**). Excess iron leads to parasite death largely through oxidative stress (Aghabi et al., 2023). To confirm that parasites were not more sensitive based on non-specific increased sensitivity to oxidative stress, we treated parasites with sodium arsenite (Ars), which has previously been used to induce oxidative stress in *T. gondii* (Aghabi et al., 2023; Augusto et al., 2021). However, overexpression of ZFT does not lead to any significant change in oxidative stress sensitivity (**Figure S2c, S2d**), suggesting the phenotype is specific to iron. We then tested how ZFT-Ty overexpression affected sensitivity to iron starvation using DFO. We found that parasites overexpressing ZFT are significantly (p = 0.0246, two-tailed unpaired t test) more resistant (EC_{50} = 22.81 μM, 95% C.I. 16.8 to 30.6 μM) to iron chelation, compared to the parental line (EC_{50} = 8.18 μM, 95% C.I. 6.94 to 9.71 μM) (**Figures 2m, S2e**). These data suggest that overexpression of ZFT leads to alterations in parasite iron sensitivity.

ZFT is essential for parasite survival

To determine the importance of ZFT in parasite replication, an inducible knockdown line was generated by replacing the promotor of ZFT with the T7S4 promoter which is inhibited in the presence of ATc (Sheiner et al., 2011) (**Figure 3a**), which was validated by PCR (**Figure S3a**). To verify conditional knockdown, we performed RT-qPCR as previously described (Sloan et al., 2025). By 24 h post induction with ATc, we found that *zft* mRNA was significantly (p = 0.0032, two-tailed unpaired t test with Welch's correction) reduced (**Figure 3b**). We epitope tagged ZFT in the T7S4-ZFT background (**Figure 3c**) and confirmed integration by PCR (**Figure S3b**). We further validated this by western blot (**Figure 3d**) and show that there is a significant ($p <$ 0.0001, two-tailed paired t test) drop in ZFT-3HA$_{zft}$ expression levels by 24 h post induction. 48 h after adding ATc, only around 10% of the protein remained (p = 0.0131, two-tailed paired t test) (**Figure 3e**). Moreover, we found that ZFT-3HA$_{zft}$ was undetectable by immunofluorescence by 24 h post induction (**Figure 3f**). Previous studies on both bacterial and eukaryotic ZIPs suggest that these proteins function in either homodimers or heterodimers (Ahern et al., 2020; Bin et al., 2011; Lin et al., 2010; Taylor et al., 2016). To determine if ZFT-3HA$_{zft}$ forms a complex in *Toxoplasma*, we performed a Blue native PAGE (BN-PAGE) and found that ZFT-3HA$_{zft}$ forms a complex of ~100 kDa, which is lost upon knockdown, suggesting the formation of a homodimer (**Figure 3g**).

Figure 3. ZFT knockdown blocks parasite replication. a. Schematic of F3ΔHX T7S4-ZFT under the T7S4 inducible promoter. **b.** RT-qPCR showing ZFT mRNA abundance, relative to 0 h, normalised to actin. Bars at mean of three independent experiments, ± SD. *p* values from two-tailed unpaired *t* test. **c.** Schematic of F3ΔHX

T7S4-ZFT-3HA$_{sag1}$ under the T7S4 inducible promoter. **d.** Western blot demonstrating successful knockdown of ZFT at 24, 48, 72 and 96 h post ATc induction. CDPK1 used as a loading control. *p* values from two-way ANOVA, Tukey corrected for multiple comparisons. **e.** Quantification of (**d**), relative to 0 h, normalised to CDPK1. Bars at mean of four independent experiments, ± SD. *p* values from two-way ANOVA, Tukey corrected for multiple comparisons. **f.** Immunofluorescence assay showing T7S4-ZFT-3HA$_{sag1}$ expression in untreated parasites and after induction with ATc for 48 h. Scale bar 5 μm. **g.** Blue native PAGE of T7S4-ZFT-3HA$_{sag1,}$ untreated and + ATc 48 h induction, showing a ZFT-HA complex at around 100 kDa. CDPK1 used as a loading control. **h.** Plaque assay of parental and T7S4-ZFT parasites untreated and ATc treated, 6 days, showing knockdown of ZFT renders the parasite unable to form plaques in the host cell monolayer. **i.** Quantification of (**h**) showing number of plaques, normalised to parental untreated. Bars at mean of three independent experiments, ± SD. *p* values from two-way ANOVA, Tukey corrected for multiple comparisons. **j.** Quantification of plaques area from (**h**). Points represent individual plaques from three independent experiments. Bars show the mean ± SD. *p* values from ordinary one-way ANOVA, Tukey corrected for multiple comparisons. **k.** Plaque assays with addition of 200 μM FAC does not rescue ZFT knockdown parasites. **l.** Quantification of plaque area generated by T7S4-ZFT parasites in untreated and high iron (200 μM FAC) conditions. Points represent individual plaque areas from three independent experiments. Bars show mean ± SD. *p* values are from two-tailed unpaired *t* test with Welch's correction. **m.** Quantification of plaque number, normalised to parental untreated. Bars are at the mean of three independent experiments, ± SD. *p* values from two-way ANOVA, Tukey corrected for multiple comparisons. **n.** Quantification of plaque area generated by parental and T7S4-ZFT parasites with and without ATc, in the presence of excess iron (200 μM FAC). Points represent individual plaque areas from three independent experiments. Bars show the mean ± SD. *p* values from ordinary one-way ANOVA, Tukey corrected for multiple comparisons.

To examine the role of ZFT, a plaque assay was performed (**Figure 3h**). Replacement of the ZFT promotor led to a significant (*p* = 0.0052, two-way ANOVA, Tukey corrected for multiple comparisons) growth defect (**Figures 3h, i, j**). This, combined with our previous study (Sloan et al., 2025), demonstrates that correct expression of ZFT is important for parasite fitness. Addition of ATc renders T7S4-ZFT parasites unable to form plaques in the host cell monolayer (**Figures 3h, i, j**), confirming that expression of ZFT is essential for the parasite. In *Plasmodium*, the growth defect upon loss of DMT1 (Loveridge and Sigala, 2024) or ZIPCO (Sahu et al., 2014) was able to be rescued by addition of exogenous substrate. To determine if loss of ZFT could be rescued by addition of iron, we treated ZFT knockdown cells with 200 μM FAC and performed a plaque assay (**Figure 3k**). We found that replacing the ZFT promoter alone led to a significant reduction in plaque area (*p* < 0.0001, two-tailed unpaired *t* test with Welch's correction) in high iron conditions (200 μM FAC) compared to untreated conditions (**Figure 3l**). However, exogenous iron was unable to rescue the growth defect, either by plaque number or area (**Figures 3k, m, n**). To determine if iron could restore growth when some ZFT remained, we performed a plaque assay with half the standard concentration of ATc (0.25 μM) (**Figure S3c**). We find that in

these conditions, ZFT KD parasites are still unable to form any plaques, even in the presence of excess iron (200 µM FAC) (**Figures S3c, d, e**). This suggests a strong irreversible growth defect and the essentiality of ZFT as the major iron transporter.

ZFT knockdown reduces mitochondrial respiration, and expression of Fe-S protein SDHB

Iron is required for proteins essential in parasitic organelles such as the apicoplast and mitochondrion (Maclean et al., 2024; Pamukcu et al., 2021; Renaud et al., 2022) and disruption of organellar function can lead to morphological changes (Mallo et al., 2021; Wu et al., 2022). To determine if ZFT knockdown was associated with changes in organellar morphology, we imaged both organelles at 24 h and 48 h of ATc. However, we saw no gross morphological defects in the apicoplast (**Figure 4a**) or mitochondria (**Figure 4b**).

Iron is required in proteins of the electron transport chain (Maclean et al., 2021). In other organisms, depletion of iron leads to defects in mitochondrial respiration (Fujimaki et al., 2019; Loveridge and Sigala, 2024), which can be assessed by quantification of oxygen consumption using a Seahorse assay (Hayward et al., 2022). We found that 48 h of ZFT knockdown significantly reduced basal mitochondrial oxygen consumption rates (mOCR) (p = 0.0233, two-tailed unpaired t test with Welch's correction) (**Figures 4c, d**), as well as maximal mitochondrial oxygen consumption rates (p = 0.0054, two-tailed unpaired t test with Welch's correction) (**Figure 4e**), compared to untreated parasites. Interestingly, we find ZFT knockdown does not significantly change the extracellular acidification rate (ECAR), a proxy for glycolysis (Hayward et al., 2022) (**Figure 4f**). The metabolic map suggests that ZFT knockdown leads to a reduction in oxidative phosphorylation and a more quiescent phenotype (**Figure 4g**). Changes in mitochondrial respiration have been linked to changes in cristae morphology (Baker et al., 2019; Huet et al., 2018). Using transmission electron microscopy (TEM) at 72 h post ATc treatment, we saw no clear changes in cristae morphology, compared to untreated parasites (**Figure 4h**). To identify a link between ZFT and mitochondrial respiration, we examined the expression of the key mitochondrial protein succinate dehydrogenase B (SDHB). SDHB has essential roles both in the TCA cycle and in complex III (Maclean et al., 2021; Silva et al., 2023) and contains three Fe-S clusters. We endogenously tagged SDHB (TGME49_215280) in the T7S4-ZFT background (**Figure S4a**), confirmed integration by PCR (**Figure S4b**), and quantified protein levels following 48 h of ZFT knockdown. Upon depletion of ZFT, SDHB-3HA expression levels were significantly (p = 0.0205, two-tailed paired t test) reduced to around 34%, compared to untreated cells (**Figure 4i, j**). As a control, we also tagged the essential cytosolic Fe-S protein ABCE1 (TGME49_216790) (Barthelme et al., 2011, 2007; Maclean et al., 2024; Renaud et al., 2025) (**Figure S4c**) and confirmed integration by PCR (**Figure S4d**). However, ABCE1-3HA expression levels remained unchanged at 48 h upon ZFT knockdown (p = 0.1758, two-tailed paired t test) (**Figure 4k, l**), demonstrating that not all Fe-S proteins are affected equally upon ZFT knockdown.

Figure 4. ZFT knockdown reduces mitochondrial respiration and the expression of Fe-S protein SDHB. a. Immunofluorescence assay of T7S4-ZFT-3HA$_{sag1}$ in the absence and presence of ATc for 24 h and 48 h, staining for CPN60 (apicoplast marker). **b.** Immunofluorescence assay as above, staining for TOM40 (mitochondrial marker). Scale bar 5 µm **c.** Mitochondrial oxygen consumption rate (mOCR) of T7S4-ZFT, untreated or ATc treated for 48 h via Seahorse assay. Quantification of basal

These data demonstrate ZFT knockdown reduces mitochondrial respiration, potentially due to loss of iron-containing proteins in the mitochondrial electron transport chain.

Knockdown of ZFT triggers partial parasite differentiation

To determine the effect of ZFT knockdown, we examined parasite replication. Interestingly, replication was not immediately inhibited, and at 48 h post ATc addition we observed large vacuoles containing many nuclei (**Figure 5a**), although these large vacuoles did not egress from the host. Around 50% of these vacuoles exhibited loss of normal vacuolar organisation (**Figure 5b**) and we saw frequent examples of asynchronous replication, with only single parasites within a vacuole containing daughter cell scaffolds (**Figure 5a, inset**).

To examine the effects of loss of ZFT in more detail, we imaged parasites by TEM. Although there were no obvious morphological changes, at 5 days post knockdown we observed a heavily vesiculated parasitophorous vacuolar membrane (PVM) with the accumulation of small vesicles in the intravacuolar space. In contrast, the PVM of untreated parasites appeared smooth, with no vesicular accumulation (**Figure 5c**). Further, we observed multiple enlarged, electron-lucent vesicles upon ZFT knockdown which were not presented in the untreated cells (**Figure 5c**). Similar large vesicles have previously been reported to contain materials that form the cyst wall (Tomita et al., 2013), a characteristic of *T. gondii* conversion to bradyzoites. Previously, we and others have demonstrated that loss of iron is sufficient to drive bradyzoite conversion (Renaud et al., 2024; Sloan et al., 2025; Ying et al., 2024). To determine if iron depletion could mimic this phenotype, parental parasites were treated with DFO for 5 days. As with the knockdown of ZFT, iron-depleted parasites also exhibit similar large vesicles close to the PV membrane (**Figure S4e**).

Figure 5. Knocking down ZFT triggers partial bradyzoite differentiation. a. Immunofluorescence of T7S4-ZFT-3HA$_{sag1}$ parasites uninduced or induced with ATc for 48 h. Highlighted is a region where parasites are disorganised and asynchronously replicating. Scale bar 5 μm. **b.** Quantification of the percentage of parasites/vacuole in parental and T7S4-ZFT parasites in the presence and absence of ATc for 48 h. 100 vacuoles for each condition were counted. Bars at mean of four independent replicates, ± SD. **c.** TEM images of T7S4-ZFT parasites with and without ATc for five days. Detail highlights regions of PVM. Yellow arrows point to enlarged vesicles in

To determine if loss of ZFT is sufficient to induce stage conversion, we examined expression bradyzoite specific markers upon ZFT knockdown. BAG1 is bradyzoite specific protein (Kannan et al., 2019) and DBL is a lectin which binds to the heavily glycosylated cyst wall (Tomita et al., 2013). At 72 and 96 h post ATc addition we saw vacuoles expressing BAG1 (**Figure 5d**) and with peripheral DBL staining around the vacuole (**Figure 5e**). We quantified this and saw a significantly higher percentage of vacuoles expressing BAG1 ($p = 0.0013$, two-way ANOVA, Tukey corrected for multiple comparisons) and DBL ($p = 0.0001$, two-way ANOVA, Tukey corrected for multiple comparisons) upon ZFT knockdown (**Figures 5f, g**). These data demonstrate ZFT depletion leads to the expression of the bradyzoite-specific markers BAG1 and DBL.

T. gondii differentiation allows cells to survive otherwise fatal stresses, and parasites can reactivate upon removal of the stress (Cerutti et al., 2020; Soete et al., 1993; Sullivan and Jeffers, 2012). To determine if parasites initiating conversion upon ZFT knockdown remained viable, we quantified parasite survival. Parasites were treated with ATc for 72 h, washed, then incubated without ATc for a further 6 days. After ATc washout, plaques formed, indicating that some parasites remained viable (**Figure 5h**). Upon quantification, around 40 % of parasites were able to reestablish growth after 72 h ZFT knockdown ($p < 0.0001$, two-tailed unpaired *t* test) (**Figure 5i**), a similar proportion to those expressing the bradyzoite marker BAG1 (**Figure 5f**). These results imply that although many Rh parasites are unable to survive a transient loss of ZFT, around 40% of vacuoles initiate transition to bradyzoites and can survive transient loss of ZFT.

ZFT knockdown cannot be rescued by excess zinc

As ZIP-domain proteins commonly transport zinc, we next aimed to assess if ZFT also plays a role in zinc uptake in *T. gondii.* Although excess iron was unable to rescue the growth phenotype upon ZFT knockdown, we were curious to see the effects of zinc on the growth phenotype. Excess zinc was unable to rescue plaque number (**Figures 6a, b**) or area (**Figure 6c**) in the absence of ZFT. As with iron, changing the promoter alone significantly ($p < 0.0001$, ordinary one-way ANOVA, Tukey corrected for multiple comparisons) sensitised the parasites to excess zinc, leading to a

decrease in plaque size (**Figure 6d**). We also tested if a zinc ionophore (2-Mercaptopyridine N-oxide) could impact growth, however the zinc ionophore was also unable to rescue the growth phenotype upon knockdown of ZFT (**Figures S5a, b**). As we had previously examined the role of iron, we next tested if both excess iron (200 µM FAC) and zinc (25 µM $ZnSO_4$) were required to rescue the growth phenotype upon ZFT knockdown (**Figure 6a**). However, addition of both metals was unable to rescue the T7S4 growth defect (**Figures 6a, b**). Interestingly, changing the promoter alone leads a significant ($p < 0.0001$, two-way ANOVA, Tukey corrected for multiple comparisons) decrease in plaque numbers (to around ~20%) compared to parental parasites in the iron and zinc conditions (**Figure S5c**), suggesting that overexpressing ZFT makes the parasites hypersensitive to both iron and zinc. We also tested the sensitivity of the ZFT-Ty overexpression line to zinc. As suggested by the plaque assays, overexpression of ZFT lead to a small but significant ($p = 0.01$, two-tailed unpaired t test) increase in zinc sensitivity (**Figures 6e, S5d**).

We next aimed to understand if the expression of ZFT is dependent on zinc availability. We treated ZFT-3HA$_{zft}$ parasites with either 50 µM $ZnSO_4$, or 5 µM TPEN, a cation chelator frequently used to reduce zinc levels (Schaefer-Ramadan et al., 2019) for 18 h and 24 h. Unlike for iron, excess zinc did not alter ZFT-3HA$_{zft}$ expression at 18 h (**Figures 6f, g**) or 24 h post treatment (**Figures 6h, i**), and we saw no change by immunofluorescence (**Figure S5e**). However, at both timepoints, TPEN treatment led to a significant reduction in ZFT ($p = 0.0191$, and $p = 0.017$, one-way ANOVA, Tukey corrected for multiple comparisons) (**Figures 6f, g, h, i**).

ZFT expression complements lack of zinc transport in *S. cerevisiae*

To investigate whether ZFT has a functional role in zinc uptake in *T. gondii*, we first codon optimized the full length ZFT sequence for expression in *Saccharomyces cerevisiae*. This was then constitutively expressed in either the parental strain (DY1457) or a strain lacking both the low and high affinity zinc transporters, Δzrt1/2 (Ullah et al., 2018). Under standard conditions, expression of TgZFT complemented the mild growth defect of the Δzrt1/2 mutant (**Figure 6j**). Upon removal of zinc by chelation by EGTA, the Δzrt1/2 mutant is unable to grow, however this is rescued by expression of TgZFT (**Figure 6k**). These data demonstrate ZFT expression complements the loss of zinc transport in the Δzrt1/2, confirming ZFT as a functional zinc transporter.

Figure 6. ZFT expression depends on the availability of zinc, and ZFT expression complements lack of zinc transport in *S. cerevisiae*. a. Plaque assays showing that the addition of 25 µM ZnSO$_4$, and the combination of 25 µM ZnSO$_4$ and 200 µM FAC, can not rescue ZFT knockdown. **b.** Number of plaques after treatments. Bars are at mean ± SD. *p* values from two-way ANOVA, Tukey corrected for multiple comparisons. **c.** Plaque areas of parasites treated with excess zinc (25 µM ZnSO$_4$). Points represent individual plaque areas from two (parental -ATc) or three (all other conditions) independent experiments. Bars show the mean ± SD. *p* values from ordinary one-way ANOVA, Tukey corrected for multiple comparisons. **d.** Plaque area from T7S4-ZFT parasites in untreated and high zinc (25 µM ZnSO$_4$) conditions. Points represent individual plaque areas from three independent experiments. Bars show the mean ± SD. *p* values are from two-tailed unpaired *t* test with Welch's correction. **e.** Graph showing mean EC$_{50}$ for zinc for parental and ZFT-Ty parasites. Bars are mean of n=3, ± SD. *p* values from two-tailed unpaired *t* test. **f.** Western blot analysis showing ZFT-3HA$_{zft}$ expression levels 18 h post infection in untreated, high zinc (50

µM ZnSO$_4$) and 5 µM TPEN conditions. CDPK1 used as a loading control. **g.** Quantification of ZFT-3HA$_{zft}$ levels at 18 h from three independent experiments. Bars are mean ± SD. *p* values from one-way ANOVA, Tukey corrected for multiple comparisons. **h.** Western blot as above, at 24 h post infection. **i.** Quantification of ZFT-3HA$_{zft}$ levels at 24 h from three independent experiments. Bars are mean ± SD. *p* values from one-way ANOVA, Tukey corrected for multiple comparisons. **j.** Spot assay showing TgZFT expression rescues growth in Δzrt1/2 *S. cerevisiae*. Representative of two independent biological experiments. **k.** Under zinc chelation (5 mM EGTA), TgZFT allows Δzrt1/2 growth. Representative of two independent biological experiments.

Knocking down ZFT results in a decrease in parasite associated metal

To determine if loss of ZFT led to changes in the metal content of parasites, we used the fluorescence iron indicator FerroOrange, which increases fluorescence in the presence of labile iron (Weber et al., 2020; Yu et al., 2022). This has not previously been used in *Toxoplasma*, so to validate, we pre-treated parasites with 200 µM FAC intracellularly for 48 h before live imaging (**Figure 7a**). Parasites preloaded with iron have a significantly higher MFI of FerroOrange compared to untreated ($p < 0.0001$, two-tailed unpaired *t* test with Welch's correction) (**Figure 7a**)**,** validating this approach. Upon 48 h KD of ZFT, the MFI of FerroOrange was significantly ($p = 0.0027$, two-tailed unpaired *t* test with Welch's correction) reduced, compared to untreated parasites (**Figure 7b**). We also used the FluoZin-3 indicator which increases in fluorescence in the presence of zinc (Li et al., 2009; Rajapakse et al., 2017). As above, to validate this approach we saw that zinc-treated parasites had significantly ($p < 0.0001$, two-tailed unpaired *t* test with Welch's correction) higher fluorescence than untreated cells (**Figure 7c**). However, upon 24 h KD of ZFT, the MFI of FluoZin-3AM was significantly reduced compared to untreated parasites ($p < 0.0001$, two-tailed unpaired *t* test with Welch's correction) (**Figure 7d**), suggesting lower parasite zinc levels in the absence of ZFT.

Figure 7. Knocking down ZFT results in a decrease in parasite associated metal.
a. Live cell imaging and quantification of FerroOrange stained parental parasites, untreated or treated with 200 μM FAC for 48 h. Scale bar 5 μm. Points represent the

MFI of FerroOrange/parasite from four independent experiments. Bars show the mean ± SD. *p* value from two-tailed unpaired *t* test with Welch's correction. **b.** Live cell imaging and quantification of FerroOrange-stained T7S4-ZFT parasites with and without ATc for 48 h. Points are MFI of FerroOrange/parasite from four independent experiments. Bars show the mean ± SD. *p* value from two-tailed unpaired *t* test with Welch's correction. **c.** Live cell imaging and quantification of FluoZin-3AM stained parental parasites treated with 50 μM $ZnSO_4$ for 24 h. Points represent MFI of FluoZin-3AM/parasite from three independent experiments. Bars show the mean ± SD. *p* value from two-tailed unpaired *t* test with Welch's correction. **d.** Live cell imaging and quantification of T7S4-ZFT parasites with and without ATc for 48 h. Scale bar 5 μm. Points represent the MFI of FluoZin-3AM per parasite from three independent experiments. Bars are mean ± SD. *p* value from two-tailed unpaired *t* test with Welch's correction. **e.** ICP-MS quantification of iron/parasite (fM) from T7S4-ZFT parasites with and without ATc at 24 h and 48 h. Bars are at the mean (n = 7 for – ATc, n = 5 for + ATc 24 h and n = 2 for + ATc 48 h) ± SD. *p* values from one-way ANOVA, Tukey corrected for multiple comparisons. **f.** ICP-MS quantification of zinc/parasite (fM) from T7S4-ZFT parasites with and without ATc 24 h and 48 h. Bars are at the mean (n = 5 for – ATc and + ATc 24 h and n = 2 for + ATc 48 h) ± SD. *p* values from one-way ANOVA, Tukey corrected for multiple comparisons. XFM of extracellular T7S4-ZFT parasites, untreated and treated for 24 and 48 h, showing phosphorus, sulphur and iron (**g**) and zinc (**h**). **i.** XFM of untreated parasites showing iron and zinc colocalisation, along with phosphorus. Quantification of Fe (**j**) and Zn (**k**) following ZFT knockdown. Bar at mean ± SD. *p* values from one-way ANOVA, Tukey corrected for multiple comparisons. **l.** Quantification of changes in Fe, Zn, S and P as a percentage of untreated cells, following the addition of ATc. Bar at mean ± SD. Scale bar for all images 5 μm.

To further examine the effect of knocking down ZFT on parasite-associated metal levels, we performed inductively coupled plasma-mass spectrometry (ICP-MS), an elemental analysis technique used to quantify elemental abundance from complex environments (Al-sandaqchi et al., 2018; Hare et al., 2012). At both 24 h and 48 h ATc treatment, we saw no significant changes in the amount of iron in the parasites (**Figure 7e**). However, iron levels were very close to the limit of detection, even in untreated cells, and iron detection by ICP-MS can be challenging due to interference (Segura et al., 2003). However, at both 24 h and 48 h ATc treatment, ZFT knockdown parasites had significantly reduced parasite associated zinc (*p* = 0.0062, and *p* = 0.0073, two-tailed paired *t* test) (**Figure 7f**).

Iron can also be quantified at subcellular resolution using X-ray fluorescence microscopy (XFM) (Aghabi et al., 2023; Miller and Ralle, 2024; Paunesku et al., 2021), which allows both spatial and quantitative data. Using XFM we were able to detect iron in extracellular parasites (**Figure 7g**), which frequently appeared localised to distinct foci in the parasites, which we believe to be the PLVAC based on our previous observations (Aghabi et al., 2023). However, by 48 h post ATc induction we were no longer able to detect iron within the parasites. We also assessed zinc localisation. In untreated cells, zinc localised to what we believe is the parasite nucleus (**Figure 7h**),

as has been seen in other organisms (Olea-Flores et al., 2022). We also saw specific foci of zinc within the cells, which, based on previous results (Chasen et al., 2019), we assume to be the PLVAC. Indeed, we found that iron and zinc, along with phosphorus frequently colocalised in extracellular parasite (**Figure 7i**), as previously has been suggested (Chasen et al., 2019). As with iron, depletion of ZFT led to a decrease in parasite associated zinc, with a loss of zinc both within the nuclei and within foci, which was very evident by 48 h post knockdown (**Figure 7h**).

Parasite associated metals were quantified in untreated parasites and at both 24 and 48 h post ZFT knockdown by XFM. At both timepoints, knockdown parasites have significantly ($p < 0.0001$, ordinary one-way ANOVA, Tukey corrected for multiple comparisons) less parasite-associated iron, from ~250 nM in untreated, to ~150 nM at 24 h and ~95 nM at 48 h (**Figure 7j**). At both 24 h and 48 h KD of ZFT, *T. gondii* parasites have significantly ($p < 0.0001$, ordinary one-way ANOVA, Tukey corrected for multiple comparisons) less parasite associated zinc, from ~6 µM in untreated, to ~3.5 µM at 24 h and ~1.7 µM at 48 h (**Figure 7k**). Interestingly, we found that the concentrations of both sulphur ($p = 0.0014$, ordinary one-way ANOVA, Tukey corrected for multiple comparisons) (**Figure S6a**), and phosphorus ($p = 0.008$, ordinary one-way ANOVA, Tukey corrected for multiple comparisons) (**Figure S6b**), were also significantly reduced at 48 h following ZFT knockdown, although the magnitude of the decrease is less than for iron and zinc. Comparison of the changes demonstrated a clear, time-dependent depletion of both iron and zinc, with much smaller drops (around 10-20%) in sulphur and phosphorus (**Figure 7l**). Both sulphur and phosphorus are highly abundant in cellular macromolecules, and this small decrease may speak to defects in protein synthesis or DNA replication upon ZFT knockdown.

Together, these data validate the role of ZFT as the major iron and zinc transporter of *T. gondii.*

Discussion

Here we have identified and characterised the first ZIP-domain containing protein in *Toxoplasma,* showing that ZFT is the major iron and zinc importer for the parasite. Sequence analysis of ZFT shows conservation of essential metal coordinating residues and identified a new motif, conserved in coccidia, in the M1 metal binding domain, although with an unknown role in transport. ZFT also encodes a leucine at position 403, previously associated with broad substrate specificity (Gyimesi et al., 2019), which fits with our data that ZFT transports both iron and zinc. However, future studies will examine the substrate preference and kinetics of this transporter in greater detail.

ZIP proteins mediate metal transport both across the plasma membrane, and from vesicles into the cytosol. The localisation of ZFT appears highly dynamic through the parasite's intracellular replication, moving from the plasma membrane to basally

clustered intracellular vesicles, after 2-3 rounds of replication. This vacuole stage-dependent localisation is highly unusual in *Toxoplasma* and has not previously been observed. Unfortunately, fluorescent tags led to significant miss-localisation of ZFT to the ER (not shown), so we were unable to observe this trafficking in live cells. Currently, our working model is that ZFT is most active at the plasma membrane, enabling rapid metal incorporation into new vacuoles, and is then internalised and later degraded, presumably to prevent toxic metal overload (Aghabi et al., 2023). This may explain the persistence of plasma-membrane localised ZFT in iron starvation, and its degradation in excess iron. However, from our current results, we cannot exclude the hypothesis that the parasite is taking up extracellular material in digestive vesicles, and ZFT transports metals freed from digested proteins into the cytosol. Future studies examining the kinetics of metal uptake, and the localisation kinetics of ZFT in intracellular parasites may be able to address this remaining question.

In *Toxoplasma,* vacuolar iron (Aghabi et al., 2023) and zinc (Chasen et al., 2019) transporters have been identified with roles in metal storage and detoxification. However, ZFT is the first *Toxoplasma* metal uptake transporter identified and characterised. This transporter is essential, highlighting the essential role of iron and zinc to the parasite. Interestingly, and unlike in *Plasmodium* (Loveridge and Sigala, 2024; Sahu et al., 2014), we were unable to complement the loss of this transporter by addition of exogenous metals. One possible explanation could be lack of metal access to the PV, however our metal dye experiments and the growth phenotype after treatment increases our confidence that exogenous metals reach the vacuole. A more compelling argument is that ZFT is both essential and sufficient for iron and zinc transport, and in its absence, there are no other transporters able to complement for its loss. As there are three other predicted ZIP transporters in *Toxoplasma*, this raises the questions of their functions within the cell, which will be addressed in future studies.

Knockdown of ZFT triggers differentiation into bradyzoites, which complements previous studies demonstrating a similar phenotype upon iron chelation (Renaud et al., 2024; Ying et al., 2024). The pathway of iron-starvation triggering differentiation is not yet known, although it is compounded by the action of BFD1 (Sloan et al., 2025). Interestingly, loss of ZFT leads to a significant inhibition of mitochondrial respiration, potentially linked with the depletion of the essential, iron-bound protein SDHB (Maclean et al., 2021). Inhibition of respiration has previously been linked to bradyzoite conversion (Tomavo and Boothroyd, 1995), however we do not yet know the signalling factors linking iron, zinc or mitochondrial function to bradyzoite differentiation. In contrast to *Plasmodium* (Loveridge and Sigala, 2024), we did not see a change in apicoplast replication. Iron is known to be required in the apicoplast (Renaud et al., 2022), although a potential role for zinc has not yet been established. It is possible that remaining iron (or zinc) is prioritised for traffic to the apicoplast, as a similar process has been observed in green plants (Blaby-Haas and Merchant, 2013), however no apicoplast metal transporters have yet been identified. While the use of iron within the parasite has been previously examined, the role of zinc in *Toxoplasma* is unknown. XFM suggests a concentration of zinc within the nucleus, where it is expected to play a role in DNA replication and gene expression (Olea-Flores et al., 2022). Beyond the nucleus, zinc is clearly concentrated within one or

two foci, in agreement with zinc storage in the PLVAC, as previously hypothesised (Chasen et al., 2019). Future studies will be needed to unpick both the specific roles of zinc in *Toxoplasma*, and the parasite's response to zinc starvation, as these are likely highly heterogenous.

ZFT appears to be highly regulated by iron availability, at the mRNA (Sloan et al., 2025), protein, and localisation levels. However, ZFT is not regulated by excess zinc, although zinc-mediated ZIP-proteins degradation has been observed in other organisms (e.g. (Pang et al., 2023)). Interestingly, there is evidence from other organisms that excess zinc modulates both iron uptake (Blaby-Haas and Merchant, 2013; Xu et al., 2019) and copper homeostasis (Xu et al., 2019). It is possible that excess zinc leads to a decrease in iron acquisition, stabilising ZFT expression, however this has not yet been assessed. Previously, we have shown that ZFT is stabilised under iron starvation (Sloan et al., 2025), however we did not see the same effect after treatment with the zinc chelator TPEN. There has not yet been any investigation into potential zinc-mediated regulation in *Toxoplasma*, and so we are currently unable to say if this is due to a lack of zinc sensing and response in these parasites, or the treatment or timings we used was insufficient to trigger such a pathway.

Here, we were able to complement growth of a yeast mutant lacking the major zinc transporters *zrt1/2* with ZFT, demonstrating sufficient expression and localisation of this transporter in yeast. However, despite multiple attempts, we were unable to express codon-optimised ZFT in an iron uptake (*fet2/fet3*) yeast mutant. The reason for this was not clear, but it is possible that the *fet2/fet3* yeast strain is less able to survive transformation, or that in this background, expression of ZFT is toxic.

The use of XFM to quantify parasite-associated metals has been vital in this work. Beyond localising iron, zinc and phosphorus to an overlapping compartment in extracellular parasites, we were able to show that knockdown of ZFT lead to significant decreases in parasite-associated iron and zinc. There are some limitations of XFM, e.g. we were unable to quantify copper or manganese using this approach, probably due to low concentrations. Further, for technical reasons we worked with chemically fixed parasites which may have led to an underestimate of metal concentrations (Jin et al., 2017). However, this provides direct evidence of changes in the elemental composition of parasites in the absence of ZFT.

In summary, here we characterise the first iron and zinc uptake transporter in *Toxoplasma*. ZFT is indispensable, highlighting the importance of iron and zinc uptake for the parasite. The identification of this transporter opens new avenues for future investigation into the essential metallobiology of these parasites.

Acknowledgments

D.A. is supported by the Wellcome IIB PhD program (218518/Z/19/Z) and E.G. is supported by the BBSRC NorthWestBio PhD program (BB/X010902/1). This work was supported by the BBSRC (BB/W014947/1) to L.P. and C.R.H. C.R.H. is funded by a Sir Henry Dale Fellowship from the Wellcome Trust and the Royal Society (213455/Z/18/Z). We acknowledge the European Synchrotron Radiation Facility (ESRF) for provision of synchrotron radiation facilities under proposal numbers LS-3200 and LS-3419, and we would like to thank Hiram Castillo-Michel for assistance and support in using beamline ID21. The authors gratefully acknowledge the MVLS Cellular Analysis facility for their support and assistance and www.ToxoDB.org, without which this work would not have been possible. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Materials and Methods

Toxoplasma gondii and host cell maintenance

Toxoplasma gondii tachyzoites were grown in human foreskin fibroblasts (HFFs) cultured in Dulbecco's modified Eagle's medium (DMEM), supplemented with 3% (D3) heat-inactivated foetal bovine serum (FBS), 2 mM L-glutamine and 50 U/mL Penicillin/Streptomycin and maintained at 37 °C with 5% CO_2.

Toxoplasma gondii strain construction

ZFT-3HA$_{zft}$ and ZFT-3HA$_{sag1}$ parasite strains were generated as previously described (Sloan et al., 2025). To generate RHtdTomato pTUB8 ZFT-Ty-DHFR parasites, ZFT was amplified from cDNA and a Ty tag was inserted at the 3' end of the gene by PCR using primers P1 and P2 (**Table 3**). The tubulin promoter TUB8 plasmid was digested with EcoRI and PacI and then assembled with the ZFT-Ty PCR product to generate the new plasmid. The DHFR sequence was amplified using primers P3 and P4. The TUB-ZFT-Ty plasmid was then digested using BbcC1 enzyme then the DHFR sequence was inserted into the plasmid. TUB8-ZFT-Ty-DHFR plasmid was linearised using enzyme ScaI and the transfected into RHtdTomato parasites using a Gene Pulse BioRad electroporator. Parasites were selected using pyrimethamine and cloned by limiting dilution. Clones were confirmed using primers P5 and P6 (**Table 3**). For the generation of the F3ΔHX T7S4-ZFT strain, the T7S4 promoter was amplified from the G13m5 plasmid (Sheiner et al., 2011) using primers P7 and P8. F3ΔHX parasites were transfected using a Gene Pulse BioRad electroporator with 30 µg of the purified repair template T7S4-ZFT PCR product and 40 µg of a plasmid containing Cas9 guide targeting the 3' end of the genomic sequence of ZFT (**Table 2**). Parasites were selected with pyrimethamine and cloned by limiting dilution. Clones were screened for 3' integration using primers P9 and P10, and 5' integration using primers P11 and P12. For the generation of the F3ΔHX T7S4-ZFT-3HA$_{sag1}$ line, ZFT was tagged as previously described (Sloan et al., 2025) in F3ΔHX T7S4-ZFT parasites, selected for with chloramphenicol, and cloned by limiting dilution. The T7S4-ZFT SDHB-3HA and T7S4-ZFT ABCE1-3HA parasite strains were generated by

tagging SDHB and ABCE1 as previously described (Maclean et al., 2024, 2021) in the F3ΔHX T7S4-ZFT parasites, selected for with chloramphenicol, and cloned by limiting dilution. T7S4-ZFT SDHB-3HA clones were screened for 5' integration using primers P13 and P15, and 3' integration using primers P14 and P16. T7S-ZFT ABCE1-3HA clones were screened for 5' integration using primers P13 and P17, and 3' integration using primers P14 and P18. Parasite lines were selected using either chloramphenicol (Sigma Aldrich, C3175), made up in MiliQ water and used at 40 µg/ml or pyrimethamine (Sigma Aldrich, 46706), made up in ethanol and used at 2 µM. All sgRNA sequences (**Table 2**) and primers (**Table 3**) used in the study are listed below.

Chemicals

Ferric ammonium citrate (FAC) (Sigma Aldrich, F5879) and Deferoxamine mesylate salt (DFO) (Sigma Aldrich, D9533), were made up in 1X phosphate buffered saline (PBS) and used as indicated. Zinc sulphate ($ZnSO_4$) (Sigma Aldrich, 307491) and 2-Mercaptopyridine N-oxide sodium salt (Sigma Aldrich, H3261), were made up in distilled water and used at indicated concentrations. N,N,N',N'-tetrakis-(2-pyridylmethyl) ethylenediamine (TPEN) (Merck, 616394), was made up in DMSO and used at 5 µM. Anhydrotetracycline hydrochloride (ATc) (abcam, ab145350), made up in ethanol and used at 0.5 µM, unless indicated.

Immunofluorescence Assays

Indirect immunofluorescence assays (IFA) were performed on infected HFF monolayers. Intracellular *T. gondii* parasites were fixed with 4% paraformaldehyde (PFA) for 20 minutes at room temperature and washed with 1X PBS. Cells were permeabilized and blocked in blocking buffer (PBS/0.2% Triton X-100/2% Bovine Serum Albumin (BSA)) for 30 minutes at room temperature, or at 4°C overnight. Coverslips were then incubated in a wet chamber with primary antibodies (rat anti-HA 1:500 (Merk 11867423001), rabbit anti-GAP45 (Plattner et al., 2008) 1:2000, a kind gift from Dominique Soldati-Favre, rabbit anti-IMC1 (Wichroski et al., 2002) 1:2000, rat anti-SAG1 1:500 (Abcam), rabbit anti-BAG1 (Kannan et al., 2019; Smith et al., 2021) 1:500, a gift from Lilach Sheiner, rabbit anti-TOM40 (Van Dooren et al., 2016) 1:1000, a kind gift from Dr. Giel Van Dooren (ANU) and mouse anti-CPN60 (Agrawal et al., 2009) 1:1000, a kind gift from Lilach Sheiner, in blocking buffer for 1 hour at room temperature. Coverslips were then washed three times with PBST (0.2% Triton X-100 in PBS). After washing, slides were incubated in a wet chamber with the secondary antibodies (Goat anti-rat AlexaFluor-594 1:1000 (Invitrogen), goat anti-mouse AlexaFluor-488 1:1000 (Invitrogen), goat anti-rabbit AlexaFluor-488 1:1000 (Invitrogen), goat anti-guinea pig-488 1:1000 (Invitrogen) or Dolichos Biflorus-488 (DBL) 1:1000, in blocking buffer for 1 h at room temperature in the dark. Coverslips were then washed with PBST (0.2% Triton X-100 in PBS), and mounted using Fluoromount with 4',6-diamidino-2-phenylindole (DAPI), to stain DNA (Southern Biotech, 0100-20).

FerroOrange Live Cell Imaging

To analyse the levels of iron in the parasites, FerroOrange (Dojindo, F374-10) was used. Parasites were untreated, treated with ATc, or 200 µM FAC for 48 h. Parasites were released, pelleted and resuspended in DMEM supplemented with 1 µM FerroOrange and 10 µM Hoechst 33342 (ThermoFisher Scientific, H3570), put onto poly-l-lysine treated live cell dishes (Cellvis, D29-20-1.5-N) and incubated at 37 °C for 10 minutes before imaging. Images were obtained using an inverted DiM8 microscope (Leica Microsystems) using LAS X Core software and processed using FIJI software. Intensity was quantified using FIJI (Schindelin et al., 2012).

FluoZin-3AM Live Cell Imaging

To analyse the levels of zinc in the parasites, a Zn^{2+} selective indicator FluoZin-3AM (ThermoFisher Scientific, F14201) was used. Parasites were untreated, treated with ATc, or 50 µM $ZnSO_4$ for 24 h. Parasites were released, pelleted and resuspended in DMEM supplemented with 2 µM FluoZin-3 and 10 µM Hoechst 33342 (ThermoFisher Scientific, H3570), put onto poly-l-lysine treated live cell dishes (Cellvis, D29-20-1.5-N) and incubated at 37 °C for 10 minutes before imaging. Images were obtained using an inverted DiM8 microscope (Leica Microsystems) using LAS X Core software and processed using FIJI software. Intensity was quantified using FIJI (Schindelin et al., 2012).

RT-qPCR

To validate the T7S4-ZFT parasites, RT-qPCR was performed to confirm successful knockdown of ZFT mRNA levels. HFFs were infected with T7S4-ZFT with and without ATc for 24, 48, 72 and 96 h before being mechanically lysed and filtered through a 3 mm filter to remove any host cell debris. Parasites were then harvested by centrifugation at 2000 x g for 10 minutes, supernatant was removed, pellet was washed once in sterile PBS before spinning down in a microcentrifuge at 7000 x rpm for 10 minutes. Supernatant was removed and pellets were frozen at -20 °C overnight. RNA was then extracted from the pellets using the RNeasy Mini Kit (Qiagen, 74104). 1 µg RNA was then treated with DNAse I 1 U/µL (ThermoFisher Scientific, 18068015) at room temperature for 15 minutes before inactivating the DNAse with 25 mM EDTA followed by DNase denaturation at 65 °C for 10 minutes. Using 1 mg of DNase I treated RNA, cDNA synthesis was then performed using the High-Capacity cDNA Reverse Transcription Kit (Applied Biosystems, A48571) according to manufacturer's instructions. RT-qPCR was carried out on the Applied Biosystems 7500 Real Time PCR machine using PowerSYBR green PCR master mix (ThermoFisher Scientific 4367659) and 2 ng of 1:10 diluted cDNAs per reaction. Primers P21 and P22 were used for actin as a housekeeping control gene and primers P23 and P24 were used for ZFT (**Table 3**). The results are from three independent biological experiments, each performed in technical triplicate. Relative fold changes for - ATc vs. + ATc samples were calculated using the Pfaffl method (Pfaffl, 2001). Graphpad Prism 10 was used to perform statistical analysis.

Transmission electron microscopy

Infected cells were fixed with 2.5% glutaraldehyde and 4% paraformaldehyde in 0.1M cacodylate buffer. Following serial washes in 0.1M cacodylate buffer, the material was post-fixed in 1% OsO_4 and 1.25% $K4[Fe(CN_6)]$ for 1 hour in the dark, and contrasted en bloc with 0.5% aqueous uranyl acetate for 1 hour at room temperature in the dark. The samples were then dehydrated in ascending acetone series and embedded in epoxy resin. Ultra-thin sections (60nm) were collected using a Leica Ultramicrotome UC6 and observed in a Jeol 1200EX transmission electron microscope (Jeol, Japan).

Plaque Assays

200, 500 or 1000 parental or ZFT-Ty/T7S4-ZFT parasites were applied onto confluent monolayers of HFF cells in a 6-well plate in D3 media, supplemented when indicated with 200 µM FAC, 25 µM $ZnSO_4$, 2 µM zinc ionophore (2-Mercaptopyridine N-oxide sodium salt (Sigma Aldrich, H3261)), or both 200 µM FAC and 25 µM $ZnSO_4$, in the presence or absence of 0.5 µM ATc, unless stated otherwise. Parasites were allowed to replicate for 7 days undisturbed, washed 3x with PBS, fixed with ice cold 100 % methanol and stained using crystal violet stain (12.5 g crystal violet in 125 ml ethanol, diluted in 500 ml 1% ammonium oxalate) for 2-3 hours at room temperature before washing with distilled H_2O, drying and imaging. For ATc washout experiments, ATc was removed from cells after 3 days. Cells were then washed 2x with PBS and fresh D3 media was added to the cells. Parasites were allowed to replicate for a further 6 days undisturbed, then fixed and stained as described above.

Fluorescent Growth Assays

Growth assays of fluorescent parasites were performed as previously (Aghabi et al., 2023). Briefly, host cells were infected with 1,000 either RHtdTomato (Elati et al., 2023) or RHtdTom TUB8 ZFT-Ty tachyzoites per well and allowed to invade for 2 hours at 37 °C. The media was then removed and supplemented with either FAC, DFO, $ZnSO_4$ or Ars (at indicated concentrations) and incubated at 37 °C with 5% CO_2 for 4 days. Total parasite fluorescence was recorded at 4 days using a PHERAstar FS microplate reader (BMG LabTech). All experiments were performed in triplicate with at least three independent biological replicates. Uninfected wells served as blanks and fluorescence values were normalised to infected, untreated wells. Dose-response curves and EC_{50} were calculated using GraphPad Prism 10.

Western blotting

Parasites were mechanically released, filtered and collected by centrifugation at 1500 x g for 10 minutes, and lysed with RIPA lysis buffer (150 mM sodium chloride, 1 % Triton X-100, 0.5 % sodium deoxycholate, 0.1 % sodium dodecyl sulphate (SDS) and 50 mM Tris, pH 8.0) for 30 minutes on ice. Samples were then resuspended with 5X Laemmli buffer (10% SDS, 50% glycerol, 300 mM Tris-HCL pH 6.8 and 0.05% bromophenol blue), boiled at 95 °C for 5 (unless stated otherwise) and separated on a 10% SDS-PAGE gel, initially at 120V for 20 minutes then 175V for 60 minutes. PageRuler Prestained Protein Ladder (ThermoFisher Scientific, 26619) was used as a molecular weight marker. Proteins were then wet transferred to nitrocellulose

membrane (0.2 µM, Cytiva, 10600004) in Towbin buffer (0.025 M Tris, 0.192 M glycine, 10 % methanol) for 60 minutes at 250 mA and blocked for at least 1 h at room temperature in blocking buffer (5 % milk in 0.1 % Tween/PBS). Blots were then stained with primary antibodies overnight at 4°C (rat anti-HA, Merk 11867423001 1:500, mouse anti-Ty (Bastin et al., 1996) 1:1000, and guinea pig anti-CDPK1 (Waldman et al., 2020) 1:10,000, both kind gifts from Sebastian Lourido,), followed by secondary fluorescent antibodies at room temperature for 1 h (rabbit anti-rat HRP 1:5000 (abcam, ab6845), goat anti-mouse HRP 1:10,000 (Promega, W4021), goat anti-mouse IRDye 680 1:10,000 (Li-Cor) and goat anti-guinea pig IRDye 680 1:10,0000 (Li-Cor)). ZFT-3HA$_{zft}$ was detected using Pierce ECL Western Blotting Substrate (ThermoFisher Scientific) and imaged using the Invitrogen iBright 1500 Imaging System. CDPK1 was detected using the LI-COR Odyssey LCX system.

For Blue native (BN) PAGE, 10^7 parasites were mechanically released, filtered and harvested by centrifugation at 1500 x g for 10 minutes. The parasite pellet was then resuspended in a BN lysis buffer (1% DDM, 750 mM aminocaproic acid, 0.5 mM EDTA and 50 mM Bis-Tris-HCL pH 7.0), and incubated on ice for 30 minutes before being centrifuged at 16,000 x g for 30 minutes at 4 °C. The supernatant containing the solubilized membrane proteins was then added to a sample buffer containing coomassie G250 (NativePAGE), with the final concentration being 0.0625% Coomassie G250 and 0.25% DDM. 10^7 parasites were loaded onto a 3-12% Native PAGE Bis-Tris gel (ThermoFisher Scientific, NP0301PK2). NativeMark^{TM} was used as the molecular marker. Proteins were then transferred onto a PVDF membrane (0.45 µM, Cytiva, 10600023) via wet transfer in Towbin buffer for 60 minutes at 100 V. Immunoblotting and chemiluminescence detection was performed as above.

Seahorse Assay / Extracellular flux analysis
The *Toxoplasma* Seahorse assay was performed as previously described (Hayward et al., 2022). Briefly, parasites were incubated in the absence or presence of ATc for 48 h, extracellular parasites were removed by washing 1X PBS, and intracellular parasites were mechanically released, filtered and harvested by spinning at 2000 x g for 10 minutes. Parasites were then resuspended Seahorse XF media (Agilent) supplemented with 5 mM glucose and 1 mM glutamine at 37 °C. 1.5×10^6 parasites in a final volume of 175 µL were added to each well of a poly-L-lysine-treated seahorse XFp cell culture miniplate (Aglient). Parasites were then centrifuged at 300 x g for 5 minutes to ensure sedimentation before the basal oxygen consumption rate (OCR) and the extracellular acidification rate (ECAR) was quantified using a Seahorse XF HS Mini Analyser (Agilent Technologies). The OCR following the addition of 1 µM atovaquone (Scientific, 17575545) was considered as the non-mitochondrial oxygen consumption rate and subtracted as background. The basal and maximal oxygen consumption rates of the parasites were calculated by subtracting the background from the raw OCR values before and after the addition of 1 µM Carbonyl cyanide 4-(trifluoromethoxy) phenylhydrazone (FCCP) (Merck, C2920) respectively. Basal ECAR values were taken as raw values. This experiment was performed in three independent biological replicates, each with three technical triplicates.

ICP-MS

Total parasite associated iron and zinc were quantified using ICP-MS. Confluent HFFs cells in T150 flasks were infected with T7S4-ZFT parasites for 24 h before the addition of ATc for a further 24 h, or infected with T7S4-ZFT parasites + ATc for 48 h to ensure efficient knockdown. Untreated (-ATc) parasites were used as a control. Parasites were harvested by scraping, syringing and filtering through either 0.3 µM filters or PD-10 columns (Cytiva, 17085101), and quantified using a haemocytometer. Parasites were then washed three times with PBS supplemented with 6% Chelex 100 sodium form (Merck, C7901), then once with 1 mM EDTA before being digested in 65% nitric acid (Merck, 225711) at 85°C for 5 h. Digested samples were then diluted 1:10 in 2% nitric acid with 0.025% Triton X-100. At least three biological replicates were collected, with three technical replicates per biological sample. Samples were quantified at the Scottish Universities Environmental Research Centre (SUREC) and processed by Dr Valerie Olive. Briefly, samples were run on an Agilent 7500ce ICP-MS, fitted with a double pass spray chamber and a self-aspire nebulizer tuned for 0.1 ml/min. A 3 ppb Indium (^{115}In) was introduced as internal standard to correct for the sensitivity drop caused by the sample matrix. Cu and Zn were taken on the masses 63 and 66 respectively and the results are an average of 10 repeats. The calibration equation was calculated from 3 points at 5, 10 and 20 ppb using Cu and Zn mono-element solutions Cu and Zn specpure ®. Iron was taken at the mass 56 using a collision cell with a flow of Hydrogen at 4 ml/min to suppress the interference of the $^{40}Ar^{16}O$. The calibration equation was calculated using a 200 ppb and 300 ppb solution of mono-element Fe specpure ®, results are an average of 20 repeats.

X-ray Fluorescence Microscopy (XFM)

T7S4-ZFT parasites were treated in the presence or absence of ATc for 24 h or 48 h. Parasites from a fully lysed dish were filtered, pelleted and resuspended in 5 mL of extracellular buffer (116 mM NaCl, 5.4 mM KCL, 0.8 mM $MgSO_4$, 5.5 mM glucose, 50 mM HEPES, pH 7.2). Approximately 500 µL of resuspended parasites were then added onto each poly-l-lysine 2 x 2 mm silicon nitride (SiN) window (Norcada, X5200D) and spun down gently at 200 x g for 2 minutes then left at room temperature for 20 minutes.

Following adhesion/infection, cells were then washed twice with 1X chelexed PBS then fixed in 4% PFA made up in 1X chelexed PBS for 20 minutes at room temperature. Cells were then washed twice in 1X chelexed PBS, twice in 100 mM ammonium acetate and finally twice in MiliQ H_2O before drying overnight. Elemental mapping was performed at the European Synchrotron Radiation Facility in Grenoble, France at beamline ID21 under proposal number LS-3419. Data is available from https://doi.esrf.fr/10.15151/ESRF-DC-2119497225. Analysis was performed using PyMca5.9.2 (Solé et al., 2007) and FIJI (Schindelin et al., 2012). The 48 h time point is representative of two independent biological experiments, whereas the 24 h time point is representative of one independent biological experiment. We would like to thank Hiram Castillo Michel for assistance and support.

Yeast strain construction

Full length ZFT was cloned into the yeast expression vector pDR195 (AddGene, #36028) using primers P19 and P20 (**Table 3**). Wildtype (DY1457) and Δzrt1/2 (Ullah et al., 2018) strains, a kind gift from Jim Dunwell (University of Reading), were transformed as previously described (Gietz and Schiestl, 2007). Positive transformations were selected on synthetic dropout agar lacking uracil (SD-Ura). Colonies were grown overnight in SD-Ura media and spotted onto SD-Ura agar, or SD-Ura agar supplemented with 5 mM EGTA, and incubated for 4 days.

Tables

gRNA Sequences	
Name	**Sequence (5' – 3')**
ZFT 3'	TGAACTGTTGCACAACCTAC
ZFT 5'	CCTTGAGACCTTTCCTCCCG
SDHB 3'	CGAGCCGAGGTTACGCAGCG
ABCE1 3'	GCTCGACGACGCTTGAAGCG

Table 2: A list of the gRNA sequences used in this study.

Primer Sequences		
Name	**Sequence (5' – 3')**	**Notes**
P1	ctcgttggcattttttcttgATGGCGACGGTCGCAGAC	ZFT amplification from cDNA Fw
P2	tgagcacaacggtgattaat*ttaatcgagcgggtcctggttcgtgtggacctc*AGCGTCTAGAGCCAGATCCAC	ZFT amplification from cDNA and Ty tag Rv
P3	CTCGGTCTTACCCTGGTGGATCTGGCTCTAGACGCaGAGGTCCACACGAACCAGGACCCGCTCGATAGCGGAGCTACTAACTTCAGC	P2A-DHFR amplification Fw
P4	GTGATTAATTTAATCGAGCGGGTCCTGGTTCGTGTGGACCgctaGACAGCCATCTCCATCTGGAT	P2A-DHFR amplification Rv
P5	ctcgttggcattttttcttgatggcgacggtcgcagac	TUB8 ZFT-Ty DHFR integration verification Fw
P6	tgagcacaacggtgattaatttaatcgagcgggtcctggttcgtgtggacctcagcgtctagagccagatccac	TUB8 ZFT-Ty DHFR integration verification Rv
P7	accgaacagggcgacgatttttcccttgagacctttcctCTTCGCCAGGCTGTAAATCC	ZFT T7S4 Fw
P8	agctggcagcagacatcacatcgtctgcgaccgtcgccatTGGTTGAAGACAGACGAAAGCAGTTG	ZFT T7S4 Rv
P9	cctgataggttctggacctccctcgc	ZFT T7S4 5' integration Fw
P10	cgtagagcagaaacgcactactaaag	ZFT T7S4 5' integration Rv
P11	gcaaagccagagggatatatgctcc	ZFT T7S4 3' integration Fw

P12	cgaggtcgacgtatcgataagcttacg	ZFT T7S4 3' integration Rv
P13	AAATAGGATGGTACATCCCCC	HA integration test Rv
P14	gaaacacacttcgtgcagc	CAT integration test Fw
P15	CTTTTCTTTTGTGGGTTGTGC	SDHB-3HA Fw
P16	CGACATTTGCATGTACATACC	SDHB-3HA UTR Rv
P17	ATCTCGTCAGCGGCATGAACC	ABCE1-3HA Fw
P18	CATACACGGACACAGATACGC	ABCE1-3HA UTR Rv
P19	aaaaaaaatatacccccagcctcgagATGGCCACCGTCGCTGATGAC	Amplification of ZFT for yeast expression Fw
P20	aagtccaaagctggatccgcctcgagTTAGTCAAGAGGGTCTTGGTTAG	Amplification of ZFT for yeast expression Rv
P21	GGGACGACATGGAGAAAATC	Actin RT-qPCR Fw
P22	AGAAAGAACGGCCTGGATAG	Actin RT-qPCR Rv
P23	CGGACAACCGAATAGTGGCT	ZFT RT-qPCR Fw
P24	CGCCAACGATCATTCCTTCG	ZFT RT-qPCR Rv

Table 3: List of primer sequences used in this study

Supplemental Figure Legends

Figure S1

a. Schematic of F3ΔHX ZFT-3HA*sag1* strain construction to tag ZFT with a 3xHA epitope tag at the C-terminal under the endogenous promoter with the sag1 3' UTR. **b.** Immunofluorescence of ZFT-3HA*sag1* demonstrating dynamic localisation throughout the lytic life cycle, showing peripheral and basal characterisation. Scale bars 5 μm. **c.** Immunofluorescence assay of ZFT-3HA*zft* 24 h post infection in untreated and low iron (100 μM DFO) conditions.

Figure S2

a. PCR of ZFT-Ty confirming the presence of the gene. **b.** Fluorescent growth curves showing the percentage of parasite survival in the presence of excess iron. **c.** Fluorescent growth curve showing the percentage of parasite survival in the presence of sodium arsenite. **d.** Graphs showing mean EC_{50} for Ars for RHtdTomato and ZFT-Ty parasites. Each points represents an independent experiment, bars are at the mean of n=3, ± SD. *p* values from two-tailed unpaired *t* test. **e.** Fluorescent growth curves showing the percentage of parasite survival where iron is depleted using the chelator DFO.

Figure S3

a. PCR of T7S4-ZFT confirming successful integration of the DHFR-T7S4 cassette at both the 5' and 3' end. **b.** PCR confirming successful tagging of ZFT with a 3HA epitope tag in the T7S4-ZFT parasite line. **c.** Plaque assays showing that the addition of 200 μM FAC and 0.25 μM ATc instead of the standard 0.5 μM ATc does not rescue the inability of ZFT knockdown parasites from forming plaques, 6 days. **d.** Quantification of plaque assay showing the number of plaques normalised to parental in untreated conditions. Bars are at the mean of three independent experiments, ± SD. *p* values from two-way ANOVA, Tukey corrected for multiple comparisons. **e.** Quantification of plaque assay showing the number of plaques normalised to parental in untreated conditions. Bars are at the mean of three independent experiments, ± SD. *p* values from two-way ANOVA, Tukey corrected for multiple comparisons.

Figure S4

a. Schematic of F3ΔHX T7S4-ZFT SDHB-3HA strain construction to endogenously tag SDHB with 3xHA epitope tags at the C-terminal under the endogenous promoter. **b.** PCR confirming successful integration of the 3HA epitope tag as well as the CAT selection cassette into the endogenous locus in the T7S4-ZFT SDHB-3HA line. Non-specific bands marked with asterisks. **c.** Schematic of F3ΔHX T7S4-ZFT ABCE1-3HA strain construction to endogenously tag ABCE1 with 3xHA epitope tags at the C-terminal under the endogenous promoter. **d.** PCR confirming successful integration of the 3HA epitope tag as well as the CAT selection cassette into the endogenous locus in the T7S4-ZFT ABCE1-3HA line. Non-specific bands marked with asterisks. **e.** Immuno-electron microscopy images of parental parasites in the presence of 100 μM DFO for five days, indicating enlarged L-vesicles in the parasites. Scale bar 500 nm.

Figure S5
a. Plaque assays showing that the addition of 2 µM Zn Ionophore does not rescue the inability of ZFT knockdown parasites from forming plaques. **b.** Quantification of plaque assay showing number of plaques, normalised to parental + Zn Ionophore. Bars are at the mean of three independent experiments, ± SD. p values from two-way ANOVA, Tukey corrected for multiple comparisons. **c.** Quantification of plaque assay showing number of plaques, normalised to parental + FAC and $ZnSO_4$. Bars are at the mean of three independent experiments, ± SD. p values from two-way ANOVA, Tukey corrected for multiple comparisons. **d.** Dose-response curves showing the percentage of both parental and ZFT-Ty parasite survival in the presence of excess zinc. **e**. Immunofluorescence of ZFT-3HA$_{zft}$ untreated and at 24 h treatment with 50 µM $ZnSO_4$ showing no change in ZFT localisation. Scale bar 5 µm.

Figure S6
Quantification of XFM levels of P (mM) (**a**) and S (mM) (**b**) per parasite following ZFT knockdown at both 24 h and 48 h ATc. Bar at mean ± SD. p values from one-way ANOVA, Tukey corrected for multiple comparisons.

References

Aghabi, D., Sloan, M., Gill, G., Hartmann, E., Antipova, O., Dou, Z., Guerra, A.J., Carruthers, V.B., Harding, C.R., 2023. The vacuolar iron transporter mediates iron detoxification in Toxoplasma gondii. Nat Commun 14, 3659. https://doi.org/10.1038/s41467-023-39436-y

Agrawal, S., van Dooren, G.G., Beatty, W.L., Striepen, B., 2009. Genetic evidence that an endosymbiont-derived endoplasmic reticulum-associated protein degradation (ERAD) system functions in import of apicoplast proteins. J Biol Chem 284, 33683–33691. https://doi.org/10.1074/jbc.M109.044024

Ahern, M.E., Bafaro, E.M., Cowan, A., Dempski, R.E., 2020. Quantifying the Oligomeric State of hZIP4 on the Surface of Cells.

Al-sandaqchi, A.T., Brignell, C., Collingwood, J.F., Geraki, K., Mirkes, E.M., Kong, K., Castellanos, M., May, S.T., Stevenson, C.W., Elsheikha, H.M., 2018. Metallome of cerebrovascular endothelial cells infected with *Toxoplasma gondii* using μ-XRF imaging and inductively coupled plasma mass spectrometry. Metallomics 10, 1401–1414. https://doi.org/10.1039/c8mt00136g

Amos, B., Aurrecoechea, C., Barba, M., Barreto, A., Basenko, E.Y., Bażant, W., Belnap, R., Blevins, A.S., Böhme, U., Brestelli, J., Brunk, B.P., Caddick, M., Callan, D., Campbell, L., Christensen, M.B., Christophides, G.K., Crouch, K., Davis, K., DeBarry, J., Doherty, R., Duan, Y., Dunn, M., Falke, D., Fisher, S., Flicek, P., Fox, B., Gajria, B., Giraldo-Calderón, G.I., Harb, O.S., Harper, E., Hertz-Fowler, C., Hickman, M.J., Howington, C., Hu, S., Humphrey, J., Iodice, J., Jones, A., Judkins, J., Kelly, S.A., Kissinger, J.C., Kwon, D.K., Lamoureux, K., Lawson, D., Li, W., Lies, K., Lodha, D., Long, J., MacCallum, R.M., Maslen, G., McDowell, M.A., Nabrzyski, J., Roos, D.S., Rund, S.S.C., Schulman, S.W., Shanmugasundram, A., Sitnik, V., Spruill, D., Starns, D., Stoeckert, C.J., Tomko, S.S., Wang, H., Warrenfeltz, S., Wieck, R., Wilkinson, P.A., Xu, L., Zheng, J., 2022. VEuPathDB: the eukaryotic pathogen, vector and host bioinformatics resource center. Nucleic Acids Research 50, D898–D911. https://doi.org/10.1093/nar/gkab929

Augusto, L., Martynowicz, J., Amin, P.H., Carlson, K.R., Wek, R.C., Sullivan, W.J., 2021. TgIF2K-B Is an eIF2α Kinase in Toxoplasma gondii That Responds to Oxidative Stress and Optimizes Pathogenicity. mBio 12, e03160-20. https://doi.org/10.1128/mBio.03160-20

Aw, Y.T.V., Seidi, A., Hayward, J.A., Lee, J., Makota, F.V., Rug, M., Van Dooren, G.G., 2021. A key cytosolic iron–sulfur cluster synthesis protein localizes to the mitochondrion of *Toxoplasma gondii*. Molecular Microbiology 115, 968–985. https://doi.org/10.1111/mmi.14651

Baker, N., Patel, J., Khacho, M., 2019. Linking mitochondrial dynamics, cristae remodeling and supercomplex formation: How mitochondrial structure can regulate bioenergetics. Mitochondrion 49, 259–268. https://doi.org/10.1016/j.mito.2019.06.003

Barthelme, D., Dinkelaker, S., Albers, S.-V., Londei, P., Ermler, U., Tampé, R., 2011. Ribosome recycling depends on a mechanistic link between the FeS cluster domain and a conformational switch of the twin-ATPase ABCE1. Proc. Natl. Acad. Sci. U.S.A. 108, 3228–3233. https://doi.org/10.1073/pnas.1015953108

Barthelme, D., Scheele, U., Dinkelaker, S., Janoschka, A., MacMillan, F., Albers, S.-V., Driessen, A.J.M., Stagni, M.S., Bill, E., Meyer-Klaucke, W., Schuönemann, V., Tampé;, R., 2007. Structural Organization of Essential Iron-Sulfur Clusters in the Evolutionarily Highly Conserved ATP-binding Cassette Protein ABCE1. Journal of Biological Chemistry 282, 14598–14607. https://doi.org/10.1074/jbc.M700825200

Barylyuk, K., Koreny, L., Ke, H., Butterworth, S., Crook, O.M., Lassadi, I., Gupta, V., Tromer, E., Mourier, T., Stevens, T.J., Breckels, L.M., Pain, A., Lilley, K.S., Waller, R.F., 2020. A Comprehensive Subcellular Atlas of the Toxoplasma Proteome via hyperLOPIT Provides Spatial Context for Protein Functions. Cell Host & Microbe 28, 752-766.e9. https://doi.org/10.1016/j.chom.2020.09.011

Bastin, P., Bagherzadeh, A., Matthews, K.R., Gull, K., 1996. A novel epitope tag system to study protein targeting and organelle biogenesis in Trypanosoma brucei. Molecular and Biochemical Parasitology 77, 235–239. https://doi.org/10.1016/0166-6851(96)02598-4

Bergmann, A., Floyd, K., Key, M., Dameron, C., Rees, K.C., Thornton, L.B., Whitehead, D.C., Hamza, I., Dou, Z., 2020. Toxoplasma gondii requires its plant-like heme biosynthesis pathway for infection. PLoS Pathog 16, e1008499. https://doi.org/10.1371/journal.ppat.1008499

Bin, B.-H., Fukada, T., Hosaka, T., Yamasaki, S., Ohashi, W., Hojyo, S., Miyai, T., Nishida, K., Yokoyama, S., Hirano, T., 2011. Biochemical Characterization of Human ZIP13 Protein. Journal of Biological Chemistry 286, 40255–40265. https://doi.org/10.1074/jbc.M111.256784

Bisetegn, H., Debash, H., Ebrahim, H., Mahmood, N., Gedefie, A., Tilahun, M., Alemayehu, E., Mohammed, O., Feleke, D.G., 2023. Global seroprevalence of *Toxoplasma gondii* infection among patients with mental and neurological disorders: A systematic review and meta-analysis. Health Science Reports 6, e1319. https://doi.org/10.1002/hsr2.1319

Blaby-Haas, C.E., Merchant, S.S., 2013. Iron sparing and recycling in a compartmentalized cell. Current Opinion in Microbiology 16, 677–685. https://doi.org/10.1016/j.mib.2013.07.019

Cerutti, A., Blanchard, N., Besteiro, S., 2020. The Bradyzoite: A Key Developmental Stage for the Persistence and Pathogenesis of Toxoplasmosis. Pathogens 9, 234. https://doi.org/10.3390/pathogens9030234

Chasen, N.M., Stasic, A.J., Asady, B., Coppens, I., Moreno, S.N.J., 2019. The Vacuolar Zinc Transporter TgZnT Protects Toxoplasma gondii from Zinc Toxicity. mSphere 4, e00086-19. https://doi.org/10.1128/mSphere.00086-19

Connolly, E.L., Fett, J.P., Guerinot, M.L., 2002. Expression of the IRT1 Metal Transporter Is Controlled by Metals at the Levels of Transcript and Protein Accumulation. Plant Cell 14, 1347–1357. https://doi.org/10.1105/tpc.001263

Conte, S.S., Walker, E.L., 2011. Transporters Contributing to Iron Trafficking in Plants. Molecular Plant 4, 464–476. https://doi.org/10.1093/mp/ssr015

Desmonts, G., Couvreur, J., 1974. Congenital Toxoplasmosis: A Prospective Study of 378 Pregnancies. N Engl J Med 290, 1110–1116. https://doi.org/10.1056/NEJM197405162902003

Dufner-Beattie, J., Langmade, S.J., Wang, F., Eide, D., Andrews, G.K., 2003. Structure, Function, and Regulation of a Subfamily of Mouse Zinc Transporter Genes. Journal of Biological Chemistry 278, 50142–50150. https://doi.org/10.1074/jbc.M304163200

Elati, H.A.A., Goerner, A.L., Martorelli Di Genova, B., Sheiner, L., De Koning, H.P., 2023. Pyrimidine salvage in Toxoplasma gondii as a target for new treatment. Front. Cell. Infect. Microbiol. 13, 1320160. https://doi.org/10.3389/fcimb.2023.1320160

Ferrer, P., Tripathi, A.K., Clark, M.A., Hand, C.C., Rienhoff, H.Y., Sullivan, D.J., 2012. Antimalarial Iron Chelator, FBS0701, Shows Asexual and Gametocyte Plasmodium falciparum Activity and Single Oral Dose Cure in a Murine Malaria Model. PLoS ONE 7, e37171. https://doi.org/10.1371/journal.pone.0037171

Frénal, K., Polonais, V., Marq, J.-B., Stratmann, R., Limenitakis, J., Soldati-Favre, D., 2010. Functional Dissection of the Apicomplexan Glideosome Molecular Architecture. Cell Host & Microbe 8, 343–357. https://doi.org/10.1016/j.chom.2010.09.002

Fujimaki, M., Furuya, N., Saiki, S., Amo, T., Imamichi, Y., Hattori, N., 2019. Iron Supply via NCOA4-Mediated Ferritin Degradation Maintains Mitochondrial Functions. Molecular and Cellular Biology 39, e00010-19. https://doi.org/10.1128/MCB.00010-19

Gaither, L.A., Eide, D.J., 2001. The Human ZIP1 Transporter Mediates Zinc Uptake in Human K562 Erythroleukemia Cells. Journal of Biological Chemistry 276, 22258–22264. https://doi.org/10.1074/jbc.M101772200

Galaris, D., Barbouti, A., Pantopoulos, K., 2019. Iron homeostasis and oxidative stress: An intimate relationship. Biochimica et Biophysica Acta (BBA) - Molecular Cell Research 1866, 118535. https://doi.org/10.1016/j.bbamcr.2019.118535

Gao, F., Dubos, C., 2021. Transcriptional integration of plant responses to iron availability. Journal of Experimental Botany 72, 2056–2070. https://doi.org/10.1093/jxb/eraa556

Gietz, R.D., Schiestl, R.H., 2007. High-efficiency yeast transformation using the LiAc/SS carrier DNA/PEG method. Nat Protoc 2, 31–34. https://doi.org/10.1038/nprot.2007.13

Gissot, M., Hovasse, A., Chaloin, L., Schaeffer-Reiss, C., Van Dorsselaer, A., Tomavo, S., 2017. An evolutionary conserved zinc finger protein is involved in *Toxoplasma gondii* mRNA nuclear export: A zinc finger protein is involved in mRNA nuclear export. Cellular Microbiology 19, e12644. https://doi.org/10.1111/cmi.12644

Guerinot, M.L., 2000. The ZIP family of metal transporters. Biochimica et Biophysica Acta (BBA) - Biomembranes 1465, 190–198. https://doi.org/10.1016/S0005-2736(00)00138-3

Gyimesi, G., Albano, G., Fuster, D.G., Hediger, M.A., Pujol-Giménez, J., 2019. Unraveling the structural elements of pH sensitivity and substrate binding in the human zinc transporter SLC39A2 (ZIP2). Journal of Biological Chemistry 294, 8046–8063. https://doi.org/10.1074/jbc.RA118.006113

Hare, D., Austin, C., Doble, P., 2012. Quantification strategies for elemental imaging of biological samples using laser ablation-inductively coupled plasma-mass spectrometry. Analyst 137, 1527. https://doi.org/10.1039/c2an15792f

Hayward, J., Rajendran, E., Makota, F., Bassett, B., Devoy, M., Neeman, T., Van Dooren, G., 2022. Real-Time Analysis of Mitochondrial Electron Transport Chain Function in Toxoplasma gondii Parasites Using a Seahorse XFe96 Extracellular Flux Analyzer. BIO-PROTOCOL 12. https://doi.org/10.21769/BioProtoc.4288

Hu, J., 2021. Toward unzipping the ZIP metal transporters: structure, evolution, and implications on drug discovery against cancer. The FEBS Journal 288, 5805–5825. https://doi.org/10.1111/febs.15658

Huet, D., Rajendran, E., Van Dooren, G.G., Lourido, S., 2018. Identification of cryptic subunits from an apicomplexan ATP synthase. eLife 7, e38097. https://doi.org/10.7554/eLife.38097

Huynh, C., Sacks, D.L., Andrews, N.W., 2006. A *Leishmania amazonensis* ZIP family iron transporter is essential for parasite replication within macrophage phagolysosomes. The Journal of Experimental Medicine 203, 2363–2375. https://doi.org/10.1084/jem.20060559

Imlay, J.A., Chin, S.M., Linn, S., 1988. Toxic DNA Damage by Hydrogen Peroxide Through the Fenton Reaction in Vivo and in Vitro. Science 240, 640–642. https://doi.org/10.1126/science.2834821

Jacques, I., Andrews, N.W., Huynh, C., 2010. Functional characterization of LIT1, the Leishmania amazonensis ferrous iron transporter. Molecular and Biochemical Parasitology 170, 28–36. https://doi.org/10.1016/j.molbiopara.2009.12.003

Jin, Q., Paunesku, T., Lai, B., Gleber, S., Chen, S., Finney, L., Vine, D., Vogt, S., Woloschak, G., Jacobsen, C., 2017. Preserving elemental content in adherent mammalian cells for analysis by synchrotron-based x-ray fluorescence microscopy. Journal of Microscopy 265, 81–93. https://doi.org/10.1111/jmi.12466

Kannan, G., Di Cristina, M., Schultz, A.J., Huynh, M.-H., Wang, F., Schultz, T.L., Lunghi, M., Coppens, I., Carruthers, V.B., 2019. Role of *Toxoplasma gondii* Chloroquine Resistance Transporter in Bradyzoite Viability and Digestive Vacuole Maintenance. mBio 10, e01324-19. https://doi.org/10.1128/mBio.01324-19

Kloehn, J., Harding, C.R., Soldati-Favre, D., 2021. Supply and demand—heme synthesis, salvage and utilization by Apicomplexa. The FEBS Journal 288, 382–404. https://doi.org/10.1111/febs.15445

Lee, W., Ha, J., Sugiyama, Y., 2020. Post-translational regulation of the major drug transporters in the families of organic anion transporters and organic anion–transporting polypeptides. Journal of Biological Chemistry 295, 17349–17364. https://doi.org/10.1074/jbc.REV120.009132

Li, L., Kaplan, J., 1998. Defects in the Yeast High Affinity Iron Transport System Result in Increased Metal Sensitivity because of the Increased Expression of Transporters with a Broad Transition Metal Specificity. Journal of Biological Chemistry 273, 22181–22187. https://doi.org/10.1074/jbc.273.35.22181

Li, Y., Shi, J., Wu, X., Luo, Z., Wang, F., Guo, Q., 2009. Tracing of intracellular zinc(II) fluorescence flux to monitor cell apoptosis by using FluoZin-3AM. Cell Biochemistry & Function 27, 417–423. https://doi.org/10.1002/cbf.1598

Lin, W., Chai, J., Love, J., Fu, D., 2010. Selective Electrodiffusion of Zinc Ions in a Zrt-, Irt-like Protein, ZIPB*. Journal of Biological Chemistry 285, 39013–39020. https://doi.org/10.1074/jbc.M110.180620

Loveridge, K.M., Sigala, P.A., 2024. Identification of a divalent metal transporter required for cellular iron metabolism in malaria parasites. Proc. Natl. Acad. Sci. U.S.A. 121, e2411631121. https://doi.org/10.1073/pnas.2411631121

Maclean, A.E., Bridges, H.R., Silva, M.F., Ding, S., Ovciarikova, J., Hirst, J., Sheiner, L., 2021. Complexome profile of Toxoplasma gondii mitochondria identifies divergent subunits of respiratory chain complexes including new subunits of cytochrome bc1 complex. PLoS Pathog 17, e1009301. https://doi.org/10.1371/journal.ppat.1009301

Maclean, A.E., Sloan, M.A., Renaud, E.A., Argyle, B.E., Lewis, W.H., Ovciarikova, J., Demolombe, V., Waller, R.F., Besteiro, S., Sheiner, L., 2024. The *Toxoplasma gondii* mitochondrial transporter ABCB7L is essential for the biogenesis of cytosolic and nuclear iron-sulfur cluster proteins and cytosolic translation. mBio 15, e00872-24. https://doi.org/10.1128/mbio.00872-24

Mahmoud, M.S., 1999. Effect of deferoxamine alone and combined with pyrimethamine on acute toxoplasmosis in mice. J Egypt Soc Parasitol 29, 791–803.

Mallo, N., Ovciarikova, J., Martins-Duarte, E.S., Baehr, S.C., Biddau, M., Wilde, M.-L., Uboldi, A.D., Lemgruber, L., Tonkin, C.J., Wideman, J.G., Harding, C.R., Sheiner, L., 2021. Depletion of a *Toxoplasma* porin leads to defects in mitochondrial morphology and contacts with the endoplasmic reticulum. Journal of Cell Science 134, jcs255299. https://doi.org/10.1242/jcs.255299

Markus, B.M., Bell, G.W., Lorenzi, H.A., Lourido, S., 2019. Optimizing Systems for Cas9 Expression in Toxoplasma gondii. mSphere 4, e00386-19. https://doi.org/10.1128/mSphere.00386-19

Miller, S., Ralle, M., 2024. Sample Preparation for X-Ray Fluorescence Microscopy of Iron Distribution in Biological Specimen. Methods Mol Biol 2839, 43–52. https://doi.org/10.1007/978-1-0716-4043-2_3

Olea-Flores, M., Kan, J., Carlson, A., Syed, S.A., McCann, C., Mondal, V., Szady, C., Ricker, H.M., McQueen, A., Navea, J.G., Caromile, L.A., Padilla-Benavides, T., 2022. ZIP11 Regulates Nuclear Zinc Homeostasis in HeLa Cells and Is Required for Proliferation and Establishment of the Carcinogenic Phenotype. Front. Cell Dev. Biol. 10, 895433. https://doi.org/10.3389/fcell.2022.895433

Pamukcu, S., Cerutti, A., Bordat, Y., Hem, S., Rofidal, V., Besteiro, S., 2021. Differential contribution of two organelles of endosymbiotic origin to iron-sulfur cluster synthesis and overall fitness in Toxoplasma. PLoS Pathog 17, e1010096. https://doi.org/10.1371/journal.ppat.1010096

Pang, C., Chai, J., Zhu, P., Shanklin, J., Liu, Q., 2023. Structural mechanism of intracellular autoregulation of zinc uptake in ZIP transporters. Nat Commun 14, 3404. https://doi.org/10.1038/s41467-023-39010-6

Paunesku, T., Gordon, A.C., White, S., Harris, K., Antipova, O., Maxey, E., Vogt, S., Smith, A., Daddario, L., Procissi, D., Larson, A., Woloschak, G.E., 2021. Use of X-Ray Fluorescence Microscopy for Studies on Research Models of Hepatocellular Carcinoma. Front. Public Health 9, 711506. https://doi.org/10.3389/fpubh.2021.711506

Pfaffl, M.W., 2001. A new mathematical model for relative quantification in real-time RT-PCR. Nucleic Acids Res 29, e45. https://doi.org/10.1093/nar/29.9.e45

Plattner, F., Yarovinsky, F., Romero, S., Didry, D., Carlier, M.-F., Sher, A., Soldati-Favre, D., 2008. Toxoplasma Profilin Is Essential for Host Cell Invasion and TLR11-Dependent Induction of an Interleukin-12 Response. Cell Host & Microbe 3, 77–87. https://doi.org/10.1016/j.chom.2008.01.001

Rajapakse, D., Curtis, T., Chen, M., Xu, H., 2017. Zinc Protects Oxidative Stress-Induced RPE Death by Reducing Mitochondrial Damage and Preventing Lysosome Rupture. Oxidative Medicine and Cellular Longevity 2017, 6926485. https://doi.org/10.1155/2017/6926485

Raventos-Suarez, C., Pollack, S., Nagel, R.L., 1982. Plasmodium Falciparum: Inhibition of in Vitro Growth by Desferrioxamine. The American Journal of Tropical Medicine and Hygiene 31, 919–922. https://doi.org/10.4269/ajtmh.1982.31.919

Renaud, E.A., Maupin, A.J.M., Berry, L., Bals, J., Bordat, Y., Demolombe, V., Rofidal, V., Vignols, F., Besteiro, S., 2025. The HCF101 protein is an important component of the cytosolic iron–sulfur synthesis pathway in Toxoplasma gondii. PLoS Biol 23, e3003028. https://doi.org/10.1371/journal.pbio.3003028

Renaud, E.A., Maupin, A.J.M., Bordat, Y., Graindorge, A., Berry, L., Besteiro, S., 2024. Iron depletion has different consequences on the growth and survival of *Toxoplasma gondii* strains. Virulence 15, 2329566. https://doi.org/10.1080/21505594.2024.2329566

Renaud, E.A., Pamukcu, S., Cerutti, A., Berry, L., Lemaire-Vieille, C., Yamaryo-Botté, Y., Botté, C.Y., Besteiro, S., 2022. Disrupting the plastidic iron-sulfur cluster biogenesis pathway in Toxoplasma gondii has pleiotropic effects irreversibly impacting parasite viability. Journal of Biological Chemistry 298, 102243. https://doi.org/10.1016/j.jbc.2022.102243

Sahu, T., Boisson, B., Lacroix, C., Bischoff, E., Richier, Q., Formaglio, P., Thiberge, S., Dobrescu, I., Ménard, R., Baldacci, P., 2014. ZIPCO , a putative metal ion transporter, is crucial for *Plasmodium* liver-stage development. EMBO Mol Med 6, 1387–1397. https://doi.org/10.15252/emmm.201403868

Schaefer-Ramadan, S., Barlog, M., Roach, J., Al-Hashimi, M., Bazzi, H.S., Machaca, K., 2019. Synthesis of TPEN variants to improve cancer cells selective killing capacity. Bioorganic Chemistry 87, 366–372. https://doi.org/10.1016/j.bioorg.2019.03.045

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D.J., Hartenstein, V., Eliceiri, K., Tomancak, P., Cardona, A., 2012. Fiji: an open-source platform for biological-image analysis. Nat Methods 9, 676–682. https://doi.org/10.1038/nmeth.2019

Segura, M., Madrid, Y., Cámara, C., 2003. Elimination of calcium and argon interferences in iron determination by ICP-MS using desferrioxamine chelating agent immobilized in sol–gel and cold plasma conditions. J. Anal. At. Spectrom. 18, 1103–1108. https://doi.org/10.1039/B301719M

Semenovskaya, K., Lévêque, M.F., Berry, L., Bordat, Y., Dubremetz, J., Lebrun, M., Besteiro, S., 2020. TgZFP2 is a novel zinc finger protein involved in coordinating mitosis and budding in *Toxoplasma*. Cellular Microbiology 22. https://doi.org/10.1111/cmi.13120

Sheiner, L., Demerly, J.L., Poulsen, N., Beatty, W.L., Lucas, O., Behnke, M.S., White, M.W., Striepen, B., 2011. A Systematic Screen to Discover and Analyze Apicoplast Proteins Identifies a Conserved and Essential Protein Import Factor. PLoS Pathog 7, e1002392. https://doi.org/10.1371/journal.ppat.1002392

Sidik, S.M., Huet, D., Ganesan, S.M., Huynh, M.-H., Wang, T., Nasamu, A.S., Thiru, P., Saeij, J.P.J., Carruthers, V.B., Niles, J.C., Lourido, S., 2016. A Genome-wide CRISPR Screen in Toxoplasma Identifies Essential Apicomplexan Genes. Cell 166, 1423-1435.e12. https://doi.org/10.1016/j.cell.2016.08.019

Silva, M.F., Douglas, K., Sandalli, S., Maclean, A.E., Sheiner, L., 2023. Functional and biochemical characterization of the Toxoplasma gondii succinate dehydrogenase complex. PLoS Pathog 19, e1011867. https://doi.org/10.1371/journal.ppat.1011867

Sloan, M.A., Scott, A., Aghabi, D., Mrvova, L., Harding, C.R., 2025. Iron-mediated post-transcriptional regulation in Toxoplasma gondii. PLoS Pathog 21, e1012857. https://doi.org/10.1371/journal.ppat.1012857

Smith, D., Kannan, G., Coppens, I., Wang, F., Nguyen, H.M., Cerutti, A., Olafsson, E.B., Rimple, P.A., Schultz, T.L., Mercado Soto, N.M., Di Cristina, M., Besteiro, S., Carruthers, V.B., 2021. Toxoplasma TgATG9 is critical for autophagy and long-term persistence in tissue cysts. eLife 10, e59384. https://doi.org/10.7554/eLife.59384

Soete, M., Fortier, B., Camus, D., Dubremetz, J.F., 1993. Toxoplasma gondii : Kinetics of Bradyzoite-Tachyzoite Interconversion in vitro. Experimental Parasitology 76, 259–264. https://doi.org/10.1006/expr.1993.1031

Solé, V.A., Papillon, E., Cotte, M., Walter, Ph., Susini, J., 2007. A multiplatform code for the analysis of energy-dispersive X-ray fluorescence spectra. Spectrochimica Acta Part B: Atomic Spectroscopy 62, 63–68. https://doi.org/10.1016/j.sab.2006.12.002

Stelzer, S., Basso, W., Benavides Silván, J., Ortega-Mora, L.M., Maksimov, P., Gethmann, J., Conraths, F.J., Schares, G., 2019. Toxoplasma gondii infection and toxoplasmosis in farm animals: Risk factors and economic impact. Food and Waterborne Parasitology 15, e00037. https://doi.org/10.1016/j.fawpar.2019.e00037

Sullivan, W.J., Jeffers, V., 2012. Mechanisms of *Toxoplasma gondii* persistence and latency. FEMS Microbiol Rev 36, 717–733. https://doi.org/10.1111/j.1574-6976.2011.00305.x

Taylor, K.M., Muraina, I.A., Brethour, D., Schmitt-Ulms, G., Nimmanon, T., Ziliotto, S., Kille, P., Hogstrand, C., 2016. Zinc transporter ZIP10 forms a heteromer with ZIP6 which regulates embryonic development and cell migration. Biochemical Journal 473, 2531–2544. https://doi.org/10.1042/BCJ20160388

Teh, M.R., Armitage, A.E., Drakesmith, H., 2024. Why cells need iron: a compendium of iron utilisation. Trends in Endocrinology & Metabolism 35, 1026–1049. https://doi.org/10.1016/j.tem.2024.04.015

Tenter, A.M., Heckeroth, A.R., Weiss, L.M., 2000. Toxoplasma gondii: from animals to humans. International Journal for Parasitology 30, 1217–1258. https://doi.org/10.1016/S0020-7519(00)00124-7

Tomavo, S., Boothroyd, J.C., 1995. Interconnection between organellar functions, development and drug resistance in the protozoan parasite, Toxoplasma gondii. International Journal for Parasitology 25, 1293–1299. https://doi.org/10.1016/0020-7519(95)00066-B

Tomita, T., Bzik, D.J., Ma, Y.F., Fox, B.A., Markillie, L.M., Taylor, R.C., Kim, K., Weiss, L.M., 2013. The Toxoplasma gondii Cyst Wall Protein CST1 Is Critical for Cyst Wall Integrity and Promotes Bradyzoite Persistence. PLoS Pathog 9, e1003823. https://doi.org/10.1371/journal.ppat.1003823

Ullah, I., Wang, Y., Eide, D.J., Dunwell, J.M., 2018. Evolution, and functional analysis of Natural Resistance-Associated Macrophage Proteins (NRAMPs) from Theobroma cacao and their role in cadmium accumulation. Sci Rep 8, 14412. https://doi.org/10.1038/s41598-018-32819-y

Van Dooren, G.G., Yeoh, L.M., Striepen, B., McFadden, G.I., 2016. The Import of Proteins into the Mitochondrion of Toxoplasma gondii. Journal of Biological Chemistry 291, 19335–19350. https://doi.org/10.1074/jbc.M116.725069

Waldman, B.S., Schwarz, D., Wadsworth, M.H., Saeij, J.P., Shalek, A.K., Lourido, S., 2020. Identification of a Master Regulator of Differentiation in Toxoplasma. Cell 180, 359-372.e16. https://doi.org/10.1016/j.cell.2019.12.013

Weber, R.A., Yen, F.S., Nicholson, S.P.V., Alwaseem, H., Bayraktar, E.C., Alam, M., Timson, R.C., La, K., Abu-Remaileh, M., Molina, H., Birsoy, K., 2020. Maintaining Iron Homeostasis Is the Key Role of Lysosomal Acidity for Cell Proliferation. Molecular Cell 77, 645-655.e7. https://doi.org/10.1016/j.molcel.2020.01.003

Wichroski, M.J., Melton, J.A., Donahue, C.G., Tweten, R.K., Ward, G.E., 2002. *Clostridium septicum* Alpha-Toxin Is Active against the Parasitic Protozoan *Toxoplasma gondii* and Targets Members of the SAG Family of Glycosylphosphatidylinositol-Anchored Surface Proteins. Infect Immun 70, 4353–4361. https://doi.org/10.1128/IAI.70.8.4353-4361.2002

Wiuf, A., Steffen, J.H., Becares, E.R., Grønberg, C., Mahato, D.R., Rasmussen, S.G.F., Andersson, M., Croll, T., Gotfryd, K., Gourdon, P., 2022. The two-domain elevator-type mechanism of zinc-transporting ZIP proteins. Sci. Adv. 8, eabn4331. https://doi.org/10.1126/sciadv.abn4331

Wong, S.Y., Remington, J.S., 1994. Toxoplasmosis in pregnancy. Clin Infect Dis 18, 853–861; quiz 862. https://doi.org/10.1093/clinids/18.6.853

Wu, Y., Ying, Z., Liu, J., Sun, Z., Li, S., Liu, Q., 2022. Depletion of Toxoplasma adenine nucleotide translocator leads to defects in mitochondrial morphology. Parasites Vectors 15, 185. https://doi.org/10.1186/s13071-022-05295-7

Xu, Z., Wang, P., Wang, H., Yu, Z.H., Au-Yeung, H.Y., Hirayama, T., Sun, H., Yan, A., 2019. Zinc excess increases cellular demand for iron and decreases tolerance to copper in Escherichia coli. Journal of Biological Chemistry 294, 16978–16991. https://doi.org/10.1074/jbc.RA119.010023

Ying, Z., Yin, M., Zhu, Z., Shang, Z., Pei, Y., Liu, J., Liu, Q., 2024. Iron Stress Affects the Growth and Differentiation of Toxoplasma gondii. IJMS 25, 2493. https://doi.org/10.3390/ijms25052493

Yu, F., Zhang, Q., Liu, H., Liu, J., Yang, S., Luo, X., Liu, W., Zheng, H., Liu, Q., Cui, Y., Chen, G., Li, Y., Huang, X., Yan, X., Zhou, J., Chen, Q., 2022. Dynamic O-GlcNAcylation coordinates ferritinophagy and mitophagy to activate ferroptosis. Cell Discov 8, 40. https://doi.org/10.1038/s41421-022-00390-6

Zhong, M., Zhou, B., 2023. Plasmodium yoelii iron transporter PyDMT1 interacts with host ferritin and is required in full activity for malarial pathogenesis. BMC Biol 21, 279. https://doi.org/10.1186/s12915-023-01776-y