Respiration shapes the neural dynamics of successful remembering in
humans.

Respiration and associative memory retrieval.

Esteban Bullón Tarrasó[1,2], Fabian Schwimmbeck[1,3,4], Marit Petzka[5,6], Tobias Staudigl[1], Bernhard P.
Staresina[7,8] & Thomas Schreiner[1*]

1. Department of Psychology, Ludwig-Maximilians-Universität München, Germany
2. Graduate School of Systemic Neurosciences, Munich, Germany
3. Department of Neurology, Christian Doppler University Hospital, Paracelsus Medical University, Salzburg, Austria.
4. Centre for Cognitive Neuroscience (CCNS), Department of Psychology, University of Salzburg, Austria
5. Institute of Psychology, University of Hamburg, Germany
6. Max Planck Research Group NeuroCode, Max Planck Institute for Human Development, Berlin, Germany
7. Department of Experimental Psychology, University of Oxford, UK
8. Oxford Centre for Human Brain Activity, Wellcome Centre for Integrative Neuroimaging, Department of
Psychiatry, University of Oxford, UK

*Correspondence: Thomas.Schreiner@psy.lmu.de

Declaration of interests

The authors declare no competing interests.

Abstract

Respiration has been shown to impact memory retrieval, yet the neural dynamics underlying

this effect remain unclear. Here, we investigated how respiration shapes both behavioral and

neural expressions of memory retrieval by re-analyzing an existing dataset where scalp

electroencephalography and respiration recordings were acquired while participants (N = 18)

performed an episodic memory task. Our results unveil that respiration influences retrieval-

related power fluctuations in the α/β band and concomitant memory reactivation. Specifically,

we found that both key neural signatures of successful remembering were co-modulated during

exhalation, with the strength of the interaction between respiration and reactivation processes

being associated with memory performance. Together, these findings suggest that respiration

may act as a scaffold for episodic memory retrieval in humans by coordinating the neural

conditions that support effective remembering.

Significance statement

Recent evidence suggests that respiration may shape neural dynamics underlying various

cognitive processes. In this study, we identify respiration as a potential pacemaker for memory

retrieval by showing that key neural signatures of effective remembering—namely, decreases in

α/β power and the reactivation of previously encoded neural representations—are tightly

synchronized with the respiratory cycle. Notably, the strength of this respiration-brain coupling

is associated with individual memory performance, underscoring the critical role and functional

significance of brain–body interactions in supporting cognitive functions.

Introduction

Compelling evidence from both rodent and human studies demonstrates that breathing might serve a purpose beyond supplying oxygen; it shapes neural activity during wake[1,2] and sleep[3,4]. Breathing-locked neural oscillations have been observed across diverse brain areas, oscillatory frequencies, and species[5–9]. While the precise mechanisms enabling this modulation remain unclear, evidence suggests that mechanoreceptors in olfactory sensory neurons, which respond to airflow, generate respiration-locked oscillations in the olfactory bulb[9–12]. These oscillations propagate along the olfactory pathway via the piriform and entorhinal cortices to neocortical and hippocampal regions[1,3]. However, since respiratory rhythm generators in the brainstem may also contribute to the entrainment of breathing-related neural oscillations[3,13–15], the preferred pathway through which breathing modulates neural activity remains uncertain.

Nevertheless, respiration has been shown - potentially through its impact on neural activity - to influence perceptual[16], motor[17,18], and cognitive processes[19] in humans. Performance is typically enhanced when stimuli are presented during inhalation[19–22], suggesting that inhalation-linked neural processes may aid perception. In contrast, mental actions such as motor and visual imagery tend to occur during exhalation[18], indicating that different phases of respiration may create time windows of opportunity for distinct cognitive functions[9]. Notably, memory processes have also been shown to be affected by the phase of respiration[23]. It has been demonstrated that retrieval-related reminder cues are more likely to trigger successful remembering if presented during inhalation as compared to exhalation[20], potentially due to stimulus processing being facilitated during inhalation. However, it is unknown whether memory-related neural processes, aside from stimulus-driven effects, might be associated with respiration, particularly during later phases of the breathing cycle, such as exhalation. Altogether, the results reported above suggest that respiration, and specifically its phase, may directly influence our ability to remember. But how could respiration facilitate effective memory retrieval?

In humans, memory retrieval has long been associated with characteristic patterns of neural activity. For instance, low-frequency power decreases in the α/β range (~8 – 30 Hz) have been consistently found to accompany successful remembering in cortical areas and the hippocampus[24–27]. The magnitude of those decreases has been proposed to reflect heightened levels of information processing and retrieval-related engagement of cortical modules[25]. In line with this assumption, it has been shown that α/β power fluctuations do not only track the quantity but also the level of detail of memory-related information represented in the brain[28–30]. Moreover, several studies have established a tight link between α/β power decreases and the emergence of retrieval-related memory reactivation[31,32]. Specifically, retrieved memory

information seems to be particularly decodable from electroencephalographic (EEG) signals

during low α/β power levels[28,33], while other studies indicated that the low frequency oscillatory

phase itself might carry a temporal signature of the reactivated memory content[34,35].

Importantly, beyond its close relationship with memory processes, α/β activity has been

demonstrated to be robustly modulated by respiration during diverse cognitive tasks (i.e.,

visuospatial, lexical, and math)[19], perception[16] and rest[6]. Hence, respiration might accomplish its

influence on the behavioral aspects of memory retrieval by modulating low-frequency

oscillatory activity accordingly. However, it remains unknown whether and to what extent

respiration might indeed coordinate retrieval-related power decreases in the α/β band and

memory reactivation.

Here, we set out to directly assess the influence of respiration on the behavioral and

neural expressions of memory retrieval using electroencephalography (EEG) and respiration

recordings acquired during an episodic memory task. First, we replicate previous findings

indicating that the phase of respiration at cue presentation influences memory performance,

with successful retrieval being more likely when reminder cues were presented close to

inhalation. We further show that this effect is potentially driven by respiration's impact on both

power modulations in the α/β band and concomitant memory reactivation. Moreover, the

strength of the interplay between respiration and retrieval-related reactivation is directly

associated with behavioral memory performance. Our results identify respiration as a potential

pacemaker for episodic memory retrieval in humans and pave the way for future research on

brain-body interactions during memory formation.

Results

To examine whether respiration affects the behavioral and neural dynamics of memory

retrieval, we analyzed EEG and respiratory recordings from 18 participants across two

experimental sessions, each involving an episodic memory task (Fig. 1a). During the learning

phase, participants were instructed to associate verbs with images of either objects or scenes,

depending on the experimental session. Memory performance was evaluated in a sequential

manner. Participants first completed a word-recognition task, classifying words (presented 3 s

each) as "old" or "new." For words recognized as "old", participants attempted to retrieve the

specific associated image (object or scene). Associative retrieval performance was normalized

by the number of accurately recognized items ("hits"). Each session concluded with an

independent "localizer task", during which participants were presented with a new set of stimuli

comprising both object and scene images. The data of the localizer task were used to train a

linear classifier in order to distinguish object- and scene-related EEG patterns during memory

retrieval. Note that parts of the data have already been published, showing that (i) slow
oscillation-spindle complexes clock memory reactivation during NREM sleep[36] and (ii) the
occurrence of these sleep-related oscillations is modulated by respiration[4].
112

113
Figure 1. Experimental procedure, behavioral results, and retrieval-related EEG results. (a) During encoding,
participants were presented with 120 verb-object or verb-scene pairs, depending on the experimental session, and
memory performance was evaluated afterward. At the end of each session, participants performed a localizer task
that involved processing a new set of object and scene images. (b) Behavioral results for both experimental sessions:
objects (light gray) and scenes (dark gray). Bar graphs show the mean percentage (±SEM) of recalled image exemplars
out of correctly recognized verbs. Dots indicate individual memory performance of participants (N = 18). No significant
difference was found in memory accuracy between experimental sessions. (c) Stimulus categories (objects vs. scenes)
could be decoded above chance from the localizer EEG data, starting around 150 ms post stimulus onset (the black
solid line indicates decoding performance (±SEM)). The horizontal dashed line indicates surrogate decoding
performance, which was estimated by shuffling the training labels 250 times. The vertical dashed line indicates
stimulus onset (time = 0 s). The lower horizontal dark gray line shows the temporal extent of significant decoding
results as derived from a dependent-samples t-test (one-sided, $p < 0.001$, corrected across time). (d) Remembered
trials were associated with stronger power decreases in the α/β band (~8-20 Hz) after cue onset (~0.5 to 1.5 s) as
compared to not remembered trials (contour lines indicate the extent of the significant cluster, $p < 0.001$, d = -1.5950;
corrected across time, frequency and electrodes). The topographical inset illustrates the sum of t-values across all
significant time points and frequencies for each electrode, illustrating that power differences were most prominent
across frontal and parietal areas. (e) Learning-related brain patterns (objects vs. scenes) were decodable during
retrieval (contour lines indicate the extent of the significant cluster, $p = 0.032$, d = 0.5972, one-sided; corrected across
times); color range (blue to yellow) represents t-values comparing the decoding performance distributions of
remembered versus not-remembered trials.

 Behavioral results and localizer-related classification

As previously reported[36], memory performance was tested before and after a 120-

minute nap (see Methods). To examine potential differences between the two learning

categories (objects vs. scenes), memory scores were averaged across pre- and post-sleep

sessions. No significant difference emerged between object and scene conditions (two-sided

paired-sample t-test, $t_{(17)}$ = 0.37, p = 0.71, d = 0.0872), suggesting that task difficulty was well

matched across image categories (see Supplementary Table 1 for a detailed overview of memory

measures). To derive the neural signatures of category-specific processing, which were used to

track memory reactivation during retrieval, we employed a localizer task at the end of each

session. During this task, participants viewed a novel set of images comprising both objects and

scenes. Category-specific EEG patterns were extracted by aggregating the localizer data across

experimental sessions and performing multivariate classification using linear discriminant

analysis (LDA). Applying fivefold cross-validation, classification accuracy significantly exceeded

chance levels between ~150 ms and 1500 ms after stimulus onset (p < 0.001, corrected across

time, Figure 1c), indicating that the classifier reliably captured category-specific neural patterns

of object and scene image processing.

Memory retrieval is associated with α/β power decreases and memory reactivation

As a first step, we set out to identify the oscillatory fingerprint of successful memory

retrieval. To extract memory-related EEG patterns, we contrasted time-frequency

representations between remembered and not-remembered items, based on associative

memory test performance. In line with previous findings[24,29,37], successful remembering was

associated with significant power decreases in the α/β range (~8 – 20 Hz), starting approximately

500 ms after cue onset across fronto-parietal areas (p < 0.001, corrected across time, frequency,

and space; see Figure 1d). Next, we assessed whether neuronal activity related to the session-

specific learning categories would be reactivated during successful retrieval[27,38]. Again, we

trained an LDA classifier on the concatenated localizer data from both experimental sessions

(objects and scenes; −0.15 to 1.5 s). The resulting training weights were then applied to retrieval

data from both sessions, centered around cue onset (-0.15 to 1.5 s). The classifier testing labels

matched the stimulus category from the preceding encoding session (object or scene), such that

above-chance classification reflects activation patterns that more closely resemble the newly

learned stimulus category than the alternative one. As shown in Figure 1e, results revealed

significantly higher classification performance for remembered as compared to not-

remembered items emerging between 0.5 and 1.5 s relative to cue onset (p = 0.032, corrected

across times; see Supplementary Fig 1c for comparisons against chance-level distributions
obtained via label shuffling).

Respiration phase modulates memory performance and EEG activity during retrieval

To assess whether the phase of respiration (mean respiratory frequency = 0.267 ± 0.041
Hz) would impact memory performance during retrieval[20], cue-centered trials (-0.5 to 3 s; see
Supplementary Fig. 1 for the cue-centered respiratory event-related potential) were segmented
into 250 ms bins. Each bin was then categorized according to its proximity to either an inhalation
peak (i.e., the inhalation-to-exhalation transition) or an exhalation trough (i.e., the exhalation-
to-inhalation transition). Memory accuracy for both categories was computed within each time
bin and normalized by the participant's individual overall performance. This yielded two time
series per participant—one for bins aligned with the inhalation peak and one with the exhalation
trough—capturing how changes in the respiratory phase over time impacted memory accuracy.
A non-parametric cluster-based permutation test revealed that respiration significantly
influenced retrieval performance (Fig. 2a, for recognition memory related results, see
Supplementary Fig 1b). Specifically, successful retrieval was more likely in trials where the
respiration cycle was closer to inhalation peaks between 0.125 and 0.625 s ($p = 0.047$, corrected
across time) and to exhalation troughs between 1.875 and 2.375 s relative to cue onset ($p =$
0.044, corrected across time). Given the cyclical nature of respiration—moving from inhalation
to exhalation—these findings suggest that trials in which participants inhaled around cue
presentation and then transitioned to exhalation (within a trial) were more likely to be
remembered.

To further corroborate this assumption, we divided the trials into two groups: one
where inhalation peaks occurred before exhalation troughs during retrieval time and another
where the order was reversed. Next, memory accuracy was computed for each of the categories.
As shown in Fig. 2b, memory performance during retrieval was significantly enhanced when
inhalation peaks preceded exhalation troughs rather than the other way around (dependent
samples t-test, two-sided, $t_{(17)} = 2.868$, $p = 0.011$). Thus, retrieval was more likely when cues
occurred prior to inhalation peaks, with an additional increase in retrieval probability around
exhalation troughs later in the trial. The follow-up sequence analysis clarified that these effects
reflect a complementary pattern: memory performance was highest when an inhalation at cue
onset was followed by exhalation thereafter.

To examine whether respiration would likewise modulate EEG activity during retrieval,
we calculated the modulation index (MI)[39] to estimate cross-frequency phase-amplitude
coupling between respiration (providing the phase) and EEG signals across all electrodes

(providing amplitude measures between 3–30 Hz). For statistical assessment, we generated an

MI-level reference distribution by randomly shifting the phase-related time series of each trial

200 times. The mean and standard deviation across the reference distribution were then used

to z-score the MI of the empirical distribution. Results revealed a significant modulation of EEG

amplitudes by the phase of respiration with local peaks in the α (9 Hz, z > 2.58 / p < 0.01) and β

range (18 Hz, z > 1.96 / p < 0.05; Fig. 2c). Next, to specifically characterize EEG fluctuations in the

α/β band across the respiratory cycle, we divided the respiratory phase of each trial into 60

evenly spaced bins (bin size = 6°). Then, the mean power in the α/β band (i.e., 8–20 Hz) was

computed for each phase bin across all trials and contrasted against a null distribution (see

Methods). The 8–20 Hz range was chosen because it corresponds to the α/β band that showed

the most consistent retrieval-related power decreases in our dataset (Fig. 1d) and in previous

studies[25,38,40]. This predefined band was therefore used to examine respiration-locked

modulation of retrieval-relevant oscillatory activity. We found respiration-locked EEG power

exhibited significant decreases around the exhalation through (114 to 138°, p = 0.023; 210 to

246°, p = 0.043) and towards the inhalation peak (288 to 330°, p = 0.003; all corrected for

multiple comparisons across phase bins and electrodes, see Fig. 2d).

221
Figure 2. Respiration's impact on memory performance and oscillatory EEG activity. (a) Successful retrieval was more
likely in trials where participants inhaled during the period of cue presentation—with inhalation peaks occurring
between 0.125 and 0.625 s (p = 0.047, d = 0.2927, corrected across time)—and subsequently exhaled — with
exhalation troughs occurring between 1.875 and 2.375 s (p = 0.044, d = -0.3061, corrected across time). Blue / orange
lines represent the difference in accuracy with respect to the participant's overall memory performance for the trials
that were closer to the inhalation peak /exhalation trough in each time bin (±SEM). The horizontal black line illustrates
the temporal extent of significance as derived from a dependent-samples t-test (two-sided, cluster corrected across
time). (b) Memory performance during retrieval was significantly enhanced when inhalation peaks preceded
exhalation troughs (mean ± SD: 70.03±13.70) rather than the other way around (mean ± SD: 64.67 ± 17.11) . Box plots
show median (±$Q_{1/3}$) accuracy of recalled image exemplars out of correctly recognized verbs. Star denotes a significant
difference as derived from a two-sided, dependent-samples t-test ($t_{(17)}$ = 2.868, p = 0.011, d = 0.6761). (c) The
modulation index (± SEM, in shade grey) suggests that the phase of respiration significantly influences EEG amplitudes
in the α/β range with local peaks at 9 Hz (z > 2.58) and 18 Hz (z > 1.96). The lower horizontal black line depicts all
significantly modulated frequencies (p < 0.05, corrected across frequencies using FDR). (d) Power in the α/β band is
modulated at different phases of the respiratory cycle (purple solid line represents the power of [8-20 Hz] band
(±SEM)). Segments in red indicate phase bins associated with significant power decreases in the α/β range as derived
from a dependent-samples t-test against a shuffled reference distribution (114 to 138°, p = 0.023, d = -1.4681; 210 to
246°, p = 0.043, d = -1.3003; 288 to 330°, p = 0.003, d = -1.4326; two-sided, cluster corrected across phase and
electrodes; inhalation peak = 0°). Light blue (orange)-shaded areas represent significant phase bins closer to the
inhalation peak (exhalation trough). The topographical insert illustrates the sum of t-values across all significant
phases for each electrode, showing that this decrease was more prominent in frontal, parietal, and right-central
cortices.

Respiration modulates retrieval-related memory reactivation and its interplay with α/β power

Having established that respiration modulates both retrieval-related memory

performance and EEG activity in the α/β range during retrieval, we finally asked whether

respiration would also impact the emergence of memory reactivation. To this end, we detected

exhalation troughs, the most prominent feature of the cue-locked respiratory ERP (see

Supplementary Fig. 1a), in the respiratory signal during retrieval (0 to 3 s relative to cue onset)

and centered the EEG data accordingly. Again, a classifier was trained on the pooled localizer

data from both learning sessions (– 0.15 to 1.5 s), but the resulting training weights were this

time specifically applied to the exhalation trough-locked EEG segments (±2 s, accounting for a

whole respiratory cycle with a mean respiratory frequency of 0.25 Hz). Results revealed

significantly higher decoding performance for remembered as compared to not-remembered

trials emerging just prior to the exhalation through (p = 0.042, corrected across times, Fig. 3a;

see Supplementary Fig 2a for remembered and not remembered against chance-level

distributions obtained via label shuffling). However, while these effects support the notion of

retrieval-related reactivation, they should nonetheless be interpreted with caution given their

moderate size.

To further assess the functional significance of the coupling between memory

reactivation and respiration, we determined its impact on the participants' memory

performance. To this end, we extracted decision values from the decoding procedure, yielding

a single-trial, time-resolved metric that reflects the classifier's confidence in assigning a specific

category (objects vs. scenes) in the cue-centered EEG data[41]. Then, for each of the remembered

trials, we extracted the respiratory phase where decision values reached their maximum. We

found that decision value maxima consistently emerged during exhalation across participants

(mean angle = 93.045º ± 0.321; Rayleigh test z = 14.58, *p* < 0.001; see polar insert in Fig. 3b).

Finally, when correlating the vector length (representing the coupling strength between

respiration and classification maxima across trials) with memory performance, we observed a

significant positive relationship (Fig. 3b; Spearman's Rank Correlation, r = 0.558, *p* = 0.018; for

results based on non-remembered trials see Supplementary Fig. 2). The distribution indicated

that the stronger the reactivation-related EEG signatures were modulated by respiration, the

better the memory performance.

So far, we have demonstrated that respiration might serve as a temporal anchor for

retrieval-related neural processes, as both characteristic power decreases in the α/β band and

memory reactivation were significantly modulated by the phase of respiration. Hence, we

hypothesized that respiration might co-modulate both key markers of successful memory

retrieval. Thus, we extracted trial-based decision values, locked to exhalation troughs, and

correlated them with α/β power values at each electrode and timepoint across trials. Again, we

focused on the predefined 8–20 Hz α/β range identified as most consistently linked to retrieval-

related power decreases (Fig. 1d), ensuring that respiration–reactivation coupling was assessed

within the same functionally relevant frequency band. When contrasting correlational values
between remembered and not-remembered trials, we observed a significant negative cluster
emerging just prior to the exhalation through (*p* = 0.023; corrected for multiple comparisons
across time and electrodes, Fig. 3c), indicating that the stronger the respiration locked α/β power
decrease, the higher the classifiers fidelity during successful remembering.

289
Figure 3. Respiration impacts memory reactivation and concomitant α/β power fluctuations. (a) Learning-related
brain patterns (objects vs. scenes) were decodable during retrieval based on exhalation trough-centered EEG data.
Contour lines indicate the extent of the significant cluster (*p* = 0.042, d = 0.8434, one-sided, corrected across times);
color range (blue to yellow) represents t-values of remembered against not-remembered distributions of decoding
performances. (b) In remembered trials, coupling strength (measured as vector length) between respiration and
classification maxima is positively correlated with memory performance (Spearman's Rank Correlation, r = 0.558,
p = 0.018). Nonlinear fit as a dashed black line (only for visualization purposes). The polar histogram illustrates the
mean respiratory phase at which decoding-derived reactivation peaks emerge during remembered trials for each
participant (mean angle = 93.045º ± 0.321; Rayleigh test z = 14.58, *p* < 0.001; inhalation peak = 0°). (c) Decoding-
derived single trial decision values and α/β power (8-20 Hz) are more negatively correlated prior to the exhalation
trough in remembered trials as compared to not-remembered trials. The green (red) solid line shows the Spearman's
correlation for (not) remembered trials (±SEM). The horizontal black line shows the temporal extent of significant
results as derived from a dependent-samples t-test (two-sided, *p* = 0.023, d = -1.0248, cluster corrected across time
and electrodes). The topographical inset illustrates the sum of t-values across all significant timepoints for each
electrode, showing that this difference was stronger in the parietal cortex.

Discussion

307   Our results unveil a role of respiration in modulating both behavioral and neural
308 components of remembering. In line with previous research, we show that the phase of
309 respiration influences memory performance (Fig. 2a), with successful retrieval being more likely

when reminder cues were presented close to inhalation. Moreover, key neural signatures of retrieval, oscillatory power decreases in α/β band (Fig. 2d) and memory reactivation (Fig. 3a), as well as their interplay (Fig. 3c) were modulated by respiration. Finally, the strength of the coupling between respiration and reactivation processes was associated with memory performance (Fig. 2b), underscoring the functional relevance of respiration–brain interactions for memory.

Extensive research in both rodents[3,11,42] and humans[2,6,20] indicates that respiration might be essential for modulating neural oscillations across various brain regions and frequency bands. Mechanistically, it has been suggested that the impact of respiration on neural activity is mediated by incoming airflow stimulating mechanoreceptors in olfactory sensory neurons[12]. Through this stimulation, breathing-locked oscillations are generated and then relayed to the olfactory cortex and spread to various target regions, and in particular to sensory areas[9,10,12]. These oscillations transiently increase neural excitability and may thus facilitate stimulus perception[16]. Beyond sensory processing, respiration has also been shown to impact cognition in various domains such as motor control[17,18] and emotion[21], often enhancing performance when stimuli are presented during inhalation[7]. In particular, a growing number of findings in humans demonstrate that breathing impacts memory processes during wake[23 (but see 43,44)]. Zelano and colleagues were the first to show that during retrieval testing, memory was better for stimuli that were presented during inhalation as compared to exhalation[20]. Interestingly, this effect became only evident when participants engaged in nasal breathing (as compared to mouth breathing), suggesting a specific role for nasal respiration in orchestrating memory-related processes. Our data remain agnostic as to whether nasal breathing is indeed crucial for impacting memory performance. Nevertheless, our finding that successful retrieval was more likely when participants inhaled during reminder cue presentation is in line with the previous result, indicating that the phase of respiration at the time of cue presentation impacts memory performance. However, while these results appear to suggest that stimulus processing might be facilitated during inhalation, the neural mechanisms underlying respiration's influence on successful memory retrieval have remained unexplored. Notably, another study examining respiration and memory[44] did not observe performance differences across the respiratory cycle. This discrepancy may relate to task design: Schaefer et al. used a recognition task without associative retrieval demands, whereas our paradigm required retrieval of specific verb–image associations. Indeed, when we restricted our analysis to recognition trials, no significant modulation by respiration phase was found (Supplementary Fig. 1b). Taken together, these observations suggest that respiration effects may emerge more prominently in tasks requiring associative retrieval or stronger mnemonic reconstruction than in simple recognition paradigms.

Cortical power decreases in the α/β band have consistently been reported to index

successful memory retrieval[24-27]. It has been suggested that these decreases might facilitate

remembering by allowing neural assemblies to express the specific neural codes associated with

stored memories[25 (but see 45)]. Indeed, α/β power modulations have been shown to track the

amount and content of information being retrieved[25,46,47], the fidelity of memory details[28-30], and

potentially provide time-windows of opportunity for memory reactivation[24,35,48]. In our data,

successful retrieval, irrespective of respiration, was associated with robust power decreases in

the α/β band across frontal and parietal areas. Two separate learning sessions with distinct

stimulus sets (objects vs. scenes; Fig. 1a) enabled us to track retrieval-related reactivation and

demonstrate that learning-specific patterns were decodable during periods of low α/β power.

Beyond these confirmatory findings concerning the role of low-frequency power

decreases and concomitant memory reactivation in memory retrieval, our results suggest that

respiration might play a crucial role in modulating those neural dynamics underlying successful

remembering. Fluctuations in α/β activity have already been demonstrated to be influenced by

respiration during diverse cognitive tasks (i.e., visuo-spatial, lexical, and math)[19], perception[16],

and rest[6]. Our results critically extend these findings by demonstrating that also retrieval-related

α/β fluctuations are tightly modulated by respiration. Specifically, we found that decreases in

α/β activity systematically emerged both around exhalation troughs and just prior to inhalation

peaks. These power reductions, synchronized with different phases of the respiratory cycle, may

serve distinct functional roles. As outlined above, our behavioral analyses revealed that

successful remembering was more likely when participants inhaled during the reminder cue

presentation and exhaled thereafter (i.e., with exhalation troughs occurring between 1.875 and

2.375 s after cue onset). Interestingly, in addition to their role in memory retrieval, low pre-

stimulus α/β levels have consistently been linked to enhanced perceptual performance[49]. Thus,

inhalation-locked decreases in low-frequency power may facilitate stimulus perception[16],

forming a fundamental foundation for cue-triggered remembering. On the contrary, signatures

of retrieval-related memory reactivation are typically observed between ~500 and 1500 ms after

reminder cue onset[38]. Accordingly, low-frequency power decreases emerging around exhalation

troughs might establish a neural environment that is optimally tuned for memory reactivation

to unfold. This interpretation is in line with our finding that not only power decreases in the α/β

band were modulated by exhalation, but that classification-derived memory reactivation, hence

the putative neural substrate of remembering[31], emerged just prior to the detected exhalation

troughs. Therefore, our results add further evidence to the notion that that different phases of

respiration might establish time windows of opportunity for distinct brain functions[9]. In addition,

respiration has recently been established to similarly influence the emergence of sleep-related

oscillations and in consequence, memory reactivation in humans[4], with reactivation processes

putatively unfolding during exhalation. Together, these results suggest that respiration might

impact memory-related neural oscillations and concomitant memory reactivation in a consistent

manner, regardless of state. One potential concern of our paradigm, however, was that

participants were required to withhold their responses for three seconds, ensuring that the

respiration effects we report occurred well before any motor output. We therefore interpret the

observed exhalation-locked neural dynamics as reflecting retrieval rather than response

execution. Nevertheless, future work without a response delay will be valuable to further

disentangle how respiration may jointly coordinate internal retrieval and overt responding.

The negative correlation between α/β power and memory reactivation signatures prior

to exhalation troughs suggests that respiration may shape the interaction between two key

neural markers of successful remembering. However, it remains unclear whether respiration

directly shapes the emergence of memory reactivation or acts indirectly via its effects on α/β

activity. Recent studies suggested that α/β power decreases may reflect eye movements rather

than memory processes per se[45,50]. In this light, our findings fit within a broader brain–body

framework, in which respiration not only modulates cortical excitability[16,51], but may also

influence, or be influenced by, oculomotor behavior[52,53], thereby contributing to fluctuations in

α/β power. To disentangle these intertwined processes, future studies combining respiration

measures, eye tracking, and neural recordings will be essential. Moreover, our design is

correlational and cannot determine whether respiration actively drives retrieval-related neural

dynamics or whether both are co-modulated by a third factor (e.g., arousal, attention, or

autonomic tone). Causal manipulations of breathing will therefore be crucial.

Complementary intracranial recordings could further test whether respiration

modulates the occurrence of hippocampal ripples during retrieval[54], which are known to trigger

memory reactivation in both hippocampal and cortical regions[55]. Given that ripple occurrence

has been shown to be modulated by respiration in rodents[3,56] and more recently in humans[57],

particularly emerging during exhalation, such work may clarify whether exhalation provides a

temporal window of reduced neural noise that facilitates mnemonic reactivation. In any case,

our finding that the strength of respiration–reactivation coupling predicts individual memory

performance provides compelling evidence for the functional relevance of this relationship,

irrespective of whether respiration exerts a direct or indirect influence on reactivation

processes. Nevertheless, while our analyses were specifically locked to the respiratory cycle, and

the observed effects align with prior reports of respiration–brain coupling[6,20], we cannot exclude

the possibility that other physiological rhythms (e.g., cardiac cycle, slow cortical potentials)

contribute to the observed modulations[58]. Future studies concurrently measuring respiration,

cardiac activity, and other slow fluctuations will be essential to disentangle their respective

influences on retrieval-related neural dynamics.

To conclude, our results reveal a critical role for respiration in modulating both the

behavioral and neural aspects of remembering. They identify respiration as a potential

pacemaker for effective remembering and highlight the importance of brain-body interactions

in human memory processing.

Methods

The current analyses are based on reference[36], where detailed information about participants,

stimuli, task, and data acquisition can be found. The data of two participants (out of 20) had to

be excluded due to low trial numbers (< 15 trials in any of the conditions). In summary, the study

involved 18 healthy participants (mean age: 20.89 ± 0.34; 15 female). Pre-study screening, which

included the Pittsburgh Sleep Quality Index (PSQI[59]), the morningness-eveningness

questionnaire[60], and a custom questionnaire assessing general health and stimulant use,

confirmed that participants were not taking any medication and had no neurological or

psychiatric conditions at the time of the experiment. The University of Birmingham Research

Ethics Committee approved the study, and all participants provided written informed consent.

They were compensated either financially or with course credit for their participation.

Experiment overview

The experiment involved two sessions per participant, spaced at least one week apart (mean

interval: 7.78 ± 0.43 days), with the session order counterbalanced across participants. On each

experimental day, participants arrived at the sleep laboratory at 11 a.m., where the set-up for

polysomnographic recordings began. Electrodes for electroencephalography (EEG),

electromyography (EMG), and electrocardiography (ECG) were applied, along with a thermistor

airflow sensor to monitor breathing. Around noon, the experiment commenced with a modified

psychomotor vigilance task (PVT[61]), followed by a memory task (see Encoding and Memory tests

below) in which participants learned to associate 120 verbs with images (either objects or

scenes[62], depending on the session). The sleep period began at approximately 1 p.m., giving

participants the opportunity to nap for ~120 minutes. After the nap, their vigilance was

reassessed using the PVT, and memory performance was tested again. A localizer task was

conducted at the end of each session (see Localizer task below). Behavioral responses and the

presentation of tasks were managed using Psychophysics Toolbox Version 3[63] and MATLAB

2018b (MathWorks, Natick, USA).

Stimuli. The experimental stimuli consisted of 360 verbs and 240 images, evenly divided between object and scene categories, presented during both experimental sessions. Object images depicted animals, food items, clothing, tools, or household items against a plain white background (e.g., a calculator). Scene images represented identifiable landscapes or locations (e.g., an island). These categories were chosen as they engage distinct brain networks, such as the lateral occipital complex for object processing[64] and the para-hippocampal place area for scene processing[65], facilitating the investigation of experience-driven memory reactivation during retrieval (see Methods: Multivariate analysis). All images were obtained from reference [62].

Familiarization. The experiment commenced with an image familiarization phase designed to (i) facilitate the learning of verb-image associations during the subsequent encoding session and (ii) establish accurate image labels for later cued recall. Each trial began with the presentation of a fixation cross for 1.5 ± 0.1 s. Participants were then shown one of 130 images depicting either objects or scenes, depending on the experimental session. Of these, 120 images were included in the learning material and accompanied by captions accurately naming the depicted exemplars. An additional 10 images, which were not used further in the experiment, were paired with incorrect captions. Each stimulus was displayed for 2.5 s. Participants were instructed to press a button upon detecting an erroneous image-caption pairing.

Encoding. Each trial began with the presentation of a fixation cross for 1.5 ± 0.1 s, followed by a verb (e.g., "jump") displayed for 1 s. Immediately after, the corresponding image to be associated with the verb was presented for 4 s. Participants were instructed to create a vivid mental image or narrative linking the verb with the object or scene depicted in the image. Following the image presentation, participants indicated whether the mental image they had formed was realistic or bizarre. Participants were informed in advance that their memory for the verb-image pairs would be tested subsequently. The learning block was repeated twice with a randomized trial order in each iteration to achieve satisfactory memory performance, as established in a prior pilot study.

Memory tests. Participants' memory for the encoded verb-image pairs was assessed both before and after the nap interval. To mitigate potential testing effects on behavioral measures of memory consolidation[66,67], only half of the learned verb-image pairs were included in each memory task, with no overlap between the two tests. Each memory test comprised 60 randomly

selected verbs from the learning phase, intermixed with 30 novel verbs ("foils") that had not been presented during learning. Each trial began with a fixation cross displayed for 1.5 ± 0.1 s, followed by the presentation of a verb on the screen. After 3 s, participants indicated within a 10 s window whether the verb was "old" (i.e., part of the learning material) or "new" (i.e., not seen during learning). For "new" responses, participants proceeded directly to the next trial. For "old" responses, participants were prompted to either type a description of the associated image or enter "do not know" if they could not recall the target image. Responses were scored as correct if (i) the typed description matched the original caption presented during the familiarization phase or (ii) the description unambiguously corresponded to the content of the target image.

Localizer task. During the localizer task, participants viewed a new set of images comprising 90 objects and 90 scenes, irrespective of the experimental session. Each trial began with a fixation cross presented for 1.5 ± 0.1 s, followed by the display of a randomly selected image (object or scene) on the computer screen for a duration between 2.5 and 10 s. Each image was presented twice during the task, and participants were instructed to indicate whether the image was being shown for the first time ("new") or the second time ("old").

EEG and respiration

A Brain Products 64-channel EEG system was used to record electroencephalography (EEG) throughout the experiment. Impedances were kept below 10 kΩ. EEG signals were referenced online to electrode FCz and sampled at a rate of 1000 Hz. Respiration was recorded using an Embla thermistor airflow sensor.

Data analysis

Behavioral. Analysis was performed based on associative memory performance, calculated as the proportion of correctly remembered images relative to the number of 'hit' trials (i.e., verbs correctly identified as old), separately for each session (object and scene) and memory test (pre- and post- sleep). To relate the respiratory phase to memory performance across trials (Fig. 2a), we first collapsed all trials from both sessions and memory tests (i.e., pre- and post-sleep). Then, trials were segmented into 250 ms bins, starting -0.5 s before cue onset. For each time bin, we categorized trials based on whether the circular mean of the respiration phase across the 250 ms of a bin aligned more closely with the inhalation peak or the exhalation trough, respectively. Memory accuracy was then computed for each category (inhalation and exhalation relatedness, respectively) within each time bin and normalized by each participant's overall memory

performance. Finally, to analyze whether specific respiration sequences (inhalation to

exhalation or exhalation to inhalation) during retrieval influence memory performance, we

categorized trials based on the presence of either an inhalation peak followed by an exhalation

trough or the reverse within -0.5 to 3 s from cue onset. Then, associative memory performance

was computed for each category.

EEG and respiration. All data were analyzed using MATLAB (2020a; MathWorks). EEG data were

preprocessed using the FieldTrip toolbox[68] for EEG/MEG analysis (v.16/07/2023). All data were

downsampled to 200 Hz and re-referenced using a common average reference (CAR). Noisy EEG

channels were identified by visual inspection, discarded, and then interpolated using a weighted

average of the neighboring channels. EEG epochs related to the retrieval task were locked to

two different events: stimulus presentation and exhalation troughs. Exhalation troughs (and

inhalation peaks for Fig. 2b) were detected using the Breathmetrics toolbox[69]. After detecting

exhalation troughs and inhalation peaks from the respiratory signal, trials were classified as INHP

→ EXHT when an inhalation peak preceded an exhalation trough within −0.5 to 3 s around cue

onset (i.e., the same time window used in Fig. 2a), and as EXHT → INHP for the reverse order.

This classification was based on detected respiratory events rather than phase estimates,

minimizing potential bias from phase interpolation. In addition, trials that contained an

exhalation trough within 3 s after cue onset were aligned to that respiratory event for analyses

shown in Fig. 3, as exhalation troughs constituted the most robust and temporally consistent

respiratory marker across trials (Supplementary Fig. 1). Trials were classified into remembered

and not-remembered categories based on associative memory performance. In the case of

localizer trials, all data were locked to the stimulus onset with a temporal window of -0.2 to 3 s.

For analyses related to respiratory phase, the phase was extracted using the double

interpolation method[4,16], unless specified otherwise.

Time-frequency analysis. Time-frequency analysis of stimuli-locked (-2 to 3 s) and respiratory

event-centered EEG segments (−5 to 5 s) was performed using FieldTrip[68]. The longer time

segments were chosen to allow for resolving low-frequency activity within the time windows of

interest for both stimuli-locked (-0.2 to 1.5 s) and respiratory event-locked (−2 to 2 s) and avoid

edge artifacts. Frequency decomposition of the data was achieved using Fourier analysis based

on sliding time windows (moving forward in 50 ms increments). The window length was set to

five cycles of a given frequency (frequency range: 1–30 Hz in 1 Hz steps). The windowed data

segments were multiplied with a Hanning taper before Fourier analysis. Afterward, power values

were z-scored across time, using a 1 s baseline window prior to stimulus onset. To analyze

oscillatory power as a function of the respiratory phase, power values were transformed from

the time axis to the phase axis. For that, the respiratory cycle was divided into 60 non-

overlapping phase bins (bin size = 6°), and after assigning all power values to the respiratory

phase bin where they occurred, the mean power was computed for each bin[16,51]. Thus, we

obtained a respiratory phase-resolved distribution, with the value at each phase bin

representing the mean of all power values that overlapped with that respiratory frequency

across all trials from 0 to 3 s.

Modulation Index. Phase amplitude coupling was assessed using the Modulation Index (MI)[39]. To

estimate the instantaneous phase of the respiration signal, we filtered the continuous

respiratory data around the peak frequency of each particular subject ± 0.05 Hz two-pass

Butterworth band-pass filter). We then extracted instantaneous amplitude data across

frequencies between 3 and 30 Hz at each electrode in steps of 2 Hz regardless of the α/β regions

identified in Fig. 1d, to avoid circularity and because respiration has been shown to modulate

large-scale neural networks. To this end, a two-pass FIR filter (order = three times the lower

frequency bound) was used to create 27 equally spaced frequency bins, with center frequencies

ranging from 3 to 30 Hz, and with fixed-frequency bandwidths of 2 Hz. The envelope of the

Hilbert-transformed band-pass filtered data was then used as an amplitude estimate. To

compute the MI (for a given frequency pair), we divided the phase signal into 20 bins, and then,

for each bin, computed the mean amplitude. This yields a distribution of amplitude as a function

of phase. The MI is defined as the Kullback–Leibler distance between that distribution and the

uniform distribution (across the same number of bins). To assess the statistical significance of

MI values, we generated participant-specific null distributions by randomly shifting the

respiration phase time series relative to the neural signal (200 iterations per participant). The

mean and standard deviation of each null distribution were used to z-score the empirical MI

values, which were then converted into p-values. This phase-shuffling procedure preserves

individual respiratory dynamics while providing within-subject reference estimates.

Additionally, we equalized the number of correct and incorrect trials within each participant via

subsampling. Results were corrected for multiple comparisons within each channel using FDR

across frequencies. The channels with frequencies that survived FDR correction were then

averaged, and frequencies laying over $z > 1.96$ (i.e., $p < 0.05$) were considered significant.

Multivariate analysis. Multivariate classification of single-trial EEG data was performed using

MVPA-Light, a MATLAB-based toolbox for multivariate pattern analysis[70]. For multivariate

classification, Linear Discriminant Analysis (LDA) was used as classifier[70]. Prior to analyses the

raw EEG data were bandpass-filtered between 0.1 and 45 Hz (4[th] order Butterworth filter) and

then z-scored the localizer data across all trials independently for each time point. Data from

both sessions were then combined and subjected to principal component analysis (PCA), which

transforms the data into linearly uncorrelated components ranked by the variance explained by

each component[71]. PCA was employed to reduce dimensionality and mitigate overfitting,

retaining the first principal components that explained 95% of the data's variance[72]. To assess

whether object and scene representations could be distinguished in the localizer, we trained

and tested a classifier to discriminate between object and scene trials. A 200 ms running average

smoothing window was applied to the data. EEG channels were treated as features, and a

separate classifier was trained and tested at each time point. Classification performance was

quantified using the area under the receiver operating characteristic curve (AUC), which

measures the probability that a randomly selected pair of Class A and Class B trials is correctly

assigned to their respective classes (0.5 = chance level; 1.0 = perfect classification). To prevent

overfitting, fivefold cross-validation was implemented, with the analysis repeated five times to

account for variability arising from random fold assignments[73]. The results were averaged across

iterations. Statistical significance was evaluated by generating surrogate decoding performance

values through 250 iterations of label shuffling, producing baseline values under the null

hypothesis of label exchangeability. Classification performance was evaluated against this null

distribution with a non-parametric cluster-based permutation test (see Statistics). This test was

performed one-sided, as the relevant hypothesis was strictly directional: we expected stronger

decodability of the current visualized category within the localizer trials.

To investigate differential evidence for object vs. scene representations as a function of prior

learning during the retrieval task, we used the temporal generalization method[74]. First, we

applied the same preprocessing described above to the concatenated localizer and retrieval

data. A classifier was trained at each time point in the localizer data and applied to each time

point during retrieval, specifically for remembered and not-remembered trials. No cross-

validation was required since localizer and retrieval datasets were independent. As a metric, we

again used the AUC. Remembered and not-remembered distributions were tested against each

other again under a unidirectional hypothesis, as based on prior work, we expected stronger

reactivation of the originally learned category during remembered compared to not-

remembered trials. We additionally tested classifier performance against label-shuffled

surrogate distributions (200 permutations) for remembered and not-remembered trials,

separately for cue- and exhalation-locked retrieval data. The surrogate data underwent identical

preprocessing, with PCA components retained up to 99 % of explained variance (instead of 95

% in the main analysis) to ensure numerical stability of the classifier during repeated training on

label-shuffled data. Decoding results were then compared with their respective null

distributions to identify significant above-chance time windows[41,75] (see Supplementary Figs. 1c,

2a).

For trial-based analyses (Fig. 3b-c), decision values[41] were extracted at each time point in the

localizer and retrieval phases for each individual trial of the retrieval task (cue-centered for Fig.

3b; exhalation trough-centered for Fig. 3c). These values were then averaged across the

significant time window in the localizer (150 to 1500 ms), resulting in a trial-by-retrieval time

matrix. The modulus of the decision values was used to represent the model's confidence at

each time point of every trial, regardless of the trial's original category. For the analysis in Fig.

3b, the maximum decision value of each trial was calculated to obtain the respiratory phase at

peak reactivation.

Statistics. Behavioral retrieval data were averaged across conditions (object- and scene-

condition, pre- and post-sleep) and subjected to a two-sided dependent-sample t-test (Fig. 1b

for category comparison). To examine the effect of the sequence order of respiratory events

(Fig. 2b) on memory performance, a two-tailed paired-sample t-test was conducted. For

correlation analyses, Spearman's correlation was used. To assess the relationship between

oscillatory power (8 to 20 Hz) and memory reactivation (decision values) across exhalation

trough-locked retrieval times, correlational values were smoothed using a 250 ms moving mean

before statistical analysis. Effect sizes from statistical tests were computed by Cohen's d.

All data were collapsed across both sessions (objects and scenes) and memory tasks (pre- and

post-sleep) per participant before entering statistical analysis. Unless stated otherwise, we used

non-parametric cluster-based permutation tests implemented in FieldTrip[68], which do not rely

on the assumption of a specific underlying distribution. A dependent-samples t-test was

conducted at the sample level to identify clusters of contiguous time points across participants,

with values thresholded at $p = 0.05$. The cluster statistic was defined as the maxsum (sum of all

t-values within a cluster), and Monte Carlo simulations were performed to determine the cluster

p-value (alpha = 0.05, two-tailed, unless stated otherwise) based on the permutation

distribution. For circular analyses, data were divided into 60 distinct phase bins (bin length = 6°),

and mean values were calculated within each bin. To assess statistical significance, the time

course values within each trial were shuffled across trials 1000 times per subject and electrode,

maintaining the original respiration phase to create a null distribution for each participant. This

conservative approach ensures that the temporal characteristics of the respiratory signal is

preserved[76]. Significance was determined by non-parametric cluster-based permutation tests

for circular data.

Data availability

All data supporting the findings of this study are publicly available at Open Data LMU

(https://data.ub.uni-muenchen.de/596).

Code availability

All code related to the analyses of the manuscript is available at GitHub

(https://github.com/EstebanBT/Bullon-Tarraso-2025)

References
671
1.   Heck, D. H. *et al.* Breathing as a Fundamental Rhythm of Brain Function. *Front. Neural*

Circuits **10**, (2017).

2.   Yuan, H., Zotev, V., Phillips, R. & Bodurka, J. Correlated slow fluctuations in respiration,

EEG, and BOLD fMRI. *NeuroImage* **79**, 81–93 (2013).

3.   Karalis, N. & Sirota, A. Breathing coordinates cortico-hippocampal dynamics in mice during

offline states. *Nat Commun* **13**, 467 (2022).

4.   Schreiner, T., Petzka, M., Staudigl, T. & Staresina, B. P. Respiration modulates sleep

oscillations and memory reactivation in humans. *Nat Commun* **14**, 8351 (2023).

5.   Biskamp, J., Bartos, M. & Sauer, J.-F. Organization of prefrontal network activity by

respiration-related oscillations. *Sci Rep* **7**, 45508 (2017).

6.   Kluger, D. S. & Gross, J. Respiration modulates oscillatory neural network activity at rest.

PLoS Biol **19**, e3001457 (2021).

7.   Brændholt, M. *et al.* Breathing in waves: Understanding respiratory-brain coupling as a

gradient of predictive oscillations. *Neuroscience & Biobehavioral Reviews* **152**, 105262

(2023).

8.   Tort, A. B. L., Brankačk, J. & Draguhn, A. Respiration-Entrained Brain Rhythms Are Global

but Often Overlooked. *Trends in Neurosciences* **41**, 186–197 (2018).

9.   Tort, A. B. L., Laplagne, D. A., Draguhn, A. & Gonzalez, J. Global coordination of brain

activity by the breathing cycle. *Nat. Rev. Neurosci.* 1–21 (2025) doi:10.1038/s41583-025-

00920-7.

10. Arshamian, A., Iravani, B., Majid, A. & Lundström, J. N. Respiration Modulates Olfactory

Memory Consolidation in Humans. *J. Neurosci.* **38**, 10286–10294 (2018).

11. Zhuang, L., Zhang, B., Qin, Z. & Wang, P. Nasal Respiration is Necessary for the Generation

of γ Oscillation in the Olfactory Bulb. *Neuroscience* **398**, 218–230 (2019).

12. Schwimmbeck, F., Bullón Tarrasó, E. & Schreiner, T. A role for respiration in coordinating

sleep oscillations and memory consolidation. *Trends in Neurosciences* **0**, (2025).

13. Herrero, J. L., Khuvis, S., Yeagle, E., Cerf, M. & Mehta, A. D. Breathing above the brain

stem: volitional control and attentional modulation in humans. *Journal of Neurophysiology*

119, 145–159 (2018).

14. Del Negro, C. A., Funk, G. D. & Feldman, J. L. Breathing matters. *Nat Rev Neurosci* **19**, 351–

367 (2018).

15. Krohn, F. *et al.* The integrated brain network that controls respiration. *eLife* **12**, e83654

(2023).

16. Kluger, D. S., Balestrieri, E., Busch, N. A. & Gross, J. Respiration aligns perception with

neural excitability. *eLife* **10**, e70907 (2021).

17. Li, S. & Rymer, W. Z. Voluntary breathing influences corticospinal excitability of

nonrespiratory finger muscles. *J Neurophysiol* **105**, 512–521 (2011).

18. Park, H.-D. *et al.* Breathing is coupled with voluntary action and the cortical readiness

potential. *Nat Commun* **11**, 289 (2020).

19. Perl, O. *et al.* Human non-olfactory cognition phase-locked with inhalation. *Nat Hum*

Behav **3**, 501–512 (2019).

20. Zelano, C. *et al.* Nasal Respiration Entrains Human Limbic Oscillations and Modulates

Cognitive Function. *J. Neurosci.* **36**, 12448–12467 (2016).

21. Mizuhara, K. & Nittono, H. Effects of respiratory phases on the processing of emotional

and non-emotional visual stimuli. *Psychophysiology* **60**, e14261 (2023).

22. Nakamura, N. H., Fukunaga, M. & Oku, Y. Respiratory modulation of cognitive

performance during the retrieval process. *PLoS ONE* **13**, e0204021 (2018).

23. Heck, D. H., Kozma, R. & Kay, L. M. The rhythm of memory: how breathing shapes memory

function. *Journal of Neurophysiology* **122**, 563–571 (2019).

24. Staresina, B. P. *et al.* Hippocampal pattern completion is linked to gamma power increases

and alpha power decreases during recollection. *eLife* **5**, e17397 (2016).

25. Hanslmayr, S., Staresina, B. P. & Bowman, H. Oscillations and Episodic Memory:

Addressing the Synchronization/Desynchronization Conundrum. *Trends Neurosci* **39**, 16–

25 (2016).

26. Treder, M. S. *et al.* The hippocampus as the switchboard between perception and

memory. *Proceedings of the National Academy of Sciences* **118**, e2114171118 (2021).

27. Chen, Z. S. & Wilson, M. A. How our understanding of memory replay evolves. *Journal of*

Neurophysiology **129**, 552–580 (2023).

28. Griffiths, B. J. *et al.* Alpha/beta power decreases track the fidelity of stimulus-specific

information. *Elife* **8**, e49562 (2019).

29. Martín-Buro, M. C., Wimber, M., Henson, R. N. & Staresina, B. P. Alpha Rhythms Reveal

When and Where Item and Associative Memories Are Retrieved. *J. Neurosci.* **40**, 2510–

2518 (2020).

30. Karlsson, A. E., Wehrspaun, C. C. & Sander, M. C. Item recognition and lure discrimination

in younger and older adults are supported by alpha/beta desynchronization.

Neuropsychologia **148**, 107658 (2020).

31. Carr, M. F., Jadhav, S. P. & Frank, L. M. Hippocampal replay in the awake state: a potential

substrate for memory consolidation and retrieval. *Nat Neurosci* **14**, 147–153 (2011).

32. O'Neill, J., Pleydell-Bouverie, B., Dupret, D. & Csicsvari, J. Play it again: reactivation of

waking experience and memory. *Trends Neurosci* **33**, 220–229 (2010).

33. Li, Y., Pazdera, J. K. & Kahana, M. J. EEG decoders track memory dynamics. *Nat Commun*

15, 2981 (2024).

34. Michelmann, S., Bowman, H. & Hanslmayr, S. The Temporal Signature of Memories:

Identification of a General Mechanism for Dynamic Memory Replay in Humans. *PLoS Biol*

14, e1002528 (2016).

35. Staudigl, T., Vollmar, C., Noachtar, S. & Hanslmayr, S. Temporal-pattern similarity analysis

reveals the beneficial and detrimental effects of context reinstatement on human

memory. *J Neurosci* **35**, 5373–5384 (2015).

36. Schreiner, T., Petzka, M., Staudigl, T. & Staresina, B. P. Endogenous memory reactivation

during sleep in humans is clocked by slow oscillation-spindle complexes. *Nat Commun* **12**,

3112 (2021).

37. Sutterer, D. W., Foster, J. J., Serences, J. T., Vogel, E. K. & Awh, E. Alpha-band oscillations

track the retrieval of precise spatial representations from long-term memory. *Journal of*

Neurophysiology **122**, 539–551 (2019).

38. Staresina, B. P. & Wimber, M. A Neural Chronometry of Memory Recall. *Trends in*

Cognitive Sciences **23**, 1071–1085 (2019).

39. Tort, A. B. L., Komorowski, R., Eichenbaum, H. & Kopell, N. Measuring Phase-Amplitude

Coupling Between Neuronal Oscillations of Different Frequencies. *Journal of*

Neurophysiology **104**, 1195–1210 (2010).

40. Schreiner, T. & Staudigl, T. Electrophysiological signatures of memory reactivation in

humans. *Phil. Trans. R. Soc. B* **375**, 20190293 (2020).

41. Linde-Domingo, J., Treder, M. S., Kerrén, C. & Wimber, M. Evidence that neural

information flow is reversed between object perception and object reconstruction from

memory. *Nat Commun* **10**, 179 (2019).

42. Yackle, K. *et al.* Breathing control center neurons that promote arousal in mice. *Science*

355, 1411–1415 (2017).

43. Thunell, E. *et al.* Nasal inhalation does not improve memory of visual repetitions.

Psychophysiology **61**, e14609 (2024).

44. Schaefer, M., Hrysanidis, C., Lundström, J. N. & Arshamian, A. Phase-locked breathing does

not affect episodic visual recognition memory but does shape its corresponding ERPs.

Psychophysiology **61**, e14493 (2024).

45. Popov, T. & Staudigl, T. Cortico-ocular coupling in the service of episodic memory
formation. *Prog Neurobiol* **227**, 102476 (2023).

46. Burgess, A. P. & Gruzelier, J. H. Short duration power changes in the EEG during
recognition memory for words and faces. *Psychophysiology* **37**, 596–606 (2000).

47. Khader, P. H. & Rösler, F. EEG power changes reflect distinct mechanisms during long-term
memory retrieval. *Psychophysiology* **48**, 362–369 (2011).

48. Jafarpour, A., Fuentemilla, L., Horner, A. J., Penny, W. & Duzel, E. Replay of very early
encoding representations during recollection. *J Neurosci* **34**, 242–248 (2014).

49. Hanslmayr, S. *et al.* Prestimulus oscillations predict visual perception performance
between and within subjects. *NeuroImage* **37**, 1465–1473 (2007).

50. Wu, X. *et al.* Low-frequency brain oscillations reflect the dynamics of the oculomotor
system: a new perspective on subsequent memory effects. Preprint at
https://doi.org/10.1101/2025.07.29.667451 (2025).

51. Kluger, D. S. *et al.* Modulatory dynamics of periodic and aperiodic activity in respiration-
brain coupling. *Nat Commun* **14**, 4699 (2023).

52. Aserinsky, E. Periodic Respiratory Pattern Occurring in Conjunction with Eye Movements
during Sleep. *Science* **150**, 763–766 (1965).

53. Rassler, B. & Raabe, J. Co-ordination of breathing with rhythmic head and eye movements
and with passive turnings of the body. *Eur J Appl Physiol* **90**, 125–130 (2003).

54. Schreiner, T. *et al.* Spindle-locked ripples mediate memory reactivation during human
NREM sleep. *Nat Commun* **15**, 5249 (2024).

55. Todorova, R. & Zugaro, M. Hippocampal ripples as a mode of communication with cortical
and subcortical areas. *Hippocampus* **30**, 39–49 (2020).

56. Liu, Y., McAfee, S. S. & Heck, D. H. Hippocampal sharp-wave ripples in awake mice are
entrained by respiration. *Sci Rep* **7**, 8950 (2017).

57. Sheriff, A. *et al.* Breathing orchestrates synchronization of sleep oscillations in the human

hippocampus. *Proceedings of the National Academy of Sciences* **121**, e2405395121 (2024).

58. Criscuolo, A., Schwartze, M. & Kotz, S. A. Cognition through the lens of a body–brain

dynamic system. *Trends in Neurosciences* **45**, 667–677 (2022).

59. Buysse, D. J., Reynolds, C. F., Monk, T. H., Berman, S. R. & Kupfer, D. J. The Pittsburgh sleep

quality index: A new instrument for psychiatric practice and research. *Psychiatry Research*

28, 193–213 (1989).

60. Horne, J. A. & Ostberg, O. A self-assessment questionnaire to determine morningness-

eveningness in human circadian rhythms. *Int J Chronobiol* **4**, 97–110 (1976).

61. Dinges, D. F. & Powell, J. W. Microcomputer analyses of performance on a portable,

simple visual RT task during sustained operations. *Behavior Research Methods,*

Instruments, & Computers **17**, 652–655 (1985).

62. Konkle, T., Brady, T. F., Alvarez, G. A. & Oliva, A. Conceptual distinctiveness supports

detailed visual long-term memory for real-world objects. *Journal of Experimental*

Psychology: General **139**, 558–578 (2010).

63. Brainard, D. H. The Psychophysics Toolbox. *Spat Vis* **10**, 433–436 (1997).

64. Malach, R. *et al.* Object-related activity revealed by functional magnetic resonance

imaging in human occipital cortex. *Proc Natl Acad Sci U S A* **92**, 8135–8139 (1995).

65. Epstein, R. & Kanwisher, N. A cortical representation of the local visual environment.

Nature **392**, 598–601 (1998).

66. Antony, J. W., Ferreira, C. S., Norman, K. A. & Wimber, M. Retrieval as a Fast Route to

Memory Consolidation. *Trends Cogn Sci* **21**, 573–576 (2017).

67. Roediger, H. L. & Karpicke, J. D. Test-enhanced learning: taking memory tests improves

long-term retention. *Psychol Sci* **17**, 249–255 (2006).

68. Oostenveld, R., Fries, P., Maris, E. & Schoffelen, J.-M. FieldTrip: Open Source Software for

Advanced Analysis of MEG, EEG, and Invasive Electrophysiological Data. *Computational*

Intelligence and Neuroscience **2011**, 1–9 (2011).

69. Noto, T., Zhou, G., Schuele, S., Templer, J. & Zelano, C. Automated analysis of breathing

waveforms using BreathMetrics: a respiratory signal processing toolbox. *Chemical Senses*

43, 583–597 (2018).

70. Treder, M. S. MVPA-Light: A Classification and Regression Toolbox for Multi-Dimensional

Data. *Front. Neurosci.* **14**, 289 (2020).

71. Jackson, J. E. *A Use's Guide to Principal Components*. (Wiley, 1991).

doi:10.1002/0471725331.

72. Grootswagers, T., Wardle, S. G. & Carlson, T. A. Decoding Dynamic Brain Patterns from

Evoked Responses: A Tutorial on Multivariate Pattern Analysis Applied to Time Series

Neuroimaging Data. *J Cogn Neurosci* **29**, 677–697 (2017).

73. Lemm, S., Blankertz, B., Dickhaus, T. & Müller, K.-R. Introduction to machine learning for

brain imaging. *NeuroImage* **56**, 387–399 (2011).

74. King, J.-R. & Dehaene, S. Characterizing the dynamics of mental representations: the

temporal generalization method. *Trends in Cognitive Sciences* **18**, 203–210 (2014).

75. Kerrén, C., Linde-Domingo, J., Hanslmayr, S. & Wimber, M. An Optimal Oscillatory Phase

for Pattern Reactivation during Memory Retrieval. *Current Biology* **28**, 3383-3392.e6

(2018).

76. Hülsemann, M. J., Naumann, E. & Rasch, B. Quantification of Phase-Amplitude Coupling in

Neuronal Oscillations: Comparison of Phase-Locking Value, Mean Vector Length,

Modulation Index, and Generalized-Linear-Modeling-Cross-Frequency-Coupling. *Front*

Neurosci **13**, 573 (2019).

846
847

848
849   Acknowledgements
850   T. Sc. is supported by the Emmy Noether program of the German Research Foundation
851   (492835154).
852

853   Contributions
854   T.Sc., M.P. and B.P.S. conceived the study and designed the experiment. T.Sc. conducted the
855   experiment. E.B.T and T.Sc. analyzed the data. E.B.T., F.S, M.P., T.St., B.P.S. and T. Sc. wrote the
856   paper.
857

858   Corresponding authors
859   Correspondence to Thomas Schreiner (thomas.schreiner@psy.lmu.de)

Supplemental Information

	Objects	Scenes	t	p
Recognition [Hits] %				
pre-sleep	74.14±3.97	73.80±4.15	0.12	0.90
post-sleep	64.54±3.98	66.05±4.45	-0.60	0.56
post- relative to pre-sleep	87.31±2.85	88.68±2.34	-0.39	0.70
Recognition [Correct Rejections] %				
pre-sleep	90.00±2.40	87.41±4.70	0.60	0.56
post-sleep	88.33±3.19	84.63±5.70	0.72	0.48
post- relative to pre-sleep	97.77±1.72	96.22±3.92	0.43	0.67
Associative Memory %				
pre-sleep	52.77±4.71	51.36±4.67	0.40	0.70
post-sleep	42.75±4.88	39.08±3.84	0.87	0.40
post- relative to pre-sleep	79.28±4.83	76.43±4.05	0.38	0.71
Associative Memory [out of hits] %				
pre-sleep	70.18±3.59	69.85±4.16	0.09	0.93
post-sleep	64.83±5.38	61.92±4.60	0.52	0.61
post- relative to pre-sleep	90.52±4.82	87.19±3.69	0.55	0.59

Supplementary Table 1. Overview of memory performance. Associative memory % refers to the percentage of correctly recalled images (relative to the total number of stimuli), while associative memory [out of hits] refers to the percentage of recalled image exemplars out of correctly recognized verbs. Statistical differences between conditions (objects vs. scenes) were assessed using dependent samples t-tests (two-sided).

869
870
Supplementary Figure 1. (a) Grand everage (±SEM) of the respiratory signal locked to the cue onset (t = 0 s, dashed
line). The respiratory signal was recorded with a thermistor. A positive slope indicates inhalation, whereas a negative
slope reflects exhalation. (b) Respiratory phase did not have any effect on recognition memory. Blue / orange lines
represent the difference in accuracy with respect to the participant's overall recognition performance for the trials
that were closer to the inhalation peak / exhalation trough in each time bin (±SEM). (c) Learning-related brain patterns
(objects vs. scenes) were decodable during retrieval, only for remembered trials. Contour lines indicate the extent of
the significant cluster (p = 0.048, d = 0.7021, one-sided; corrected across times); color range (blue to yellow)
represents t-values of (not) remembered against null distributions of decoding performances.

Supplementary Figure 2. (a) Learning-related brain patterns (objects vs. scenes) were decodable during retrieval based on exhalation trough-centered EEG data, only for remembered trials. Contour lines indicate the extent of the significant cluster (p = 0.041, d = 1.0675, one-sided, corrected across times); color range (blue to yellow) represents t-values of (not) remembered against null distributions of decoding performances. (b) Coupling strength (measured as vector length) between respiration and not remembered trials' maximum decision values from the LDA model (Spearman's Rank Correlation, r = 0.2816, *p* = 0.2577).