Phenotype–genotype discordance in antimicrobial resistance profiles of Gram-negative uropathogens recovered from catheter-associated urinary tract infections in Egypt

Mohamed Eladawy[1, 2], Nathan Heslop[1], David Negus[1], Jonathan C. Thomas[1#], Lesley Hoyles[1#]

[1] Department of Biosciences, School of Science and Technology, Nottingham Trent University, Nottingham NG11 8NS, UK.

[2] Department of Microbiology and Immunology, Faculty of Pharmacy, Mansoura University, Egypt.

Mohamed Eladawy ORCiD 0000-0003-0200-0752

Nathan Heslop ORCiD 0009-0005-4369-2005

David Negus ORCiD 0000-0001-9047-4565

Jonathan Thomas ORCiD 0000-0002-1599-9123

Lesley Hoyles ORCiD 0000-0002-6418-342X

[#]Corresponding authors: jonathan.thomas@ntu.ac.uk, lesley.hoyles@ntu.ac.uk

Keywords: uropathogenic strains, bacterial genomics, resistance phenotypes, UPEC, EUCAST.

Abbreviations: AMR, antimicrobial resistance; CARD, Comprehensive Antibiotic Resistance Database; CAUTI, catheter-associated urinary tract infection; DDT, disc diffusion test; ESBL, extended spectrum β-lactamase; HAI, healthcare-associated infection; ICU, intensive care unit; MDR, multidrug-resistant; ME, major error; MLST, multilocus sequence type; VME, very major error; UPEC, uropathogenic *Escherichia coli*; UTI, urinary tract infection; XDR, extensively drug-resistant.

Data availability: The sequence data included in the study are available under BioProject PRJNA1208035.

ABSTRACT

Catheter-associated urinary tract infections (CAUTIs) are among the most common healthcare-associated infections in low- and middle-income countries (LMICs). In this study, phenotypic (EUCAST AMR profiles) and genomic data were generated for 132 isolates (67 *Escherichia coli*, 14 *Pseudomonas aeruginosa*, 11 *Klebsiella pneumoniae*, 9 *Proteus mirabilis*, 8 *Providencia* spp., 5 *Enterobacter hormaechei*, and 18 rare uropathogens) recovered from CAUTIs in an Egyptian hospital. Among the *Escherichia coli* isolates, phylogroup B2 was most represented (53.7 %), followed by B1 (19.4 %), A (11.9 %), D (7.4 %), F (5.9 %) and C (1.4 %). Several (22/132, 16.6 %) isolates were multidrug-resistant, while almost half (62/132, 46.9 %) were extensively drug-resistant. Comparison of phenotypic data with genotypic data from three different AMR-profiling tools (ResFinder, CARD, AMRFinder) highlighted phenotype–genotype discordance as an important consideration in resistome studies in Egypt, with a total concordance of 91.1 %, 85.7 %, and 80.3 % for ResFinder, CARD and AMRFinder, respectively. *Pseudomonas,* at the species level, exhibited the greatest discordance. At the antimicrobial level, meropenem was subject to greatest discordance. In addition to the findings from our comparative analyses, new AMR variants are reported for Egypt for *Pseudomonas* (*OXA-486*, *OXA-488*, *OXA-905*, *IMP-43*, *PDC-35*, *PDC-45*, *PDC-201*) and *Escherichia coli* (*TEM-176*, *TEM-190*). In summary, we show that there is phenotype–genotype discordance in AMR profiling among CAUTI isolates, highlighting the need for comprehensive approaches in resistome studies. We also show the genomic diversity of Gram-negative uropathogens contributing to disease burden in a little-studied LMIC setting.

INTRODUCTION

Urinary tract infections (UTIs) are an increasing public health problem caused by a range of uropathogens. The frequency of healthcare-associated UTIs is 12.9 %, 19.6 % and 24 %, respectively, in the United States, Europe and developing countries [1]. In 2019, bacterial antimicrobial resistance (AMR) contributed to 48,700 deaths associated with UTIs out of 541,000 cases of infectious syndromes [2]. Resistance to almost all antibiotics in healthcare-associated UTIs is above 20 %, with significant geographical variation [1]. However, meta-analysis and surveillance data are scarce from low- and middle-income countries (LMICs) [3, 4]. Egypt initiated a national surveillance program for healthcare-associated infections (HAIs) with international collaborators and national entities to establish benchmarks to outline the microbiological profiles and reduce AMR in clinical settings. Phase one of the National Action Plan showed nearly 50 % of *Escherichia* (*Esc.*) *coli* and 75 % of *Klebsiella pneumoniae* isolates from urine were resistant to quinolones and cephalosporins, respectively [5]. Across 27 countries, Egypt came in third place for total consumption of antibacterials (1355.4 tonnes) in 2020 [6].

In Egypt, catheter-associated UTIs (CAUTIs) are among the most common types of HAIs affecting patients in intensive care units (ICUs) [7-10]. Between 2011 and 2012, 15 % (69/472) of HAIs were UTIs and the rate of CAUTIs was 1.2 per 1,000 device-days [7]. Another surveillance showed that UTIs represented 15 % (406/2688) of HAIs, the rate of CAUTI development increased up to 1.9/1,000 device-days and CAUTIs represented 98.3 % of UTIs [8]. An epidemiological study (between 2011 and 2017) of carbapenem-resistant *Enterobacteriaceae* in Egyptian ICUs showed that UTIs accounted for 14.8 % (236/1598) of HAIs, with 63.8 % of UTIs catheter-related [10]. The second most common HAIs were UTIs (12.2 %) in Egyptian clinical settings [9].

There are few reports on the genomic diversity of uropathogens found in Egyptian clinical settings [11-16]. However, high rates of AMR are observed in Egypt beside documented transmission of carbapenem-resistant strains from Egypt to other countries [17-19]. We collected a range of Gram-negative uropathogens (n = 132 isolates) from an Egyptian hospital over a 9-month period, and generated genotypic and phenotypic data for them. Our aim was to investigate the AMR patterns associated with UTIs in an Egyptian clinical setting and provide evidence that could inform the development of effective antimicrobial surveillance plans and targeted infection

control policies. In addition, we assessed the agreement/concordance between phenotypic and
genotypic data to evaluate the usefulness of such data from a diagnostic perspective.

METHODS

Recovery of isolates

Anonymized isolates (n = 132) with no patient data were recovered from urinary
catheters between December 2020 and August 2021 by staff at the Urology and Nephrology
Center, Mansoura, Egypt during routine diagnostic procedures as described previously [16]. The
study of anonymized clinical isolates beyond the diagnostic requirement was approved by the
Urology and Nephrology Center, Mansoura, Egypt. No other ethical approval was required for
the use of the clinical isolates.

Antimicrobial susceptibility testing

Antimicrobial susceptibility testing was performed using the disc diffusion test (DDT) on
Mueller-Hinton agar (Oxoid Ltd, UK) [16]. Twelve different antimicrobial discs (Oxoid Ltd, UK)
were used: penicillins – piperacillin/tazobactam (30/6 μg); cephalosporins – ceftazidime (10 μg),
cefepime (30 μg); monobactams – aztreonam (30 μg); carbapenems – doripenem (10 μg),
meropenem (10 μg); aminoglycosides – tobramycin (10 μg), amikacin (30 μg); quinolones –
ciprofloxacin (5 μg), levofloxacin (5 μg); miscellaneous – nitrofurantoin (100 μg),
sulfamethoxazole/trimethoprim (1.25–23.75 μg).

Results were interpreted according to breakpoints of EUCAST guidelines version 12.0,
2022 (http://www.eucast.org/clinical_breakpoints/), with EUCAST reference strains - *Esc. coli*
ATCC 25922 and *Pse. aeruginosa* ATCC 27853 - used for quality control purposes.

DNA extraction and whole-genome sequencing

DNA was extracted from isolates and their genomes were sequenced and assembled as
described previously [16, 20].

Bioinformatic analyses

Bakta v1.5.1 (database v4.0) was used for annotating genes within genomes [21]. sourmash

v4.8.4 [22] was used to compare and identify (with Genome Taxonomy Database release 214)

species for all genomic data (**Supplementary Table 1**). Genomes that could not be assigned to a

species by sourmash were identified using ribosomal multilocus sequence type (rMLST)

(https://pubmlst.org/species-id; [23]), and by comparison with relevant reference sequences

(**Supplementary Table 1**) using FastANI v1.33 [24].

Serotypes were determined using ECTyper v1.0.0 (database v1.0) [25]. Phylogrouping of

Esc. coli was achieved using Ezclermont v0.7.0 [26]. MLST of each isolate was determined using

mlst v2.23.0 (https://github.com/tseemann/mlst) and relevant schema from http://pubmlst.org [27-

33].

AMR genes were identified using (accessed February 2025) the Resistance Gene

Identifier v6.0.3 of the Comprehensive Antibiotic Resistance Database (CARD) v4.0.0 [34],

ResFinder v4.6.0 (ResFinder database v2.4.0, PointFinder database v4.1.1) [35], and AMRFinder

v4.0.3 (database version 2024-12-18.1) [36]. Genes responsible for efflux pumps were excluded

from this study [37]. Only AMR genes that showed perfect or strict matches (identity ≥ 90 %) were

reported.

For the *Esc. coli* genomic data, a phylogenetic tree was constructed using core genes

identified using Panaroo v1.3.4 [38] (strict mode, identity threshold 98 %, alignment option

"core"). Core genes were present in all 67 *Esc. coli*, and were aligned with prank (v170427) [39].

Gene alignments were concatenated with FASconCAT-G (v1.02) into a single alignment [40].

Phylogenetic analysis was performed using the concatenated sequences and RaxML v1.2.0

(GTR+G8+IO) [41]. Node support was determined using 1000 non-parametric bootstrap replicates.

The phylogenetic tree was visualized using iTOL v6.9 [42]. Virulence traits encoded in genomes

were detected via BLASTP (≥90 % coverage, ≥70 % identity) with the core protein sequence

dataset from the Virulence Factors Database (VFDB) [43].

Comparison of AMR genotypes and phenotypes

According to EUCAST guidelines, "*susceptible, standard dosing regimen*" (S) and "*susceptible, increased exposure*" (I) categories are grouped under the term "*susceptible*". Using CARD, ResFinder, and AMRFinder, genomic data of isolates were compared with DDT results for 132 isolates against 12 antimicrobials [except when certain species lacked breakpoints; i.e. nitrofurantoin was used only for *Esc. coli* (n = 67)*,* sulfamethoxazole/trimethoprim was used only for *Enterobacterales* (n = 114)]. For each combination, concordance was considered positive if a) WGS data were predicted to encode AMR genes and the isolate had a phenotypic resistant profile "true resistant" (WGS-R/DDT-R) or b) WGS data were not predicted to encode AMR genes and the isolate had a phenotypic susceptible profile "true susceptible" (WGS-S/DDT-S) [44, 45]. Discordance was considered positive where major errors (MEs) or very major errors (VMEs) were detected. MEs (i.e. false positives; WGS-R/DDT-S) are defined as a resistant genotype and susceptible phenotype. VMEs (i.e. false negatives; WGS-S/DDT-R) are defined as a susceptible genotype and resistant phenotype. To detect AMR gene mutations, PointFinder tool for ResFinder and Point mutations tool for AMRFinder results were also included in concordance analyses.

RESULTS AND DISCUSSION

The main aim of this study was to characterize a collection of Gram-negative CAUTI-associated bacteria (n = 132 isolates) isolated in Egypt to investigate AMR genotype–phenotype concordance in relation to gene data available in widely used, public databases. We also sought to understand the genomic diversity of the dominant species contributing to CAUTIs in an Egyptian clinical setting.

Genome characterization

Summary statistics for the 132 high-quality genomes assembled in this study can be found in **Supplementary Table 1**. All genomes were assigned to the class *Gammaproteobacteria*, with the exception of Me47, which was found to represent a novel member of the family *Alcaligenaceae* (**Figure (1A)**). Among our isolates, UPEC dominated

(50.7 %, 67/132), followed by *Pseudomonas aeruginosa* (10.6 %, 14/132), *K. pneumoniae* (8.3

%, 11/132), other uropathogens (6.8 %, 9/132), *Proteus mirabilis* (6.8 %, n = 9/132),

Providencia spp. (6 %, n = 8/132), *Enterobacter hormaechei* (3.7 %, n = 5/132), *Alcaligenes* spp.

(3 %, n = 4/132), *Serratia* spp. (2.2 %, n = 3/132) and *Pseudomonas vlassakiae* (1.51 %, n =

2/132). UPEC was previously reported as the main cause of UTIs for adults and neonates in

Egypt, followed by other *Enterobacterales* or Gram-positive bacteria [46-49]. *Providencia* – once

considered a rare pathogen – is increasingly recognized as an opportunistic pathogen capable of

causing serious UTIs [50]; we found it among the main taxa represented in our study.

170 Antimicrobial susceptibility of isolates

According to Egyptian Urological guidelines, symptomatic CAUTIs are treated like

complicated UTIs with second- or third-generation cephalosporins, while uncomplicated

pyelonephritis is treated with a short course of fluoroquinolones as the first-line therapy [51]. AMR

for quinolones (ciprofloxacin, levofloxacin) and cephalosporins (cefepime, ceftazidime) was >64

% among MDR uropathogens in Egyptian clinical settings [52]. More than half of UPEC isolates

(57.8 %) showed high-level ciprofloxacin resistance and *gyrA* mutations were detected in 76.7 %

of isolates in a previous study, leading to recommendations for revision of empirical antibiotic

treatment of UTIs [53].

Many of our isolates were resistant to ceftazidime (68.9 %, 91/132) and ciprofloxacin

(62.1 %, 82/132). Few isolates were resistant to nitrofurantoin (5.9 %, 4/67) and meropenem

(18.2 %, 24/132) (**Figure (1B)**; **Supplementary Table 2**). The prevalence of susceptible, R1 and

R2 categories [54, 55] were 19.6 %, 7.6 % and 9.1 % respectively; 16.6 % (n = 22/132) of isolates

were MDR, while 46.9 % of isolates (n=62/132) were XDR (**Figure (1C)**). *Ent. hormaechei* (60

%, 3/5) and *K. pneumoniae* (100 %, n = 11) were associated with the highest prevalence of MDR

and XDR, respectively.

187 AMR determinants

A wide range of AMR determinants for β-lactams, quinolones, and aminoglycosides have

been reported in different uropathogens [56-59]. The main AMR genes encoded in our genomes

were predicted using ResFinder, CARD and AMRFinder (**Supplementary Table 3**). New AMR

variants are reported here for uropathogens in Egypt: *OXA-486*, *OXA-488*, *OXA-905*, *IMP-43*,

PDC-35, *PDC-45*, and *PDC-201* for *Pse. aeruginosa*, and *TEM-176* and *TEM-190* for *Esc. coli*.

WGS-based diagnostics may soon take over phenotypic testing for surveillance purposes,

particularly in situations where the low error rate has minimal consequences [60]. Comparison of

WGS data and DDT results (with respect to predicted AMR genes and empirical resistance

phenotypes) yielded a total concordance of 91.1 %, 85.7 % and 80.3 % for ResFinder, CARD

and AMRFinder, respectively. The highest concordance was recorded for *Alcaligenes* (n = 4)

species: 97.7 %, AMRFinder; 97.2 %, ResFinder; 100 % CARD (**Supplementary Table 4**).

Variations in discordance were detected for different databases, where the MEs (WGS-

R/DDT-S) were 3.9 %, 11.9 % and 15.7 %, while VMEs (WGS-S/DDT-R) were 4.9 %, 2.2 %

and 3.8% for ResFinder, CARD, and AMRFinder, respectively (**Figure (2)**). The use of an

ineffective therapeutic agent in treatment due to a VME could result in treatment failure, while

an ME may restrict therapeutic choices and create complications in the treatment process. The

more severe impact of VMEs is evident in FDA regulations for the approval of diagnostic tests,

where the FDA mandates that VMEs be kept below 1.5 % and MEs below 3 % for the approval

of a new AMR diagnostic test or device [61].

Concordance for aminoglycosides, quinolones, and miscellaneous antibiotics exceeded

89.3 % in all databases. Meropenem was associated with highest discordance in AMRFinder

(48.4 %) and CARD (47.7 %), but showed 10.6 % discordance in ResFinder. The range of total

discordance of β-lactam antibiotics for the three databases was 10.6–20.4 % ResFinder, 9–47.7

% CARD and 15.1–48.4 % AMRFinder (**Supplementary Table 5**).

In a comparison of CARD and ResFinder for two global collections and a total of 2,587

isolates comprising five pathogens of medical importance, ME rates were higher in CARD (42

%) than ResFinder (25 %). However, CARD showed almost no VMEs (1.1 %) compared with

ResFinder (4.4 %) [37]. In comparison, another report showed an overall concordance of 91 % for

ResFinder while ME and VME rates were 6.2 % and 2.1 %, respectively, for 488 different

isolates, where *Pse. aeruginosa* had the highest percentage of discordant results (44.4 %) [44]. In

agreement with previous studies [16, 44, 60], *Pseudomonas* showed the highest discordance here, and

across the three databases used in our study: 31.2 % for AMRFinder, 28.9 % for ResFinder, and
21.2 % for CARD.

In this study, 34.5 % of MEs tested phenotypically "*susceptible, increased exposure*" and
were therefore considered as not resistant due to the EUCAST update of susceptibility definitions
2019 "*by the pooling of S and I isolates together into one category*" [62]. Notably in another study,
22.7 % of MEs tested phenotypically as "*susceptible at higher exposure*" in 234 non-duplicate
Esc. coli strains [45]. Discordant bioinformatic predictions of AMR from WGS have previously
been shown through analysis of seven clinical isolates with three different databases (ResFinder,
CARD and ARG-ANNOT) and where the overall ME "false positives" and VME "false
negatives" were 20 % and 14 %, respectively [63]. In the current study, the overall MEs and VME
in three databases were 10.9 % and 3.6 %, respectively. The reduction in MEs and VMEs
compared to earlier study [63] may in part reflect improvements in curation and coverage of AMR
genes through regular database updates over time. The high MEs rate may be also due to low
expression levels of enzymes, insufficient understanding of gene regulation, mutations in
intergenic regions and *de novo* rRNA mutations.

ResFinder/PointFinder (overall concordance, 91.1 %) showed better *in silico* AMR
prediction than CARD (85.7 %) and AMRFinder with Point mutations (80.3 %) (**Figure (2)**).
PointFinder tackles chromosome mutation-mediated resistance only in *Esc. coli* and *Klebsiella.*
A combined AMR prediction from ResFinder and PointFinder for ciprofloxacin was shown to
result in improved prediction performance and a reduced VME rate [37].

For accurate *in silico* AMR predictions, our understanding of phenotypic–genotypic
correlations must be improved to reach FDA requirements for clinical microbiology diagnostic
testing. Excluding genes associated with efflux pump mechanisms from detected AMR genes has
been recommended [37]. Identity thresholds for AMR gene detection should be ≥ 90 % for different
databases [63]. Point mutations leading to AMR should be considered alongside acquired resistance
determinants, especially if they are reported from specific clinical sources such as urine, as the
most common site for discordance was urine for both *Enterobacterales* and *Pse. aeruginosa*, and
mutations in *gyrA* are associated with quinolone resistance of *Enterobacteriaceae* in urine [64-66].
Databases should replace general terminology (β-lactams, quinolones, etc.) of antimicrobial class
as a prediction to confer resistance to all members of the same antimicrobial class, so *in silico*

AMR prediction should be tailored to antibiotics themselves rather than antibiotic classes to
avoid MEs. For example, the aminoglycoside gene *aac(6′)-1* leads to amikacin and tobramycin
resistance, while *aac(3)-IIa* leads to gentamicin and tobramycin resistance [67]. This was reported
recently as poor correlation (85.9 %) for aminoglycosides between EUCAST and WGS data
using ARG-ANNOT and CARD databases, and the authors developed a phenotype-based
algorithm to reflect mechanisms responsible for the corresponding aminoglycoside resistance
phenotypes [68].

Curation of *in silico* databases is required. Some AMR genes listed in databases are not
truly responsible for AMR. For example, the *crpP* gene identified by ResFinder and AMRFinder
was previously known as a novel ciprofloxacin-modifying enzyme [69]. However, recent studies
showed that *crpP* presence is not always associated with ciprofloxacin resistance [70-72]. In this
study *crpP* was responsible for 7/8 MEs generated by AMRFinder and all seven MEs generated
by ResFinder when analysing ciprofloxacin resistance. *crpP* was detected in only one strain
(Me73B) resistant to ciprofloxacin and was the only quinolone resistance determinant detected
by ResFinder for that strain. However, AMRFinder also detected a point mutation in *gyrA*, T83I,
which most likely is the true explanation of ciprofloxacin resistance in Me73B.

Variations between databases can also be found. For example, the AMR phenotype
associated with the gene *armA* by ResFinder was amikacin and tobramycin, while AMRFinder
predicted resistance to gentamicin. Moreover, the phenotypic prediction for *aac(6′)-Ic* in
ResFinder and CARD is resistance to amikacin and tobramycin, while AMRFinder only provides
aminoglycoside as subclass.

The *smeF* gene in *Stenotrophomonas maltophilia* is responsible for quinolones resistance
[73] and was detected by all databases. However, its classification varied: *smeF* was excluded from
both CARD and ResFinder as it is considered an efflux pump gene, whereas AMRFinder
included it as a resistance determinant for quinolones. This discrepancy contributed to the
difference in MEs between ResFinder and AMRFinder for ciprofloxacin in our dataset, with
ResFinder showing seven MEs and AMRFinder eight MEs (**Figure (2)**). Although both
ResFinder and AMRFinder identified the gene from the exact same contig region with identical
coverage, the predicted gene names and associated resistance prediction differed in isolate Me21.

ResFinder annotated the gene as *aac(6')-Ib3*, predicting resistance to amikacin and tobramycin,

while AMRFinder identified it as *aac(6')-Ib*, predicting resistance to gentamicin.

Detailed analyses of UPEC isolates

Pangenomic approaches highlight genetic diversity, revealing adaptations, virulence

factors and resistance traits among collections of genomes. Several pangenome studies of UPEC

strains have been performed for both global collections [74] and individual countries, including

Ireland and Bangladesh [75, 76], with the smallest pangenome consisting of at least 16,797 genes in

total, the number of core genes ranging from 2,926 to 2,945, and the number of unique genes

ranging from 2,900 to 15,266. The UPEC pangenome (n = 67 isolates) here consisted of 10,453

genes: 3,136 (30 %) core; 6,303 (60.3 %) accessory; 1014 (9.7 %) unique.

UPEC isolates from phylogroup B2 were responsible for most of the reported UTIs (n =

36/67, 53.7 %) **(Figure (3))**, in agreement with a previous Egyptian study [77]. We identified 29

different serotypes (**Supplementary Table 1**), with phyologroup B2 serotypes O25:H4 (n =

12/67, 17.9 %) and O75:H5 (n = 11/67, 16.4 %) most common. Although phylogroup B2 strains

are uncommon among the commensal microbiota, they are highly virulent when present; B2

strains have been shown to persist in the intestinal microbiota of infants [78], while also being

associated with wildlife and livestock [79], and herbivorous and omnivorous mammals [80]. A

previous Egyptian study showed the most prevalent phylogroup in the country was A followed

by B2 [81]. Other phylogroups detected by us were B1 (19.4 %), A (11.9 %), D (7.4 %), F (5.9 %)

and C (1.4 %).

The principal virulence genes associated with all UPEC strains (n = 67) were type 1

fimbriae (*fimH*), outer membrane protein (*ompA*) and enterobactin (*entS*). Curli fibers (*csgA* 98

%, n = 66), common pilus (*ecp* 95 %, n = 64), nutritional and metabolic factors (*fyuA* 79 %, n =

53, *chuA* 67 %, n = 45), K1 capsule type (*kpsM* 66 %, n = 44), aerobactin (*iutA* 60 %, n = 40), P

fimbriae (*papC* 52 %, n = 35, *papG* 33 %, n = 22), α-hemolysin and iron siderophores (*hlyA* and

iroN 24 %, n = 16), exotoxins (*cnf1* 13 %, n = 9), and S fimbriae and F1C fimbriae (*sfa* and *foc* 6

%, n = 4) were also represented (**Figure (3)**; **Supplementary Table 6**). *fimH* and *sfaX* were

present in all phylogroups; all other *sfa* genes were only associated with phylogroup B2. *entD*

was only in phylogroup B1. *cnf-1* was present in all phylogroup B2 isolates and only one isolate
(Me87) from B1.

fim, ecp and *csg* gene families have previously been found to be core to the UPEC
pangenome [76]. In contrast, a PCR-based study from Mansoura reported a much higher prevalence
of afimbrial adhesins such as *afa/dra* (14 %) and S-fimbriae (*sfaS*, 60.6 %) [82], which were largely
absent from our dataset. Additionally, our data show a higher prevalence of P fimbriae genes
(*papG*, 33 %; *papC*, 52 %) compared to [82], where genes were found only in only 21.3 % and 49.3
% of isolates, respectively. Similarly, our detection of iron acquisition genes *iutA* (60 %) and
chuA (67 %) exceeds their reported values of 34.6 % and 54 % [82]. Notably, *cnf1* (cytotoxic
necrotizing factor 1) was much rarer in our dataset (13 %) compared to 42 % in [82], suggesting
potential differences in strain pathogenicity. Overall, our results highlight a strong presence of
core adhesion and iron acquisition systems in UPEC strains, with notable variations in toxin-
associated genes compared to previous studies. These findings emphasize the genetic diversity of
UPEC virulence factors and the importance of comparing detection methods when assessing
pathogenic potential.

The β-lactam resistance genes bla_{OXA-1} and bla_{TEM-1} were present in all phylogroups,
$bla_{TEM-176}$ was restricted to phylogroup B1, while other bla_{TEM} variants were present in only
phylogroup A. *dha* gene variants were restricted to phylogroup B2, while bla_{NDM-5} was absent
from B2. $bla_{CTX-M-15}$ was present in all phylogroups, while $bla_{CTX-M-14}$ and $bla_{CTX-M-16}$ were
present in phylogroup F only. In Egypt, $bla_{CTX-M-15}$ is the most predominant ESBL-producing
type among uropathogenic and diarrheagenic *Esc. coli* strains [83]. In terms of quinolone
resistance, *qnrB4* was present in phylogroup B2 only, while *qnrS1* was present in all
phylogroups. Mutations of *gyrA* and *parC* genes were present in all phylogroups. Genes of
aminoglycoside-modifying enzymes were present in all phylogroups.

UPEC isolates were most susceptible to nitrofurantoin (3 % resistance). Nitrofurantoin
was reported as a drug of choice for treating UTIs in previous Egyptian studies [84, 85], but it has
therapeutic limitations due to side-effects in certain patient groups [86]. *In silico* AMR prediction
using CARD showed 95.5 % concordance with phenotypic results (**Figure (2)**). In a recent
Egyptian study, the resistance level to nitrofurantoin was 51 % in *Klebsiella* spp., but it remains
effective for *Esc. coli* UTIs in males and females (92 % sensitivity). In comparison, *Esc. coli* and

Klebsiella spp. showed elevated resistance rates to cephalosporins: 75 % and 81 %, respectively
[87].

Conclusion

Our study provides crucial insights into the genomic and AMR landscape of CAUTI-
associated uropathogens in Egypt, highlighting the importance of understanding AMR
phenotype–genotype discordance in this setting. These findings reinforce the need for integrated
phenotypic and genomic surveillance strategies to improve AMR diagnostics, guide treatment
decisions, and inform infection control policies in resource-limited settings.

ACKNOWLEDGEMENTS

We would like to Dr Essam Elsawy and staff of Urology and Nephrology Centre,
Mansoura, Egypt for providing the clinical isolates used in this study.

ME – did all phenotypic work; extracted DNA for sequencing; all bioinformatics
associated with clinical isolates; characterized the AMR and virulence genes encoded by the
isolates and their plasmids; MLST analysis and summary; interpreted virulence and AMR data;
pangenome analysis. JCT – pangenome analysis and associated supervision. NH, DN – library
preparation and genome sequencing of some clinical isolates; LH – sourmash analyses;
supervised the study. ME, JCT and LH wrote the original version of the manuscript, and all
authors approved the final version.

FUNDING

This work was supported by The Egyptian Ministry of Higher Education & Scientific
Research represented by The Egyptian Bureau for Cultural & Educational Affairs in London. NH
was funded through a placement studentship provided by Department of Biosciences,
Nottingham Trent University. Computing resources used in this study were funded through the
Research Contingency Fund of Nottingham Trent University.

TRANSPARENCY DECLARATIONS

The authors declare that there are no conflicts of interest.

REFERENCES

1. Tandogdu Z, Wagenlehner FME. Global epidemiology of urinary tract infections. *Current Opinion in Infectious Diseases* 2016; **29**: 73-9.

2. European Antimicrobial Resistance C. The burden of bacterial antimicrobial resistance in the WHO European region in 2019: a cross-country systematic analysis. *Lancet Public Health* 2022; **7**: e897-e913.

3. Allegranzi B, Bagheri Nejad S, Combescure C et al. Burden of endemic health-care-associated infection in developing countries: systematic review and meta-analysis. *Lancet (London, England)* 2011; **377**: 228-41.

4. Öztürk R, Murt A. Epidemiology of urological infections: a global burden. *World Journal of Urology* 2020; **38**: 2669-79.

5. WHO. Global antimicrobial resistance and use surveillance system (GLASS) report: early implementation 2016-2017. In: Organization WH, ed: World Health Organization, 2018.

6. WHO. Global antimicrobial resistance and use surveillance system (GLASS) report: 2022. World Health Organization, 2022.

7. See I, Lessa FC, ElAta OA et al. Incidence and pathogen distribution of healthcare-associated infections in pilot hospitals in Egypt. *Infect Control Hosp Epidemiol* 2013; **34**: 1281-8.

8. Talaat M, El-Shokry M, El-Kholy J et al. National surveillance of health care-associated infections in Egypt: Developing a sustainable program in a resource-limited country. *Am J Infect Control* 2016; **44**: 1296-301.

9. El-Kholy A, El-Mahallawy HA, Elsharnouby N et al. Landscape of Multidrug-Resistant Gram-Negative Infections in Egypt: Survey and Literature Review. *Infect Drug Resist* 2021; **14**: 1905-20.

10. Kotb S, Lyman M, Ismail G et al. Epidemiology of Carbapenem-resistant Enterobacteriaceae in Egyptian intensive care units using National Healthcare-associated Infections Surveillance Data, 2011-2017. *Antimicrob Resist Infect Control* 2020; **9**: 2.

11. Zakaria AS, Edward EA, Mohamed NM. Genomic Insights into a Colistin-Resistant Uropathogenic Escherichia coli Strain of O23:H4-ST641 Lineage Harboring mcr-1.1 on a Conjugative IncHI2 Plasmid from Egypt. *Microorganisms* 2021; **9**.

12. Zakaria AS, Edward EA, Mohamed NM. Pathogenicity Islands in Uropathogenic Escherichia coli Clinical Isolate of the Globally Disseminated O25:H4-ST131 Pandemic Clonal Lineage: First Report from Egypt. *Antibiotics (Basel)* 2022; **11**.

13. Sherif M, Palmieri M, Mirande C et al. Whole-genome sequencing of Egyptian multidrug-resistant Klebsiella pneumoniae isolates: a multi-center pilot study. *European Journal of Clinical Microbiology & Infectious Diseases* 2021; **40**: 1451-60.

14. Edward EA, Mohamed NM, Zakaria AS. Whole Genome Characterization of the High-Risk Clone ST383 Klebsiella pneumoniae with a Simultaneous Carriage of bla(CTX-M-14) on IncL/M Plasmid and bla(CTX-M-15) on Convergent IncHI1B/IncFIB Plasmid from Egypt. *Microorganisms* 2022; **10**.

15. Baiomy AA, Serry FE, Kadry AA et al. Genome Analysis of Pseudomonas aeruginosa Strains from Chronically Infected Patients with High Levels of Persister Formation. *Pathogens* 2023; **12**.

16. Eladawy M, Thomas JC, Hoyles L. Phenotypic and genomic characterization of Pseudomonas aeruginosa isolates recovered from catheter-associated urinary tract infections in an Egyptian hospital. *Microb Genom* 2023; **9**.

17. Kohler P, Tijet N, Kim HC et al. Dissemination of Verona Integron-encoded Metallo-beta-lactamase among clinical and environmental Enterobacteriaceae isolates in Ontario, Canada. *Sci Rep* 2020; **10**: 18580.

18.   van Duin D, Perez F, Rudin SD et al. Surveillance of carbapenem-resistant Klebsiella pneumoniae:
tracking molecular epidemiology and outcomes through a regional network. *Antimicrob Agents*
Chemother 2014; **58**: 4035-41.
19.   Hassuna NA, Khairalla AS, Farahat EM et al. Molecular characterization of Extended-spectrum β
lactamase- producing E. coli recovered from community-acquired urinary tract infections in Upper Egypt.
Sci Rep 2020; **10**: 2772.
20.   Negus D, Foster G, Hoyles L. Lelliottia amnigena recovered from the lung of a harbour porpoise,
and comparative analyses with Lelliottia spp. *Access Microbiol* 2023; **5**.
21.   Schwengers O, Jelonek L, Dieckmann MA et al. Bakta: rapid and standardized annotation of
bacterial genomes via alignment-free sequence identification. *Microb Genom* 2021; **7**.
22.   Pierce NT, Irber L, Reiter T et al. Large-scale sequence comparisons with sourmash. *F1000Res*
2019; **8**: 1006.
23.   Jolley KA, Bliss CM, Bennett JS et al. Ribosomal multilocus sequence typing: universal
characterization of bacteria from domain to strain. *Microbiology (Reading, England)* 2012; **158**: 1005-15.
24.   Jain C, Rodriguez-R LM, Phillippy AM et al. High throughput ANI analysis of 90K prokaryotic
genomes reveals clear species boundaries. *Nature communications* 2018; **9**: 1-8.
25.   Bessonov K, Laing C, Robertson J et al. ECTyper: in silico Escherichia coli serotype and species
prediction from raw and assembled whole-genome sequence data. *Microbial genomics* 2021; **7**.
26.   Beghain J, Bridier-Nahmias A, Le Nagard H et al. ClermonTyping: an easy-to-use and accurate in
silico method for Escherichia genus strain phylotyping. *Microbial genomics* 2018; **4**.
27.   Jolley KA, Bray JE, Maiden MCJ. Open-access bacterial population genomics: BIGSdb software,
the PubMLST.org website and their applications. *Wellcome open research* 2018; **3**: 124.
28.   Wirth T, Falush D, Lan R et al. Sex and virulence in Escherichia coli: an evolutionary perspective.
Molecular microbiology 2006; **60**: 1136-51.
29.   Bai L, Xia S, Lan R et al. Isolation and characterization of cytotoxic, aggregative Citrobacter
freundii. *PloS one* 2012; **7**: e33054.
30.   Diancourt L, Passet V, Verhoef J et al. Multilocus sequence typing of Klebsiella pneumoniae
nosocomial isolates. *J Clin Microbiol* 2005; **43**: 4178-82.
31.   Curran B, Jonas D, Grundmann H et al. Development of a multilocus sequence typing scheme for
the opportunistic pathogen Pseudomonas aeruginosa. *J Clin Microbiol* 2004; **42**: 5644-9.
32.   Diancourt L, Passet V, Nemec A et al. The population structure of Acinetobacter baumannii:
expanding multiresistant clones from an ancestral susceptible genetic pool. *PloS one* 2010; **5**: e10034.
33.   Miyoshi-Akiyama T, Hayakawa K, Ohmagari N et al. Multilocus sequence typing (MLST) for
characterization of Enterobacter cloacae. *PloS one* 2013; **8**: e66358.
34.   McArthur AG, Waglechner N, Nizam F et al. The comprehensive antibiotic resistance database.
Antimicrob Agents Chemother 2013; **57**: 3348-57.
35.   Florensa AF, Kaas RS, Clausen P et al. ResFinder - an open online resource for identification of
antimicrobial resistance genes in next-generation sequencing data and prediction of phenotypes from
genotypes. *Microbial genomics* 2022; **8**.
36.   Feldgarden M, Brover V, Haft DH et al. Validating the AMRFinder Tool and Resistance Gene
Database by Using Antimicrobial Resistance Genotype-Phenotype Correlations in a Collection of Isolates.
Antimicrob Agents Chemother 2019; **63**.
37.   Mahfouz N, Ferreira I, Beisken S et al. Large-scale assessment of antimicrobial resistance marker
databases for genetic phenotype prediction: a systematic review. *Journal of Antimicrobial Chemotherapy*
2020; **75**: 3099-108.
38.   Tonkin-Hill G, MacAlasdair N, Ruis C et al. Producing polished prokaryotic pangenomes with the
Panaroo pipeline. *Genome biology* 2020; **21**: 180.
39.   Loytynoja A. Phylogeny-aware alignment with PRANK. *Methods Mol Biol* 2014; **1079**: 155-70.

40. Kuck P, Longo GC. FASconCAT-G: extensive functions for multiple sequence alignment
preparations concerning phylogenetic studies. *Front Zool* 2014; **11**: 81.
41. Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
phylogenies. *Bioinformatics (Oxford, England)* 2014; **30**: 1312-3.
42. Letunic I, Bork P. Interactive Tree Of Life (iTOL) v4: recent updates and new developments.
Nucleic Acids Res 2019; **47**: W256-W9.
43. Chen L, Yang J, Yu J et al. VFDB: a reference database for bacterial virulence factors. *Nucleic Acids*
Res 2005; **33**: D325-8.
44. Rebelo AR, Bortolaia V, Leekitcharoenphon P et al. One Day in Denmark: Comparison of
Phenotypic and Genotypic Antimicrobial Susceptibility Testing in Bacterial Isolates From Clinical Settings.
Front Microbiol 2022; **13**: 804627.
45. Vanstokstraeten R, Pierard D, Crombe F et al. Genotypic resistance determined by whole genome
sequencing versus phenotypic resistance in 234 Escherichia coli isolates. *Sci Rep* 2023; **13**: 449.
46. Mohamed ES, Khairy RMM, Abdelrahim SS. Prevalence and molecular characteristics of ESBL and
AmpC β -lactamase producing Enterobacteriaceae strains isolated from UTIs in Egypt. *Antimicrob Resist*
Infect Control 2020; **9**: 198.
47. Shrief R, El-Ashry AH, Mahmoud R et al. Effect of Colistin, Fosfomycin and
Meropenem/Vaborbactam on Carbapenem-Resistant Enterobacterales in Egypt: A Cross-Sectional Study.
Infect Drug Resist 2022; **15**: 6203-14.
48. Baz AMK, El-Agamy OAE, Ibrahim AM. Incidence of urinary tract infection in neonates with
significant indirect Hyperbilirubinemia of unknown etiology: case-control study. *Ital J Pediatr* 2021; **47**:
35.
49. Hassan R, El-Gilany AH, Abd Elaal AM et al. An overview of healthcare-associated infections in a
tertiary care hospital in Egypt. *Infect Prev Pract* 2020; **2**: 100059.
50. Rajni E, Jain A, Garg VK et al. Providencia Causing Urinary Tract Infections: Are We Reaching a
Dead End? *Indian J Crit Care Med* 2022; **26**: 446-51.
51. Nouh K, Kasem A, Shaher H et al. *THE EGYPTIAN UROLOGICAL GUIDELINES, First Edition, Chapter*
XI: Urinary Tract Infections Guidelines., 2021.
52. Elshamy AA, Aboshanab KM, Yassien MA et al. Prevalence of plasmid-mediated resistance genes
among multidrug-resistant uropathogens in Egypt. *African health sciences* 2020; **20**: 190-8.
53. Esmaeel NE, Gerges MA, Hosny TA et al. Detection of Chromosomal and Plasmid-Mediated
Quinolone Resistance Among Escherichia coli Isolated from Urinary Tract Infection Cases; Zagazig
University Hospitals, Egypt. *Infect Drug Resist* 2020; **13**: 413-21.
54. Magiorakos AP, Srinivasan A, Carey RB et al. Multidrug-resistant, extensively drug-resistant and
pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired
resistance. *Clin Microbiol Infect* 2012; **18**: 268-81.
55. Bitew A, Tsige E. High Prevalence of Multidrug-Resistant and Extended-Spectrum beta-
Lactamase-Producing Enterobacteriaceae: A Cross-Sectional Study at Arsho Advanced Medical
Laboratory, Addis Ababa, Ethiopia. *J Trop Med* 2020; **2020**: 6167234.
56. Jomehzadeh N, Saki M, Ahmadi K et al. The prevalence of plasmid-mediated quinolone
resistance genes among Escherichia coli strains isolated from urinary tract infections in southwest Iran.
Mol Biol Rep 2022; **49**: 3757-63.
57. Tickler IA, Kawa D, Obradovich AE et al. Characterization of Carbapenemase- and ESBL-Producing
Gram-Negative Bacilli Isolated from Patients with Urinary Tract and Bloodstream Infections. *Antibiotics*
(Basel) 2023; **12**.
58. de Sousa T, Hebraud M, Alves O et al. Study of Antimicrobial Resistance, Biofilm Formation, and
Motility of Pseudomonas aeruginosa Derived from Urine Samples. *Microorganisms* 2023; **11**.

59. Zeng Q, Xiao S, Gu F et al. Antimicrobial Resistance and Molecular Epidemiology of
Uropathogenic Escherichia coli Isolated From Female Patients in Shanghai, China. *Frontiers in cellular and*
infection microbiology 2021; **11**: 653983.
60. Ellington M, Ekelund O, Aarestrup FM et al. The role of whole genome sequencing in
antimicrobial susceptibility testing of bacteria: report from the EUCAST Subcommittee. *Clinical*
microbiology and infection 2017; **23**: 2-22.
61. Jorgensen JH, Ferraro MJ. Antimicrobial susceptibility testing: a review of general principles and
contemporary practices. *Clinical infectious diseases : an official publication of the Infectious Diseases*
Society of America 2009; **49**: 1749-55.
62. Nabal Diaz SG, Algara Robles O, Garcia-Lechuz Moya JM. New definitions of susceptibility
categories EUCAST 2019: clinic application. *Rev Esp Quimioter* 2022; **35 Suppl 3**: 84-8.
63. Doyle RM, O'Sullivan DM, Aller SD et al. Discordant bioinformatic predictions of antimicrobial
resistance from whole-genome sequencing data of bacterial isolates: an inter-laboratory study. *Microb*
Genom 2020; **6**.
64. Shenagari M, Bakhtiari M, Mojtahedi A et al. High frequency of mutations in gyrA gene
associated with quinolones resistance in uropathogenic Escherichia coli isolates from the north of Iran.
Iran J Basic Med Sci 2018; **21**: 1226-31.
65. Ku PM, Hobbs DA, Gilmore M et al. 1234. Can Susceptibility to One Carbapenem be Conferred to
Another? Frequency of Discordance in Gram-negative Clinical Isolates. *Open Forum Infectious Diseases*,
p. S706-S7, Oxford University Press US.
66. Thompson D, Xu J, Ischia J et al. Fluoroquinolone resistance in urinary tract infections:
Epidemiology, mechanisms of action and management strategies. *BJUI Compass* 2024; **5**: 5-11.
67. Krause KM, Serio AW, Kane TR et al. Aminoglycosides: An Overview. *Cold Spring Harb Perspect*
Med 2016; **6**.
68. Mancini S, Marchesi M, Imkamp F et al. Population-based inference of aminoglycoside resistance
mechanisms in Escherichia coli. *EBioMedicine* 2019; **46**: 184-92.
69. Chávez-Jacobo VM, Hernández-Ramírez KC, Romo-Rodríguez P et al. CrpP Is a Novel
Ciprofloxacin-Modifying Enzyme Encoded by the Pseudomonas aeruginosa pUM505 Plasmid. *Antimicrob*
Agents Chemother 2018; **62**.
70. Hernández-García M, García-Castillo M, García-Fernández S et al. Presence of Chromosomal
crpP-like Genes Is Not Always Associated with Ciprofloxacin Resistance in Pseudomonas aeruginosa
Clinical Isolates Recovered in ICU Patients from Portugal and Spain. *Microorganisms* 2021; **9**.
71. López M, Rojo-Bezares B, Chichón G et al. Resistance to Fluoroquinolones in Pseudomonas
aeruginosa from Human, Animal, Food and Environmental Origin: The Role of CrpP and Mobilizable ICEs.
Antibiotics 2022; **11**: 1271.
72. Zubyk HL, Wright GD. CrpP Is Not a Fluoroquinolone-Inactivating Enzyme. *Antimicrob Agents*
Chemother 2021; **65**: e0077321.
73. Jia W, Wang J, Xu H et al. Resistance of Stenotrophomonas maltophilia to Fluoroquinolones:
Prevalence in a University Hospital and Possible Mechanisms. *Int J Environ Res Public Health* 2015; **12**:
5177-95.
74. Fatima S, Akbar A, Irfan M et al. Virulence Factors and Antimicrobial Resistance of Uropathogenic
Escherichia coli EQ101 UPEC Isolated from UTI Patient in Quetta, Balochistan, Pakistan. *Biomed Res Int*
2023; **2023**: 7278070.
75. Whelan S, Bottacini F, Buttimer C et al. Whole genome sequencing of uropathogenic E. coli from
Ireland reveals diverse resistance mechanisms and strong correlation with phenotypic (EUCAST)
susceptibility testing. *Infect Genet Evol* 2024; **121**: 105600.
76. Hossain M, Tabassum T, Rahman A et al. Genotype-phenotype correlation of beta-lactamase-
producing uropathogenic Escherichia coli (UPEC) strains from Bangladesh. *Sci Rep* 2020; **10**: 14549.

77. Farahat EM, Hassuna NA, Hammad AM et al. Distribution of integrons and phylogenetic groups
among Escherichia coli causing community-acquired urinary tract infection in Upper Egypt. *Canadian*
Journal of Microbiology 2021; **67**: 451-63.
78. Nowrouzian FL, Adlerberth I, Wold AE. Enhanced persistence in the colonic microbiota of
Escherichia coli strains belonging to phylogenetic group B2: role of virulence factors and adherence to
colonic cells. *Microbes Infect* 2006; **8**: 834-40.
79. Aguirre-Sanchez JR, Valdez-Torres JB, Del Campo NC et al. Phylogenetic group and virulence
profile classification in Escherichia coli from distinct isolation sources in Mexico. *Infect Genet Evol* 2022;
106: 105380.
80. Gordon DM, Cowling A. The distribution and genetic structure of Escherichia coli in Australian
vertebrates: host and geographic effects. *Microbiology (Reading, England)* 2003; **149**: 3575-86.
81. El-Mahdy R, Mahmoud R, Shrief R. Characterization of E. coli Phylogroups Causing Catheter-
Associated Urinary Tract Infection. *Infect Drug Resist* 2021; **14**: 3183-93.
82. El-Baz R, Said HS, Abdelmegeed ES et al. Characterization of virulence determinants and
phylogenetic background of multiple and extensively drug resistant Escherichia coli isolated from
different clinical sources in Egypt. *Appl Microbiol Biotechnol* 2022; **106**: 1279-98.
83. Abdelrahim SS, Fouad M, Abdallah N et al. Comparative Study of CTX-M-15 Producing
Escherichia coli ST131 Clone Isolated from Urinary Tract Infections and Acute Diarrhoea. *Infect Drug*
Resist 2021; **14**: 4027-38.
84. Shash RY, Elshimy AA, Soliman MY et al. Molecular Characterization of Extended-Spectrum β-
Lactamase Enterobacteriaceae Isolated from Egyptian Patients with Community- and Hospital-Acquired
Urinary Tract Infection. *Am J Trop Med Hyg* 2019; **100**: 522-8.
85. Elzayat M, Barnett-Vanes A, Dabour MF et al. Prevalence of undiagnosed asymptomatic
bacteriuria and associated risk factors during pregnancy: a cross-sectional study at two tertiary centres in
Cairo, Egypt. *BMJ Open* 2017; **7**: e013198.
86. Recht J, Chansamouth V, White NJ, Ashley EA. Nitrofurantoin and glucose-6-phosphate
dehydrogenase deficiency: a safety review. JAC Antimicrob Resist. 2022 May 3;4(3):dlac045. doi:
10.1093/jacamr/dlac045. PMID: 35529053; PMCID: PMC9070801.

87. Tarek A, Abdalla S, Dokmak NA et al. Bacterial Diversity and Antibiotic Resistance Patterns of
Community-Acquired Urinary Tract Infections in Mega Size Clinical Samples of Egyptian Patients: A Cross-
Sectional Study. *Cureus* 2024; **16**: e51838.

A)

B)

C)

Figure (1): AMR profiles of uropathogens (n = 132 isolates) included in this study. A) Dendrogram generated from comparison of

sourmash signatures for uropathogens showing their species affiliations and whether they are MDR or XDR. Phylogroup affiliations

are given for *Esc. coli* isolates. B) Resistance summary for the 132 uropathogens using seven different antimicrobial classes according

to EUCAST guidelines. (C) Data presented using the category designations suggested by [54, 55]. Penicillins: piperacillin/tazobactam.

Monobactams: aztreonam. Cephalosporins: ceftazidime and cefepime. Carbapenems: doripenem and meropenem. Quinolones:

ciprofloxacin and levofloxacin. Aminoglycosides: amikacin and tobramycin. Miscellaneous: sulfamethoxazole/trimethoprim and

nitrofurantoin.

Figure (2): AMR genotype–phenotype discordance summarized for the 132 isolates included in this study. (A, B) Percentage and number of genotype–phenotype concordances and discordances are shown, when comparing WGS data and DDT results using ResFinder, CARD, and AMRFinder. The bars are coloured according to Concordance or Discordance (refer to colour legend at the top of the figure). White numbers on top of the bars correspond to the respective number of results. (A) Species-level data, (B) antimicrobial-level data. Numbers in parentheses in (A) correspond to the number of isolates per species / the number of isolate-antimicrobial combinations. (A) Others: *Paenalcaligenes suwonensis*, *Klebsiella ornithinolytica, Citrobacter portucalensis, Leclercia adecarboxylata, Morganella morganii, Stenotrophomonas maltophilia, Acinetobacter baumannii,* and *Achromobacter xylosoxidans*. (B) Nitrofurantoin concordance was analyzed only via CARD. Levofloxacin and doripenem were not analyzed with ResFinder/PointFinder. Piperacillin/tazobactam (TZP), aztreonam (ATM), ceftazidime (CAZ), cefepime (FEP), doripenem (DOR), meropenem (MEM), ciprofloxacin (CIP), levofloxacin (LEV), amikacin (AK), tobramycin (TOB), sulfamethoxazole/trimethoprim (SXT), and nitrofurantoin (NIF). Major errors (ME) are WGS-R/DDT-S, while very major error (VME) are WGS-S/DDT-R.

Figure (3): Phylogenetic tree for 67 *Esc. coli* isolates constructed using pangenome analysis of core genes, via panaroo and RaxML. UPEC strains are grouped into clusters (A, B1, B2, C, D, F) depending on EzClermont phylogroups. The maximum-likelihood tree was annotated using iTOL, with bootstrap values expressed as a percentage of 1000 replicates. Scale bar, nucleotide substitutions per site. Virulence genes were identified through BLASTp with VFDB based on coverage >= 90 % and identity >=70 %.