**Title: An optogenetic toolkit for robust activation of FGF, BMP, & Nodal signaling in zebrafish**

**Running title:** Optogenetic signaling activation

**Key words:** molecular optogenetics, zebrafish, signaling, FGF, BMP, Nodal, embryogenesis

Leanne E. Iannucci[1†], Velanganni Selvaraj Maria Thomas[1†], Micaela R. Murphy[1], Caitlin E.T. Donahue[1], William K. Anderson[1], Catherine E. Rogers[1], Allison J. Saul[1,2], Katherine W. Rogers[1*]

[†]These authors contributed equally

[1]Unit on Developmental Signaling, Division of Developmental Biology, *Eunice Kennedy Shriver* National Institute of Child Health and Human Development, NIH, Bethesda, MD 20814, USA

[2]Current address: School of Biological Sciences, University of Utah, Salt Lake City, UT 84112, USA

[*]Correspondence: katherine.rogers@nih.gov

**ABSTRACT** Cell signaling regulates a wide range of biological processes including development, homeostasis, and disease. Accessible technologies to precisely manipulate signaling have important applications in basic and translational research. Here, we introduce an optogenetic toolkit comprised of 1) a zebrafish-optimized FGF signaling activator, 2) a single-transcript Nodal signaling activator, and 3) a previously established BMP signaling activator. We thoroughly characterize this suite of tools in zebrafish embryos and show that they provide tunable, light-dependent spatiotemporal control of signaling *in vivo*. In response to blue light (~455 nm), receptor kinase domains fused to blue light-dimerizing LOV domains enable robust signaling activation with minimal ectopic activity in the dark or at wavelengths over 495 nm. Optogenetic activation by each tool is pathway-specific and results in increased expression of known target genes. Signaling is activated with rapid on/off kinetics, and activation strength depends on light irradiance. Finally, we demonstrate spatially localized signaling activation with our optimized FGF activator. Together, our results establish this optogenetic toolkit as a potent experimental platform to rapidly, directly, and adjustably activate FGF, BMP, and Nodal signaling in zebrafish embryos.

## INTRODUCTION

Many biological processes are regulated by cell signaling, from development to homeostasis to disease. Signaling dynamics (i.e., levels and durations) play a crucial role in orchestrating diverse biological processes including embryonic patterning (Bosman and Sonnen, 2022; Hill, 2022; Kicheva and Briscoe, 2023; Sagner and Briscoe, 2017). For example, cell fate decisions can differ dramatically based on a single pathway's signaling frequency (Purvis and Lahav, 2013). Spatiotemporally dynamic signaling gradients in developing tissues are known to coordinate patterning in multiple contexts across the animal kingdom (Guillotin and Birnbaum, 2020; Hill, 2022; Jones and Mullins, 2022; Liberali and Schier, 2024; Sagner and Briscoe, 2017). During early vertebrate development, dynamic signaling gradients generated by fibroblast growth factor (FGF), bone morphogenetic protein (BMP), and Nodal are established (Hill, 2022; Jones and Mullins, 2022; Liberali and Schier, 2024). These pathways signal via receptor kinases and remain at the center of many active developmental biology investigations due to their key roles in patterning the embryonic body axes. Established methods to manipulate FGF, BMP, and Nodal signaling such as mutating pathway genes, using heat shock promoters to drive genetically encoded agonists or antagonists, recombinant proteins, and small molecule drugs have provided foundational information about their roles. However, these methods can be cumbersome and slow. Precision molecular technologies to experimentally activate receptor kinase-based signaling *in vivo* can open new avenues of study.

Molecular optogenetic strategies use light to manipulate biological processes and have yielded novel insights across fields including developmental biology (Johnson and Toettcher, 2018; Rogers and Müller, 2020). Optogenetics has been used to control processes as diverse as gene expression (LaBelle et al., 2021; Reade et al., 2017), apoptosis (Mruk et al., 2020), protein localization (Buckley, 2019; Buckley et al., 2016), and signaling (Benman et al., 2022; Čapek et al., 2019; Kainrath et al., 2017; McNamara et al., 2025; Rogers et al., 2020; Sako et al., 2016; Saul, Rogers et al., 2023). This technology has generated considerable enthusiasm, reflected in the nearly exponential increase in publications involving molecular optogenetic approaches since the early 2000s (Kolar et al., 2018). Research exploring signaling is well suited for molecular optogenetic approaches because this technology can enable precise spatiotemporal control over signaling (Farahani et al., 2021). Applying these strategies to experimentally manipulate FGF, BMP, and Nodal signaling in vertebrate embryos is appealing because optogenetic methods to

63    activate receptor kinase-regulated pathways such as these have been demonstrated (Chang et al.,

64    2014; Crossman and Janovjak, 2022; Grusch et al., 2014; Humphreys et al., 2020; Kainrath et al.,

65    2017; Kim et al., 2014; McNamara et al., 2025; Rogers et al., 2020; Sako et al., 2016).

66           Optogenetic tools have already helped reveal key insights into the roles of signaling in

development across a range of model systems (Farahani et al., 2021; Rogers and Müller, 2020),

including the vertebrate zebrafish embryo. This widely used model is well suited to address many

developmental biology questions because it is externally fertilized, translucent, and microscopy-

friendly (Hill, 2022; Jones and Mullins, 2022; Liberali and Schier, 2024; Mullins et al., 2021)—

all attributes that make zebrafish an ideal system for optogenetic approaches. Optogenetic

manipulation of Nodal signaling in zebrafish has been used to define developmental windows

during which Nodal is actively patterning the germ layers (Reade et al., 2017; Vopalensky et al.,

2018), to investigate how Nodal and BMP signaling are interpreted (Rogers et al., 2020; Sako et

al., 2016), and to examine the spatial requirements for Nodal-mediated patterning (McNamara et

al., 2025). Optogenetic approaches have also provided insights into how Nodal and non-canonical

Wnt signaling influence cell migration in zebrafish embryos (Čapek et al., 2019; Emig et al., 2024).

Finally, optogenetic tools that activate extracellular signal-regulated kinase (ERK) have been

demonstrated in zebrafish (Benman et al., 2022; Kainrath et al., 2017; Patel et al., 2019) and used

to investigate ERK's influence on morphogenetic movements (Patel et al., 2019) and cell size/cycle

regulation in the epidermis (Ramkumar et al., 2025). These compelling early studies showcase the

novel research avenues that optogenetic technology can enable. However, to address gaps in our

understanding of dynamic signal interpretation, optogenetic experiments introducing precise

signaling inputs are ideal, but require in-depth knowledge of tool kinetics, responses to different

light irradiances, and spatial activation properties. Therefore, a robustly characterized toolkit of

optogenetic signaling activators with a unified design under identical conditions is valuable for

use in developmental signaling investigations.

88           Toward these goals, here we present a comprehensive, direct comparison of an optogenetic

toolkit to activate FGF, BMP, and Nodal signaling during zebrafish development. Our toolkit

comprises a novel zebrafish-optimized FGF activator, a single-transcript Nodal activator, and an

established BMP activator (Rogers et al., 2020). Here we show that light-mediated signaling

activation by these tools *in vivo* is wavelength- and pathway-specific and leads to activation of

known pathway target genes. We demonstrate that these tools activate signaling with distinct, rapid

on/off kinetics and that signaling activation strength depends on light irradiance. Finally, we

demonstrate localized control of signaling with our novel FGF activator via spatially restricted

light exposure. Together, our work provides a resource for zebrafish biologists in search of

technology that enables straightforward, dynamic experimental activation of FGF, BMP, or Nodal

signaling with high spatiotemporal control.

**RESULTS**

**Optogenetic receptor kinase activator strategy**

The optogenetic tools characterized here use the general activation strategy of light-

mediated receptor kinase dimerization described in pioneering studies from 2014 (Chang et al.,

2014; Crossman and Janovjak, 2022; Grusch et al., 2014; Kim et al., 2014). FGF, BMP, and Nodal

pathways are all amenable to this strategy because they signal via a similar mechanism: Ligand

binding to extracellular receptor domains initiates interactions between intracellular receptor

kinases, leading to phosphorylation of an effector that translocates to the nucleus and regulates

target gene expression together with other transcription factors (Derynck and Budi, 2019; Heldin

and Moustakas, 2016; Ornitz and Itoh, 2015). To couple signaling activation to light rather than

ligand exposure, intracellular receptor kinase domains can be fused to light-responsive dimerizing

proteins. Light exposure can now activate signaling by inducing receptor kinase interactions.

Because these fusions lack the receptor extracellular ligand binding domain, they are insensitive

to extracellular ligand. The optogenetic tools characterized here are based on constructs developed

by (Sako et al., 2016) and contain the flavin mononucleotide (FMN)-binding, blue light

homodimerizing "light oxygen voltage-sensing" (LOV) domain AUREOCHROME variant from

the algae *Vaucheria frigida* (VfAU1) (Takahashi et al., 2007; Toyooka et al., 2011) fused to

zebrafish receptor kinase domains (Fig. 1). VfAU1 produced the most robust, reversible light-

dependent activation compared to other LOV variants tested in cell culture in a pioneering study

(Grusch et al., 2014). The constructs used here contain a myristoylation motif for membrane

targeting and a C-terminal HA or FLAG tag for immuno-detection. We refer to these constructs as

"bOpto" tools for **b**lue light **Opto**genetic signaling activators.

We first sought to create a zebrafish-optimized bOpto-FGF activator. Activation of FGF

receptor kinases engages downstream signaling mediators, including SOS and MEK, culminating

in ERK phosphorylation (Ornitz and Itoh, 2015). Optogenetic ERK activators have been described

in many systems (Choi et al., 2021; Csanaky et al., 2019; Dine et al., 2018; Grusch et al., 2014;

Kim et al., 2014; Krishnamurthy et al., 2020; Pal et al., 2023; Rauschendorfer et al., 2021; Yadav

et al., 2021). To our knowledge, three previous optogenetic ERK activators have been reported in

zebrafish: 1) mFGFR1-MxCBD, a mouse FGF receptor intracellular domain fused to a cobalamin

binding domain—a homodimer in the dark that dissociates with green light exposure (Kainrath et

al., 2017), 2) BcLOV-SOS$_{cat}$, a fusion of the human RAS activator SOS$_{cat}$ (Toettcher et al., 2013)

with the temperature-sensitive blue light homodimerizing domain BcLOV4 (Benman et al., 2022),

3) and psMEK$^{E203K}$, a constitutively active hamster MEK1 fused to 400/500 nm photoswitchable

Dronpa monomers modified from (Zhou et al., 2017) by the addition of the cancer-associated

activating E203K substitution (Patel et al., 2019; Ramkumar et al., 2025). These ERK activators

feature human-, mouse-, or hamster-derived signaling components that may behave differently in

different species. For example, in the process of creating optimized hamster-derived psMEK$^{E203K}$,

preliminary experiments showed that a non-optimized psMEK tool (Zhou et al., 2017) had strong

activity in *Drosophila* but weak activity in zebrafish (Patel et al., 2019). This could reflect low

sensitivity of zebrafish to ectopic ERK signaling (Patel et al., 2019), or sub-optimal interactions

between the hamster-derived kinase and zebrafish pathway components. In addition, BcLOV-

SOS$_{cat}$ and psMEK$^{E203K}$ act downstream of the receptor kinase (Benman et al., 2022; Patel et al.,

2019; Ramkumar et al., 2025), which is useful for interrogating the roles of a specific pathway

branch but may not provide information about the entire FGF ligand-activated signaling cascade.

Also, for practical reasons, we sought to develop a tool reversibly activated by blue light, rather

than a tool that is active in the dark and inhibited by green light (Kainrath et al., 2017). Therefore,

our objective was to design a bOpto tool that 1) uses a zebrafish-specific receptor kinase to

recapitulate normal pathway interactions, 2) activates all pathway branches regulated by FGF

ligand to mimic endogenous signaling, and 3) is activated by blue light (~450 nm), and reverts to

the inactive state when returned to dark. In zebrafish, the five FGF receptors function largely

redundantly, although phenotypes are uniquely observed when *fgfr1a* is knocked down in

combination with any other receptor, suggesting that this receptor is a key FGF signaling mediator

in the early embryo (Leerberg et al., 2019). We therefore fused everything downstream of the

putative transmembrane domain from zebrafish Fgfr1a (zFgfr1a) to LOV to generate bOpto-FGF

(Fig. 1A,A', Addgene #232639).

Next, we created a single-transcript version of a previously published optogenetic Nodal activator. The original LOV-based bOpto-Nodal activator is composed of two constructs: VfAU1 fused to the serine/threonine kinase domains of the zebrafish type I receptor Acvr1ba and the type II receptor Acvr2ba, respectively (Sako et al., 2016). To minimize potential for dark leakiness (McNamara et al., 2025) and simplify handling, we created a single-transcript construct that codes for both components. In "bOpto-2A-Nodal", a viral PTV-1 2A peptide sequence separates the two coding sequences (Fig. 1C,C', Addgene #232640) (Kemmler et al., 2023; Ryan et al., 1991). Ribosomes that encounter this 2A sequence during translation fail to form a peptide bond, leading to release of the previously translated polypeptide; translation then continues after the 2A site (Donnelly et al., 2001a; Donnelly et al., 2001b). We therefore expect bOpto-2A-Nodal to produce two separate type I-LOV and type II-LOV proteins from a single mRNA. This approach has been successful with other optogenetic systems (Humphreys et al., 2023; Krishnamurthy et al., 2020; Ueda et al., 2022). In addition, we replaced the HA tag on the type II-LOV component with a FLAG tag to facilitate immuno-detection of individual components (Fig. 1C).

The third tool in our kit is a previously established bOpto-BMP activator (Fig. 1B,B') (Rogers et al., 2020; Saul, Rogers et al., 2023). This is a three-component system composed of VfAU1 fused to the cytoplasmic domains of the zebrafish BMP receptors Bmpr1aa, Acvr1l, and Bmpr2a. Previous work demonstrated robust BMP activation in zebrafish with minimal dark leakiness (Rogers et al., 2020).

Together, bOpto-FGF, -BMP, and -2A-Nodal comprise the toolkit of zebrafish-specific receptor kinase-based optogenetic signaling activators that we characterize herein.

**Wavelength dependent optogenetic activators of FGF, BMP, and Nodal signaling**

We first sought to determine whether bOpto tools activate their respective downstream signaling pathways specifically in response to blue light exposure (455 nm). The VfAU1 domain used here should homodimerize only in response to blue light (Mitra et al., 2012; Takahashi et al., 2007; Toyooka et al., 2011). To empirically test the wavelength dependence of bOpto tools and define "safe handling" conditions, we injected zebrafish embryos at the one-cell stage with mRNA encoding *bOpto-FGF*, *bOpto-BMP*, or *bOpto-2A-Nodal*. mRNA amounts were titrated in earlier experiments to optimize dynamic range (not shown). We then either maintained embryos in the dark, exposed them to 455 nm light (50 W/m$^2$, Supp. Fig. 1A), or exposed them to 495+ nm light

(495-2200 nm, 18.51 W/m$^2$, Supp. Fig. 1B) from 2 hours post-fertilization (hpf) to 1 day post-

fertilization (dpf). Ectopic BMP signaling results in ventralization (V1-V4) at 1dpf (Kishimoto et

al., 1997; Nguyen et al., 1998; Schmid et al., 2000), whereas embryos experiencing strong ectopic

FGF or Nodal signaling have developmental defects and often lyse by 1 dpf (Feldman et al., 2002;

Fürthauer et al., 1997; Fürthauer et al., 2004; Rebagliati et al., 1998; Rogers et al., 2017). We used

these phenotypes as a convenient, rapid readout of ectopic signaling activation. At 1 dpf uninjected

controls appeared morphologically normal in all light conditions (Fig. 2A, Supp. Fig. 2),

suggesting minimal phototoxicity. Injected embryos maintained in the dark or exposed to 495+ nm

light were mostly morphologically indistinguishable from uninjected siblings. In contrast, injected

embryos exposed to 455 nm light had phenotypes consistent with strong ectopic activation of the

corresponding pathway. Our phenotype data indicates that exposure to 455 nm light robustly

activates bOpto tools, whereas we did not observe obvious ectopic signaling activation in the dark

or in response to 495+ nm light.

To directly assay activation of each pathway at an earlier developmental stage, we then

used hybridization chain reaction immunofluorescence (HCR-IF) (Schwarzkopf et al., 2021) to

assess phosphorylated effectors of FGF (ppERK), BMP (pSmad1), and Nodal (pSmad2) pathways.

We assessed signaling in *bOpto*-injected and uninjected embryos exposed to dark, 455 nm light,

or 495+ nm light for 30 minutes starting at early gastrulation (50% epiboly – shield) (Fig. 2B-D

and B'-D', Supp. Figs. 3 & 4). To account for non-uniform *bOpto* mRNA levels and distribution,

we include mRNA encoding *GFP* in our *bOpto* injection mixes (Lim et al., 2009). The mixture of

both *bOpto* and *GFP* mRNAs should be dispersed with the same relative distribution; as such,

GFP signal can be used as a proxy for bOpto tool distribution. Phosphorylated effector

fluorescence intensity measured by HCR-IF was normalized pixel-wise to GFP fluorescence

intensity to account for heterogeneity.

We did not notice differences in ppERK, pSmad1, or pSmad2 signal in uninjected embryos

across light conditions (Fig. 2B-D, Supp. Fig. 4), indicating that light exposure does not affect

these signaling pathways in this context. For all tools, there was a significant effect of wavelength

on normalized phosphorylated signaling effector levels ($p < 0.0001$) (Fig. 2B'-D', Supp. Figs. 3 &

4, Supplementary Materials). bOpto-FGF embryos exposed to 455 nm light showed a 2.9-fold

increase in normalized ppERK levels compared to dark ($p < 0.0001$), and a 3.5-fold increase when

compared to 495+ nm ($p < 0.0001$) (Fig. 2B'). In contrast, there were no differences between

bOpto-FGF embryos exposed to 495+ nm light versus dark ($p = 0.4099$). Similarly, bOpto-2A-

Nodal embryos exposed to 455 nm light showed a 4.9-fold increase in normalized pSmad2 levels

compared to dark ($p < 0.0001$), and a 4.9-fold increase when compared to 495+ nm ($p < 0.0001$)

(Fig. 2D'). There were no differences between bOpto-2A-Nodal embryos exposed to 495+ nm

light versus dark ($p = 0.9997$). Finally, bOpto-BMP embryos exposed to 455 nm light similarly

showed a 1.6-fold increase in normalized pSmad1 levels compared to dark ($p < 0.0001$), and a 1.5-

fold increase compared to 495+ nm ($p < 0.0001$). Levels in embryos exposed to 495+ nm were not

different from dark exposed embryos ($p = 0.9492$) (Fig. 2C'). However, bOpto-BMP showed a

less clear increase in signaling in response to blue light compared to -FGF and -2A-Nodal, possibly

due to the presence of the endogenous ventral-to-dorsal pSmad1 signaling gradient (Rogers et al.,

2020; Tucker et al., 2008).

Together, our phenotype and HCR-IF results demonstrate robust bOpto-mediated signaling

activation with 455 nm light exposure and negligible activation when exposed to wavelengths

above 495nm (at 18.51 $W/m^2$) or when maintained in the dark. Based on these results, to safely

handle embryos expressing bOpto tools without inadvertent photoactivation of signaling

pathways, we use a light source with a peak wavelength well above 495 nm (~600 nm, Supp. Fig.

1C).

**bOpto tools activate expression of known pathway target genes**

Signaling activation leads to expression of pathway-specific target genes. To determine

whether bOpto-FGF, -BMP, and -2A-Nodal activate downstream target genes, we used

hybridization chain reaction fluorescence *in situ* hybridization (HCR-FISH) (Choi et al., 2018) to

assess expression of known pathway target genes. We exposed *bOpto*-injected and uninjected

embryos to dark or 455 nm light (50 $W/m^2$) for two hours starting before gastrulation (dome - 30%

epiboly stage), then fixed embryos and performed multiplexed HCR-FISH (Fig. 3). We chose a

two-hour exposure to allow enough time for ample transcript accumulation, and selected three

genes per pathway based on robust evidence implicating them as pathway targets: FGF = *spry4*,

*dusp6*, and *il17rd* (Fig. 3A) (Fürthauer et al., 2002; Fürthauer et al., 2001; Molina et al., 2007);

BMP = *sizzled*, *gata2a*, and *bambia* (Fig. 3B) (Greenfeld et al., 2021; Rogers et al., 2020); Nodal

= *lefty1*, *goosecoid*, and *noto* (Fig. 3C) (Bennett et al., 2007; Bisgrove et al., 1999). In all cases,

gene expression was similar in uninjected dark embryos, uninjected 455 nm light exposed

embryos, and injected dark embryos, suggesting negligible bOpto leakiness in the dark. In contrast,

455 nm light exposure activated gene expression in *bOpto*-expressing embryos. Together, our

results demonstrate robust optogenetic activation of known target genes by bOpto-FGF, -BMP, and

-2A-Nodal in zebrafish embryos, establishing that this suite of optogenetic activators provides

experimental access to major signaling pathways during zebrafish development.

**bOpto tools are pathway-specific**

Receptors can engage in cross-talk, in which pathway components canonically associated

with one signaling pathway are involved in transducing signaling from one or more additional

pathways (Derynck and Budi, 2019; Ramachandran et al., 2018). We therefore designed an

experiment to investigate whether cross-talk is associated with this suite of optogenetic signaling

activators. We injected zebrafish embryos at the one-cell stage with mRNA encoding *bOpto-FGF*,

*-BMP*, or *-2A-Nodal* and reared them in the dark until early gastrulation (50% epiboly – shield).

We then exposed half of the injected embryos, as well as uninjected control siblings, to 455 nm

light (50 W/m$^2$) for 30 minutes. To simultaneously detect phosphorylated signaling effectors from

all three pathways (FGF (ppERK), BMP (pSmad1), and Nodal (pSmad2)), we then performed

triple IF staining by modifying the dual IF protocol from (Soh et al., 2020).

In uninjected controls, signaling differences were not observed in dark compared to blue

light exposed embryos (Fig. 4), consistent with earlier HCR-IF experiments (Fig. 2). We also noted

minimal differences between uninjected embryos and injected embryos maintained in the dark,

again suggesting negligible leaky dark activation. Blue light-exposed bOpto-FGF embryos showed

strong ectopic ppERK activation, but no obvious ectopic pSmad1 or pSmad2 activity in any

condition (Fig. 4A). Similarly, blue light-exposed bOpto-BMP embryos had higher levels of

pSmad1 but no evident ectopic ppERK or pSmad2 activity in any condition (Fig. 4B). Embryos

expressing bOpto-2A-Nodal exposed to blue light had strong ectopic pSmad2 activation and

showed no changes in pSmad1 levels between conditions (Fig. 4C). We occasionally observed

embryos in bOpto-2A-Nodal experiments with possible ectopic ppERK activity (Supp. Fig. 5),

though this appears to be an IF or imaging artifact. Overall, bOpto-FGF, -BMP, and -2A-Nodal

generate robust pathway-specific optogenetic activation at the level of signaling in gastrulation

stage zebrafish.

**bOpto tools have rapid on/off kinetics**

Speed and reversibility are two benefits of optogenetic strategies (Rogers and Müller, 2020). These features are especially important in studies investigating how cells interpret signaling (Farahani et al., 2021) to make fate decisions during early embryogenesis, where key patterning events can occur in a matter of hours or even minutes (Huang et al., 2017; Sako et al., 2016). To determine the on/off kinetics of bOpto-BMP, -FGF, and -2A-Nodal, we injected mRNA encoding each respective tool together with *GFP* mRNA into zebrafish embryos at the one cell stage. At early gastrulation stage (50% epiboly – shield) we exposed embryos to 455 nm light (50 W/m$^2$) for 30 minutes, then removed light for a defined tool-dependent dark recovery period. We fixed embryos at multiple time points throughout the exposure and recovery phases, together with time-matched injected controls that were maintained in the dark the entire time. We then used HCR-IF to quantify phosphorylated signaling effector levels corresponding to the stimulated pathway (Fig. 5). GFP signal was used to normalize *bOpto* mRNA amount and distribution. In addition, normalized effector signal was background subtracted against dark controls (Supp. Figs. 6 & 7). We observed clear signaling activation within the 30 min light exposure phase and a return to baseline after light removal for all tools (Fig. 5A-C, A'-C').

First, to discern differences in "on" kinetics of the tools, we analyzed the light exposure phase of the time course (Fig. 5A"-C"). Three-parameter logistic regressions were fit to the data. From the model, three values were calculated approximating the time to reach a percentage of the curve's upper asymptote: tON$_{20}$ predicting initial activation, tON$_{50}$ predicting half-maximum, and tON$_{90}$ predicting saturation (Supplementary Materials). bOpto-FGF (curve fit R$^2$ = 0.503) had the fastest on kinetics, reaching saturation within 2 minutes (Fig. 5A"). Because our assayed time points were determined *a priori*, there was insufficient coverage between 0 and 2 min for confident predictions of tON$_{20}$, tON$_{50}$, and tON$_{90}$. However, it is clear that initiation and saturation occur between 0 and 2 min for bOpto-FGF. bOpto-BMP (curve fit R$^2$ = 0.534) had a tON$_{20}$ of 1.78 min (CI: -4.97-8.53 min), a tON$_{50}$ of 6.36 min (CI: 1.35-11.37 min), and tON$_{90}$ of 13.62 min (CI: 0.75-26.49 min) (Fig. 5B"). Of the three constructs, bOpto-2A-Nodal (curve fit R$^2$ = 0.966) had the slowest on kinetics: it had a tON$_{20}$ of 16.12 min (CI: 12.25-19.99 min), a tON$_{50}$ of 19.78 min (CI: 16.13-23.43 min), and tON$_{90}$ of 25.59 min (CI: 14.04-37.14 min) (Fig. 5C"). Comparing initiation times (tON$_{20}$), and assuming that this occurs for bOpto-FGF between 0-2 minutes, bOpto-2A-Nodal is ~14 min slower than bOpto-FGF; bOpto-BMP and bOpto-FGF both likely initiate

activation before 2 minutes of light exposure. Comparing the tools' time to saturation ($tON_{90}$), again assuming that this occurs for bOpto-FGF within the first 2 minutes, bOpto-FGF is ~12 minutes faster than bOpto-BMP and ~24 minutes faster than bOpto-2A-Nodal. It also appears that the tools that clearly reach saturation prior to end of the 30 minute light exposure (bOpto-FGF and -BMP) maintain peak signaling levels until light removal.

Next, to discern differences in "off" kinetics of the tools, we analyzed the dark recovery phase of the time courses (Fig. 5A'''-C'''). In this phase, embryos that had been exposed to the 30 minute, 455 nm light pulse were incubated in the dark for an additional 55 (bOpto-FGF) or 90 (bOpto-BMP and -2A-Nodal) minutes. The same curve fitting approach that was used to determine the on kinetics was also used here to assess the off kinetics with a conceptual alteration: We calculated the time to a defined percentage *decrease* from the upper asymptote to reflect the progressive return to baseline (Supplementary Materials). For each tool we calculated $tOFF_{20}$ predicting time to initial deactivation, $tOFF_{50}$ predicting time to half-maximum, and $tOFF_{90}$ predicting approximate time to return to baseline. The off kinetics of bOpto-FGF (curve fit $R^2 =$ 0.877) were the most rapid of the tested tools, with a $tOFF_{20}$ of 1.76 min (CI: -9.05-12.56 min), $tOFF_{50}$ of 5.30 min (CI: 2.72-7.88 min), and a $tOFF_{90}$ of 10.92 min (CI: -37.37-59.20 min). bOpto-BMP (curve fit $R^2 = 0.667$) was determined to have a $tOFF_{20}$ of 20.81 min (CI: 8.47-33.15 min), a $tOFF_{50}$ of 27.11 min (CI: 19.56-34.67 min), and a $tOFF_{90}$ of 37.10 min (CI: 20.99-53.21 min). Finally, bOpto-2A-Nodal (curve fit $R^2 = 0.626$) had the slowest off kinetics, with a $tOFF_{20}$ of 10.63 min (CI: -4.34-25.61 min), a $tOFF_{50}$ of 32.67 min (CI: 22.22-43.12 min), and a $tOFF_{90}$ of 67.60 min (CI: 33.56-101.63 min). Comparing the tools' time to initial deactivation ($tOFF_{20}$), bOpto-FGF began to decrease first, 9 min before bOpto-2A-Nodal and 19 min before bOpto-BMP. In contrast to bOpto-FGF and -2A-Nodal-induced signaling, which started to decrease relatively soon after light removal, bOpto-BMP-induced signaling was sustained into the dark period and then rapidly progressed towards baseline. Comparing time to return to baseline ($tOFF_{90}$), bOpto-2A-Nodal had the slowest return time, ~31 minutes slower than bOpto-BMP and ~57 minutes slower than bOpto-FGF.

Each tool's time to reach saturation in the light exposure phase ($tON_{90}$) was faster than its time to return to baseline after the light had been removed ($tOFF_{90}$). Additionally, the relative kinetics between the tools (e.g., bOpto-FGF fastest, bOpto-2A-Nodal slowest) was overall

consistent for both on and off kinetics. In summary, these data demonstrate that all tools in the

bOpto toolkit activate signaling rapidly and reversibly.

**Optogenetic signaling activation is dependent on irradiance**

Another key advantage of optogenetic tools is the ability to adjust activation strength by

modulating light irradiance (Rogers and Müller, 2020). This ability is particularly useful because

different signaling levels can have distinct biological consequences (Johnson and Toettcher, 2019).

To characterize the irradiance dependence of bOpto tools, we injected embryos with mRNA

encoding *bOpto* constructs and *GFP* at the one-cell stage, then at early gastrulation (50% epiboly

– shield) exposed embryos to 455 nm light (Supp. Fig. 1A) at eight different irradiances ranging

from 0.02 to 50 W/m$^2$, changing roughly 3-fold between steps (Fig. 6A). Because bOpto-BMP and

-2A-Nodal have slower times to saturation than bOpto-FGF (Fig. 5), we chose exposure durations

of 5 minutes for bOpto-FGF and 25 minutes for bOpto-BMP and -2A-Nodal. We fixed embryos

immediately after exposure and used HCR-IF to quantify the relevant phosphorylated signaling

effector (Fig. 6), normalizing for mRNA level and distribution using GFP signal and background

subtracting based on dark controls (Supp. Figs. 8 & 9).

For all three constructs, levels of phosphorylated signaling effectors increased with

increasing light irradiance (Fig. 6). To characterize each tool's irradiance dependence, three-

parameter logistic regressions were fit to the data. From the model, three values were calculated

approximating the irradiance at which a percentage of the curve's upper asymptote is reached: $I_{20}$

predicting initial response, $I_{50}$ predicting half-maximum, and $I_{90}$ predicting saturation

(Supplementary Materials). bOpto-FGF (curve fit $R^2 = 0.748$) was most sensitive to irradiance

with an $I_{20}$ of 0.05 W/m$^2$ (CI: 0-0.11 W/m$^2$), an $I_{50}$ of 0.09 W/m$^2$ (CI: 0.03-0.15 W/m$^2$), and an $I_{90}$

of 0.15 W/m$^2$ (CI: 0.05-0.25 W/m$^2$). bOpto-BMP (curve fit $R^2 = 0.678$) had an $I_{20}$ of 0.16 W/m$^2$

(CI: -0.03-0.35 W/m$^2$), an $I_{50}$ of 0.35 W/m$^2$ (CI: 0.17-0.53 W/m$^2$), and an $I_{90}$ of 0.67 W/m$^2$ (CI:

0.26-1.08 W/m$^2$). bOpto-2A-Nodal (curve fit $R^2 = 0.578$) was the least sensitive to irradiance with

an $I_{20}$ of 3.90 W/m$^2$ (CI: -0.27-8.06 W/m$^2$), an $I_{50}$ of 11.00 W/m$^2$ (CI: 5.54-16.45 W/m$^2$), and an $I_{90}$

of 22.25 W/m$^2$ (CI: 6.12-38.38 W/m$^2$). We speculate that the bOpto-2A-Nodal curve plateaus over

the range of tested irradiances; however, additional coverage between 10-100 W/m$^2$ is needed to

make a definite conclusion.

To directly compare the light sensitivity of these tools, we first need to account for the fact that exposure durations differed by tool. We therefore converted the $I_{20}$ and $I_{90}$ to dosages ($J/m^2$) by multiplying them by the duration of exposure (bOpto-FGF = 5 min; -BMP and -2A-Nodal = 25 min) and calculating fold changes between tools. Based on their dosage-adjusted $I_{20}$ values, bOpto-FGF was activated at a 14-fold lower light dosage than -BMP and a 356-fold lower light dosage than -2A-Nodal. Similarly, based on dosage-adjusted $I_{90}$ values, bOpto-FGF saturates at a 36-fold lower light dosage than -BMP and a 1211-fold lower light dosage than -2A-Nodal. Together, our results demonstrate tunable, light irradiance-dependent activation of FGF, BMP, and Nodal signaling with this bOpto toolkit in zebrafish embryos.

**Spatially localized optogenetic signaling activation**

A valuable opportunity enabled by optogenetic tools is the ability to manipulate biological processes with spatial precision (Rogers and Müller, 2020). Spatially resolved Nodal activation was recently achieved in zebrafish using a CRY2-based optogenetic activator (McNamara et al., 2025) and, at the level of gene expression, with the two-transcript LOV-based Nodal activator (Sako et al., 2016). Localized signaling activation has also been demonstrated for bOpto-BMP in zebrafish (Rogers et al., 2020). To our knowledge, FGF remains the only pathway activated by our toolkit without evidence for local control via receptor kinase-based optogenetics in zebrafish.

We therefore co-injected *bOpto-FGF* and *nls-Kaede* mRNA at the one-cell stage. Kaede is a green-to-red photoconvertible fluorescent protein that photoconverts at 405 nm (Ando et al., 2002). *nls-Kaede + bOpto-FGF* expressing embryos were reared in the dark, then at early gastrulation stage (shield and older) were irradiated with 445 nm light for 5 minutes using a laser scanning confocal microscope within a 75 x 600 μm region of interest. During the first 3 minutes of exposure, embryos simultaneously received 405 nm light in the same region of interest to photoconvert Kaede-expressing cells. HCR-IF was then performed to detect the FGF signaling effector ppERK. Ectopic ppERK signal overlapped the photoconverted Kaede signal, demonstrating spatially localized FGF activation (Fig. 7, Supp. Fig. 10). In some embryos we noted ERK phosphorylation in a dumbbell-like pattern around the edges of the illumination region. We speculate that, due the sensitivity of bOpto-FGF, this may be caused by out-of-plane light from our choice of illumination method—a confocal laser scanning microscope—that projects a cone-like light volume onto the sample. Dumbbell-like activation was not observed in other illumination

paradigms that project non-angular pencil beam-like light (McNamara et al., 2025), and as such

might be reduced with different hardware. In summary, we demonstrate optogenetically activated

localized ERK phosphorylation in gastrulation stage zebrafish embryos using spatially restricted

light exposure.

**DISCUSSION**

**Rapid, spatially resolved optogenetic signaling activation in zebrafish**

Here we introduce a zebrafish-optimized optogenetic toolkit and provide the systematic in-

depth characterization necessary to use this suite of tools in a variety of experiments. The toolkit

is comprised of a novel FGF activator, a modified single-transcript Nodal activator, and a

previously established BMP activator. This head-to-head *in vivo* characterization enables direct

performance comparisons between the optogenetic signaling activators. We performed these

characterizations in a relevant developmental context—the gastrulating zebrafish—where all three

pathways are synergistically acting to pattern the embryonic axes. We show that bOpto tools enable

wavelength- and pathway-specific activation of FGF, BMP, and Nodal signaling (Figs. 2 & 4) and

target genes (Fig. 3) with the simple, easily controlled input of 455 nm light. Signaling can be

activated with rapid on/off kinetics (Fig. 5), ectopic signaling strength can be tuned by modulating

light irradiance (Fig. 6), and activation can be localized with spatially restricted illumination (Fig.

7). This optogenetic toolkit can therefore be a highly useful resource for investigations using

zebrafish as a model system.

**FGF, BMP, and Nodal pathway cross-talk**

We did not observe obvious, direct cross-activation of non-target pathways with bOpto

tools using triple ppERK, pSmad1, pSmad2 IF (Fig. 4). However, we only assessed FGF, BMP,

and Nodal signaling, and it is possible that other pathways could be inadvertently targeted by

bOpto tools. Here we assessed direct effects by focusing on a relatively short 30 minute light

exposure, during which signaling was ectopically activated for 20-30 minutes (Fig. 5). However,

with sufficiently long signaling durations, we would expect to observe inter-pathway effects

because FGF, BMP, and Nodal are known to engage in many downstream interactions. For

example, Nodal activates expression of the BMP inhibitor Chordin (Koshida et al., 2002; Kudoh

et al., 2004) and promotes transcription of FGF ligand (Bennett et al., 2007; Mathieu et al., 2004;

van Boxtel et al., 2018). FGF inhibits BMP transcription (Fürthauer et al., 1997) and can block

nuclear translocation of pSmad1 (Kretzschmar et al., 1997).

We occasionally observed possible ectopic activation of FGF signaling in bOpto-2A-Nodal

experiments (Supp. Fig. 5), which appears to be an IF or imaging artifact. However, other

possibilities include leaky Nodal activation leading to *fgf* transcription. Another confounding

factor is the observation that ERK phosphorylation can be caused by mechanical damage even at

early stages (Supp. Fig. 11). FGF/ERK signaling is known to be involved in injury and

regeneration responses in zebrafish adults (De Simone et al., 2021; Owlarn et al., 2017; Poss et al.,

2000; Whitehead et al., 2005), *Xenopus* embryos (Christen and Slack, 1999; LaBonne and

Whitman, 1997; Li et al., 2013), and *Drosophila* embryos (Belacortu and Paricio, 2011). We do

not believe this is a likely explanation, but we present this as an important consideration in the

interpretation of ppERK IF data in early zebrafish embryos.

In summary, although bOpto tools are pathway-specific, optogenetic experiments

involving long-term signaling activation may affect non-targeted pathways due to downstream

pathway interactions, though this caveat is not exclusive to optogenetic strategies.

**bOpto tools have distinct on/off kinetics and irradiance sensitivities**

Our measurements revealed the temporal activation and deactivation properties of bOpto

tools in zebrafish (Fig. 5, Supp. Figs. 6 & 7). All constructs in this toolkit have relatively fast on

kinetics, in the order of minutes. Signaling activation is reversible, decreasing to baseline after

light is removed. In addition, each tool takes longer to deactivate than to activate. Despite these

similarities, the tools also have distinct kinetic properties. bOpto-FGF is the fastest signaling

activator and bOpto-2A-Nodal is the slowest. bOpto-BMP on/off kinetics fall in between the other

two tools, although we observed a unique reversion feature: Signaling was briefly sustained after

light removal before rapidly returning to baseline. In contrast, there was a shorter lag between light

removal and the decrease in ectopic signaling for bOpto-FGF and -2A-Nodal.

Overall, the signaling kinetics we measured for bOpto-BMP and bOpto-2A-Nodal in

zebrafish are consistent with prior literature. As our zebrafish-optimized bOpto-FGF tool is novel,

no literature comparisons are available. The kinetics we measured for bOpto-BMP are roughly

consistent with those measured in the establishing study (Rogers et al., 2020). bOpto-2A-Nodal

signaling kinetics appeared to be similar to those measured for the two-component bOpto-Nodal parent constructs (McNamara et al., 2025). bOpto-2A-Nodal on kinetics also appear roughly similar to those measured for a recent CRY2-based Nodal activator, though the CRY2-based construct may have faster off kinetics (Emig et al., 2024; McNamara et al., 2025). However, differences in exposure conditions (e.g., wavelength, irradiance) complicate direct comparisons.

We observed similar trends with responses to irradiance (Fig. 6, Supp. Figs. 8 & 9). Signaling activation by all tools increased as irradiance increased. bOpto-FGF activity was initiated with very low light dosages, whereas activation from bOpto-2A-Nodal required much higher light dosages, and bOpto-BMP was responsive to dosages between those that activated the FGF and Nodal tools. Previous literature also reported light dosage-dependence for bOpto-BMP (Rogers et al., 2020), the two-component bOpto-Nodal parent constructs, and the CRY2-based Nodal activator (McNamara et al., 2025). The striking differences in light dosage sensitivity we demonstrate here illustrate the importance of controlling and reporting light irradiances when using the bOpto toolkit.

Although further molecular characterization would be required to explain the differences in on/off kinetics and irradiance sensitivity between the bOpto tools, we speculate that several factors may contribute. First, the phosphorylation kinetics of the receptor kinase domains may differ, as well as the kinetics of phosphatases targeting pathway effectors. The kinase domain in the FGF activator is a receptor tyrosine kinase, whereas the BMP and Nodal activators are serine/threonine kinase-based. Receptor tyrosine kinase activity can be rapid (Farahani et al., 2023), and zebrafish embryos treated with a MEK inhibitor show strong reduction in ERK phosphorylation within just 10-20 minutes (Wilcockson et al., 2023). Another possible factor is the number of components in each bOpto activator: bOpto-FGF is a one-component system, whereas bOpto-2A-Nodal is composed of a type I and type II construct, and bOpto-BMP comprises two type I and one type II constructs (Fig. 1). This may influence the fraction of productive interactions formed upon light exposure. Finally, in the case of bOpto-2A-Nodal, the use of the 2A peptide might contribute to its observed properties. bOpto-2A-Nodal consists of two Myr-tagged constructs (Sako et al., 2016) separated by a 2A peptide that in principle produces two separate membrane-localized proteins (Fig. 1C). However, the localization of these components has not been directly tested. A previous report showed that a Myr-tagged construct downstream of a 2A peptide failed to localize to the membrane in HEK293T cells (Hadpech et al., 2018), suggesting

that the type II-LOV component of bOpto-2A-Nodal may be cytoplasmic. If the type II-LOV

construct is in fact located in the cytoplasm, activation kinetics could be delayed due to the larger

search space of the cytoplasm vs. cell membrane. Conversely, other optogenetic receptor kinase-

based activators in which one component is cytoplasmic have been demonstrated to robustly

activate signaling with an improved signal to noise ratio (Krishnamurthy et al., 2020; McNamara

et al., 2025). We observed minimal dark leakiness with bOpto-2A-Nodal. If the type II-LOV

component of bOpto-2A-Nodal is cytoplasmic, this might explain the apparent improvement in

dynamic range compared to previous observations from parent constructs (McNamara et al., 2025).

To determine the subcellular localization of each component of bOpto-2A-Nodal, future studies

could use immuno-detection of the HA tag on the type I-LOV component and the FLAG tag on

the type II-LOV component.

Overall, our measurements demonstrate that signaling activation with the bOpto toolkit is

rapid, reversible, and tunable in gastrulation-stage zebrafish. This characterization therefore

provides a range of useful parameters that can be experimentally manipulated to generate signaling

pulses with different durations and amplitudes.

**Strategies to optimize use of bOpto signaling activators**

Based on our work, we suggest four key practical recommendations for zebrafish

researchers planning to incorporate bOpto signaling activators into their studies. First, it is crucial

to inject the appropriate amount of functional mRNA to avoid weak or leaky activation. Controls

should be included in every experiment as a quality assessment (Saul, Rogers et al., 2023). In

addition, mRNA injections can lead to heterogeneous distribution of bOpto tools. Two strategies

to mitigate these effects are 1) addition of GFP mRNA to injection mixes (as done here), or 2)

fusing fluorescent proteins to bOpto tools. Fluorescent read outs can be used to identify embryos

injected with appropriate mRNA amounts and to normalize for differences in distribution or levels.

In experiments involving signaling quantification, we found that GFP normalization improved the

signal to noise ratio and aided interpretability.

Second, we show that signaling was not detectibly activated by light over 495 nm (Fig. 2,

Supp. Figs. 2, 3, & 4). Our 495+ nm light source (18.51 W/m$^2$) emitted light at a lower irradiance

than our 455 nm source (50 W/m$^2$) (Supp. Fig. 1). However, >20 hours of 495+ nm light exposure

at this relatively lower irradiance was insufficient to activate signaling based on phenotypes (Fig.

2A, Supp. Fig. 2). In contrast, as little as 0.15 W/m$^2$ 455 nm light for 5 minutes robustly activated

ectopic FGF signaling (Fig. 6, Supp. Fig. 9). We therefore recommend using light sources well

above 495 nm to safely handle embryos. This can be achieved with bespoke lightbulbs, or by

covering white light sources (e.g., microscope stages) with red filters.

Third, our characterization of on/off kinetics and irradiance responses indicate that bOpto

tools have distinct properties that should be considered when determining light exposure

conditions in experimental designs. For example, bOpto-FGF is several hundred-fold more

sensitive to light dosage and is also activated 10-20 minutes faster than bOpto-2A-Nodal (Figs. 5

& 6). Therefore, light exposure conditions that saturate signaling from bOpto-FGF may not even

initiate signaling from bOpto-2A-Nodal.

Fourth, a potential confounding factor in optogenetic experiments is light-induced

dysregulation of development. Circadian oscillations have been characterized in many organisms

including zebrafish (Dekens et al., 2003; Froland Steindal and Whitmore, 2019; Krylov et al.,

2021; Laranjeiro and Whitmore, 2014; Tamai et al., 2004; Whitmore et al., 2000; Ziv and Gothilf,

2006). Of the four circadian genes that we assayed in uninjected gastrulating zebrafish, three

showed significant light-mediated increases in gene expression by HCR-FISH (*cry2*, *per2*, *tefa*;

Supp. Fig. 12). It is unclear whether these changes in circadian gene expression affect

developmental signaling. However, we did not detect obvious impacts of 455 nm light exposure

on FGF, BMP, or Nodal signaling (Figs. 2B-D, B'-D', Fig. 4, Supp. Fig. 4), gene expression (Fig.

3), or morphology at 1 dpf (Fig. 2A, Supp. Fig. 2). We therefore speculate that the effects of light

exposure on circadian gene expression did not confound the interpretation of our results. However,

additional studies are needed to probe the effects of circadian gene responses in the context of

optogenetic experiments. We suggest that blue (e.g., 455 nm) light exposed uninjected controls

should be included in optogenetic experiments as a precaution.

**Prospects for optogenetic technology in developmental biology**

We foresee at least three major uses of bOpto tools to address outstanding questions. First,

to examine how cells decode these signaling features in a variety of contexts, our characterization

of bOpto on/off kinetics and light irradiance dependence can help researchers create bespoke

signaling inputs with desired amplitudes and durations. Second, it may be possible to multiplex

these tools and jointly manipulate multiple pathways—for example, to generate simultaneous

pulses of BMP and FGF signaling. This could facilitate studies examining how cells interpret combinatorial signaling, which appears to be a key patterning strategy (Briscoe and Small, 2015; Kicheva and Briscoe, 2023). Third, it will be valuable to use bOpto tools to impose tailored experimental signaling inputs in pathway loss-of-function mutants, as in (McNamara et al., 2025; Vopalensky et al., 2018). This could help define which features of signaling gradients are important for their patterning functions—e.g., slope, shape, dynamics—among other applications.

Finally, several technical advances would make optogenetic strategies even more accessible. A current challenge with bOpto tools is that they are introduced by mRNA injection, which is transient and can lead to inconsistent distribution. Transgenic lines expressing bOpto constructs would obviate these challenges. Converting the two-transcript bOpto-Nodal tool (Sako et al., 2016) into the single-transcript construct used here facilitates the generation of transgenics since only a single genomic insertion is necessary. In addition, we demonstrated local signaling activation via spatially restricted light exposure in gastrulation-stage zebrafish on the order of 10-100 μm. The resolution of optogenetic control is in principle limited only by the combination of focusing optics used in illumination hardware and the underlying biology of the optogenetic tool. With further experimental refinements it may be possible to activate with even finer spatial resolution. For example, cellular precision has been achieved in intact *Drosophila* embryos (Guglielmi et al., 2015; Izquierdo et al., 2018) and cell culture (Benedetti et al., 2018; Donahue et al., 2021). Finally, tools to manipulate additional pathways, inhibit signaling, and orthogonal tools activated by different wavelengths would unlock further experimental possibilities and provide more opportunities to explore unprecedented research avenues.

In summary, the suite of optogenetic FGF, BMP, and Nodal activators characterized here represents a powerful experimental platform for developmental biologists and beyond. This toolkit provides a path for exploring key developmental biology questions such as how signaling dynamics and spatial gradients regulate embryogenesis. We anticipate that this work will motivate a fruitful expansion of optogenetic technology in developmental biology research.

**Acknowledgements:** We thank the NIH zebrafish facility staff for maintaining our fish stocks and Matthew T. Monaghan for helping to test immunofluorescence staining strategies. We also thank Jeffrey A. Farrell and his lab, Pedro P. Rocha, Sarah E. Sheppard, Leah F. Rosin, Harry A. Burgess, Patrick Müller, and the NICHD Aquatics Affinity Group for helpful discussions.

**Competing interests:** No competing interests declared.

**Funding:** This work was supported by NIH Intramural funding ZIAHD009002-01 to KWR.

**MATERIALS & METHODS**

**Zebrafish husbandry and mRNA injections**

Zebrafish husbandry and research protocols were approved by the NICHD Animal Care and Use Committee in accordance with the Guide for the Care and Use of Laboratory Animals of the National Institutes of Health. The AB zebrafish wild type strain was used in all experiments. Embryos were incubated at 28 °C in embryo medium (reverse osmosis water, Instant Ocean Sea Salt (0.25 g/L), and $NaHCO_3$ (~0.06 g/L, depending on initial pH)). For injections, mRNA was synthesized from NotI-linearized pCS2+ plasmids (see Construct Sequences, Supplementary Materials) using SP6 mMessage mMachine kits (Invitrogen AM1340). 1 nl of injection mix containing mRNA and phenol red tracer was injected through the chorion at the one-cell stage. For experiments in Figs. 2, 5, and 6, and Supp. Figs. 6, 7, 8, and 9, 20 pg *GFP* mRNA (modified from (Lim et al., 2009)) was included in the same injection mix as *bOpto* mRNA. For all experiments, embryos were incubated at 28 °C immediately after injection. Between 1.5 - 2 hpf, healthy, fertilized embryos were transferred to 6-well dishes used in experiments, wrapped in aluminum foil to prevent light exposure (or immediately exposed to light for phenotyping at 1 dpf, see below and Fig. 2A and Supp. Fig. 2), and returned to 28 °C (Saul, Rogers et al., 2023). At 1 dpf, embryos were scored on Leica M80 stereoscopes. Representative phenotype images were acquired on Leica M80 stereoscopes with Flexacam C1 or C5 camera attachments (Fig. 2A and Supp. Fig. 2). The mRNA amounts injected for each optogenetic tool are listed below:

bOpto-FGF (Addgene #232639): 3.5 pg

bOpto-BMP (combined in injection mix)

bOpto-Bmpr1aa (Addgene # 207614): 23.4 pg

bOpto-Acvr1l (Addgene # 207615): 23.4 pg

bOpto-Bmpr2a (Addgene # 207616): 40.2 pg

bOpto-2A-Nodal (Addgene #232640): 10 pg

We have found that mRNA amounts are crucial: Inject too little, and optogenetic activation is negligible, inject too much, and leaky ectopic dark activation will occur (Saul, Rogers et al., 2023). In addition, we have anecdotally observed a decay in mRNA aliquot efficacy over time. We therefore included phenotype controls in all experiments. From each experiment, a subset of embryos was exposed to dark or 455 nm light starting at ~2 hpf and phenotypes were scored at 1 dpf. We only proceeded with experiments in which these controls demonstrated that the appropriate amount of functional mRNA had been delivered. The following criteria were implemented for inclusion:

*bOpto-FGF and -2A-Nodal experiments*
Proceed if at least 60% of injected, 455 nm-exposed embryos are severely deformed or lysed. In addition to the criteria above, do not proceed if the dark injected condition exhibits more than 20% of the following phenotypes: mild-severe deformity, lysed.

*bOpto-BMP experiments*
Proceed if, in the injected, 455 nm-exposed condition, 1) there are V2 or higher embryos present, and 2) at least 60% of embryos show V1-V4 phenotypes (Kishimoto et al., 1997; Nguyen et al., 1998; Schmid et al., 2000). In addition to the criteria above, do not proceed if the dark injected condition exhibits more than 20% of the following phenotypes: V1-V4, mild-severe deformity, lysed.

In addition, rare experiments in which uninjected embryos in both dark and light conditions had high numbers of defects (over ~30%) were excluded.

**Safe embryo handling**

To protect embryos from light in "dark" conditions, 6-well dishes containing embryos were wrapped in aluminum foil. To handle light-sensitive embryos without incurring inadvertent photoactivation, in a windowless room with the door closed we used a ~600 nm red light source (Supp. Fig. 1C) (A19 9W Equivalent 60W, E26 Red LED Colored Light Bulb, UNILAMP, Cat. No. B0C7YZ4KSY). In experiments requiring manipulation or visualization of light-sensitive embryos on a dissecting scope, inadvertent photoconversion was prevented by covering the scope base/white light source with red gel filter paper (#E106, Rosco, Cat. No. 110084014805-E106). In experiments requiring fixation, embryos were fixed either under these red light conditions, or, for some light exposed embryos, in the LED incubator in which they were exposed to light.

**Whole embryo light exposure**

Embryos were exposed to 455 nm light using the "LED incubators" described previously (Saul, Rogers et al., 2023). To create a 495+ nm light source (18.51 W/m$^2$, Fig. 2, Supp. Fig. 2 & 12) a broadband (400 - 2200 nm) Quartz-Tungsten Halogen white light source (Thor Labs, Cat. No. QTH10/M) was used as the base emitter. A 2" x 2" square glass 495 nm long-pass filter (Thor Labs, Cat. No: FGL495S) was mounted to the source via a custom 3D printed adapter (design modified from: Thor Labs, Cat. No. SM2FH) to restrict emission wavelengths to 495+ nm. Emitted spectra from each light source was measured over the 200-1000 nm range using a compact spectrometer (Thor Labs, Cat. No. CCS200) with a cosine corrector (Thor Labs, Cat. No. CCSB1)

Desired irradiance was confirmed prior to all experiments. Power at the sample plane across the 350-1100 nm range was measured with a digital optical power meter (Thor Labs, Cat. No. PM100D) coupled to a microscope slide sensor head (Thor Labs, Cat. No. S170C). Measured power was converted to irradiance by dividing the power by the sensor head's active detector area.

For the light irradiance experiments (Fig. 6), neutral density filters (Norman Pack of 3 Neutral Density (ND) 810551 Filters (5"), B & H Foto & Electronics Corp., Cat. No. NONDS5) were taped over the light output of the LED incubators (Saul, Rogers et al., 2023). Combinations of ND filters and adjustments to the LED power settings were tuned to achieve the desired irradiances.

**Spatial activation experiments**

For spatial activation experiments (Fig. 7, Supp. Fig 10), embryos were dechorionated using pronase at the 1 cell stage before being injected with *bOpto-FGF* (3.5 pg) + *nlsKaede* (25 pg) mRNA. Post-dechorionation and injection, embryos were incubated at 28 °C on agarose-coated petri dishes in the dark. Upon reaching 50% epiboly, embryos were carefully moved under a safe handling light source to a fresh agarose-coated dish which had wells cast into the agarose to maintain the embryos at a stable orientation (animal pole up) during light exposure (Zaucker et al., 2021).

To restrict light to a small (75 x 600 µm) area within the embryo, an upright laser scanning confocal microscope (Nikon AX-FN AXR-MP) was used as the illumination source. The light path consisted of the emission from a combination of laser lines passing through a BS20/80 transmission/reflectance mirror and a Nikon CF145 W Apochromat 20x/1.0 Objective (Mat. No. MRD77226) before reaching the sample plane. A 594 nm laser was used to position the sample in the center of the field of view. A consistent focal plane for illumination was identified by finding the apex of the animal pole and then moving the z-position 125 µm down towards the center of the embryo. A 75 x 600 µm region of interest (ROI) was defined in the center of the 815 x 815 µm field-of-view. Stimulation to the ROI was controlled using the AX and ND Stimulation modules in Nikon Elements and was two consecutive blocks totaling a 5-minute exposure period: 1) 405 nm + 445 nm light scanning the ROI for 3 minutes and 2) 445 nm light only for 2 minutes. Restricting 405 nm exposure to the first 3-minute block allowed sufficient Kaede photoconversion while minimizing any potential phototoxicity. Laser power was set to 25% for 405 nm laser and 15% for 445 nm laser. Power measured at the sample plane was 5.6 µW for block 1 and 2.1 µW for block 2. It took 1.02s to irradiate the 45,000 µm² ROI. Consequently, effective irradiance at the sample plane over the stimulation period was 126.93 W/m² in block 1 and 47.60 W/m² in block 2. The irradiance and dosage due to 445 nm only over the whole exposure period was 47.60 W/m² and 14280.00 J/m², respectively.

One embryo at a time was subjected to the exposure paradigm above. After exposure, embryos were immediately fixed for downstream processing. Time-matched unexposed and uninjected embryos were also fixed as controls.

**Triple pSmad1 + pSmad2 + ppERK immunofluorescence staining protocol**

To simultaneously detect phosphorylated effectors of FGF (ppERK), BMP (pSmad1), and
Nodal (pSmad2) signaling, we developed a triple immunofluorescence (IF) staining protocol based
on the pSmad1 + pSmad2 double IF protocol from (Soh et al., 2020). Note that primary antibodies
for pSmad1 and pSmad2 are both rabbit and detected with anti-rabbit secondaries; however, lower
pSmad2 epitope abundance and/or antibody affinity combined with amplification of the pSmad2
signal are thought to prevent significant crosstalk.

Embryos were fixed overnight at 4 °C in 4% formaldehyde in 1x PBS (phosphate buffered
saline). After washing 3x with 1x PBST (1x PBS + 0.1% Tween 20), embryos were manually
dechorionated using forceps. Embryos were washed 3x in MeOH and stored at -20 °C overnight
or longer. Embryos were then washed 4x with PBST, incubated at room temperature (RT) for 20
min in ice cold acetone, washed again with 4x with PBST, then blocked in 10% fetal bovine serum
(FBS) in 1x PBST at RT for ≥1 hour. Embryos were then incubated at 4 °C overnight in 1:5000
rabbit anti-pSmad2/3 primary antibody (Cell Signaling Technology #8828S) in 10% FBS. The
following day embryos were washed 5x in 1x PBST, re-blocked in 10% FBS for ≥1 hour at RT,
then incubated at 4 °C overnight in 1:500 goat anti-rabbit HRP secondary antibody (Jackson
ImmunoResearch Laboratories #111-035-003) in 10% FBS. Next, embryos were washed ≥5x with
1x PBST, incubated in Tyramide Signal Amplification (TSA) buffer (Akoya Biosciences
NEL744001KT) for ≥15 min at RT, and incubated in 1:75 fluorescein:amplification buffer dilution
for 45 min at RT. Embryos were then washed 4x with 1x PBST, and peroxidase was inactivated by
incubation with 1% $H_2O_2$ at RT for 25 min. After ≥5x washes with 1x PBST, embryos were
incubated in 10% FBS for ≥1 hr at RT and then in 1:5000 mouse anti-ppERK1/2 primary antibody
(Sigma-Aldrich #M8159) in 10% FBS at 4 °C overnight with rocking. Embryos were then washed
≥5x in 1x PBST, incubated for ≥1 hr in 10% FBS at RT, then incubated at 4 °C overnight in 1:500
donkey anti-mouse HRP secondary antibody (Jackson ImmunoResearch Laboratories #715-035-
150) in 10% FBS. Following ≥5x PBST washes and a ≥15 min incubation in TSA amplification
buffer at RT, embryos were incubated for 45 min in a 1:75 Cy3:amplification buffer dilution. After
4x 1x PBST washes, embryos were washed 2x with MeOH, incubated at RT for ≥3 hours, re-
blocked in 10% FBS for ≥1 hour at RT, and then incubated in 1:100 rabbit anti-pSmad1/5/9
primary antibody (Cell Signaling Technology #13820S) at 4 °C overnight. The following day
embryos were washed ≥5x in 1x PBST, re-blocked in 10% FBS at RT for ≥1 hr, then incubated in
1:100 goat anti-rabbit Alexa647 secondary antibody (Invitrogen #A32733) and 1:5000 DAPI at 4

°C overnight. Embryos were washed ≥5x with 1x PBST, followed by 2x PBS washes, and imaged

within 1 month of staining completion.

**Hybridization Chain Reaction fluorescence *in situ* hybridization (HCR-FISH, mRNA**

**detection)**

Our hybridization chain reaction fluorescence *in situ* hybridization (HCR-FISH) protocol

is based on (Choi et al., 2018). Briefly, embryos were fixed in 4% formaldehyde at 4 °C overnight.

After the overnight fixation, the embryos were washed 3x with 1x PBST (1x PBS + 0.1% Tween

20) to stop fixation. Embryos were manually dechorionated in 1x PBST. The dechorionated

embryos were washed with 100% MeOH 3x, then stored overnight or longer at -20 °C.

Embryos were rehydrated briefly with 50% MeOH in PBST followed by 4x washes using

1x PBST. The residual PBST was carefully removed without disturbing the embryos. The embryos

were pre-hybridized at 37 °C with shaking at 300 rpm for 30 minutes in pre-warmed hybridization

buffer (Molecular Instruments). The HCR was multiplexed for simultaneous detection of three

different genes per pathway. Three different initiators (B1, B2, and B3) were used in combination

with three different pathway specific gene probes (Supp. Tables 11 & 12). The probe solution was

prepared by adding equal volumes of 1 uM probe solution to the pre-warmed hybridization buffer

(Molecular Instruments). Following pre-hybridization, embryos were incubated with the probe

hybridization solution at 37 °C for ≥12 hours with shaking at 300 rpm. After hybridization, excess

probe was removed by washing with probe wash buffer (Molecular Instruments) for 1 hour,

followed by washing the embryos with 5x SSCT (5x Saline-sodium citrate + 0.1% Tween 20) for

not more than 20 minutes.

Next, the pre-amplification was done by adding pre-warmed amplification buffer

(Molecular Instruments) and incubating embryos at room temperature for 30 minutes. The

amplification solution was prepared by snap cooling 30 pmol each of hairpin 1 and hairpin 2 for

three HCR amplifiers (B1-488, B2-546, and B3-647) in separate PCR tubes and combining them

with amplification buffer. The embryos were incubated with the amplification solution overnight

at room temperature with gentle rocking.

After amplification the embryos were washed briefly for 10 minutes with 5x SSCT and

then for a total of an hour. Embryos were then washed briefly with 1x PBST and stained with the

nuclear stain DAPI (1:5000) for 2 hours at room temperature. The embryos were then washed with

5x PBST at room temperature over 1 hour, then washed briefly 3x with 1x PBS. Embryos were

mounted using 1% LMP (Low melting point agarose) and imaged within two weeks after

processing.

**Hybridization Chain Reaction Immunofluorescence (HCR-IF, protein detection)**

776           Our hybridization chain reaction immunofluorescence (HCR-IF) protocol is based on

(Schwarzkopf et al., 2021). This protocol utilizes HCR-based signal amplification to detect

protein. In brief, the embryos were fixed with 4% formaldehyde in 1x PBS overnight at 4 °C. The

fixation was stopped by washing the embryos 4x with 1x PBST. Then embryos were washed with

100% MeOH 3x at room temperature to remove residual PBST. Embryos were permeabilized in

100% MeOH at -20 °C overnight or longer. The embryos were rehydrated by washing 4x using 1x

PBST. Then the embryos were blocked with antibody buffer (Molecular Instruments) for either ~4

hours at 4 °C or 1-4 hours at room temperature. The following primary antibodies were used: 1:100

rabbit anti-pSmad1/5/9 primary antibody (Cell Signaling Technology #13820S), 1:5000 rabbit

anti-pSmad2/3 primary antibody (Cell Signaling Technology #8828S) and 1:5000 mouse anti-

ppERK1/2 primary antibody (Sigma-Aldrich #M8159). Embryos were incubated in primary

antibody in antibody buffer overnight at 4 °C. The embryos were washed with 1x PBST for $\geq$2

hours with gentle rocking. Next, embryos were incubated in 1:1000 donkey-anti rabbit-B1

secondary antibody (Molecular Instruments) against rabbit anti-pSmad1, or 1:500 donkey-anti

rabbit-B1 secondary antibody (Molecular Instruments) against rabbit anti-pSmad2, or 1:500

donkey-anti mouse-B5 secondary antibody (Molecular Instruments) against mouse anti-ppERK

for 3 hours at room temperature. After incubation in secondary antibody, the embryos were washed

with 1x PBST for >30 minutes. After a brief rinse with 1x SSCT embryos were incubated in

amplification buffer (Molecular Instruments) for 30 minutes. The amplifier solution was prepared

by snap cooling hairpin 1 and hairpin 2 for corresponding HCR amplifiers (B1-647 or B5-647)

and combining them after snap cooling in amplification buffer. Embryos were incubated in HCR

amplification solution overnight at room temperature with gentle rocking, then washed with 5x

SSCT for 40 minutes and rinsed briefly with 1x PBST (For pSmad2 HCR-IF experiments in Figs.

5 and 6 and Supp. Figs. 6-9, amplification occurred at 4 °C. We found this led to better signal to

noise ratio for pSmad2). Embryos were then incubated in 1:5000 DAPI for 2.5 hours and washed

with 1x PBST for 1 hour, followed by a 1x PBS for 15 minutes and stored in PBS. Note that GFP

survives the MeOH incubation and HCR-IF protocol and was imaged directly.

**Confocal imaging**

Embryos that underwent HCR-FISH, HCR-IF, or triple immunofluorescence staining were

transferred to melted 1-1.2% low melting agarose (LMA) (in embryo medium) using a glass

pipette. Embryos were then immediately removed from the LMA, mounted on a plastic petri dish,

and oriented with the animal cap up using a stainless-steel needle. LMA was allowed to solidify

and 1x PBS was added to the petri dish before imaging the embryos.

IF, HCR-IF, and HCR-FISH-processed embryos were imaged using a Zeiss LSM 800

upright confocal laser scanning microscope. The light path consisted of the emission from one or

a combination of laser lines passing a W N-Achroplan 10x/0.3 objective before reaching the

sample plane. Reflected light was collected with 2 GaAsP-PMT and an Airyscan63 detectors. The

details of the laser lines and emission detection ranges used for all experiments are given in

Supplementary Table 1. Images were acquired with 0.7x zoom in bidirectional scanning mode with

16-bit depth, with a resolution of 512x512 pixels using ZEN microscopy software. For all

experiments, z-stacks were acquired with a 9 μm interval between slices, except for spatial

activation experiments in which a 10 μm interval was used. Remaining imaging settings (e.g., laser

power, gain, digital gain, line averaging, pixel dwell time) were consistently maintained for all

samples within a biological replicate (where there existed an internal dark control) for each

experiment and bOpto tool.

**Image Processing**

*Visualization Pipeline*

Triple IF images from pathway specificity experiments (Fig. 4 and Supp. Fig. 5) and HCR-

FISH gene expression experiments (Fig. 3) were processed using the same custom FIJI-based

pipeline (Supplementary Materials, Codes folder). In brief, for each sample, a maximum intensity

z-projection was generated, and lookup table for each acquired channel was adjusted to the defined

minimum and maximum values for that experiment.

HCR-IF images from wavelength dependence (Fig. 2), on/off kinetics (Fig. 5, Supp. Fig.

6), irradiance dependence (Fig. 6, Supp. Fig. 8), and spatial activation (Fig. 7, Supp. Fig. 10)

experiments, and HCR-FISH images from circadian experiments (Supp. Fig. 12) were processed
using a custom MATLAB-based pipeline (Supplementary Materials, Codes folder). The user is
prompted to supply lookup table bounds for visualization; this was kept consistent within each
experiment and pathway. Maximum intensity z-projections were then generated for each sample
and channel. Additionally, the user is prompted to identify a threshold for nuclear segmentation
based on the DAPI or Kaede Green channels. Afterwards, a 3D mask defining the nuclear region
is generated (hereby termed "DAPI region") and applied voxelwise to the effector signal channel
image volume. A "nuclear" maximum intensity projection is then generated.

*Quantification Pipeline*

For all HCR-IF experiments needing downstream effector signal normalization and
quantification (wavelength dependence (Fig. 2, Supp. Figs. 3 & 4), on/off kinetics (Fig. 5, Supp.
Figs. 6 & 7), irradiance dependence (Fig. 6, Supp. Figs. 8 & 9)), the MATLAB-pipeline then
prompts the user to specify a threshold for the GFP channel. Like nuclear thresholding, a 3D mask
defining the "GFP region" is generated. The intersection between the DAPI and GFP regions was
also calculated as the "DAPI+GFP region". The effector signal channel image volume is divided
voxelwise by the GFP channel image volume, creating a "normalized signal" data volume. The
following median values are then calculated for the imaged volume: 1) raw effector signal within
the DAPI region, 2) GFP-normalized effector signal within the DAPI region, 3) raw effector signal
within the GFP region, 4) GFP-normalized effector signal within the GFP region, 5) raw effector
signal within the DAPI+GFP region, and 6) GFP-normalized effector signal within the DAPI+GFP
region. Maximum z-projections of masks are exported for quality control assessment. Nuclear and
GFP thresholds were kept consistent for all samples within a biological replicate (where there
existed an internal dark control) for each experiment and bOpto tool. For FGF and BMP, the
median GFP-normalized effector signal within the GFP region was used in all downstream
analyses. As Nodal effector, pSmad2, is largely nuclear the median GFP-normalized effector signal
within the DAPI+GFP region was used.

*Quality Control*

Output data was then assessed for quality by 1-2 authors before additional processing. If
any of the following criteria were met through screening maximum intensity projections, that

sample was thrown out: 1) damage was apparent in the DAPI channel spanning 1/3 of the embryo,

2) any part of the embryo was out of frame, 3) a portion of an errant second embryo was in frame,

4) the embryo was mounted on an angle (>40 degrees off axis), 5) z-stack does not encompass

entire embryo. If the following criteria was met through screening maximum intensity projections,

that entire exposure group within the biological replicate was removed: saturation-induced

"streaking" present in any one channel. If the following criteria was met through screening masks,

that sample was removed: <5% of the embryo was covered by the appropriate GFP mask,

indicating a sample did not receive enough mRNA for analysis.

871            Any embryo imaged that passed all inclusion criteria was defined as a technical replicate

within an exposure condition and biological replicate for the appropriate experiment. Sufficient

"trials" were run for each experiment such that 3 biological replicates were acquired for all assayed

conditions within each experiment. GFP-normalized signal for all technical replicates within an

exposure condition, biological replicate, and bOpto tool were averaged together and processed as

follows for the corresponding experiments:

*Wavelength Dependence*

879            For each bOpto tool, the mean GFP-normalized signal for all samples within each

wavelength (Dark, 495+ nm, 455nm) and biological replicate was fit by a linear mixed-effects

model (Fixed Effect: Wavelength, Random Effect: Biological Replicate; Personality: Standard

Least Squares, Method: REML). Predicted least square means and standard error were calculated

for each wavelength. Where warranted by a significant ($p < 0.05$) fixed effect, *post hoc*

comparisons were performed between all exposure combinations and corrected for multiple

comparisons with Tukey's HSD method where $p < 0.05$ was considered to be statistically

significant (Supp. Tables 2-4).

*On/Off Kinetics*

889            For each bOpto tool, the mean GFP-normalized signal for all 455 nm exposed samples

within each time point (FGF: 0, 2, 8, 15, 22, 30, 35, 50, 60, 70, and 85 min; BMP/Nodal: 0, 2, 8,

18, 30, 35, 50, 60, 70, 85, and 120 min) and biological replicate was subtracted by its time-matched

dark control. The dark-subtracted, normalized data was then analyzed by a one-way ANOVA with

a fixed effect of time. Where warranted by a significant ($p < 0.05$) fixed effect, *post hoc*

comparisons were performed between each time point and time = 0 min with Dunnett's method
for multiple comparison correction where $p < 0.05$ was considered to be statistically significant
(Supp. Tables 6-7).

Next, models were fit to the dark-subtracted, GFP-normalized dataset to calculate estimates
for on and off kinetic parameters. One model was generated for the "On" portion of the data (time
<= 30) and a second model was fit to the "Off" portion of the data (time >= 30). We chose to fit a
three-parameter logistic regression to each data subset. The equation for this model is defined as:

$$y = \frac{c}{1 + e^{(-a(x-b))}} \qquad \text{(Eq.1)}$$

where y is the dark-subtracted normalized signal [AU], x is time [min], a is the growth rate [min$^-$
$^1$], b is the inflection point [min], and c is the upper asymptote [AU].
This curve fit has two underlying assumptions: 1) the general behavior of the bOpto tools will
follow a tri-phasic lag-growth/decay-plateau behavior, and 2) the lower asymptote will be 0 (i.e.,
for "ON" the dark subtracted, normalized signal begins at zero; for "OFF" the dark subtracted,
normalized signal will eventually return to zero at some point in time). $R^2$ was reported as the
goodness of fit for each model. The predicted value of each factor (a, b, c) and its corresponding
95% confidence interval is also reported (Supp. Tables 8-9).

Additionally for the model fitting the "on" data, we used the predicted equation to solve
for time at various points along the curve to more easily describe the activation kinetics.
(Supplementary Materials) We took the value for the upper asymptote, c, and calculated three
different percentages (20%, 50%, and 90%) of that value. We then used the curve fit equation to
predict the time it takes from the start to reach those values. These conceptually represent: $tON_{20}$
predicting initial activation, $tON_{50}$ predicting half-maximum, and $tON_{90}$ predicting saturation.
Additionally, for the model fitting the "off" data, we performed an analogous analysis with one
modification: we calculated the time from light removal to reach three different percentage
*decreases* of the upper asymptote (20%, 50%, and 90% decreases). These conceptually represent:
$tOFF_{20}$ predicted time to initial deactivation, $tOFF_{50}$ predicted time to half-maximum, and $tOFF_{90}$
predicted approximate time to return to baseline. The predicted value of each tON and tOFF factor
and its corresponding 95% confidence interval was reported in (Supp. Tables 8-9).

*Irradiance Dependence*

For each bOpto tool, the mean GFP-normalized signal for all samples within each irradiance (0, 0.02, 0.05, 0.15, 0.46, 1.39, 4.17, 12.50, 50.00 $W/m^2$) and biological replicate was subtracted by its round-matched dark control (Supp. Table 10). Data were fit with the same three-parameter logistic regression model used in the on/off kinetics experiments. (Eq. 1) The predicted value of each factor (a, b, c) and its corresponding 95% confidence interval was reported. From the model, three values were calculated approximating the irradiance at which a percentage (20%, 50%, and 90%) of the curve's upper asymptote is reached: $I_{20}$ predicting initial response, $I_{50}$ predicting half-maximum, and $I_{90}$ predicting saturation. The predicted value of each factor and its corresponding 95% confidence interval was reported in (Supp. Table 10).

**References:**

**Ando, R., Hama, H., Yamamoto-Hino, M., Mizuno, H. and Miyawaki, A.** (2002). An optical marker based on the UV-induced green-to-red photoconversion of a fluorescent protein. *Proc Natl Acad Sci U S A* **99**, 12651-12656.

**Belacortu, Y. and Paricio, N.** (2011). Drosophila as a model of wound healing and tissue regeneration in vertebrates. *Dev Dyn* **240**, 2379-2404.

**Benedetti, L., Barentine, A. E. S., Messa, M., Wheeler, H., Bewersdorf, J. and De Camilli, P.** (2018). Light-activated protein interaction with high spatial subcellular confinement. *Proc Natl Acad Sci U S A* **115**, E2238-E2245.

**Benman, W., Berlew, E. E., Deng, H., Parker, C., Kuznetsov, I. A., Lim, B., Siekmann, A. F., Chow, B. Y. and Bugaj, L. J.** (2022). Temperature-responsive optogenetic probes of cell signaling. *Nat Chem Biol* **18**, 152-160.

**Bennett, J. T., Joubin, K., Cheng, S., Aanstad, P., Herwig, R., Clark, M., Lehrach, H. and Schier, A. F.** (2007). Nodal signaling activates differentiation genes during zebrafish gastrulation. *Dev Biol* **304**, 525-540.

**Bisgrove, B. W., Essner, J. J. and Yost, H. J.** (1999). Regulation of midline development by antagonism of lefty and nodal signaling. *Development* **126**, 3253-3262.

**Bosman, S. L. and Sonnen, K. F.** (2022). Signaling oscillations in embryonic development. *Curr Top Dev Biol* **149**, 341-372.

**Briscoe, J. and Small, S.** (2015). Morphogen rules: design principles of gradient-mediated embryo patterning. *Development* **142**, 3996-4009.

**Buckley, C. E.** (2019). Optogenetic Control of Subcellular Protein Location and Signaling in Vertebrate Embryos. *Methods Mol Biol* **1920**, 143-162.

**Buckley, C. E., Moore, R. E., Reade, A., Goldberg, A. R., Weiner, O. D. and Clarke, J. D. W.** (2016). Reversible optogenetic control of subcellular protein localization in a live vertebrate embryo. *Dev Cell* **36**, 117-126.

**Čapek, D., Smutny, M., Tichy, A. M., Morri, M., Janovjak, H. and Heisenberg, C. P.** (2019). Light-activated Frizzled7 reveals a permissive role of non-canonical wnt signaling in mesendoderm cell migration. *Elife* **8**, e42093.

**Chang, K. Y., Woo, D., Jung, H., Lee, S., Kim, S., Won, J., Kyung, T., Park, H., Kim, N., Yang, H. W., et al.** (2014). Light-inducible receptor tyrosine kinases that regulate neurotrophin signalling. *Nat Commun* **5**, 4057.

**Choi, H. M. T., Schwarzkopf, M., Fornace, M. E., Acharya, A., Artavanis, G., Stegmaier, J., Cunha, A. and Pierce, N. A.** (2018). Third-generation in situ hybridization chain reaction: multiplexed, quantitative, sensitive, versatile, robust. *Development* **145**.

**Choi, I. Y., Lim, H., Huynh, A., Schofield, J., Cho, H. J., Lee, H., Andersen, P., Shin, J. H., Heo, W. D., Hyun, S. H., et al.** (2021). Novel culture system via wirelessly controllable optical stimulation of the FGF signaling pathway for human and pig pluripotency. *Biomaterials* **269**, 120222.

**Christen, B. and Slack, J. M.** (1999). Spatial response to fibroblast growth factor signalling in Xenopus embryos. *Development* **126**, 119-125.

**Crossman, S. H. and Janovjak, H.** (2022). Light-activated receptor tyrosine kinases: Designs and applications. *Curr Opin Pharmacol* **63**, 102197.

**Csanaky, K., Hess, M. W. and Klimaschewski, L.** (2019). Membrane-Associated, Not Cytoplasmic or Nuclear, FGFR1 Induces Neuronal Differentiation. *Cells* **8**.

**De Simone, A., Evanitsky, M. N., Hayden, L., Cox, B. D., Wang, J., Tornini, V. A., Ou, J., Chao, A., Poss, K. D. and Di Talia, S.** (2021). Control of osteoblast regeneration by a train of Erk activity waves. *Nature*.

**Dekens, M. P., Santoriello, C., Vallone, D., Grassi, G., Whitmore, D. and Foulkes, N. S.** (2003). Light regulates the cell cycle in zebrafish. *Curr Biol* **13**, 2051-2057.

**Derynck, R. and Budi, E. H.** (2019). Specificity, versatility, and control of TGF-beta family signaling. *Sci Signal* **12**.

**Dine, E., Gil, A. A., Uribe, G., Brangwynne, C. P. and Toettcher, J. E.** (2018). Protein phase separation provides long-term memory of transient spatial stimuli. *Cell Syst* **6**, 655-663 e655.

**Donahue, C. E. T., Siroky, M. D. and White, K. A.** (2021). An Optogenetic Tool to Raise
Intracellular pH in Single Cells and Drive Localized Membrane Dynamics. *J Am Chem Soc* **143**,
18877-18887.

**Donnelly, M. L. L., Hughes, L. E., Luke, G., Mendoza, H., Ten Dam, E., Gani, D. and Ryan,
M. D.** (2001a). The 'cleavage' activities of foot-and-mouth disease virus 2A site-directed mutants
and naturally occurring '2A-like' sequences. *J Gen Virol* **82**, 1027-1041.

**Donnelly, M. L. L., Luke, G., Mehrotra, A., Li, X., Hughes, L. E., Gani, D. and Ryan, M. D.**
(2001b). Analysis of the aphthovirus 2A/2B polyprotein 'cleavage' mechanism indicates not a
proteolytic reaction, but a novel translational effect: a putative ribosomal 'skip'. *J Gen Virol* **82**,
1013-1025.

**Emig, A. A., Hansen, M., Grimm, S., Coarfa, C., Lord, N. D. and Williams, M. K.** (2024).
Temporal dynamics of BMP/Nodal ratio drive tissue-specific gastrulation morphogenesis.
*Development*.

**Farahani, P. E., Reed, E. H., Underhill, E. J., Aoki, K. and Toettcher, J. E.** (2021). Signaling,
Deconstructed: Using Optogenetics to Dissect and Direct Information Flow in Biological
Systems. *Annu Rev Biomed Eng* **23**, 61-87.

**Farahani, P. E., Yang, X., Mesev, E. V., Fomby, K. A., Brumbaugh-Reed, E. H., Bashor, C.
J., Nelson, C. M. and Toettcher, J. E.** (2023). pYtags enable spatiotemporal measurements of
receptor tyrosine kinase signaling in living cells. *Elife* **12**.

**Feldman, B., Concha, M. L., Saude, L., Parsons, M. J., Adams, R. J., Wilson, S. W. and
Stemple, D. L.** (2002). Lefty antagonism of squint is essential for normal gastrulation. *Current
Biology* **12**, 2129-2135.

**Froland Steindal, I. A. and Whitmore, D.** (2019). Circadian Clocks in Fish-What Have We
Learned so far? *Biology (Basel)* **8**.

**Fürthauer, M., Lin, W., Ang, S. L., Thisse, B. and Thisse, C.** (2002). Sef is a feedback-
induced antagonist of Ras/MAPK-mediated FGF signalling. *Nat Cell Biol* **4**, 170-174.

**Fürthauer, M., Reifers, F., Brand, M., Thisse, B. and Thisse, C.** (2001). sprouty4 acts in vivo
as a feedback-induced antagonist of FGF signaling in zebrafish. *Development* **128**, 2175-2186.

**Fürthauer, M., Thisse, C. and Thisse, B.** (1997). A role for FGF-8 in the dorsoventral
patterning of the zebrafish gastrula. *Development* **124**, 4253-4264.

**Fürthauer, M., Van Celst, J., Thisse, C. and Thisse, B.** (2004). Fgf signalling controls the
dorsoventral patterning of the zebrafish embryo. *Development* **131**, 2853-2864.

**Greenfeld, H., Lin, J. and Mullins, M. C.** (2021). The BMP signaling gradient is interpreted
through concentration thresholds in dorsal-ventral axial patterning. *PLoS Biol* **19**, e3001059.

**Grusch, M., Schelch, K., Riedler, R., Reichhart, E., Differ, C., Berger, W., Ingles-Prieto, A.**
**and Janovjak, H.** (2014). Spatio-temporally precise activation of engineered receptor tyrosine
kinases by light. *EMBO J* **33**, 1713-1726.

**Guglielmi, G., Barry, J. D., Huber, W. and De Renzis, S.** (2015). An optogenetic method to
modulate cell contractility during tissue morphogenesis. *Dev Cell* **35**, 646-660.

**Guillotin, B. and Birnbaum, K. D.** (2020). Just passing through: The auxin gradient of the root
meristem. *Curr Top Dev Biol* **137**, 433-454.

**Hadpech, S., Jinathep, W., Saoin, S., Thongkum, W., Chupradit, K., Yasamut, U.,**
**Moonmuang, S. and Tayapiwatana, C.** (2018). Impairment of a membrane-targeting protein
translated from a downstream gene of a "self-cleaving" T2A peptide conjunction. *Protein Expr*
*Purif* **150**, 17-25.

**Heldin, C. H. and Moustakas, A.** (2016). Signaling Receptors for TGF-beta Family Members.
*Cold Spring Harb Perspect Biol* **8**.

**Hill, C. S.** (2022). Establishment and interpretation of NODAL and BMP signaling gradients in
early vertebrate development. *Curr Top Dev Biol* **149**, 311-340.

**Huang, A., Amourda, C., Zhang, S., Tolwinski, N. S. and Saunders, T. E.** (2017). Decoding
temporal interpretation of the morphogen Bicoid in the early Drosophila embryo. *Elife* **6**,
e26258.

**Humphreys, P. A., Woods, S., Smith, C. A., Bates, N., Cain, S. A., Lucas, R. and Kimber, S.**
**J.** (2020). Optogenetic Control of the BMP Signaling Pathway. *ACS Synth Biol*.

**Humphreys, P. E. A., Woods, S., Bates, N., Rooney, K. M., Mancini, F. E., Barclay, C.,**
**O'Flaherty, J., Martial, F. P., Domingos, M. A. N. and Kimber, S. J.** (2023). Optogenetic
manipulation of BMP signaling to drive chondrogenic differentiation of hPSCs. *Cell Rep* **42**,
113502.

**Izquierdo, E., Quinkler, T. and De Renzis, S.** (2018). Guided morphogenesis through
optogenetic activation of Rho signalling during early Drosophila embryogenesis. *Nat Commun* **9**,
2366.

**Johnson, H. E. and Toettcher, J. E.** (2018). Illuminating developmental biology with cellular
optogenetics. *Curr Opin Biotechnol* **52**, 42-48.

**Johnson, H. E. and Toettcher, J. E.** (2019). Signaling dynamics control cell fate in the early
*Drosophila* embryo. *Developmental Cell* **48**, 361-370.e363.

**Jones, W. D. and Mullins, M. C.** (2022). Cell signaling pathways controlling an axis organizing
center in the zebrafish. *Curr Top Dev Biol* **150**, 149-209.

**Kainrath, S., Stadler, M., Reichhart, E., Distel, M. and Janovjak, H.** (2017). Green-light-
induced inactivation of receptor signaling using cobalamin-binding domains. *Angew Chem Int*
*Ed Engl* **56**, 4608-4611.

**Kemmler, C. L., Moran, H. R., Murray, B. F., Scoresby, A., Klem, J. R., Eckert, R. L.,**
**Lepovsky, E., Bertho, S., Nieuwenhuize, S., Burger, S., et al.** (2023). Next-generation
plasmids for transgenesis in zebrafish and beyond. *Development* **150**.

**Kicheva, A. and Briscoe, J.** (2023). Control of Tissue Development by Morphogens. *Annu Rev*
*Cell Dev Biol* **39**, 91-121.

**Kim, N., Kim, J. M., Lee, M., Kim, C. Y., Chang, K. Y. and Heo, W. D.** (2014).
Spatiotemporal control of fibroblast growth factor receptor signals by blue light. *Chem Biol* **21**,
903-912.

**Kishimoto, Y., Lee, K. H., Zon, L., Hammerschmidt, M. and Schulte-Merker, S.** (1997). The
molecular nature of zebrafish swirl: BMP2 function is essential during early dorsoventral
patterning. *Development* **124**, 4457-4466.

**Kolar, K., Knobloch, C., Stork, H., Znidaric, M. and Weber, W.** (2018). OptoBase: A web
platform for molecular optogenetics. *ACS Synth Biol* **7**, 1825-1828.

**Koshida, S., Shinya, M., Nikaido, M., Ueno, N., Schulte-Merker, S., Kuroiwa, A. and**
**Takeda, H.** (2002). Inhibition of BMP activity by the FGF signal promotes posterior neural
development in zebrafish. *Dev Biol* **244**, 9-20.

**Kretzschmar, M., Doody, J. and Massague, J.** (1997). Opposing BMP and EGF signalling
pathways converge on the TGF-beta family mediator Smad1. *Nature* **389**, 618-622.

**Krishnamurthy, V. V., Fu, J., Oh, T. J., Khamo, J., Yang, J. and Zhang, K.** (2020). A
Generalizable Optogenetic Strategy to Regulate Receptor Tyrosine Kinases during Vertebrate
Embryonic Development. *J Mol Biol* **432**, 3149-3158.

**Krylov, V. V., Izvekov, E. I., Pavlova, V. V., Pankova, N. A. and Osipova, E. A.** (2021).
Circadian rhythms in zebrafish (Danio rerio) behaviour and the sources of their variability. *Biol*
*Rev Camb Philos Soc* **96**, 785-797.

**Kudoh, T., Concha, M. L., Houart, C., Dawid, I. B. and Wilson, S. W.** (2004). Combinatorial
Fgf and Bmp signalling patterns the gastrula ectoderm into prospective neural and epidermal
domains. *Development* **131**, 3581-3592.

**LaBelle, J., Ramos-Martinez, A., Shen, K., Motta-Mena, L. B., Gardner, K. H., Materna, S.**
**C. and Woo, S.** (2021). TAEL 2.0: An Improved Optogenetic Expression System for Zebrafish.
*Zebrafish* **18**, 20-28.

**LaBonne, C. and Whitman, M.** (1997). Localization of MAP kinase activity in early Xenopus
embryos: implications for endogenous FGF signaling. *Dev Biol* **183**, 9-20.

**Laranjeiro, R. and Whitmore, D.** (2014). Transcription factors involved in retinogenesis are co-opted by the circadian clock following photoreceptor differentiation. *Development* **141**, 2644-2656.

**Leerberg, D. M., Hopton, R. E. and Draper, B. W.** (2019). Fibroblast Growth Factor Receptors Function Redundantly During Zebrafish Embryonic Development. *Genetics* **212**, 1301-1319.

**Li, J., Zhang, S., Soto, X., Woolner, S. and Amaya, E.** (2013). ERK and phosphoinositide 3-kinase temporally coordinate different modes of actin-based motility during embryonic wound healing. *J Cell Sci* **126**, 5005-5017.

**Liberali, P. and Schier, A. F.** (2024). The evolution of developmental biology through conceptual and technological revolutions. *Cell*.

**Lim, S. M., Pereira, L., Wong, M. S., Hirst, C. E., Van Vranken, B. E., Pick, M., Trounson, A., Elefanty, A. G. and Stanley, E. G.** (2009). Enforced expression of Mixl1 during mouse ES cell differentiation suppresses hematopoietic mesoderm and promotes endoderm formation. *Stem Cells* **27**, 363-374.

**Mathieu, J., Griffin, K., Herbomel, P., Dickmeis, T., Strahle, U., Kimelman, D., Rosa, F. M. and Peyrieras, N.** (2004). Nodal and Fgf pathways interact through a positive regulatory loop and synergize to maintain mesodermal cell populations. *Development* **131**, 629-641.

**McNamara, H. M., Guyer, A. M., Jia, B. Z., Parot, V. J., Dobbs, C. D., Schier, A. F., Cohen, A. E. and Lord, N. D.** (2025). Optogenetic control of Nodal signaling patterns. *Development*.

**Mitra, D., Yang, X. and Moffat, K.** (2012). Crystal structures of Aureochrome1 LOV suggest new design strategies for optogenetics. *Structure* **20**, 698-706.

**Molina, G. A., Watkins, S. C. and Tsang, M.** (2007). Generation of FGF reporter transgenic zebrafish and their utility in chemical screens. *BMC Dev Biol* **7**, 62.

**Mruk, K., Ciepla, P., Piza, P. A., Alnaqib, M. A. and Chen, J. K.** (2020). Targeted cell ablation in zebrafish using optogenetic transcriptional control. *Development* **147**.

**Mullins, M. C., Acedo, J. N., Priya, R., Solnica-Krezel, L. and Wilson, S. W.** (2021). The zebrafish issue: 25 years on. *Development* **148**.

**Nguyen, V. H., Schmid, B., Trout, J., Connors, S. A., Ekker, M. and Mullins, M. C.** (1998). Ventral and lateral regions of the zebrafish gastrula, including the neural crest progenitors, are established by a bmp2b/swirl pathway of genes. *Dev Biol* **199**, 93-110.

**Ornitz, D. M. and Itoh, N.** (2015). The Fibroblast Growth Factor signaling pathway. *Wiley Interdiscip Rev Dev Biol* **4**, 215-266.

**Owlarn, S., Klenner, F., Schmidt, D., Rabert, F., Tomasso, A., Reuter, H., Mulaw, M. A., Moritz, S., Gentile, L., Weidinger, G., et al.** (2017). Generic wound signals initiate regeneration in missing-tissue contexts. *Nat Commun* **8**, 2282.

**Pal, A. A., Benman, W., Mumford, T. R., Huang, Z., Chow, B. Y. and Bugaj, L. J.** (2023).
Optogenetic clustering and membrane translocation of the BcLOV4 photoreceptor. *Proc Natl*
*Acad Sci U S A* **120**, e2221615120.

**Patel, A. L., Yeung, E., McGuire, S. E., Wu, A. Y., Toettcher, J. E., Burdine, R. D. and**
**Shvartsman, S. Y.** (2019). Optimizing photoswitchable MEK. *Proc Natl Acad Sci U S A* **116**,
25756-25763.

**Poss, K. D., Shen, J., Nechiporuk, A., McMahon, G., Thisse, B., Thisse, C. and Keating, M.**
**T.** (2000). Roles for Fgf signaling during zebrafish fin regeneration. *Dev Biol* **222**, 347-358.

**Purvis, J. E. and Lahav, G.** (2013). Encoding and decoding cellular information through
signaling dynamics. *Cell* **152**, 945-956.

**Ramachandran, A., Vizan, P., Das, D., Chakravarty, P., Vogt, J., Rogers, K. W., Müller, P.,**
**Hinck, A. P., Sapkota, G. P. and Hill, C. S.** (2018). TGF-β uses a novel mode of receptor
activation to phosphorylate SMAD1/5 and induce epithelial-to-mesenchymal transition. *Elife* **7**,
e31756.

**Ramkumar, N., Richardson, C., O'Brien, M., Butt, F. A., Park, J., Chao, A. T., Bagnat, M.,**
**Poss, K. D. and Di Talia, S.** (2025). Phased ERK responsiveness and developmental robustness
regulate teleost skin morphogenesis. *Proc Natl Acad Sci U S A* **122**, e2410430122.

**Rauschendorfer, T., Gurri, S., Heggli, I., Maddaluno, L., Meyer, M., Ingles-Prieto, A.,**
**Janovjak, H. and Werner, S.** (2021). Acute and chronic effects of a light-activated FGF
receptor in keratinocytes in vitro and in mice. *Life Sci Alliance* **4**.

**Reade, A., Motta-Mena, L. B., Gardner, K. H., Stainier, D. Y., Weiner, O. D. and Woo, S.**
(2017). TAEL: a zebrafish-optimized optogenetic gene expression system with fine spatial and
temporal control. *Development* **144**, 345-355.

**Rebagliati, M. R., Toyama, R., Fricke, C., Haffter, P. and Dawid, I. B.** (1998). Zebrafish
nodal-related genes are implicated in axial patterning and establishing left-right asymmetry. *Dev*
*Biol* **199**, 261-272.

**Rogers, K. W., ElGamacy, M., Jordan, B. M. and Müller, P.** (2020). Optogenetic investigation
of BMP target gene expression diversity. *Elife* **9**, e58641.

**Rogers, K. W., Lord, N. D., Gagnon, J. A., Pauli, A., Zimmerman, S., Aksel, D. C., Reyon,**
**D., Tsai, S. Q., Joung, J. K. and Schier, A. F.** (2017). Nodal patterning without Lefty inhibitory
feedback is functional but fragile. *Elife* **6**, e28785.

**Rogers, K. W. and Müller, P.** (2020). Optogenetic approaches to investigate spatiotemporal
signaling during development. In *Gradients and Tissue Patterning*, pp. 37-77.

**Ryan, M. D., King, A. M. and Thomas, G. P.** (1991). Cleavage of foot-and-mouth disease virus
polyprotein is mediated by residues located within a 19 amino acid sequence. *J Gen Virol* **72 ( Pt**
**11)**, 2727-2732.

**Sagner, A. and Briscoe, J.** (2017). Morphogen interpretation: concentration, time, competence,
and signaling dynamics. *Wiley Interdiscip Rev Dev Biol* **6**, e271.

**Sako, K., Pradhan, S. J., Barone, V., Inglés-Prieto, Á., Müller, P., Ruprecht, V., Čapek, D.,**
**Galande, S., Janovjak, H. and Heisenberg, C. P.** (2016). Optogenetic control of Nodal
signaling reveals a temporal pattern of Nodal signaling regulating cell fate specification during
gastrulation. *Cell Rep* **16**, 866-877.

**Saul, A. J.\*, Rogers, C. E.\*, Garmendia-Cedillos, M., Pohida, T. and Rogers, K. W.** (2023).
Optogenetic Signaling Activation in Zebrafish Embryos. *Journal of Visualized Experiments*.

**Schmid, B., Furthauer, M., Connors, S. A., Trout, J., Thisse, B., Thisse, C. and Mullins, M.**
**C.** (2000). Equivalent genetic roles for bmp7/snailhouse and bmp2b/swirl in dorsoventral pattern
formation. *Development* **127**, 957-967.

**Schwarzkopf, M., Liu, M. C., Schulte, S. J., Ives, R., Husain, N., Choi, H. M. T. and Pierce,**
**N. A.** (2021). Hybridization chain reaction enables a unified approach to multiplexed,
quantitative, high-resolution immunohistochemistry and in situ hybridization. *Development* **148**.

**Soh, G. H., Pomreinke, A. P. and Muller, P.** (2020). Integration of Nodal and BMP Signaling
by Mutual Signaling Effector Antagonism. *Cell Rep* **31**, 107487.

**Takahashi, F., Yamagata, D., Ishikawa, M., Fukamatsu, Y., Ogura, Y., Kasahara, M.,**
**Kiyosue, T., Kikuyama, M., Wada, M. and Kataoka, H.** (2007). AUREOCHROME, a
photoreceptor required for photomorphogenesis in stramenopiles. *Proc Natl Acad Sci U S A* **104**,
19625-19630.

**Tamai, T. K., Vardhanabhuti, V., Foulkes, N. S. and Whitmore, D.** (2004). Early embryonic
light detection improves survival. *Current Biology* **14**, R104-R105.

**Toettcher, J. E., Weiner, O. D. and Lim, W. A.** (2013). Using optogenetics to interrogate the
dynamic control of signal transmission by the Ras/Erk module. *Cell* **155**, 1422-1434.

**Toyooka, T., Hisatomi, O., Takahashi, F., Kataoka, H. and Terazima, M.** (2011).
Photoreactions of aureochrome-1. *Biophys J* **100**, 2801-2809.

**Tucker, J. A., Mintzer, K. A. and Mullins, M. C.** (2008). The BMP signaling gradient patterns
dorsoventral tissues in a temporally progressive manner along the anteroposterior axis. *Dev Cell*
**14**, 108-119.

**Ueda, Y., Miura, Y., Tomishige, N., Sugimoto, N., Murase, M., Kawamura, G., Sasaki, N.,**
**Ishiwata, T. and Ozawa, T.** (2022). Mechanistic insights into cancer drug resistance through
optogenetic PI3K signaling hyperactivation. *Cell Chem Biol* **29**, 1576-1587 e1575.

**van Boxtel, A. L., Economou, A. D., Heliot, C. and Hill, C. S.** (2018). Long-range signaling
activation and local inhibition separate the mesoderm and endoderm lineages. *Dev Cell* **44**, 179-
191 e175.

**Vopalensky, P., Pralow, S. and Vastenhouw, N. L.** (2018). Reduced expression of the Nodal
co-receptor Oep causes loss of mesendodermal competence in zebrafish. *Development* **145**,
dev.158832.

**Whitehead, G. G., Makino, S., Lien, C. L. and Keating, M. T.** (2005). fgf20 is essential for
initiating zebrafish fin regeneration. *Science* **310**, 1957-1960.

**Whitmore, D., Foulkes, N. S. and Sassone-Corsi, P.** (2000). Light acts directly on organs and
cells in culture to set the vertebrate circadian clock. *Nature* **404**, 87-91.

**Wilcockson, S. G., Guglielmi, L., Araguas Rodriguez, P., Amoyel, M. and Hill, C. S.** (2023).
An improved Erk biosensor detects oscillatory Erk dynamics driven by mitotic erasure during
early development. *Developmental Cell*.

**Yadav, V., Tolwinski, N. and Saunders, T. E.** (2021). Spatiotemporal sensitivity of mesoderm
specification to FGFR signalling in the Drosophila embryo. *Sci Rep* **11**, 14091.

**Zaucker, A., Mitchell, C. A., Coker, H. L. E. and Sampath, K.** (2021). Tools to Image
Germplasm Dynamics During Early Zebrafish Development. *Front Cell Dev Biol* **9**, 712503.

**Zhou, X. X., Fan, L. Z., Li, P., Shen, K. and Lin, M. Z.** (2017). Optical control of cell
signaling by single-chain photoswitchable kinases. *Science* **355**, 836-842.

**Ziv, L. and Gothilf, Y.** (2006). Circadian time-keeping during early stages of development. *Proc*
*Natl Acad Sci U S A* **103**, 4146-4151.
1216

**Figure 1: Optogenetic signaling activator toolkit. A,B,C)** Schematic of constructs used here to activate FGF (A), BMP (B), and Nodal (C) signaling. Myr = myristoylation motif, GS = glycine/serine linker, LOV = light oxygen voltage-sensing domain, HA = hemagglutinin tag, FLAG = FLAG epitope. The single-transcript bOpto-2A-Nodal construct (C) follows the same design as -FGF and -BMP, except the type I (Acvr1ba) and type II (Acvr2ba) components are connected via a 2A peptide sequence (gray). **A',B',C')** Optogenetic strategy to activate FGF (A'), BMP (B'), and Nodal (C') signaling. Blue light-dimerizing LOV domains are fused to myristoylated receptor kinase domains. Blue light exposure should lead to receptor kinase interactions, signaling effector phosphorylation, and activation of target genes.

**Figure 2: Wavelength-dependent activation of FGF, BMP, & Nodal signaling. A)** Uninjected (-) embryos and embryos injected (+) at the one-cell stage with the indicated mRNA were exposed to dark, 495+ nm light (18.51 W/m$^2$), or 455 nm light (50 W/m$^2$) starting ~2 hours post-fertilization (hpf). Phenotypes were scored at 1 day post-fertilization (dpf) (N = 3; Supp. Fig. 2; Mean +/- SD). **B,C,D)** Uninjected embryos and embryos injected with the indicated *bOpto + GFP* mRNA were exposed to dark, 495+ nm light, or 455 nm light starting at early gastrulation (50% epiboly - shield) for 30 min. HCR-IF was used to detect phosphorylated signaling effectors (ppERK, pSmad1, and pSmad2 reflect FGF, BMP, and Nodal signaling, respectively). Scale bar is 200 *μ*m. **B',C',D')** Quantification of experiments shown in B,C,D. Linear-mixed model-predicted least squared means of GFP-normalized phosphorylated effector signal +/- SEM. D' shows nuclear signal only. (N = 3, * indicates *p* < 0.05; Supp. Figs. 3 & 4).

**Figure 3: bOpto tools activate target gene expression.** Uninjected embryos (-) and embryos injected (+) at the one-cell stage with mRNA encoding bOpto-FGF **(A)**, bOpto-BMP **(B)**, or bOpto-2A-Nodal **(C)** were exposed to 455 nm light (50 W/m$^2$, bottom rows) or dark (top rows) for 2 hours starting before gastrulation (dome - 30% epiboly). Multiplexed HCR-FISH was used to detect expression of the indicated pathway target genes. (N = 3; Scale bar is 200 $\mu$m).

**Figure 4: Pathway-specific optogenetic activation of FGF, BMP, & Nodal signaling.** Uninjected embryos (-) and embryos injected (+) at the one-cell stage with mRNA encoding bOpto-FGF **(A)**, bOpto-BMP **(B)**, or bOpto-2A-Nodal **(C)** were exposed to dark (left panel) or 455 nm light (50 W/m², right panel) for 30 minutes starting at gastrulation (50% epiboly - shield). Triple IF was used to simultaneously detect phosphorylated signaling effectors (ppERK, pSmad1, and pSmad2 reflect FGF, BMP, and Nodal signaling, respectively). (N = 3; Supp. Fig. 5; Scale bar is 200 $\mu$m).

**Figure 5: On/off kinetics of optogenetic signaling activator toolkit.** Embryos were injected at the one-cell stage with mRNA encoding *GFP* and either *bOpto-FGF*, *bOpto-BMP*, or *bOpto-2A-Nodal* . **A,B,C)** Starting at early gastrulation (50% epiboly - shield), embryos were either kept in the dark (top row) or exposed to 455 nm light (50 W/m$^2$) light for 30 min (bottom row) and fixed during and after exposure. HCR-IF was used to detect activated signaling effectors (ppERK, pSmad1, and pSmad2 reflect FGF, BMP, and Nodal signaling, respectively). (N = 3; Scale bar is 200 $\mu$m). **A',B',C')** Quantification of experiments shown in A-C. Signal was GFP-normalized and subtracted against dark time-matched controls (N = 3; each N indicated by matched shape; Supp. Fig. 6 & 7; Mean +/- SD; * indicates $p < 0.05$ when compared to time = 0 min). **A''- C'', A'''-C''')** A three-parameter logistic regression was fit to the 0-30 min data (A''-C'') or the >= 30 min data (A'''-C'''). (N = 3; Mean +/- SD; Supplementary Materials; solid line represents predicted curve fit +/- 95% CI).

**A**

| Irradiance (W/m²) | All | 0.02 | 0.05 | 0.15 | 0.46 | 1.39 | 4.17 | 12.50 | 50.00 |
|---|---|---|---|---|---|---|---|---|---|
| Dosage (J/m²) — FGF (5 min) | | 4.63 | 13.89 | 46.30 | 138.89 | 416.67 | 1250.00 | 3750.00 | 15000.00 |
| BMP, Nodal (25 min) | | 23.15 | 69.44 | 231.48 | 694.44 | 2083.33 | 6250.00 | 18750.00 | 75000.00 |

**Figure 6: Irradiance sensitivity of optogenetic toolkit. A)** Table of irradiance and corresponding light dosage values used in B-D'. **B,C,D)** Embryos were injected at the one-cell stage with mRNA encoding *GFP* and either *bOpto-FGF* (B), *bOpto-BMP* (C), or *bOpto-2A-Nodal* (D). Starting at early gastrulation (50% epiboly - shield), embryos were exposed to 455 nm light at the indicated irradiance for 5 min (bOpto-FGF, B) or 25 min (bOpto-BMP and -2A-Nodal, C and D). HCR-IF was used to detect phosphorylated signaling effectors (ppERK, pSmad1, and pSmad2 reflect FGF, BMP, and Nodal signaling, respectively; Scale bar is 200 $\mu$m). **B',C',D')** Quantification of experiments in B-D. Signal was GFP-normalized and subtracted against dark controls. A three-parameter logistic regression was fit to the data. (N = 2-3; each N indicated by matched shape; Mean +/- SD; Supp. Fig. 8 & 9; solid line represents predicted curve fit +/- 95% CI; D' shows nuclear signal only). Tables indicate goodness of fit ($R^2$), the predicted irradiance (with 95% CI) at which 20 ($I_{20}$), 50 ($I_{50}$), and 90% ($I_{90}$) of the curve's upper asymptote is reached (Supplementary Materials).

**Figure 7: Spatially localized FGF signaling activation.** Embryos were injected at the one-cell stage with mRNA encoding the green-to-red photoconvertible fluorescent protein *nls-Kaede* and *bOpto-FGF*. At early gastrulation (shield) embryos were either kept in the dark **(A)** or illuminated locally with 405 and 445 nm confocal lasers **(B,B')**. HCR-IF was used to detect ppERK. Dotted white line in B' outlines photoconverted Kaede in embryo shown in B. (Supp. Fig. 10; Scale bar is 200 µm).

**Supplementary Figure 1: Light source spectra.** Measured spectra for **A)** the 455 nm light used in uniform blue light exposure experiments (all figures except Fig. 7), **B)** the 495+ nm used in Figs. 2 and Supp. Figs. 2, 3, 4, & 12, and **C)** the 600 nm light source used to avoid inadvertent signaling activation during handling of embryos expressing bOpto constructs.

**Supplementary Figure 2: Wavelength-dependent activation of FGF, BMP, & Nodal signaling experiment repeats. A-C")** Uninjected embryos (-) and embryos injected (+) at the one-cell stage with the *bOpto-FGF* (A-A"), *bOpto-BMP* (B-B"), or *bOpto-2A-Nodal* (C-C") mRNA were exposed to dark, 495+ nm light (18.51 W/m²), or 455 nm light (50 W/m²) starting 1.5-2 hours post-fertilization. Phenotypes were scored at 1 day post-fertilization. Individual repeats are shown. Numbers indicate the total number of embryos scored in each condition for that experiment. Figure 2A shows the combined data from these three repeats. Phenotype legend images shown here are the same images shown in Figure 2A.

**Supplementary Figure 3: Wavelength-dependent activation of FGF, BMP, & Nodal signaling quantification I. A-C')** Embryos injected at the one-cell stage with the indicated mRNA (+ *GFP* mRNA) were exposed to dark, 495+ nm light (18.51 W/m²), or 455 nm light (50 W/m²) starting at early gastrulation (50% epiboly - shield) for 30 min. HCR-IF was used to detect activated signaling effectors (ppERK, pSmad1, and pSmad2 reflect FGF, BMP, and Nodal signaling, respectively). A,B,C) Raw phosphorylated signaling effector intensity was measured in each DAPI-positive nuclear pixel. Each dot represents the median nuclear pixel intensity in one embryo. Black lines represent the mean nuclear intensity of all embryos in the indicated condition. A', B') GFP-normalized phosphorylated effector signal in each GFP-positive pixel. Each dot represents the median GFP-normalized pixel signal in one embryo. Black lines represent the mean GFP-normalized signal of all embryos in the indicated condition. C') pSmad2 intensity in each DAPI + GFP-positive pixel was divided by the corresponding GFP intensity. Each dot represents the median GFP-normalized nuclear pixel signal in one embryo. Black lines represent the mean GFP-normalized nuclear signal of all embryos in the indicated condition.

**Supplementary Figure 4: Wavelength-dependent activation of FGF, BMP, & Nodal signaling quantification II.** Uninjected embryos (-) and embryos injected (+) at the one-cell stage with the indicated mRNA + *GFP* mRNA were exposed to dark, 495+ nm light, or 455 nm light starting at early gastrulation (50% epiboly - shield) for 30 min. HCR-IF was used to detect phosphorylated signaling effectors (ppERK, pSmad1, and pSmad2 reflect FGF, BMP, and Nodal signaling, respectively). **A,B,C)** Raw phosphorylated signaling effector intensity was measured in each DAPI-positive nuclear pixel. Each symbol (square, circle, triangle) represents the averaged median intensities from all embryos in one experimental repeat. **A',B')** Phosphorylated signaling effector intensity in each GFP-positive pixel was divided by the corresponding GFP intensity. Each symbol (square, circle, triangle) represents the averaged median GFP-normalized signal from all embryos in one experimental repeat. **C')** pSmad2 intensity in each DAPI + GFP-positive pixel was divided by the corresponding GFP intensity. Each symbol (square, circle, triangle) represents the averaged median GFP-normalized signal from all embryos in one experimental repeat.

**Supplementary Figure 5: bOpto-2A-Nodal triple immunofluorescence repeats. A-E)** Embryos injected at the one-cell stage with mRNA encoding *bOpto-2A-Nodal* exposed to dark (left panel) or 455 nm light (50 W/m², right panel) starting at early gastrulation (50% epiboly - shield stage) for 30 min. Triple IF staining was used to simultaneously detect activated signaling effectors (pSmad1, ppERK, and pSmad2 reflecting BMP, FGF, and Nodal signaling, respectively). Only ppERK and pSmad2 shown here. Three bOpto-2A-Nodal-expressing embryos per condition from five experimental repeats are shown. (Scale bar is 200 $\mu$m).

**Supplementary Figure 6: Co-injected GFP signal used for quantification of on / off kinetics. A-C)** Embryos were injected at the one-cell stage with mRNA encoding *GFP* and either *bOpto-FGF* (A), *bOpto-BMP* (B), or *bOpto-2A-Nodal* (C). Starting at early gastrulation (50% epiboly - shield), embryos were exposed to 455 nm light (50 W/m$^2$) for 30 min and fixed at different time points during and after exposure. HCR-IF staining used to detect activated signaling effectors is shown in Fig. 5 (ppERK, pSmad1, and pSmad2 reflect FGF, BMP, and Nodal signaling, respectively). HCR-IF signal was normalized voxel-wise based on co-injected GFP intensity shown here (see Supp. Fig. 7). GFP images correspond to representative IF images shown in Fig. 5. (Scale bar is 200 $\mu$m).

**Supplementary Figure 7: Quantification of optogenetic signaling activator toolkit on/off kinetics.** Embryos were injected at the one-cell stage with mRNA encoding *GFP* and either *bOpto-FGF* (A,A'), *bOpto-BMP* (B,B'), or *bOpto-2A-Nodal* (C,C'). Starting at early gastrulation (50% epiboly - shield), embryos were exposed to 455 nm light (50 W/m$^2$) for 30 min and fixed during and after exposure. Three full repeats were performed across 3-5 trials. HCR-IF staining used to detect phosphorylated signaling effectors is shown in Fig. 5 (ppERK, pSmad1, and pSmad2 reflect FGF, BMP, and Nodal signaling, respectively). **A,B,C)** Raw phosphorylated signaling effector intensity was measured in each DAPI-positive nuclear pixel. Each dot represents the median nuclear pixel intensity in one embryo. Black lines represent the mean nuclear intensity of all embryos in the indicated condition. **A',B')** Phosphorylated siganling effector intensity in each GFP-positive pixel was divided by the corresponding GFP intensity. Each dot represents the median GFP-normalized pixel signal in one embryo. Black lines represent the mean GFP-normalized signal of all embryos in the indicated condition. **C')** pSmad2 intensity in each DAPI + GFP-positive pixel was divided by the corresponding GFP intensity. Each dot represents the median GFP-normalized nuclear pixel signal in one embryo. Black lines represent the mean GFP-normalized nuclear signal of all embryos in the indicated condition.

**Supplementary Figure 8: Co-injected GFP signal used for quantification of irradiance dependence HCR-IF. A-C)** Embryos were injected at the one-cell stage with mRNA encoding *GFP* and either *bOpto-FGF* (A), *bOpto-BMP* (B), or *bOpto-2A-Nodal* (C). Starting at early gastrulation (50% epiboly - shield), embryos were exposed to 455 nm light (50 W/m$^2$) at the indicated irradiances for 5 min (bOpto-FGF) or 25 min (bOpto-BMP and -2A-Nodal). HCR-IF staining used to detect phosphorylated signaling effectors is shown in Fig. 6 (ppERK, pSmad1, and pSmad2 to reflect FGF, BMP, and Nodal signaling, respectively). HCR-IF signal was normalized based on co-injected GFP intensity shown here; GFP images correspond to representative images shown in Fig. 6. (Scale bar is 200 $\mu$m).

**Supplementary Figure 9: Quantification of light irradiance dependence.** Embryos were injected at the one-cell stage with mRNA encoding *GFP* and either *bOpto-FGF* (A,A'), *bOpto-BMP* (B,B'), or *bOpto-2A-Nodal* (C,C'). At early gastrulation (50% epiboly - shield), embryos were exposed to 455 nm light (50 W/m²) at the indicated irradiance for 5 min (bOpto-FGF) or 25 min (bOpto-BMP and -2A-Nodal). HCR-IF was used to detect phosphorylated signaling effectors (ppERK, pSmad1, and pSmad2 reflect FGF, BMP, and Nodal signaling, respectively). **A,B,C)** Raw phosphorylated signaling effector intensity was measured in each DAPI-positive nuclear pixel. Each dot represents the median nuclear pixel intensity in one embryo. Black lines represent the mean nuclear intensity of all embryos in the indicated condition. **A',B')** Phosphorylated siganling effector intensity in each GFP-positive pixel was divided by the corresponding GFP intensity. Each dot represents the median GFP-normalized pixel signal in one embryo. Black lines represent the mean GFP-normalized signal of all embryos in the indicated condition. **C')** pSmad2 intensity in each DAPI + GFP-positive pixel was divided by the corresponding GFP intensity. Each dot represents the median GFP-normalized nuclear pixel signal in one embryo. Black lines represent the mean GFP-normalized nuclear signal of all embryos in the indicated condition.

**Supplementary Figure 10: Spatially localized FGF signaling activation repeats.** Embryos were injected at the one-cell stage with mRNA encoding the green-to-red photoconvertible fluorescent proteins *nls-Kaede* and *bOpto-FGF*. At early gastrulation embryos were either kept in the dark (**A-C, J-L, S-U**) or illuminated locally with 405 and 445 nm confocal lasers. HCR-IF was used to detect ppERK. Two representative exposed embryos from three repeats shown (**D-I', M-R', V-AA'**, respectively). Dotted white lines outline photoconverted Kaede from images to the left. A-F' are shown in Fig. 7. (Scale bar is 200 *μ*m).

**Supplementary Figure 11: Damage-activated ERK phosphorylation at gastrulation stage. A-E)** Wild type, uninjected zebrafish embryos were dechorionated with pronase at the one-cell stage. At early gastrulation (~50% epiboly), embryos were left undamaged (B,C,D,E) or firmly poked with a metal probe (A,B',C',D',E') and fixed 20 min later. Triple IF + DAPI staining was used to detect nuclei (B, B') and phosphorylated signaling effectors ppERK (C,C'), pSmad1 (D,D') and, pSmad2 (E,E'). Damage visible in the DAPI channel correlates with ERK phosphorylation (arrowheads). (Scale bar is 200 $\mu$m).

**Supplementary Figure 12: Circadian gene responses to light exposure in gasturation stage embryos. A-D)** Uninjected AB wild type zebrafish embryos were reared in the dark. At early gastrulation (shield stage), a subset of embryos were exposed to 495+ nm light (18.51 W/m$^2$) or 455 nm light (50 W/m$^2$) for 30 min, then dark for 20 min. A third set of embryos were maintained in the dark throughout the experiment. HCR-FISH was performed for the indicated circadian-related genes. Each panel represents four experimental repeats for the indicated gene. (Scale bar is 200 $\mu$m). **A'-D')** Mean HCR-FISH intensities from experiments shown in A-D. Each symbol represents mean intensity from one embryo. **A''-D'')** Quantification of experiments shown in A-D. Linear-mixed model-predicted least squared means of HCR-FISH signal +/- SEM (N = 4; LMM - Fixed Effect: Wavelength, Random Effect: Replicate; post hoc Tukey's HSD where * indicates p < 0.05).