Recombination alters the furin cleavage site in novel bat-borne HKU5-CoV-2 coronavirus

Ting-Yu Yeh[a,b,m], Vincent Tsai[a,c,k], Samuel M. Liao[a,d,k], Chia-En Hong[a,e,k], Feng-Yu Kuo[a,f,k], Yen Chun Wang[a,g,k], Michael C. Feehley[a,b,h,k], Patrick J. Feehley[a,b,h,k], Yi-Chen Lai[i,j,k], Gregory P. Contreras[a,b,k]

[a]Auxergen Inc., Rita Rossi Colwell Center, Baltimore, Maryland, MD 21202, USA
[b]Auxergen S.r.l., Tecnopolis Science and Technology Park of the University of Bari, University of Bari, Valenzano BA, Italy
[c]Current address: Taipei Wego Private Senior High School, Beitou District, Taipei city, Taiwan
[d]Current address: Davis Senior High School, Davis, CA, USA
[e]Current address: Henry M. Gunn High School, Palo Alto, CA, USA
[f]Current address: Taipei Municipal Chien Kuo High School, Taipei city, Taiwan
[g]Current address: Taipei Municipal Chenggong High School, Taipei city, Taiwan
[h]Current address: Department of Biophysics, Johns Hopkins University, Baltimore, MD 21218, USA
[i]Department of Radiology, Taipei Veterans General Hospital, Taipei city, Taiwan
[j] School of Medicine, National Yang Ming Chiao Tung University, Taipei City, Taiwan
[k] These authors contributed equally
[m]Corresponding author

Corresponding author: Ting-Yu Yeh, MD
yehty@auxergen.com

Abstract

HKU5-CoV-2 is a new bat-infecting coronavirus phylogenetically related to MERS-CoV. It has recently been confirmed that HKU5-CoV-2 can enter human cells and organoids *in vitro* via the ACE2 receptor, raising concerns about its pandemic potential if zoonotic spillover leads to sustained transmission in human populations. Whether recombination influenced HKU5-CoV-2 is completely unclear to date. Here we report the first evidence of HKU5-CoV-2 viral recombination, in association with the mutations at the S1/S2 furin cleavage site (FCS). Using linkage disequilibrium and haploblock analysis, we identify that recombination at the receptor binding domain (RBD) and FCS of the viral spike protein. SNP23016/23043/23064/23156/23285 at the RBD and SNP23833/23847 at the FCS are recombinant hotspots. Our results suggest recombination may lead to Ser723 deletion/insertion and Ser729Ala substitution, both of which could change the furin cleavage activity and host tropism of HKU5-CoV-2.

Introduction

HKU5-CoV-2 merbecovirus is a newly discovered coronavirus from *Pipistrellus* spp. bats in China. Phylogenetically related to MERS-CoV, this virus can enter host cells by using the ACE2 receptor present in many birds and mammals, including humans. [1] Although there is no evidence of human transmission to date, this virus still poses a potential risk of zoonotic spillover and pandemics. It is well known that recombination quickly shapes pathogenesis and evolution of coronaviruses like SARS-CoV-2, even within 3 weeks of human transmission.[2] However, whether recombination has influenced HKU5-CoV-2 is completely unknown. Here we report the first evidence that recombination drove HKU5-CoV-2 diversity and led to alteration in the S1/S2 furin cleavage site (FCS).

Results and Discussion

Six HKU5-CoV-2 sequences (153, 155, 023, 028, 381, 441) are available in GenBase (C-AA08189 to C-AA08194).[1] Linkage disequilibrium (LD) analysis of 765 single nucleotide polymorphisms (SNPs) detected 3896 recombinant SNP pairs and 27 haploblocks. Recombination events have occurred in the receptor binding domain (RBD) of the spike protein between SNP23001 and SNP23016, which encodes Tyr448 that contacts the ACE2 receptor.[1] SNP23016, SNP23043, SNP23064, SNP23156, and SNP23285 in the RBD are also hotspots for intradomain or long-range recombination (Figure 1A-1C).

It has been suggested that the FCS appeared independently multiple times in the evolution of the coronavirus spike protein.[3] We find that the HKU5-CoV-2 FCS contains several mutations (Figure 1D). Haploblock analysis revealed that SNP23830 and SNP23833 are recombinant breakpoints at the FCS (haploblock 17/18, Figure 1E). Moreover, SNP23833 and SNP23847 also form 9 and 41 recombinant SNP pairs, respectively, indicating that the FCS has been a crossover hotspot (Figure 1C and 1F). Additionally, the FCS is often lost during virus adaptation to cell culture by either deletions or point mutations.[4] Taken together, these results suggest that recombination may cause Ser723 deletion/insertion and Ser729Ala substitution, both of which could change furin cleavage activity (e.g. C_AA085189/190/194 versus C_AA085191/192/193).

Another non-LD algorithm, Recco, identified 10 putative recombination breakpoints in NSP4, NSP13, NSP14, and the transmembrane domain of the spike protein (Table 1). Other breakpoints (SNP18065, 19205, 25521) were also confirmed by LD and haploblocks analysis. It is reasonable to speculate that recombination of HUK5-CoV-2 virus is underestimated. It has been reported that recombination events alter host tropism of MERSr-COVs and BtCov-422 with broader receptor usage.[5] Therefore, the FCS variations caused by recombination and

mutations could significantly affect viral entry and infectivity. Genomic surveillance of HUK5-CoV-2 and its recombinants in wild bat populations could be the key strategy to mitigate zoonotic transmission and future outbreak.

Materials and Methods

Genomic sequences of HKU-CoV-2 (BtHKU5-CoV-2-153, BtHKU5-CoV-2-155, BtHKU5-CoV-2-023, BtHKU5-CoV-2-028, BtHKU5-CoV-2-381, and BtHKU5-CoV-2-441) were obtained from Genbase database (National Genomics Data Center, China, https://ngdc.cncb.ac.cn/genbase/?lang=en). Accession numbers: C_AA085189.1, C_AA085190.1, C_AA085191.1, C_AA085192.1, C_AA085193.1, and C_AA085194.1. The nucleotide and SNP number are based on C_AA085191.1 sequence in this study. The sequences alignment and linkage disequilibrium (LD) analysis using HaploView (https://www.broadinstitute.org/haploview/haploview) are described in detail as previously. [6,7] Only mutations shown in the same allele at more than two sequences, but not the singletons, were considered SNPs and analyzed in this study. D' value calculated by normalizing the degree of nonrandom association between two alleles (D), with its maximum possible value D_{max}. Haplotype blocks with recombination measurement were determined with the solid spine of LD ($D' > 0.8$). [7]

The viral sequences were analyzed by Recco program, which scores the cost of obtaining one of the sequences from the others by mutation and recombination.[8] The recombinant fragment and breakpoints were measured by the parameter α weighting recombination cost against mutation cost. Only the fragment with significant mutation cost (>5) and p value <0.05 are considered as recombinant fragments.

References

1. Chen J, Zhang W, Li Y, et al. Bat-infecting merbecovirus HKU5-CoV lineage 2 can use human ACE2 as a cell entry receptor. Cell 2025; 188(6):1729-1742.e16.

2. Yeh TY, Contreras GP. Viral transmission and evolution dynamics of SARS-CoV-2 in shipboard quarantine. Bull World Health Organ. 2021;99(7):486-495.

3. Carabelli AM, Peacock TP, Thorne LG, et al. SARS-CoV-2 variant biology: immune escape, transmission and fitness. Nat Rev Microbiol 2023;21(3):162-177.

4. Vu MN, Lokugamage KG, Plante JA, et al. QTQTN motif upstream of the furin-cleavage site plays a key role in SARS-CoV-2 infection and pathogenesis. Proc Natl Acad Sci USA. 2022; 119(32), e2205690119

5. Tolentino JE, Lytras S, Ito J, Holmes EC, Sato K. Recombination as an evolutionary driver of MERS-related coronavirus emergence. Lancet Infect Dis. 2024;24(9):e546.

6. Yeh TY, Feehley PJ, Feehley MC, Chen CF, Tsai TY, Cheng HL, Contreras GP. Rapidly evolving recombinant monkeypox virus strains compound the challenges of the 2024 mpox outbreak. Lancet Microbe. 2025;6(2):101017.

7. Barrett JC, Fry B, Maller J, Daly MJ. Haploview: analysis and visualization of LD and haplotype maps. Bioinformatics. 2005; 21:263–265.

8. Maydt J, Lengauer T. Recco: recombination analysis using cost optimization Bioinformatics. 2006;22(9):1064-71.

Declaration of interest

TYY is the CSO of Auxergen Inc and Auxergen s.r.l.. GPC is the CEO of Auxergen Inc and Auxergen s.r.l.. Other authors have no conflict of financial interest.

Funding
No funding

Ethical approval

None declared.

Author Contributions: Yeh and Contreras had full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

Concept and design: Yeh

Acquisition, analysis, or interpretation of data: Yeh, Tsai, Laio, Hong, Kuo, Wang, Feehley MC, Feehley PJ, Lai

Drafting of the manuscript: Yeh, Contreras

Critical revision of the manuscript for important intellectual content: Yeh, Contreras

Statistical analysis: Yeh, Tsai, Laio, Hong, Kuo, Wang, Feehley MC, Feehley PJ, Lai

Supervision: Yeh, Contreras

Figure 1. Recombination of HKU-CoV-2 RBD and FCS. (A) Haplotype block (14 to 17) organization of HKU-CoV-2 RBD sequences. The Hedrick's multi-allelic D' statistics in the crossing areas represents the recombination level between two haploblocks. Haplotype frequencies are shown as the numbers next to each haplotype block, or > 10% (thick lines) or > 1% (thin lines). (B) Alignment of the adjacent sequence of recombinant breakpoint SNP23016 (red). ACE2 receptor binding residues Y448 and S453 are colored in orange. (C) Pairwise LD plots of recombinant SNP pairs (white squares) in RBD, FCS (SNP23847, green), or the intergenic region between spike and E protein (SNP26093). The recombinant breakpoints in haploblock 15, 16, and 17 are colored in red. (D) Alignment of FCS sequences and SNPs locations. Furin cleavage site (purple arrow), sequences with Ser272/273 deletion (yellow), or S729A substitution (green) are shown. (E) Haplotype block organization of (D). (F) HaploView LD map of FCS and other SNPs. Red squares indicate high levels of LD, and white and blue squares represent low levels of LD. Recombinant SNP pairs are marked with asterisks.

Table 1. Recco analysis of HKU-CoV-2 sequences*. The recombination breakpoints are verified by LD (recombinant SNP pairs#) and haploblock analysis (between or at the end of haploblocks).

Recombinant fragment (nucleotide)*	Mutation cost*	SNP (recombinant SNP pair numbers)#	Location	Haploblock
8893-8997	381.2 (p=0.04)	8892 (0)	Polyprotein 1ab, AA 2879-2913, NSP4 transmembrane domain	10
18066-18397	410.2 ($p<10^{-3}$)	18065 (86)	Polyprotein 1ab, AA 5937-6046, NSP13 Helicase, NSP14 ExoN domain	12
19206-19292	410.2 ($p<10^{-3}$)	19205 (9)	Polyprotein 1ab, AA 6317-6345, NSP14, N7-MTase active site	12 (end)
25522-25563	394.2 (p=0.017)	25521 (41)	spike protein AA 1286-1299, transmembrane domain	19, 20