Sex-dimorphic effects of neuromelanin buildup in rodent nigral dopamine neurons: implications for sex-biased vulnerability in Parkinson's disease

Sebastian Luca D'Addario[1,2,3], Mariangela Massaro Cenere[1,3], Francesca Cossa[1,3], Silvia Scaricamazza[1], Valentina Nesci[1], Ginevra D'Ottavio[1,4], Daniele Caprioli[1,4], Alberto Ferri[1,5], Miquel Vila[3,6,7], Ada Ledonne[#1,3,8*], Nicola Biagio Mercuri[#1,3,9]

[1] Fondazione Santa Lucia IRCCS, Department of Experimental Neuroscience, Rome, Italy.
[2] Uninettuno Telematic International University, Faculty of Psychology, Rome, Italy.
[3] Aligning Science Across Parkinson's (ASAP) Collaborative Research Network; Chevy Chase, Maryland USA.
[4] Sapienza University of Rome, Department of Physiology and Pharmacology, Rome, Italy.
[5] Institute of Traslational Pharmacology IFT-CNR, Rome, Italy.
[6] Vall d'Hebron Research Institute (VHIR)-Center for Networked Biomedical Research on Neurodegenerative Diseases (CIBERNED), Autonomous University of Barcelona (UAB), Barcelona, Spain.
[7] Catalan Institution for Research and Advanced Studies (ICREA), Barcelona, Spain.
[8] University of Rome Tor Vergata, Department of Systems Medicine, Pharmacology Unit, Rome, Italy.
[9] University of Rome Tor Vergata, Department of Systems Medicine, Neurology Unit, Rome, Italy.

Equal senior authors

*Corresponding author:

Ada Ledonne, PhD
University of Rome Tor Vergata
Department of System Medicine
Via Montpellier n°1, 00133, Rome, Italy
Email: ada.ledonne@uniroma2.it

Abstract

Neuromelanin (NM) is a dark pigment accumulating with age in human substantia nigra pars compacta (SNpc) dopamine (DA) neurons, conferring the dark look that inspired nigral area's name. Despite NM has long been associated with Parkinson's disease (PD), as melanized neurons favorably degenerate during disease development, NM functions within SNpc DA neurons are still mostly elusive. Here, by exploiting an NM-producing rat model generated by viral vector-induced expression of human Tyrosinase (hTyr), we inspected NM impact on nigral DA neurons' survival and activity, on mitochondrial functionality of SNpc, and behaviors resembling non-motor and motor PD symptoms. Our data reveal sex dimorphism in NM effects on nigrostriatal dopamine circuit, with sex-biased alterations in neuronal firing activity and underlying intrinsic currents, nigral mitochondrial functions, and non-motor PD symptoms (anxiety). In conclusion, this study discloses unrealized NM effects within nigral DA neurons, advancing our comprehension of sex-specific features shaping sex-biased vulnerability to PD.

Keywords: Tyrosinase, substantia nigra, firing activity, non-motor symptoms, motor symptoms, anxiety, sex dimorphism.

Introduction

Parkinson's disease (PD) is characterized by a multifaceted symptomatology including motor anomalies (rigidity, resting tremor, postural instability, and bradykinesia) and a spectrum of non-motor symptoms, counting sleep disturbances, autonomic dysfunctions, cognitive deficits, psychiatric manifestations like depression, anxiety, and apathy associated to prodromal stages, and psychosis and dementia arising at late diseases stages[1,2,3]. Although PD diagnosis relies on appearance of cardinal motor deficits, it is now recognized that prodromal non-motor symptoms can precede them of several years[2]. Thus, disclosure of mechanisms instrumental to both motor and non-motor symptoms is key to comprehending PD etiology and progression.

Increasing evidence indicates that biological sex is an important factor influencing PD development and phenotypic expression[4]. The risk of developing PD is higher in men than in women, with an estimated global prevalence of 1.4:1 for males versus females. Nonetheless, women manifest a faster pathological progression, associated with a higher mortality rate. Sex-related differences also involve symptomatology - with some symptoms being more prevalent in one sex than the other - and response to pharmacological and surgical PD therapy[4,5]. Despite this extensive evidence, the investigation of sex dimorphism in PD animal models remains mostly neglected, failing to provide essential information to understand processes governing sex-related vulnerability in PD patients[4,5].

The degeneration of dopamine (DA) neurons of the substantia nigra pars compacta (SNpc) is the main hallmark of PD. While it is recognized that multiple pathological processes can undermine nigral DA neuron viability (i.e., misfolded α-synuclein (α-syn) aggregation, oxidative stress, mitochondrial dysfunction, impairment of the ubiquitin-proteasome system, and neuroinflammation), the precise cascade of events liable for their degeneration in PD remains unidentified[2]. A key part could be played by neuromelanin (NM)[6], a dark brown cytoplasmatic pigment naturally produced in human catecholaminergic neurons with highest accumulation in SNpc DA neurons. NM appears around the third year of life in human SNpc and progressively accumulates with age[7] conferring the dark appearance that inspired nigral area's nomenclature. NM is a complex polymeric molecule, made of a melanin structure bound with lipids, peptides, and metals[8]; it is synthesized as by-product of DA metabolism, with non-enzymatic DA autoxidation led by ferric ion producing quinones, that by multiple steps of polymerization and aggregation with proteins, lipids, and metals constitute the final NM pigment[9]. Beyond DA autoxidation, NM biosynthesis may rely on an enzymatic pathway lead by Tyrosinase (Tyr) starting from tyrosine oxidation. Tyr is responsible for the production of peripheral melanins in skin and hairs[10], while due to limited protein amounts detected in human SNpc[11,12,13], its contribution to NM production remains still debated.

A causal relationship between NM buildup and PD pathogenesis has been inferred based on earlier evidence that NM-containing neurons preferentially degenerate during PD progression[14] and intracellular inclusions resembling Lewy bodies (LBs) mainly form in proximity to NM granules[15]. Later advances on understanding of NM functions in nigral DA neurons have been limited by the absence

of adequate preclinical models, as NM is not endogenously produced in rodents, opposite to hu-
mans. The recent introduction of first NM-producing rodent models, generated by inducing overex-
pression of human Tyr (hTyr) by adeno-associated viral vector (AAV) or genetic manipulations, pro-
vided direct evidence on NM-dependent histopathological alterations in the nigrostriatal DA circuits,
corroborating the idea that abnormal NM levels contribute to PD development[16,17,18,19]. Nevertheless,
multiple aspects inherent to NM accumulation within nigral DA neurons are still enigmatic: it is un-
known whether (and how) NM affects SNpc DA neurons' activity, if the vulnerability of nigral DA
neurons to NM buildup is sex-biased, and there is sex dimorphism in NM roles in promoting non-
motor and motor PD symptoms. To increase knowledge of these aspects, in this study we generated
a novel NM-producing rat model by inducing AAV-dependent hTyr expression in the SNpc. We pro-
vide original evidence on sex-biased NM roles in shaping nigral DA neurons' functions and fostering
emergence of non-motor PD symptoms, thus gaining insight into sex-specific processes that could
influence sex-biased vulnerability to PD development.

87
88
Results

AAV-induced hTyr expression triggers NM production in nigral DA neurons

To evaluate NM-induced effects on nigral DA neurons, we generated a model of hTyr overexpression
based on bilateral injection of serotype 9 AAV carrying hTyr (AAV-hTyr) above the SNpc of rats of
both sexes **(Fig. 1A).** After 4 weeks, NM production - visible to the naked eye as a dark brown area
entirely covering the SNpc - appeared in AAV-hTyr-injected rats but not in control animals inoculated
with an AAV empty vector (AAV-null) **(Fig. 1B).** Quantification of hTyr by immunofluorescence **(Fig.**
1C) revealed a similar number of hTyr-expressing nigral DA neurons among sexes (**Fig. 1D**). hTyr-
positive DA neurons were also observed in the ventral tegmental area (VTA) (**Extended Data Fig.**
1A). Masson-Fontana staining confirmed NM accumulation at cellular level in SNpc (**Fig. 1E**) and
VTA neurons **(Extended Data Fig. 1B)**. Patchy pigmentation also appeared in substantia nigra pars
reticulata (SNpr), despite confined to cells with typical morphology of DA neurons (**Extended Data**
Fig. 1B). To assess whether there is sex-biased proneness in NM accumulation, we measured NM
levels within SNpc DA neurons in female- and male AAV-hTyr-injected rats, reporting higher intra-
cellular NM levels in males. **(Fig. 1F).** Collectively, these findings demonstrate fast NM production
by AAV9-induced hTyr overexpression in rats, with a slight sex dimorphism in intracellular NM
buildup.

106
107
NM effects on the nigrostriatal dopamine circuit

To disclose whether nigral NM buildup induces early histopathological alterations in the nigrostriatal
DA circuit, first we performed stereological cell counts of SNpc tyrosine hydroxylase positive (TH+)
neurons **(Fig. 2A).** We found a reduction in SNpc TH+ cells in male AAV-hTyr-injected rats compared

to controls **(Fig. 2B)**. To decipher if this relies on TH down-regulation (a typical adaptation of nigral

DA neurons to toxic insults) or real neuronal degeneration, we quantified total DA neurons, that is

TH+ plus TH negative/NM positive (TH- NM+) cells. We found a slight, but not significant, decrease

of total DA neurons, implying that NM at this stage does not cause substantial neuronal death, but

just TH down-regulation **(Fig. 2C)**. Likewise, females showed fewer TH+ cells **(Fig. 2D)**, but unal-

tered total DA neurons **(Fig. 2E)**. Next, we evaluated NM-induced effects on VTA DA neurons, ob-

serving equal number of TH+ neurons and total DA neurons either in male- **(Fig. 2F-G)** and female

(Fig. 2I-J) AAV-hTyr-injected rats and respective controls. The percentage of NM-producing cells

was analogous between SNpc and VTA in both sexes **(Fig. 2H-K)**, further supporting a preferential

vulnerability of SNpc (over VTA) DA neurons to NM buildup.

Moreover, we quantified TH density in the dorsal striatum (dST) in three fields along the rostro-

caudal axes (**Extended Data Fig. 2A-C**) reporting regular levels both in males (**Extended Data Fig.**

2D-F) and females (**Extended Data Fig. 2G-I**). Analyses of TH density in the ventral striatum (i.e.,

nucleus accumbens (NAc) core and NAc shell (dorsal and ventral parts) proved male AAV-hTyr-

injected rats have regular TH levels in NAc core **(Extended Data Fig. 3A)** and ventral NAc shell

(Extended Data Fig. 3C), and a slight, but not statistically significant, reduction in the dorsal NAc

shell **(Extended Data Fig. 3B)**. Female AAV-hTyr-injected rats displayed unaltered TH density in

the analyzed NAc parts **(Extended Data Fig. 3D-F).**

130
131
NM accumulation affects the mitochondrial function in SNpc in a sex-specific manner

Mitochondria are key to proper cellular bioenergetics and their dysfunction contributes to PD devel-

opment[20]. To assess whether NM overload damages mitochondrial functionality, we evaluated mi-

tochondrial respiration in freshly dissected SNpc punches from AAV-hTyr-injected rats and controls

(Fig. 3A). Real-time measurement of oxygen consumption rate (OCR) in male NM-producing rats

(Fig. 3B) revealed unaltered basal respiration and ATP production but increased maximal respiration

and spare capacity **(Fig. 3C)**. Thus, NM buildup in males appears to strain mitochondrial function.

Differently, NM overload did not affect nigral mitochondrial functionality in female AAV-hTyr-injected

rats, whose OCR **(Fig. 3D)**, basal and maximal respiration, ATP production, and spare capacity **(Fig.**

3E) are comparable to the control. These data prove that NM load within SNpc DA neurons promotes

sex-dimorphic nigral mitochondrial adaptations.

143
144
NM buildup induces sex-specific alterations of nigral DA neurons' activity

To reveal the impact of NM buildup on functional features of nigral DA neurons, we performed patch-

clamp recordings in midbrain slices from AAV-hTyr-injected rats and controls 4 weeks post-AAVs

injection. First, we analyzed nigral DA neurons' function in female rats **(Fig. 4A)**. NM production was

visible in medial SNpc within single DA neurons of AAV-hTyr-injected rats, as dark aggregates close

to plasmatic membranes or in the cytosol **(Fig. 4B)**; this allowed precise selection of NM-containing

neurons to compare with control NM-lacking cells. We found altered passive membrane properties in female NM-producing SNpc DA neurons, displaying reduced membrane capacitance (C_m), normal membrane resistance (R_m), and a trend to hyperpolarization (reduced I_{hold} at V_H = -60 mV) **(Fig. 4C)**. Moreover, NM accumulation deeply perturbed spontaneous firing activity, influencing firing pattern and frequency. As known, nigral DA neurons can exhibit different firing patterns (regular pacemaker, irregular spiking, and bursting) or barely absence of firing (silent neurons) due to influence of multi-faceted intrinsic and extrinsic factors. We observed that while in female controls most SNpc DA neurons fired as regular pacemakers and a minority in bursting mode **(Fig. 4D)**, NM-producing DA neurons displayed reduced incidence of pacemaker firing neurons and increase of bursting neurons; few cells were irregularly firing or silent **(Fig. 4D).** Analysis of the coefficient of variation of the inter-spike interval (CV-ISI) confirmed this NM-induced shift in firing patterns, with increased firing irregu-larity **(Fig. 4E)** and higher bursting activity **(Fig. 4F)** in SNpc DA neurons of AAV-Tyr-injected rats. Additionally, the pacemaker population exhibited a trend toward hyperactivity **(Fig. 4G)**.

Next, we verified whether NM produces similar functional alterations in male rats **(Fig. 4H).** Male NM-producing DA neurons displayed regular passive membrane properties (normal C_m and R_m) and a trend to hyperpolarization **(Fig. 4I).** NM buildup did not induce substantial alterations in the spon-taneous firing pattern of nigral DA neurons, as incidence of regular pacemaker, bursting, irregular, and silent neurons was similar among AAV-hTyr-injected rats and controls **(Fig. 4J).** Coherently, CV-ISIs indicated an identical distribution among regular pacemakers or irregular/bursting cells, with a prevalence of regular pacemakers in both groups **(Fig. 4K).** Remarkably, the pacemaker popula-tion displayed a trend toward hypoactivity, oppositely to females **(Fig. 4L).** Collectively, these data prove sex-biased NM roles in shaping nigral DA neurons' activity.

Sex-dimorphic effects of NM buildup on intrinsic currents of SNpc DA neurons

To identify mechanisms underlying NM-induced changes of nigral DA neuron activity, we analyzed two intrinsic currents known to shape firing pattern, pacemaker fidelity, and rate: 1) the hyperpolari-zation-activated current (I_h) mediated by hyperpolarization-activated cyclic nucleotide-gated (HCN) channels, and 2) the afterhyperpolarization-associated current (I_{AHP}), underlying the afterhyperpo-larization phase of the action potential[21]. We found that NM buildup did not affect I_h in female nigral DA neurons, showing regular I_h peak amplitude and density **(Fig. 5A,B)**. Differently, NM blunts I_{AHP} area in female SNpc DA neurons **(Fig. 5B);** such decrease was maintained even after normalization over the C_m **(Fig. 5C).**

Specular investigations in male rats demonstrated that neither I_h **(Fig. 5D)** nor I_{AHP} **(Fig. 5E)** were altered in NM-producing SNpc DA neurons. Thus, NM triggers sex-biased changes in ionic mecha-nisms that govern neuronal activity. Importantly, I_{AHP} reduction can increase spontaneous firing rate and burstiness, thus explaining the NM-induced shifts of nigral DA neuron activity observed exclu-sively in females.

189
190
Nigral NM accumulation induces non-motor PD symptoms in male rats

To evaluate whether NM accumulation within SNpc DA neurons contributes to early insurgence of

non-motor PD symptoms, we conducted a comprehensive behavioral analysis of AAV-hTyr-injected

rats and controls **(Fig. 6A)**. Inspection of anxiety domain in the elevated plus maze (EPM) revealed

that male NM-producing rats spent less time in the open arms (an index of anxiety) **(Fig. 6B)** and

explored less the central area of an open field (OF) (likewise indicative of higher anxiety) compared

to controls **(Fig. 6C).** To assess whether nigral NM buildup also promotes social anxiety, male rats

were tested in the social avoidance test (SAT). While, as expected, control rats exhibited social

preference (that is higher interaction with a social target (ST) over a non-social target (NT)), AAV-

hTyr-injected rats displayed social avoidance, spending less time in the interaction zone (IZ) close

to ST **(Fig. 6D).** Next, as further scrutiny of anxiety domain, we evaluated compulsive behavior in

the marble burying test (MBT), observing a trend, still not significant, to increased number of marbles

buried by AAV-hTyr-injected rats **(Fig. 6E).**

To assess whether NM buildup also affected the motivational domain, influencing goal-oriented be-

haviors, rats were subjected to a two-bottle choice test (TBC) for 3 days and to a sucrose self-

administration protocol for 8 days **(Fig. 6A)**. TBC revealed sucrose preference over water in both

groups **(Fig. 6f, Extended Data Fig. 4A)**, also displaying comparable water consumption and body

weight gain **(Extended Data Fig. 4B,C)**. This suggests that both groups exhibit preserved hedonic

reactions. Outcomes from the self-administration protocol indicated no differences among groups in

instrumental responding (number of lever presses and rewards earned) **(Fig. 6G and Extended**

Data Fig. 4D,E) nor in motivation, assessed using progressive ratio **(Fig. 6H)**. This means that nigral

NM buildup at early stages does not affect the motivational domain.

Additionally, we looked over early appearance of motor anomalies in male NM-producing rats. In the

grip strength test, AAV-hTyr-injected rats showed greater force than controls **(Fig. 6I),** suggestive of

increased rigidity or dysregulation in force's control. Notably, locomotor activity in an OF was regular,

with comparable distance moved in both groups **(Fig.6J)**. Altogether, these data demonstrate that

NM accumulation within nigral DA neurons promotes early anxiety appearance and slight motor al-

terations in male rats.

219
220
Female rats are more resilient to nigral NM-induced behavioral alterations

To disclose whether there is sex-biased vulnerability toward NM-induced behavioral alterations, fe-

males AAV-hTyr-injected rats and controls were exposed to specular tasks of males **(Fig. 7A).** Fe-

male NM-producing rats did not exhibit early signs of anxiety, as proved by equal time spent in EPM

open arms **(Fig. 7B)** or in OF center **(Fig. 7C)** with respect of controls. Likewise, female AAV-hTyr-

injected rats displayed regular social preference **(Fig. 7D)**, and no sign of compulsive behaviors in

the MBT **(Fig. 7E)**, further confirming absence of anxiety. Additionally, female NM-producing rats

exhibited regular sucrose preference **(Fig. 7F, Extended Data Fig. 5A)** and normal performance in sucrose self-administration task, with no differences in lever press, reward delivery **(Fig. 7G, Extended Data Fig. 5B-E),** or progressive ratio test **(Fig. 7H),** thus confirming that nigral NM buildup is influential on goal-oriented behaviors, such as natural rewards seeking. Notably, female NM-producing rats displayed a trend toward increased grip strength **(Fig. 7I)**, along with regular locomotion during OF exploration **(Fig. 7J)** compared with controls. Collectively, these findings indicate female rats are less susceptible to early emergence of non-motor PD symptoms during initial stages of nigral NM buildup.

Sustained NM buildup impairs motor coordination irrespective of sex

To assess whether nigral NM buildup at more advance stages can impair motor function, rats were examined in a rotarod test (measuring motor coordination and learning) 8 weeks after AAV-hTyr injection **(Extended Data Fig. 6A).** Remarkably, prolonged NM overload worsened motor performances on rotarod irrespective of sex **(Extended Data Fig. 6B,C)**. Besides demonstrating that protracted NM accumulation significantly impairs motor function, this evidence suggests that female-biased resilience might be restricted to the initial phases of NM overload, with a later attenuation of sex dimorphism at more advanced stages.

Discussion

This study shows that NM buildup within SNpc DA neurons triggers multifaceted and sex-specific cellular and behavioral alterations. Specifically, NM affects nigral DA neurons' activity, mitochondrial functions in SNpc, and fosters non-motor alterations in a sex-dependent manner. Beyond revealing unrealized effects of NM accumulation within nigral DA neurons, our evidence advances the understanding of sex-specific features underlying sex-biased vulnerability in PD.

NM effects on the nigrostriatal DA circuit

Disclosure of factual NM influence on the nigrostriatal DA system is key to comprehend PD development. While earlier observation that melanized nigral DA neurons preferentially degenerate during PD progression[14] provided initial clues on NM association to PD, latest evidence envisages multi-layered NM effects (protective or detrimental) drawing a more complex picture on NM impact on nigral DA neurons[6,22]. NM at low levels is believed to be neuroprotective, by detoxification of oxidized DA species (an intrinsic process of NM biosynthesis) or metals' chelation, thus blunting ROS levels and metal-mediated oxidations[22,23]. Otherwise, excessive NM amounts can damage nigral DA neurons, by troubling several cellular processes[6]. Indeed, PD-related dysfunctions in NM-producing rodents and non-human primates (NHP) infer NM contribution to PD development[16,17,18,19,24]. Still, various NM-related aspects, including its impact on neuronal activity or existence of sex-specific factors shaping its roles, need to be clarified. Here, we reveal early sex dimorphic NM effects on nigrostriatal DA system by using a hTyr-based model displaying significant NM levels within nigral DA neurons four weeks post AAV-hTyr injection. Biological sex seems to affect NM biosynthetic rate, as intracellular NM levels are higher in male SNpc DA neurons than in females. In contrast, AAV-hTyr-injected NHP seem to display higher NM levels in female SNpc DA neurons[24]. In our model, nigral NM production induces TH down-regulation irrespective of sex, without significant SNpc DA neurons' degeneration. A causal link between NM accumulation and TH down-regulation is also supported by evidence showing that pigmentation is associated with fewer TH+ neurons in rodents, NHP and humans[16,17,18,24,25].

Mitochondrial dysfunction is among the key determinants of PD pathology; impaired mitochondrial respiration in parallel with mitochondrial DNA (mtDNA) deletions has been steadily reported in SNpc in PD patients and models[20,26,27,28,29]. Nevertheless, knowledge about mitochondrial functionality during prodromal PD stages is scarce. We show that NM induces early sex-specific mitochondrial alterations in SNpc, namely increased spare capacity and a trend toward higher maximal respiration in males, but regular functioning in females. This male-specific mitochondrial dysfunction may rely on an adaptation of nigral DA neurons to preserve proper bioenergetic levels to cope with NM burden, as energy-demanding cellular processes (proteasomal and lysosomal degradation) may feasibly trigger to offset NM levels. Notably, similar mitochondrial dysfunctions occur in peripheral blood mon-

onucleate cells (PBMCs)[30], immortalized lymphocytes[31], and fibroblasts from PD patients with idio-
pathic or familiar PD forms[32], indicating commonality amongst peripheral cells and SNpc. Nonethe-
less, additional in-deep analyses during prodromal and late PD phases require to better disclose
mitochondrial alterations developing during PD progression. Assuming that the mitochondrial alter-
ations herein described refer to SNpc DA neurons, input of other nigral cellular types, including as-
trocytes and microglia, cannot be excluded. Although glial cells may be activated by NM released to
the extracellular space from dying DA neurons[33], NM intracellular buildup in initial stages does not
induce neurodegeneration; this narrows the involvement of other cellular types, ascribing mitochon-
drial dysfunctions to nigral DA neurons, which intrinsically produce NM. Of note, intact mitochondrial
function in female neurons, despite significant NM buildup, supports lesser vulnerability compared
to males. NM role in fostering mitochondrial dysfunction is reinforced by evidence that pigmented
DA neurons of human SNpc show the highest levels of mitochondrial DNA deletions (a vulnerability
trait for PD development)[34] and that NM promotes oxidative stress and impair mitochondrial functions
in SH-SY5Y cells, used as proxy of DA neurons[16,35,36].

Sex-biased roles for NM in shaping nigral DA neurons' activity
Adequate comprehension of NM roles in PD pathogenesis cannot overlook the key question on
whether (and how) NM buildup influences nigral DA neuron activity. To directly face this unsolved
point, we analyzed functional features of NM-producing SNpc DA cells, providing the first evidence
of sex-dysmorphic NM impact on nigral DA neuron's function. Specifically, female NM-accumulating
DA neurons display altered passive membrane properties (reduced C_m), a trend to hyperpolarization,
and robust adjustments in the spontaneous firing activity, with a switch in firing patterns from regular
pacemaker to bursting. Differently, male NM-producing DA neurons show slighter functional
changes, without gross adjustments in the firing pattern, but, oppositely to females, a trend toward
hypoactivity.
Regarding NM impact on membrane properties, since C_m depends on membrane size and compo-
sition, lower values in pigmented female neurons could rely on cell shrinkage and/or altered lipidic
bilayer composition or thickness due to NM intercalation. Based on its intrinsic lipidic content, NM
can aggregate with lipidic cellular compartments; indeed, NM closeness to DA neurons' plasma
membrane is evident in AAV-hTyr-injected rats. Future studies may gain insight into sex-specific
features underlying milder NM influence on C_m in male neurons (despite higher NM levels), disclos-
ing whether differences arise from sex dimorphism in membrane composition, favoring NM's mem-
brane anchoring in females.
NM-producing SNpc DA neurons display a trend toward hyperpolarization. While this could appear
at odds with hyperactivity/burstiness of female neurons, a certain level of hyperpolarization needs to
allow bursts' trigger in SNpc DA neurons *ex vivo* slice preparations[37,38]; in this view, mild hyperpolar-

ization stands as a permissive factor to bursts' onset. Importantly, NM load induces sex-biased op-

posite effects on spontaneous firing activity of SNpc DA neurons, namely hyperactivity and higher

burstiness in females and a trend to hypoactivity in males. To unveil mechanisms underlying such

sex-dimorphic NM effects we analyzed two intrinsic currents - I_h and I_{AHP} – known to shape firing

rate, regularity, and pattern of nigral DA neurons[21]. Regardless of sex, NM does not alter I_h in nigral

DA neurons. This aligns with regular I_h in SNpc DA neurons in PD models based on α-syn accumu-

lation[39,40,41,42]. Otherwise, I_{AHP} is selectively reduced in female NM-producing neurons, while it is un-

altered in males. Thus, NM load induces sex-biased I_{AHP} reduction, and this fits with hyperactivity

and burstiness of female NM-loaded neurons. Of note, I_{AHP} decrease associated with hyperactiv-

ity/burstiness also occur in aged α-syn overexpressing rats[41], thus denoting shared alterations within

PD models. The exact mechanism by which NM affects I_{AHP} remains to be specified; theoretically,

NM may inhibit I_{AHP}-mediating channels, namely small conductance- and big conductance K^+ chan-

nels (SK and BK) either by a direct binding or by indirect interference with channels' regulatory

mechanisms, i.e., regulatory subunits or intracellular Ca^{2+} fluctuations.

Sex-dimorphic effects of NM on PD symptomatology

Which is the behavioral phenotype associated with NM buildup within SNpc DA neurons? Do the

sex-dimorphic adaptations of nigral DA neurons lie behind sex-biased vulnerability to develop PD

symptoms? As known, PD is characterized by a prolonged prodromal period with a high prevalence

of neuropsychiatric symptoms, like anxiety, social phobia, depression, and apathy[2]. We show sex-

specific vulnerability to develop non-motor PD symptoms during early NM buildup. Increased anxiety

is exclusively evidenced in male NM-producing rats. Early appearance of anxiety has been seen in

other PD models characterized by blunted DA levels in SNpc projections areas[43,44,45,46] and structural

and functional alterations of SNpc correlate with anxiety in traumatic brain injury[47]. Coherently with

females' resilience to NM-induced anxiety, female *Pink1* KO rats (a genetic PD model) do not display

anxiety before ageing[48]. A link between NM and anxiety is further suggested by evidence that nigral

load of iron (that is primarily bound to NM) correlates with anxiety in PD patients[49].

Social anxiety is another common prodromal PD symptom. Nigral NM overload causes social avoid-

ance selectively in males, while females exhibit regular social preference. Such sex-dimorphic social

avoidance reflects clinical PD phenotypes, with higher prevalence of social phobia in males[50]. Other

evidence advises SNpc DA nucleus' dysregulation can be instrumental to social deficits, as fewer

nigral DA neurons have been reported in a model of autism spectrum disorder[51], and social avoid-

ance has been correlated with nigral mitochondrial dysfunctions in male mice[52].

Apathy, anhedonia, and depression can develop concurrently with anxiety in PD patients. Irrespec-

tive of sex, NM-producing rats display normal hedonic response and motivation for natural rewards,

as proved by regular scores in the TBC and sucrose self-administration protocols. Previous evidence

suggests that moderate loss of SNpc DA neurons need to impair goal-oriented behaviors, as reduced

sucrose self-administration has been reported in PD models with partial lesions of SNpc DA neurons[53,54], while sucrose preference is preserved in a α-syn-based PD model displaying slight SNpc DA neurons' loss[55]. Coherently, initial NM buildup, which does not induce nigral DA neurons' degeneration, does not affect motivated behaviors.

Nigral NM overload induced motor anomalies overt as increased grip strength of NM-producing rats of both sexes. This aligns with other PD rodent models with unilateral lesions[56,57,58] and with PD patients, exhibiting higher grip force in early pathological stages[59,60,61]. Abnormal grip force can rely on inability to release an applied force due to dysregulation of basal ganglia circuitry, thus promoting rigidity, a cardinal PD hallmark[62,63]. Notably, nigral NM buildup do not influence basal locomotion at initial stages while impair motor coordination at more advanced phases (~ 8 weeks) irrespective of sex. Thus, the female-specific resilience appears locked to initial phases of NM overload, with a later attenuation of sex dimorphism.

Male sex is the second most relevant risk factor for PD, beyond aging. Aside to a higher prevalence, PD manifests earlier and with greater DA denervation in men. Sex dimorphism is replicated in PD models, as male rats and primates are more susceptible to toxin-induced SNpc DA neurons' degeneration than females[64,65,66]. Sex-specific features dictating males' vulnerability are mostly unknown. Estrogens are commonly considered key factors shaping sex-biased liability to PD, based on their neurotrophic, neuroprotective, and modulatory effects on female nigrostriatal DA circuit[67,68,69,70]. Sex chromosome genes can also underpin sex-biased vulnerability of the nigrostriatal DA circuit during PD development[70,71]. Microarray analyses of single DA neurons from human SNpc sections revealed higher levels of PD-associated genes (α-syn and PINK-1)[72] and lower expression of genes involved in oxidative phosphorylation and synaptic transmission in males in respect with females[73]. Thus, male SNpc DA neurons own a pattern of gene expression that could foster PD development. The Y-chromosome gene (sex-determining region Y, SRY) has been proposed as a male-specific factor contributing to DA system's vulnerability. SRY is expressed within a subset of DA neurons of the ventral midbrain of males[74], wherein influences gene transcription of key enzymes of DA metabolism, such us TH, monoamine oxidase A (MAO_A), and DA decarboxylase[74,75,76]. SRY is upregulated in 6-OHDA- and rotenone-based PD models, whereas its downregulation is sufficient to mitigate SNpc DA neurons' degeneration and motor deficits of male PD rodents[77]. Future investigations require to identify key factors accounting for sex-biased NM roles on SNpc DA neurons fostering PD development.

Conclusions and open issues

This study provides novel insights on NM impact on nigral DA neurons, revealing sex-biased functional adaptations contributing to male-specific susceptibility to develop prodromal PD symptoms during initial NM buildup. Hyperactivity and burstiness of nigral DA neurons emerge as adaptations that females, but not males, are able to trigger during NM buildup. Whether this sex dimorphism

relies on female-specific features (naturally lacking in males) or on mechanisms intrinsic to both sexes but impaired in males due to their higher vulnerability to NM, remains to be determined. Noteworthy, sex-specific changes of SNpc DA neurons' activity associate with sex-biased behaviors of NM-producing rats. Bursting activity of female SNpc DA neurons could ensure sufficient DA levels to allow DA-dependent functions, explaining why females are more resistant to early manifestation of PD symptoms during initial NM buildup. Otherwise, the trend toward hypofunction of male SNpc DA neurons, in addition to TH down-regulation (affecting DA biosynthetic rate), could reduce DA tone in projection areas exposing males to earlier manifestation of PD symptoms.

Chronic NM production induces behavioural alterations also in female rats, revealing an attenuation of sex-dimorphism at advanced stages of NM buildup. Time-dependent acceleration of pathological signs in females during prolonged NM buildup would reflect clinical evidence in PD patients, as females, despite lower initial incidence, display faster disease progression than males. By reproducing male-specific vulnerability observed in PD patients in early stages as well as females' pathological recruitment along disease progression, this NM-producing rat model represents a precious tool to explore PD etiology and to probe sex-tailored innovative therapeutic approaches.

Conclusively, beyond corroborating the relationship between NM and PD, this study advances the understanding of the determinants of sex-biased vulnerability observed in PD patients. Increased recognition of sex differences could aid in the development of customized approaches to better manage PD symptomatology.

Author contributions

Conceptualization: A.L., S.L.D, M.M.C, M.V., and N.B.M.

Methodology: S.L.D, M.M.C, A.L., D.C., A.F., and M.V.

Investigation: S.L.D, M.M.C., A.L., F.C., S.S., V.N., and G.D.

Project administration: A.L.

Supervision: A.L.

Funding acquisition: N.B.M. and M.V.

Visualization: A.L, S.L.D, and M.M.C.

Writing of the manuscript: A.L. and S.L.D.

All authors revised the manuscript and approved the submitted version.

Acknowledgements
This research was funded by Aligning Science Across Parkinson's [ASAP-020505] through the Michael J. Fox Foundation for Parkinson's Research (MJFF). For open access, the author has applied a CC BY public copyright license to all Author Accepted Manuscripts arising from this submission.

430 The study was also supported by NEXTGENERATIONEU (NGEU) and funded by the Ministry of
431 University and Research (MUR), National Recovery and Resilience Plan (NRRP), project MNESYS
432 (PE0000006) – A Multiscale integrated approach to the study of the nervous system in health and
433 disease (DN. 1553 11.10.2022). Sponsors did not play a role in study design, data collection, anal-
434 ysis and interpretation, or writing of this manuscript.
435
436
Data availability
Source data and key resources used in the current study are available as indicated in the Key Re-
source table posted at the Zenodo repository at https://zenodo.org/records/15528775.
No code was generated for this study; all data cleaning, preprocessing, analysis, and visualization
was performed using OriginPro 2024 (OriginLab), Prism10 (GraphPad), Stereoinvestigator 2019.1.3
(MicroBright-Field), ImageJ (NIH), pClamp 10.3 (Molecular Devices), EthoVision XT17 (Noldus).
Additional request of information should be addressed to the corresponding author Ada Ledonne
(ada.ledonne@uniroma2.it).
445
446
447 **Disclosure of conflict of interest**
448 The authors declare no competing interests.
449

References

References

1. Sveinbjornsdottir, S. The clinical symptoms of Parkinson's disease. *J Neurochem* 139, 318-324
(2016).

2. Poewe, W., Seppi, K., Tanner, C.M., Halliday, G.M., Brundin, P., Volkmann, J., et al. Parkinson
disease. *Nat Rev Dis Primers* 3, 17013 (2017).

3. Hussein, A., Guevara, C.A., Del Valle, P., Gupta, S., Benson D.L., & Huntley G.W. Non-Motor
Symptoms of Parkinson's Disease: The Neurobiology of Early Psychiatric and Cognitive Dysfunc-
tion. *Neuroscientist* 29, 97-116 (2023).

4. Meoni, S., Macerollo, A., & Moro, E. Sex differences in movement disorders. *Nature Reviews*
Neurology 16, 84–96 (2020).

5. Cerri S, Mus L, & Blandini F. Parkinson's Disease in Women and Men: What's the Difference? *J*
Parkinsons Dis 9, 501-515 (2019).

6. Vila, M. Neuromelanin, aging, and neuronal vulnerability in Parkinson's disease. *Mov Disord*, 34,
1440-1451 (2019).

7. Fedorow, H., Halliday, G.M., Rickert, C.H., Gerlach, M., Riederer, P., & Double, K.L. Evidence for
specific phases in the development of human neuromelanin. *Neurobiol Aging* 27, 506-12 (2006).

8. Zucca, F.A., Vanna, R., Cupaioli, F.A., Bellei. C., De Palma, A., Di Silvestre, D., et al. Neuro-
melanin organelles are specialized autolysosomes that accumulate undegraded proteins and lipids
in aging human brain and are likely involved in Parkinson's disease. *NPJ Parkinsons Dis* 4, 17
(2018).

9. Sulzer, D., Bogulavsky, J., Larsen, K.E., Behr. G., Karatekin, E., Kleinman, M.H., et al. Neuro-
melanin Biosynthesis Is Driven by Excess Cytosolic Catecholamines Not Accumulated by Synaptic
Vesicles. *PNAS* 97, 11869–74 (2000).

10. Sánchez-Ferrer, A., Rodríguez-López, J.N., García-Cánovas, F., & García-Carmona, F. Tyrosi-
nase: a comprehensive review of its mechanism. *Biochim Biophys Acta* 1247, 1-11 (1995).

11. Miranda, M., Botti, D., Bonfigli, A., Ventura, T., & Arcadi, A. Tyrosinase-like activity in normal
human substantia nigra. *Gen Pharmacol* 15, 541-4 (1984).

12. Greggio, E., Bergantino, E., Carter, D., Ahmad, R., Costin, G.E., Hearing, V.J., et al. Tyrosinase
exacerbates dopamine toxicity but is not genetically associated with Parkinson's disease. *J Neuro-
chem* 93, 246-56 (2005).

13. Xu, Y., Stokes, A.H., Freeman, W.M., Kumer, S.C., Vogt, B.A., & Vrana, K.E. Tyrosinase mRNA
is expressed in human substantia nigra. *Brain Res Mol Brain Res* 45, 159-62 (1997).

14. Hirsch, E., Graybiel, A.M., & Agid, Y.A. Melanized dopaminergic neurons are differentially sus-
ceptible to degeneration in Parkinson's disease. *Nature* 334, 345-8 (1988).

15. Braak, E., Sandmann-Keil, D., Rüb, U., Gai, W.P., de Vos, R.A., Steur, E.N., et al. alpha-synu-
clein immunopositive Parkinson's disease-related inclusion bodies in lower brain stem nuclei. *Acta*
Neuropathol 101, 195–201 (2001).

16. Carballo-Carbajal, I., Laguna, A., Romero-Giménez, J., Cuadros, T., Bové, J., Martinez-Vicente,
M., et al. Brain tyrosinase overexpression implicates age-dependent neuromelanin production in
Parkinson's disease pathogenesis. *Nat Commun* 10, 973 (2019).

17. Gonzalez-Sepulveda, M., Compte, J., Cuadros, T., Nicolau, A., Guillard-Sirieix, C., Peñuelas, N., et al. M. In vivo reduction of age-dependent neuromelanin accumulation mitigates features of Parkinson's disease. *Brain* 146, 1040-1052 (2023).

18. Laguna, A., Peñuelas, N., Gonzalez-Sepulveda, M., Nicolau, A., Arthaud, S., Guillard-Sirieix, C., et al. Modelling human neuronal catecholaminergic pigmentation in rodents recapitulates age-related neurodegenerative deficits. *Nat Commun* 15, 8819 (2024).

19. Garcia-Gomara, M., Juan-Palencia, A., Alfaro, M., Cuadrado-Tejedor, M., & Garcia-Osta, A. Neuroprotective Effects of Dexamethasone in a Neuromelanin-Driven Parkinson's Disease Model. *J Neuroimmune Pharmacol* 20, 2 (2024).

20. Grünewald, A., Kumar, K.R., & Sue, C.M. New insights into the complex role of mitochondria in Parkinson's disease. *Prog Neurobiol* 177, 73-93 (2019).

21. Gantz, S.C., Ford, C.P., Morikawa, H., & Williams, J.T. The Evolving Understanding of Dopamine Neurons in the Substantia Nigra and Ventral Tegmental Area. *Annu Rev Physiol* 80, 219-241 (2018).

22. Zucca, F.A., Basso, E., Cupaioli, F.A., Ferrari, E., Sulzer, D., Casella, L., et al. Neuromelanin of the human substantia nigra: an update. *Neurotox Res* 25, 13-23 (2014).

23. Segura-Aguilar, J., Paris, I., Muñoz, P., Ferrari, E., Zecca, L., & Zucca, F.A. Protective and toxic roles of dopamine in Parkinson's disease. *J Neurochem* 129, 898-915 (2014).

24. Chocarro, J., Rico, A.J., Ariznabarreta, G., Roda, E., Honrubia, A., Collantes, M., et al. Neuromelanin accumulation drives endogenous synucleinopathy in non-human primates. *Brain* 146, 5000-5014 (2023).

25. McCormack, A.L., Di Monte, D.A., Delfani, K., Irwin, I., DeLanney, L.E., Langston, W.J., et al. Aging of the nigrostriatal system in the squirrel monkey. *J Comp Neurol* 471, 387-95 (2004).

26. Schapira, A.H., Cooper, J.M., Dexter, D., Jenner, P., Clark, J.B., & Marsden, C.D. Mitochondrial complex I deficiency in Parkinson's disease. *Lancet* 1, 1269 (1989).

27. Janetzky, B., Hauck, S., Youdim, M.B., Riederer, P., Jellinger, K., Pantucek, F, et al. Unaltered aconitase activity, but decreased complex I activity in substantia nigra pars compacta of patients with Parkinson's disease. *Neurosci Lett* 169, 126–8 (1994).

28. Bender, A., Krishnan, K.J., Morris, C.M., Taylor, G.A., Reeve, A.K., Perry, R.H., et al. High levels of mito-chondrial DNA deletions in substantia nigra neurons in aging and Parkinson disease. *Nat Genet* 38, 515–7 (2006).

29. Gonzalez-Rodriguez, P., Zampese, E., Stout, K.A., Guzman, J.N., Ilijic, E., Yang, B., et al. Disruption of mitochondrial complex I induces progressive parkinsonism. *Nature* 599, 650–6 (2021).

30. Schirinzi, T., Salvatori, I., Zenuni, H., Grillo, P., Valle, C., Martella, G., et al. Pattern of Mitochondrial Respiration in Peripheral Blood Cells of Patients with Parkinson's Disease. *Int J Mol Sci* 23, 10863 (2022).

31. Annesley, S.J., Lay, S.T., De Piazza, S.W., Sanislav, O. Hammersley, E., Allan, C.Y., et al. Immortalized Parkinson's dis-ease lymphocytes have enhanced mitochondrial respiratory activity. *Dis. Model. Mech* 9, 1295–1305 (2016).

32. Antony, P.M., Kondratyeva, O., Mommaerts, K., Ostaszewski, M., Sokolowska, K., Bau-muratov,
A.S., et al. Fibroblast mi-tochondria in idiopathic Parkinson's disease display morphological changes
and enhanced resistance to depolarization. *Sci. Rep* 10, 1569 (2020).

33. Zecca, L., Wilms, H., Geick, S., Claasen, J-H., Brandenburg, L-O., Holzknecht, C., et al. Human
neuromelanin induces neuroinflammation and neurodegen-eration in the rat substantia nigra: impli-
cations for Parkinson's disease. *Acta Neuropathol* 116, 47-55 (2008).

34. Elstner, M., Müller, S.K., Leidolt, L., Laub, C., Krieg, J., Schlaudraff, F., et al. Neuromelanin,
neurotransmitter status and brain-stem location determine the differential vulnerability of catechola-
minergic neurons to mitochondrial DNA deletions. *Mol Brain* 4, 43 (2011).

35. Shamoto-Nagai, M., Maruyama, W., Yi, H., Akao, Y., Tribl, F., Gerlach, M., et al. Riederer, P.,
Naoi, M. Neuromelanin induces oxidative stress in mitochondria through release of iron: mechanism
behind the inhibition of 26S proteasome. *J Neural Transm (Vienna)* 113, 633-44 (2006).

36. Lawana, V., Um, S.Y., Foguth, R.M., & Cannon, J.R. Neuromelanin formation exacerbates HAA-
induced mi-tochondrial toxicity and mitophagy impairments. *Neurotoxicology* 81, 147-160 (2020).

37. Johnson, S.W., Seutin, V., & North, R.A. Burst firing in dopamine neurons induced by N-methyl-
D-aspartate: role of electrogenic sodium pump. *Science* 258, 665–667 (1992).

38. Seutin, V., Johnson, S.W., & North, R.A. Apamin increases NMDA-induced burst-firing of rat
mesencephalic dopamine neurons. *Brain Res* 630, 341–344 (1993).

39. Subramaniam, M., Althof, D., Gispert, S., Schwenk, J., Auburger, G., Kulik, A., et al. Mutant
alpha-synuclein enhances firing frequencies in dopa-mine substantia nigra neurons by oxidative im-
pairment of A-type potassium channels. *J Neurosci* 34, 13586–13599 (2014).

40. Tozzi, A., Sciaccaluga, M., Loffredo, V., Megaro, A., Ledonne, A., Cardinale, A., et al. Dopamine-
dependent early synaptic and motor dysfunctions induced by α-synuclein in the nigrostriatal circuit.
Brain 144, 3477-3491 (2021).

41. Ledonne, A., Massaro Cenere, M., Paldino, E., D'Angelo, V., D'Addario, S.L., Casadei, N., et al.
Morpho-Functional Changes of Nigral Dopamine Neurons in an α-Synuclein Model of Parkinson's
Disease. *Mov Disord* 38, 256-266 (2023).

42. Guatteo, E., Berretta, N., Monda, V., Ledonne, A., & Mercuri, N.B. Pathophysiological features
of nigral dopaminergic neurons in animal models of Parkinson's disease. *Int J Mol Sci* 23, 4508
(2022).

43. Zhu, X.R., Maskri, L., Herold, C., Bader, V., Stichel, C.C., Güntürkün, O., et al. Non-motor be-
havioural impairments in parkin-deficient mice. *Eur J Neurosci* 26, 1902-11. (2007).

44. Tadaiesky, M.T., Dombrowski, P.A., Figueiredo, C.P., Cargnin-Ferreira, E., Da Cunha, C., &
Takahashi, R.N. Emotional, cognitive and neurochemical alterations in a premotor stage model of
Parkinson's disease. Neuroscience 156, 830-40 (2008). Erratum in: Neuroscience 480, 246-247
(2022).

45. Taylor, T.N., Caudle, W.M., Shepherd, K.R., Noorian, A., Jackson, C.R., Iuvone, P.M., et al.
Nonmotor symptoms of Parkinson's disease revealed in an animal model with reduced monoamine
storage capacity. *J Neurosci* 29, 8103-13 (2009).

46. Vieira, J.C.F., Bassani, T.B., Santiago, R.M., de O Guaita, G., Zanoveli, J.M., da Cunha, C., et
al. Anxiety-like behavior induced by 6-OHDA animal model of Parkinson's disease may be related

to a dysregulation of neurotransmitter systems in brain areas related to anxiety. *Behav Brain Res* 371, 111981 (2019).

47. Gao, L., Xue, Q., Gong, S., Li, G., Tong, W., Fan, M., et al. Structural and functional alterations of sub-stantia nigra and associations with anxiety and depressive symptoms following traumatic brain injury. *Front Neurol* 3, 719778 (2022).

48. Marquis, J.M., Lettenberger, S.E., & Kelm-Nelson, C.A. Early-onset Parkinsonian behaviors in female Pink1-/- rats. Behav Brain Res 377, 112175 (2020).

49. An, H., Zeng, X., Niu, T., Li, G., Yang, J., Zheng, L., et al. Quantifying iron deposition within the substantia nigra of Parkinson's disease by quantitative susceptibility mapping. *J Neurol Sci* 386, 46-52 (2018).

50. Gultekin, B.K., Ozdilek, B., & Bestepe, E.E. Social phobia in Parkinson's disease: Prevalence and risk factors. *Neuropsychiatr Dis Treat* 10, 829-34 (2014).

51. Zarate-Lopez, D., Garzón-Partida, A.P., Gonzalez-Perez, O., & Gálvez-Contreras, A.Y. Sex differences in autism-like behavior and dopaminergic neurons in substantia nigra of juvenile mice prenatally exposed to valproate. *Dev Psychobiol* 66, e22469 (2024).

52. Rosenberg, A.M., Saggar, M., Monzel, A.S., Devine, J., Rogu, P., Limoges, A., et al. Brain mitochondrial diversity and network organization predict anxiety-like behavior in male mice. *Nat Commun* 14, 4726 (2023).

53. Drui, G., Carnicella, S., Carcenac, C., Favier, M., Bertrand, A., Boulet, S., et al. Loss of dopaminergic nigrostriatal neurons accounts for the motivational and affective deficits in Parkinson's disease. *Mol Psychiatry* 19, 358-67 (2014).

54. Carnicella, S., Drui, G., Boulet, S., Carcenac, C., Favie,r M., Duran, T., et al. Implication of dopamine D3 receptor activation in the reversion of Parkinson's disease-related motivational deficits. Transl Psychiatry 4, e401 (2014).

55. Massaro Cenere, M., Tiberi, M., Paldino, E., D'Addario, S.L., Federici, M., Giacomet, C., et al. Systemic inflammation accelerates neurodegeneration in a rat model of Parkinson's disease over-expressing human alpha synuclein. *NPJ Parkinsons Dis* 10, 213 (2024).

56. Dunnett, S.B., Torres, E.M., & Annett, L.E. A lateralised grip strength test to evaluate unilateral nigrostriatal lesions in rats. *Neurosci Lett* 246,1–4 (1998).

57. Jeyasingham, R.A., Baird, A.L., Meldrum, A., & Dunnett, S.B. Differential effects of unilateral striatal and nigrostriatal lesions on grip strength, skilled paw reaching and drug-induced rotation in the rat. *Brain Res Bull* 55, 541-8 (2001).

58. Ma, Y., Zhan, M., OuYang, L., Li, Y., Chen, S., Wu, J., et al. The effects of unilateral 6-OHDA lesion in medial forebrain bundle on the motor, cognitive dysfunctions and vulnerability of different striatal interneuron types in rats. *Behav Brain Res* 266, 37-45 (2014).

59. Fellows, S.J., Noth, J., & Schwarz, M. Precision grip and Parkinson's disease. *Brain* 121, 1771–84 (1998).

60. Wenzelburger, R., Zhang, B.R., Poepping, M., Schrader, B., Muller, D., Kopper, F., et al. Dyskinesias and grip control in Parkinson's disease are normalized by chronic stimulation of the subthalamic nucleus. *Ann Neurol* 52, 240–3 (2002).

61. Pradhan, S., Scherer, R., Matsuoka, Y., & Kelly, V.E. Grip force modulation characteristics as a
marker for clinical disease progression in individuals with Parkinson disease: case-control study.
Phys Ther 95, 369-79 (2015).

62. Prodoehl, J., Corcos, D.M., & Vaillancourt, D.E. Basal Ganglia Mechanisms Underlying Precision
Grip Force Control. *Neurosci Biobehav Rev* 33, 900–908 (2009).

63. Rodriguez-Oroz, M.C., Jahanshahi, M., Krack, P., Litvan, I., Macias, R., Bezard E., et al. Initial
clinical manifestations of Parkinson's disease: features and pathophysiological mechanisms. *Lancet*
Neurol 8, 1128-113 (2009).

64. Murray, H.E., Pillai, A.V., McArthur, S.R., Razvi, N., Datla, K,P,, Dexter, D,T., et al. Dose- and
sex-dependent effects of the neurotoxin 6-hydroxydopamine on the nigrostriatal dopaminergic path-
way of adult rats: differential actions of estrogen in males and females. *Neuroscience* 116, 213-22
(2003).

65. Leranth, C., Roth, R.H., Elsworth, J.D., Naftolin, F., Horvath, T.L., & Redmond, D.E. Jr. Estrogen
is essential for maintaining nigrostriatal dopamine neurons in primates: implications for Parkinson's
disease and memory. *J Neurosci* 20, 8604-9 (2000).

66. De Miranda, B.R., Fazzari, M., Rocha, E.M., Castro, S., & Greenamyre, J.T. Sex Differences in
Rotenone Sensitivity Reflect the Male-to-Female Ratio in Human Parkinson's Disease Incidence.
Toxicol Sci 170, 133-143 (2019).

67. Sawada, H., & Shimohama, S. Neuroprotective effects of estradiol in mesencephalic dopamin-
ergic neurons. *Neurosci Biobehav Rev* 24, 143-7 (2000).

68. Kishi, Y., Takahashi, J., Koyanagi, M., Morizane, A., Okamoto, Y., Horiguchi, S., et al. Estrogen
promotes differentiation and survival of dopaminergic neurons derived from human neural stem cells.
J Neurosci Res 79, 279-86 (2005).

69. Bourque, M., Dluzen, D.E., & Di Paolo, T. Neuroprotective actions of sex steroids in Parkinson's
disease. *Front Neuroendocrinol* 30, 142-157 (2009).

70. Jurado-Coronel, J.C., Cabezas, R., Rodríguez, M.F.A., Echeverria, V., García-Segura, L.M., &
Barreto, G.E. Sex differences in Parkinson's disease: Features on clinical symptoms, treatment out-
come, sexual hormones and genetics. *Review Front Neuroendocrinol* 50, 18-30 (2018).

71. McCarthy, M.M., & Arnold, A.P. Reframing sexual differentiation of the brain. *Nat Neurosci* 14,
677-83 (2011).

72. Cantuti-Castelvetri, I., Keller-McGandy, C., Bouzou, B., Asteris, G., Clark, T.W., Frosch, M.P., et
al. Effects of gender on nigral gene expression and parkinson disease. *Neurobiol Dis* 26, 606-14
(2007).

73. Simunovic, F., Yi, M., Wang, Y., Stephens, R., & Sonntag, K.C. Evidence for gender-specific
transcriptional profiles of nigral dopamine neurons in Parkinson disease. *PLoS One* 5, e8856 (2010).

74. Czech, D.P., Lee, J., Sim, H., Parish, C.L., Vilain, E., & Harley, V.R. The human testis-determin-
ing factor SRY localizes in midbrain dopamine neurons and regulates multiple components of cate-
cholamine synthesis and metabolism. *J Neurochem* 122, 260-71 (2012).

75. Wu, J.B., Chen, K., Li, Y., Lau, Y.F.C., & Shih, J.C. Regulation of monoamine oxidase A by the
SRY gene on the Y chromosome. *FASEB J*, 23, 4029-38 (2009).

76. Tao. Q., Fan, X., Li, T., Tang, Y., Yang, D., & Le, W. Gender segregation in gene expression
and vulnerability to oxidative stress induced injury in ventral mesencephalic cultures of dopamine
neurons. *J Neurosci Res* 90, 167-78 (2012).

77. Lee, J., Pinares-Garcia, P., Loke, H., Ham, S., Vilain, E., & Harley, V.R. Sex-specific neuropro-
tection by inhibition of the Y-chromosome gene, SRY, in experimental Parkinson's disease. *Proc*
Natl Acad Sci USA 116, 16577-16582 (2019).

Additional References in Methods

78. D'Addario, S.L., Rosina, E., Massaro Cenere, M., Bagni, C., Mercuri, N.B., & Ledonne, A. ErbB
inhibition rescues nigral dopamine neuron hyperactivity and repetitive behaviors in a mouse model
of fragile X syndrome. *Mol Psychiatry* (2024). Doi: 10.1038/s41380-024-02831-y

79. Ledonne, A., Nobili, A., Latagliata, E.C., Cavallucci, V., Guatteo, E., Puglisi-Allegra, S., et al.
Neuregulin 1 signalling modulates mGluR1 function in mesencephalic dopaminergic neurons. *Mol*
Psychiatry 20, 959-73 (2015).

80. Ledonne, A., & Mercuri, N.B. mGluR1-Dependent Long Term Depression in Rodent Midbrain
Dopamine Neurons Is Regulated by Neuregulin 1/ErbB Signaling. *Front Mol Neurosci* 11:346 (2018).

81. Scaricamazza, S., Salvatori, I., Giacovazzo, G., Loeffler, J.P., Renè, F., Rosina, M., et al. Skel-
etal-Muscle Metabolic Reprogramming in ALS-SOD1G93A Mice Predates Disease Onset and Is A
Promising Therapeutic Target. *iScience* 23, 101087 (2020).

82. Di Segni, M., Andolina, D., D'Addario, S.L., Babicola, L., Ielpo, D., Luchetti, A., et al. Sex-de-
pendent effects of early unstable post-natal environment on response to positive and negative stimuli
in adult mice. *Neuroscience* 413, 1-10 (2019).

83. Costa-Ferro, Z.S.M., Schmidt do Prado-Lima, P.A., Onsten, G.A., Oliveira, G.N., Brito, G.C.,
Ghilardi, I.M., et al. Bone marrow mononuclear cell transplant prevents rat depression and modu-
lates inflammatory and neurogenic molecules. *Prog Neuropsychopharmacol Biol Psychiatry* 113,
110455 (2022).

84. Richardson, N.R., & Roberts, D.C. Progressive ratio schedules in drug self-administration studies
in rats: a method to evaluate reinforcing efficacy. *J Neurosci Methods* 66, 1-11 (1996).

Methods

Animals

All procedures were carried out following the guidelines on the ethical use of animals from the Council Directive of the European Communities (2010/63/EU) and were approved by the Italian Ministry of Health (Authorization N°78-2023). Wild-type rats (Sprague-Dawley background; RRID:MGI:5651135) were bred in our facility and housed in a temperature- (23 ± 1 ∘C) and humidity-controlled environment (45 %–60 % relative humidity), with a 12 h light/dark cycle. Male and female rats were subjected to surgical procedures for virus injection at 2 months of age and used for different experiments after 4-8 weeks.

Stereotaxic virus injection

Rats (250-300 g; 2-month-old) were anesthetized with xylazine (Rompun, 20 mg/ml; 0.5 ml/kg; Bayer) and tiletamine/zolazepam (Zoletil, 100 mg/ml; 0.5 ml/kg; Virbac) and then mounted in a stereotaxic frame (David Kopf Instruments, Tujunga, CA, USA) equipped with a rat adapter. Rats were stereotaxically injected into the SNpc (AP = -4.9 mm, ML=1.8 ± mm, DV = -7 mm) with the viral vectors AAV9-CMV-htyr (AAV-htyr) or AAV9-CMV-null (AAV-null) (2 µl for side, 0.4 µL/min, titer 9.11 X10^{12} gc/mL) produced at the Viral Vector Production Unit of the Autonomous University of Barcelona (UPV-UAB, Spain). After injection, the needle was left in place for an additional 5 min to prevent backflow. Rats were used for different experiments 4-8 weeks post-injections.

Histopathology of nigrostriatal circuit
Brain preparation

Rats were intraperitoneally injected with xylazine (Rompun; 20 mg/ml, 0.5 ml/kg; Bayer) and tiletamine/zolazepam (Zoletil; 100mg/ml, 0.5 ml/kg; Virbac). Rats were transcardially perfused with 1% heparin in 0.1 M sodium phosphate buffer (PB) to remove blood-derived antigens that may interfere with target antigens' detection, followed by 4% paraformaldehyde (PFA) in phosphate buffer (PB; 0.1 M, pH 7.4). Brain samples were postfixed in 4% paraformaldehyde (PFA) in phosphate buffer (PB; 0.1 M, pH 7.4) at 4 °C. To allow deep perfusion throughout the tissue, brain samples were maintained overnight in 4% PFA. Then samples were rinsed three times in PB solution and put in 30% sucrose solution at 4 °C until sinking for cryoprotection. Next, brains were embedded into Tissue-tek O.C.T. (Optimal Cutting Temperature) compound and frozen in isopentane, cooled into liquid nitrogen, and stored at -80 °C. Coronal sections (30 µm) were cut from the anterior part to the posterior part of the brain with a cryostat (Leica) at -20°C. Slices were stored at 4°C in 0.1 M PB containing 0.02% sodium azide before the staining.

Immunofluorescence

To verify the AAV-hTyr transduction at 4 weeks post-AAV injection, nine coronal midbrain sections with an interval of five sections (to cover the entire SNpc) were stained by double immunofluorescence with antibodies against tyrosine hydroxylase (TH) and human tyrosinase (hTyr) as follows. First, sections were incubated with a blocking solution made with 5% donkey serum and 0.3% TritonX-100 (Sigma) in a PB solution to prevent the non-specific binding of antibodies. Then, primary antibodies anti-TH antibody (1:800, Novus Cat# NB 300-108, RRID:AB_350436) and anti-hTyr (1:250, Thermo Fisher Scientific Cat# MA5-14177, RRID:AB_10980000), diluted in blocking solution, were incubated together over-weekend at 4°C and then incubated for 2h at room temperature with an adequate Alexa Fluor 488 (1:200, Thermo Fisher Scientific Cat# A-21206, RRID:AB_2535792) and 555 secondary antibodies (1:200, Thermo Fisher Scientific Cat# A-31570, RRID:AB_2536180).

Nuclei were stained with DAPI (4',6-diamidin-2-fenilindole). The specificity of the immunofluorescence labeling was confirmed by omitting primary antibodies and using normal serum instead (negative control). Confocal laser scanner microscopy (Zeiss, Oberkochen, Germany LSM800) was used to acquire images.

Immunohistochemistry

For stereological cell count in the SNpc, nine slices, with an interval of five sections, and 12 striatal sections per animal were processed for chromogenic immunohistochemistry. First, the endogenous peroxidase was neutralized with a 0.3% H_2O_2 solution in PB. Sections were then incubated with a blocking solution made with 5% normal goat serum and 0.3 % Triton X-100 (Sigma) in a PB solution for one hour. Free-floating sections were incubated with the primary anti-TH antibody (1:800, Novus Cat# NB 300-108, RRID:AB_350436), diluted in PB containing 0.3% Triton X-100 overnight at 4°C. After three rinses, sections were incubated with a biotinylated secondary antibody (Millipore Cat# AP132B, RRID:AB_92488) followed by the incubation with extravidin-peroxidase (1:1000, Sigma-Aldrich Cat# e2886, RRID:AB_2620165), and visualized by Vector Blue substrate Kit (Vector Laboratories Cat# SK-4700, RRID:AB_2314425). Slides were then dehydrated in a series of alcohols (50%, 70%, 95%, 100%), cleared in xylenes, and coverslipped with Entellan mounting medium (Sigma).

Stereological cell count

The three-dimensional optical fractionator stereological probe was used to estimate the number of TH+, Tyr+, and NM+ neurons in SNpc and VTA. We used the Stereo Investigator System (Micro-Bright-Field, Vermont, USA). An optical microscope (Axio Imager M2, Zeiss, Oberkochen, Germany) equipped with a motorized stage, a MAC 6000 controller (Ludl Electronic Products, Ltd) and a camera are connected to software Stereoinvestigator 2019.1.3 (RRID:SCR_018948; https://www.mbfbioscience.com/stereo-investigator). The region of interest was defined by TH staining and according to the rat brain Paxinos and Franklin's atlas and outlined with a 5x objective[41]. The three-dimensional optical fractionator stereological probe (x, y, z dimension of 50×50×30 μm, with a guard zone of 5 μm along the z axis) was used. Cells were marked with a 100X oil objective.

Masson-Fontana

Midbrain slices were fixed in 4% PFA in 0.1 M PB overnight at 4 °C and then immersed in 30% sucrose solution. Slices were cut into 5 μm-thick sections by a freezing microtome, and NM granules were stained by the Masson-Fontana Staining Kit (DiaPath #010228). This procedure is based on the ability of NM to chelate metals by reducing silver nitrate to a visible metallic state. Staining was performed by incubating the sections with ammoniac solution for 40 min at 56ºC, followed by sodium thiosulphate for 2 min and a final counterstain with Kernechtrot for 7 min. Between each step, samples were rinsed in distilled water. Slides were then dehydrated in a series of alcohols (50%, 70%, 95%, 100%), cleared in xylenes, and coverslipped with Entellan mounting medium (Sigma).

Intracellular NM quantification

30 μm-thick unstained sections representative of the SNpc from different animals were used for intracellular NM quantification. The visualization of unstained NM brown pigment identified melanized catecholaminergic neurons. Bright-field pictures from unstained sections were taken with an optical microscope (Axio Imager M2, Zeiss, Oberkochen, Germany) coupled to a camera. All SNpc NM+ neurons were analysed by means of optical densitometry using ImageJ software (NIH, USA; RRID:SCR_003070; https://imagej.net/) to quantify the intracellular optical density (OD) of NM,

855 as previously reported[18]. Each cell was manually outlined excluding the nucleus; the integrated den-
856 sity value per cell was measured and corrected for non-specific background by subtracting values
857 obtained from the neuropil in the same images. All quantifications were performed in blind for the
858 different experimental groups.
859

Tyrosine hydroxylase fibers' quantification
Sections were photographed with a light microscope (Axio Imager M2, Zeiss, Oberkochen, Ger-
many). Densitometric analysis (OD) of dorsal striatum and nucleus accumbens (core and shell) TH+
fibers were analyzed using the Java image processing and plugin analysis program in ImageJ (NIH,
USA; RRID:SCR_003070; https://imagej.net/). Densitometry values were corrected for non-specific
background staining by subtracting densitometric values measured from the nearby cortex. The area
distinction was manually outlined according to the rat brain Paxinos and Franklin's atlas. All quanti-
fications were performed blindly for the experimental groups.
868
869

Analyses of mitochondrial function
Tissue punches
Rats were anesthetized with isoflurane, decapitated, and the brain was dissected as previously re-
ported[79]. The brain was placed in ice-cold oxygenated artificial cerebrospinal fluid (aCSF: 120 mM
NaCl, 3.5 mM KCl, 1.3 $CaCl_2$, 1 mM $MgCl_2$, 0.4 mM H_2PO_4, 5 mM HEPES, 10 mM D-glucose, pH
7.4), saturated with 95% O_2 and 5% CO_2 for 30 minutes. Coronal brain sections (180 µm thickness)
were obtained with a vibratome (VT1200S, Leica, Nussloch, Germany) in ice-cold aCSF under con-
tinuous oxygenation. A reusable biopsy punch (0.35 mm diameter) (World Precision Instrument, FL,
USA) was gently pressed on the brain slices to collect punches including the SNpc.
879

Seahorse XFe96 analysis
The punches of SNpc were carefully placed in the bottom center of a well of XFe96 Cell Culture
Microplate (Agilent Technologies, CA; RRID:SCR_019545; https://www.agilent.com/) filled with 180
µl aCSF with 1 mM sodium pyruvate solution (Gibco™,Grand Island, NY and Scotland, UK). Next,
the XFe96 Cell Culture Microplate was placed in a non-CO_2 incubator at 37°C for 30 minutes. For
the calibration, each injection port (A, B, and C) of the XFe96 cartridge was differently loaded with:
oligomycin 5 µM (Sigma Aldrich, USA), FCCP (2-[2-[4- (trifluoromethoxy) phenyl] hydrazinylidene]-
propanedinitrile) 1 µM (Sigma Aldrich, USA) + Pyruvate 0.75 µM (Gibco™,Grand Island, NY and
Scotland, UK), or rotenone 1 µM (Sigma Aldrich USA) /antimycin A 1 µM (Sigma Aldrich, USA). As
soon as the calibration was completed, the plate was extracted and the XFe96 Cell Culture Micro-
plate containing tissue punches was inserted to start the assay of mitochondrial functionality. The
test consisted of 4 cycles of measurements (each consisting of: 2 minutes of mix, 1 minute of wait,
and 3 minutes of measurement) to assess the oxygen consumption rate (OCR) before (baseline)
and after injection of different drugs.
894

Electrophysiology
Slice preparation
Acute midbrain slices containing the SNpc were used for electrophysiological patch-clamp record-
ings of nigral DA neurons of AAV-null and AAV-hTyr-injected rats. Midbrain slices were obtained
following published procedures[41] with minor changes. Briefly, rats were anesthetized with isoflurane
and decapitated. The brain was rapidly removed from the skull, and a tissue block containing the
midbrain was immersed in a low-sodium N-methyl-D-glucamine (NMDG)-based artificial cerebrospi-
nal fluid (aCSF) at 2-4 °C. The NMDG-based solution contained (in mM): 92 NMDG, 2.5 KCl, 1.25

NaH$_2$PO$_4$, 30 NaHCO$_3$, 20 HEPES, 25 glucose, 2 thiourea, 5 Na-ascorbate, 3 Na-pyruvate, 0.5
CaCl$_2$, and 10 MgSO$_4$, saturated with 95% O$_2$–5% CO$_2$ (pH 7.3). Horizontal slices (250 µm) were cut
using a vibratome (Leica VT1200S, Leica Microsystems, Wetzlar, Germany). Slices were maintained
in NMDG-based aCSF at 33.0 ± 0.5°C for 15 min before adding increasing volumes of a sodium-
spike solution (2M NaCl in NMDG-based aCSF) every 5 min for 40 min, then were transferred in a
HEPES-based aCSF for long-term storage, containing (in mM): 92 NaCl, 2.5 KCl, 1.25 NaH$_2$PO$_4$, 30
NaHCO$_3$, 20 HEPES, 25 glucose, 2 thiourea, 5 Na-ascorbate, 3 Na-pyruvate, 2 CaCl$_2$, and 2 MgSO$_4$
(pH 7.3). After 1h of recovery, slices were transferred in the recording chamber and perfused at 2.5–
3.0 mL/min with aCSF (33.0 ± 0.5°C) containing (in mM): NaCl 126, NaHCO$_3$ 24, glucose 10, KCl
2.5, CaCl$_2$ 2.4, NaH$_2$PO$_4$ 1.2 and MgCl$_2$ 1.2 (95% O$_2$–5% CO$_2$, pH 7.3).
913

Electrophysiological recordings
Patch-clamp recordings from SNpc DA neurons were performed with setups equipped with upright
microscopes (Nikon Eclipse FN1, RRID:SCR_014995; BX51WI Olympus, RRID:SCR_015801) and
infrared video-cameras (CoolSNAP Photometrics, and Evolve Photometrics, Arizona). Identification
of SNpc DA neurons was done based on their morphology (fusiform, tightly packed, and medium to
large-sized cell bodies (diameter ≥ 20 µm)), localization (in the medial SNpc, close to the medial
terminal nucleus of the accessory optic tract, MT), and their typical electrophysiological features, as
slow spontaneous firing (1–8 Hz) in cell-attached configuration, large action potential (> 2ms) with a
prominent afterhyperpolarization, or the presence of a prominent hyperpolarization-activated inward
current (I$_h$) in response to hyperpolarizing voltage steps[41,79,80].
Cell-attached and whole-cell patch-clamp recordings were performed with glass borosilicate pipettes
(4–7 MΩ) pulled with a PP-83 Narishige puller and filled with a solution containing (in mM): 125 K-
gluconate, 10 KCl, 10 HEPES, 2 MgCl$_2$, 4 ATP-Mg$_2$, 0.3 GTP-Na$_3$, 0.75 EGTA, 0.1 CaCl$_2$, 10 Phos-
phocreatine-Na$_2$ (pH 7.3 with 2 M KOH). Recordings were made with a Multiclamp 700B amplifier
(Molecular Devices, USA; RRID:SCR_018455) using Clampex 10.3 software (pClamp, Molecular
Devices, USA; RRID:SCR_011323, http://www.moleculardevices.com/products/soft-
ware/pclamp.html) and the A/D converter Digidata 1440A (Molecular Devices, USA;
RRID:SCR_021038) connected to a computer.
The spontaneous firing activity was recorded in cell-attached configuration. The mean firing fre-
quency and coefficient of variation of the interspike intervals (CV-ISIs) were measured in at least 2-
3 min of recordings. Membrane resistance (R$_m$) and capacitance (C$_m$) were measured with the "mem-
brane test" protocol (Clampex 10.3) consisting of a 5 mV hyperpolarizing step from -60 mV, at
33.3 Hz by averaging 100 consecutive measurements to calculate final values. R$_m$ and C$_m$ were
measured within 1 min after membrane rupture. Membrane access resistance was repeatedly mon-
itored and recordings in which it exceeded 25 MΩ were discarded. No liquid junction potential cor-
rection was applied.
I$_h$ was elicited by a voltage-clamp protocol of four hyperpolarizing voltage steps (-60/-120 mV, 20
mV increment, for 1 sec, V$_H$= -60 mV) applied to SNpc DA neurons. I$_h$ amplitude was the difference
between instantaneous and steady-state current at each voltage step. The afterhyperpolarization-
associated current (I$_{AHP}$) was induced by applying a depolarizing voltage step (from V$_H$ = -60 mV to
0 mV, 100 ms) and measured as I$_{AHP}$ peak and area under the outward current which follows the
depolarization.

Behavior
Behavioral analyses was conducted following the new Research Domain Criteria initiative of National
Institute of Mental Health (NIMH RDoC, https://www.nimh.nih.gov/research/research-funded-by-

 nimh/rdoc/index.shtml). Behavioural tests were conducted between 9:00 am and 16:00 pm after 1
951 hour of room acclimation.

Grip strength

954 The test was conducted using the hanging grid as previously described[81]. The average value ob-
955 tained from three trials was normalized on animal weight to determine the final score.

Marble burying test

958 Marble burying test (MBT) was conducted as previously reported[78], with minor modifications. Briefly,
959 rats were placed individually in a novel cage (43 cm X 26 cm X 20 cm) containing 5 cm of regular
960 bedding and fifteen marbles (15 mm diameter) and left undisturbed for 20 min. The number of mar-
961 bles buried was estimated by considering buried a marble covered ≥ 2/3 with bedding. Marbles were
962 cleaned with 50% ethanol before each session.

Elevated plus maze

Elevated plus maze (EPM) was carried out as previously described[82]. The apparatus consisted of 2
closed arms and 2 open arms (each 51 cm × 11 cm). The percentage of entries in the open arms
(open entries / open + closed × 100) and the percentage of time spent in the open arms (time in
open / open + closed × 100) during 5 min of exploration were collected and analysed by the auto-
mated video-tracking system EthoVision XT17 (Noldus, The Netherlands; RRID:SCR_000441;
https://www.noldus.com/ethovision).

Social avoidance test

The social avoidance test (SAT) was conducted following published procedures[83], with minor
changes. A rat was placed in a circular open field arena (88 cm diameter) containing a protective
cage (13 cm × 13 cm) placed on a side of the arena, defined as interaction zone (IZ). The test
consists of two sessions of exploration of 5 min each. The first session with an empty cage (no target,
NT) and the second session with a social target (ST) represented by a novel rat matched for age
and sex. The total time spent in the IZ was recorded and calculated by the fully automated video
tracking system EthoVision XT17 (Noldus, The Netherlands; RRID:SCR_000441; https://www.nol-
dus.com/ethovision).

Open field

982 Locomotor activity was assessed in an open field (OF) arena (diameter 100 cm), by evaluating total
983 distance moved and time spent in the arena center during 30 min of exploration.

Sucrose self-administration

Self-administration apparatus: Rats were trained in self-administration chambers located inside
sound-attenuating cubicles, fitted with an electric fan and controlled by a custom-made system. Each
self-administration chamber was equipped with a stainless-steel grid floor and two operant panels
placed on the left and right walls. The right panel was equipped with the palatable solution-paired
active (retractable) lever. Responses on this lever activated the infusion pump and the three-light
cue located above the lever. The palatable solution was delivered to a receptacle located near the
solution-paired lever, connected with a silicon tubing to a syringe that contained the palatable solu-
tion (see **Fig. 6a**).

Two-bottle choice: The training procedure was as previously described[53]. Two-bottle choice (TBC)
started 3 days before self-administration procedures. The first day rats were individually housed with
two bottles containing water (habituation). After habituation, one bottle was replaced with another

one containing a sucrose solution at 2.5%. The intake of each animal was measured for two con-
secutive days and then normalized for the body weight.

Self-administration training: Rats were trained to self-administer the sucrose solution 1 hour/day for
8 days (first and second day were trained for 3h) at fixed ratio 1 (FR1) with no timeout[53]. The training
sessions started with the illumination of the house light and the insertion of the sucrose solution-
paired lever; responses on this lever resulted in the delivery of the sucrose solution (0.2 mL for 4 s)
paired with the illumination of the three-light cue (for 20 s).

Progressive ratio test: After the self-administration training, rats were tested for progressive ratio
(PR), an index of motivation. In this procedure, the number of lever presses (ratio value) required for
drug delivery was incremented within sessions and different days following this progression: 1, 2, 4,
6, 9, 12, 15, 20, 25, 32, 40, 50, 62, 77, 95, 118, 145, 178, 219, 268, 328, 402, 492, 603, etc.[84]. The
completion of each ratio requirement resulted in the delivery of the sucrose solution (0.2 mL). Ses-
sions lasted 3h; then, the levers were retracted. Rats were excluded when 1 hour passed from the
last reward earned.
1011

Rotarod test
The test was performed as previously described[55] with minor modifications. Animals were accus-
tomed to being placed on the rod rotating at low speed (4 rpm) for at least 30 seconds before the
first session of the test. The test consisted of three sessions performed on three consecutive days;
each session consisted of three trials with an accelerating rod from 4 to 40 rpm in 300 s, and at least
5 min of rest between trials. The time in which rats fell from the rod was recorded for each trial and
the average of the three daily trials was used as final value.

1019

Statistical analysis
Statistical analyses were conducted using OriginPro 2024 (OriginLab; RRID:SCR_014212;
http://www.originlab.com/) and Prism10 (GraphPad; RRID:SCR_002798;
http://www.graphpad.com/). The normality of data sets was evaluated with Shapiro-Wilk test and/or
D'Agostino & Pearson test. For normally distributed data, comparisons were performed with two-
tailed Student's t-test (simple or with Welch's correction), repetitive measures two-way ANOVA, as
appropriate. When normality was violated, data were analyzed with Mann-Whitney test, Chi-Square
test, Kruskal-Wallis ANOVA followed by post hoc Dunn's multiple-comparison, Friedman ANOVA
followed by post hoc Wilcoxon signed-rank test (for within paired comparisons) and Kolmogorov–
Smirnov test, or Mann-Whitney test with Holm-Sidak multiple-comparison method (for between un-
paired comparisons).

Figures and figure legends

Figure 1. NM production within nigral DA neurons by local AAV-induced human tyrosinase (hTyr) expression

A, Scheme of the experimental design and timeline used to induce *in vivo* NM production within nigral DA neurons of female and male rats. **B,** Representative unstained rat brains mounted on a cryostat, showing that a dark brown area entirely covering the SNpc (black arrow) can be macro-scopically detected in AAV-hTyr-injected rats, but not in control AAV-null animals, four weeks post-AAVs injection. **C,** Representative double-labelled confocal images of coronal SNpc sections im-munostained for TH (green) and hTyr (red) (scale bar: 100 µm). **D,** Quantification of hTyr-expressing neurons in male and female AAV-hTyr-injected rats, measured as total number of TH+hTyr+ neurons (left; male *n*=7; female *n*= 7; Student's *t*-test, P = 0.517) and as percentage of hTyr+ neurons over total TH+ neurons (right; male *n* = 7; female *n* = 7; Student's *t*-test, P = 0.211). **E,** Representative micrographs of Masson-Fontana staining (showing NM as dark brown granules) in 5-µm-thick SNpc sections from an AAV-hTyr-injected rat (scale bar: 200 µm, 20 µm, and 10 µm). **F,** Representative images of NM-containing neurons from unstained SNpc sections (left; scale bar: 10 µm), and quan-tification of intracellular NM optical density in male and female AAV-hTyr-injected rats (right; Mann-

1050 Whitney test; P = 0.044). Data are expressed as mean ± SEM. **(D, F)**; dots represent individual
1051 animals **(D)** or individual cells **(F)**. *P < 0.05.

1052

1053 **Figure 2. Nigral NM buildup downregulates tyrosine hydroxylase in SNpc DA neurons inde-**
1054 **pendently from sex.**

A, Qualitative mouse brain atlas showing selected SNpc serial sections for quantitative histological analysis. **B,** Representative SNpc images (left, scale bar: 200 μm), and quantification (right) of TH+ neurons in male rats (AAV-null $n = 6$; AAV-hTyr $n = 7$; Student's t-test, P = 0.018). **C,** TH-immunostained SNpc showing TH−NM+ (black arrowhead) and TH+NM+ (white arrowhead) neurons (left, scale bar: 10μm), and histogram (right) of total DA neurons in SNpc of male rats (AAV-null $n = 6$; AAV-hTyr $n = 7$; Mann-Whitney test, P = 0.073). **D,** Representative SNpc images (left, scale bar: 200 μm), and quantification (right) of TH+ neurons in female rats (AAV-null $n = 6$; AAV-hTyr $n = 7$; Student's t-test with Welch's correction, P = 0.020). **E,** TH-immunostained SNpc showing TH−NM+ (black arrowhead) and TH+NM+ (white arrowhead) neurons (left, scale bar: 10 μm), and histogram (right) of total SNpc DA neurons of female rats (AAV-null $n = 6$; AAV-hTyr $n = 7$; Student's t-test with Welch's correction, P = 0.135). **F,** Representative VTA images (left, scale bar: 200 μm), and quantification (right) of TH+ neurons in male rats (AAV-null $n = 6$; AAV-hTyr $n = 6$; Student's t-test, P = 0.261). **G,** Count of total VTA DA neurons of male rats (AAV-null $n = 6$; AAV-hTyr $n = 6$; Student's t-test, P = 0.656). **H,** Quantification of total NM-containing neurons (as % of AAV-null TH-immunopositive neurons) in the SNpc and VTA of male AAV-hTyr rats (SNpc $n = 6$; VTA $n = 6$; Student's t-test, P = 0.344). **I,** Representative VTA images (left, scale bar: 200 μm), and quantification (right) of TH+ neurons in female rats (AAV-null $n = 6$; AAV-hTyr $n = 7$; Student's t-test, P = 0.640). **J,** Count of total VTA DA neurons of female rats (AAV-null $n = 6$; AAV-hTyr $n = 7$; Student's t-test, P = 0.948). **K,** Quantification of total NM-containing neurons (as % of AAV-null TH-immunopositive neurons) in the SNpc and VTA of female AAV-hTyr rats (SNpc $n = 6$; VTA $n = 6$; Student's t-test, P = 0.128). Data are expressed as mean ± SEM **(B-K)**; dots represent individual animals. *P < 0.05.

1077

Figure 3 Nigral NM buildup induces sex-specific changes of the mitochondrial functionality in SNpc.

A, Scheme of the protocol used to evaluate the mitochondrial function on SNpc punches obtained by midbrain rat slices. **B,** Metabolic profile of the Mitochondrial Stress Test showing real-time measurements of the oxygen consumption rate (OCR) after the injection of 1) Oligomycin (Oligo), 2) FCCP pyruvate (FCCP+P), and 3) Rotenone/Antimycin (Rot/Ant) in SNpc punches of male animals. **C,** Mitochondrial parameters obtained from Seahorse mitochondrial stress test in male AAV-hTyr-injected rats (n = 5) and AAV-null rats (n = 5): basal respiration (P = 0.494, Student's t-test), ATP production (P = 0.690, Mann-Whitney test), maximal respiration (P = 0.055, Mann-Whitney test), and spare respiratory capacity (P = 0.047, Student's t-test). Data are represented as median ± quartiles. *P < 0.05. **D,** Metabolic profile of Mitochondrial Stress Test showing real-time measurements of the OCR after the injection of Oligo, FCCP+P, and Rot/Ant in SNpc punches of female rats. **E,** Mitochondrial parameters obtained from Seahorse mitochondrial stress test in female AAV-hTyr-injected rats (n = 7) and AAV-null rats (n = 7): basal respiration (P = 0.997, Student's t-test), ATP production (P = 0.766, Student's t-test), maximal respiration (P = 0.620, Student's t-test), spare respiratory capacity (P = 0.527, Student's t-test). Data are expressed as median ± quartiles.

Figure 4. NM induces sex-specific alterations of nigral DA neurons' activity.

A, Experimental design used to evaluate NM effects on SNpc DA neurons' function by *ex vivo* elec-
trophysiology. **B,** Example images of midbrain slices acquired through infrared microscopy showing
NM buildup in SNpc (up, 4X magnification) and within nigral DA neurons of AAV-hTyr-injected rats

(bottom, 40X magnification). Intracellular NM aggregates are indicated by arrows. **C,** Plots of passive
membrane properties of female nigral DA neurons, namely membrane capacitance (C_m) (left), mem-
brane resistance (R_m) (middle), and holding current (I_{hold}) to V_H = -60 mV (right) in AAV-hTyr-injected
rats (n = 47 cells/3 rats) and AAV-null rats (n = 44 cells/3 rats), Mann-Whitney test, C_m: P = 6.22 X
10^{-4}; R_m: P = 0.387; I_{hold}: P = 0.011. **D,** Up, pie diagrams indicating incidence (%) of different firing
patterns (regular pacemaker, irregular, bursting, and silent) of female nigral DA neurons; Bottom,
Example cell-attached firing traces, with indication of relative prevalence (cells in each firing pat-
tern/total cells). **E,** Histogram of coefficient of variation of interspike intervals (CV-ISIs) showing
higher irregularity in female NM-producing DA neurons (AAV-null n = 28 and AAV-hTyr n = 30, Mann-
Whitney test, P = 0.054). **F,** Proportion of regular pacemaker and bursty neurons (χ^2 test, χ^2 (1,
N=56), P=0.041); **G,** Quantification of firing frequencies of female pacemaker neurons showing a
trend to hyperactivity in NM-producing cells (AAV-null n = 23 cells/3 rats and AAV-hTyr n = 16 cells/3
rats; Student's t-test, P = 0.065). **H,** Experimental design for *ex vivo* electrophysiology of male nigral
DA neurons. **I,** Left, Example images of NM-expressing or control SNpc DA neurons during the
patch-clamp procedure; Right, Plots of passive membrane properties (C_m, R_m, and I_{hold} at V_H = -60
mV) of male nigral DA neurons (AAV-null n = 49 cells/4 rats and AAV-hTyr n = 45 cells/4 rats; Mann-
Whitney test, C_m: P = 0.215; R_m: P = 0.591; I_{hold}: P = 0.014). **J,** Up, Pie diagrams showing incidence
(%) of different firing modes; Bottom, Example firing traces and relative prevalence (cells in each
firing mode/total cells). **K,** Plot of CV-ISIs indicating similar temporal spike distribution among male
groups (AAV-null n = 23 cells and AAV-hTyr n = 28 cells, Mann-Whitney test, P = 0.632). **L,** Histo-
gram of firing frequency of male pacemaker neurons showing a trend toward hypoactivity of NM-
producing SNpc DA neurons (AAV-null n = 17 cells/4 rats; AAV-hTyr n = 19 cells/4 rats; Student's t-
test, P = 0.073). Data are expressed as mean ± SEM (**C, F, G, I, L**); dots represent individual neu-
rons. *P < 0.05 and ***P < 0.001.

1124

Figure 5. Sex-specific effects of NM buildup on intrinsic currents of SNpc DA neurons.

A, Example traces of hyperpolarization-activated current (I_h) (left); quantification of I_h amplitude (middle) or I_h density (right) of female nigral DA neurons of AAV-null (n = 46 cells/3 rats) and AAV-hTyr-injected rats (n = 42 cells/3 rats); RM two-way ANOVA, I_h amplitude: treatment X voltage P = 0.485. I_h density: treatment X voltage P = 0.112. **B,** Representative traces of afterhyperpolarization-associated current (I_{AHP}) (left); plots of I_{AHP} peak (middle) and I_{AHP} area (right) in female SNpc DA neurons of AAV-null (n = 36 cells/3 rats) and AAV-hTyr rats (n = 36 cells/3 rats); Mann-Whitney test; I_{AHP} peak: P = 0.088; I_{AHP} area: P = 8.33 X 10^{-4}). **C,** Quantification of I_{AHP} peak and area normalized on C_m (I_{AHP} peak density and area density, respectively) (AAV-null n = 35 cells/3 rats; AAV-hTyr n = 35 cells/3 rats; Mann-Whitney test, I_h peak density: P = 0.241; I_h area density: P = 0.021. **D,** Example I_h traces (left) and quantification of I_h amplitude (right) of male nigral DA neurons of AAV-null (n = 38 cells/4 rats) and AAV-hTyr rats (n = 41 cells/4 rats); RM two-way ANOVA, Treatment X Voltage, P = 0.118). **E,** Representative I_{AHP} traces (left) and plots of I_{AHP} peak (middle) and I_{AHP} area (right) in male nigral DA neurons of AAV-null (n = 40 cells/4 rats) and AAV-hTyr (n = 38 cells/4 rats); Mann-Whitney test, I_{AHP} peak: P = 0.599; I_{AHP} area: P = 0.292). **A-E,** Data are expressed as mean ± SEM; dots represent individual neurons. *P < 0.05 and ***P < 0.001.

1141

Fig. 6 Nigral NM buildup induces non-motor PD symptoms in male rats.

A, Experimental timeline and design used to evaluate male AAV-hTyr-injected and control rats in
different behavioral domains (anxiety, motivational and motor domains). **B,** Heatmaps (left) and
quantification (right) of the time spent in the elevated plus maze (EPM) (as % of the total time) show-
ing that AAV-hTyr-injected rats spent less time in the open arms (AAV-null n = 10; AAV-hTyr n = 10;
Mann-Whitney test, P = 0.002). **C,** Heatmaps (left) and quantification (right) of the time spent in the
center of an open field (AAV-null n = 10 and AAV-hTyr-injected rats n = 10, Mann-Whitney test, P =
0.011). **D,** Heatmaps (left) and quantification (right) of the time spent in the interaction zone (IZ)
during the social avoidance test; NM-producing rats do not exhibit preference for the social target
(ST) over an empty non-social target (NT) (AAV-null n = 10; AAV-hTyr n = 10; RM two-way ANOVA
followed by Sidak's test, AAV-null: NT *vs* ST P = 0.003, *AAV*-hTyr: NT *vs* ST P = 0.012). **E**, Repre-
sentative images of marble burying test (MBT) (left) and number of marbles buried (right) (AAV-null
n = 10 and AAV-hTyr n = 10, Student's *t*-test, P = 0.080). **F,** Quantification of the sucrose intake
normalized on the body weight of each animal (ml/g) indicating regular sucrose preference in both
groups (AAV-null n = 8; AAV-hTyr n = 8; Kruskal-Wallis followed by Dunn's test, H_2O *vs* Sucrose:
AAV-null P = 0.018; AAV hTyr P = 0.001). **G,** Plots of the number of lever press (left) and reward
delivery (right) obtained by AAV-null (n = 8) and AAV-hTyr-injected rats (n = 8). Data are mean
values of the last 3 days of the self-administration protocol; comparisons were with Student's *t*-test
(lever press, P = 0.882) or Mann-Whitney test (reward delivery, P = 0.959). **H,** Histogram of the
progressive ratio test (AAV-null n = 8; AAV-hTyr n = 8; Student's *t*-test, P = 0.413). **I,** Quantification
of grip strength normalized on the body weight of each animal (AAV-null n = 10; AAV-hTyr n = 10;
Student's *t*-test, P = 0.046). **J,** Representative traces (left) and quantification of the distance moved
in an open field (right) by AAV-null (n = 10) and AAV-hTyr-injected rats (n = 10); Mann-Whitney test
with Holm-Sidak's correction: 0-5 min P = 0.538; 5-10 min P = 0.779; 10-15 min P = 0.40; 15-20 min
P = 0.895; 20-25 min P = 0.895; 25-30 min P = 0.911). Data are expressed as median ± quartiles
(B-I) or mean ± SEM **(J)**; dots represent individual animals. *P<0.05, **P<0.01, ***P<0.001.

1168

1169 **Fig. 7 Nigral NM buildup does not induce behavioral alterations in female rats.**

1170

A, Experimental timeline and design used to evaluate female AAV-hTyr-injected and control rats in
different behavioral domains (anxiety, motivational and motor domains). **B,** Heatmaps (left) and
quantification (right) of the time spent in the elevated plus maze (EPM) (as % of the total time) by
AAV-null (n = 10) and AAV-hTyr-injected rats (n = 11). Student's t-test, P = 0.297. **C,** Heatmaps (left)
and quantification (right) of the time spent in the center of an open field (right) by AAV-null (n = 10)
and AAV-hTyr-injected rats (n = 11); Mann-Whitney test, P = 0.557. **D,** Heatmaps (left) and quanti-
fication (right) of the time spent in the interaction zone (IZ) during the social avoidance test, indicating
regular preference for the social target (ST) over empty non-social target (NT) in both groups (AAV-
null n = 10 and AAV-hTyr n = 11; RM two-way ANOVA followed by Sidak's test. AAV-null: NT *vs* ST
P = 0.006*; AAV-*hTyr*:* NT *vs* ST P = 0.018). **E**, Example images of marble burying test (MBT) (left)
and number of marbles buried (right) by AAV-null (n = 10) and AAV-hTyr-injected rats (n = 11);
Mann-Whitney test (P = 0.845). D, Quantification of the sucrose intake normalized on the body
weight of each animal (ml/g) indicating regular sucrose preference in both groups (AAV-null n = 9;
AAV-hTyr n = 11; Kruskal-Wallis followed by Dunn's test. H_2O vs Sucrose: AAV-null P = 0.0004;
AAV-hTyr P = 0.001). **G,** Plots of the number of lever press (left) and reward delivery (right) obtained
by AAV-null (n = 9) and AAV-hTyr-injected rats (n = 11). Data are mean values of the last 3 days of
the self-administration protocol; comparisons were with Student's *t*-test (lever press, P = 0.236) or
Mann-Whitney test (reward delivery, P = 0.710). **H,** Histogram of the progressive ratio test (AAV-null
n = 9; AAV-hTyr n = 11; Student's *t*-test, P = 0.265). **I,** Quantification of grip strength normalized on
the body weight of each animal (AAV-null n = 10; AAV-hTyr n = 11; Mann-Whitney test, P = 0.06).
J, Representative traces (left) and quantification (right) of the distance moved in an open field by
female AAV-null (n = 10) and AAV-hTyr-injected rats (n = 11); Mann-Whitney test with Holm-Sidak's
correction (0-5 min P = 0.975; 5-10 min P = 0.823; 10-15 min P = 0.993; 15-20 min P = 0.999; 20-
25 min P = 0.913; 25-30 min P = 0.999). Data are expressed as median ± quartiles **(B-I)** or mean ±
SEM **(J)**; dots represent individual animals. *P < 0.05, **P < 0.01, ***P < 0.001.

Extended Data Figures

1197

1198

Extended Data Fig. 1. AAV-induced hTyr expression and NM accumulation in VTA and SNpr

A, Representative double-labeled confocal images of coronal VTA sections immunostained for TH
(green) and hTyr (red) (scale bar: 100 μm). **B,** Representative micrographs of Masson-Fontana
staining in midbrain sections from an AAV-hTyr-injected rat, showing NM accumulation (in dark
brown) in VTA and SNpr four weeks post-AAV-hTyr injection (scale bar: 200 μm, 20 μm, and 10
μm).

1205

1206

1207

Extended Data Fig. 2. Nigral NM buildup does not impair dST dopaminergic fibers.

A-C, Qualitative rat brain atlas showing analysed bregma ranges of coronal serial sections of dorsal
striatum (dST) for quantitative histological analysis. **D-F,** Representative TH-immunolabeled striatal
coronal sections (left, scale bar: 500 μm), and quantification (right) of TH fibers optical density (as
% of AAV-null) in male rats (AAV-null $n = 6$; AAV-hTyr $n = 6$; d: Mann-Whitney test, P = 0.562; e:
Student's *t*-test, P = 0.398; f: Student's *t*-test, P = 0.417. **G-I,** Representative TH-immunolabeled
striatal coronal sections (left, scale bar: 500 μm), and quantification (right) of TH fibers optical density
(as % of AAV-null) in female AAV-null ($n = 5$) and AAV-hTyr-injected rats ($n = 7$); g: Student's *t*-test,
P = 0.712; h: Student's *t*-test, P = 0.967; i: Student's *t*-test, P = 0.929. Data are expressed as mean
± SEM **(D-I)**; dots represent individual animals.

1218
1219

Extended Data Fig. 3. Nigral NM buildup does not impair NAc dopaminergic fibers.

A-C, Representative TH-immunolabeled coronal sections of NAc Core and NAc Shell (left, scale bar:
500µm), and quantification (right) of TH fibers optical density (as % of AAV-null) in male rats (AAV-
null *n* = 6; AAV-hTyr *n* = 6; a: Student's *t*-test, P = 0.211; b: Student's *t*-test, P = 0.087; c: Student's
t-test, P = 0.818). **D-F,** Representative TH-immunolabeled striatal coronal sections (left, scale bar:
500 µm) and quantification (right) of TH fibers optical density (as % of AAV-null) in female rats (AAV-
null *n* = 5; AAV-hTyr *n* = 6; d: Student's *t*-test, P = 0.708; e: Student's *t*-test, P = 0.920; f: Student's
t-test, P = 0.685). Data are expressed as mean ± SEM **(A-F)**; dots represent individual animals.

1228

1229

1230 **Extended Data Fig. 4. Additional data related to the sucrose self-administration in male rats**
1231 **presented in Fig. 6.**

1232

A, Quantification of water intake normalized on body weight of each animal (ml/g) (AAV-null $n = 8$;
AAV-hTyr $n = 8$; Mann-Whitney test, P = 0.130). **B,** Quantification of sucrose intake (as % of total
intake) of AAV-null ($n = 8$) and AAV-hTyr-injected rats ($n = 8$); Mann-Whitney test (P = 0.234). **A-B)**
Data are median ± quartiles and dots represent individual animals. **C**, Plot of mean body weight of
AAV-null ($n = 8$) and AAV-hTyr-injected rats ($n = 8$) during different experimental phases: two-bottle
choice (TBC, days -1 to -4), training phase (days 1 to 8), and progressive ratio (PR) (day 9); RM two-
way ANOVA followed by Sidak's test, days: -1 (P = 0.979), -2 (P = 0.973), -3 (P = 0.949), -4 (P =
0.992), 1 (P = 0.995), 2 (P = 0.988), 3 (P = 0.974), 4 (P = 0.991), 5 (P = 0.992), 6 (P = 0.997), 7 (P
= 0.975), 8 (P = 0.994) and 9 (P = 0.990). **D**, Chart of mean number of lever press across days (AAV-
null $n = 8$; AAV-hTyr $n = 8$; Mann-Whitney test with Holm-Sidak's correction, days: 1 (P = 0.963), 2
(P = 0.871), 3 (P = 0.163), 4 (P = 0.967), 5 (P = 0.642), 6 (P = 0.967), 7 (P = 0.910), and 8 (P =
0.910). **E**, Plot of mean number of rewards delivery across days (AAV-null $n = 8$; AAV-hTyr $n = 8$;
Mann-Whitney test with Holm-Sidak's correction; days: 1 (P = 0.989), 2 (P = 0.989), 3 (P = 0.718),
4 (P = 0.987), 5 (P = 0.761), 6 (P = 0.989), 7 (P = 0.872) and 8 (P = 0.985). **C-E)** Data are expressed
as mean ± SEM. *P < 0.05, **P < 0.01.

1248

Extended Data Fig. 5. Additional data related to the sucrose self-administration in female rats presented in Fig. 7.

A, Quantification of water intake normalized on body weight of each animal (ml/g) (AAV-null n = 9; AAV-hTyr n = 11; Mann-Whitney test, P = 0.564). **B,** Quantification of sucrose intake (as % of total intake) of AAV-null (n = 9) and AAV-hTyr-injected rats (n = 11); Mann-Whitney test, P = 0.067. **(a,b)** Data are median ± quartiles and dots represent individual animals. **C**, Plot of mean body weight of AAV-null (n = 9) and AAV-hTyr-injected rats (n = 11) during different experimental phases: two-bottle choice (TBC, days -1 to -4), training phase (days 1 to 8), and progressive ratio (PR) (day 9). Mann-Whitney test with Holm-Sidak's correction, days: -1 (P = 0.985), -2 (P = 0.985), -3 (P = 0.977), -4 (P = 0.982), 1 (P = 0.985), 2 (P = 0.985), 3 (P = 0.985), 4 (P = 0.985), 5 (P = 0.985), 6 (P = 0.985), 7 (P = 0.985), 8 (P = 0.985), and 9 (P = 0.985). **D**, Chart of mean number of lever press across days (AAV-null n = 9; AAV-hTyr n = 11; Mann-Whitney test with Holm-Sidak's correction; days: 1 (P = 0.999), 2 (P = 0.999), 3 (P = 0.998), 4 (P = 0.999), 5 (P = 0.998), 6 (P = 0.999), 7 (P = 0.983), and 8 (P = 0.672). **E**, Plot of mean number of rewards delivery across days (AAV-null n = 9; AAV-hTyr n = 11; Mann-Whitney test with Holm-Sidak's correction; days: 1 (P = 0.999), 2 (P = 0.999), 3 (P = 0.945), 4 (P = 0.996), 5 (P = 0.999), 6 (P = 0.995), 7 (P = 0.995) and 8 (P = 0.821). **C-E)** Data are expressed as mean ± SEM. #P < 0.05 (between factors); **P < 0.01 (within factors AAV-null).

1267

1268

Extended Data Fig. 6 Sustained NM buildup impairs motor coordination irrespective of sex

A, Experimental timeline and design used to evaluate rats' motor coordination with the rotarod test
at advanced stages of NM buildup (8 weeks post-AAV-hTyr injection). B, Plot of latency to fall from
rotarod during three daily sessions by male AAV-null (n = 10) and AAV-hTyr-injected rats (n = 10),
showing that AAV-hTyr-injected rats fall earlier than controls. RM two-way ANOVA followed by
Sidak's test, days 1 (P = 0.005), 2 (P = 0.015) and 3 (P = 0.011). **C,** Plot of latency to fall from rotarod
across different sessions by female rats indicating that AAV-hTyr-injected rats (n = 9) fall earlier from
rotarod compared to AAV-null (n = 10). Multiple Mann-Whitney test with Holm-Sidak correction, days
1 (P = 0.109), 2 (P = 0.030) and 3 (P = 0.109). **B-C,** Data are expressed as mean ± SEM. *P < 0.05.