†*Cryptovaranoides* is not a squamate

Michael W. Caldwell,[1,2*] Chase D. Brownstein,[3,4] Dalton L. Meyer,[5] Simon G. Scarpetta[6,7],

Michael S.Y. Lee[8,9], Tiago R. Simões[10]

[1]Department of Biological Sciences, University of Alberta, Edmonton, AB, Canada

[2]Department of Earth and Atmospheric Sciences, University of Alberta, Edmonton, AB, Canada

[3]Department of Ecology and Evolutionary Biology, Yale University, New Haven CT

[4]Stamford Museum and Nature Center, Stamford, CT

[5] Department of Earth and Planetary Sciences, Yale University, New Haven CT

[6] Museum of Vertebrate Zoology, Department of Integrative Biology, University of California,

Berkeley, CA

[7]Department of Environmental Science, University of San Francisco, San Francisco, CA

[8]College of Science and Engineering, Flinders University, Adelaide 5001, Australia

[9]Earth Sciences Section, South Australian Museum, North Terrace, Adelaide 5000, Australia.

[10]Department of Ecology and Evolutionary Biology, Princeton University, Princeton NJ, 08544,

USA

*Corresponding author: mw.caldwell@ualberta.ca

Abstract.

Accurate reconstruction of the timescale of organismal evolution requires placement of extinct representatives among living branches. In this way, the fossil record has the capacity to revise hypotheses of organismal evolution by producing representatives of clades that far pre-date the age of the clade inferred using phylogenies built from molecular data and previous fossil calibrations. Recently, one fossil with the potential to drastically change current understanding surrounding the timescale of reptile diversification was described from Triassic fissure-fill deposits in the United Kingdom. This taxon, †*Cryptovaranoides microlanius*, was originally placed deep within the squamate crown clade, suggesting that many lineages of living lizards and snakes must have appeared by the Triassic and implying long ghost lineages that paleontologists and molecular phylogeneticists have failed to detect using all other available data. Our team challenged this identification and instead suggested †*Cryptovaranoides* had unclear affinities to living reptiles, but a crown-squamate interpretation was later re-iterated by the team that

originally described this species. Here, we again challenge the morphological character codings

used to support a crown squamate affinity for †*Cryptovaranoides microlanius* and illustrate

several empirical problems with analyses that find this taxon is a crown squamate. Our analyses

emphasize the importance of stringency in constructing hypodigms of fossils, particularly when

they may be key for proper time calibration of the Tree of Life.

Introduction.

Paleontology has found an important role in the era of widespread genome sequencing of

living phylogenetic diversity: providing justification for the placement of fossil calibrations

along molecular phylogenies [1–6]. Placing fossils on the Tree of Life is common practice, and

new discoveries [7–10] and reshuffling of hypothesized phylogenetic relationships [11] are

constantly revising what fossils are best to use as prior calibration constraints. These factors

make the robust placement of key fossil taxa essential for properly calibrating phylogenies in

time, which themselves form a foundation for modern evolutionary biology [12].

Recently, Whiteside et al. [13] described †*Cryptovaranoides microlanius* based on a

partially articulated skeleton and a collection of referred material from the Carnian [14] to

Norian-Rhaetian [13,15–17] (237-201.5 million years ago) fissure fill deposits of England, UK.

In a paper [18] published at the end of 2023 we refuted the affinities of †*C. microlanius* to

Anguimorpha, a deeply nested crown squamate clade, proposed by Whiteside et al. [13]. In their

response to our paper, Whiteside et al. [19] disagree with many of our anatomical observations

and restate their position on the affinities of †*C. microlanius.*

Whiteside et al. [19] referred additional Late Triassic fossils to †*Cryptovaranoides microlanius*

in support of Whiteside et al. [13] and presented phylogenetic results that this taxon is a crown

group squamate (squamate hereafter). Whiteside et al. [19] is also an impassioned rebuttal to

Brownstein et al. [18], who substantially revised the description of †*Cryptovaranoides* and upon

correction of numerous character scorings in Whiteside et al. [13], found this taxon to be "*either*

an archosauromorph or an indeterminate neodiapsid" and not a lepidosaur, much less a crown

squamate.

Here we provide point-by-point refutations of the interpretation of Whiteside et al.

[13,19] and describe what we consider to be major methodological errors in the comparative

anatomical work and phylogenetic analyses they conduct. We also emphasize that both

Whiteside et al. [13,19] papers fail to replicate the inferred position of †*C. microlanius* across

phylogenetic analyses of the same and across different morphological datasets. We also highlight

here where Whiteside et al. [19] should have reported their recovered synapomorphies, they

elected instead to report results from other studies as though they had been recovered in the

analyses conducted by Whiteside et al. [19].

Results.

Factual Errors in Whiteside et al. [19].

Whiteside et al. [13,19] include numerous substantive comparative anatomy errors that can

broadly be grouped into two categories: (i) anatomical interpretation, and (ii) translating these

interpretations into scorings in different phylogenetic datasets. In Brownstein et al. [18], we

identified 22 errors in the study of Whiteside et al. [13] of type (i) ("Results" in Brownstein et al.

[18]) and several more of type (ii) ("Supplementary Material" in Brownstein et al. [18]). In turn,

Whiteside et al. [19] claimed that there were five observational errors in Brownstein et al. [18].

Four of these are supposed observational errors (type (i)) and one concerns a difference in how

Brownstein et al. [18] and Whiteside et al. [19] score a character (type (ii)). This implies that

Whiteside et al. [19] admitted that the other 18 errors type (i) produced by their previous study

were correctly identified by Brownstein et al. [18]. Yet, Whiteside et al. [19] discuss additional

characters in sections of their study and provide different interpretations of the anatomy of

†*Cryptovaranoides microlanius* than do Brownstein et al. [18], but without clear justification.

Here we focus on the four supposed observational errors (type (i)) listed by Whiteside et al. [19].

Entepicondylar and ectepicondylar foramina of humerus. Whiteside et al. [19] provide additional

photographs of features on the distal ends of the humeri referred to †*Cryptovaranoides*

microlanius that they maintain are homologous with the entepicondylar and ectepicondylar

foramina observed in some, but not all, squamates. We disagree about the identity of the features

that Whiteside et al. [19] identify as these paired distal foramina for the following reasons:

(i) First, the features on the humeri that Whiteside et al. [19] figure are not foramina, but

fossae on the posterodistal surface of the humerus that are filled in with sediment. Although

Whiteside et al. [19] claim to observe this in a third isolated and referred humerus (NHMUK

R38929), they neither figure this third humeral fragment, nor do they figure the internal structure

of any of the supposed humeral heads, nor the evidence for referring these isolated elements to

†*C. microlanius*. Contrary to Whiteside et al. [19], computed tomography scans provide essential

information about the structure of these fossae and show that, when infill is removed, foramina

are absent (figure 4 in Brownstein et al. [18]).

(ii) The structures that Whiteside et al. [19] interpret to be the entepicondylar and

ectepicondylar foramina on the holotype and referred humeri are also in the wrong place on the

bone to be homologized as such. In all crown reptiles that possess these foramina, they are

placed low on the anterior surface (entepicondylar) and high on the distal surface

(ectepicondylar) of the humerus (e.g., see [8,20] for examples in squamates, [21] for examples in

turtles, and finally [22,23] for examples in rhynchocephalians) and are dissimilar in shape and

size (the entepicondylar foramen is elongated along the long axis of the humerus; the

ectepicondylar foramen is circular). The features that Whiteside et al. [19] figure on the holotype

and referred humeri of †*C. microlanius* are very similar in shape and size, are oddly both placed

on the same side of the bone, and differ in placement from the foramina observed in other

exemplar fossils and living species of reptiles [8,20–23]. The morphology of the fossae observed

on the distal end of the humeri referred to †*C. microlanius* by Whiteside et al. [19] are, however,

similar in placement, size, and shape to the fossae described on the distal ends of the humeri of

some archosauromorphs, including the azendohsaurid †*Puercosuchus traverorum* (figure 13a in

[24]).

(iii) We reiterate that the presences of these foramina are not an unambiguous

synapomorphy of crown Squamata, and are present across living reptilian diversity [8,20,22]. We

note that, even if their interpretation is accurate, although the ectepicondylar foramen is variably

present across squamates, the entepicondylar foramen is always absent in all crown squamates

[8,20]. Among lepidosaurs, the presence of both foramina would be found only among

sphenodontians and stem lepidosaurs, such as †*Palaeagama* [8,26]. Further, these foramina are

also present in many non-lepidosaurian reptiles, including captorhinids (e.g. †*Captorhinus*),

younginiforms (e.g., †*Hovasaurus* and †*Youngina*), *Claudiosaurus*, Acleistorhinidae

(*Delorhynchus*), *Mesosaurus*, the possible stem turtle †*Eunotosaurus africanus*, and in some

sauropterygians (e.g., *Serpianosaurus* and *Lariosaurus*)[8,34,35]. In conclusion, if anything, the

presence of both foramina would support the assignment of †*Cryptovaranoides* as outside of

crown Squamata.

Absence of jugal posterior process. Whiteside et al. [19] state: "*Except for a very few fossil taxa,* *including two polyglyphanodontians,* † Tianyusaurus *and* † Polyglyphanodon*, ... squamates* *lack a posterior process on the jugal.*" This is entirely incorrect. The two extinct taxa noted by Whiteside et al. [19] here are only unusual among squamates in the fact that they have a complete lower temporal bar, and thus an elongated jugal posterior process [20,27–29]. However, most families of squamates include species with a jugal posterior process (Figure 1)[8,20,28,30,31], even if a complete lower temporal bar is not present. Thus, the presence of a jugal posterior process was incorrectly scored in the datasets used by Whiteside et al. [19] (type (ii) error).

Anterior emargination of the maxillary nasal process. Whiteside et al. [19] state that Brownstein et al. [18] *"seemingly relied on the anteriorly broken left maxilla"* and "*The anterior margin of* *the maxillary nasal process of the right maxilla tapers anteriorly…with no evidence of the type of* *emargination suggested"*. The character in question (ch. 18 in the dataset used by Whiteside et al. [19]), is one of the original characters from [8], which is the basis for the dataset [32] used by Whiteside et al. [19]. In both datasets, this character is described as "*Maxillae,* **posterior** **emargination***, between nasal and orbital processes*"[boldface added], and this character is extensively described in [8]. Therefore, Whiteside et al. [19] incorrectly assess the anterior margin of the maxilla instead of the posterior margin.

Expanded radial condyle of the humerus. Whiteside et al. [19] (p. 3) state that Whiteside et al. [13] "*noted an expanded radial condyle on the humerus.*" Whiteside et al. [19] (p.3, fig. 1b) write in their figure caption regarding another isolated fragment: *"(b) NHMUK PV 38911* *isolated larger specimen of the distal end of left humerus of* †Cryptovaranoides microlanius *in* *(above) anterior and (below) posterior views* **showing similar features except the condyle of the** **capitellum.**" (boldface added). Then, in Whiteside et al. [19] (subsection 2.3), the authors state that this fragment does not have a radial condyle/capitellum of the humerus, which they defend as follows: "*This is reinforced by the larger humerus (figure 1b) which is missing the condyle,* ***as*** ***is typical in the preservation of fissure lepidosaurs*** *(e.g.* †Clevosaurus; *[12, fig. 29b]) but the*

cavity in which it sat clearly indicates a substantial condyle in life.” (boldface added). What

Whiteside et al. [19] highlight is that there is a radial condyle (i.e., capitellum) preserved insome

specimens, but not in others, , deferring such difference among the materials referred to

†*Cryptovaranoides* to poor preservation without justification beyond *“as is typical ….of fissure*

lepidosaurs…” . Taphonomy and poor preservation cannot be used to justify the inference that

anatomical feature was present when it is not preserved and there is no evidence of postmortem

damage. In such a situation when a feature's absence is potentially ascribable to preservation, its

presence should be considered ambiguous—or that some of these materials may simply not

belong to the same taxon, which is the point raised previously by Browenstein et al. [18]).

Finally, even in the case of the isolated humerus with a preserved radial condyle (illustrated by

Whiteside et al. [19]), thisis fairly small compared to that of squamates, either crown memebs or

the earliest known pan-squamates, such as †*Megachirella wachtleri* (Fig.1 in [8]).

Evaluation of character state determinations provided by Whiteside et al. (2024).

In this section, we revisit each alternative interpretation of the anatomy of

†*Cryptovaranoides microlanius* provided by Whiteside et al. [19], who discussed 26 characters

that they suggest have bearing on the placement of this taxon within Lepidosauria, Pan-

Squamata, crown Squamata, and successive clades within the crown. We note first that these

characters do not correspond to optimized character states in the phylogenies that Whiteside et al.

[13,19] infer, but instead to an assemblage of character state optimizations presented in various

papers in the literature (e.g., [20,30,33]). This is not an issue *per se* for a referral of †*C.*

microlanius to Squamata, but it does mean that these characters do not clearly support the

position(s) of †*C. microlanius* among reptiles that Whiteside et al. [13,19] recover in the

phylogenies they present. With this noted, we provide a point-by-point discussion of character

interpretations in Brownstein et al. [18] that Whiteside et al. [19] challenge.

Preservation of the septomaxilla. Whiteside et al. [13,19] claim that a small, disarticulated piece

of bone preserved anterolateral to the vomer in block containing the holotype of

†*Cryptovaranoides microlanius* is the septomaxilla. They further suggest that a portion of the

medial surface on a maxilla referred to this taxon but clearly from a much larger reptile provides

additional support for the presence of a septomaxilla in †*C. microlanius.* The CT scans published

by Whiteside et al. [13,19] show that the bone in the holotype NHMUK PV R36822 is isolated

and with no clear morphological affinities to the septomaxillae in squamates (see CT scans in

[20]). Whiteside et al. [19] also suggest that the identity of this bone as the septomaxilla is

supported by its placement between the maxilla and premaxilla of the holotype, but given the

level of disarticulation of the holotype and the morphology of the bone fragment, a similar

argument could be made that it is a portion of the anterior end of one of the vomers, or

potentially, the contralateral premaxilla in dorsal view. In any case, we feel it is premature to

code †*C. microlanius* for any character related to the morphology of the septomaxilla based on

this disarticulated and damaged bone fragment.

Expanded radial condyle of the humerus. Whiteside et al. [19] provide additional justification of

the presence of an expanded radial condyle of the humerus by stating that the projection of the

radial condyle above the adjacent region of the distal anterior extremity of †*Cryptovaranoides*

microlanius supports their choice to score the expanded radial condyle as present. The projection

of the radial condyle above the adjacent region of the distal anterior extremity is not the

condition specified in either of the morphological character sets that Whiteside et al. [19] cite

[18,32]. The condition specified in those studies is the presence of a distinct condyle that is

expanded. The feature described in Whiteside et al. [19] does not correspond to the character

scored in the phylogenetic datasets (see also additional considerations for this character in the

previous section). .

Absence of the posterior process of the jugal. Whiteside et al. [19] suggest that the absence of the

posterior process of the jugal in †*Cryptovaranoides microlanius* supports its affinities to

Squamata. The absence of the jugal posterior process, which we have argued is not clear in the

holotype of this species and may be worn off [18], is a notoriously variable character among

lepidosaurs that has caused substantial confusion about the placement of fossils. The new figure

provided by Whiteside et al. [19] (figure 1f) of the jugal is pixelated and unclear. The presence of

a jugal posterior process, including its development into a complete jugal bar, is documented for

numerous pan-squamate, pan-lepidosaur, and extinct crown lepidosaur species

[8,20,29,32,36,37]. Notably, and as stated by [19], complete loss of the temporal bar potentially

diagnoses all lepidosaurs (with presence in rhynchocephalians being a reversal), so absence of

the jugal process in †*Cryptovaranoides microlanius* (if confirmed) cannot be used to place it

within Squamata The lower temporal bar is, however also incomplete, and the jugal posterior

process variably developed, in numerous clades within Archosauromorpha, including the

†*Azendohsauridae* [38–41], †*Prolacerta broomi* [42,43], †*Teyujagua paradoxa* [44], and

'†*Protorosauria*' [45–50].. The development and enclosure of the temporal fenestra in reptiles

has been linked to the expression of two genes, *Runx2* and *Msx2*, in *in vivo* studies [51]. We

suspect that the variable extent of this feature in many early-diverging lepidosaur and

archosauromorph lineages might be an example of the 'zone of variability' [52], in which the

canalization of development and other constraints (e.g., functionality; see [53]) had not yet

completely acted to 'fix' the morphology of the posterior process. This hypothesis will of course

require further experimental study of living model systems.

Anterior emargination of the maxillary nasal process. Whiteside et al. [19] flag our interpretation

of the anterior maxillary nasal process as emarginated, which we found united

†*Cryptovaranoides microlanius* with archosauromorphs. Although we still interpret this state as

such based on the computed tomography scan data, we again note than none of our analyses [18]

unambiguously place †*Cryptovaranoides microlanius* within Archosauromorpha and that, while

optimized as such, this single character necessarily has limited utility for placing †*C. microlanius*

among reptiles.

Subdivision of the metotic fissure. Whiteside et al. [19] claim that the division of the metotic

fissure into the vagus foramen and recessus scalae tympani by the crista tuberalis, a key

squamate feature, can be scored for †*Cryptovaranoides microlanius*, yet paradoxically suggest

the presence of this condition is only inferable based on other observations of the anatomy of the

holotype and referred specimens. Actually, Whiteside et al. [19] argue that because another

character relating to a different part of the structure of the metotic fissure can be inferred based

on the presence of the crista tuberalis, that the division of the metotic fissure may also be inferred

by the presence of the crista tuberalis. This logic would imply that no reptile taxon should exist

that possesses solely either a crista tuberalis or a subdivided metotic fissure, even when this is an

observed state combination in squamates scored for the matrices that both our teams have used to

analyze the phylogenetic position of †*Cryptovaranoides microlanius* [8,32]. This inference

resultantly lacks any justification. In an attempt to verify that †*C. microlanius* possessed a vagus

foramen, Whiteside et al. [19] state that they searched for isolated otoccipital fragments from the

same locality and found an otoccipital fragment (NHMUK PV R36822) with a vagus foramen

that they refer to †*C. microlanius*. It is unclear to us how Whiteside et al. [19] accounted for

confirmation bias when conducting this collections search, or on what basis they refer this

isolated bone to †*C. microlanius*. In any case, the presence of the lateral opening of the recessus

scalae tympani is not observable in the fragment, and thus the division of the metotic fissure is

not demonstrated in the referred fragment.

Fusion of exoccipitals and opisthotics. As Whiteside et al. [19] note, we concur with their

identification of an otoccipital in †*Cryptovaranoides microlanius* formed by the fusion of the

exoccipitals with the opisthotics. Our concern is with the use of this feature to assign †*C.*

microlanius to Squamata. Whiteside et al. [19] cite de Queiroz and Gauthier [33], who list the

presence of an otoccipital as a distinguishing feature of squamates. However, Whiteside et al.

[19] fail to provide any phylogenetic evidence supporting the optimization of this feature as a

squamate synapomorphy in phylogenies including †*C. microlanius*, nor do they recognize that an

otoccipital is present in numerous non-squamate reptiles, including numerous archosauromorphs

[24,54,55]. For these reasons, the presence of an otoccipital cannot be used to assign †*C.*

microlanius to Squamata or even Lepidosauria instead of Archosauromorpha or other clades of

reptiles known from the Permo-Triassic, except probably the turtle total clade (†*Eunotosaurus*

africanus possesses unfused exoccipitals and opisthotics [56]). We acknowledge that just

because a character state is widespread among reptiles, its optimization along a phylogeny is

what is of primary importance for referring taxa to a particular clade. To this end, we reiterate

that Whiteside et al. [19] do not show that the presence of an otoccipital optimizes as an

ambiguous or unambiguous synapomorphy of Lepidosauria, Pan-Squamata, or Squamata in any

of their phylogenies. For characters that show evidence of homoplastic evolution like the

presence of an otoccipital (and indeed, the presence of a jugal posterior process; see above),

phylogenetic character optimization is essential.

Enclosed vidian canal exiting anteriorly at base of each basipterygoid process. In our restudy of

the holotype of †*Cryptovaranoides microlanius*, Brownstein et al. [18] were unable to verify the

presence of an enclosed vidian canal exiting the sphenoid (=parabasisphenoid) via the base of the

corresponding basipterygoid process. Whiteside et al. [19] take issue with Brownstein et al. [18]

interpretation of this region of the braincase and suggest that a larger, abraded, and isolated

sphenoid (NHMUK PV R 37603a) that they refer to †*C. microlanius* without justification also

supports their interpretation of the anatomy of this region of the braincase, yet they also say that

this fragment is of only limited informativeness and appear to agree with us that the best course

of action is to score this character as missing data when including †*C. microlanius* in

phylogenetic analyses.

Development of the choanal fossa of the palatine. Squamata includes multiple instances where

the complexity of the bony palate increases dramatically, an innovation that is related to

chemosensory evolution and the integration of different bones that form this region of the skull

[20,20,30,31,57–61]. One feature recognized as phylogenetically informative is the development

of the choanal fossa on the ventral surface of the palatine (also known as the palatine

sulcus)[18,20,31,62]. Whiteside et al. [19] claim that the development of the palatine choanal

fossa in †*Cryptovaranoides microlanius* is comparable to the development of this feature in

living squamates, and cite the living iguanian genus *Ctenosaura* as an example of a squamate

with similar anteroposterior development of this feature as †*C. microlanius.* Importantly, most

iguanians appear to show an apparently plesiomorphic condition where the choanal fossa is

anteriorly restricted on the palatine [20]; this feature has actually contributed to debates about

whether iguanians form the living sister to all other crown squamates (as found in some

morphological character-based phylogenies; [20,30]) or are deeply nested within the crown clade

and secondarily convergent with rhynchocephalians (as implied by phylogenies based on DNA

sequence data [63–69] alone or with morphological data [8]). Some crown squamates almost

completely lack the fossa (Fig 3), further complicating the argument that the choanal fossa in †*C.*

microlanius represents the plesiomorphic squamate condition; a proper phylogenetic analysis is

needed.

308   Secondly, Whiteside et al. [19] take issue with Brownstein et al. [18] contention that the

palatine fossa is present, albeit variably, across a wide swath of reptilian diversity, including

many early-diverging archosauromorph clades. They explicate this concern by distinguishing

between the narrow channel found in taxa like †*Tanystropheus* (figure 11 in Whiteside et al. [19])

and the wider fossa found in †*C. microlanius*. We agree that the fossa in †*C. microlanius* is more

developed than in some early-diverging lepidosaurs, such as †*Marmoretta oxoniensis* [37].

However, it is clear that the feature in †*Cryptovaranoides* falls within the variation in the choanal

fossa length and depth (Figure 2; Figure 3) that represents the ancestral condition in lepidosaurs

[20] and is exemplified across archosauromorph (see, for example [24]) and indeed diapsid [70–

72] diversity. Finally, Whiteside et al. [19] cite the discovery of a large, isolated palatine that

they state confirms the presence of a "squamate-type" choanal fossa in †*C. microlanius*, except

they provide no justification for why this isolated bone should be referred to this species. In any

case, the morphology of that bone is also unlike those of squamates (Figure 2).

Vomer ventral ridges and dentition. Whiteside et al. [19] reiterate their earlier characterization

[13] of the vomer of †*Cryptovaranoides microlanius* as ridged, and suggest that the morphology

of the vomerine ridges and vomerine teeth in †*C. microlanius* is comparable to the condition in

the anguid anguimorph *Pseudopus apodus*. We dispute this on the basis that the vomer of †*C.*

microlanius as figured by Whiteside et al. [19] shows no clear ridges equivalent to those in

Pseudopus apodus [73] or other squamates with ridged vomers, including extinct forms such as

†*Eoscincus ornatus* [31] (Figure 4). Instead, the new photograph of the holotype specimen of †*C.*

microlanius provided in Whiteside et al. [19] shows that the vomer is indeed toothed, but no

ridges are figured or visible. We once again re-examined the CT scan data and failed to find any

structure resembling the ridges present in some anguimorphs, although we acknowledge that the

nature of the scan data (43 microns per segment) means that the vomer surface on the scan

segmentations is coarse. In any case, no ridges equivalent in shape or size to those found in

anguimorph squamates are present in the holotype of †*C. microlanius.*

The presence of teeth on the vomer is itself rare among squamates; only in the pan-

scincoid †*Eoscincus ornatus* [31] and some [73] anguid and varanoid [74] species are vomerine

teeth documented. Whiteside et al. [13,19] appear to suggest that the presence of vomerine ridges

and a row of vomerine teeth therefore allies †*C. microlanius* with anguimorphs, even though

none of the phylogenies that they infer (and indeed, no phylogeny to our knowledge) optimizes

the presence of vomerine teeth as a synapomorphy of crown Anguimorpha. Furthermore, the

structure of the vomer in †*C. microlanius* is fundamentally unlike those of anguimorph

squamates, which are elongate and possess pronounced ridges that are placed medially on the

ventral surface of the main body of each vomer and each house only one row of teeth along their

posterior third [73,75]. In contrast, the condition that Whiteside et al. [19] figure for †*C.*

microlanius shows multiple, parallel rows of vomerine teeth on either side of the vomer that run

across the entire length of the bone. This morphology is even unlike that observed in the only

squamate known where multiple vomerine tooth rows are present, †*Eoscincus ornatus* [31]

(Figure 4). Multiple rows of vomerine teeth are common in reptiles outside of Squamata [76]; the

presence of only one row is restricted to a handful of clades, including millerettids [77,78],

†*Tanystropheus* [49], and some [79], but not all [71,80] choristoderes. In †*Eoscincus ornatus*, all

vomerine teeth are restricted to a raised surface at the posterior end of the ventral surface of the

main body of the vomer [31]. Thus, Whiteside et al. [19] have provided no evidence that

vomerine ridges are present in †*C. microlanius* nor that the presence of ridges and teeth is a

synapomorphy of Anguimorpha. We note again that vomerine teeth are widespread across

amniote diversity outside Squamata [76–78] and appear in many lineages of Triassic

archosauromorphs.

Lacrimal arches dorsally over lacrimal duct and floors lacrimal duct with medial process

posteriorly. Brownstein et al. [18] argued that this feature was unobservable in the holotype of

†*Cryptovaranoides microlanius* [18]. Whiteside et al. [19] dispute this by suggesting that their

CT scan data does not reliably show this feature and instead provide a photograph of the

holotype skull that they state shows this morphology (figure 2a-c in [19]). This figure shows no

curvature to the lacrimal, which appears as a flat element, whereas an arch dorsally and a floor

ventrally would indicate the presence of a foramen. We are also confused by Whiteside et al. [19]

statement that they did not code this feature (dorsal arcuation of the lacrimal over the lacrimal

duct, which is posteriorly floored by the medial process of the lacrimal) for †*C. microlanius* but

use it to refer this species to the Anguimorpha based on its optimization as a plesiomorphy of

that clade in phylogenetic analyses that place †*C. microlanius* within Anguimorpha. This appears

to suggest that, despite never conducting an analysis where this feature is coded as 'present' for

†*C. microlanius*, Whiteside et al. [19] used the presence of this feature to ally †*C. microlanius*

with Anguimorpha not as a recovered synapomorphy but merely deciding it was so. This action

would be an arbitrary unification of a species with a clade based on a selected character state and

would deny the equal possibility that this character state is absent due to secondary reversal in

†*C. microlanius.* In sum, the use of lacrimal morphology to ally †*C. microlanius* with

anguimorphs is apparently not based on direct character state optimization, i.e., a test of

congruence.

Distinct quadratojugal absent. The quadratojugal cannot be located in the holotype of

†*Cryptovaranoides microlanius* [13,19]. Brownstein et al. [18] argued that this might be due to

postmortem disarticulation and damage to the skull, and also noted that a complete ontogenetic

series would *ideally* be required to test whether the quadratojugal is modified throughout

ontogeny [18]. Whiteside et al. [19] focused on our comment about the ideal situation of having

an ontogenetic series for †*C. microlanius* to assess the development of the quadratojugal, which

we restate would be helpful to understand how this bone transforms through ontogeny given the

complex restructuring to this region of the skull that occurs throughout the evolution of

lepidosaurs [36]. However, Whiteside et al. [19] state "we argue based on juvenile and adult

specimens and the absence of a quadratojugal facet on the quadrate."

First, as highlighted by Brownstein et al. [18] , the region of the quadrate that would

articulate with the quadratojugal is not preserved in the holotype of †*C. microlanius*, so it is

impossible to tell whether a facet for the quadratojugal is present. Second, Whiteside et al. [19]

fail to provide any justification for the referral of the isolated partial quadrate NHMUK PVR

37606 to †*C. microlanius*. Third, Whiteside et al. [19] fail to identify the quadratojugal facet on

this isolated quadrate when they figure this bone; indeed, based on the figure, the process that

would have housed the quadratojugal facet is also missing from this quadrate (NHMUK PVR

37606). It is impossible to tell whether †*C. microlanius* lacked a quadratojugal based on the

current data. Instead, all that can be said is that a quadratojugal is not preserved in the holotype

of †*C. microlanius*, and the region housing the articular facet for the quadratojugal is not

preserved in either the holotype or referred quadrate.

Pterygoid/quadrate overlap. Whiteside et al. [19] restate their original interpretation of

Whiteside et al. [13] that the pterygoid and quadrate have a short overlap in †*Cryptovaranoides*

microlanius. Whiteside et al. [19] oddly restrict comparisons of the morphology of the quadrate

in †*Cryptovaranoides microlanius*, which they agree is damaged, to the morphology present in

rhynchocephalians, and suggest based on this comparison that their interpretation is correct. We

restate that this is not possible to verify without more complete, articulated or semi-articulated

palates assignable to †*Cryptovaranoides microlanius* based on apomorphic character states and

combinations.

Fusion of the premaxillae and single median tooth. Whiteside et al. [19] suggest that Brownstein

et al. [18] incorrectly characterized the nature of fusion of the premaxillae into a single median

element in †*Cryptovaranoides microlanius*. However, we reiterate that no justification has been

given in their studies [13,19] for referring these isolated premaxillae to †*C. microlanius.*

Whiteside et al. [19] focus on differentiating these isolated large premaxillae from the

premaxillae of two other fissure fill lepidosaurs, †*Gephyrosaurus bridensis* and

†*Diphydontosaurus avonis*, apparently without considering the possibility that additional taxa are

present in the assemblage. Scoring fused premaxillae as present for †*C. microlanius* despite the

presence of unfused, paired premaxillae in the holotype defines (1) ontogenetic character state

transformations solely based on the relative size of the holotype and larger isolated bones

referred without justification and (2) which character state among those present in ontogeny is

the phylogenetically informative one. Defining both of these requires a robust ontogenetic series,

which is not available for †*C. microlanius* at present. Observations of the development of living

squamates also suggest that these larger fused premaxillae should not be referred to †*C.*

microlanius. In squamates with a median premaxilla, the premaxilla is invariably a single

element upon first appearance very early in embryonic development [81–85]. In contrast, the

holotype of †*C. microlanius*, which is very clearly a juvenile (i.e., post-embryonic) specimen,

possesses paired premaxillae. The ontogenetic series that Whiteside et al. [19] suggest for †*C.*

microlanius therefore differs from any known squamate with a single median premaxilla.

Peg-in-notch articulation of quadrate with rod-shaped squamosal. Whiteside et al. [19] suggest

that both their study and Brownstein et al. [18] are in agreement about the presence of a peg-in-

notch articulation between the quadrate and squamosal. However, Brownstein et al. [18]

suggested (and we reiterate herein) that the presence of this type of articulation is unclear based

on the available data for the holotype, as these bones are both damaged in the relevant sections

and disarticulated (Figure 5).

Frontal underlaps parietal laterally on frontoparietal suture. Whiteside et al. [19] confirm that

the presence of this feature in †*Cryptovaranoides microlanius* is entirely based on the

morphology of a referred isolated frontal that they state matches the corresponding articular

portion of the prefrontal in the holotype. However, we do not understand how this element could

match the corresponding articular surface unless it was from the same individual or an animal of

the same size. Unless Whiteside et al. [19] were able to articulate these bones, it is unclear how

this inference can be made. Further, we note that the posterior process of the prefrontal is broken

off in the holotype of †*Cryptovaranoides microlanius.* In squamates and other lepidosaurs, this

process abuts the lateral surface of the frontal along its anteroposterior axis. Two of us (C.D.B.

and D.L.M.) worked to rearticulate the holotype of †*Eoscincus ornatus*, which involved

rearticulating the prefrontal and frontal [31]. Rearticulating these bones is simply impossible

without the complete posterior process of the prefrontal, as the orientation of this process must

match the curvature of the lateral margin of the frontal. Because the isolated frontal is not figured

in either paper by Whiteside et al. [13,19], it is impossible for us to verify whether this bone

actually matches the corresponding articulation surface on the prefrontal in the holotype. In any

case, we regard it a best practice to exclude this isolated frontal from discussions of the affinities

of †*C. microlanius*.

Medial process of the articular and prearticular. Whiteside et al. [19] revise their initial [13]

characterization of the medial process and refer to it as 'rudimentary,' scoring it as unobservable.

We determined that this process is absent [18] and offer no further comment besides that it

appears the distinction between our interpretation of this feature and that of Whiteside et al. [19]

is, as they explicate, largely arbitrary.

Bicapitate cervical ribs and cervical ribs with an anteriorly oriented process. Whiteside et al.

[19] took issue with Brownstein et al. [18] characterization of the morphology of the cervical ribs

in †*Cryptovaranoides microlanius*, and compared the bicapitate morphology of the cervical ribs

in †*C. microlanius* to two anguimorph squamates, *Pseudopus apodus* and *Varanus* spp. However,

the cervical ribs of *P. apodus* are not bicapitate, as indeed shown by the figure from [86] that

Whiteside et al. [19] cited. Similarly, the ribs of *Varanus* are unicapitate, not bicapitate, as shown

by [87]. We assume that Whiteside et al. [19] referred to the morphology of the cervical rib head

in *P. apodus* because of the presence of a posterior process on the rib head [86]. This is not the

bicapitate condition and is not homologous with the morphology of the cervical rib heads in

either †*C. microlanius* or archosauromorphs, where an anterior process is offset from the process

formed by the capitulum and tuberculum [54,88]. We reiterate that the condition in †*C.*

microlanius is identical to that in archosauromorphs (figure 2c, figure 3 in [18]); the comparisons

made by Whiteside et al. [19] are across non-homologous structures and result from a

misinterpretation of cervical rib anatomy. This confusion in part stems from the lack of a fixed

meaning for uni- and bicapitate rib heads; in any case, †*C. microlanius* possesses a condition

identical to archosauromorphs as we have shown.

Cervical and dorsal vertebral intercentra. Whiteside et al. [19] suggest that Brownstein et al.

[18] inferred the presence of cervical and dorsal intercentra in †*Cryptovaranoides microlanius*;

however, we did not propose this in that study. In fact, Whiteside et al. [13] state in their original

paper that there "are gaps between the vertebrae indicating that intercentra were present (but

displaced in the specimen) on CV3 and posteriorly. Some images of bones on the scans are

identified as intercentra" (p. 11, [13]). The statement in [19] consequently represents an incorrect

attribution and an incorrect characterization of our reexamination, as we determined that no

cervical and dorsal intercentra were preserved or present in the holotype. Whiteside et al. [19]

justify their position that cervical intercentra are present in †*C. microlanius* based on the

morphology of another isolated bone that they refer to this taxon without justification. In any

case, the absence of dorsal intercentra is not a distinguishing feature of squamates because

several lineages of squamates, including lacertids, xantusiids, gekkotans, and the stem-squamate

†*Bellairsia gracilis* all possess cervical and dorsal intercentra [32,89,90].

Anterior dorsal vertebrae, diapophysis fuses to parapophysis. Whiteside et al. [19] concur with

Brownstein et al. [18] that the diapophyses and parapophyses are unfused in the anterior dorsals

of the holotype of †*Cryptovaranoides microlanius*, and restate that fusion of these structures is

based on the condition they observed in isolated vertebrae (e.g., NHMUK PV R37277) that they

refer to †*C. microlanius* based on general morphological similarity and without reference to

diagnostic characters of †*C. microlanius*. This feature should not be scored as present for †*C.*

microlanius.

Zygosphene–zygantrum in dorsal vertebrae. Whiteside et al. [19] again claim that the

zygosphene-zygantrum articulation is present in the dorsal series of †*Cryptovaranoides*

microlanius based on the presence of 'rudimentary zygosphenes and zygantra' in isolated

vertebrae (NHMUK PV R37277) that they again refer to this species without justification. The

structures that Whiteside et al. [19] label as the zygosphenes and zygantra (figure 3 in [19]) are

clearly not, however, zygosphenes and zygantra, as the former is by definition a centrally located

wedge-like process that fits into the zygantrum, which is a fossa on the following vertebra. The

structures labeled zygosphenes and zygantra by Whiteside et al. [19] are just the dorsal surfaces

of the prezygapophyses and the medial margins of the postzygapophyses.

Anterior and posterior coracoid foramina/fenestra. Whiteside et al. [19] use isolated coracoids

(e.g., NHMUK PV R37960) referred to †*Cryptovaranoides microlanius* without justification to

support their claim that the upper 'fenestra' (foramen) of the coracoid in the holotype specimen is

indeed a foramen. Confusingly, Whiteside et al. [19] label this feature as the 'primary coracoid

fenestra' in their figure 3i-j, even though it is clearly the coracoid foramen. Whiteside et al. [19]

appear to have confused the coracoid foramen, which is completely bounded by bone and placed

inside the coracoid, with the coracoid fenestra, which is bounded *in part* by the coracoid (the

coracoid margin is curved to form part of the bounding region in species with the coracoid

fenestra) but also by the interclavicle (see figures in [18,20]). The coracoid fenestra is not

contained within the coracoid. This feature is also not shown to be optimized as a pan-squamate

synapomorphy in any phylogeny including †*Cryptovaranoides microlanius* [13,19], so its

bearing on the identification of †*C. microlanius* as a pan-squamate is unclear to us. Finally, we

note here that Whiteside et al. [19] appear to have labeled a small piece of matrix attached to a

coracoid that they refer to †*C. microlanius* as the supracoroacoid [*sic*] foramen in their figure 3,

although this labeling is inferred because only "suc, supracoroacoid [*sic*]" is present in their

figure 3 caption.

Atlas pleurocentrum fused to axis pleurocentrum. Whiteside et al. [19] reiterate their claim that the atlas and axis pleurocentra are present in †*Cryptovaranoides microlanius* based on their identification of an isolated, globose bone fragment (NHMUK PV 38897) as the atlas intercentrum, but provide no additional justification for this identification and refer the reader to their original paper for a figure of this bone [13], which is only visible on CT scans due to its entombment within the matrix that includes the holotype. To this end, Whiteside et al. [19] revise their initial interpretation of the what they identify as the preserved atlas-axis region, but again without any justification or figures detailing how they reidentified a bone fragment that they had initially believed was a cervical intercentrum as the atlas centrum and intercentrum 2. Thus, Whiteside et [19] do not provide any additional description of how they reidentified these bones or a figure illustrating their revised interpretation, and so we cannot comment on the strength of their revised interpretation. Again, we note that we could not identify the morphology of the atlas and axis with any confidence in the holotype of †*C. microlanius*, and so we again believe any relevant characters should be scored as missing data for this taxon.

Midventral crest of presacral vertebrae. Whiteside et al. [19] appear to agree with Brownstein et al. [18]assertion that a midventral crest is not present on the presacral vertebrae [18]. We never challenged their scoring of keels on the caudal centra as missing data. Rather, we stated that keels are clearly present on the cervical vertebrae [18].

Angular does not extend posteriorly to reach articular condyle. Whiteside et al. [19] state that the posterior portion of the angular is present medially on the right mandible in the holotype, and provide an interpretation of the anatomy of this region. They cite figure 3a in Whiteside et al. [19], but this shows the mandible in lateral view and only a small portion of the angular is visible. As such, it is not clear to us what they interpret to be the posterior portion of the angular. In any case, we could not identify the feature that Whiteside et al. [19] consider to be the posterior angular extent anywhere on the CT scans or on the accessible portions of the real specimen (e.g., figure 2b in [19]). Whiteside et al. [19] describe the posterior extent of the angular as a 'contoured feature,' but we are entirely unclear about what this actually describes.

557 *Ulnar patella.* The only disagreement between Brownstein et al. [18]interpretation and that made

558 by Whiteside et al. [13,19] concerns whether the ulnar patella is absent due to the ontogenetic

559 state or preservation status of the holotype [13,19] or an ontogenetically invariable feature of the

560 anatomy of †*Cryptovaranoides microlanius* [18]; we suggested the latter based on our

561 observation that the forelimb of the holotype is articulated and mostly complete.

Overarching Empirical Problems in Whiteside et al. [19].

564 In this section, we focus on broader empirical issues in Whiteside et al. [19] that directly impact
565 conclusions regarding the taxonomy and phylogenetic placement of *Cryptovaranoides*.
566
Unassignable specimens and "hypodigm inflation." Whiteside et al. [13] and Whiteside et al.

[19] describe and defend their addition of isolated elements to fill in gaps in the holotype concept

of †*Cryptovaranoides* to build, in our view, an unjustifiably inflated hypodigm. Whiteside et al.

[19] (p. 15) try to ameliorate this problem in their Section (5.5) where they state that: "*We*

emphasize that we always match isolated bones with their equivalents on the holotype" and then

state in contradiction that: "*We consider it **remiss** not to include isolated bones as they provide*

details which no scan can. Furthermore, as the holotype is a juvenile, they give additional

*information from larger, and presumably older individuals, **on characters of the holotype***

***otherwise unavailable or uncertain**.*" (boldface added). In summary, Whiteside et al. [19]

claimed to only match isolated bones with equivalent ones in the holotype but also

acknowledged use of larger elements that are not comparable with the ones in the holotype, but

the morphological justification for those referrals (e.g. shared autapomorphies) was not always

explicit. This lack of consistency is a serious empirical issue.

Apomorphic characters not empirically obtained. Whiteside et al. [19] discusses nearly 30

anatomical characters (Sections 2-6), most of which are used to support their core hypothesis that

†*Cryptovaranoides* is a squamate. Yet, none of these 30 characters or conditions are actually

found by Whiteside et al. [19] to be synapomorphies of squamates in their various phylogenetic

analyses. Section 6, Whiteside et al. [19] (p. 16) indicate they modified the datasets of

Brownstein et al. [18] and [32], but preferred the results of [11] and performed two different

analyses based on [32]: one using only morphological data, and one using morphological data

with several constraints from a molecular backbone. The former was tested using maximum

parsimony and the latter using Bayesian inference. What is not clear from either Section 6 or 7, is

which phylogenetic analysis was used by Whiteside et al. [19] to "*review the apomorphy*

distribution". But this uncertainty is inconsequential as the apomorphy distributions discussed in

Section 7 of Whiteside et al. [19] were not derived from their phylogenetic analyses. Instead,

Whiteside et al. [19] (p. 19) exclusively extracted characters from the literature to support their

identification of †*Cryptovaranoides* as a squamate, rather than basing this inference primarily on

their own phylogenetic analyses. These literature sourced characters include *"two diagnostic*

characters of Squamata (=crown-clade Squamata)" and *"further eight squamate*

synapomorphies" listed by Whiteside et al. [19] (Section 7, p. 19). As stated by the authors, "*we*

give the citation for the squamate synapomorphy at the end of each character," indicating that

these diagnostic characters or synapomorphies were picked from the literature and not derived

from ancestral state reconstructions based on their phylogenetic results.

In order to check the characters listed by Whiteside et al. [19] (p.19) as "*two diagnostic*

characters" and "*eight synapomorphies*" in support of a squamate identity for

†*Cryptovaranoides*, we conducted a parsimony analysis of the revised version of the dataset [32]

provided by Whiteside et al. [19] in TNT v 1.5 [91]. We used Whiteside et al.'s [19] own data

version as stated by them, i.e., with (0) scored for character 1, and not including character 383.

We recovered eight apomorphies at the squamate node of which only three were recovered as

unambiguous synapomorphies (boldface indicates the state as scored by Whiteside et al. [19] for

†*Cryptovaranoides*):

Ch. 138. Basisphenoid (or fused parabasisphenoid), ventral aspect, shape, concavity:

single (0)/divided (1)/ absent (**2**). 0 -> 2

Ch. 142. Prootics, alar crest: absent (0)/ present (**1**). 0->1

Ch. 347. Prefrontal/palatine antorbital contact: absent (0) / narrow, forming less than 1/3

the transverse distance between the orbits (1) / contact broad, forming at least 1/2 the distance

between the orbits (**2**). {01}->2

The eight features described by Whiteside et al. [19] (p.19) as synapomorphies for

†*Cryptovaranoides* + Squamata were also not inferred at their recovered crown squamate node

from their own results. Rather, they were copied nearly verbatim from principally two sources

[20,92] and presented as if they were recovered as synapomorphies by Whiteside et al. [19] (p.

19). In Table 1, the italicized text is from Whiteside et al. [19] (p.19), including their literature

source for the supposed synapomorphy. The non-italicized text is the character number and state

from [11] as recovered by us from the TNT analysis conducted (for this reassessment of

Whiteside et al. [19]. We also include where in our phylogeny the character state is recovered as

synapomorphic (Table 1).

Our reanalysis shows that Whiteside et al. [19] did not diagnose †*Cryptovaranoides* +

crown Squamata based on synapomorphies found by their own analyses as there were only three

recovered in their results (see above). Instead, Whiteside et al. [19] provided ad hoc

mischaracterizations of diagnostic features of crown Squamata from studies that did not include

†*Cryptovaranoides* and then discussed and reviewed synapomorphies of Squamata from these

sources [20,92]. Because each additional taxon has the possibility of inducing character

reoptimization, an empirical analysis of character optimization that includes a given new taxon is

necessary to support referring said taxon to any particular clade. If †*Cryptovaranoides* was a

crown squamate, it could plausibly show character distributions not previously sampled among

the crown or stem group.

Discussion and Conclusions.

As we noted in Brownstein et al. [18], the anatomy of †*Cryptovaranoides* is similar to

many Late Triassic crown reptiles. For example, the quadrates of †*Cryptovaranoides* are closely

comparable to those of archosauromorphs such as †*Prolacerta* [42], †*Macrocnemus* [45] and

†*Malerisaurus* [93], with which †*Cryptovaranoides* shares the presence of a notch on the

quadrate for an immobile articulation with the squamosal. Whiteside et al.'s [19] inference that

the quadrate, the notch, and the articulation with the squamosal was mobile, i.e., streptostylic as

in squamates, is unfounded.

Similarly, the vertebrae and cervical ribs of the holotype of †*Cryptovaranoides* do not

resemble those of squamates but are instead comparable to those of non-squamate neodiapsids,

especially archosauromorphs (a point left unaddressed by Whiteside et al. [19]). For example,

whereas fusion of the neural arches to the centra occurs during embryonic ossification in

squamates [94], in †*Cryptovaranoides* unfused bony neural arches and centra are present, as

commonly observed in archosaurs and other non-lepidosauromorph neodiapsids [95,96].

Several errors in Whiteside et al. [13] that we identified in Brownstein et al. [18] were challenged recently by Whiteside et al. [19]. The anatomical interpretations of Brownstein et al. [18] that are challenged in Whiteside et al. [19] are erroneously disputed in the latter study. Whiteside et al. [19] incorrectly interpreted neodiapsid and lepidosaur anatomy and consequently provide clearly erroneous scorings for †*Cryptovaranoides* for morphological character matrices. Whiteside et al. [19] also make several empirical errors, including the unjustified inflation of the †*Cryptovaranoides* hypodigm based on contradictory arguments, and analytical problems (e.g., selecting apomorphic characters from the literature supporting their preferred placement of †*Cryptovaranoides* instead of using characters empirically obtained from phylogenetic ancestral-state reconstruction).

We end this comment with a perspective on the fossil record of neodiapsid evolution. The anatomies and morphologies that diagnose crown group squamates are many and varied, and, except for a few features, hardly universally distributed amongst the living and fossil members of the crown. They are themselves the product of some 250 million years of evolutionary time and would not have evolved in a linear fashion. Rather, phylogenetic analyses of diverse extinct and living reptile clades have shown that the squamate *bauplan* originated in the context of extensive mosaic and homoplastic osteological character evolution [8,31,35,62,72]. We do not doubt that members of Pan-Squamata were present during the Triassic; this is supported by the fossil record [8] as well as numerous time-calibrated phylogenies based on genomic [63,64], morphological [31,32,62], and total-evidence [8,35] data. However, Whiteside et al. [13] and Whiteside et al. [19] have provided no compelling osteological evidence that supports the presence of crown Squamata or any of its inclusive clades in the Triassic.

References

1. Inoue J, Donoghue PCJ, Yang Z. 2010 The Impact of the Representation of Fossil Calibrations on Bayesian Estimation of Species Divergence Times. *Syst. Biol.* **59**, 74–89. (doi:10.1093/sysbio/syp078)

2. Heath TA, Huelsenbeck JP, Stadler T. 2014 The fossilized birth–death process for coherent calibration of divergence-time estimates. *Proc. Natl. Acad. Sci.* **111**. (doi:10.1073/pnas.1319091111)

3. Near TJ. 2004 Estimating divergence times of notothenioid fishes using a fossil-calibrated molecular clock. *Antarct. Sci.* **16**, 37–44. (doi:10.1017/S0954102004001798)

4. Near TJ, Bolnick DI, Wainwright PC. 2005 Fossil calibrations and molecular divergence
time estimates in centrarchid fishes (Teleostei: Centrarchidae). *Evolution* **59**, 1768–1782.
(doi:10.1111/j.0014-3820.2005.tb01825.x)

5. Parham JF *et al.* 2012 Best Practices for Justifying Fossil Calibrations. *Syst. Biol.* **61**, 346–
359. (doi:10.1093/sysbio/syr107)

6. Pennington PT, Cronk QCB, Richardson JA, Near TJ, Sanderson MJ. 2004 Assessing the
quality of molecular divergence time estimates by fossil calibrations and fossil–based model
selection. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* **359**, 1477–1483.
(doi:10.1098/rstb.2004.1523)

7. Field DJ, Benito J, Chen A, Jagt JWM, Ksepka DT. 2020 Late Cretaceous neornithine from
Europe illuminates the origins of crown birds. *Nature* **579**, 397–401. (doi:10.1038/s41586-
020-2096-0)

8. Simões TR, Caldwell MW, Tałanda M, Bernardi M, Palci A, Vernygora O, Bernardini F,
Mancini L, Nydam RL. 2018 The origin of squamates revealed by a Middle Triassic lizard
from the Italian Alps. *Nature* **557**, 706–709. (doi:10.1038/s41586-018-0093-3)

9. Brownstein CD. 2023 Syngnathoid Evolutionary History and the Conundrum of Fossil
Misplacement. *Integr. Org. Biol.* **5**, obad011. (doi:10.1093/iob/obad011)

10. Velazco PM, Buczek AJ, Hoffman E, Hoffman DK, O'Leary MA, Novacek MJ. 2022
Combined data analysis of fossil and living mammals: a Paleogene sister taxon of Placentalia
and the antiquity of Marsupialia. *Cladistics* **38**, 359–373. (doi:10.1111/cla.12499)

11. Budd GE, Mann RP. 2023 Two Notorious Nodes: A Critical Examination of Relaxed
Molecular Clock Age Estimates of the Bilaterian Animals and Placental Mammals. *Syst.*
Biol. , syad057. (doi:10.1093/sysbio/syad057)

12. Smith SD, Pennell MW, Dunn CW, Edwards SV. 2020 Phylogenetics is the New Genetics
(for Most of Biodiversity). *Trends Ecol. Evol.* **35**, 415–425. (doi:10.1016/j.tree.2020.01.005)

13. Whiteside DI, Chambi-Trowell SAV, Benton MJ. 2022 A Triassic crown squamate. *Sci. Adv.*
8, eabq8274. (doi:10.1126/sciadv.abq8274)

14. Simms MJ, Drost K. 2024 Caves, dinosaurs and the Carnian Pluvial Episode: Recalibrating
Britain's Triassic karst 'fissures'. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **638**, 112041.
(doi:10.1016/j.palaeo.2024.112041)

15. Walkden GM, Fraser NC, Simms MJ. 2021 The age and formation mechanisms of Late
Triassic fissure deposits, Gloucestershire, England: Comments on Mussini, G. et al. (2020).
Anatomy of a Late Triassic Bristol fissure: Tytherington fissure 2. *Proc. Geol. Assoc.* **132**,
127–137. (doi:10.1016/j.pgeola.2020.10.006)

16. Chambi-Trowell S, Whiteside D, Benton M. 2019 Diversity in rhynchocephalian
Clevosaurus skulls based on CT reconstruction of two late triassic species from great britain.
Acta Palaeontol. Pol. **64**. (doi:10.4202/app.00569.2018)

17. Butler RJ, Meade LE, Cleary TJ, McWhirter KT, Brown EE, Kemp TS, Benito J, Fraser NC.
2024 *Hwiccewyrm trispiculum* gen. et sp. nov., a new leptopleuronine procolophonid from
the Late Triassic of southwest England. *Anat. Rec.* **307**, 1390–1420. (doi:10.1002/ar.25316)

18. Brownstein CD, Simões TR, Caldwell MW, Lee MSY, Meyer DL, Scarpetta SG. 2023 The
affinities of the Late Triassic *Cryptovaranoides* and the age of crown squamates. *R. Soc.*
Open Sci. **10**, 230968. (doi:10.1098/rsos.230968)

19. Whiteside DI, Chambi-Trowell SAV, Benton MJ. 2024 Late Triassic †*Cryptovaranoides*
microlanius is a squamate, not an archosauromorph. *R. Soc. Open Sci.* **11**, 231874.
(doi:10.1098/rsos.231874)

20. Gauthier JA, Kearney M, Maisano JA, Rieppel O, Behlke ADB. 2012 Assembling the
Squamate Tree of Life: Perspectives from the Phenotype and the Fossil Record. *Bull.*
Peabody Mus. Nat. Hist. **53**, 3–308. (doi:10.3374/014.053.0101)

21. Hermanson G, Arnal FAM, Szczygielski T, Evers SW. 2024 A systematic comparative
description of extant turtle humeri, with comments on humerus disparity and evolution based
on fossil comparisons. *Anat. Rec.* **307**, 3437–3505. (doi:10.1002/ar.25450)

22. Simões TR, Kinney-Broderick G, Pierce SE. 2022 An exceptionally preserved *Sphenodon*-
like sphenodontian reveals deep time conservation of the tuatara skeleton and ontogeny.
Commun. Biol. **5**, 1–19. (doi:10.1038/s42003-022-03144-y)

23. DeMar Jr. DG, Jones MEH, Carrano MT. 2022 A nearly complete skeleton of a new
eusphenodontian from the Upper Jurassic Morrison Formation, Wyoming, USA, provides
insight into the evolution and diversity of Rhynchocephalia (Reptilia: Lepidosauria). *J. Syst.*
Palaeontol. **20**, 1–64. (doi:10.1080/14772019.2022.2093139)

24. Marsh AD, Parker WG, Nesbitt SJ, Kligman BT, Stocker MR. 2022 *Puercosuchus*
traverorum n. gen. n. sp.: a new malerisaurine azendohsaurid (Archosauromorpha:
Allokotosauria) from two monodominant bonebeds in the Chinle Formation (Upper Triassic,
Norian) of Arizona. *J. Paleontol.* **96**, 1–39. (doi:10.1017/jpa.2022.49)

25. Simões TR, Pyron RA. 2021 The Squamate Tree of Life. *Bull. Mus. Comp. Zool.* **163**, 47–95.
(doi:10.3099/0027-4100-163.2.47)

26. Simões TR, Caldwell MW, Pierce SE. 2020 Sphenodontian phylogeny and the impact of
model choice in Bayesian morphological clock estimates of divergence times and
evolutionary rates. *BMC Biol.* **18**, 191. (doi:10.1186/s12915-020-00901-5)

27. Simões TR, Funston GF, Vafaeian B, Nydam RL, Doschak MR, Caldwell MW. 2016
Reacquisition of the lower temporal bar in sexually dimorphic fossil lizards provides a rare
case of convergent evolution. *Sci. Rep.* **6**, 24087. (doi:10.1038/srep24087)

28. Gao K, Norell MA. 2000 Taxonomic composition and systematics of Late Cretaceous lizard
assemblages from Ukhaa Tolgod and adjacent localities, Mongolian Gobi Desert. *Bull. Am.*
Mus. Nat. Hist. **2000**, 1–118. (doi:10.1206/0003-0090(2000)249<0001:TCASOL>2.0.CO;2)

29. Mo J-Y, Xu X, Evans SE. 2010 The evolution of the lepidosaurian lower temporal bar: new
perspectives from the Late Cretaceous of South China. *Proc. R. Soc. B Biol. Sci.*
(doi:10.1098/rspb.2009.0030)

30. Estes R, de Queiroz K, Gauthier J. 1988 Phylogenetic relationships within Squamata. In
Phylogenetic Relationships of the lizard families, pp. 119–281. Stanford, California: Stanford
University Press.

31. Brownstein CD, Meyer DL, Fabbri M, Bhullar B-AS, Gauthier JA. 2022 Evolutionary
origins of the prolonged extant squamate radiation. *Nat. Commun.* **13**, 7087.
(doi:10.1038/s41467-022-34217-5)

32. Tałanda M, Fernandez V, Panciroli E, Evans SE, Benson RJ. 2022 Synchrotron tomography
of a stem lizard elucidates early squamate anatomy. *Nature* **611**, 99–104.

33. de Queiroz K, Gauthier JA. 2020 *Squamata*. CRC Press. See
http://repository.si.edu/xmlui/handle/10088/107192.

34. Hill RV. 2005 Integration of Morphological Data Sets for Phylogenetic Analysis of Amniota:
The Importance of Integumentary Characters and Increased Taxonomic Sampling. *Syst. Biol.*
54, 530–547. (doi:10.1080/10635150590950326)

35. Simões TR, Kammerer CF, Caldwell MW, Pierce SE. 2022 Successive climate crises in the
deep past drove the early evolution and radiation of reptiles. *Sci. Adv.* **8**, eabq1898.
(doi:10.1126/sciadv.abq1898)

36. Ford DP, Evans SE, Choiniere JN, Fernandez V, Benson RBJ. 2021 A reassessment of the
enigmatic diapsid *Paliguana whitei* and the early history of Lepidosauromorpha. *Proc. R.*
Soc. B Biol. Sci. **288**, 20211084. (doi:10.1098/rspb.2021.1084)

37. Griffiths EF, Ford DP, Benson RBJ, Evans SE. 2021 New information on the Jurassic
lepidosauromorph *Marmoretta oxoniensis*. *Pap. Palaeontol.* **7**, 2255–2278.
(doi:10.1002/spp2.1400)

38. Sengupta S, Bandyopadhyay S. 2021 The osteology of *Shringasaurus indicus*, an
archosauromorph from the Middle Triassic Denwa Formation, Satpura Gondwana Basin,
Central India. *J. Vertebr. Paleontol.* **41**, e2010740. (doi:10.1080/02724634.2021.2010740)

39. Sengupta S, Ezcurra MD, Bandyopadhyay S. 2017 A new horned and long-necked
herbivorous stem-archosaur from the Middle Triassic of India. *Sci. Rep.* **7**, 8366.
(doi:10.1038/s41598-017-08658-8)

40. Flynn JJ, Nesbitt SJ, Michael Parrish J, Ranivoharimanana L, Wyss AR. 2010 A new species
of *Azendohsaurus*(Diapsida: Archosauromorpha) from the Triassic Isalo Group of

southwestern Madagascar: cranium and mandible. *Palaeontology* **53**, 669–688. (doi:10.1111/j.1475-4983.2010.00954.x)

41. Sen K. 2003 *Pamelaria dolichotrachela*, a new prolacertid reptile from the Middle Triassic of India. *J. Asian Earth Sci.* **21**, 663–681. (doi:10.1016/S1367-9120(02)00110-4)

42. MODESTO SP, SUES H-D. 2004 The skull of the Early Triassic archosauromorph reptile *Prolacerta broomi* and its phylogenetic significance. *Zool. J. Linn. Soc.* **140**, 335–351. (doi:10.1111/j.1096-3642.2003.00102.x)

43. Sobral G. 2023 The holotype of *Prolacerta broomi* Parrington, 1935 revisited. *Acta Palaeontol. Pol.* **68**. (doi:10.4202/app.01067.2023)

44. Pinheiro FL, França MAG, Lacerda MB, Butler RJ, Schultz CL. 2016 An exceptional fossil skull from South America and the origins of the archosauriform radiation. *Sci. Rep.* **6**, 22817. (doi:10.1038/srep22817)

45. Miedema F, Spiekman SNF, Fernandez V, Reumer JWF, Scheyer TM. 2020 Cranial morphology of the tanystropheid *Macrocnemus bassanii* unveiled using synchrotron microtomography. *Sci. Rep.* **10**, 12412. (doi:10.1038/s41598-020-68912-4)

46. Spiekman SNF, Fraser NC, Scheyer TM. 2021 A new phylogenetic hypothesis of Tanystropheidae (Diapsida, Archosauromorpha) and other "protorosaurs", and its implications for the early evolution of stem archosaurs. *PeerJ* **9**, e11143. (doi:10.7717/peerj.11143)

47. Spiekman SNF, Neenan JM, Fraser NC, Fernandez V, Rieppel O, Nosotti S, Scheyer TM. 2020 The cranial morphology of *Tanystropheus hydroides* (Tanystropheidae, Archosauromorpha) as revealed by synchrotron microtomography. *PeerJ* **8**, e10299. (doi:10.7717/peerj.10299)

48. Nosotti S. 2007 *Tanystropheus longobardicus* (Reptilia Protorosauria): Re-interpretations of the anatomy based on new specimens from the Middle Triassic of Besano (Lombardy, northern Italy). *Mem. Soc. Ital. Sci. Nat.* **35**, 1–88.

49. Spiekman SNF, Neenan JM, Fraser NC, Fernandez V, Rieppel O, Nosotti S, Scheyer TM. 2020 Aquatic Habits and Niche Partitioning in the Extraordinarily Long-Necked Triassic Reptile *Tanystropheus*. *Curr. Biol.* **30**, 3889-3895.e2. (doi:10.1016/j.cub.2020.07.025)

50. Spiekman SNF, Wang W, Zhao L, Rieppel O, Fraser NC, Li C. 2023 *Dinocephalosaurus orientalis* Li, 2003: a remarkable marine archosauromorph from the Middle Triassic of southwestern China. *Earth Environ. Sci. Trans. R. Soc. Edinb.* **114**, 218–250. (doi:10.1017/S175569102400001X)

51. Tokita M, Chaeychomsri W, Siruntawineti J. 2013 Skeletal gene expression in the temporal region of the reptilian embryos: implications for the evolution of reptilian skull morphology. *SpringerPlus* **2**, 336. (doi:10.1186/2193-1801-2-336)

52. Bever GS, Gauthier JA, Wagner GP. 2011 Finding the frame shift: digit loss, developmental
variability, and the origin of the avian hand. *Evol. Dev.* **13**, 269–279. (doi:10.1111/j.1525-
142X.2011.00478.x)

53. Schaerlaeken V, Herrel A, Aerts P, Ross CF. 2008 The functional significance of the lower
temporal bar in *Sphenodon punctatus*. *J. Exp. Biol.* **211**, 3908–3914.
(doi:10.1242/jeb.021345)

54. Ezcurra MD. 2016 The phylogenetic relationships of basal archosauromorphs, with an
emphasis on the systematics of proterosuchian archosauriforms. *PeerJ* **4**, e1778.
(doi:10.7717/peerj.1778)

55. Pinheiro FL, De Simão-Oliveira D, Butler RJ. 2020 Osteology of the archosauromorph
Teyujagua paradoxa and the early evolution of the archosauriform skull. *Zool. J. Linn. Soc.*
189, 378–417. (doi:10.1093/zoolinnean/zlz093)

56. Bever GS, Lyson TR, Field DJ, Bhullar B-AS. 2015 Evolutionary origin of the turtle skull.
Nature **525**, 239–242. (doi:10.1038/nature14900)

57. Rieppel O, Kley NJ, Maisano JA. 2009 Morphology of the skull of the white-nosed
blindsnake, *Liotyphlops albirostris* (Scolecophidia: Anomalepididae). *J. Morphol.* **270**, 536–
557. (doi:10.1002/jmor.10703)

58. Lee MSY. 1997 The phylogeny of varanoid lizards and the affinities of snakes. *Philos. Trans.*
R. Soc. Lond. B. Biol. Sci. **352**, 53–91. (doi:10.1098/rstb.1997.0005)

59. Abramyan J, Richman JM. 2015 Recent insights into the morphological diversity in the
amniote primary and secondary palates. *Dev. Dyn.* **244**, 1457–1468.
(doi:10.1002/dvdy.24338)

60. Strong CRC, Scherz MD, Caldwell MW. 2022 Convergence, divergence, and
macroevolutionary constraint as revealed by anatomical network analysis of the squamate
skull, with an emphasis on snakes. *Sci. Rep.* **12**, 14469. (doi:10.1038/s41598-022-18649-z)

61. Watanabe A, Fabre A-C, Felice RN, Maisano JA, Müller J, Herrel A, Goswami A. 2019
Ecomorphological diversification in squamates from conserved pattern of cranial integration.
Proc. Natl. Acad. Sci. **116**, 14688–14697. (doi:10.1073/pnas.1820967116)

62. Meyer D, Brownstein CD, Jenkins KM, Gauthier JA. 2023 A Morrison stem gekkotan
reveals gecko evolution and Jurassic biogeography. *Proc. R. Soc. B*
(doi:10.1098/rspb.2023.2284)

63. Title PO *et al.* 2024 The macroevolutionary singularity of snakes. *Science* **383**, 918–923.
(doi:10.1126/science.adh2449)

64. Burbrink FT *et al.* 2020 Interrogating Genomic-Scale Data for Squamata (Lizards, Snakes,
and Amphisbaenians) Shows no Support for Key Traditional Morphological Relationships.
Syst. Biol. **69**, 502–520. (doi:10.1093/sysbio/syz062)

65. Townsend TM, Larson A, Louis E, Macey JR. 2004 Molecular Phylogenetics of Squamata:
The Position of Snakes, Amphisbaenians, and Dibamids, and the Root of the Squamate Tree.
Syst. Biol. **53**, 735–757. (doi:10.1080/10635150490522340)

66. Pyron RA, Burbrink FT, Wiens JJ. 2013 A phylogeny and revised classification of Squamata,
including 4161 species of lizards and snakes. *BMC Evol. Biol.* **13**, 93. (doi:10.1186/1471-
2148-13-93)

67. Streicher JW, Wiens JJ. 2017 Phylogenomic analyses of more than 4000 nuclear loci resolve
the origin of snakes among lizard families. *Biol. Lett.* **13**, 20170393.
(doi:10.1098/rsbl.2017.0393)

68. Zheng Y, Wiens JJ. 2016 Combining phylogenomic and supermatrix approaches, and a time-
calibrated phylogeny for squamate reptiles (lizards and snakes) based on 52 genes and 4162
species. *Mol. Phylogenet. Evol.* **94**, 537–547. (doi:10.1016/j.ympev.2015.10.009)

69. Singhal S, Colston TJ, Grundler MR, Smith SA, Costa GC, Colli GR, Moritz C, Pyron RA,
Rabosky DL. 2021 Congruence and Conflict in the Higher-Level Phylogenetics of Squamate
Reptiles: An Expanded Phylogenomic Perspective. *Syst. Biol.* **70**, 542–557.
(doi:10.1093/sysbio/syaa054)

70. Dudgeon TW, Maddin HC, Evans DC, Mallon JC. 2020 The internal cranial anatomy of
Champsosaurus (Choristodera: Champsosauridae): Implications for neurosensory function.
Sci. Rep. **10**, 7122. (doi:10.1038/s41598-020-63956-y)

71. Brownstein CD. 2022 High morphological disparity in a bizarre Paleocene fauna of
predatory freshwater reptiles. *BMC Ecol. Evol.* **22**, 34. (doi:10.1186/s12862-022-01985-z)

72. Jenkins KM, Foster W, Napoli JG, Meyer DL, Bever GS, Bhullar B-AS. In press. Cranial
anatomy and phylogenetic affinities of *aursaurus major*, with new information on the unique
bolosaurid feeding apparatus and evolution of the impedance-matching ear. *Anat. Rec.* **n/a**.
(doi:10.1002/ar.25546)

73. Klembara J, Dobiašová K, Hain M, Yaryhin O. 2017 Skull Anatomy and Ontogeny of
Legless Lizard *Pseudopus apodus* (Pallas, 1775): Heterochronic Influences on Form. *Anat.
Rec.* **300**, 460–502. (doi:10.1002/ar.23532)

74. Yi H-Y, Norell MA. 2013 New Materials of *Estesia mongoliensis* (Squamata: Anguimorpha)
and the Evolution of Venom Grooves in Lizards. *Am. Mus. Novit.* **2013**, 1–31.
(doi:10.1206/3767.2)

75. Krister T. Smith, Jörg Habersetzer. 2021 The anatomy, phylogenetic relationships, and
autecology of the carnivorous lizard *"Saniwa" feisti* Stritzke, 1983 from the Eocene of
Messel, Germany. *Comptes Rendus Palevol* **20**, 441–506.

76. Matsumoto R, Evans SE. 2017 The palatal dentition of tetrapods and its functional
significance. *J. Anat.* **230**, 47–65. (doi:10.1111/joa.12534)

77. Jenkins XA *et al.* 2025 New information on the anatomically derived millerettid Milleretta
rubidgei from the latest Permian based on μCT data. *Zool. J. Linn. Soc.* **203**, zlaf004.
(doi:10.1093/zoolinnean/zlaf004)

78. Jenkins XA, Benson RBJ, Ford DP, Browning C, Fernandez V, Griffiths E, Choiniere J,
Peecook BR. 2025 Cranial osteology and neuroanatomy of the late Permian reptile
Milleropsis pricei and implications for early reptile evolution. *R. Soc. Open Sci.* **12**, 241298.
(doi:10.1098/rsos.241298)

79. Dudgeon TW, Maddin HC, Evans DC, Mallon JC. 2020 Computed tomography analysis of
the cranium of Champsosaurus lindoei and implications for the choristoderan neomorphic
ossification. *J. Anat.* **236**, 630–659. (doi:10.1111/joa.13134)

80. Matsumoto R, Evans SE. 2016 Morphology and function of the palatal dentition in
Choristodera. *J. Anat.* **228**, 414–429. (doi:10.1111/joa.12414)

81. Khannoon ER, Evans SE. 2020 Embryonic skull development in the gecko, *Tarentola*
annularis (Squamata: Gekkota: Phyllodactylidae). *J. Anat.* **237**, 504–519.
(doi:10.1111/joa.13213)

82. Hernández-Jaimes C, Jerez A, Ramírez-Pinilla MP. 2012 Embryonic development of the
skull of the Andean lizard Ptychoglossus bicolor (Squamata, Gymnophthalmidae). *J. Anat.*
221, 285–302. (doi:10.1111/j.1469-7580.2012.01549.x)

83. Skawiński T, Skórzewski G, Borczyk B. 2021 Embryonic development and perinatal
skeleton in a limbless, viviparous lizard, *Anguis fragilis* (Squamata: Anguimorpha). *PeerJ* **9**,
e11621. (doi:10.7717/peerj.11621)

84. Skawiński T, Kaczmarek P, Borczyk B. 2023 Embryonic development of the skull in a
parthenogenetic lizard, the mourning gecko (*Lepidodactylus lugubris*). *J. Anat.* **243**, 618–
629. (doi:10.1111/joa.13871)

85. Ollonen J, Da Silva FO, Mahlow K, Di-Poï N. 2018 Skull Development, Ossification
Pattern, and Adult Shape in the Emerging Lizard Model Organism *Pogona vitticeps*: A
Comparative Analysis With Other Squamates. *Front. Physiol.* **9**.
(doi:10.3389/fphys.2018.00278)

86. Čerňanský A, Yaryhin O, Ciceková J, Werneburg I, Hain M, Klembara J. 2019 Vertebral
Comparative Anatomy and Morphological Differences in Anguine Lizards With a Special
Reference to *Pseudopus apodus*. *Anat. Rec.* **302**, 232–257. (doi:10.1002/ar.23944)

87. Cieri RL. 2018 The axial anatomy of monitor lizards (Varanidae). *J. Anat.* **233**, 636–643.
(doi:10.1111/joa.12872)

88. Nesbitt SJ. 2011 The Early Evolution of Archosaurs: Relationships and the Origin of Major
Clades. *Bull. Am. Mus. Nat. Hist.* **2011**, 1–292. (doi:10.1206/352.1)

89. Barbadillo LJ, Martínez-Solano I. 2002 Vertebral Intercentra in Lacertidae: Variation and
Phylogenetic Implications. *Copeia* **2002**, 208–212.

90. Hoffstetter RJ. 1969 Vertebrae and ribs of modern reptiles. In: Gans, Bellairs and Parsons,
Biology of the Reptilia. , 201–310.

91. Goloboff PA, Catalano SA. 2016 TNT version 1.5, including a full implementation of
phylogenetic morphometrics. *Cladistics* **32**, 221–238. (doi:10.1111/cla.12160)

92. de Queiroz K, Gauthier JA. 2020 Squamata. In *Phylonyms: a companion to the PhyloCode*,
pp. 1093–1101. CRC Press.

93. Nesbitt SJ, Stocker MR, Chatterjee S, Horner JR, Goodwin MB. 2021 A remarkable group of
thick-headed Triassic Period archosauromorphs with a wide, possibly Pangean distribution.
J. Anat. **239**, 184–206. (doi:10.1111/joa.13414)

94. Winchester L, Bellairs A d'A. 1977 Aspects of vertebral development in lizards and snakes.
J. Zool. **181**, 495–525. (doi:10.1111/j.1469-7998.1977.tb03259.x)

95. Brochu CA. 1996 Closure of neurocentral sutures during crocodilian ontogeny: Implications
for maturity assessment in fossil archosaurs. *J. Vertebr. Paleontol.* **16**, 49–62.
(doi:10.1080/02724634.1996.10011283)

96. Griffin CT, Stocker MR, Colleary C, Stefanic CM, Lessner EJ, Riegler M, Formoso K,
Koeller K, Nesbitt SJ. 2021 Assessing ontogenetic maturity in extinct saurian reptiles. *Biol.*
Rev. **96**, 470–525. (doi:10.1111/brv.12666)

Supplementary Figures

Figure 1. Comparison of jugal morphologies among living lepidosaurs. Note the variability in
the presence and development of the posterior process, as well as the presence of an ossified
jugal itself. CT scan images are from digimorph.org.

Figure 2. Comparison of palatine morphologies. Blue shading indicates choanal fossa. Top

image of †*Cryptovaranoides* referred left palatine is from Whiteside et al. [19]. Middle is the left

palatine of †*Helioscopos dickersonae* (Squamata: Pan-Gekkota) from the Late Jurassic Morrison

Formation [62]. Bottom is the right palatine of †*Eoscincus ornatus* (Squamata: Pan-Scincoidea)

from the Late Jurassic Morrison Formation [31].

Prominent choanal fossa on anterior margin of ventral surface of palatine

| 0: absent | 1: present, limited to anterior third of palatine | 2: present and extending over halfway along palatine | considered "prominent" by WEA24 |

Sphenodon punctatus
after Gauthier et al. (2012)

†*Helioscopos dickersonae*
after Meyer et al. (2023)

†*Eoscincus ornatus*
after Brownstein et al. (2022)

†*Cryptovaranoides*
Holotype

984 **Figure 3.** Choanal fossa character scoring.

988 **Figure 4.** Comparison of vomer morphologies. Top image is †*Eoscincus ornatus* from

989 Brownstein et al. (2022). Note that the rows of vomerine teeth are posteriorly placed, and the

990 vomerine ridges are large and laterally placed. Bottom image of †*Cryptovaranoides* holotype is

991 from Whiteside et al. [19].

Cephalic head of (mobile) quadrate with notch for the squamosal, peg-in-notch articulation

0: absent   1: present   considered "present" by WEA24

feature WEA24 identify as notch

quadrate
squamosal

notch

Sphenodon punctatus
after Gauthier et al. (2012)

Amphiglossus splendidus
after Gauthier et al. (2012)

†*Cryptovaranoides*
Holotype

995 **Figure 5.** Quadrate-squamosal articulation character scoring.

Vomer and maxilla meet at anterior margin of fenestra exochoanalis

0: absent   1: present   considered "present" by WEA24

····· damaged

vomer
premaxilla
maxilla

Sphenodon punctatus
after Gauthier et al. (2012)

Pogona vitticeps
after Gauthier et al. (2012)

†*Cryptovaranoides*
Holotype

999 **Figure 6.** Anterior articulation of vomer and maxilla character scoring.

1000

1001

Celestus enneagrammus
after Gauthier et al. (2012)

position of foramen
according to WEA24

†*Cryptovaranoides*
Holotype

Figure 7. Presence of a medially positioned posterior mylohyoidal foramen on the mandible. As shown, there is no identifiable foramen on the mandible of †*Cryptovaranoides*. Bottom image of †*Cryptovaranoides* holotype skull is from Whiteside et al. [19].

†*Cryptovaranoides*
Holotype

†*Peurcosuchus traverorum*
after Marsh et al. (2022)

1013

Figure 8. Comparison of the humerus of †*Cryptovaranoides microlanius* (from Whiteside et al. [19]) to †*Puercosuchus traverorum* (from [24]). Left image of †*Cryptovaranoides* holotype humerus is from Whiteside et al. [19]. Ent.for, entepicondylar foramen, ect.for, ectepicondylar foramen, tr.cv, trochlear cavity, cap, capitellum, cbf, cuboid fossa, g, groove.

1018

1019

1020

1021 **Table 1.**

1022

Character	Number in Talanda et al. (2022)	Study Cited in Whiteside et al. (2024)	Synapomorphy of	Unambiguous Optimization

Cephalic head of (mobile) quadrate with notch for the squamosal, peg-in-notch articulation with rod-shaped squamosal	123	de Queiroz and Gauthier (2020)	Pan-Squamata	1→0
Vomer and maxilla meet at anterior margin of fenestra exochoanalis	371	Gauthier et al. (2012)	N/A	N/A
Prominent choanal fossa on anterior margin of ventral surface of palatine	100	Gauthier et al. (2012)	Lepidosauria	N/A
Subdivision of embryonic metotic fissure by the crista tuberalis into vagus (jugular) foramen and recessus scala tympani	382	Simões et al. (2018); de Queiroz and Gauthier (2020)	N/A	N/A
No quadrate foramen	118	Gauthier et al. (2012)	Lepidosauria	1→0
Medially positioned posterior mylohyoidal foramen on mandible	163	Gauthier et al. (2012)	N/A	N/A
Fusion of exoccipitals and opisthotics forming an otoccipital	151	Gauthier et al. (2012); de Queiroz and Gauthier (2020)	N/A	N/A
Trunk vertebrae lack intercentra	237	de Queiroz and Gauthier (2020)	Pan-Unidentata	1→0

1023

1024

1025

1026

1027