Cryptovaranoides is not a squamate

Michael W. Caldwell,[1,2*] Chase D. Brownstein,[3,4] Dalton L. Meyer,[5] Simon G. Scarpetta[6,7], Michael S.Y. Lee[8,9], Tiago R. Simões[10]

[1]Department of Biological Sciences, University of Alberta, Edmonton, AB, Canada

[2]Department of Earth and Atmospheric Sciences, University of Alberta, Edmonton, AB, Canada

[3]Department of Ecology and Evolutionary Biology, Yale University, New Haven CT

[4]Stamford Museum and Nature Center, Stamford, CT

[5] Department of Earth and Planetary Sciences, Yale University, New Haven CT

[6] Museum of Vertebrate Zoology, Department of Integrative Biology, University of California, Berkeley, CA

[7]Department of Environmental Science, University of San Francisco, San Francisco, CA

[8]College of Science and Engineering, Flinders University, Adelaide 5001, Australia

[9]Earth Sciences Section, South Australian Museum, North Terrace, Adelaide 5000, Australia.

[10]Department of Ecology and Evolutionary Biology, Princeton University, Princeton NJ, 08544, USA

*Corresponding author: mw.caldwell@ualberta.ca

Abstract.

Accurate reconstruction of the timescale of organismal evolution requires knowledge of the placement of extinct representatives among living branches. The fossil record has the capacity to reframe hypotheses of organismal evolution by producing representatives of clades that predate previously known fossils or node ages. Recently, one fossil with the potential to drastically change current consensus surrounding the timescale of reptile diversification was described from Triassic fissure-fill deposits in the United Kingdom. This taxon, *Cryptovaranoides microlanius*, was originally placed deep within the squamate crown clade, suggesting that many lineages of lizards and snakes must have appeared by the Triassic and implying long ghost lineages that paleontologists and molecular phylogeneticists have failed to detect using all other available data. Our team questioned this identification and instead suggested *Cryptovaranoides* had unclear affinities to living reptiles, but this alternative interpretation was again challenged by the

team that originally described this species. Here, we dissect the morphological character codings used to support a crown squamate affinity for *Cryptovaranoides microlanius* and illustrate empirical problems with analyses that hypothesize this taxon to be a crown squamate. Our analyses emphasize the importance of stringency in constructing hypodigms of fossils, particularly when they may be key for proper time calibration of the Tree of Life.

Keywords: *Cryptovaranoides,* Triassic, fossil, Squamata, Archosauromorpha.

Introduction.

Paleontology has found a new role in the era of widespread genome sequencing of living phylogenetic diversity: providing justification for the placement of fossil calibrations along molecular phylogenies [1–6]. Placing fossils on the Tree of Life has become common practice, and new discoveries [7–10] and reshuffling of hypothesized phylogenetic relationships [11] often revise which fossils are best to use as prior constraints. These factors make the robust placement of potential fossil calibrations essential for calibrating phylogenies to absolute time, which themselves are a foundation of modern evolutionary biology [12].

Recently, Whiteside et al. [13], described *Cryptovaranoides microlanius* based on a partially articulated skeleton and a collection of referred material from the Carnian [14] to Norian-Rhaetian [13,15–17] (237-201.5 million years ago) fissure fill deposits of England, UK. In a subsequent study, we [18] refuted the affinities of *C. microlanius* to Anguimorpha as a deeply nested crown squamate that was proposed by Whiteside et al. [13] based on a re-examination of the CT scan data of the holotype and referred specimens. This prompted substantial edits to how this taxon was scored in the morphological data matrix used to assess its phylogenetic relationships, finding *C. microlanius* to be "…*either an archosauromorph or an indeterminate neodiapsid…*" and not a lepidosaur, much less a crown squamate. In an impassioned response to our study, Whiteside et al. [19] disagreed with many of our anatomical observations and restated their position on the affinities of *C. microlanius.* Whiteside et al. [19] also referred additional Late Triassic fossils to *Cryptovaranoides microlanius* and presented putative phylogenetic results that this taxon is a crown group squamate (squamate hereafter).

Here we provide point-by-point re-examination of the new evidence and interpretations of Whiteside et al. [19] regarding the anatomy and phylogenetic affinities of *C. microlanius*. We identify and describe what we consider to be major methodological errors in the comparative anatomical work and phylogenetic analyses of Whiteside et al. [19], and present the results of our reanalysis of their actual data matrices and the recovered synapomorphies ignored by those authors.

Results.

Errors and Untested Synapomorphies

Both Whiteside et al. [13] and Whiteside et al. [19] include numerous comparative anatomy errors that can broadly be grouped into two categories: (i) anatomical interpretation, and (ii) translating these interpretations into scorings in different phylogenetic datasets. In Brownstein et al. [18], we identified 22 errors in the study of Whiteside et al. [13] of type (i) ("Results" in Brownstein et al. [18]) and several more of type (ii) ("Supplementary Material" in Brownstein et al. [18]). In Whiteside et al. [19], the authors claim that there were five observational errors in Brownstein et al. [18]. Four of these are supposed observational errors (type (i)) and one concerns a difference in how Brownstein et al. [18] and Whiteside et al. [19] score a character (type (ii)). In turn, this implies that Whiteside et al. [19] acknowledge that the other 18 type (i) errors produced by their previous study were correctly identified by Brownstein et al. [18]. Yet, Whiteside et al. [19] discuss additional characters in sections of their study and provide different interpretations of the anatomy of *Cryptovaranoides microlanius* than do Brownstein et al. [18], but without clear justification.

In this section we also revisit each alternative interpretation of the anatomy of *Cryptovaranoides microlanius* provided by Whiteside et al. [19], who discussed 26 characters that they suggest have bearing on the placement of this taxon within Lepidosauria, Pan-Squamata, crown Squamata, and successive clades within the crown. We note first that these characters do not correspond to optimized character states in the phylogenies that Whiteside et al. [13] and WEA24 inferred, but instead to an assemblage of character state optimizations presented in various papers in the literature (e.g., [20–22]). This is not an issue *per se* for a referral of *C. microlanius* to Squamata, but it does mean that these characters are of unclear relevance to the actual support for the position(s) of *C. microlanius* among reptiles that Whiteside et al. [13] and Whiteside et al. [19] recovered in the phylogenies that they presented. With this noted, we provide a point-by-point discussion of character interpretations in Brownstein et al. [18] that were challenged by Whiteside et al. [19].

Entepicondylar and ectepicondylar foramen of humerus. The features on the humeri that Whiteside et al. [19] figure and claim are the ente- and ectepicondylar foramina, are in fact, fossae that are filled in with sediment. Although they claim to observe this in yet another referred

humerus, they neither figure the internal structure of the bone nor provide any evidence for its

referral to *Cryptovaranoides microlanius*. Contrary to the assertion of Whiteside et al. [19], CT

scans do in fact provide essential information about the structure of these fossae and show that,

when infill is removed, foramina are absent (Figure 4 in Brownstein et al. [18]). Further, these

structures are in the wrong place to be the claimed foramina. In all diapsids that possess these

foramina, they are placed low on the anterior surface (entepicondylar) and high on the distal

surface (ectepicondylar) of the humerus (e.g., see [8,21] for examples in squamates, [23] for

turtles, and [24,25] for sphenodontians) and are dissimilar in shape and size (the entepicondylar

foramen is elongated along the long axis of the humerus; the ectepicondylar foramen is circular).

The features that Whiteside et al. [19] figure are similar in shape and size, are placed on the same

side of the bone, and differ in placement from humeral foramina observed in other exemplar

fossils and living reptile species [8,21,23–25]. The morphology of the fossae observed on the

distal end of the humeri referred to *C. microlanius* by Whiteside et al. [19] are, however, similar

in placement, size, and shape to fossae described on the distal ends of the humeri of some

archosauromorphs, including the azendohsaurid *Puercosuchus traverorum* (Figure 8; also see

fig.13a in [26]). We further note that, even if WEA24's interpretation was accurate, the

entepicondylar foramen is always absent in all crown squamates [8,21,27]. Among lepidosaurs,

the presence of both entepicondylar and ectepicondylar foramina is found only among

sphenodontians, stem lepidosaurs, and stem squamates [8,28]. Further, these foramina are also

present in many non-lepidosaurian reptiles, including captorhinids (e.g. *Captorhinus*),

younginiforms (e.g., *Hovasaurus* and *Youngina*), Claudiosaurus, Acleistorhinidae

(Delorhynchus), *Mesosaurus*, the possible stem turtle *Eunotosaurus africanus*, and in some

sauropterygians (e.g., *Serpianosaurus* and *Lariosaurus*)[8,29,30]. If anything, the presence of

both foramina would support the assignment of *Cryptovaranoides* as outside of crown Squamata,

not within the clade nor as a squamate synapomorphy. The position taken by Whiteside et al. [13]

and Whiteside et al. [19] on this feature is puzzling.

Absence of jugal posterior process. WEA24 state: "*Except for a very few fossil taxa, including*

two polyglyphanodontians,Tianyusaurus and Polyglyphanodon, … squamates lack a posterior

process on the jugal…". This is entirely incorrect. The two extinct taxa noted by WEA24 here

are only unusual among squamates in the fact that they have a complete lower temporal bar, and

thus an elongated jugal posterior process [21,31–33]. However, most families of squamates

include species with a jugal posterior process (Figure 1)[8,21,22,32,34], even if a complete lower

temporal bar is not present. The high degree of observed variability in the development of the

jugal posterior process and lower temporal bar (e.g., Figure 1) means that the placement of

Cryptovaranoides microlanius within any pan-lepidosaur clade based on this feature should be

viewed with caution. The development and enclosure of the temporal fenestra in reptiles has

been linked to the expression of two genes, *Runx2* and *Msx2*, in *in vivo* studies [35]. We suspect

that the variable extent of this feature in many early-diverging lepidosaur and archosauromorph

lineages might be an example of the 'zone of variability' [36], in which the canalization of

development and other constraints (e.g., functionality; see [37]) had not yet completely acted to

'fix' the morphology of the posterior process. This hypothesis will of course require further

experimental study of living model systems. Together, these observations suggest that the

presence of a jugal posterior process was incorrectly scored in the datasets used by WEA24 (type

(ii) error.

Anterior emargination of the maxillary nasal process. Whiteside et al. [19] state that Brownstein

et al. [18] *"seemingly relied on the anteriorly broken left maxilla"* and that "*The anterior margin*

of the maxillary nasal process of the right maxilla tapers anteriorly…with no evidence of the type

of emargination suggested". The character in question (ch. 18 in the dataset used by Whiteside et

al. [19]), is one of the original characters from [8], which is the basis for the dataset used by

Whiteside et al. [19] [38]. In both datasets, this character is described as "*Maxillae, **posterior***

***emargination**, between nasal and orbital processes*"[boldface added], and this character is

extensively described in [8]. Therefore, Whiteside et al. [19] incorrectly assessed the anterior

margin of the maxilla instead of the posterior margin, which is another type (ii) error.

Expanded radial condyle of the humerus. Whiteside et al. [19] (p. 3) state that Whiteside et al.

[13] "*noted an expanded radial condyle on the humerus.*" Yet, nowhere in Whiteside et al. [13]

do the authors mention or figure such a structure. Whiteside et al. [19] (p.3, fig. 1b) write in their

figure caption regarding another isolated fragment: *"(b) NHMUK PV 38911 isolated larger*

specimen of the distal end of left humerus of Cryptovaranoides microlanius *in (above) anterior*

*and (below) posterior views **showing similar features except the condyle of the capitellum**."*

(boldface added). Then, in Whiteside et al. [19] (subsection 2.3), the authors state that this

fragment does not have a radial condyle/capitellum of the humerus, which they defend as

follows: "*This is reinforced by the larger humerus (figure 1b) which is missing the condyle, **as is***

typical in the preservation of fissure lepidosaurs** (e.g. Clevosaurus; [12, fig. 29b]) but the **cavity

***in which it sat clearly indicates a substantial condyle in life.*"** (boldface added). What WEA24

highlight is that there is no condyle in this humerus or the holotype specimen and defer it to poor

preservation without justification beyond *"...as is typical ....of fissure lepidosaurs..."* as a reason

to conclude the condyle was present and to explain this inconsistency among the bones referred

to *Cryptovaranoides.* Taphonomy and poor preservation cannot be used to infer the presence of

an anatomical feature that is absent. Finally, even in the case of the isolated humerus with a

preserved capitulum, the condyle illustrated by Whiteside et al. [19] is fairly small compared to

even the earliest known pan-squamates, such as *Megachirella wachtleri* (Figure 4).

Preservation of the septomaxilla. Whiteside et al. [13] and Whiteside et al. [19] claim that a

small, disarticulated piece of bone that is preserved anterolateral to the vomer in the block

containing the holotype of *Cryptovaranoides microlanius* is the septomaxilla. They further

suggest that a portion of the medial surface on a maxilla referred to this taxon that is likely from

a much larger reptile provides additional support for the presence of a septomaxilla in *C.*

microlanius. Simply put, the CT scans published by Whiteside et al. [13] and Whiteside et al.

[19] show that the bone in the holotype is isolated and with no clear morphological affinities to

the septomaxillae in squamates (see CT scans in [21]). Whiteside et al. [19] also suggest that the

identity of this bone as the septomaxilla is supported by its placement between the maxilla and

premaxilla of the holotype but given the level of disarticulation of the holotype and the

morphology of the bone fragment, a similar argument could be made that it is a portion of the

anterior end of one of the vomers. In any case, we feel it is premature to code *C. microlanius* for

any character related to the morphology of the septomaxilla based on this disarticulated and

damaged bone fragment.

Expanded radial condyle of the humerus. Whiteside et al. [19] provided additional justification

of the presence of an expanded radial condyle of the humerus by stating that the projection of the

radial condyle above the adjacent region of the distal anterior extremity of *Cryptovaranoides*

microlanius supports their choice to score the expanded radial condyle as present. This is not the

condition specified in either of the morphological character sets that they cite [18,38] – the

presence of a distinct condyle that is expanded – and is by their own description not homologous

to the condition in other squamates.

Anterior emargination of the maxillary nasal process. Whiteside et al. [19] flagged our

interpretation of the anterior maxillary nasal process as emarginated, which we found united

Cryptovaranoides microlanius with archosauromorphs. Although we still interpret this state as

such based on the computed tomography scan data, we again note than none of our analyses [18]

unambiguously place *Cryptovaranoides microlanius* within Archosauromorpha and that, while

optimized as such, this single character necessarily has limited utility for placing *C. microlanius*

among reptiles.

Subdivision of the metotic fissure. Whiteside et al. [19] claim that the division of the metotic

fissure into the vagus foramen and recessus scala tympani by the crista tuberalis, a key squamate

feature, can be scored for *Cryptovaranoides microlanius*, yet paradoxically suggest the presence

of this condition is only inferable based on other observations of the anatomy of the holotype and

referred specimens. In fact, Whiteside et al. [19] argue that because another character relating to

a different part of the structure of the metotic fissure can be inferred based on the presence of the

crista tuberalis, that the division of the metotic fissure may also be inferred by the presence of the

crista tuberalis. This logic would imply that no reptile taxon should exist that possesses solely

either a crista tuberalis or a subdivided metotic fissure, even when this is an observed state

combination in squamates scored for the matrices that both our teams have used to analyze the

phylogenetic position of *Cryptovaranoides microlanius* [8,38]. Thus this inference lacks

justification. To verify that *C. microlanius* possessed a vagus foramen, Whiteside et al. [19] state

that they searched for isolated otoccipital fragments from the same locality and found an

otoccipital fragment with a vagus foramen that they refer to *C. microlanius* without other

justification. It is unclear to us how Whiteside et al. [19] accounted for confirmation bias when

conducting this collections search, or on what basis they refer this isolated bone to *C.*

microlanius. In any case, the presence of the lateral opening of the recessus scala tympani is not

figured in the fragment, and thus the division of the metotic fissure is not demonstrated in the

referred fragment.

Fusion of exoccipitals and opisthotics. As Whiteside et al. [19] note, we concur with their

identification of an otoccipital in *Cryptovaranoides microlanius* formed by the fusion of the

exoccipitals with the opisthotics. Our concern is with the use of this feature to assign *C.*

microlanius to Squamata. Whiteside et al. [19] cite de Queiroz and Gauthier [20], who list the

presence of an otoccipital as a distinguishing feature of squamates. However, Whiteside et al.

[19] fail to provide any phylogenetic evidence supporting the optimization of this feature as a

squamate synapomorphy in phylogenies including *C. microlanius*, nor do they recognize that an

otoccipital is present in numerous non-squamate reptiles, including numerous archosauromorphs

[26,39,40]. For these reasons, the presence of an otoccipital alone cannot be used to assign *C.*

microlanius to Squamata or even Lepidosauria instead of Archosauromorpha or other clades of

reptiles known from the Permo-Triassic, except to distinguish the turtle total clade

(*Eunotosaurus africanus* possesses unfused exoccipitals and opisthotics [41]). We acknowledge

that with respect to a character state widespread among reptiles, optimization along a phylogeny

is what is of primary importance for referring taxa to a particular clade. To this end, we reiterate

that Whiteside et al. [19] did not show that the presence of an otoccipital optimizes as an

ambiguous or unambiguous synapomorphy of Lepidosauria, Pan-Squamata, or Squamata in any

of their phylogenies. For characters that show evidence of homoplastic evolution like the

presence of an otoccipital (and indeed, the presence of a jugal posterior process; see above),

phylogenetic character optimization is essential.

Enclosed vidian canal exiting anteriorly at base of each basipterygoid process. In our restudy of

the holotype of *Cryptovaranoides microlanius*, we were unable to verify the presence of an

enclosed vidian canal exiting the sphenoid via the base of the corresponding basipterygoid

process. Whiteside et al. [19] take issue with our interpretation of this region of the braincase and

suggest that a larger, abraded, and isolated sphenoid that they refer to *C. microlanius* also

supports their interpretation of the anatomy of this region of the braincase. Yet, they stated that

this fragment is of limited informativeness and appear to agree with us that the best course of

action is to score this character as missing data when including *C. microlanius* in phylogenetic

analyses.

Development of the choanal fossa of the palatine. Squamata includes multiple instances where

the complexity of the bony palate increases dramatically, an innovation that is related to

chemosensory evolution and the integration of different bones that form this region of the skull

[21,21,22,34,42–46]. One feature recognized as phylogenetically informative is the development

of the choanal fossa on the ventral surface of the palatine (also known as the palatine sulcus)

[18,21,34,47]. Whiteside et al. [19] claim that the development of the palatine choanal fossa in

Cryptovaranoides microlanius is comparable to the development of this feature in living

squamates, and cite the living iguanian genus *Ctenosaura* as an example of a squamate with

similar anteroposterior development of this feature as *C. microlanius.* Importantly, most

iguanians appear to show the plesiomorphic condition where the choanal fossa is anteriorly

restricted on the palatine [21]; this feature has contributed to debates about whether Iguania

forms the living sister group of all other crown squamates (as found in many morphological

character-based phylogenies; [21,22]) or is deeply nested within the crown clade and secondarily

convergent with rhynchocephalians (as implied by phylogenies made using DNA sequence data

[48–54] and some made using morphological data [8]). As such, this comparison is not salient to

the discussion about whether the development of the choanal fossa in *C. microlanius* represents

the squamate condition. Secondly, Whiteside et al. [19] take issue with our contention that the

palatine fossa is present, albeit variably, across a wide swath of reptilian diversity, including

many early-diverging archosauromorph clades. They explicate this concern by distinguishing

between the narrow channel found in taxa like *Tanystropheus* (figure 11 in Whiteside et al. [19])

and the wider fossa found in *C. microlanius*. We agree that the fossa in *C. microlanius* is more

developed than in some early-diverging lepidosaurs, such as *Marmoretta oxoniensis* [55].

However, it is clear that the feature in *Cryptovaranoides* falls within the variation in the choanal

fossa length and depth (Figure 2; Figure 3) that represents the ancestral condition in lepidosaurs

[21] and is exemplified across archosauromorph (see, for example [26]) and indeed diapsid [56–

58] diversity. Finally, Whiteside et al. [19] cite the discovery of a large, isolated palatine that they

state confirms the presence of a "squamate-type" choanal fossa in *C. microlanius*, except they

provide no justification for why this isolated bone should be referred to this species. In any case,

the morphology of that bone is also unlike those of squamates (Figure 2).

Vomer ventral ridges and dentition. Whiteside et al. [19] reiterated their earlier characterization

[13] of the vomer of *Cryptovaranoides microlanius* as ridged, and suggest that the morphology

of the vomerine ridges and vomerine teeth in *C. microlanius* is comparable to the condition in the

anguid anguimorph *Pseudopus apodus*. We dispute this on the basis that the vomer of *C.*

microlanius as figured by Whiteside et al. [19] shows no clear ridges equivalent to those in

Pseudopus apodus [59] or other squamates with ridged vomers, including extinct forms such as

Eoscincus ornatus [34] (Figure 4). Instead, the new photograph of the holotype specimen of *C.*

microlanius provided in Whiteside et al. [19] shows that the vomer is indeed toothed, but no

ridges are figured or visible. We once again reexamined the CT scan data and failed to find any

structure resembling the ridges present in some anguimorphs, although we acknowledge that the

nature of the scan data (43 microns per segment) means that the vomer surface on the scan

segmentations is coarse. In any case, no ridges equivalent in shape or size to those found in

anguimorph squamates are present in the holotype of *C. microlanius.* The presence of teeth on

the vomer is itself rare among squamates; only in the pan-scincoid *Eoscincus ornatus* [34] and

some [59] anguid and varanoid [60] species are vomerine teeth documented. Whiteside et al.

[13,19] appear to suggest that the presence of vomerine ridges and a row of vomerine teeth

therefore allies *C. microlanius* with anguimorphs. The structure of the vomer in *C. microlanius*,

however, is fundamentally unlike those of anguimorph squamates, which are elongate and

possess pronounced ridges that are placed medially on the ventral surface of the main body of

each vomer and each house only one row of teeth along their posterior third [59,61]. In contrast,

the condition that Whiteside et al. [19] figure for *C. microlanius* shows multiple, parallel rows of

vomerine teeth on either side of the vomer that run across the entire length of the bone. This

morphology is even unlike that observed in the only squamate known where multiple vomerine

tooth rows are present, *Eoscincus ornatus* [34] (Figure 4). In that species, all vomerine teeth are

restricted to a raised surface at the posterior end of the ventral surface of the vomer main body

[34]. Thus, Whiteside et al. [19] have provided no evidence that vomerine ridges are present in

C. microlanius. We note again that vomerine teeth are widespread across neodiapsid diversity

outside Squamata [62] and appear in many lineages of Triassic archosauromorphs.

Lacrimal arches dorsally over lacrimal duct and floors lacrimal duct with medial process
posteriorly. We argued that this feature was unobservable in the holotype of *Cryptovaranoides*
microlanius [18]. Whiteside et al. [19] dispute this by suggesting that the CT scan data does not
reliably show the feature, and instead provide a photograph of the holotype skull that was stated
to show this morphology (figure 2a-c in [19]). This figure shows no curvature to the lacrimal,
which appears as a flat element, whereas an arch dorsally and a floor ventrally would indicate the
presence of a foramen. Whiteside et al. [19] did not code this feature (dorsal arcuation of the
lacrimal over the lacrimal duct, which is posteriorly floored by the medial process of the
lacrimal) for *C. microlanius*, but still use it to refer *C. microlanius* to Anguimorpha based on
optimization as a plesiomorphy of that clade in phylogenetic analyses that place *C. microlanius*
within Anguimorpha. This appears to suggest that, despite never conducting an analysis where
this feature is coded as 'present' for *C. microlanius*, Whiteside et al. [19] used the presence of the
feature to ally *C. microlanius* with Anguimorpha, not as a recovered synapomorphy, but stating
that it was so. This action would be an arbitrary unification of a species with a clade based on a
selected character state and would deny the equal possibility that this character state is absent
due to secondary reversal in *C. microlanius*. In sum, the use of lacrimal morphology to ally *C.*
microlanius with anguimorphs is apparently not based on direct character state optimization, i.e.,
a test of congruence.

Distinct quadratojugal absent. The quadratojugal cannot be located in the holotype of
Cryptovaranoides microlanius [13,19]. We argued that this might be due to postmortem
disarticulation and damage to the skull, and also noted that a complete ontogenetic series would
ideally be required to test whether the quadratojugal is modified throughout ontogeny [18].
Whiteside et al. [19] focused on our comment about the ideal situation of having an ontogenetic
series for *C. microlanius* to assess the development of the quadratojugal, which we restate would
be helpful to understand how this bone transforms through ontogeny given the complex
restructuring to this region of the skull that occurs throughout the evolution of lepidosaurs [63].
However, Whiteside et al. [19] state "we argue based on juvenile and adult specimens and the
absence of a quadratojugal facet on the quadrate." First, the region of the quadrate that would
articulate with the quadratojugal is not preserved in the holotype of *C. microlanius*, so it is

impossible to tell whether a facet for the quadratojugal is present. Second, Whiteside et al. [19]

fail to provide any justification for the referral of the isolated partial quadrate NHMUK PVR

37606 to *C. microlanius*. Third, Whiteside et al. [19] did not identify the quadratojugal facet on

this isolated quadrate when they figure this bone; indeed, based on the figure the process that

would have housed the quadratojugal facet is also missing from this quadrate (NHMUK PVR

37606). It is impossible to tell whether *C. microlanius* lacked a quadratojugal based on the

current data. Instead, all that can be said is that a quadratojugal is not preserved in the holotype

of *C. microlanius*, and the region housing the articular facet for the quadratojugal is not

preserved in either the holotype or referred quadrate.

Pterygoid/quadrate overlap. Whiteside et al. [19] restate the interpretation of Whiteside et al.

[13] that the pterygoid and quadrate have a short overlap in *Cryptovaranoides microlanius.*

Whiteside et al. [19] only compared the morphology of the quadrate in *Cryptovaranoides*

microlanius, which they agree is damaged, to the morphology present in rhynchocephalians, and

suggest based on this comparison that their interpretation is correct. We restate that this is not

possible to verify without more complete, articulated or semi-articulated palates assignable to

Cryptovaranoides microlanius based on apomorphic character states and combinations.

Fusion of the premaxillae and single median tooth. Whiteside et al. [19] suggest that Brownstein

et al. [18] incorrectly characterized the nature of fusion of the premaxillae into a single median

element in *Cryptovaranoides microlanius.* However, we reiterate that no justification has been

given in any paper [13,19] for referring these isolated premaxillae to *C. microlanius.* Whiteside

et al. [19] focus on differentiating these isolated large premaxillae from the premaxillae of two

other fissure fill lepidosaurs, *Gephyrosaurus bridensis* and *Diphydontosaurus avonis*, without

considering the possibility that additional taxa are present in the assemblage. Scoring fused

premaxillae as present for *C. microlanius* despite the presence of unfused, paired premaxillae in

the holotype defines (1) ontogenetic character state transformations solely based on the relative

size of the holotype and larger isolated bones referred without justification and (2) which

character state among those present in ontogeny is the phylogenetically informative one.

Defining both of these requires a robust ontogenetic series, which is not available for *C.*

microlanius at present. Observations of the development of living squamates also suggest that

these larger fused premaxillae should not be referred to *C. microlanius.* In crown squamates with
a median premaxilla, the premaxilla is invariably a single element upon first appearance very
early in embryonic development [64–68]. In contrast, the holotype of *C. microlanius*, which is
very clearly a juvenile and not an embryonic specimen, possesses paired premaxillae. The
ontogenetic series that Whiteside et al. [19] suggest for *C. microlanius* therefore differs from any
known squamate with a single median premaxilla.

Peg-in-notch articulation of quadrate with rod-shaped squamosal. Whiteside et al. [19] suggest
that both our teams are in agreement about the presence of a peg-in-notch articulation between
the quadrate and squamosal. We reiterate that we believe the presence of this type of articulation
is unclear based on the available data for the holotype, as these bones are both damaged in the
relevant sections and disarticulated (Figure 5).

Frontal underlaps parietal laterally on frontoparietal suture. Whiteside et al. [19] confirm that
the presence of this feature in *Cryptovaranoides microlanius* is based on the morphology of a
referred isolated frontal that they state matches the corresponding articular portion of the
prefrontal in the holotype. However, we do not understand how this bone could possibly match
the corresponding articular surface unless it was from the same individual or an animal of
exactly the same size. Unless Whiteside et al. [19] were actually able to articulate these bones, it
is unclear how this inference can be made. Finally, we note that the posterior process of the
prefrontal is broken off in the holotype of *Cryptovaranoides microlanius.* In squamates and other
lepidosaurs, this process abuts the lateral surface of the frontal along its anteroposterior axis.
Several of us (C.D.B. and D.L.M.) worked to rearticulated the holotype of *Eoscincus ornatus*,
which involved rearticulating the prefrontal and frontal. Rearticulating these bones is simply
impossible without the complete posterior process of the prefrontal, as the orientation of this
process must match the curvature of the lateral margin of the frontal. Because the isolated frontal
is not figured in either paper by Whiteside et al. [13,19], it is impossible for us to verify whether
this bone actually matches the corresponding articulation surface on the prefrontal in the
holotype. In any case, we regard it a best practice to exclude this isolated frontal from
discussions of the affinities of *C. microlanius.*

Medial process of the articular and prearticular. Whiteside et al. [19] revise their initial [13]

characterization of the medial process and refer to it as 'rudimentary,' scoring it as unobservable.

We determined that this process is absent [18] and offer no further comment besides that it

appears the distinction between our interpretation of this feature and that of Whiteside et al. [19]

is, as they explicate, largely arbitrary.

Bicapitate cervical ribs and cervical ribs with an anteriorly oriented process. Whiteside et al.

[19] took issue with our characterization of the morphology of the cervical ribs in

Cryptovaranoides microlanius, and compared the bicapitate morphology of the cervical ribs in

C. microlanius to two anguimorph squamates, *Pseudopus apodus* and *Varanus* spp. However, the

cervical ribs of *P. apodus* are not bicapitate, as shown by the very figure from [69] that Whiteside

et al. [19] cited. Similarly, the ribs of *Varanus* are unicapitate, not bicapitate, as shown in [70].

We assume that Whiteside et al. [19] referred to the morphology of the cervical rib head in *P.*

apodus because of the presence of a posterior process on the rib head [69]. This is not the

bicapitate condition and is not homologous with the morphology of the cervical rib heads

in either *C. microlanius* or archosauromorphs, where an anterior process is offset from the

process formed by the capitulum and tuberculum [39,71]. We reiterate that the condition in *C.*

microlanius is identical to that in archosauromorphs (figure 2c, figure 3 in [18]); the comparisons

made by Whiteside et al. [19] are across non-homologous structures and result from a

misinterpretation of cervical rib anatomy.

Cervical and dorsal vertebral intercentra. Whiteside et al. [19] suggest that we inferred the

presence of cervical and dorsal intercentra in *Cryptovaranoides microlanius*; however, we did

not propose this in our paper. In fact, Whiteside et al. [13] state in their original paper that there

"are gaps between the vertebrae indicating that intercentra were present (but displaced in the

specimen) on CV3 and posteriorly. Some images of bones on the scans are identified as

intercentra" (p. 11, [13]). The statement in [19] consequently represents an incorrect attribution

and an incorrect characterization of our reexamination, as we determined that no cervical and

dorsal intercentra were preserved or present in the holotype. Whiteside et al. [19] justify their

position that cervical intercentra are present in *C. microlanius* based on the morphology of

another isolated bone that they refer to this taxon. In any case, the absence of dorsal intercentra is

not a distinguishing feature of squamates because several lineages of squamates, including

lacertids, xantusiids, gekkotans, and the stem-squamate *Bellairsia gracilis* all possess cervical

and dorsal intercentra [38,72,73].

Anterior dorsal vertebrae, diapophysis fuses to parapophysis. Whiteside et al. [19] concur with

Brownstein et al. [18] that the diapophyses and parapophyses are unfused in the anterior dorsals

of the holotype of *Cryptovaranoides microlanius*, and restate that fusion of these structures is

based on the condition they observed in isolated vertebrae that they again refer to *C. microlanius*

without justification. As such, this feature should not be scored as present for *C. microlanius.*

Zygosphene–zygantrum in dorsal vertebrae. Whiteside et al. [19] again claim that the

zygosphene-zygantrum articulation is present in the dorsal series of *Cryptovaranoides*

microlanius based on the presence of 'rudimentary zygosphenes and zygantra' in isolated

vertebrae that they again refer to this species without justification. The structures that Whiteside

et al. [19] label as the zygosphenes and zygantra (figure 3 in [19]) are clearly not, however,

zygosphenes and zygantra, as the former is by definition a centrally located wedge-like process

that fits into the zygantrum, which is a fossa on the following vertebra. The structures labeled

zygosphenes and zygantra by Whiteside et al. [19] are the dorsal surfaces of the

prezygopophyses and the medial margins of the postzygapophyses.

Anterior and posterior coracoid foramina/fenestra. Whiteside et al. [19] use isolated coracoids

referred to *Cryptovaranoides microlanius* to support their claim that the upper 'fenestra'

(foramen) of the coracoid in the holotype specimen is indeed a foramen. Confusingly, Whiteside

et al. [19] label this feature as the 'primary coracoid fenestra' in their figure 3i-j, even though it is

clearly the coracoid foramen. Whiteside et al. [19] appear to have confused the coracoid

foramen, which is completely bounded by bone and placed inside the coracoid, with the coracoid

fenestra, which is bounded *in part* by the coracoid (the coracoid margin is curved to form part of

the bounding region in species with the coracoid fenestra) but also by the interclavicle (see

figures in [18,21]. The coracoid fenestra is not contained within the coracoid. This feature is also

not shown to be optimized as a pan-squamate synapomorphy in any phylogeny including

Cryptovaranoides microlanius [13,19], so its bearing on the identification of *C. microlanius* as a

pan-squamate is unclear to us. Finally, we note here that Whiteside et al. [19] appear to have

labeled a small piece of matrix attached to a coracoid that they refer to *C. microlanius* as the

supracoroacoid [*sic*] foramen in their figure 3, although this labeling is inferred because only

"suc, supracoroacoid [*sic*]" is present in their figure 3 caption.

Atlas pleurocentrum fused to axis pleurocentrum. Whiteside et al. [19] reiterate their claim that

the atlas and axis pleurocentra are present in *Cryptovaranoides microlanius* based on their

identification of an isolated, globose bone fragment as the atlas intercentrum, but provide no

additional justification for this identification and refer the reader to their original paper for a

figure of this bone [13], which is only visible on CT scans due to its entombment within the

matrix that includes the holotype. To this end, Whiteside et al. [19] revise their initial

interpretation of the what they identify as the preserved atlas-axis region, but again without any

justification or figures detailing how they reidentified a bone fragment that they had initially

believed was a cervical intercentrum as the atlas centrum and intercentrum 2. Thus, Whiteside et

[19] do not provide any additional description of how they reidentified these bones or a figure

illustrating their revised interpretation, and so we cannot comment on the strength of their

revised interpretation. Again, we note that we were not able to identify the morphology of the

atlas and axis with any confidence in the holotype of *C. microlanius*, and so we again believe any

relevant characters should be scored as missing data for this taxon.

Midventral crest of presacral vertebrae. Whiteside et al. [19] appear to agree with our assertion

that a midventral crest is not present on the presacral vertebrae [18]. We never challenged their

scoring of keels on the caudal centra as missing data. Rather, we stated that keels are clearly

present on the cervical vertebrae [18].

Angular does not extend posteriorly to reach articular condyle. Whiteside et al. [19] state that

the posterior portion of the angular is present medially on the right mandible in the holotype, and

provide an interpretation of the anatomy of this region. They cite figure 3a in Whiteside et al.

[19], but this shows the mandible in lateral view and only a small portion of the angular is

visible. As such, it is not clear to us what they interpret to be the posterior portion of the angular.

In any case, we could not identify anywhere on the CT scans or on the accessible portions of the

real specimen (e.g., figure 2b in [19]) the feature that Whiteside et al. [19] consider to be the

posterior angular extent. Whiteside et al. [19] describe the posterior extent of the angular as a

'contoured feature,' but we are entirely unclear about what this actually describes.

Ulnar patella. The only disagreement between our interpretation [18] and that made by

Whiteside et al. [13,19] concerns whether the ulnar patella is absent due to the ontogenetic state

or preservation status of the holotype [13,19] or an ontogenetically invariable feature of the

anatomy of *Cryptovaranoides microlanius* [18]; we suggested the latter based on our observation

that the forelimb of the holotype is articulated and mostly complete.

Overarching Empirical Problems in Whiteside et al. [19].

Unassignable specimens and "hypodigm inflation." Whiteside et al. [13] and Whiteside et al.

[19] describe and defend their addition of isolated elements to fill in gaps in the holotype concept

of *Cryptovaranoides* to build, in our view, an unjustifiably inflated hypodigm. WEA24 (p. 15)

try to ameliorate this problem in their Section (5.5) where they state that: "*We emphasize that we*

always match isolated bones with their equivalents on the holotype..." and then state in

contradiction that: "*We consider it **remiss** not to include isolated bones as they provide details*

which no scan can. Furthermore, as the holotype is a juvenile, they give additional information

*from larger, and presumably older individuals, **on characters of the holotype otherwise***

***unavailable or uncertain**.*" (boldface added). In summary, Whiteside et al. [19] claimed to only

match isolated bones with equivalent ones in the holotype but also acknowledged use of

elements that are not comparable with the ones in the holotype. This lack of consistency is a

serious empirical issue.

Apomorphic characters not empirically obtained. Whiteside et al. [19] discussed nearly 30

anatomical characters (Sections 2-6), most of which are used to support their core hypothesis that

Cryptovaranoides is a squamate. Yet, none of these 30 characters or conditions are actually

found by Whiteside et al. [19] to be synapomorphies of Squamata in their various phylogenetic

analyses. In other words, they did not present the results of their Tests of Congruence, but rather

simply elected to argue that untested interpretations of anatomy were apomorphies,

synapomorphies, or shared characters.This is a non-trivial substantive methodological flaw in

Whiteside et al. [13], that was perpetuated in Whiteside et al. [19], and that nullifies the bulk of

the arguments presented repeatedly in both studies.

In Section 6, Whiteside et al. [19] (p. 16) indicate they modified the datasets of

Brownstein et al. [18] and [38], but preferred the results of [11] and performed two different

analyses based on [38]: one using only morphological data, and one using morphological data

with several constraints from a molecular backbone. The former was tested using maximum

parsimony and the latter using Bayesian inference. What is not clear from either Section 6 or 7,

is which phylogenetic analysis was used by WEA24 to "*review the apomorphy distribution*". But

this uncertainty is inconsequential as the apomorphy distributions discussed in Section 7 of

Whiteside et al. [19] were not derived from their phylogenetic analyses. Instead, Whiteside et al.

[19] (p. 19) exclusively extracted characters from the literature to support their identification of

Cryptovaranoides as a squamate, rather than basing this inference primarily on their own

phylogenetic analyses. These literature sourced characters include *"two diagnostic characters of*

Squamata (=crown-clade Squamata)" and *"...further eight squamate synapomorphies"* listed by

WEA24 (Section 7, p. 19). As stated by the authors, "*we give the citation for the squamate*

synapomorphy at the end of each character"— indicating that these diagnostic characters or

synapomorphies were picked from the literature and not derived from ancestral state

reconstructions based on their phylogenetic results.

550   In order to check the characters listed by Whiteside et al. [19] (p.19) as "*two diagnostic*

characters" and "*eight synapomorphies*" in support of a squamate identity for *Cryptovaranoides*,

we conducted parsimony analysis of the revised version of the dataset [38] provided by WEA24

in TNT v 1.5 [74]. We used Whiteside et al.'s [19] own data version—e.g., with (0) scored for

character 1 and not including character 383. We recovered eight apomorphies at the squamate

node of which only three were recovered as unambiguous synapomorphies (boldface indicates

the state as scored by Whiteside et al. [19]for *Cryptovaranoides*):

557   Ch. 138. Basisphenoid (or fused parabasisphenoid), ventral aspect, shape, concavity:

558   single (0)/divided (1)/ absent (**2**). 0 -> 2

559   Ch. 142. Prootics, alar crest: absent (0)/ present (**1**). 0->1

560   Ch. 347. Prefrontal/palatine antorbital contact: absent (0) / narrow forming less than 1/3

561   the transverse distance between the orbits (1) / contact broad, forming at least 1/2 the distance

562   between the orbits (**2**). 01->2

563   The eight features described by Whiteside et al. [19] (p.19) as synapomorphies for

Cryptovaranoides + Squamata were also not inferred at their recovered crown squamate node

from their own results. Rather, they were copied nearly verbatim from principally two sources

[21,75] and presented as if they were recovered as synapmorphies by Whiteside et al. [19] (p.

19). In Table 1, the italicized text is from Whiteside et al. [19] (p.19), including their literature

source for the supposed synapomorphy. The non-italicized text is the character number and state

from [11] as recovered by us from the TNT analysis conducted for this reassessment of

Whiteside et al. [19]. We also include where in our phylogeny the character state is recovered as

synapomorphic (Table 1).

Our reanalysis shows that Whiteside et al. [19] did not diagnose *Cryptovaranoides* +

crown Squamata based on synapomorphies found by their own analyses as there were only three

(see our results above). Instead, Whiteside et al. [19] provided ad hoc mischaracterizations of

diagnostic features of crown Squamata from studies that did not include *Cryptovaranoides* and

then discussed and reviewed synapomorphies of Squamata from these sources [21,75]. Because

each additional taxon has the possibility of inducing character reoptimization, an empirical

analysis of character optimization that includes a given new taxon is necessary to support

referring said taxon to any particular clade. If *Cryptovaranoides* was indeed a Triassic crown

squamate, it could plausibly show character distributions not previously sampled among known

members of the crown or stem group.

Discussion and Conclusions.

As we noted above and in Brownstein et al. [18], the anatomy of *Cryptovaranoides* is

similar to many Late Triassic neodiapsid reptiles. For example, the quadrates of

Cryptovaranoides are closely comparable to those of archosauromorphs such as *Prolacerta* [76],

Macronemus [77] and *Malerisaurus* [78], with which *Cryptovaranoides* shares the presence of a

notch on the quadrate for an immobile articulation with the squamosal. Furthermore, WEA24's

inference that the quadrate, the notch, and the articulation with the squamosal was mobile, i.e.,

streptostylic as in squamates, is unfounded.

Similarly, the vertebrae and cervical ribs of the holotype of *Cryptovaranoides* do not

resemble those of squamates, but in fact share important features with non-squamate

neodiapsids, especially archosauromorphs (a point left unaddressed by Whiteside et al. [19]). For

example, whereas fusion of the neural arches to the centra occurs during embryonic ossification

in squamates [79], in *Cryptovaranoides* unfused bony neural arches and centra are present, as

commonly observed in archosaurs and other non-lepidosauromorph neodiapsids [80,81]—we

provide a long-form review of these and other features in *Cryptovaranoides* that compare

favorably with non-squamate reptiles in Supplementary Material.

Several errors in Whiteside et al. [13] identified in Brownstein et al. [18], and subsequent

errors in Whiteside et al. [19] that were discussed above result in the incorrectly interpreted

neodiapsid and lepidosaur anatomy of *Cryptovaranoides* and consequently provide erroneous

scorings for *Cryptovaranoides* for morphological character matrices. Whiteside et al. [19] also

make additional errors on an alpha taxonomic level, such as the unjustified inflation of the

Cryptovaranoides hypodigm based on contradictory arguments pertaining to the assignment of

fragmentary and isolated elements to the holotype. Finally, we highlight here that where

Whiteside et al. [13] and Whiteside et al. [19] should have reported their own recovered

synapomorphies, that is, the actual results of their phylogenetic analyses, rather than listing a

substantive number of supposed apomorphies that the authors argue support the squamate

affinities of *Cryptovaranoides* broadly, and anguimorph affinities more specifically. On a

methodological and analytical level, we do not consider that manually selecting apomorphic

characters supporting their preferred placement of *Cryptovaranoides* instead of using characters

empirically obtained from phylogenetic ancestral state reconstruction of their phylogenetic

results, stands as valid support of a phylogenetic hypothesis.

We end with a perspective on the fossil record of neodiapsid evolution. The anatomies

and morphologies that diagnose crown group squamates are many and varied, and except for a

few features, hardly universally distributed amongst the living and fossil members of the crown.

They are themselves the product of some 250 million years of evolutionary time and would not

have evolved in a linear fashion. Rather, phylogenetic analyses of diverse extinct and living

reptile clades have shown that the squamate *bauplan* originated in the context of extensive

mosaic and homoplastic osteological character evolution [8,30,34,47,58]. We do not doubt that

members of Pan-Squamata were present during the Triassic; this is supported by the fossil record

[8] as well as numerous time-calibrated phylogenies based on genomic [48,49] morphological

[34,38,47] and total-evidence [8,30] data. However, Whiteside et al. [13] and Whiteside et al.

[19] have provided no direct osteological evidence that supports the presence of crown Squamata

or any of its inclusive clades in the Triassic.

References

1.   Inoue J, Donoghue PCJ, Yang Z. 2010 The Impact of the Representation of Fossil
Calibrations on Bayesian Estimation of Species Divergence Times. *Syst. Biol.* **59**, 74–89.
(doi:10.1093/sysbio/syp078)

2.   Heath TA, Huelsenbeck JP, Stadler T. 2014 The fossilized birth–death process for coherent
calibration of divergence-time estimates. *Proc. Natl. Acad. Sci.* **111**.
(doi:10.1073/pnas.1319091111)

3. Near TJ. 2004 Estimating divergence times of notothenioid fishes using a fossil-calibrated
molecular clock. *Antarct. Sci.* **16**, 37–44. (doi:10.1017/S0954102004001798)

4. Near TJ, Bolnick DI, Wainwright PC. 2005 Fossil calibrations and molecular divergence
time estimates in centrarchid fishes (Teleostei: Centrarchidae). *Evolution* **59**, 1768–1782.
(doi:10.1111/j.0014-3820.2005.tb01825.x)

5. Parham JF *et al.* 2012 Best Practices for Justifying Fossil Calibrations. *Syst. Biol.* **61**, 346–
359. (doi:10.1093/sysbio/syr107)

6. Pennington PT, Cronk QCB, Richardson JA, Near TJ, Sanderson MJ. 2004 Assessing the
quality of molecular divergence time estimates by fossil calibrations and fossil–based model
selection. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* **359**, 1477–1483.
(doi:10.1098/rstb.2004.1523)

7. Field DJ, Benito J, Chen A, Jagt JWM, Ksepka DT. 2020 Late Cretaceous neornithine from
Europe illuminates the origins of crown birds. *Nature* **579**, 397–401. (doi:10.1038/s41586-
020-2096-0)

8. Simões TR, Caldwell MW, Tałanda M, Bernardi M, Palci A, Vernygora O, Bernardini F,
Mancini L, Nydam RL. 2018 The origin of squamates revealed by a Middle Triassic lizard
from the Italian Alps. *Nature* **557**, 706–709. (doi:10.1038/s41586-018-0093-3)

9. Brownstein CD. 2023 Syngnathoid Evolutionary History and the Conundrum of Fossil
Misplacement. *Integr. Org. Biol.* **5**, obad011. (doi:10.1093/iob/obad011)

10. Velazco PM, Buczek AJ, Hoffman E, Hoffman DK, O'Leary MA, Novacek MJ. 2022
Combined data analysis of fossil and living mammals: a Paleogene sister taxon of Placentalia
and the antiquity of Marsupialia. *Cladistics* **38**, 359–373. (doi:10.1111/cla.12499)

11. Budd GE, Mann RP. 2023 Two Notorious Nodes: A Critical Examination of Relaxed
Molecular Clock Age Estimates of the Bilaterian Animals and Placental Mammals. *Syst.
Biol.* , syad057. (doi:10.1093/sysbio/syad057)

12. Smith SD, Pennell MW, Dunn CW, Edwards SV. 2020 Phylogenetics is the New Genetics
(for Most of Biodiversity). *Trends Ecol. Evol.* **35**, 415–425. (doi:10.1016/j.tree.2020.01.005)

13. Whiteside DI, Chambi-Trowell SAV, Benton MJ. 2022 A Triassic crown squamate. *Sci. Adv.*
8, eabq8274. (doi:10.1126/sciadv.abq8274)

14. Simms MJ, Drost K. 2024 Caves, dinosaurs and the Carnian Pluvial Episode: Recalibrating
Britain's Triassic karst 'fissures'. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **638**, 112041.
(doi:10.1016/j.palaeo.2024.112041)

15. Walkden GM, Fraser NC, Simms MJ. 2021 The age and formation mechanisms of Late
Triassic fissure deposits, Gloucestershire, England: Comments on Mussini, G. et al. (2020).
Anatomy of a Late Triassic Bristol fissure: Tytherington fissure 2. *Proc. Geol. Assoc.* **132**,
127–137. (doi:10.1016/j.pgeola.2020.10.006)

16. Chambi-Trowell S, Whiteside D, Benton M. 2019 Diversity in rhynchocephalian
Clevosaurus skulls basedon CT reconstruction of two Late Triassic speciesfrom Great
Britain. *Acta Palaeontol. Pol.* **64**. (doi:10.4202/app.00569.2018)

17. Butler RJ, Meade LE, Cleary TJ, McWhirter KT, Brown EE, Kemp TS, Benito J, Fraser NC.
2024 Hwiccewyrm trispiculum gen. et sp. nov., a new leptopleuronine procolophonid from
the Late Triassic of southwest England. *Anat. Rec.* **307**, 1390–1420. (doi:10.1002/ar.25316)

18. Brownstein CD, Simões TR, Caldwell MW, Lee MSY, Meyer DL, Scarpetta SG. 2023 The
affinities of the Late Triassic Cryptovaranoides and the age of crown squamates. *R. Soc.*
Open Sci. **10**, 230968. (doi:10.1098/rsos.230968)

19. Whiteside DI, Chambi-Trowell SAV, Benton MJ. 2024 Late Triassic Cryptovaranoides
microlanius is a squamate, not an archosauromorph. *R. Soc. Open Sci.* **11**, 231874.
(doi:10.1098/rsos.231874)

20. de Queiroz K, Gauthier JA. 2020 *Squamata*. CRC Press. See
http://repository.si.edu/xmlui/handle/10088/107192.

21. Gauthier JA, Kearney M, Maisano JA, Rieppel O, Behlke ADB. 2012 Assembling the
Squamate Tree of Life: Perspectives from the Phenotype and the Fossil Record. *Bull.*
Peabody Mus. Nat. Hist. **53**, 3–308. (doi:10.3374/014.053.0101)

22. Estes R, de Queiroz K, Gauthier J. 1988 Phylogenetic relationships within Squamata.
Phylogenetic Relatsh. Lizard Fam. , 119–281.

23. Hermanson G, Arnal FAM, Szczygielski T, Evers SW. 2024 A systematic comparative
description of extant turtle humeri, with comments on humerus disparity and evolution based
on fossil comparisons. *Anat. Rec.* **307**, 3437–3505. (doi:10.1002/ar.25450)

24. Simões TR, Kinney-Broderick G, Pierce SE. 2022 An exceptionally preserved Sphenodon-
like sphenodontian reveals deep time conservation of the tuatara skeleton and ontogeny.
Commun. Biol. **5**, 1–19. (doi:10.1038/s42003-022-03144-y)

25. DeMar Jr. DG, Jones MEH, Carrano MT. 2022 A nearly complete skeleton of a new
eusphenodontian from the Upper Jurassic Morrison Formation, Wyoming, USA, provides
insight into the evolution and diversity of Rhynchocephalia (Reptilia: Lepidosauria). *J. Syst.*
Palaeontol. **20**, 1–64. (doi:10.1080/14772019.2022.2093139)

26. Marsh AD, Parker WG, Nesbitt SJ, Kligman BT, Stocker MR. 2022 Puercosuchus
traverorum n. gen. n. sp.: a new malerisaurine azendohsaurid (Archosauromorpha:
Allokotosauria) from two monodominant bonebeds in the Chinle Formation (Upper Triassic,
Norian) of Arizona. *J. Paleontol.* **96**, 1–39. (doi:10.1017/jpa.2022.49)

27. Simões TR, Pyron RA. 2021 The Squamate Tree of Life. *Bull. Mus. Comp. Zool.* **163**, 47–95.
(doi:10.3099/0027-4100-163.2.47)

28. Simões TR, Caldwell MW, Pierce SE. 2020 Sphenodontian phylogeny and the impact of
model choice in Bayesian morphological clock estimates of divergence times and
evolutionary rates. *BMC Biol.* **18**, 191. (doi:10.1186/s12915-020-00901-5)

29. Hill RV. 2005 Integration of Morphological Data Sets for Phylogenetic Analysis of Amniota:
The Importance of Integumentary Characters and Increased Taxonomic Sampling. *Syst. Biol.*
54, 530–547. (doi:10.1080/10635150590950326)

30. Simões TR, Kammerer CF, Caldwell MW, Pierce SE. 2022 Successive climate crises in the
deep past drove the early evolution and radiation of reptiles. *Sci. Adv.* **8**, eabq1898.
(doi:10.1126/sciadv.abq1898)

31. Simões TR, Funston GF, Vafaeian B, Nydam RL, Doschak MR, Caldwell MW. 2016
Reacquisition of the lower temporal bar in sexually dimorphic fossil lizards provides a rare
case of convergent evolution. *Sci. Rep.* **6**, 24087. (doi:10.1038/srep24087)

32. Gao K, Norell MA. 2000 Taxonomic composition and systematics of Late Cretaceous lizard
assemblages from Ukhaa Tolgod and adjacent localities, Mongolian Gobi Desert. *Bull. Am.*
Mus. Nat. Hist. **2000**, 1–118. (doi:10.1206/0003-0090(2000)249<0001:TCASOL>2.0.CO;2)

33. Mo J-Y, Xu X, Evans SE. 2010 The evolution of the lepidosaurian lower temporal bar: new
perspectives from the Late Cretaceous of South China. *Proc. R. Soc. B Biol. Sci.*
(doi:10.1098/rspb.2009.0030)

34. Brownstein CD, Meyer DL, Fabbri M, Bhullar B-AS, Gauthier JA. 2022 Evolutionary
origins of the prolonged extant squamate radiation. *Nat. Commun.* **13**, 7087.
(doi:10.1038/s41467-022-34217-5)

35. Tokita M, Chaeychomsri W, Siruntawineti J. 2013 Skeletal gene expression in the temporal
region of the reptilian embryos: implications for the evolution of reptilian skull morphology.
SpringerPlus **2**, 336. (doi:10.1186/2193-1801-2-336)

36. Bever GS, Gauthier JA, Wagner GP. 2011 Finding the frame shift: digit loss, developmental
variability, and the origin of the avian hand. *Evol. Dev.* **13**, 269–279. (doi:10.1111/j.1525-
142X.2011.00478.x)

37. Schaerlaeken V, Herrel A, Aerts P, Ross CF. 2008 The functional significance of the lower
temporal bar in Sphenodon punctatus. *J. Exp. Biol.* **211**, 3908–3914.
(doi:10.1242/jeb.021345)

38. Tałanda M, Fernandez V, Panciroli E, Evans SE, Benson RJ. 2022 Synchrotron tomography
of a stem lizard elucidates early squamate anatomy. *Nature* **611**, 99–104.

39. Ezcurra MD. 2016 The phylogenetic relationships of basal archosauromorphs, with an
emphasis on the systematics of proterosuchian archosauriforms. *PeerJ* **4**, e1778.
(doi:10.7717/peerj.1778)

40. Pinheiro FL, De Simão-Oliveira D, Butler RJ. 2020 Osteology of the archosauromorph
Teyujagua paradoxa and the early evolution of the archosauriform skull. *Zool. J. Linn. Soc.*
189, 378–417. (doi:10.1093/zoolinnean/zlz093)

41. Bever GS, Lyson TR, Field DJ, Bhullar B-AS. 2015 Evolutionary origin of the turtle skull.
Nature **525**, 239–242. (doi:10.1038/nature14900)

42. Rieppel O, Kley NJ, Maisano JA. 2009 Morphology of the skull of the white-nosed
blindsnake, Liotyphlops albirostris (Scolecophidia: Anomalepididae). *J. Morphol.* **270**, 536–
557. (doi:10.1002/jmor.10703)

43. Lee MSY. 1997 The phylogeny of varanoid lizards and the affinities of snakes. *Philos. Trans.*
R. Soc. Lond. B. Biol. Sci. **352**, 53–91. (doi:10.1098/rstb.1997.0005)

44. Abramyan J, Richman JM. 2015 Recent insights into the morphological diversity in the
amniote primary and secondary palates. *Dev. Dyn.* **244**, 1457–1468.
(doi:10.1002/dvdy.24338)

45. Strong CRC, Scherz MD, Caldwell MW. 2022 Convergence, divergence, and
macroevolutionary constraint as revealed by anatomical network analysis of the squamate
skull, with an emphasis on snakes. *Sci. Rep.* **12**, 14469. (doi:10.1038/s41598-022-18649-z)

46. Watanabe A, Fabre A-C, Felice RN, Maisano JA, Müller J, Herrel A, Goswami A. 2019
Ecomorphological diversification in squamates from conserved pattern of cranial integration.
Proc. Natl. Acad. Sci. **116**, 14688–14697. (doi:10.1073/pnas.1820967116)

47. Meyer D, Brownstein CD, Jenkins KM, Gauthier JA. 2023 A Morrison stem gekkotan
reveals gecko evolution and Jurassic biogeography. *Proc. R. Soc. B*
(doi:10.1098/rspb.2023.2284)

48. Title PO *et al.* 2024 The macroevolutionary singularity of snakes. *Science* **383**, 918–923.
(doi:10.1126/science.adh2449)

49. Burbrink FT *et al.* 2020 Interrogating Genomic-Scale Data for Squamata (Lizards, Snakes,
and Amphisbaenians) Shows no Support for Key Traditional Morphological Relationships.
Syst. Biol. **69**, 502–520. (doi:10.1093/sysbio/syz062)

50. Townsend TM, Larson A, Louis E, Macey JR. 2004 Molecular Phylogenetics of Squamata:
The Position of Snakes, Amphisbaenians, and Dibamids, and the Root of the Squamate Tree.
Syst. Biol. **53**, 735–757. (doi:10.1080/10635150490522340)

51. Pyron RA, Burbrink FT, Wiens JJ. 2013 A phylogeny and revised classification of Squamata,
including 4161 species of lizards and snakes. *BMC Evol. Biol.* **13**, 93. (doi:10.1186/1471-
2148-13-93)

52. Streicher JW, Wiens JJ. 2017 Phylogenomic analyses of more than 4000 nuclear loci resolve
the origin of snakes among lizard families. *Biol. Lett.* **13**, 20170393.
(doi:10.1098/rsbl.2017.0393)

53. Zheng Y, Wiens JJ. 2016 Combining phylogenomic and supermatrix approaches, and a time-
calibrated phylogeny for squamate reptiles (lizards and snakes) based on 52 genes and 4162
species. *Mol. Phylogenet. Evol.* **94**, 537–547. (doi:10.1016/j.ympev.2015.10.009)

54. Singhal S, Colston TJ, Grundler MR, Smith SA, Costa GC, Colli GR, Moritz C, Pyron RA,
Rabosky DL. 2021 Congruence and Conflict in the Higher-Level Phylogenetics of Squamate
Reptiles: An Expanded Phylogenomic Perspective. *Syst. Biol.* **70**, 542–557.
(doi:10.1093/sysbio/syaa054)

55. Griffiths EF, Ford DP, Benson RBJ, Evans SE. 2021 New information on the Jurassic
lepidosauromorph *Marmoretta oxoniensis*. *Pap. Palaeontol.* **7**, 2255–2278.
(doi:10.1002/spp2.1400)

56. Dudgeon TW, Maddin HC, Evans DC, Mallon JC. 2020 The internal cranial anatomy of
Champsosaurus (Choristodera: Champsosauridae): Implications for neurosensory function.
Sci. Rep. **10**, 7122. (doi:10.1038/s41598-020-63956-y)

57. Brownstein CD. 2022 High morphological disparity in a bizarre Paleocene fauna of
predatory freshwater reptiles. *BMC Ecol. Evol.* **22**, 34. (doi:10.1186/s12862-022-01985-z)

58. Jenkins KM, Foster W, Napoli JG, Meyer DL, Bever GS, Bhullar B-AS. In press. Cranial
anatomy and phylogenetic affinities of , with new information on the unique bolosaurid
feeding apparatus and evolution of the impedance-matching ear. *Anat. Rec.* **n/a**.
(doi:10.1002/ar.25546)

59. Klembara J, Dobiašová K, Hain M, Yaryhin O. 2017 Skull Anatomy and Ontogeny of
Legless Lizard Pseudopus apodus (Pallas, 1775): Heterochronic Influences on Form. *Anat.*
Rec. **300**, 460–502. (doi:10.1002/ar.23532)

60. Yi H-Y, Norell MA. 2013 New Materials of Estesia mongoliensis (Squamata: Anguimorpha)
and the Evolution of Venom Grooves in Lizards. *Am. Mus. Novit.* **2013**, 1–31.
(doi:10.1206/3767.2)

61. 2021 The anatomy, phylogenetic relationships, and autecology of the carnivorous lizard
"Saniwa" feisti Stritzke, 1983 from the Eocene of Messel, Germany. *Comptes Rendus*
Palevol **20**, 441–506.

62. Matsumoto R, Evans SE. 2017 The palatal dentition of tetrapods and its functional
significance. *J. Anat.* **230**, 47–65. (doi:10.1111/joa.12534)

63. Ford DP, Evans SE, Choiniere JN, Fernandez V, Benson RBJ. 2021 A reassessment of the
enigmatic diapsid Paliguana whitei and the early history of Lepidosauromorpha. *Proc. R.*
Soc. B Biol. Sci. **288**, 20211084. (doi:10.1098/rspb.2021.1084)

64. Khannoon ER, Evans SE. 2020 Embryonic skull development in the gecko, Tarentola
annularis (Squamata: Gekkota: Phyllodactylidae). *J. Anat.* **237**, 504–519.
(doi:10.1111/joa.13213)

65. Hernández-Jaimes C, Jerez A, Ramírez-Pinilla MP. 2012 Embryonic development of the
skull of the Andean lizard Ptychoglossus bicolor (Squamata, Gymnophthalmidae). *J. Anat.*
221, 285–302. (doi:10.1111/j.1469-7580.2012.01549.x)

66. Skawiński T, Skórzewski G, Borczyk B. 2021 Embryonic development and perinatal
skeleton in a limbless, viviparous lizard, Anguis fragilis (Squamata: Anguimorpha). *PeerJ* **9**,
e11621. (doi:10.7717/peerj.11621)

67. Skawiński T, Kaczmarek P, Borczyk B. 2023 Embryonic development of the skull in a
parthenogenetic lizard, the mourning gecko (Lepidodactylus lugubris). *J. Anat.* **243**, 618–
629. (doi:10.1111/joa.13871)

68. Ollonen J, Da Silva FO, Mahlow K, Di-Poï N. 2018 Skull Development, Ossification
Pattern, and Adult Shape in the Emerging Lizard Model Organism Pogona vitticeps: A
Comparative Analysis With Other Squamates. *Front. Physiol.* **9**.
(doi:10.3389/fphys.2018.00278)

69. Čerňanský A, Yaryhin O, Ciceková J, Werneburg I, Hain M, Klembara J. 2019 Vertebral
Comparative Anatomy and Morphological Differences in Anguine Lizards With a Special
Reference to Pseudopus apodus. *Anat. Rec.* **302**, 232–257. (doi:10.1002/ar.23944)

70. Cieri RL. 2018 The axial anatomy of monitor lizards (Varanidae). *J. Anat.* **233**, 636–643.
(doi:10.1111/joa.12872)

71. Nesbitt SJ. 2011 The Early Evolution of Archosaurs: Relationships and the Origin of Major
Clades. *Bull. Am. Mus. Nat. Hist.* **2011**, 1–292. (doi:10.1206/352.1)

72. Barbadillo LJ, Martínez-Solano I. 2002 Vertebral Intercentra in Lacertidae: Variation and
Phylogenetic Implications. *Copeia* **2002**, 208–212.

73. Hoffstetter R, Jean. 1969 Vertebrae and ribs of modern reptiles. In: Gans, Bellairs and
Parsons, Biology of the Reptilia. , 201–310.

74. Goloboff PA, Catalano SA. 2016 TNT version 1.5, including a full implementation of
phylogenetic morphometrics. *Cladistics* **32**, 221–238. (doi:10.1111/cla.12160)

75. de Queiroz K, Gauthier JA. 2020 Squamata. *Phylonyms Companion PhyloCode*

76. Modesto SP, Sues H-D. 2004 The skull of the Early Triassic archosauromorph reptile
Prolacerta broomi and its phylogenetic significance. *Zool. J. Linn. Soc.* **140**, 335–351.

77. Miedema F, Spiekman SN, Fernandez V, Reumer JW, Scheyer TM. 2020 Cranial
morphology of the tanystropheid Macrocnemus bassanii unveiled using synchrotron
microtomography. *Sci. Rep.* **10**, 12412.

78. Nesbitt SJ, Stocker MR, Chatterjee S, Horner JR, Goodwin MB. 2021 A remarkable group of
thick-headed Triassic Period archosauromorphs with a wide, possibly Pangean distribution.
J. Anat. **239**, 184–206. (doi:10.1111/joa.13414)

79. Winchester L, Bellairs A d'A. 1977 Aspects of vertebral development in lizards and snakes.
J. Zool. **181**, 495–525. (doi:10.1111/j.1469-7998.1977.tb03259.x)

80. Brochu CA. 1996 Closure of neurocentral sutures during crocodilian ontogeny: Implications
for maturity assessment in fossil archosaurs. *J. Vertebr. Paleontol.* **16**, 49–62.
(doi:10.1080/02724634.1996.10011283)

81. Griffin CT, Stocker MR, Colleary C, Stefanic CM, Lessner EJ, Riegler M, Formoso K,
Koeller K, Nesbitt SJ. 2021 Assessing ontogenetic maturity in extinct saurian reptiles. *Biol.*
Rev. **96**, 470–525. (doi:10.1111/brv.12666)

Figures

871 **Figure 1.** Comparison of jugal morphologies among living lepidosaurs. Note the variability in

872 the presence and development of the posterior process, as well as the presence of an ossified

873 jugal itself. CT scan images are from digimorph.org.

Figure 2. Comparison of palatine morphologies. Blue shading indicates choanal fossa. Top

image of *Cryptovaranoides* referred palatine is from Whiteside et al. [19].

Prominent choanal fossa on anterior margin of ventral surface of palatine

0: absent

1: present, limited to anterior third of palatine

2: present and extending over halfway along palatine

considered "prominent" by WEA24

Sphenodon punctatus
after Gauthier et al. (2012)

†*Helioscopos dickersonae*
after Meyer et al. (2023)

†*Eoscincus ornatus*
after Brownstein et al. (2022)

†*Cryptovaranoides*
Holotype

887 **Figure 3.** Choanal fossa character scoring.

891 **Figure 4.** Comparison of vomer morphologies. Top image is *Eoscincus ornatus* from Brownstein

892 et al. (2022). Note that the rows of vomerine teeth are posteriorly placed, and the vomerine

893 ridges are large and laterally placed. Bottom image of *Cryptovaranoides* holotype is from

894 Whiteside et al. [19].

Cephalic head of (mobile) quadrate with notch for the squamosal, peg-in-notch articulation

0: absent   1: present   considered "present" by WEA24

quadrate
squamosal

Sphenodon punctatus
after Gauthier et al. (2012)

Amphiglossus splendidus
after Gauthier et al. (2012)

†*Cryptovaranoides*
Holotype

896
897
898   **Figure 5.** Quadrate-squamosal articulation character scoring.
899
900

Vomer and maxilla meet at anterior margin of fenestra exochoanalis

0: absent   1: present   considered "present" by WEA24

····· damaged

vomer
premaxilla
maxilla

Sphenodon punctatus
after Gauthier et al. (2012)

Pogona vitticeps
after Gauthier et al. (2012)

†*Cryptovaranoides*
Holotype

901
902   **Figure 6.** Anterior articulation of vomer and maxilla character scoring.
903
904

Celestus enneagrammus
after Gauthier et al. (2012)

position of foramen
according to WEA24

†*Cryptovaranoides*
Holotype

Figure 7. Presence of a medially positioned posterior mylohyoidal foramen on the mandible. As shown, there is no identifiable foramen on the mandible of *Cryptovaranoides*. Bottom image of *Cryptovaranoides* holotype skull is from Whiteside et al. [19].

†*Cryptovaranoides*
Holotype

†*Peurcosuchus traverorum*
after Marsh et al. (2022)

917 **Figure 8.** Comparison of the humerus of *Cryptovaranoides microlanius* to *Puercosuchus*

918 *traverorum* [26]. Left image of *Cryptovaranoides* holotype humerus is from Whiteside et al.

919 [19].

924   **Table 1.**

Character	Number in Talanda et al. (2022)	Study Cited in WEA24	Synapomorphy of	Unambiguous Optimization
Cephalic head of (mobile) quadrate with notch for the squamosal, peg-in-notch articulation with rod-shaped squamosal	123	de Queiroz and Gauthier (2020)	Pan-Squamata	1→0
Vomer and maxilla meet at anterior margin of fenestra exochoanalis	371	Gauthier et al. (2012)	N/A	N/A
Prominent choanal fossa on anterior margin of ventral surface of palatine	100	Gauthier et al. (2012)	Lepidosauria	N/A
Subdivision of embryonic metotic fissure by the crista tuberalis into vagus (jugular) foramen and recessus scala tympani	382	Simões et al. (2018); de Queiroz and Gauthier (2020)	N/A	N/A
No quadrate foramen	118	Gauthier et al. (2012)	Lepidosauria	1→0
Medially positioned posterior mylohyoidal foramen on mandible	163	Gauthier et al. (2012)	N/A	N/A
Fusion of exoccipitals and opisthotics forming an otoccipital	151	Gauthier et al. (2012); de Queiroz and Gauthier (2020)	N/A	N/A
Trunk vertebrae lack intercentra	237	de Queiroz and Gauthier (2020)	Pan-Unidentata	1→0