Are all waterholes equal from a lion's view? Exploring the role of prey abundance and catchability

in waterhole visitation patterns in a savannah ecosystem

Romain Dejeante[1*]; Andrew J. Loveridge[2,3]; David W. Macdonald[2]; Daphine Madhlamoto[4]; Simon

Chamaillé-Jammes[1,5,6,#] ; Marion Valeix[1,5,7,#]

1. CEFE, Univ Montpellier, CNRS, EPHE, IRD, Montpellier, France

2. Wildlife Conservation Research Unit, Department of Biology, University of Oxford, Oxford OX1 3EL,

UK

3. Panthera, 104 West 40th Street, New York, NY 10018, USA.

4. Zimbabwe Parks and Wildlife Management Authority, Main Camp Research, Hwange National Park,

Zimbabwe

5. LTSER France, Zone Atelier 'Hwange', Hwange National Park, Zimbabwe

6. Mammal Research Institute, Department of Zoology and Entomology, University of Pretoria, Pretoria,

South Africa

7. REHABS/SRU, International Research Lab, CNRS-NMU, George Campus, Nelson Mandela

University, 6530, George, South Africa

* Corresponding author: dejeante.romain@laposte.net

shared last authorship

Abstract

Prey abundance and catchability shape the spatial ecology of predators. Predators can select habitats

where prey are more abundant to maximize encounter rate with prey or habitats where prey are more

catchable to maximize prey capture. These hypotheses are commonly referred to as prey-abundance and

prey-catchability hypotheses. Although these hypotheses are often tested at the landscape scale, little is

known about how between-patch variations in prey abundance and catchability determine the space use of

predators. In many savannah ecosystems, large herbivores aggregate around waterholes, which become

hotspots of prey and their selection by predators is classically interpreted as supporting the prey-

abundance hypothesis. Here, we investigated whether between-waterhole variations in prey abundance

and catchability influence the frequency and duration of lion visits to waterholes, testing the prey-

abundance and prey-catchability hypotheses at the resource-patch scale. We combined datasets on (1) lion

movements recorded from GPS collars deployed on 20 adult males and 16 adult females between 2002

and 2015, (2) prey abundance evaluated from long-term, regular monitoring of waterholes and (3) prey

catchability evaluated from remote-sensing satellite imagery of vegetation cover around waterholes in

Hwange National Park (Zimbabwe). Lions did not use all waterholes in their territory equally: there was a

high variability in the frequency and duration of visits. Surprisingly, between-waterhole variations in prey

abundance and catchability only slightly explained these variations in frequency – and even less in

duration – of lion visits to waterholes. Yet, the frequency of lion visits to waterholes decreased with the

number of waterholes within their territory, and male lions more frequently visited the waterholes

surrounded by more open habitats. We discuss the limits of our work, but also the ecological mechanisms

that may explain these findings. First, lions and their prey are involved in a 'shell-game' that leads them

to adopt unpredictable movement strategies. Second, lions have only access to a limited number of

waterholes amongst which to distribute their hunting effort. Lastly, lions visit waterholes not only to hunt

but also to interact with social mates and competitors. This work challenges the implicit assumption that

all waterholes are the same from a lion's view and calls for further studies investigating the drivers of the

variability in lion visits at the resource-patch scale.

Introduction

The study of how predator-prey interactions shape the spatial behaviour of both predators and prey has a long history in the ecological literature (Creel et al., 2005; Fortin et al., 2005; Gaynor et al., 2019; Gilliam & Fraser, 1987; Palmer et al., 2022). Predators and prey are involved in a 'shell game' in which they endlessly adjust their behaviour in response to one another (Mitchell & Lima, 2002; Sih, 2005; Simon et al., 2019) with prey attempting to avoid predators and predators attempting to be unpredictable for prey (Patin et al., 2020). From a predation perspective, predators are classically expected to use habitats more intensively where prey are more abundant, seeking to maximize the encounter rate with prey (i.e., "prey-abundance hypothesis") (Davidson et al., 2012; Kittle et al., 2017; Zabihi-Seissan et al., 2022). They are also expected to use habitats more intensively where prey are more catchable, seeking to maximize prey capture (i.e., "prey-catchability hypothesis") (Balme et al., 2007; Hopcraft et al., 2005; Zabihi-Seissan et al., 2022). However, these hypotheses are not mutually exclusive and predators should adjust their spatial behaviour to variations in both prey abundance and catchability, i.e., prey availability (Smith et al., 2020). Despite this recognition, the roles of prey abundance and prey catchability on predator spatial behaviour have largely been examined separately, limiting our understanding of their relative importance or their possible interactive influence on spatial ecology of predators. For example, do predators consistently favour habitats where prey are more catchable, or do they seek to maximize prey-catchability solely where prey are locally scarce?

Unfortunately, ecologists often lack knowledge on prey abundance and catchability at the resource-patch scale to investigate the intricacies between these two measures. Instead, prey abundance and catchability are commonly evaluated at the landscape scale by considering some habitat types as a proxy for prey abundance, and others as a proxy for prey catchability (see for example Balme et al. 2007; Kauffman et al. 2007; Smith et al. 2020). In savannah ecosystems, waterholes and open grasslands are intensively used by large herbivores (Courbin et al., 2019; Smit et al., 2007; Valeix, Loveridge, et al., 2009) and their selection by predators is interpreted as supporting the prey-abundance hypothesis

(Davidson et al., 2012; Valeix et al. 2010). Conversely, tall grasslands, woody vegetation, thickets or

rocky outcrops offer numerous ambush sites for stalk-and-ambush predators and their selection by such

predators is interpreted as supporting the prey-catchability hypothesis (Davidson et al., 2012; Davies et

al., 2016; Hopcraft et al., 2005). However, comparing a predator's selection for several habitat types does

not allow ecologists to address how predators respond to between-patch variation in prey abundance and

catchability. Indeed, prey abundance and catchability likely vary spatially from one site to another – see

for example Chamaillé-Jammes et al. (2016) for between-waterhole variations in herbivore distribution –

and predators may seek to maximize differently prey capture or prey encounter rate depending on these

local variations.

An important gap in our understanding of the spatial ecology of predators is therefore predator

responses to between-patch variations in prey abundance and catchability. Here, to explore the prey-

abundance and prey-catchability hypotheses at the between-patch scale, we investigated the spatial

responses of African lions (*Panthera leo*) to between-waterhole variations in prey abundance and

catchability in a savannah ecosystem, Hwange National Park, Zimbabwe, where the role of waterholes on

the spatial ecology of lions has been extensively studied.

In this ecosystem, waterholes are known to attract large herbivores, particularly in the dry season

(Valeix, Loveridge, et al., 2009; Valeix, 2011), and thus can be considered as prey hotspots, and

ultimately key hunting sites for lions (Davidson et al., 2012; Valeix et al., 2010; Valeix, Fritz, et al.,

2009). They even shape lion territory size (Loveridge et al., 2009; Valeix et al., 2012) and the way lions

use their territory (Dejeante, Andrew J. Loveridge, et al., 2025). However, most of these studies did not

consider heterogeneity in the characteristics of waterholes. The abundance of herbivores varies, spatially,

from one waterhole to another, and between-waterhole variations in prey abundance are consistent over

years and substantially explain the spatial structure of herbivores communities (Chamaillé-Jammes et al.,

2016). Further, whereas some waterholes are located in large open grasslands, others are located in

woodlands or bushlands, influencing the catchability of herbivores near these waterholes, as lions are

stalk-and-ambush hunters expected to have a higher hunting success in environments where vegetation

can provide concealment (Davies et al., 2016; Funston et al., 2001; Schaller, 1972). How lions adjust their

space use to between-waterhole variations in both prey abundance and catchability remains unknown.

The intensity of use of a location – although commonly evaluated in movement ecology, for

example to estimate an animal's home range or an animal's habitat selection – has the limitation of

aggregating how frequently and for how long an individual visits a location, which restricts our

understanding of the animal's spatial ecology (Bastille-Rousseau et al., 2024; Benhamou & Riotte-

Lambert, 2012). Here, we therefore investigated how between-waterhole differences in prey abundance

and catchability influence the frequencies and durations of lion visits to the waterholes available in their

territory. In particular, following the *prey-abundance hypothesis*, we predicted that lions would more

frequently use waterholes with a higher prey abundance and for a longer duration. However, such a

response may depend on the average prey abundance within lion territories: for example, lions may spend

more time close to a waterhole with a higher prey abundance when prey are generally scarce within their

territory, but may not do so if prey are abundant throughout their territory *(functional response in prey-*

abundance hypothesis). Following the *prey-catchability hypothesis*, because lions are predators that

commonly hunt by stalking and ambushing their prey (Funston et al., 2001; Schaller, 1972), we predicted

that lions would more frequently use those waterholes with a higher prey catchability (i.e., less open

vegetation or short distance between vegetation cover and waterhole bank) and for a longer duration.

Similarly, we tested whether lions respond more strongly to between-waterhole variations in prey

catchability when prey were scarce in their territories *(functional response in prey-catchability*

hypothesis). Finally, because lions may adjust how often they visit a waterhole and the time they spend

close to this waterhole to both the local variations in prey abundance and catchability simultaneously, we

also predicted that they should maximize the use of waterholes with both high prey abundance and high

prey catchability *(functional response in both prey-abundance and prey-catchability hypotheses)*.

Methods

Study area

The study was conducted in the north-eastern area of Hwange National Park, Zimbabwe. The park covers

14,600 km² of semi-arid savannah. In this study, we used data collected during the late dry season, from

July to mid-October, when natural rain-fed pans are dry and surface water remains available only in

artificial permanent waterholes in which underground water is pumped (Chamaillé-Jammes et al., 2007).

A GIS layer of all artificial permanent waterholes that retain water throughout the dry season was

available.

Lion visits to waterholes

We investigated lion visits to waterholes from movement data collected from 20 adult male lions and 16

adult female lions between 2002 and 2015. Lions were monitored with GPS collars, collecting hourly or

two-hourly locations. As lions were commonly equipped over more than one year, we used animal-year

as our sampling unit, resulting in 38 animal-year units for males, and 35 animal-year units for females.

Because prey-abundance data (see the following section) was only available during the dry season, only

movement data from the late dry season, between July and mid-October, were analysed. Lion handling

and care protocols were consistent with guidelines provided in the 'Code of Practice for Biologists using

Animals', Department of Biology, University of Oxford and approved by University of Oxford,

Biomedical Sciences, Animal Welfare and Ethical Review Body. Lion handling procedures were carried

out by project staff trained and certified by the Zimbabwe Veterinary Association, Wildlife Group

(Certificate numbers: Davidson: 09/03; Elliot: 9/10; Hunt: A11/04, 005/09, 5/14; Loveridge: 6/2000,

20/36 (2007), 6/14; Stapelkamp: 34/2008) in accordance with Statutory Instrument 409 of 1999 (Clause

21A to 21J) amending the Regulations of 1975 to the Dangerous Drugs Act, Zimbabwe.

For each animal-year, we evaluated the frequency and the duration of visits to each waterhole of a

lion territory using the R package *recurse* (Bracis et al., 2018) to count, respectively, the number of times

a collared lion entered a circular buffer around a waterhole – over 100 days – and to measure the time

spent within this buffer per visit. Following Bracis et al., (2018), the entrance and exit times for each visit

were calculated by linear interpolation between the trajectory locations inside and outside the buffer. We

performed our analyses using both a 1km and a 2km radius around each waterhole. These distances are

commonly employed to investigate the behaviour of lions in the vicinity of waterholes in the study

ecosystem (Courbin et al., 2016; Davidson et al., 2013; Dejeante et al., 2024; Valeix et al., 2010; Valeix,

Fritz, et al., 2009). We used the two distance thresholds to test the sensitivity of our results to the chosen

value. Results did not differ qualitatively, and we present results obtained with the 1km radius threshold

in the main text. Results from the analyses with the 2km radius threshold are available in the

Supplementary Information (Tables S1 and S2). Because the number of days of tracking during the dry

season varied from one animal-year to another – from 60 to 105 days – we re-calculated the number of

visits to a waterhole over a period of 100 days using a proportional relationship. This allowed us to

compare the number of visits from one animal-year to another.

We estimated the frequency and the duration of lion visits to each waterhole within their territory,

as delineated by the minimum convex polygon created from 100% of lion's movement data collected

during the late dry season. Although using kernel-based methods is commonly a more revealing approach

to delineate an animal's home range (Börger et al., 2006; Nilsen et al., 2008), here we used a minimum

convex polygon to consider as available all waterholes potentially reachable by an individual, even those

rarely or never visited. Contrary to kernel-based methods, drawing a minimum convex polygon does not

allow us to measure differences in the intensity of use. However, these relative differences in the intensity

of use of each available waterhole within a lion territory were explored from the differences in the

frequency and duration of visits. In the following analyses, we seek to explain differences in the

frequency and duration of lion visits to each waterhole of their territory by between-waterhole variations

in prey abundance and catchability.

Prey abundance data

To evaluate prey abundance at each waterhole, we used a long-term, regular monitoring of waterholes

conducted by the Wildlife Environment Zimbabwe (WEZ) since 1972. Every year, at the end of the dry

season, the WEZ records the animals coming to drink to each waterhole of the park over a 24h period (all

waterholes with water are monitored). This monitoring provides information on the total number of herds

and the average herd size that visited a waterhole for all large herbivore species. Importantly, Chamaillé-

Jammes et al. (2016) showed that the spatial structure of the herbivore community is mostly explained by

between-waterhole variations and that differences in prey abundance between waterholes are consistent

over years. Although the park has a wide diversity of large herbivores, here, we focused only on the main

prey species of lions (Davidson et al., 2013): African buffalo (*Syncerus caffer*), plains zebra (*Equus*

quagga), blue wildebeest (*Connochaetes taurinus*), roan antelope (*Hippotragus equines*), sable antelope

(*Hippotragus niger*), greater kudu (*Tragelaphus strepsiceros*) and eland (*Taurotragus oryx*). For species

other than buffalo, because encounter rates between predators and prey are better predicted by prey herd

density than by prey individual density (Fryxell et al., 2007), and because their herd size does not show a

high variability in Hwange National Park (Supplementary Information Table S3), we used the total

number of herds drinking at a waterhole (pooling all species) as a proxy for medium-sized prey

abundance at this waterhole. Hunting medium-sized prey is generally different from hunting larger prey,

such as buffalo. Large prey are indeed harder and more dangerous to catch but provide more food

(Funston et al., 1998, 2001). Besides, buffaloes in the study ecosystem are mostly found in a few very

large groups (hundreds of individuals; with groups as large as ~1000 individuals) that have fixed and

large territories (Wielgus, 2020). Hence although they regularly visit the waterholes in their territory, the

likelihood of not observing them during a 24h period is high. As buffaloes are a particularly important

prey species for lions and given this low probability of detection, we chose not to aggregate buffaloes and

other prey, but to average buffalo data over a 5-year window, to maximize the chance of the large buffalo

herd detection at this waterhole. As the buffalo herds range over stable spatial areas over time, we believe

that using a 5-year window allowed us to minimize the detection bias.

Variables related to medium-sized prey or buffalo abundance were centred and scaled per animal-

year territory (hereafter 'Herd' and 'Buffalo' variables) to test whether, within a territory, lions visit more

waterholes with higher abundance in herds of medium-sized prey or in buffalos ('prey-abundance

hypothesis'). In addition, we also calculated the mean herd and buffalo abundance per animal-year within

each lion territory ('Herd Territory' and 'Buffalo Territory') and then centred and scaled this variable

across all animal-year territories. Adding an interaction term between the 'Herd' and 'Herd Territory'

variables allowed us to test whether lions visit more waterholes with higher herd abundance specifically

when their territory is poor or rich in herds ('functional response in prey-abundance hypothesis').

Prey catchability data

In Hwange National Park, the vegetation is dominated by bushlands and woodlands interspersed with

patches of grasslands surrounding waterholes (Arraut et al., 2018). In this analysis, we considered two

scales. First, we used the distance between the waterhole bank and the vegetation cover (measured using

Google Earth Pro © for each cardinal direction (North, South, East and West) and then averaging these 4

distances for each waterhole) as a proxy for the fine-scale prey catchability at a waterhole (Figure 1;

'Distance to cover' variable). Second, we used the mean vegetation openness within the 1km surrounding

each waterhole as a proxy for the large-scale prey catchability around a waterhole ('vegetation openness'

variable). To do so, we used the 30-m resolution vegetation map produced by Arraut et al. (2018) to

calculate, for each pixel, the proportion of open vegetation (category 'grassland' and 'open bushlands' in

the original map) in a radius of 250 m. We then averaged the proportion of open vegetation within a

radius of 1km around each waterhole. For each metric, we assumed that prey in more open areas (i.e.,

high distance between the waterhole and the vegetation cover or higher proportion of open vegetation) are

less catchable because of the better visibility conditions to detected an approaching lion – and that lions

should therefore more frequently visit (or over longer periods) the waterholes surrounded by less open

areas if prey catchability drives their movement decisions at the patch-scale.

Data analysis

We investigated how between-waterhole variations in prey abundance and catchability influenced the

differences in the frequency and duration of lion visits to the different waterholes of their territory. Since

the frequency and duration of lion visits to waterholes exhibited overdispersion in Poisson models, we

used negative binomial regressions to model these response variables. For each waterhole within a lion

territory, we tested the influence of the explanatory variables related to prey abundance and prey

catchability on the frequency or on the duration of visits to the corresponding waterhole by the

corresponding lion. Because of the structure of the data, all models included a random intercept with

animal-year identity. Model sets are presented in Tables 1 and 2 and are grouped according to the tested

hypotheses (see Introduction). Because the frequency of visits to a waterhole and the duration of these

visits may not be related solely to prey availability but also to the number of waterholes within the lion

territory, we controlled for this effect by including the number of waterholes within the territory of an

animal-year as a fixed effect in all tested models.

To determine the most parsimonious model explaining the differences of frequency and duration

of lion visits to a waterhole, we used the Akaike information criterion corrected for small sample size

(AICc) (Burnham & Anderson, 2004). In the following, we present results obtained from the most

parsimonious models, i.e., with both a ΔAICc < 2 and the lowest number of explanatory variables

(Arnold, 2010). Selected models show no multicollinearity between predictors, as indicated by variance

inflation factors (VIF) lower than 2. Analyses were conducted using the R package *glmmTMB* (Brooks et

al., 2017). We used the R package *performance* (Lüdecke et al., 2021) to measure the VIF scores and the

goodness-of-fit measure of the models as estimated by the marginal pseudo-R^2 for generalized linear

mixed models (Nakagawa et al., 2017).

Results

Influence of prey abundance and catchability on the frequency of lion visits to a waterhole

On average, lions visited a given waterhole in their territory 8 times (for female) or 9 times (for males)

over a 100-day period of the late dry season, but a very large variability was observed, and some

waterholes could be visited by the same lion more than 50 times (Figure 2.a-b). For male lions, the most

parsimonious model explaining the variability of the frequency of visits to a waterhole included the

number of waterholes in the territory, the number of buffalo herds visiting the waterhole and the

vegetation openness in the surroundings of the waterhole (Table 1; *Buffalo + Vegetation openness*).

Details of model coefficients and errors are available in the Supplementary Information (Table S4). Male

lions more frequently visited waterholes surrounded by open vegetation and those more frequently used

by buffalo herds, relative to other waterholes available in their territories (Figure 3). Unsurprisingly, they

also less frequently visited each waterhole when their territory encompassed a larger number of

waterholes. Overall, this model explained 28% of the variability in the number of visits to a waterhole.

The vegetation openness was the main variable explaining this variability. Indeed, the null model – that

solely included the number of waterholes – explained only 6% of this variability and only 8% when also

accounting for buffalo abundance against 25% when accounting for the vegetation openness (and not

accounting for buffalo abundance) (Table 1).

For female lions, the most parsimonious model explaining the variability of the frequency of

visits to a waterhole included the number of waterholes in the territory, the number of herds of medium-

sized prey and the vegetation openness in the surroundings of the waterhole (Table 1; *Herd + Vegetation*

openness). Details of model coefficients and errors are available in the Supplementary Information (Table

S5). Female lions more frequently visited waterholes surrounded by open vegetation and those more

frequently used by herds of medium-sized prey (Figure 3). As for male lions, they also less frequently

visited each waterhole when the number of waterholes in their territory increased. Although this model

explained 36% of the variability in the frequency of visits to a waterhole by female lions, the metrics

related to prey abundance and catchability explained only a small amount of this variability. Indeed, the

null model – that included the number of waterholes only – explained 30% of total variability. This result

suggests that the vegetation openness and the number of medium-sized prey herds only slightly affect the

frequency of visits to a waterhole by female lions.

Influence of prey abundance and catchability on the duration of lion visits to a waterhole

On average, lions visited a given waterhole in their territory for 7 hours (for females) or 6 hours (for

males) per visit, but with high variability between waterholes, as some visits could last more than 24

hours (Figure 2.c-d). The relationship between durations and frequencies of lion visits to a given

waterhole of their territories is available in the Supplementary Information (Figure S1). For both male and

female lions, all models tested only slightly explained the variability in the duration of visits to a

waterhole. For female lions, the most parsimonious model included solely the number of waterholes in

the territory (Table 2; *null hypothesis*) and only explained 3% of the variability in the duration of visits

(Figure 4). Details of model coefficients and errors are available in the Supplementary Information (Table

S6). For male lions, the most parsimonious model, which included the number of waterholes in the

territory and the vegetation openness in the surroundings of the waterhole (Table 2; *Vegetation openness*)

– only explained 9% of this variability. Details of model coefficients and errors are available in the

Supplementary Information (Table S7). Although male lions stayed longer times at a waterhole where the

vegetation openness was greater in the surroundings (Figure 4), the low pseudo-R^2 of this model suggests

that this variable only slightly affect the duration of visits.

Discussion

In savannah ecosystems, waterhole areas are key foraging patches for predators and their prey (Davidson

et al., 2012; Mosser et al., 2015; Valeix et al., 2010; Valeix, Fritz, et al., 2009; Valeix, Loveridge, et al.,

2009). However, whether all waterholes are similarly attractive from a predator's viewpoint or whether

predators favour some waterhole areas rather than others remains unknown. Here, we investigated how

between-waterhole variations in prey abundance and catchability influenced the frequency and duration of

lion visits to a waterhole. Surprisingly, we found that the prey-related variables considered in this work

only slightly affected the frequency – and even less the duration – of visits to a waterhole by male and

female lions. Our main findings are that (1) lions do not use equally (different frequencies and durations

of visits) the waterholes in their territory, (2) the duration of visits of waterholes by lions is not influenced

by the associated prey abundance and vegetation openness, (3) lions of both sexes tend to more frequently

visit waterholes in their territory when there are fewer waterholes, (4) prey abundance (as measured in

this work) does not affect the frequency of visits at a waterhole by lions, and (5) male lions tend to more

often visit waterholes surrounded by open vegetation. The ecological mechanisms and the limits of our

work that may explain these findings are discussed below.

A dynamic predator-prey space race

In ecosystems with defined prey hotspots, predators are expected to employ a 'leap-frogging' behaviour

(Sih, 2005) whereby their spatial patterns are shaped by these prey hotspots. This has been documented

for wolves (Kittle et al., 2017) and lions (Kittle et al., 2016; for Hwange: Dejeante, Loveridge,

Macdonald, Madhlamoto, Chamaillé-Jammes, et al., 2025). Although predators can rely on cues of higher

prey abundance or higher prey catchability to move between these prey hotspots (Balme et al., 2007;

Davidson et al., 2012; Hopcraft et al., 2005), predator and prey are involved in a 'shell game' in which

they endlessly adjust their behaviour in relation to one another to minimize the predictability of their

presence (Gaynor et al., 2019; Mitchell, 2009; Mitchell & Lima, 2002). Indeed, if predators are too

predictable, prey can proactively increase their vigilance when foraging close to well-known ambush site

or can avoid foraging nearby. Because of this predator-prey space race, prey can seek to remain

unpredictable to predators and can be reluctant to use some waterholes despite their importance in the

landscape or using them during the day when the risk of predation is lower (Courbin et al., 2019).

However, waterholes remain key hunting sites for lions and some waterholes remain consistently more

visited by herbivores than others (Chamaillé-Jammes et al., 2016) and may therefore be seen as sites with

more predictable prey presence by lions. As demonstrated by Patin et al. (2020), predators should

therefore seek to be unpredictable to optimize their hunting success, integrating some level of randomness

in their movement behaviour to maximize their chances of encountering and capturing prey. Regarding

this predator-prey space race, lions would therefore benefit from not visiting too regularly the same

waterholes in order to be less predictable by prey. Lions in Hwange National Park are known to make

most of their kills close to a waterhole and to commonly depart from a kill site to visit another waterhole

(Valeix et al., 2011). Altogether, this and our findings suggest that lions may spread out their hunting

grounds in an unpredictable manner. Even in the absence of kills, lions need to depart sufficiently quickly

from a given hunting patch to counter-balance the anti-predator responses adopted by prey once they

detect their presence there (Charnov et al. 1976; Kotler 1992). One of the anti-predator responses adopted

by prey is to move away from the site of an encounter with lions (Courbin et al., 2016; Martin & Owen-

Smith, 2016; Say-Sallaz et al., 2023), possibly leading to a depression in prey abundance in the vicinity of

a waterhole following a lion visit. Here, unfortunately, our prey-related variables do not allow us to draw

inference about these short-term variations in prey abundance as they cannot be used to predict the

number of prey at waterhole at the specific time of the visit by a lion and may therefore not perfectly

reflect prey abundance from a lion's viewpoint. Indeed, although some waterholes have consistently

greater herbivore abundance than others in Hwange National Park (Chamaillé-Jammes et al., 2016), lions

may use other cues to infer instantaneous prey abundance. Because of a short-term depression in prey

abundance, one waterhole having consistently greater herbivore abundance than others may offer – at a

given time, for example following an attempted hunt – less opportunities for lions to encounter prey than

an undisturbed waterhole having on average a lower number of herbivores in its vicinity. Being territorial

animals (Schaller, 1972), lions have access to only a limited number of waterholes to spread out their

hunting effort. Therefore, even if they seek preferentially to visit undisturbed feeding patches, where prey

are most available, lions may be constrained to visit patches with low prey abundance or poor prey

catchability if those patches are the ones least disturbed by their previous presence.

A territorial and social species

Although this study focuses on the importance of prey-related variables to explain between-waterhole

variations in the duration and frequency of visits to a waterhole, lions are not solely predators but also

territorial and social animals (Schaller, 1972). Other variables, unrelated to prey, may therefore partly

explain differences in the frequency and duration of visits to a waterhole. In particular, beyond being key

hunting patches, waterholes are key marking places for lions (Dejeante, Andrew J. Loveridge, et al.,

2025). For example, lions vocalize more frequentlyly near water sources (Wijers et al. 2021), and male

lions more frequently encounter their rivals near waterholes (Dejeante et al. 2024). The distribution of

these waterholes greatly structures the size and shape of lion territories (Valeix et al., 2012) and drives

their territorial movements in the core and in the periphery of the territory (Dejeante, Andrew J.

Loveridge, et al., 2025). We would therefore expect that lions' territorial behaviours – and not solely

predator-prey interactions – influence the duration and the frequency of lion visits to a waterhole. As

demonstrated for another territorial species (wolves that integrate space and time to visit territorial

borders; Schlägel et al. 2017), lions might, for example, be more likely to visit a waterhole not visited for

a long time to renew their marking signals at this place. Interestingly, here, we found a great influence of

the vegetation openness around a waterhole: male lions were more likely to frequently visit a waterhole if

it was surrounded by open habitats. This result is not intuitively supportive of the prey-catchability

hypothesis, but prey catchability may still matter in these large open areas, but at a finer scale than

captured by our metrics. In addition, because a more open habitat is likely associated with a higher

probability of encountering large herds of large herbivores in Hwange National Park, this result could be

shaped by a long-term cue of high prey abundance, not captured by the data available from waterhole

counts. However, other explanations are possible. In particular, a territorial-related explanation is that

male lions spend more time with their pride in open areas because the visibility is greater in these

habitats, allowing them to detect more easily intruders and so to less frequently patrol their territory

(Dejeante et al., 2024; Funston et al., 1998). If male lions favour visiting waterhole surrounded by open

habitats for territorial reasons not related to prey, these prey could, however, become reluctant to use

these waterholes. This would lead to a mismatch between prey abundance and the patterns of lion visits to

a waterhole. At the interspecific level, the spatio-temporal use of waterholes may also be partly driven by

carnivore intraguild interactions. In the Hwange ecosystem, this is for example the case with lions often

encountering spotted hyaenas around waterholes and leaving the waterhole area after such encounters

(Périquet et al., 2021).

Conclusion

Many studies investigating the role of waterholes, i.e., prey patches, in the spatial behaviour of lions

implicitly assumed that all waterholes are the same (Davidson et al., 2012, 2013; Valeix et al., 2010;

Valeix, Fritz, et al., 2009). Here – to our knowledge for the first time – we investigated whether this

assumption was true. Clearly there is an heterogeneity in the frequency and duration of lion visits at

waterholes, suggesting that not all waterholes are the same from a lion's viewpoint. Although lions did

not equally use all waterholes within their territory, we were able to provide little evidence that the

variability of the frequency and duration of lion visits to a waterhole was determined by between-

waterhole variations in prey abundance and our measures of its catchability. Three, non-exclusive, main

points can explain this result: (1) lions and their prey are involved in a "shell-game" that leads them to

adopt unpredictable movement strategies, and the proxies used in this work did not take into account

these dynamics, (2) lions have access to a limited number of waterholes amongst which to distribute their

hunting effort, and this leads them to visit some waterholes with lower prey abundance or catchability

solely because those are the least disturbed, (3) lions do not visit waterholes solely to hunt but also to

interact with conspecifics (both social mates or rivals) for example to frequently renew their territorial

signals at these places, whatever the abundance and catchability of prey is there.

Declaration of interest

388   The authors declare that they have no conflict of interest.

Acknowledgments

We thank the Zimbabwe Parks and Wildlife Management Authority for permission to carry out this

research. The Hwange Lion Project was funded by grants from the Robertson Foundation, the Recanati-

Kaplan Foundation, the Darwin Initiative for Biodiversity grant 162/09/015, a CV Starr Scholarship, The

Eppley Foundation, Disney Foundation, Marwell Preservation Trust, Regina B. Frankenburg Foundation,

Rufford Maurice Laing Foundation, Panthera Foundation and the generosity of Joan and Riv Winant. We

are deeply grateful to Jane Hunt, Zeke Davidson, Nicholas Elliot, Brent Stapelkamp, Dan Parker, Agrippa

Moyo, Lovemore Sibanda, Moreangels Mbizah, Andrea Sibanda and Liomba Mathe for their roles in the

collection of lion GPS data. We are also indebted to Wildlife Environment Zimbabwe (WEZ) for their

invaluable long-term dataset on waterhole use by herbivores, and we sincerely thank Yolan Richard for

measuring the distances between the waterholes' bank and the vegetation cover. Finally, we thank Kirby

Mills and Andrew Kittle for their comments on an earlier version of this manuscript.

Data Availability

Data and codes used in this study are publicly available in a figshare repository (Dejeante, Andrew J

Loveridge, et al., 2025)

References

Arnold, T. W. (2010). Uninformative Parameters and Model Selection Using Akaike's Information

Criterion. *The Journal of Wildlife Management*, *74*(6), 1175–1178.

https://doi.org/10.1111/j.1937-2817.2010.tb01236.x

Arraut, E. M., Loveridge, A. J., Chamaillé-Jammes, S., Valls-Fox, H., & Macdonald, D. W. (2018). The

2013-2014 vegetation structure map of Hwange National Park, Zimbabwe, produced using free

satellite images and software. *Koedoe*, *60*(1), 1–10. https://doi.org/10.4102/koedoe.v60i1.1497

Balme, G., Hunter, L., & Slotow, R. (2007). Feeding habitat selection by hunting leopards Panthera

pardus in a woodland savanna: Prey catchability versus abundance. *Animal Behaviour*, *74*, 589–

598. https://doi.org/10.1016/j.anbehav.2006.12.014

Bastille-Rousseau, G., Crews, S. A., Donovan, E. B., Egan, M. E., Gorman, N. T., Pitman, J. B., Weber,

A. M., Audia, E. M., Larreur, M. R., Manninen, H., Blake, S., Eichholz, M. W., Bergman, E., &

Rayl, N. D. (2024). A multi-property assessment of intensity of use provides a functional

understanding of animal movement. *Methods in Ecology and Evolution*, *15*(2), 345–357.

https://doi.org/10.1111/2041-210X.14274

Benhamou, S., & Riotte-Lambert, L. (2012). Beyond the Utilization Distribution: Identifying home range

areas that are intensively exploited or repeatedly visited. *Ecological Modelling*, *227*, 112–116.

https://doi.org/10.1016/j.ecolmodel.2011.12.015

Börger, L., Franconi, N., De Michele, G., Gantz, A., Meschi, F., Manica, A., Lovari, S., & Coulson, T.

(2006). Effects of Sampling Regime on the Mean and Variance of Home Range Size Estimates.

Journal of Animal Ecology, *75*(6), 1393–1405.

Bracis, C., Bildstein, K. L., & Mueller, T. (2018). Revisitation analysis uncovers spatio-temporal patterns

in animal movement data. *Ecography*, *41*(11), 1801–1811. https://doi.org/10.1111/ecog.03618

Brooks, M. E., Kristensen, K., van Benthem, K. J., Magnusson, A., Berg, C. W., Nielsen, A., Skaug, H.

J., Mächler, M., & Bolker, B. M. (2017). *glmmTMB Balances Speed and Flexibility Among*

Packages for Zero-inflated Generalized Linear Mixed Modeling. *9*.

Burnham, K. P., & Anderson, D. R. (2004). Multimodel Inference: Understanding AIC and BIC in Model

Selection. *Sociological Methods & Research*, *33*(2), 261–304.

https://doi.org/10.1177/0049124104268644

Chamaillé-Jammes, S., Charbonnel, A., Dray, S., Madzikanda, H., & Fritz, H. (2016). Spatial Distribution

of a Large Herbivore Community at Waterholes: An Assessment of Its Stability over Years in

Hwange National Park, Zimbabwe. *PLOS ONE*, *11*(4), e0153639.

https://doi.org/10.1371/journal.pone.0153639

Chamaillé-Jammes, S., Fritz, H., & Murindagomo, F. (2007). Climate-driven fluctuations in surface-

water availability and the buffering role of artificial pumping in an African savanna: Potential

implication for herbivore dynamics. *Austral Ecology*, *32*(7), 740–748.

https://doi.org/10.1111/j.1442-9993.2007.01761.x

Charnov, E. L., Orians, G. H., & Hyatt, K. (1976). Ecological Implications of Resource Depression. *The*

American Naturalist, *110*(972), 247–259.

Courbin, N., Loveridge, A. J., Fritz, H., Macdonald, D. W., Patin, R., Valeix, M., & Chamaillé-Jammes,

S. (2019). Zebra diel migrations reduce encounter risk with lions at night. *Journal of Animal*

Ecology, *88*(1), 92–101. https://doi.org/10.1111/1365-2656.12910

Courbin, N., Loveridge, A. J., Macdonald, D. W., Fritz, H., Valeix, M., Makuwe, E. T., & Chamaillé-

Jammes, S. (2015). Reactive responses of zebras to lion encounters shape their predator-prey

space game at large scale. *Oikos*, *125*(6). https://ora.ox.ac.uk/objects/uuid:63cebb0e-a530-45a0-

8234-cd4464f77328

Courbin, N., Loveridge, A. J., Macdonald, D. W., Fritz, H., Valeix, M., Makuwe, E. T., & Chamaillé-

Jammes, S. (2016). Reactive responses of zebras to lion encounters shape their predator–prey

space game at large scale. *Oikos*, *125*(6), 829–838. https://doi.org/10.1111/oik.02555

Creel, S., Winnie Jr., J., Maxwell, B., Hamlin, K., & Creel, M. (2005). Elk Alter Habitat Selection as an

Antipredator Response to Wolves. *Ecology*, *86*(12), 3387–3397. https://doi.org/10.1890/05-0032

Davidson, Z., Valeix, M., Kesteren, F. V., Loveridge, A. J., Hunt, J. E., Murindagomo, F., & Macdonald,

D. W. (2013). Seasonal Diet and Prey Preference of the African Lion in a Waterhole-Driven

Semi-Arid Savanna. *PLOS ONE*, *8*(2).

Davidson, Z., Valeix, M., Loveridge, A. J., Hunt, J. E., Johnson, P. J., Madzikanda, H., & Macdonald, D.

W. (2012). Environmental determinants of habitat and kill site selection in a large carnivore:

Scale matters. *Journal of Mammalogy*, *93*(3), 677–685. https://doi.org/10.1644/10-MAMM-A-

424.1

Davies, A. B., Tambling, C. J., Kerley, G. I. H., & Asner, G. P. (2016). Effects of Vegetation Structure on

the Location of Lion Kill Sites in African Thicket. *PLOS ONE*, *11*(2), e0149098.

https://doi.org/10.1371/journal.pone.0149098

Dejeante, R., Loveridge, Andrew J, Macdonald, D. W., Madhlamoto, D., Chamaillé, S., & Valeix, M.

(2025). *Data from: Are all waterholes equal from a lion's view? Exploring the role of prey*

abundance and catchability in waterhole visitation patterns in a savannah ecosystem (p. 503717

Bytes) [Dataset]. figshare. https://doi.org/10.6084/M9.FIGSHARE.28825691

Dejeante, R., Loveridge, Andrew J., Macdonald, D. W., Madhlamoto, D., Chamaillé-Jammes, S., &

Valeix, M. (2025). Territorial movements of African lions: Can waterholes shape multiple

central-place territories in a savannah ecosystem? *Animal Behaviour*, *224*, 123193.

https://doi.org/10.1016/j.anbehav.2025.123193

Dejeante, R., Loveridge, A. J., Macdonald, D. W., Madhlamoto, D., Valeix, M., & Chamaillé-Jammes, S.

(2024). Counter-strategies to infanticide: The importance of cubs in determining lion habitat

selection and social interactions. *Journal of Animal Ecology*, *93*(2), 159–170.

https://doi.org/10.1111/1365-2656.14045

Fortin, D., Beyer, H. L., Boyce, M. S., Smith, D. W., Duchesne, T., & Mao, J. S. (2005). Wolves

Influence Elk Movements: Behavior Shapes a Trophic Cascade in Yellowstone National Park.

Ecology, *86*(5), 1320–1330. https://doi.org/10.1890/04-0953

Fryxell, J. M., Mosser, A., Sinclair, A. R. E., & Packer, C. (2007). Group formation stabilizes predator–

prey dynamics. *Nature*, *449*(7165), 1041–1043. https://doi.org/10.1038/nature06177

Funston, P. J., Mills, M. G. L., & Biggs, H. C. (2001). Factors affecting the hunting success of male and

female lions in the Kruger National Park. *Journal of Zoology*, *253*(4), 419–431.

https://doi.org/10.1017/S0952836901000395

Funston, P. J., Mills, M. G. L., Biggs, H. C., & Richardson, P. R. K. (1998). Hunting by male lions:

Ecological influences and socioecological implications. *Animal Behaviour*, *56*(6), 1333–1345.

https://doi.org/10.1006/anbe.1998.0884

Gaynor, K. M., Brown, J. S., Middleton, A. D., Power, M. E., & Brashares, J. S. (2019). Landscapes of

Fear: Spatial Patterns of Risk Perception and Response. *Trends in Ecology & Evolution*, *34*(4),

355–368. https://doi.org/10.1016/j.tree.2019.01.004

Gilliam, J. F., & Fraser, D. F. (1987). Habitat Selection Under Predation Hazard: Test of a Model with

Foraging Minnows. *Ecology*, *68*(6), 1856–1862. https://doi.org/10.2307/1939877

Hopcraft, J. G. C., Sinclair, A. R. E., & Packer, C. (2005). Planning for success: Serengeti lions seek prey

accessibility rather than abundance. *Journal of Animal Ecology*, *74*(3), 559–566.

https://doi.org/10.1111/j.1365-2656.2005.00955.x

Kauffman, M. J., Varley, N., Smith, D. W., Stahler, D. R., MacNulty, D. R., & Boyce, M. S. (2007).

Landscape heterogeneity shapes predation in a newly restored predator–prey system. *Ecology*

Letters, *10*(8), 690–700. https://doi.org/10.1111/j.1461-0248.2007.01059.x

Kittle, A. M., Anderson, M., Avgar, T., Baker, J. A., Brown, G. S., Hagens, J., Iwachewski, E., Moffatt,

S., Mosser, A., Patterson, B. R., Reid, D. E. B., Rodgers, A. R., Shuter, J., Street, G. M.,

Thompson, I. D., Vander Vennen, L. M., & Fryxell, J. M. (2017). Landscape-level wolf space use

is correlated with prey abundance, ease of mobility, and the distribution of prey habitat.

Ecosphere, *8*(4), e01783. https://doi.org/10.1002/ecs2.1783

Kittle, A. M., Bukombe, J. K., Sinclair, A. R. E., Mduma, S. A. R., & Fryxell, J. M. (2016). Landscape-

level movement patterns by lions in western Serengeti: Comparing the influence of inter-specific

competitors, habitat attributes and prey availability. *Movement Ecology*, *4*(1), 17.

https://doi.org/10.1186/s40462-016-0082-9

Kotler, B. P. (1992). Behavioral resource depression and decaying perceived risk of predation in two

species of coexisting gerbils. *Behavioral Ecology and Sociobiology*, *30*(3), 239–244.

https://doi.org/10.1007/BF00166708

Loveridge, A. J., Valeix, M., Davidson, Z., Murindagomo, F., Fritz, H., & Macdonald, D. W. (2009).

Changes in home range size of African lions in relation to pride size and prey biomass in a semi-

arid savanna. *Ecography*, *32*(6), 953–962.

Lüdecke, D., Ben Shachar, M., Patil, I., Waggoner, P., & Makowski, D. (2021). performance: An R

Package for Assessment, Comparison and Testing of Statistical Models. *The Journal of Open*

Source Software, *6*, 3139. https://doi.org/10.21105/joss.03139

Martin, J., & Owen-Smith, N. (2016). Habitat selectivity influences the reactive responses of African

ungulates to encounters with lions. *Animal Behaviour*, *116*, 163–170.

https://doi.org/10.1016/j.anbehav.2016.04.003

Mitchell, W. A. (2009). Multi-behavioral strategies in a predator–prey game: An evolutionary algorithm

analysis. *Oikos*, *118*(7), 1073–1083. https://doi.org/10.1111/j.1600-0706.2009.17204.x

Mitchell, W. A., & Lima, S. L. (2002). Predator-prey shell games: Large-scale movement and its

implications for decision-making by prey. *Oikos*, *99*(2), 249–259. https://doi.org/10.1034/j.1600-

0706.2002.990205.x

Mosser, A. A., Kosmala, M., & Packer, C. (2015). Landscape heterogeneity and behavioral traits drive

the evolution of lion group territoriality. *Behavioral Ecology*, *26*(4), 1051–1059.

https://doi.org/10.1093/beheco/arv046

Nakagawa, S., Johnson, P. C. D., & Schielzeth, H. (2017). The coefficient of determination R2 and intra-

class correlation coefficient from generalized linear mixed-effects models revisited and expanded.

Journal of The Royal Society Interface, *14*(134), 20170213.

https://doi.org/10.1098/rsif.2017.0213

Nilsen, E. B., Pedersen, S., & Linnell, J. D. C. (2008). Can minimum convex polygon home ranges be

used to draw biologically meaningful conclusions? *Ecological Research*, *23*(3), 635–639.

https://doi.org/10.1007/s11284-007-0421-9

Palmer, M. S., Gaynor, K. M., Becker, J. A., Abraham, J. O., Mumma, M. A., & Pringle, R. M. (2022).

Dynamic landscapes of fear: Understanding spatiotemporal risk. *Trends in Ecology & Evolution*,

37(10), 911–925. https://doi.org/10.1016/j.tree.2022.06.007

Patin, R., Fortin, D., & Chamaillé-Jammes, S. (2020). *A theory of the use of information by enemies in the*

predator-prey space race. https://doi.org/10.1101/2020.01.25.919324

Périquet, S., Fritz, H., Revilla, E., Macdonald, D. W., Loveridge, A. J., Mtare, G., & Valeix, M. (2021).

Dynamic interactions between apex predators reveal contrasting seasonal attraction patterns.

Oecologia, *195*(1), 51–63. https://doi.org/10.1007/s00442-020-04802-w

Say-Sallaz, E., Chamaillé-Jammes, S., Périquet, S., Loveridge, A. J., Macdonald, D. W., Antonio, A.,

Fritz, H., & Valeix, M. (2023). Large carnivore dangerousness affects the reactive spatial

response of prey. *Animal Behaviour*, *202*, 149–162.

https://doi.org/10.1016/j.anbehav.2023.05.014

Schaller, G. (1972). *Serengeti Lion: A Study of Predator-Prey Relations (Wildlife behavior and ecology)*.

University of Chicago Press. https://www.biblio.com/book/serengeti-lion-study-predator-prey-

relations/d/460478448

Schlägel, U. E., Merrill, E. H., & Lewis, M. A. (2017). Territory surveillance and prey management:

Wolves keep track of space and time. *Ecology and Evolution*, *7*(20), 8388–8405.

https://doi.org/10.1002/ece3.3176

Sih, A. (2005). Predator-Prey Space Use as an Emergent Outcome of a Behavioral Response Race. In P.

Barbosa & I. Castellanos (Eds.), *Ecology Of Predator-Prey Interactions* (pp. 240–255). Oxford

University PressNew York, NY. https://doi.org/10.1093/oso/9780195171204.003.0011

Simon, R. N., Cherry, S. G., & Fortin, D. (2019). Complex tactics in a dynamic large herbivore–carnivore

spatiotemporal game. *Oikos*, *128*(9), 1318–1328. https://doi.org/10.1111/oik.06166

Smit, I. P. J., Grant, C. C., & Devereux, B. J. (2007). Do artificial waterholes influence the way

herbivores use the landscape? Herbivore distribution patterns around rivers and artificial surface

water sources in a large African savanna park. *Biological Conservation, Special Section: Coastal*

Sandplains, *136*(1), 85–99. https://doi.org/10.1016/j.biocon.2006.11.009

Smith, J. A., Donadio, E., Bidder, O. R., Pauli, J. N., Sheriff, M. J., Perrig, P. L., & Middleton, A. D.

(2020). Where and when to hunt? Decomposing predation success of an ambush carnivore.

Ecology, *101*(12), e03172. https://doi.org/10.1002/ecy.3172

Valeix, M. (2011). Temporal dynamics of dry-season water-hole use by large African herbivores in two

years of contrasting rainfall in Hwange National Park, Zimbabwe. *Journal of Tropical Ecology*,

27(2), 163–170. https://doi.org/10.1017/S0266467410000647

Valeix, M., Chamaillé-Jammes, S., Loveridge, A. J., Davidson, Z., Hunt, J. E., Madzikanda, H., &

Macdonald, D. W. (2011). Understanding Patch Departure Rules for Large Carnivores: Lion

Movements Support a Patch-Disturbance Hypothesis. *The American Naturalist*, *178*(2), 269–275.

https://doi.org/10.1086/660824

Valeix, M., Fritz, H., Loveridge, A. J., Davidson, Z., Hunt, J. E., Murindagomo, F., & Macdonald, D. W.

(2009). Does the risk of encountering lions influence African herbivore behaviour at waterholes?

Behavioral Ecology and Sociobiology, *63*(10), 1483–1494. https://doi.org/10.1007/s00265-009-

0760-3

Valeix, M., Loveridge, A. J., Chamaillé-Jammes, S., Davidson, Z., Murindagomo, F., Fritz, H., &

Macdonald, D. W. (2009). Behavioral adjustments of African herbivores to predation risk by

lions: Spatiotemporal variations influence habitat use. *Ecology*, *90*(1), 23–30.

https://doi.org/10.1890/08-0606.1

Valeix, M., Loveridge, A. J., Davidson, Z., Madzikanda, H., Fritz, H., & Macdonald, D. W. (2010). How

key habitat features influence large terrestrial carnivore movements: Waterholes and African

lions in a semi-arid savanna of north-western Zimbabwe. *Landscape Ecology*, *25*(3), 337–351.

https://doi.org/10.1007/s10980-009-9425-x

Valeix, M., Loveridge, A. J., & Macdonald, D. W. (2012). Influence of prey dispersion on territory and

group size of African lions: A test of the resource dispersion hypothesis. *Ecology*, *93*(11), 2490–

2496. https://doi.org/10.1890/12-0018.1

Wielgus, E. (2020). *The social dynamics of the Cape buffalo and the epidemiological implications.*

Wijers, M., Trethowan, P., du Preez, B., Chamaillé-Jammes, S., Loveridge, A. J., Macdonald, D. W., &

Markham, A. (2021). The influence of spatial features and atmospheric conditions on African lion

vocal behaviour. *Animal Behaviour*, *174*, 63–76. https://doi.org/10.1016/j.anbehav.2021.01.027

Zabihi-Seissan, S., Prokopenko, C. M., & Vander Wal, E. (2022). Wolf spatial behavior promotes

encounters and kills of abundant prey. *Oecologia*, *200*(1), 11–22. https://doi.org/10.1007/s00442-

022-05218-4

Table 1. Summary of the candidate models explaining the variability of the frequency (i.e., the number of

visits over 100 days) of lion visits to a waterhole. All models included the number of waterholes within

the territory of each animal-year and a random effect with the identity of each animal-year. The most

parsimonious models (ΔAICc < 2 and lowest number of parameters K) are shown in bold. For these

models, we used a distance threshold of 1km to consider that lions visited a waterhole.

Frequency of visits to a waterhole – 1km		Female lion			Male lion		
Candidate models	**K**	**AICc**	**ΔAICc**	**R²**	**AICc**	**ΔAICc**	**R²**
Null hypothesis							
Number of waterholes	4	1752.1	16.5	0.30	2413.8	71.4	0.06
Prey abundance hypothesis							
Herd	5	1738.8	3.2	0.35	2391.4	49.0	0.14
Buffalo	5	1747.1	11.5	0.32	2410.3	67.9	0.08
Herd + Buffalo	6	1739.1	3.4	0.35	2393.1	50.7	0.14
Functional response in prey-abundance hypothesis							
Herd × Herd Territory	7	1741.5	5.9	0.35	2391.9	49.6	0.15
Buffalo × Buffalo Territory	7	1742.7	7.1	0.35	2412.1	69.7	0.09
Herd × Herd Territory + Buffalo × Buffalo Territory	10	1742.9	7.3	0.37	2396.3	54.0	0.16
Prey catchability hypothesis							
Distance to cover	5	1753.4	17.8	0.30	2398.3	55.9	0.13
Vegetation openness	5	1742.9	7.2	0.32	2350.9	8.5	0.25
Distance to cover + Vegetation openness	6	1744.0	8.4	0.33	2352.8	10.4	0.26
Functional response in prey-catchability hypothesis							
Distance to cover × Herd Territory	7	1757.3	21.7	0.30	2398.6	56.2	0.14
Vegetation openness × Herd Territory	7	1741.7	6.0	0.34	2351.3	9.0	0.26
Distance to cover × Buffalo Territory	7	1754.3	18.7	0.31	2393.8	51.5	0.17
Vegetation openness × Buffalo Territory	7	1738.4	2.8	0.36	2346.3	3.9	0.28
Distance to cover × (Herd Territory + Buffalo Territory)	9	1757.9	22.2	0.32	2395.1	52.8	0.18

Vegetation openness × (Herd Territory + Buffalo Territory)	9	1740.1	4.4	0.36	2347.2	4.9	0.28
Vegetation openness + Distance to cover × (Herd Territory + Buffalo Territory)	10	1746.4	10.7	0.35	2347.7	5.3	0.29
Distance to cover + Vegetation openness × (Herd Territory + Buffalo Territory)	10	1741.4	5.7	0.37	2349.3	7.0	0.28
Prey-abundance + Prey-catchability hypothesis							
Herd + Distance to cover	6	1740.7	5.0	0.35	2380.8	38.4	0.18
Herd + Vegetation openness	6	**1736.3**	**0.6**	**0.36**	2346.0	3.6	0.27
Herd + Distance to cover + Vegetation openness	7	1737.8	2.2	0.36	2348.0	5.6	0.27
Buffalo + Distance to cover	6	1748.9	13.3	0.32	2395.1	52.7	0.16
Buffalo + Vegetation openness	6	1738.6	2.9	0.35	**2344.2**	**1.8**	**0.28**
Buffalo + Distance to cover + Vegetation openness	7	1739.0	3.3	0.35	2346.0	3.7	0.29
Herd + Buffalo + Distance to cover	7	1741.0	5.4	0.35	2382.3	39.9	0.18
Herd + Buffalo + Vegetation openness	7	1735.9	0.2	0.36	2343.8	1.5	0.28
Herd + Buffalo + Distance to cover + vegetation openness	8	1737.0	1.4	0.37	2345.8	3.4	0.28
Prey abundance × Prey catchability hypothesis							
Herd × Distance to cover	7	1740.1	4.4	0.35	2377.8	35.4	0.20
Herd × Vegetation openness	7	1735.6	0.0	0.36	2347.3	4.9	0.27
Herd × (Distance to cover + Vegetation openness)	9	1737.8	3.1	0.36	2351.0	8.6	0.27
Buffalo × Distance to cover	7	1750.2	14.5	0.33	2393.6	51.3	0.17
Buffalo × Vegetation openness	7	1740.5	4.9	0.35	2343.6	1.2	0.29
Buffalo × (Distance to cover + Vegetation openness)	9	1741.7	6.1	0.36	2345.7	3.3	0.29
Herd × Distance to cover + Vegetation openness	8	1737.8	2.2	0.36	2349.1	6.7	0.27
Herd × Vegetation openness + Distance to cover	8	1737.2	1.6	0.36	2349.4	7.0	0.27
Buffalo × Distance to cover + vegetation openness	8	1739.6	4.0	0.36	2344.3	2.0	0.29
Buffalo × Vegetation openness + Distance to cover	8	1741.0	5.3	0.35	2345.5	3.1	0.29
(Herd + Buffalo) × Distance to cover + vegetation openness	10	1736.3	0.6	0.38	2343.6	1.2	0.30
(Herd + Buffalo) × Vegetation openness + Distance to cover	10	1736.6	0.9	0.38	2342.4	0.0	0.30
(Herd + Buffalo) × (Vegetation openness + Distance to cover)	12	1738.7	3.1	0.38	2343.9	1.5	0.30

601   **Table 2.** Summary of the candidate models explaining the variability of the duration of lion visits to a

602   waterhole. All models included the number of waterholes within the territory of each animal-year and a

603   random effect with the identity of each animal-year. The most parsimonious model (ΔAICc < 2 and

604   lowest number of parameters *K*) is shown in bold. For these models, we used a distance threshold of 1km

605   to consider that lions visited a waterhole.

Duration of visit to a waterhole – 1km		Female lion			Male lion		
Candidate models	**K**	**AICc**	**ΔAICc**	**R²**	**AICc**	**ΔAICc**	**R²**
Null hypothesis							
Number of waterholes	4	**4026.6**	**0.0**	**0.03**	5020.0	6.1	0.06
Prey abundance hypothesis							
Herd	5	4028.5	1.9	0.03	5019.1	5.1	0.07
Buffalo	5	4028.4	1.8	0.03	5018.0	4.1	0.07
Herd + Buffalo	6	4030.0	3.4	0.03	5018.8	4.9	0.08
Functional response in prey-abundance hypothesis							
Herd × Herd Territory	7	4030.1	3.5	0.04	5020.4	6.5	0.08
Buffalo × Buffalo Territory	7	4031.8	5.2	0.03	5021.5	7.6	0.07
Herd × Herd Territory + Buffalo × Buffalo Territory	10	4035.6	9.0	0.05	5024.6	10.7	0.09
Prey catchability hypothesis							
Distance to cover	5	4028.2	1.6	0.03	5020.3	6.3	0.06
Vegetation openness	5	4028.0	1.4	0.03	**5015.8**	**1.9**	**0.08**
Distance to cover + Vegetation openness	6	4030.0	3.4	0.03	5017.9	4.0	0.08
Functional response in prey-catchability hypothesis							
Distance to cover × Herd Territory	7	4030.3	3.8	0.04	5022.1	8.2	0.07
Vegetation openness × Herd Territory	7	4028.4	1.8	0.05	5017.5	3.5	0.08
Distance to cover × Buffalo Territory	7	4031.7	5.1	0.03	5021.6	7.7	0.07
Vegetation openness × Buffalo Territory	7	4027.5	0.9	0.05	5016.0	2.1	0.09
Distance to cover × (Herd Territory + Buffalo Territory)	9	4034.2	7.6	0.04	5023.3	9.4	0.08

Model	df	AIC	Δ	w	AIC	Δ	w
Vegetation openness × (Herd Territory + Buffalo Territory)	9	4030.0	3.4	0.06	5016.2	2.3	0.10
Vegetation openness + Distance to cover × (Herd Territory + Buffalo Territory)	10	4036.2	9.6	0.04	5021.1	7.2	0.10
Distance to cover + Vegetation openness × (Herd Territory + Buffalo Territory)	10	4032.1	5.5	0.06	5018.3	4.4	0.10
Prey abundance + Prey catchability hypothesis							
Herd + Distance to cover	6	4030.2	3.6	0.03	5019.6	5.6	0.07
Herd + Vegetation openness	6	4030.0	3.4	0.03	5016.8	2.8	0.08
Herd + Distance to cover + Vegetation openness	7	4032.0	5.4	0.03	5018.8	4.9	0.08
Buffalo + Distance to cover	6	4029.9	3.3	0.03	5018.6	4.6	0.08
Buffalo + Vegetation openness	6	4029.7	3.1	0.03	5013.9	0.0	0.09
Buffalo + Distance to cover + Vegetation openness	7	4031.7	5.1	0.03	5016.0	2.1	0.09
Herd + Buffalo + Distance to cover	7	4031.5	4.9	0.03	5019.5	5.5	0.08
Herd + Buffalo + Vegetation openness	7	4031.6	5.0	0.03	5015.8	1.9	0.09
Herd + Buffalo + Distance to cover + vegetation openness	8	4033.6	7.0	0.03	5017.9	4.0	0.09
Prey abundance × Prey catchability hypothesis							
Herd × Distance to cover	7	4032.1	5.5	0.03	5021.2	7.3	0.07
Herd × vegetation openness	**7**	4030.1	3.5	0.04	5018.8	4.9	0.08
Herd × (Distance to cover + Vegetation openness)	9	4034.2	7.6	0.04	5021.6	7.6	0.08
Buffalo × Distance to cover	7	4031.8	5.2	0.03	5019.5	5.6	0.08
Buffalo × Vegetation openness	7	4030.2	3.6	0.04	5016.0	2.1	0.09
Buffalo × (Distance to cover + Vegetation openness)	9	4033.2	6.6	0.04	5018.7	4.8	0.09
Herd × Distance to cover + Vegetation openness	8	4034.0	7.4	0.03	5019.9	5.9	0.08
Herd × Vegetation openness + Distance to cover	8	4032.1	5.5	0.04	5020.9	7.0	0.08
Buffalo × Distance to cover + Vegetation openness	8	4033.6	7.0	0.03	5016.8	2.9	0.09
Buffalo × Vegetation openness + Distance to cover	8	4032.2	5.6	0.04	5018.1	4.1	0.09
(Herd + Buffalo) × Distance to cover + Vegetation openness	10	4036.9	10.3	0.04	5020.3	6.4	0.10
(Herd + Buffalo) × Vegetation openness + Distance to cover	10	4035.5	8.9	0.04	5022.1	8.1	0.09
(Herd + Buffalo) × (Vegetation openness + Distance to cover)	12	4038.7	12.1	0.05	5023.5	9.6	0.10

606 **Figures**

609 **Figure 1.** A patch of grassland surrounding a waterhole in Hwange National Park, Zimbabwe. Photo

610 Google Earth Pro ©

Figure 2. Distribution of the frequencies and durations of lion visits to a specific waterhole during the dry

season. We used a 1km distance threshold to define lion visits to a waterhole. For male lions, one visit to

a waterhole lasting 76h is not shown in the plot. Waterholes along the x-axis are ordered by the average

frequency of visits by female lions (a) and male lions (b) or by the average duration of visits by female

lions (c) or male lions (d). The x-axis shows the name of each waterhole. Each dot shows the frequency or

duration of visits to the corresponding waterhole for one animal-year.

Figure 3. Influence of prey abundance and catchability on the frequency of visits to a specific waterhole

by female (yellow) and male (red) lions. Ribbon extremities show 95% confidence interval, whereas lines

show the mean value of the response variable. Predictions are made from models using a 1km distance

threshold to consider that lions visit a waterhole. The dotted horizontal line shows the frequency of visits

to a waterhole expected from the null hypothesis (i.e., the frequency of lion visits to a waterhole solely

depends on the number of waterholes within their territory). Prey-related metrics are centred and scaled

per animal-year territory to compare the relative abundance and the relative catchability of prey from one

waterhole to another waterhole available in the animal-year territory.

Figure 4. Influence of the number of waterholes within lions territories and of the relative vegetation openness in the surroundings of a waterhole on the durations of visits to a specific waterhole by female (yellow) and male (red) lions. Ribbon extremities show 95% confidence interval, whereas lines show the mean value of the response variable. Predictions are made from models using a 1km distance threshold to consider that lions visit a waterhole. The dotted horizontal line shows the duration of visits to a waterhole expected from the null hypothesis (i.e., the duration of lion visits to a waterhole solely depends on the number of waterholes within their territory). For male lions, one visit to a waterhole lasting 76h is not shown in the plot.

 Supplementary Information

Table S1. Results using a 2km distance threshold to define lion visits to a waterhole. Summary of the

candidate models explaining the variability in the frequency of lion visits to a waterhole. The frequency

of lion visits to a waterhole is calculated as the number of visits over 100 days (between early July and

mid-October; i.e., during the late dry season). All models included the number of waterholes within the

territory of each animal-year and a random effect with the identity of each animal-year. The most

parsimonious model (ΔAICc < 2 and lowest number of parameters K) is shown in bold.

Frequency of visits to a waterhole – 2km		Female lion			Male lion		
Candidate models	**K**	**AICc**	**ΔAICc**	**R²**	**AICc**	**ΔAICc**	**R²**
Null hypothesis							
Number of waterholes	4	1939,6	17,7	0,32	2652,8	64,8	0,06
Prey abundance hypothesis							
Herd	5	1933,9	12,0	0,35	2643,5	55,4	0,10
Buffalo	5	1937,9	16,0	0,33	2651,3	63,2	0,07
Herd + Buffalo	6	1935,0	13,1	0,35	2645,0	56,9	0,10
Functional response in prey-abundance hypothesis							
Herd × Absolute Herd	7	1936,4	14,5	0,35	2644,8	56,7	0,11
Buffalo × Absolute Buffalo	7	1937,2	15,3	0,34	2653,9	65,8	0,08
Herd × Absolute Herd + Buffalo × Absolute Buffalo	10	1939,6	17,7	0,36	2649,6	61,6	0,11
Prey catchability hypothesis							
Distance to cover	5	1939,1	17,1	0,32	2647,3	59,3	0,09
Vegetation openness	5	1924,5	2,6	0,35	2593,7	5,7	0,23
Distance to cover + Vegetation openness	6	1926,6	4,6	0,35	2594,5	6,4	0,23
Functional response in prey-catchability hypothesis							
Distance to cover × Absolute Herd	7	1941,6	19,7	0,33	2648,9	60,9	0,10
Vegetation openness × Absolute Herd	7	1927,0	5,0	0,36	2595,0	6,9	0,23
Distance to cover × Absolute Buffalo	7	1942,1	20,2	0,33	2645,1	57,0	0,11
Vegetation openness × Absolute Buffalo	7	1928,0	6,1	0,35	2591,2	3,1	0,25
Distance to cover × (Absolute Herd + Absolute Buffalo)	9	1944,0	22,1	0,34	2645,4	57,3	0,13

Model	k						
Vegetation openness × (Absolute Herd + Absolute Buffalo)	9	1930,0	8,1	0,36	2591,1	3,0	0,26
Vegetation openness + Distance to cover × (Absolute Herd + Absolute Buffalo)	10	1930,2	8,3	0,37	2589,7	1,6	0,27
Distance to cover + Vegetation openness × (Absolute Herd + Absolute Buffalo)	10	1932,1	10,2	0,36	2592,4	4,3	0,26
Prey abundance + Prey catchability hypothesis							
Herd + Distance to cover	6	1933,9	12,0	0,35	2640,6	52,5	0,11
Herd + Vegetation openness	6	**1923,6**	**1,7**	**0,36**	2593,5	5,4	0,24
Herd + Distance to cover + Habitat openness	7	1925,7	3,8	0,36	2594,3	6,2	0,24
Buffalo + Distance to cover	6	1938,1	16,1	0,33	2646,2	58,2	0,10
Buffalo + Vegetation openness	6	**1923,6**	**1,7**	**0,36**	**2589,8**	**1,8**	**0,25**
Buffalo + Distance to cover + Vegetation openness	7	1925,5	3,6	0,36	2590,2	2,1	0,25
Herd + Buffalo + Distance to cover	7	1935,3	13,4	0,35	2642,0	54,0	0,12
Herd + Buffalo + Vegetation openness	7	1924,3	2,4	0,36	2591,5	3,4	0,25
Herd + Buffalo + Distance to cover + Vegetation openness	8	1926,3	4,4	0,36	2591,9	3,8	0,25
Prey abundance × Prey catchability hypothesis							
Herd × Distance to cover	7	1935,4	13,5	0,35	2637,4	49,3	0,13
Herd × Habitat openness	7	1921,9	0	0,37	2595,4	7,4	0,24
Herd × (Distance to cover + Vegetation openness)	9	1925,9	3,9	0,37	2598,1	10,0	0,24
Buffalo × Distance to cover	7	1939,0	17,1	0,34	2646,1	58,1	0,11
Buffalo × Vegetation openness	7	1925,5	3,6	0,35	2591,0	2,9	0,25
Buffalo × (Distance to cover + Vegetation openness)	9	1927,1	5,2	0,36	2590,1	2,0	0,26
Herd × Distance to cover + Vegetation openness	8	1927,7	5,7	0,36	2596,1	8,0	0,24
Herd × Vegetation openness + Distance to cover	8	1924,0	2,1	0,37	2596,1	8,1	0,24
Buffalo × Distance to cover + Vegetation openness	8	1925,6	3,6	0,36	2588,1	0	0,26
Buffalo × Vegetation openness + Distance to cover	8	1927,4	5,5	0,35	2591,4	3,4	0,25
(Herd + Buffalo) × Distance to cover + Habitat openness	10	1927,7	5,8	0,37	2590,8	2,7	0,27
(Herd + Buffalo) × Vegetation openness + Distance to cover	10	1926,4	4,5	0,37	2593,7	5,6	0,26
(Herd + Buffalo) × (Vegetation openness + Distance to cover)	12	1928,5	6,6	0,38	2594,4	6,3	0,27

 Table S2. Results using a 2km distance threshold to define lion visits to a waterhole. Summary of the

650   candidate models explaining the variability in the duration of lion visits to a waterhole. All models

651   included the number of waterholes within the territory of each animal-year and a random effect with the

652   identity of each animal-year. The most parsimonious model (ΔAICc < 2 and lowest number of parameters

653   *K*) is shown in bold.

Duration of visit to a waterhole – 2km		Female lion			Male lion		
Candidate models	**K**	**AICc**	**ΔAICc**	**R²**	**AICc**	**ΔAICc**	**R²**
Null hypothesis							
Number of waterholes	4	**4684,2**	**5,0**	**0,0**	5742,3	6,5	0,07
Prey abundance hypothesis							
Herd	5	4684,6	5,5	0,01	5740,2	4,3	0,08
Buffalo	5	4685,4	6,3	0,0	5744,4	8,5	0,07
Herd + Buffalo	6	4686,5	7,3	0,01	5742	6,1	0,08
Functional response in prey-abundance hypothesis							
Herd × Absolute Herd	7	4684,3	5,2	0,04	5741,2	5,4	0,10
Buffalo × Absolute Buffalo	7	4687,1	7,9	0,02	5747,8	12,0	0,07
Herd × Absolute Herd + Buffalo × Absolute Buffalo	10	4689,5	10,3	0,04	5747,2	11,3	0,10
Prey catchability hypothesis							
Distance to cover	5	4684,3	5,2	0,01	5743,5	7,6	0,07
Vegetation openness	5	4684,6	5,5	0,01	**5736,0**	**0,1**	**0,09**
Distance to cover + Habitat openness	6	4681,4	2,2	0,03	5737,7	1,8	0,09
Functional response in prey-catchability hypothesis							
Distance to cover × Absolute Herd	7	4684,8	5,6	0,03	5744,2	8,3	0,09
Vegetation openness × Absolute Herd	7	4684,2	5,1	0,04	5737,0	1,1	0,11
Distance to cover × Absolute Buffalo	7	4687,0	7,8	0,01	5746,8	10,9	0,08
Vegetation openness × Absolute Buffalo	7	4683,1	3,9	0,03	5739,0	3,1	0,10
Distance to cover × (Absolute Herd + Absolute Buffalo)	9	4688,2	9,0	0,04	5748	12,1	0,09

Vegetation openness × (Absolute Herd + Absolute Buffalo)	9	4684,6	5,4	0,05	5740,6 4,7	0,11
Vegetation openness + Distance to cover × (Absolute Herd + Absolute Buffalo)	10	4684,1	5,0	0,06	5742,1 6,2	0,11
Distance to cover + Vegetation openness × (Absolute Herd + Absolute Buffalo)	10	4680,0	0,8	0,08	5742,4 6,5	0,11

Prey abundance + Prey catchability hypothesis

Herd + Distance to cover	6	4684,7	5,5	0,01	5741,7 5,8	0,08
Herd + Vegetation openness	6	4685,5	6,4	0,01	5735,9 0,0	0,10
Herd + Distance to cover + Vegetation openness	7	4682,5	3,4	0,03	5737,6 1,7	0,10
Buffalo + Distance to cover	6	4685,2	6,0	0,01	5745,5 9,6	0,07
Buffalo + Vegetation openness	6	4685,8	6,7	0,01	5738,0 2,1	0,09
Buffalo + Distance to cover + Vegetation openness	7	4681,5	2,3	0,03	5739,7 3,8	0,09
Herd + Buffalo + Distance to cover	7	4686,3	7,1	0,01	5743,4 7,5	0,08
Herd + Buffalo + Vegetation openness	7	4687,3	8,1	0,01	5737,9 2,0	0,10
Herd + Buffalo + Distance to cover + Vegetation openness	8	4683,4	4,3	0,03	5739,7 3,8	0,10

Prey abundance × Prey catchability hypothesis

Herd × Distance to cover	7	4685,9	6,7	0,02	5743,7 7,8	0,08
Herd × Habitat openness	**7**	4683,1	3,9	0,03	5737,6 1,7	0,10
Herd × (Distance to cover + Vegetation openness)	9	4682,7	3,5	0,04	5741,5 5,6	0,10
Buffalo × Distance to cover	7	4687,3	8,1	0,01	5747,6 11,7	0,07
Buffalo × Vegetation openness	7	4683,9	4,7	0,02	5739,9 4,0	0,09
Buffalo × (Distance to cover + Vegetation openness)	9	4681,2	2,0	0,05	5743,7 7,8	0,09
Herd × Distance to cover + Vegetation openness	8	4683,4	4,2	0,03	5739,6 3,7	0,10
Herd × Habitat openness + Distance to cover	8	4680,6	1,4	0,04	5739,4 3,5	0,10
Buffalo × Distance to cover + Vegetation openness	8	4683,6	4,4	0,03	5741,7 5,8	0,09
Buffalo × Vegetation openness + Distance to cover	8	4679,2	0	0,05	5741,6 5,7	0,09
(Herd + Buffalo) × Distance to cover + Vegetation openness	10	4686,1	7,0	0,04	5743,8 7,9	0,10
(Herd + Buffalo) × Vegetation openness + Distance to cover	10	4681,7	2,5	0,05	5743,5 7,6	0,10
(Herd + Buffalo) × (Vegetation openness + Distance to cover)	12	4685,4	6,2	0,06	5747,7 11,8	0,10

655 **Table S3.** Herd size of medium-sized prey in Hwange National Park. These sizes were calculated with

656 both adult and juvenile individuals.

Species	Mean	Lower CI 95%	Upper CI 95%
Eland	6.3	4.5	8.0
Greater kudu	4.4	4.3	4.5
Roan antelope	3.8	3.5	4.1
Sable antelope	4.7	4.3	5.1
Blue wildebeest	18.2	16.1	20.3
Plains zebra	7.3	7.1	7.5

660 **Table S4.** Coefficients, standard errors and p-values for the negative binomial regressions testing the

661 influence of prey abundance and catchability on the frequency of visits to waterholes by male lions.

662 Positive coefficients indicate for higher frequencies of visits.

Predictor	Estimate	Std. Error	z value	p-value
(Intercept)	2.121	0.048	44.38	< 0.001
Number of waterholes	-0.263	0.054	-4.86	< 0.001
Buffalo	0.149	0.051	2.93	< 0.05
Habitat openness	0.417	0.049	8.54	< 0.001

665 **Table S5.** Coefficients, standard errors and p-values for the negative binomial regressions testing the

666 influence of prey abundance and catchability on the frequency of visits to waterholes by female lions.

Predictor	Estimate	Std. Error	z value	p-value
(Intercept)	1.773	0.067	26.27	< 0.001
Number of waterholes	-0.642	0.075	-8.53	< 0.001
Herd	0.158	0.053	2.95	< 0.05
Habitat openness	0.114	0.053	2.17	< 0.05

667 Positive coefficients indicate for higher frequencies of visits.

670 **Table S6.** Coefficients, standard errors and p-values for the negative binomial regressions testing the

671 influence of prey abundance and catchability on the duration of visits to waterholes by female lions.

672 Positive coefficients indicate for higher durations of visits.

Predictor	Estimate	Std. Error	z value	p-value
(Intercept)	6.114	0.060	102.52	< 0.001
Number of waterholes	0.155	0.069	2.25	< 0.05

675 **Table S7**. Coefficients, standard errors and p-values for the negative binomial regressions testing the

676 influence of prey abundance and catchability on the duration of visits to waterholes by male lions.

677 Positive coefficients indicate for higher durations of visits.

Predictor	Estimate	Std. Error	z value	p-value
(Intercept)	5.803	0.050	115.42	< 0.001
Number of waterholes	-0.203	0.058	-3.52	< 0.001

Predictor	Estimate	Std. Error	z value	p-value
Habitat openness	0.110	0.044	2.51	< 0.05

Figure S1. Relationship between the duration and the frequency of lion visits to a specific waterhole for females (yellow) and males (red). The duration of visits to a waterhole is in minutes but log-transform for graphical representations. Vertical dashed lines show visits to a waterhole lasting 1h, 6h and 24h.