Ratiometric signaling produces robust temporal integration for accurate cellular gradient sensing

Debraj Ghose*

Wyss Institute for Biologically Inspired Engineering,

Harvard University, Boston, MA, 02215, USA

James Nolen*

Department of Mathematics, Duke University, Durham, NC, 27708, USA

Kaiyun Guan

Curriculum in Bioinformatics and Computational Biology,

University of North Carolina at Chapel Hill, Chapel Hill, NC, 27599, USA

Timothy C. Elston

Department of Pharmacology and Computational Medicine Program,

University of North Carolina at Chapel Hill, Chapel Hill, NC, 27599, USA

Daniel J. Lew†

Department of Biology, Massachusetts Institute of Technology, Cambridge, MA, 02139, USA

(Dated: March 2025)

Abstract

Cells excel at interpreting noisy chemical gradients to guide fertilization, development, and immune responses, but the mechanisms underlying this remarkable ability remain poorly understood. Previous work showed that some G protein signaling pathways can overcome challenges from uneven receptor distribution by using a ratiometric signaling strategy. In this mechanism, G proteins receive information from both bound and unbound receptors, unlike classical signaling where only bound receptors contribute. Here, we show that ratiometric signaling also provides an unexpected ability to suppress noise from low receptor numbers. The benefit stems from each G protein remembering the last receptor state it encountered, so that at any instant, ratiometric G protein collectives reflect time-averaged receptor activity. Unlike classical signaling, this averaging remains unbiased and accurate across the varying ligand concentrations present in a spatial gradient. Using theory and simulations, we demonstrate that this averaging mechanism allows cells to surpass theoretical limits for gradient detection from instantaneous receptor information alone. Our findings reveal how ratiometric biochemical architectures enable robust temporal integration across spatially varying signals, providing cells with enhanced directional accuracy under noisy conditions.

I.   INTRODUCTION

Cells are exceptional at extracting directional information from their chemical environment, often under complex and noisy conditions that demand high accuracy. Some of the earliest direct experimental evidence that cells respond to chemical gradients emerged in the late 19th century, with studies involving bacteria clustering near oxygen-rich zones [1, 2] and plant spermatozoids finding female reproductive organs [3]. Since then, chemotactic or chemotropic responses by cells or subcellular components have been documented in pollen tube guidance [4, 5], protozoa [6–8], fungi [9–14], immune cells [15, 16], animal sperm [17–19], neurons [20, 21], developmental processes [22], cancer cells [23–25], and many other systems. Together, these observations broadly illustrate the diverse scenarios under which gradient-driven cell guidance occurs.

---
* These authors contributed equally to this work.

† djlew@mit.edu

In all cases, cells detect chemical signals through binding to cell-surface receptors that engage internal signaling pathways [26, 27]. Using these receptors, a cell can in principle estimate the direction of a ligand gradient by measuring the difference in chemical concentration across its width. However, the rapid movement of fast-swimming cells like bacteria or sperm causes them to collide with ligand molecules more often at their front than at their back. This differential collision rate can make the concentration gradients they perceive across their width appear hundreds of times steeper than the actual environmental gradients [28]. Such cells have developed "temporal sensing" strategies that monitor concentration changes over time as they move through space. In contrast, a majority of larger eukaryotic cells, which are slower moving or non-motile, employ a "spatial sensing" strategy that compares concentrations of chemicals on different sides of the cell to infer gradient direction. Once a direction is established, eukaryotic cells can polarize and move or grow toward (or away from) higher ligand concentrations. Even so, these spatial measurements can be limited by gradient measurements with low signal to noise, especially for smaller cells. These limits have been theoretically studied for both absolute ligand concentration measurement [28–30] and gradient measurement [31, 32], grounded in considerations of ligand-binding noise. A well-appreciated strategy to enhance low signal is to use time-averaging of receptor signals, which can significantly improve both concentration measurement and gradient detection. However, molecular mechanisms for averaging receptor signals and improving gradient detection remain poorly understood.

In classical G protein signaling, G proteins are activated by ligand-bound receptors and deactivated at a uniform rate. Here, inspired by the mating response of budding yeast, we examined an alternate strategy: ratiometric signaling, in which G proteins are activated by bound receptors and deactivated by unbound receptors [33, 34]. This architecture means that downstream signaling depends on the ratio of active to inactive receptor levels rather than absolute number of active receptors.

We discovered that ratiometric signaling enables robust time averaging that significantly outperforms classical signaling. This mechanism arises from collectives of downstream G proteins that remember past receptor states. While classical signaling fails to maintain accurate averaging across the varying concentrations present in a spatial gradient, ratiometric signaling automatically adjusts to different concentrations. Furthermore, this mechanism allows cells to surpass theoretical limits for instantaneous gradient sensing, revealing how

60 cells achieve their remarkable ability to interpret noisy directional signals. These results
61 reveal design principles underlying robust spatial gradient sensing and suggest strategies for
62 engineering synthetic systems that can navigate complex chemical environments.

63 II. RESULTS

64 A. Instantaneous limits for gradient estimation by receptors

To evaluate the potential benefits or pitfalls of a ratiometric sensing strategy, we need
a metric that tells us how accurately a theoretical cell can infer the direction of a ligand
gradient. Previous studies have compared the ligand concentrations sensed by the front
and back halves of a cell, thereby determining whether the difference (signal) exceeds the
noise [28, 31]. However, splitting the cell in half presupposes prior knowledge of the axis
along which concentration changes occur. Because a real cell generally does not have that
information, other work has instead estimated the cell's "best guess" of the gradient direction
by summing the vectors from the cell center to all active receptors [32, 35] (Fig. 1A).

The resulting active receptor vector, V_{RL}, provides an estimate of the gradient direction
pointing towards the highest ligand concentration. Comparing V_{RL} with the actual gradient
direction provides an angular measure of how closely the cell's guess matches reality. In
principle, one could also compute the gradient direction from inactive receptors alone to
produce a "best guess" vector V_R pointing towards the lowest ligand concentration. To
combine information from both active and inactive receptors, we can calculate $V_C = V_{RL} - V_R$
(Fig. 1A).

In what follows, we simulate a spherical cell in three dimensions, with N_R receptors ran-
domly distributed over its surface. As receptor diffusion on the membrane is slow, especially
in budding yeast ($\sim 0.0005 \mu m^2/s$) [33, 36], we neglect it. Receptors can switch between ac-
tive and inactive states at rates that depend on local ligand concentration $C(x)$; in statistical
equilibrium, the probability of a receptor being active at a given time is

$$\mathbb{P}\big(\text{receptor at } x \text{ is active}\big) = \frac{C(x)}{C(x) + K_d}. \tag{1}$$

85 We expose the spherical cell to linear ligand gradients of the form

$$C(x, \theta) = C_0(1 + b(x \cdot \theta)/L), \tag{2}$$

where θ is the direction vector, C_0 is the mid-point (average) ligand concentration, and x is a point on the sphere of radius L centered at the origin. We first assume the gradient is centered around K_d (i.e., $C_0 = K_d$). Molecular noise from ligand binding/unbinding causes estimates of the gradient direction θ to fluctuate over time. We quantify directional accuracy by examining the empirical cumulative distribution functions for the angular deviation of $\bar{V}_{RL}$, $-\bar{V}_R$, and $\bar{V}_C$ (the normalized summed vectors) relative to θ across many independent realizations (Fig. 1 B i-iii).

As $\bar{V}_c$, $\bar{V}_{RL}$, and $-\bar{V}_R$ are each statistical estimators of gradient direction, we can compare them to the Cramér-Rao bound [37], which provides a rigorous lower bound on the variance of $\bar{V}$ from the true direction θ, relative to the signal strength $\|\mathbb{E}[\bar{V}]\|$. Accuracy of the direction estimate is quantified by a "noise-to-signal ratio" (NSR), and the statistical bound takes the form

$$NSR \geq \mathrm{trace}\big(\mathcal{I}_R^{-1}\big) \tag{3}$$

where $\mathcal{I}_R$ is a Fisher information matrix that can be computed numerically. In the weak-signal regime (small slope, b) this bound is approximated by

$$NSR \geq \mathrm{trace}\big(\mathcal{I}_R^{-1}\big) \;\approx\; \frac{6}{N_R\,b^2}\,\frac{\big(C_0 + K_d\big)^2}{K_d\,C_0}, \tag{4}$$

which shows that the minimal achievable variance in the estimated direction decreases proportionally to $1/N_R$ and depends inversely on b^2, thereby highlighting the advantage of stronger gradients and higher receptor number (details of this bound and a derivation of (4) appear in the Supplementary Information). A related bound is derived in [32] for a different model.

A comparison with this theoretical lower bound (Equation 3) shows that the estimates $\bar{V}_c$, $\bar{V}_{RL}$, and $-\bar{V}_R$ are indeed noisy, leaving ample room for improvement, especially at low N_R (Fig. 1 C). At high N_R, the variance of $\bar{V}_c$, $-\bar{V}_R$, and $\bar{V}_{RL}$ decrease along with the Cramér-Rao bound, which scales as $1/N_R$ (Fig. 1C). For $C_0 = K_d$, the variance of the combined estimate $\bar{V}_C$ is roughly half that of either $\bar{V}_{RL}$ or $\bar{V}_R$, as expected from the $1/N_R$ scaling of the optimal variance and the fact that roughly half of the receptors are active in this regime. Effectively, $\bar{V}_C$ draws on twice the data that either $\bar{V}_{RL}$ or $-\bar{V}_R$ alone can provide, demonstrating that incorporating both ligand-bound and unbound receptors significantly enhances directional sensing.

To determine whether these conclusions extend to other gradients, we simulated a range of linear gradients covering different mean ligand concentrations (Fig. 1D). As the ligand concentration rises, saturation becomes more pronounced, so a larger fraction of receptors are active and the signal is diminished. Consistent with that expectation, V_{RL} performs increasingly poorly at higher concentrations (Figure 1D i-iii). Though $-V_R$ also degrades at high concentrations, it does so differently from V_{RL}. When the gradient is centered below K_d, V_{RL} outperforms $-V_R$ (Fig. 1D i); at K_d, V_{RL} and $-V_R$ perform similarly (Fig. 1D iii); above K_d, $-V_R$ becomes superior (Fig. 1D iii). This pattern suggests that at low ligand concentrations, the scarce active receptors provide more information than the many inactive ones, whereas at high ligand concentrations, the fewer inactive receptors carry more information. The combined vector V_C performs as well as or better than either V_{RL} or $-V_R$ across the same tested range (Fig. 1 D i-iii). Hence, pooling signals from both active and inactive receptors broadens the range of gradients that can be decoded accurately.

B. Ratiometric signaling biochemistry greatly enhances gradient estimation

Until now, we have imagined that a cell could apply some procedure to calculate V_C (and extract the direction $\bar{V}_C$), but we neglected the actual biochemical processes by which such calculations necessarily arise. Indeed, any downstream signaling step cannot preserve 100% of the information available at an earlier step, as noise in the interpretation mechanism will inevitably degrade the signal. For the ubiquitous and large family of G protein coupled receptors (GPCRs) that signal to downstream heterotrimeric G proteins, the biochemical mechanisms have been elucidated, and we consider here a simplified scheme in which a cell contains N_G G proteins that diffuse on the 2D surface of the spherical cell and can become activated or inactivated. Activation occurs with some probability upon collision between a diffusing inactive G protein and a ligand-bound receptor.

Following [33], we explore two different models of G protein inactivation (Fig. 2). In the *classical* model, inactivation occurs at a spatially uniform rate, k_i', that is independent of receptor occupancy (Fig. 2A). In contrast, in the *ratiometric* model, inactivation occurs with some probability upon collision between a diffusing active G protein and an unbound receptor (Fig. 2B). We adjust the uniform inactivation rate of the classical model so that matched classical and ratiometric simulations have the same steady-state abundance of

active G proteins, thereby permitting fair comparisons [33]. Note that unlike the ratiometric model, the classical model does not incorporate information about the spatial distribution of inactive receptors.

In a ratiometric system, because ligand binding simultaneously increases the number of active receptors and decreases the number of inactive receptors, these opposing effects can amplify fluctuations due to molecular noise [38]. Indeed, [39] suggests that for a fixed ligand concentration, a cell's measurement of ligand levels is noisier under a ratiometric (or "concerted") mechanism than under the classical mechanism. Nevertheless, this need not imply that direction sensing becomes less reliable. For example, suppose (F_r, B_r) and (F_c, B_c) are pairs of random variables representing the active G protein concentration near the cell's front (F) and back (B), under the ratiometric (r) or classical (c) mechanisms. It may be that $CV(F_c) < CV(F_r)$ and $CV(B_c) < CV(B_r)$, yet $F_r - B_r > F_c - B_c$ (or $F_r/B_r > F_c/B_c$) still holds with high probability. (CV denotes coefficient of variation.) In other words, although the "signal" at any given location may be noisier in the ratiometric mechanism, the overall direction information (e.g., the difference $F - B$ or ratio F/B) can remain more robust compared to the classical system (see Section 5 and Fig. S6–S8 in the Supplement).

To compare direction sensing between classical and ratiometric systems, we calculated resultant vectors for active G proteins in a manner analogous to the previously discussed active receptor vector V_{RL} (Fig. 1A). We denote the resultant vectors of active G proteins under the ratiometric and classical models by V_G^r and V_G^c, respectively, and their normalized directions by $\bar{V}_G^r$ and $\bar{V}_G^c$. For ligand gradients near the receptor K_d, the ratiometric model (V_G^r) yields a significantly more accurate estimate of the gradient direction than the classical model (V_G^c) as demonstrated by temporal traces (Fig. 2C) (Movies S1-S6) and cumulative distribution functions (Fig. 2D) of V_G^c direction relative to gradient direction. Consequently, ratiometric sensing markedly enhances the accuracy of spatial gradient detection, even though it may sacrifice some accuracy in estimating the overall ligand concentration.

We next compared the performance of V_G^r to that of the combined receptor vector V_C. In simulations with many receptors (large N_R), the receptor-based vector V_C performed better than V_G^r (Fig.3A). This was expected, as the smaller number of G proteins ($N_G = 2500$) cannot retain all directional information from the larger number of receptors ($N_R = 6000$). However, as N_R decreased, V_C became increasingly susceptible to stochastic noise. Re-

markably, V_G^r retained its performance at lower N_R values (Fig.3A, B), so that below some threshold in receptor count (approx. $N_R < 1280$), V_G^r actually outperformed V_C (Fig. 3C). Even more unexpectedly, for sufficiently low N_R (approx. $N_R < 800$), V_G^r outperformed the accuracy limit given by the theoretical Cramér–Rao bound for V_C with the same number of receptors (Fig.3C). The Cramér-Rao bound (3) applies only to direction estimates based on instantaneous receptor data (i.e., V_C, V_R, V_{RL}) and need not apply to V_G^r or V_G^c; our finding shows that the accuracy of V_G^r in this regime cannot be matched by any unbiased estimator based only on instantaneous receptors states at a single fixed time. While these simulations were carried out using custom code written in MATLAB 2024a, the surprising results were verified independently using the particle-based simulator SMOLDYN (Figures S1 and S2). Thus, ratiometric G protein signaling can continue to provide accurate directional information despite receptor scarcity.

C. Ratiometric sensing's enhanced gradient estimation arises from robust collective G protein memory

In all parameter regimes that we explored with our simulations, the ratiometric model V_G^r is more accurate in direction estimation than the classical model V_G^c. In some parameter regimes explored in our simulations, the ratiometric model V_G^r also outperforms both V_{RL} and V_C, and it even performs better than the theoretical bound for estimation based on instantaneous receptor states (Fig. 3C). Here we explore the reasons for this superior performance of the ratiometric model.

The observation that V_G^r may perform better than both V_{RL} and V_C, even better than the theoretical bound for estimation based on instantaneous receptor states, suggests that the distribution of active G proteins must carry more information than the instantaneous distribution of receptors alone. Because receptors and G proteins occupy binary states, the G protein population can only carry additional information if $N_G > N_R$. Indeed, reducing the number of G proteins lowers the accuracy of V_G^r (Fig. 4A).

We propose that when $N_G > N_R$, the collection of G proteins captures additional information by repeatedly sampling receptor states over time, thus storing past receptor states. The accuracy of the direction estimate V_G^r depends on the level of correlation in G protein states, which is governed by two key time scales. The first is the characteristic timescale of

receptor state switching, λ^{-1}, where

$$\lambda = k_{\text{on}} C + k_{\text{off}}, \tag{5}$$

k_{on} and k_{off} are on/off rates of ligand binding, and C is the local ligand concentration. The second is the timescale of diffusive encounter, τ_{dif}, which is the mean time between G protein-receptor collisions. This depends on the G protein diffusion constant (D_G) and receptor number (N_R):

$$\tau_{\text{dif}} \propto \frac{L^2}{N_R D_G} \ln\left(\frac{\delta}{r_*}\right), \tag{6}$$

where L is the cell's linear dimension (e.g., radius), r_* is the receptor interaction radius, and δ is the typical receptor spacing. In the ratiometric model, it is the product $\lambda\tau_{\text{dif}}$ that determines the degree to which the G protein states are correlated.

   We consider a simplified model with n G proteins diffusing in the neighborhood of a single receptor that stochastically switches between active and inactive states, switching to active at rate $k_{\text{on}}C$ and to inactive at rate k_{off} (Fig.4B). As the receptor is switching states, the n G proteins encounter the receptor at random intervals with mean collision time τ_{dif}. In the ratiometric version of this simplified model, G proteins activate upon encountering an active receptor but deactivate only upon collision with an inactive receptor. In this case, the mean fraction of active G proteins (A_n) is the same as the fraction of time the receptor is on: $\mathbb{E}[A_n] = p_R$, but the variance of A_n changes with τ_{dif} and λ. A lower value of λ would lead to the receptor spending more time in a given state, causing states of G proteins to become more correlated. We derive an expression for variance of A_n (see Supplement):

$$\text{Var}(A_n) = p_R(1 - p_R)\left[\frac{1}{n} + \left(1 - \frac{1}{n}\right)\frac{1}{1 + \lambda\tau_{\text{dif}}}\right]. \tag{7}$$

Thus, faster receptor switching (larger λ) or slower G protein diffusion (larger τ_{dif}) raises $\lambda\tau_{\text{dif}}$, which lowers $\text{Var}(A_n)$ and improves the accuracy of estimating p_R. In Section 4 of the Supplement, we explain how this mechanism works when there are multiple receptors distributed across the cell surface, and we show how the timescales λ^{-1} and τ_{dif} affect the reliability of the direction estimate V_G^r. In our simulations, increasing λ (by increasing k_{off} while keeping K_d constant) leads to improved gradient sensing accuracy (lower NSR) (4C left), while increasing D_G leads to degraded accuracy (higher NSR) (4C right).

   This analysis supports the striking observation that the noise-to-signal ratio of V_G^r is relatively insensitive to N_R across a range of values of N_R that extends well below N_G.

Figure 2E, Figure 2C, and Figure 3D each demonstrate this insensitivity over a broad range. In Figure 3D, at low receptor number, V_G^r maintains a low noise-to-signal ratio—even below what is theoretically achievable for V_C at the same N_R (i.e., below the Cramér–Rao bound for V_C). This occurs because historical sampling of multiple receptor states enables V_G^r to encode more information than is available in instantaneous receptor states. In Figure 4A, as N_G increases (with N_R fixed), the estimate V_G^r becomes more accurate than V_C, the noise-to-signal ratio of V_G^r ultimately dropping below the Cramér–Rao bound for V_C.

In principle, the classical model V_G^c could also benefit from this time averaging, yet we observe that V_G^c always performs worse than V_G^r. The reason for this difference in performance has to do with a dose-response bias that is inherent in the classical model. For concentration estimation (rather than gradient direction estimation), this bias was observed already in [35]. For the ratiometric model, each G protein's state corresponds to the state of the receptor it most recently encountered. Consequently, the probability $p_G^r(x)$ that a G protein at position x is active (in the ratiometric model) corresponds approximately to $p_R(x)$, the probability that a receptor at x is active. For the classical model, where a G protein's state is reset to inactivate at rate k_{gi}, the probability $p_G^c(x)$ that a G protein at x is active depends on how many receptors it encountered since its last resetting event. This number of encounters depends on k_{gi} and on the diffusive time scale τ_{dif}. In the supplement we derive the formula

$$p_G^c(x) \approx \frac{p_R(x)\mathbb{E}[N]}{1 + p_R(x)\mathbb{E}[N]} \tag{8}$$

where $\mathbb{E}[N] = \tau_{\text{off}}/\tau_{\text{dif}}$ is the mean number of encounters, and $\tau_{\text{off}} = k_{gi}^{-1}$. A consequence of this is that p_G^c is not equal to p_R across the cell, and typically the gradient of p_G^c is significantly flatter than that of p_R. Since the NSR for direction estimation is inversely proportional to the square of the ligand gradient (e.g., $1/b^2$ in (4)), this flattening of the gradient of p_G^c relative to that of p_R implies a loss of accuracy in direction estimation for V_G^c.

To see how this bias effect may affect gradient sensing, we compared simulations of the ratiometric and classical versions of the simplified model described above. In the classical version, the n G proteins become activated upon collision with an active receptor and deactivate spontaneously at a baseline rate k_{gi}. To compare these models, we examined the active G protein fraction (A_n) at three positions across the cell corresponding to different ligand concentrations: back (low concentration, $p_R = 0.14$), middle ($p_R = 0.5$), and front (high concentration, $p_R = 0.75$) (Fig. 4D). At the middle position, where $p_R = 0.5$, we can tune

k_{gi} so that the classical model matches the ratiometric model, with both accurately reflecting the receptor state. However, at the front and back, where receptors experience higher or lower ligand concentrations, the classical model with this fixed k_{gi} deviates from p_R, while the ratiometric model maintains an unbiased estimate across all three regions without any parameter adjustment. Consequently, the ratiometric model produces a steeper gradient from front to back than the classical case.

This observation suggests the possibility of optimally tuning k_{gi} to improve gradient sensing in the classical model. We therefore swept through different values of k_{gi} in our full cellular gradient-sensing simulations (Fig. 4E). While we identified an optimal k_{gi} value that minimized the noise-to-signal ratio, this optimized classical model still performed worse than the combined receptor estimate V_C and failed to surpass the instantaneous theoretical bound, which is markedly inferior to the ratiometric model (Fig. 3D). Thus, ratiometric signaling enables robust collective memory through unbiased local averaging of receptor states, allowing cells to surpass the instantaneous theoretical limit for gradient detection.

D.   Responsiveness to changing gradients

In the preceding discussion, we assumed that ligand gradients were stable over time. However, this is not always the case: neutrophils chase motile bacteria, social amoebae collectively generate fruiting bodies by following dynamically shifting cAMP signals, and budding yeast track their mating partners as both cells grow [10, 40, 41]. While the robust collective memory encoded in G protein states by ratiometric signaling improves accuracy at steady state, it may slow responses to changing gradient directions.

To assess how different parameters affect a cell's ability to adapt to changing gradients, we performed simulations in which we allowed V_G^r to reach steady state in a gradient and then instantaneously reversed the gradient direction by 180 degrees (Fig. 5A). We quantified responsiveness by measuring the time required for V_G^r to cross the 90-degree threshold toward the new gradient direction (Fig. 5B).

We first examined the effect of receptor switching kinetics by varying both ligand off-rate k_{off} and on-rate k_{on} while maintaining a constant K_d (Fig. 5C). While k_{on} is physically capped by the rate of diffusional encounters, our parameters remain biologically plausible; at $K_d = 6 \, \text{nM}$, the maximum $k_{off} = 0.1 \, \text{s}^{-1}$ corresponds to $k_{on} \approx 1.67 \times 10^7 \, \text{M}^{-1}\text{s}^{-1}$, which is well

within the Smoluchowski limit. Increasing k_{off} from 0.005 to 0.1 s^{-1} reduced both the time to adapt and the steady-state noise-to-signal ratio. These dual benefits can be understood through the receptor switching rate λ (Equation 5): increasing k_{off} increases λ, which directly accelerates receptor state transitions and reduces the time to achieve statistical equilibrium after the gradient direction is reversed. Simultaneously, higher λ increases the product $\lambda\tau_{\text{dif}}$; as we have discussed, this reduces correlation in downstream G protein states, and thus improves steady-state directional accuracy.

Increasing receptor density also accelerated adaptation to new gradient directions (Fig. 5D). Higher N_R reduces the mean time between G protein–receptor encounters (τ_{dif}), allowing G proteins to update their states more frequently and thereby adapt more quickly to the reversed gradient. As we have already noted, the noise-to-signal ratio of V_G^r (at steady state) may be relatively insensitive to changes in N_R over a certain range, depending on N_G (e.g., as N_R increased from 375 to 6000 in Fig. 5D).

In contrast to receptor kinetics and density, increasing the G protein diffusion constant D_G presented a trade-off between adaptation speed and steady-state accuracy (Fig. 5E). Faster diffusion reduces τ_{dif}, enabling more frequent G protein–receptor encounters and thereby accelerating adaptation to the new gradient direction. However, unlike increasing N_R, faster diffusion of G protein increases correlation in G protein states and increases the noise-to-signal ratio of V_G^r at steady state.

Together, these results reveal distinct strategies by which cells could tune their responsiveness to changing gradients. Faster receptor kinetics provide the most favorable outcome, simultaneously enhancing both steady-state accuracy and adaptation speed. Increased receptor density primarily accelerates adaptation with minimal impact on steady-state performance. Higher G protein diffusion offers faster adaptation but compromises spatial precision—a trade-off that cells may exploit depending on whether their environment demands accurate gradient tracking or rapid reorientation.

 III.   DISCUSSION

   A.   Ratiometric signaling in the budding yeast system

    Gradient decoding is most challenging when cells are small, gradients are shallow, and chemical concentrations are low, but yeast cells find mates despite encountering all of these challenges. An effective strategy to extract signals from noisy measurements would be to integrate concentration measurements over time, allowing time averaging to reduce the noise (fluctuations) without affecting the signal (gradient). However, once a chemical has bound to a cell-surface receptor, diffusion of the receptor-ligand complex would degrade the infor- mation about the location where the binding took place. Such blurring can limit the interval over which time-averaging is effective for directional sensing[35]. This is particularly prob- lematic for systems like yeast pheromone sensing, where ligand-receptor complexes can per- sist for several minutes. Interestingly, yeast cells severely limit receptor diffusion ($D < 0.0005$ $\mu m^2/s$), perhaps mitigating this problem [33]. However, with very slow receptor diffusion, receptors often accumulate non-uniformly on one side of the cell [33, 42–44]. Non-uniform receptor distributions in turn can lead to spatial patterns of ligand-receptor complexes that do not accurately reflect the spatial distribution of ligands in the environment.

    Yeast cells appear to use a ratiometric sensing strategy to compensate for non-uniform receptor distributions [33]. Ligand binding to receptors promotes activation of G proteins that are then turned off by RGS proteins (Sst2 in yeast). In classical signaling models, G protein inactivation occurs in a receptor-independent manner, but in yeast cells the RGS protein is bound to inactive receptors [42]. Thus, whereas active receptors promote signal- ing, inactive receptors inhibit signaling, and the net signal reflects the ratio of active to inactive receptors [34]. If receptors are concentrated in some region of the cell surface, both active and inactive receptors may be locally enriched, so ratiometric sensing can allow the cell to accurately infer the local ligand distribution by compensating for the local receptor abundance. Interestingly, we find that, in addition to extracting information from both active and inactive receptors, ratiometric sensing could also better enable a cell to interpret temporally averaged receptor states over time.

B. Noise in concentration measurement from ratiometric signaling

In principle, ratiometric signaling could be a widely applicable strategy to improve the detection of spatial gradients, especially when gradient concentrations are near the ligand-receptor K_d. It seems intuitively plausible that a cell capable of using information about the spatial distributions of inactive as well as active receptors would have a firmer basis to infer the spatial distribution of ligand than a cell whose only information comes from the active receptors. However, as discussed in the Supplement and shown in [38, 39], ratiometric signaling can be less accurate at sensing absolute concentration. Nevertheless, we find that despite being worse at sensing ligand concentration, the ratiometric strategy provides more accurate gradient sensing by enabling acquisition of information from both active and inactive receptors as well as locally averaging receptor signals over time.

We show that ratiometric signaling outperforms classical signaling, even at very low receptor numbers when less than 0.02% of the cell surface is covered in receptors. Notably, these conditions resemble scenarios where cells initially populate their membranes with specific receptors—such as in lymphocytes like B cells, which upregulate receptors upon transitioning between germinal center zones; and in various developmental contexts where cells switch states and begin expressing new receptors [45, 46]. In such scenarios, ratiometric signaling could enable reliable gradient detection with lower receptor density, all while resisting misdirection by local fluctuations in receptor localization or ligand availability.

C. Mechanism underlying gradient detection accuracy by ratiometric signaling

One of our most intriguing observations was that ratiometric signaling could outperform the theoretical bound for gradient direction estimates that use only receptor snapshots at a single moment in time. This surprising advantage arises because downstream molecules—specifically, diffusing G proteins—act as a form of collective molecular memory. As each G protein updates its activity state only upon encountering a receptor, it encodes receptor information from an earlier time point. A population of G proteins that do this for both active and inactive receptors can smooth out short-lived fluctuations and enhance the fidelity of direction estimates. In effect, G proteins collectively time-average receptor states, allowing the cell to extract more robust spatial information than would be possible

from instantaneous receptor states alone.

In ratiometric signaling, two parameters critically balance directional accuracy and adaptive speed: the switching rate (λ) of receptor-ligand binding and the diffusion coefficient (D_G) of G proteins. Increasing λ enhances collective local averaging by G proteins (Fig. 4E (left) and 5C), improving steady-state gradient detection, and promotes faster adaptation to shifting gradients (Fig. 5C). In contrast, raising D_G weakens local averaging and diminishes steady state-gradient detection (Fig. 4E (right) and 5E). But higher D_G accelerates adaptation to changing signals, as more frequent receptor–G protein collisions swiftly update G protein states in response to new external conditions (Fig. 5C). Together, these trade-offs highlight how ratiometric signaling can be tuned to optimize either accuracy or speed, as required by the biological context.

D. Broader relevance of ratiometric signaling in responding to directional cues

It is not known whether receptors other than GPCRs can engage ratiometric signaling pathways. However, the principle underlying the averaging of upstream signals by more abundant downstream components may extend beyond receptor signaling, particularly in systems where both active and inactive states of upstream proteins convey meaningful information. In MAPK cascades, downstream kinases often exceed the abundance of their upstream regulators, exemplified by Raf in mammalian ERK pathways [47], B-Raf in cancer signaling contexts [48], and Ste11 in yeast mating responses [49]. These systems possess features that suggest the potential for ratiometric measurement-driven averaging to enhance accuracy and robustness.

Overall, our work suggests that ratiometric signaling mechanisms provide a powerful framework for cells to interpret directional receptor information and mitigate noise. The ability to sample both active and inactive states, combined with downstream emergent molecular memory, enables more accurate direction finding than that achieved by classical signaling mechanisms. These results offer testable predictions for experimental systems and may help clarify why some cells have a higher abundance of downstream proteins in signaling cascades.

IV. ACKNOWLEDGMENTS

This work was supported by NIH/NIGMS grant R35GM122488 to D.J.L. D.G. was supported by the Alfred P. Sloan Foundation's Matter-to-Life program.

V. DATA AVAILABILITY STATEMENT

The data and simulation code that support the findings of this study are openly available in the following GitHub repository: https://github.com/DebrajGhose/ratiometric-sensing-robust.

Appendix A: Simulation Framework

The simulation models the interactions between receptors and G proteins on a cell surface, taken to be a sphere $\mathcal{S}$ in three dimensions. The simulation focuses on their activation and inactivation dynamics under a pheromone gradient.

On the surface of the sphere, receptors are positioned randomly; their locations are fixed and do not change over time. Receptors also switch state randomly and independently of each other; they can be either ligand-bound (active) or unbound (inactive). The rate of activation depends on the local ligand concentration, which is taken to be time-independent.

The G proteins are also positioned randomly over the surface of the sphere, independently and uniformly distributed. Unlike the receptors, however, each G protein performs a diffusive random walk along the surface of $\mathcal{S}$. G proteins in the simulation can exist in either an active or inactive state. The state of a G protein is influenced by its interaction with nearby receptors and its inherent deactivation rate. G proteins switch state according to either the classical model or the ratiometric model. In both models, a G protein becomes activated upon encountering an active receptor. The difference in the two models lies in the mechanism for deactivation. In the classical model, a G protein becomes deactivated randomly at the inherent deactivation rate. In the ratiometric model, a G protein becomes deactivated by encountering an inactive receptor.

We use the following nomenclature:

$\mathcal{S}$	cell surface, a sphere in $\mathbb{R}^3$, centered at the origin
L	radius of the cell (2.5 μm for a yeast cell)
x	cartesian coordinate for a position on the sphere $\mathcal{S}$
θ	true direction of pheromone concentration gradient, $\theta \in \mathbb{R}^3$, $\|\theta\| = 1$
$C(x)$	local ligand concentration at position x on cell surface
b	gradient steepness parameter for ligand concentration, $b \in (0, 1)$
K_d	dissociation constant for pheromone-receptor binding (6 nM for yeast)
k_{off}	rate of receptor deactivation or ligand unbinding (typically $0.1 s^{-1}$ in our simulations)
k_{on}	rate constant for receptor activation, $k_{\text{on}} = k_{\text{off}}/K_d$
N_R	total number of receptors (from 375 to 6000 in our simulations)
$X_1, \ldots, X_{N_R}$	locations of the receptors on $\mathcal{S}$
$R_1, \ldots, R_{N_R}$	states of receptors, either 0 (inactive, unbound) or 1 (active, bound)
N_G	total number of G proteins on $\mathcal{S}$ (from 181 to 5000 in our simulations)
$Y_1, \ldots, Y_{N_G}$	locations of the G proteins on $\mathcal{S}$
$G_1, \ldots, G_{N_G}$	states of the G proteins, $G_j \in \{0, 1\}$.
r_*	activation radius for activation of G protein by an active receptor (5 nm)
D_G	diffusion constant for G protein (from 0.001 to 0.008 $\mu m^2/s$ in our simulations)
$k_i = 0$	G protein inactivation rate constant for ratiometric model
k_i'	G protein inactivation rate constant for nonratiometric model

1. Receptor dynamics

The receptor configuration at time t is characterized by the pairs $\{(X_k, R_k(t))\}_{k=1}^{N_R}$. The positions $\{X_k\}_{k=1}^{N_R}$ are sampled independently and uniformly on $\mathcal{S}$. The receptor positions are assumed to not change with time (we neglect receptor diffusion). Given a receptor at position $X_k = x$, the receptor state $R_k(t) \in \{0, 1\}$ changes as a Markov process with rates:

$$0 \mapsto 1 \qquad \text{at rate } k_{\text{on}}C(x), \tag{A1}$$

$$1 \mapsto 0 \qquad \text{at rate } k_{\text{off}}. \tag{A2}$$

The receptor states are initialized according to the stationary distribution. Therefore, at any time, the probability that a receptor at location $X_k = x$ is active is:

$$p_R(x) = \mathbb{P}(R_k(t) = 1 \mid X_k = x) = \frac{C(x)}{C(x) + K_d}, \qquad K_d = k_{\text{off}}/k_{\text{on}}.$$

The ligand concentration depends on the direction vector θ, the true gradient direction. We will write $C = C(x, \theta)$ to emphasize this dependence on θ. We take C to have the linear form

$$C(x, \theta) = C_0(1 + b(x \cdot \theta)/L), \quad x \in \mathcal{S}, \tag{A3}$$

where $C_0 > 0$ and $b \in (0, 1)$, so that $(1 - b)C_0 \leq C(x, \theta) \leq (1 + b)C_0$. Consequently, the probability $p_R(x)$ also depends on θ, and we emphasize this by writing $p_R(x, \theta)$.

As we will explain later, the rate parameter

$$\lambda_0 = k_{\text{on}}C_0 + k_{\text{off}}$$

plays an important role in the model; λ_0^{-1} has the units of $(time)$ and we refer to λ_0^{-1} as the receptor turnover time.

We consider values of N_R in the range 375 to 6000 in our simulations. For a cell radius $L = 2.5\mu m$ and receptor activation radius $r_* = 5nm$, the fraction of area covered by receptor is in the range of 0.0375% to 0.6%.

2. G protein dynamics

The G proteins are initialized independently, uniformly at random on the sphere, at positions $Y_1(0), \ldots, Y_{N_G}(0)$. These positions diffuse over $\mathcal{S}$ in time, independently and with diffusion constant D_G; statistically, their joint spatial distribution is invariant in time. An inactive G protein (state $G_j(t) = 0$) at location $Y_j(t)$ is activated upon interaction with an active receptor. Such activation occurs when $Y_j(t)$ diffuses within radius r_* of an active receptor. The simulation uses a spatial partitioning algorithm (KD-tree) to identify, at each time step, G proteins within a certain reaction radius of any receptor.

Once active ($G_j = 1$), the mechanism for de-activation of G protein is either the ratiometric model or the classical model. In the classical model, an active G protein deactivates at a fixed rate k_i', the inherent deactivation rate, but an active G protein is unaffected by encountering inactive receptors. In the ratiometric model, however, an active G protein deactivates upon diffusing within radius r_* of an inactive receptor and there is no basal deactivation ($k_i = 0$ in the ratiometric model). To compare the ratiometric and classical models, we choose k_i' so that at steady state, the mean fraction of active G protein is the

same as that for the ratiometric model. In particular, we adjust the parameter k_i' for each ligand concentration profile to maintain this comparison.

Depending on the random receptor configuration, it is possible for a G protein to be within the reaction radius of two or more receptors simultaneously. For the ratiometric model, when a G protein is near multiple receptors simultaneously the state (at each time step) of the G protein is matched that of a neighboring receptor chosen uniformly at random (from among those receptors within the reaction radius). Hence, if the majority of nearby receptors are active, the probability increases for the G protein to become active. Conversely, if most nearby receptors are inactive, the likelihood increases for the G protein to become inactive. This dynamic represents a ratiometric response of the G protein to the local receptor state.

3. Simulation methodology and verification

Simulations were performed using custom solvers in MATLAB (v2021b to v2024a) on a Linux-based computing system (Longleaf cluster at UNC Chapel Hill). Simulations were initialized by randomly and uniformly distributing receptor and G protein populations on the surface of a spherical cell. Receptor positions were kept stationary while G proteins performed random walks (Brownian motion) across the cell's surface with a fixed time step and diffusion constant. Receptor activation followed a first-order reaction, with an activation rate determined by the local pheromone concentration prescribed along the cell's x-axis to establish a gradient (Equation (A2) and A1). Activation and deactivation events were implemented via Monte Carlo sampling. In the classical receptor model, G proteins became active upon encountering active receptors within a specified reaction distance and deactivated at a constant rate. In the ratiometric model, G proteins were activated upon encountering active receptors within the same reaction distance and deactivated upon encountering inactive receptors. In the presence of multiple receptors, the ratio of active to inactive receptors governed the probability of G-protein activation or deactivation. For each set of parameters, 1000 simulations were run, tracking molecular states, positions, and time-resolved changes in receptor and G-protein activation. All MATLAB code for these simulations is available at: https://github.com/DebrajGhose/ratiometric-sensing-robust

To validate our solver, MATLAB simulations from Figures 1, 2D and 3C were replicated in Smoldyn (v2.71) with continuous space and discretized time intervals on a Linux-based

computing system (Longleaf cluster at UNC Chapel Hill). Periodic boundary conditions were applied in both spatial directions. Molecules were modeled as point particles without volumes. Brownian motion of the molecules was simulated using the Euler-Maruyama method. We used a time step of 0.005 secs and recorded the coordinates of receptors and G proteins every 10 secs over 20000 secs. The simulation methods used were consistent with the customized code, except for the setup of the pheromone gradient. As Smoldyn does not support continuously spatially dependent concentration, the cell sphere was divided into 11 vertical compartments along the gradient direction, with each compartment assigned a local pheromone concentration defined as:

$$P_{local,i} = P_{low} + (i-1)\frac{P_{high} - P_{low}}{N-1}$$

where P_{low} and P_{high} represent the lowest and highest pheromone concentrations, respectively. i is the compartment index. $N = 11$ is the total number of compartments. Accordingly, we modeled the receptor activation as a first-order reaction with a local activation rate in the ith compartment defined as:

$$k_{a,i} = \frac{k_d}{K_d} \times P_{local,i}$$

488 where $k_d = 0.1s^{-1}$ is the dissociation rate between the receptor and the ligand.

489 All code for Smoldyn simulations are available at: https://github.com/DebrajGhose/ratiometric-
490 sensing-robust Figure 6 and 7 show that Smoldyn simulations produce similar results to
491 those of our custom code.

Appendix B: Estimation model

At a given time t, we use the receptor data and the G protein data to compute various
estimates of the direction parameter θ, treated as an unknown quantity. The receptor
data at a fixed time are random variables $\{(X_k, R_k(t))\}_{k=1}^{N_R}$. These variables are pairwise
independent, and their distribution depends on the parameter θ. Thus, the receptor data
may be regarded as N_R independent points in $\mathcal{S} \times \{0,1\}$, and their joint density on $(\mathcal{S} \times$
$\{0,1\})^{N_R}$ is

$$\rho(\{(X_k, R_k)\}, \theta) = \prod_{k=1}^{N_R} \frac{1}{4\pi L^2} \left(p_R(X_k, \theta)\mathbb{I}_{R_k=1} + (1 - p_R(X_k, \theta))\mathbb{I}_{R_k=0} \right). \tag{B1}$$

The factor $4\pi L^2$ is the surface area of $\mathcal{S}$. The terms $\mathbb{I}_{R=1}$ and $\mathbb{I}_{R=0}$ are indicators of the events $R = 1$ and $R = 0$, respectively.

Using active receptor data, we define

$$V_{RL} = \sum_{k=1}^{N_R} X_k \mathbb{I}_{R_k=1} \tag{B2}$$

where $\mathbb{I}_{R_k=1}$ is the indicator of the event $R_k = 1$. Thus, V_{RL} is the sum of the positions of active receptors only. Similarly, using inactive receptors, we define

$$V_R = \sum_{k=1}^{N_R} X_k \mathbb{I}_{R_k=0}, \tag{B3}$$

and we regard the reflected vector $(-V_R)$ as an estimate of θ, since the gradient of inactive receptor will typically be in a direction anti-aligned with θ. The combined receptor estimate is

$$V_C = V_{RL} - V_R = \sum_{k=1}^{N_R} X_k \left(\mathbb{I}_{R_k=1} - \mathbb{I}_{R_k=0} \right).$$

These V_{RL}, $-V_R$, and V_C are vectors in $\mathbb{R}^3$. From these vectors, direction information is computed by normalizing

$$\bar{V}_C = \frac{V_C}{\|V_C\|}, \qquad \bar{V}_{RL} = \frac{V_{RL}}{\|V_{RL}\|}, \qquad -\bar{V}_R = \frac{-V_R}{\|V_R\|}.$$

Each of these direction estimates is a random unit vector, depending on time (because the receptor states change in time). To compute the distribution of these estimates, we obtain independent samples by re-initializing independently all of the receptor positions and states.

Using the G protein data, we compute the vector

$$V_G = \sum_{j=1}^{N_G} Y_j \mathbb{I}_{G_j=1} \tag{B4}$$

which is the sum of positions of all active G proteins. From this a direction estimate is computed by normalizing:

$$\bar{V}_G = \frac{V_G}{\|V_G\|}.$$

Where appropriate, we will use superscripts V_G^r and V_G^c to further distinguish the cases where the vector (B4) is computed under the ratiometric model versus the classical model, respectively. Like the estimates $\bar{V}_C$, $\bar{V}_{RL}$ and $-\bar{V}_R$, the estimate $\bar{V}_G$ is a random unit vector that depends on time. However, $\bar{V}_G(t)$ is not a function of the receptor data $\{X_k, R_k(t)\}$

at any fixed time; $\bar{V}_G(t)$ depends on receptor information from previous times, due to the way in which G proteins interact with receptors. We obtain n approximately independent estimates for $\bar{V}_G$ by computing $\bar{V}_G$ at sufficiently spaced time points $t_1 < t_2 < \cdots < t_n$ where $t_{k+1} - t_k \geq 1s$.

To compare these direction estimates with the true gradient direction θ, we compute the angular deviation $\arccos(\bar{V} \cdot \theta)$ of the estimate $\bar{V}$ $(= \bar{V}_C, \bar{V}_{RL}, -\bar{V}_R, \bar{V}_G)$ from the true direction θ. The empirical cumulative distributions of these angular deviations are shown in (Fig. 1B, 1D, 2D, 3C) of the main text – this function of $\alpha \in [0, 180]$ is the fraction of the random estimates $\bar{V}$ within α degrees of the true direction θ. In (Fig. 2C, 3A, 3B) of the main text, we also show the fluctuation of these estimates over time, while the receptor and G protein states change asynchronously (but receptor positions are fixed).

1. Cramér-Rao bound

The estimates $\bar{V}_C$, $\bar{V}_{RL}$ and $-\bar{V}_R$ are random variables; in fact, they are functions of the receptor data $\{(X_k, R_k)\}_{k=1}^{N_R}$ at a fixed time. As such, we examine their efficiency by comparing their variance to theoretical lower bounds provided by the Cramér-Rao theory. The present setting is complicated by the fact that the parameter space for the direction parameter θ is the unit sphere $\mathcal{S}_1$, which is not a linear space. Nevertheless, the variance of an estimator on the sphere can be bounded from below in terms of the inverse $\mathcal{I}_R^{-1}$ of the Fisher information matrix

$$\mathcal{I}_R = \mathbb{E}_\theta \left[(\mathrm{grad}_\theta \log \rho(Z,\theta))(\mathrm{grad}_\theta \log \rho(Z,\theta))^T \right], \tag{B5}$$

where $\rho(Z,\theta)$ is given by (B1) with $Z = \{X_k, R_k\}_{k=1}^{N_R}$, and $\mathbb{E}_\theta$ denotes integration with respect to density $\rho(\cdot,\theta)$, parameterized by θ. This matrix $\mathcal{I}$ should be regarded as a symmetric 2×2 matrix acting on the tangent space at $\theta \in \mathcal{S}_1$. An applicable Cramér-Rao type bound is given in [37]. (See also Chapter 13 of [50]). It follows from Theorem 3.2 and Example 4.2 of [37] that the estimate $\bar{V}$ must satisfy

$$NSR = \frac{\mathbb{E}_\theta \left[\|\theta^\perp \bar{V}\|^2 \right]}{\|\mathbb{E}_\theta \left[\bar{V} \right] \|^2} \geq \mathrm{trace}(\mathcal{I}_R^{-1}) \tag{B6}$$

where $\theta^\perp Y = (I - \theta\theta^T)Y$ is the projection of vector $Y \in \mathbb{R}^3$ onto the subspace orthogonal to the vector θ. By definition $\bar{V}$ is normalized so that $\|\bar{V}\| = 1$; however, $\|\mathbb{E}_\theta[\bar{V}]\| < 1$. The

law of large numbers implies that as $N_R \to \infty$, $\mathbb{E}_\theta[\bar{V}] \to \theta$ and $\|\mathbb{E}_\theta[\bar{V}]\| \to 1$. For a weak signal and with few receptors, however, $\|\mathbb{E}_\theta[\bar{V}]\|$ may be small. Nevertheless, (B6) holds as long as the estimator $\bar{V}$ is unbiased and $\|\mathbb{E}_\theta[\bar{V}]\| \neq 0$, which is the case here for each of the estimators $\bar{V} = \bar{V}_C, \bar{V}_{RL}, \bar{V}_R$, by the symmetry of our choice of $C(x, \theta)$. In particular, $\|\mathbb{E}_\theta[\bar{V}]\| = |\mathbb{E}_\theta[\bar{V} \cdot \theta]|$ holds in our setting. In the ratio appearing in (B6), $\|\theta^\perp \bar{V}\|^2$ is zero if and only if $\bar{V}$ lies in the line spanned by the true direction θ; on the other hand, $\|\theta^\perp \bar{V}\|^2$ is maximal when $\bar{V}$ is orthogonal to θ. We also note that $\|\theta^\perp \bar{V}\|^2 = \sin^2(\alpha)$ where $\alpha = \arccos(\bar{V} \cdot \theta)$ is the angular deviation between $\bar{V}$ and θ.

We refer to the ratio in (B6) as the "noise to signal" ratio (NSR) for the estimate V. It is similar to a squared coefficient of variation $(CV)^2$ for the normalized vector $\bar{V}$: the numerator is the variance of the normalized vector $\bar{V}$ in the directions orthogonal to θ, while the denominator is the mean of $\bar{V}$ squared. On the other hand, this ratio differs from the usual notion of coefficient of variation in that the quantity $\bar{V}$ is vector-valued, and the denominator $\mathbb{E}_\theta[\bar{V}]$ is not the mean of $\theta^\perp \bar{V}$. Another Cramér-Rao type bound is given by [51] (see Theorem 2 and Corollary 2):

$$\mathbb{E}_\theta\left[d^2(\bar{V}, \theta)\right] \geq \text{trace}(\mathcal{I}_R^{-1}) + \text{curvature terms}.$$

Here $d(u, v) = \arccos(u \cdot v)$ is the standard Riemannian distance between points u and v on the unit sphere, and the curvature terms (due to the positive curvature of $\mathcal{S}_1$) are negligible when $\mathbb{E}_\theta\left[d^2(\bar{V}, \theta)\right]$ is small (when N_R is sufficiently large).

The Fisher information matrix can be computed, using the fact that the joint density ρ in (B1) has a product structure (the pairs $\{(X_k, R_k)\}$ are independent and identically distributed). One finds that

$$\mathcal{I}_R = \frac{N_R}{4\pi L^2} \int_\mathcal{S} (1 - p_R(x, \theta)) p_R(x, \theta) (\text{grad}_\theta \log C(x, \theta))(\text{grad}_\theta \log C(x, \theta))^T \, dx \tag{B7}$$

$$= \frac{N_R}{4\pi L^2} \int_\mathcal{S} \frac{(\text{grad}_\theta p_R(x, \theta))(\text{grad}_\theta p_R(x, \theta))^T}{(1 - p_R(x, \theta)) p_R(x, \theta)} \, dx \tag{B8}$$

Assuming p_R is a function of $x \cdot \theta$, then this symmetry of p_R about θ implies that $\mathcal{I}_R$ is a multiple of the 2×2 identity matrix. For various choices of $C(x, \theta)$, we evaluate the integral in (B8) numerically, and then compute $\text{trace}(\mathcal{I}_R^{-1})$ appearing in (B6). For $\bar{V} = \bar{V}_C, \bar{V}_{RL}, -\bar{V}_R$, we approximate the ratio

$$NSR = \frac{\mathbb{E}_\theta\left[\|\theta^\perp \bar{V}\|^2\right]}{\|\mathbb{E}_\theta[\bar{V}]\|^2} \tag{B9}$$

and compare with the lower bound (B6). The expectations are approximated by generating independent samples of $\bar{V}$ (as described above) and averaging over these samples. The comparisons are shown in (Fig. 1C) of the main text for different total receptor numbers N_R.

The G protein configuration at time t is described by pairs $\{Y_j(t), G_j(t)\}_{j=1}^{N_G}$. These variables and the corresponding direction estimate $\bar{V}_G$ are not a function of the receptor data at any fixed time: they depend on the history of the receptors up to time t. Nevertheless, we may ask how the variance of the direction estimate $\bar{V}_G$ compares to the estimates $\bar{V}_C$, $\bar{V}_{RL}$ and $-\bar{V}_R$, based on the receptor data at a fixed time. To make this comparison, we compute the ratio

$$\frac{\mathbb{E}_\theta\left[\|\theta^\perp \bar{V}_G\|^2\right]}{\|\mathbb{E}_\theta\left[\bar{V}_G\right]\|^2}, \tag{B10}$$

approximating the expectation by averaging over approximately independent samples of $\bar{V}_G$. This ratio is shown in (Fig. 2E, 3D, 4A, 4E) of the main text along with the same statistic for the direction estimates $\bar{V}_C$, $\bar{V}_{RL}$ and $-\bar{V}_R$. We see that when N_R is small ($N_R = 375$) and when using the ratiometric model for G protein deactivation, the ratio (B10) is below the lower bound trace($\mathcal{I}_R^{-1}$) in (B6), below the theoretical lower bound for the estimates $\bar{V}_C$, $\bar{V}_{RL}$, and $-\bar{V}_R$ which use receptor data at a fixed time only. This is because $\bar{V}_G$ is more concentrated near the true direction θ than are the other estimates, even more concentrated than what the theoretical limit allows for any estimator based only on receptor data at a fixed time. The estimator $\bar{V}_G$ is not a function of the receptor data at any fixed time, so the theoretical bound in (B6) does not apply to the variance ratio (B10) for $\bar{V} = \bar{V}_G$. We propose that the fact that the active G protein estimator violates this bound indicates that the G protein configuration is effectively averaging receptor data over a period of time, incorporating an effectively larger amount of receptor information than what is available at a fixed time.

2. Weak signal behavior of Fisher information

Let us suppose $C(x, \theta)$ is given by (A3) with $b \ll 1$ small. Then $\mathrm{grad}_\theta \log C \approx b\theta^\perp \bar{x}$ where $\bar{x} = x/|x| = x/L$. In this case, the Fisher information $\mathcal{I}_R$ is close to a multiple of the

$2x2$ identity matrix, $\mathcal{I}_R \approx N_R \kappa^2 I$, with

$$\kappa^2 = \frac{b^2}{2|4\pi L^2|} \int_{\mathcal{S}} \|\theta^\perp \bar{x}\|^2 (1 - p_R(x,\theta)) p_R(x,\theta)\, dx. \tag{B11}$$

When b is small, then $p_R(y,\theta) \approx C_0/(C_0 + K_d)$. Hence,

$$\kappa^2 \approx b^2 \frac{C_0 K_d}{(C_0 + K_d)^2} \frac{1}{2|\mathcal{S}_1|} \int_{\mathcal{S}_1} \|\theta^\perp \cdot y\|^2\, dy \,,$$

where $\mathcal{S}_1$ is the unit sphere. By symmetry, we have $\int_{\mathcal{S}_1} \|\theta^\perp \cdot y\|^2\, dy = \int_{\mathcal{S}_1}(y_1)^2 + (y_2)^2\, dy = \frac{2}{3}|\mathcal{S}_1|$. Thus, we have

$$\mathrm{trace}(\mathcal{I}_R{}^{-1}) \approx 2(N_R \kappa^2)^{-1} \approx \frac{6}{N_R b^2} \frac{(C_0 + K_d)^2}{K_d C_0}. \tag{B12}$$

This has the same form as the analogous expression in the two-dimensional model of [32] (c.f. equation (6) for σ_ϕ^2 therein, with $p = 2b$). In particular, as observed there, this quantity is minimized when $C_0 = K_d$, suggesting that sensing is most effective when $C_0 = K_d$.

Notice that the expression in (B12) diverges as $N_R b^2 \to 0$, while the the numerator in (B6) always satisfies $\mathbb{E}_\theta\left[\|\theta^\perp \bar{V}\|^2\right] \leq 1$. Thus, it must also be true that the denominator in (B6) satisfies $\|\mathbb{E}_\theta\left[\bar{V}\right]\|^2 \leq O(N_R b^2)$ as $b \to 0$. (This can be justified directly, by approximating the distribution of V (unnormalized) by a Gaussian with mean of $b\theta$ and covariance $N_R^{-1} I_{3\times3}$.) On the other hand, as $Nb^2 \to \infty$, $\|\mathbb{E}_\theta\left[\bar{V}\right]\| \to 1$. So, (B12) and (B6) together imply that when b is small and $N_R b^2 \to \infty$,

$$\mathbb{E}_\theta\left[\|\theta^\perp \bar{V}\|^2\right] \geq \frac{6}{N_R b^2} \frac{(C_0 + K_d)^2}{K_d C_0}(1 + o(1)). \tag{B13}$$

In particular, the mean squared angular deviation of $\bar{V}$ is of the order $1/(N_R b^2)$. Nevertheless, the bound (B6) holds even when $N_R b^2$ is not large.

3. Covariance and reliability thresholds

How large must N_R be in order that V_{RL}, $-V_R$, or V_C be a reliable estimate of the direction θ? The Cramér-Rao bound shows that the mean squared angular deviation of these estimates must be at least of order $N_R^{-1} b^{-2}$. Here we consider an upper bound on angular deviation of these estimates, and we derive an explicit threshold on N_R such that when N_R is above this threshold, the median angle deviation of these estimates will be

within a certain tolerance. As we will show, this framework also extends the estimate V_G based on G protein, and it is a useful perspective for comparing V_{RL}, $-V_R$, and V_C with V_G.

Given the true direction $\theta \in S$ of the ligand gradient, and given a tolerance $\delta > 0$, we say that a vector V (either $V = V_{RL}$, $-V_R$, or V_C) is a δ-**reliable** direction estimate for θ if

$$\mathbb{P}\left(\arccos\left(\frac{V}{|V|}\cdot\theta\right)\leq\delta\right)\geq\frac{1}{2}. \tag{B14}$$

Note that $\arccos(\frac{V}{|V|}\cdot\theta)$ is the angle between the vector θ and the vector V, even if V is not a unit vector. So, condition (B14) is equivalent to saying that the median angle deviation of V from θ is less than δ. Define the cone of angle δ around θ:

$$\mathcal{C}_{\delta,\theta} = \left\{x \in \mathbb{R}^3 \mid \arccos\left(\frac{x}{\|x\|},\theta\right)\leq\delta\right\}. \tag{B15}$$

Since the cone $\mathcal{C}_{\delta,\theta}$ is invariant under scaling $x \mapsto rx$ for any $r > 0$, $V \in \mathcal{C}_{\delta,\theta}$ if and only if $\bar{V} = V/|V| \in \mathcal{C}_{\delta,\theta}$. Thus, the reliability condition (B14) is equivalent to

$$\mathbb{P}\left(V \in \mathcal{C}_{\delta,\theta}\right)\geq\frac{1}{2}. \tag{B16}$$

For a fixed deviation tolerance (e.g. $\delta = 30°$), how large must N_R be in order that the estimates V_{RL}, $-V_R$, or V_C be reliable in this sense? While the exact distributions of V_{RL}, $-V_R$, and V_C are not explicit, we can estimate $\mathbb{P}\left(V \in \mathcal{C}_{\delta,\theta}\right)$ in terms of the means and covariances of $V = V_{RL}, -V_R, V_C$; this will provide a sufficient condition for δ-reliability. We say that a random vector V is unbiased if its mean $\mu = \mathbb{E}[V]$ is aligned with θ, which means that $\mu = |\mu|\theta$.

Proposition B.1 *Suppose that a random vector V has mean $\mu = \mathbb{E}[V]$ and covariance matrix $\Sigma = Cov(V)$. Suppose that V is unbiased: $\mu = |\mu|\theta$. If*

$$trace(\Sigma) \leq |\mu|^2\sin^2(\delta)/2, \tag{B17}$$

then V is δ-reliable estimate of θ.

Proof: If $r \leq |\mu|\sin(\delta)$, then the ball of radius r centered at μ is contained in the cone $\mathcal{C}_{\delta,\theta}$. See Figure 8. In other words, the condition

$$|V - \mu| < |\mu|\sin(\delta)$$

implies $V \in \mathcal{C}_{\delta,\theta}$. By Chebyshev's inequality:

$$\mathbb{P}(|V - \mu| \geq |\mu| \sin(\delta)) \leq \frac{\mathbb{E}[|V - \mu|^2]}{|\mu|^2 \sin^2(\delta)} = \frac{\text{trace}(\Sigma)}{|\mu|^2 \sin^2(\delta)}.$$

If (B17) holds, then the right side is less than $1/2$. Hence

$$\mathbb{P}\left(V \in \mathcal{C}_{\delta,\theta}\right) \geq \mathbb{P}(|V - \mu| < |\mu| \sin(\delta)) = 1 - \mathbb{P}\left(|V - \mu| \geq |\mu| \sin(\delta)\right) \geq 1/2. \qquad \text{(B18)}$$

Thus, V is δ-reliable.

The condition (B17) is a sufficient condition for V to be δ-reliable, but it is not a necessary condition. Nevertheless, condition (B17) is easier to verify than computing the probability in (B16), and it is not based on any approximation.

We now apply this to the particular cases $V = V_{RL}, -V_R, V_C$. Let μ_{RL} denote the mean of V_{RL}:

$$\mu_{RL} = \mathbb{E}[V_{RL}] = N_R \mu_1, \qquad \mu_1 = \mathbb{E}[R_1 X_1] = \frac{1}{|S|} \int_S p_R(x) x \, dx.$$

The mean of $-V_R$ is the same, since $\mathbb{E}[X_i] = 0$:

$$\mu_R = \mathbb{E}[-V_R] = N_R \mathbb{E}[(R_1 - 1)X_1] = N_R \mathbb{E}[R_1 X_1] - N_R \mathbb{E}[X_1] = N_R \mu_1 = \mathbb{E}[V_{RL}].$$

Since the pairs $\{(X_k, R_k)\}_{k=1}^{N_R}$ are independent, the covariance matrix of V_{RL} is

$$\Sigma_{RL} = \text{Cov}(V_{RL}) = N_R \Sigma_{RL}^1,$$

where Σ_{RL}^1 is the covariance of $R_1 X_1$:

$$\Sigma_{RL}^1 = \text{Cov}(R_1 X_1) = \frac{1}{|S|} \int_S p_R(x) x x^T \, dx - \mu_1 (\mu_1)^T. \qquad \text{(B19)}$$

Similarly, the covariance matrix of $-V_R$ is

$$\Sigma_R = \text{Cov}(-V_R) = N_R \Sigma_R^1,$$

where Σ_R^1 is the covariance of $(R_1 - 1)X_1$:

$$\Sigma_R^1 = \text{Cov}((R_1 - 1)X_1) = \frac{1}{|S|} \int_S (1 - p_R(x)) x x^T \, dx - \mu_1 (\mu_1)^T. \qquad \text{(B20)}$$

The mean of μ_C of V_C is exactly twice the mean of V_{RL}:

$$\mu_C = \mathbb{E}[V_C] = N_R \mathbb{E}[(2R_1 - 1)X_1] = \frac{N_R}{|S|} \int_S (2p_R(x) - 1)x \, dx = N_R 2\mu_1 = 2\mu_{RL}.$$

The covariance of V_C is

$$\Sigma_C = N_R \Sigma_C^1$$

where Σ_C^1 is the covariance of $(2R - 1)X_1$:

$$\Sigma_C^1 = \text{Cov}((2R_1 - 1)X_1) = \frac{1}{|S|} \int_S x x^T \, dx - 4\mu_1(\mu_1)^T. \tag{B21}$$

Notice that the relation

$$\Sigma_C^1 = \Sigma_{RL}^1 + \Sigma_R^1 - 2\mu_1 \mu_1^T$$

always holds. The estimates V_{RL}, $-V_R$, and V_C are all unbiased if μ_1 is aligned with θ:

$\mu_1 = |\mu_1|\theta$.

By applying Proposition B.1 with $V = V_{RL}, -V_R, V_C$ we obtain the following thresholds for δ-reliability:

Corollary B.1 *Suppose that the estimates V_{RL}, $-V_R$, and V_C are unbiased: $\mu_1 = |\mu_1|\theta$. The reliability condition* (B17) *for V_{RL} is*

$$trace(\Sigma_{RL}) \leq |\mu_{RL}|^2 \sin^2(\delta)/2$$

or, equivalently,

$$N_R \geq \frac{2 \cdot trace(\Sigma_{RL}^1)}{(\mu_1 \cdot \theta)^2 \sin^2(\delta)}. \tag{B22}$$

The reliability condition (B17) *for $-V_R$ is*

$$trace(\Sigma_R) \leq |\mu_R|^2 \sin^2(\delta)/2$$

or, equivalently,

$$N_R \geq \frac{2 \cdot trace(\Sigma_R^1)}{(\mu_1 \cdot \theta)^2 \sin^2(\delta)}. \tag{B23}$$

The reliability condition (B17) *for V_C is*

$$trace(\Sigma_C) \leq |\mu_C|^2 \sin^2(\delta)/2$$

or, equivalently,

$$N_R \geq \frac{2 \cdot trace(\Sigma_C^1)}{(2\mu_1 \cdot \theta)^2 \sin^2(\delta)}. \tag{B24}$$

We may interpret the quantity $\mu_1 \cdot \theta$ to be the "signal strength"; this calculation shows that reliability of V_{RL}, $-V_R$ and V_C requires that N_R is of the order $(\mu_1 \cdot \theta)^{-2}$, which is large when the signal is weak (shallow ligand gradient). Suppose that the gradient is weak: $C(x) = C_0(1 + bx \cdot \theta)$ with $b > 0$ small (here we assume $L = 1$). Then $p_R(x)$ is close to linear in x:

$$p_R(x) = \frac{C(x)}{C(x) + K_D} = p_R(0) + bp_R(0)(1 - p_R(0))x \cdot \theta + O(b^2)$$

and thus

$$(\mu_1 \cdot \theta) \approx b\, p_R(0)(1 - p_R(0))c_1, \qquad c_1 = \frac{1}{|\mathcal{S}_1|}\int_{\mathcal{S}_1}(x_1)^2\, dx = \frac{1}{3}, \qquad \Sigma_{RL}^1 \approx p_R(0)c_1 I_3, \qquad \Sigma_R^1 \approx (1 - p_R(0))c_1 I$$

where I_3 is the 3×3 identity matrix. Hence, in this weak gradient regime, the sufficient condition (B22) for V_{RL} to be δ-reliable reduces to

$$N_R \geq \frac{2 \cdot \text{trace}(\Sigma_{RL}^1)}{(\mu_1 \cdot \theta)^2 \sin^2(\delta)} \approx \frac{p_R(0)}{b^2(p_R(0))^2(1 - p_R(0))^2} \cdot \frac{18}{\sin^2(\delta)}. \qquad (B25)$$

The condition (B23) for $-V_R$ to be δ-reliable reduces to

$$N_R \geq \frac{2 \cdot \text{trace}(\Sigma_R^1)}{(\mu_1 \cdot \theta)^2 \sin^2(\delta)} \approx \frac{(1 - p_R(0))}{b^2(p_R(0)^2(1 - p_R(0))^2} \cdot \frac{18}{\sin^2(\delta)}. \qquad (B26)$$

The condition (B24) for V_C to be δ-reliable reduces to

$$N_R \geq \frac{2 \cdot \text{trace}(\Sigma_C^1)}{(2\mu_1 \cdot \theta)^2 \sin^2(\delta)} \approx \frac{1}{b^2(p_R(0))^2(1 - p_R(0))^2} \cdot \frac{9}{2\sin^2(\delta)}. \qquad (B27)$$

Comparing (B25), (B26), and (B27), we see that when $p_R(0) \in (1/4, 3/4)$, V_C has the lowest threshold for reliability. When $p_R(0) = 1/2$ (i.e. $C_0 = K_d$), the thresholds (B25) and (B26) for V_{RL} and $-V_R$ are the same, and the threshold (B27) for V_C is exactly $1/2$ of (B25) or (B26). In this scenario, V_C is effectively using twice the data as V_{RL} or $-V_R$. For $p_R(0)$ outside of the range $(1/4, 3/4)$, V_C is not the most reliable of the three estimates. When $p_R(0) < 1/4$ (i.e. $C_0 < (1/3)K_d$), the reliability threshold for V_{RL} ((B25)) is lowest among the three; when $p_R(0) > 3/4$ (i.e. $C_0 > 3K_d$), the reliability threshold for $-V_R$ ((B25)) is lowest among the three. The intuition about why V_C would perform worse than V_{RL} when $p_R(0) < 1/4$ is that in this regime the majority of receptors are inactive and their conditional angular distribution is more uniform than the conditional distribution of active receptor. Indeed, the conditional density of a receptor's location, conditioned on it being inactive, is proportional to $1 - p_R(x) = K_d/(K_d + C(x))$, and this density is more uniform when C_0/K_d is small. So, when C_0/K_d is small, the estimate V_C is dominated by

 the contribution from inactive receptors which are approximately uniformly distributed on
648 the sphere.

649 Another approach to estimating the probability in (B16) would be to approximate V by
650 a Gaussian random vector having the same mean and covariance (justified by the central
651 limit theorem when N_R is large). If V has mean $|\mu|\theta$ and covariance matrix $\frac{\sigma^2}{N_R} I_{3\times3}$, then
652 under this approximation

$$\mathbb{P}\left(V \in \mathcal{C}_{\delta,\theta}\right) \approx \mathbb{P}\left(|\mu|\theta + \frac{\sigma}{\sqrt{N_R}}Z \in \mathcal{C}_{\delta,\theta}\right) = \mathbb{P}\left(\frac{\sqrt{N_R}|\mu|}{\sigma}\theta + Z \in \mathcal{C}_{\delta,\theta}\right)$$

653 where Z is a standard Gaussian vector in $\mathbb{R}^3$ with mean zero and covariance $I_{3\times3}$. Thus, V
654 is δ-reliable once $N_R > \frac{\sigma^2}{|\mu|^2}c_\delta^2$, where the constant $c_\delta > 0$ is the smallest value r such that
655 $\mathbb{P}\left(r\theta + Z \in \mathcal{C}_{\delta,\theta}\right) \geq 1/2$. Applied to V_{RL}, $-V_R$, and V_C, this approach yields thresholds with
656 the same scaling as (B25), (B26), (B27), but with the less explicit universal constant c_δ^2.

Appendix C: Time averaging

In the ratiometric model, the instantaneous G protein states (active/inactive) are derived
from historical receptor states. Each G protein's state is determined by its most recent
interaction with a receptor, either active or inactive (Figure 9). If $N_G > N_R$, this mechanism
may allow the G protein states to represent more information than what is encoded by the
receptor states at a single time. Here we explain this mechanism. Throughout this section
we consider only the ratiometric model for G protein dynamics.

1. Sampling the state of a single receptor at multiple random times

For simplicity, we first consider the case of a single receptor, and n G proteins interacting
with it, and we ignore the spatial aspect of the model for now. The state of a given G
protein is determined by the state of the receptor at the time of the protein's most recent
encounter with that receptor. If the times at which the n proteins most recently interacted
with that receptor are $t_1, t_2, \ldots, t_n$ then the states of the G proteins represent n samples of
the historical state of the (single) receptor. These historical states are correlated, however;
the correlation is higher when the interaction times are close together. If the interaction
times are far apart, then sampling the state of the receptor at n different times can be similar
to sampling n independent receptors.

The receptor state is a two-state, continuous-time Markov process $R(t)$; the two states

are the active state (1) and the inactive state (0). The transition rates for this chain are

given by (A1), where $x \in \mathcal{S}$ is the fixed location of the receptor. For clarity in notation we

write $\lambda_+ = k_{\mathrm{on}} C(x)$ for the activation rate, $\lambda_- = k_{\mathrm{off}}$ for the inactivation rate, and we define

$\lambda = \lambda_+ + \lambda_-$. The stationary distribution for this process is $\nu = (\nu(0), \nu(1))$, with

$$\nu(0) = \frac{\lambda_-}{\lambda_+ + \lambda_-}, \qquad \nu(1) = \frac{\lambda_+}{\lambda_+ + \lambda_-}.$$

So, $\nu(0)$ and $\nu(1)$ are the steady-state probabilities of finding the receptor in states 0 or

1, respectively. Let $p_t(z, y)$ denote the transition probabilities: $p_t(z, y)$ is the probability

that the state at time $t > 0$ is y, given that the state at time 0 is z, $y, z \in \{0, 1\}$. These

probabilities satisfy the system

$$\frac{d}{dt} p_t(z, 0) = -\lambda_+ p_t(z, 0) + \lambda_- p_t(z, 1)$$
$$\frac{d}{dt} p_t(z, 1) = \lambda_+ p_t(z, 0) - \lambda_- p_t(z, 1)$$

for $x \in \{0, 1\}$. This has an explicit solution. We find that the transition probabilities are

$$p_t(0, 0) = \nu(0) \left(1 - e^{-\lambda t}\right) + e^{-\lambda t},$$
$$p_t(0, 1) = \nu(1) \left(1 - e^{-\lambda t}\right),$$
$$p_t(1, 0) = \nu(0) \left(1 - e^{-\lambda t}\right),$$
$$p_t(1, 1) = \nu(1) \left(1 - e^{-\lambda t}\right) + e^{-\lambda t}.$$

In particular, regardless of the initial state x, $p_t(z, y)$ converges exponentially fast to $\nu(y)$,

at rate $\lambda = \lambda_+ + \lambda_-$. Thus, $\lambda^{-1} = (k_{\mathrm{on}} C + k_{\mathrm{off}})^{-1}$ is the time scale for "loss of memory" or

mixing of the receptor state.

Now, suppose the receptor process R has achieved stationarity, and suppose there are n

G proteins that read the state of this single receptor at random times $t_1, \ldots, t_n$ (unordered).

Assume these times are independent of each other and of the receptor process. Suppose

that for each k, t_k is exponentially distributed with mean $\tau = \mathbb{E}[t_k]$. We will use $G_i \in \{0, 1\}$

to denote the state of the ith G protein (i.e. $G_i = R(t_i)$). Although the times $\{t_k\}_{k=1}^n$ are

independent, the random variables $(G_1, \ldots, G_n) = (R(t_1), \ldots, R(t_n))$ are dependent, since

the data come from the same receptor. However, the level of dependence is controlled by

the dimensionless parameter $\lambda \tau$, as we now show. Let A_G denote the fraction of receptor

readings that are in the active state:

$$A_G = \frac{1}{n} \sum_{i=1}^{n} \mathbb{I}_{G_i=1} = \frac{1}{n} \sum_{k=1}^{n} R(t_k).$$

Let $p_R = \nu(1) = \frac{\lambda_+}{\lambda_+ + \lambda_-} = \mathbb{P}(R(t) = 1)$ be the probability that the receptor is active (R is statistically stationary, so this probability does not change with time). The autocorrelation function for the receptor state is: $\mathbb{E}[R(t)R(t+s)] = p_R^2 + p_R(1 - p_R)e^{-\lambda s}$.

Proposition C.1 *The mean and variance of A_G are $\mathbb{E}[A_G] = p_R$ and*

$$Var(A_G) = \frac{1}{n}p_R(1 - p_R) + \left(1 - \frac{1}{n}\right) \frac{p_R(1 - p_R)}{(1 + \lambda\tau)}. \tag{C1}$$

The first term in this expression, $p_R(1 - p_R)/n$, would be the variance if the n readings were independent (e.g. taken separately from n independent receptors); in the limit $n \to \infty$ this term vanishes. The second term in the variance is the result of temporal correlation in the receptor process. As $\lambda\tau \to \infty$, the last term vanishes, leaving only $p_R(1 - p_R)/n$.

2. Effective receptor number

Let us compare this result to an alternative model, which will motivate the notion of **effective receptor number**. Instead of there being a single receptor with state changing in time, let us now suppose there is a cluster of n_R receptors with independent states that do not change in time. Let p_R be the probability that a given receptor in the cluster is active. Suppose that each of the n G proteins samples a state from this cluster: each protein chooses one of the n_R receptors at random and records its state. So, two proteins that choose the same receptor must record the same state. As before, let A_G be the fraction of readings that are in the active state:

$$A_G = \frac{1}{n} \sum_{i=1}^{n} \mathbb{I}_{G_i=1}.$$

Proposition C.2 *The mean and variance of A_G are $\mathbb{E}[A_G] = p_R$ and*

$$Var(A_G) = \frac{1}{n}p_R(1 - p_R) + \left(1 - \frac{1}{n}\right) \frac{p_R(1 - p_R)}{n_R}. \tag{C2}$$

Both models have the same mean p_R for all parameter choices. Comparing (C1) and (C2), we see that the variance in the two models is the same when $(1 + \lambda\tau) = n_R$. Thus, we

709 could think of $(1 + \lambda\tau)$ as an **effective number of receptors**: when the G proteins read a
710 single receptor at independent random times, it is as if each G protein reads a state chosen
711 randomly and independently from a group of $(1 + \lambda\tau)$ independent receptors. The fact that
712 the G proteins read historical states effectively expands the pool of independent receptors.
713 Interestingly, the quantity $n_R = (1 + \lambda\tau)$ does not depend on n, assuming $n > 1$.

3.   Derivation of Proposition C.1 and Proposition C.2

For both models, the variance of A_G is

$$
\mathrm{Var}(A_G) = \sum_{i=1}^{n} \mathrm{Var}(\mathbb{I}_{G_i=1}) + \sum_{i \neq k} \mathrm{Cov}(\mathbb{I}_{G_i=1}, \mathbb{I}_{G_k=1})
$$

$$
= n p_R (1 - p_R) + (n^2 - n) \mathrm{Cov}(\mathbb{I}_{G_1=1}, \mathbb{I}_{G_2=1}), \tag{C3}
$$

and the covariance term is

$$
\mathrm{Cov}(\mathbb{I}_{G_1=1}, \mathbb{I}_{G_2=1}) = \mathbb{P}(G_1 = 1, G_2 = 1) - \mathbb{P}(G_1 = 1)\mathbb{P}(G_2 = 1) \tag{C4}
$$

$$
= \mathbb{P}(G_1 = 1, G_2 = 1) - p_R^2. \tag{C5}
$$

For the model considred in Proposition C.1, the joint density of the independent times
(t_1, t_2) is $\rho(t_1, t_2) = \alpha^2 e^{-\alpha(t_1+t_2)}$ for $t_1, t_2 \geq 0$. Therefore,

$$
\mathbb{P}(G_1 = 1, G_2 = 1) = \int_0^\infty \int_0^\infty \mathbb{P}(R(t_1) = 1, \quad R(t_2) = 1)\rho(t_1, t_2)\, dt_1\, dt_2
$$

$$
= 2 \iint_{t_1 \leq t_2} \mathbb{P}(R(t_2) = 1 \mid R(t_1) = 1)\mathbb{P}R(t_1) = 1)\alpha^2 e^{-\alpha(t_1+t_2)}\, dt_1\, dt_2
$$

$$
= 2 p_R \iint_{t_1 \leq t_2} \left(e^{-\lambda(t_2-t_1)} + p_R(1 - e^{-\lambda(t_2-t_1)}) \right) \alpha^2 e^{-\alpha(t_1+t_2)}\, dt_1\, dt_2
$$

$$
= p_R^2 + 2 p_R(1 - p_R) \iint_{t_1 \leq t_2} e^{-\lambda(t_2-t_1)} \alpha^2 e^{-\alpha(t_1+t_2)}\, dt_1\, dt_2
$$

$$
= p_R^2 + p_R(1 - p_R) \left(\frac{\alpha}{\lambda + \alpha} \right). \tag{C6}
$$

715 Therefore,

$$
\mathrm{Cov}(\mathbb{I}_{G_1=1}, \mathbb{I}_{G_2=1}) = p_R(1 - p_R) \left(\frac{1}{(\lambda/\alpha) + 1} \right),
$$

and

$$
\mathrm{Var}(A_G) = n p_R(1 - p_R) + (n^2 - n) p_R(1 - p_R) \left(\frac{1}{(\lambda/\alpha) + 1} \right). \tag{C7}
$$

For the model considered in Proposition C.2,

$$\mathbb{P}(G_1 = 1, G_2 = 1) = p_R \mathbb{P}(\text{two proteins choose the same receptor})$$

$$+ p_R^2 \mathbb{P}(\text{two proteins choose different receptors})$$

$$= p_R \frac{1}{n_R} + p_R^2 \left(1 - \frac{1}{n_R} \right). \tag{C8}$$

Therefore,

$$\text{Cov}(\mathbb{I}_{G_1=1}, \mathbb{I}_{G_2=1}) = \frac{p_R(1 - p_R)}{n_R}.$$

Appendix D: Spatially extended, ratiometric model

In the analysis above we considered a population of G proteins interacting with a single receptor. Here we show how these ideas extend to the case when there are multiple receptors distributed spatially across the cell surface, and the G proteins interact with them, according to the ratiometric model, as the proteins diffuse over the surface. Throughout this section, we assume the ratiometric model, so $V_G = V_G^r$.

1. G protein ages

Considering the times at which a G protein interacts with a receptor, we define the **age** of a G protein at time t to be $s \geq 0$ if the last interaction between that G protein and some receptor occured at time $t - s$. If the G protein has age s, then that G protein has been in the same state for at least time $t - s$. Here we say "at least" time $t - s$ because it could be that the G protein was already in that state just before this most recent encounter with a receptor. For example, when an already-active G protein encounters an active receptor, the G protein's age will reset to zero but its state does not change. (See Figure 9). We will say that a G protein at time t is **linked** to a particular receptor if that G protein's last interaction was with that particular receptor – if that G protein's state at time t was recorded from that particular receptor (at time $t - s$, where s is the protein's age). Each G protein is linked to only one receptor at a time. Two or more G proteins may be linked to the same receptor, which creates dependence between those G protein states. (See Figure 10). In fact, this linking is the only source of statistical dependence between G protein states: G protein states are dependent only through interaction with the same receptor. Conditioned

on being linked to two different receptors, two G protein states are statistically independent. However, for G proteins linked to the same receptor, their ages may be different, and the observations above suggest that if these ages are sufficiently far apart, then the recorded states are approximately independent. As explained above in Proposition C.1, the level of dependence/independence between proteins linked to the same receptor is controlled by the ratio of two times scales – the mean G protein age and the receptor turnover time.

We can estimate the typical age of a G protein. If there are a total of N_R receptors, then the typical distance between receptors is of the order $\delta = \sqrt{(4\pi L^2/N_R)}$ where L is the cell radius. This is because $4\pi L^2/N_R$ is the surface area per receptor, and the surface is two-dimensional (hence, the square root). Therefore, a G protein diffusing on the cell surface is navigating among receptors that are typically distance δ apart. If we want to know when a given G protein last interacted with some receptor (any receptor), it is the same as looking *backwards* in time and asking when does a Brownian motion first hit an r_*-neighborhood of some receptor, among the N_R receptors distributed across the surface approximately δ apart. If the receptors have an interaction radius of $r_* \ll \delta$ and the G protein diffusion coefficient is D_G, then the typical **age** of a G protein will be of the order

$$\tau = \frac{\delta^2}{D_G} \ln(\delta/r_*) = \frac{L^2}{N_R D_G} \ln(\delta/r_*). \tag{D1}$$

We may also describe this quantity as the time scale of diffusive encounters between a G protein and receptors. This is motivated by thinking about a periodic array of discs in $\mathbb{R}^2$, having radius r_* and with centers δ apart. The quantity $\frac{\delta^2}{D_G} \ln(\delta/r_*)$ is the typical time it takes for a Brownian motion (with diffusion coefficient D_G) to hit one of the discs (any disc), when started from a random location. (The logarithm is an artifact of the 2-dimensional nature of the cell surface and of the Green's function in 2-dimensions.) Here and elsewhere we drop constants of proportionality, such as 4π, that do not depend on the parameters, since we are mainly interested in the scaling with parameters. This time τ is not the typical time required for a given G protein to find a *particular* receptor. Rather, it is the time required (looking backward in time) to encounter any one of the receptors – this later quantity is what determines the age of a G protein. The relation in (D1) can be derived as in Appendix B of [28]; ignoring constants of proportionality that do not depend on the parameters, (D1) above has the same form as Eq. 14 from [28]. See also [52], [53], and references therein for a broader discussion of first-passage time calculations. This suggests that the age of a G

768 protein is a random variable that is approximately exponentially distributed with mean τ.

769 If the number of G proteins is $N_G > N_R$, then on average there are (N_G/N_R) G proteins

770 per receptor. So, if the receptors and proteins are uniformly distributed on the sphere, we

771 should expect that for each of the N_R receptors there are about (N_G/N_R) G proteins that

772 are **linked** with that receptor, in the sense defined above.

2. Covariance for G protein, fixed receptor locations

As we did with the simple single-receptor model above, we now consider how the time τ

effects the variation of the G protein signal. Specifically, we show how for the ratiometric

model the covariance of V_G depends on the time scale τ and the time scale of receptor

dynamics. Here we suppose that the receptor locations are fixed at $X_1, \ldots, X_{N_R}$. The

receptor states are changing randomly, but their locations are fixed. A receptor at X_i

activates at rate $k_{\mathrm{on}}C(X_i)$ and deactivates at rate k_{off}; thus, the time scale for receptor

dynamics at location X_i is $\lambda^{-1}(X_i) = (k_{\mathrm{on}}C(X_i) + k_{\mathrm{off}})^{-1}$. Suppose there are N_G proteins

that are placed at locations $\{Y_j\}_{j=1}^{N_G}$, which are independent and uniformly distributed on

the sphere. We consider their states $\{G_j\}_{j=1}^{N_G}$ at a fixed time, where $G_j \in \{0,1\}$. We are

interested in the vector

$$V_G = \sum_{j=1}^{N_G} G_j Y_j$$

784 which is the sum of the positions of active G proteins. Let $\mu_G = \mathbb{E}[V_G]$ be the mean of V_G,

785 and let $\Sigma_G = \mathrm{Cov}(V_G)$ be the covariance matrix. As in (B16), we say that the G protein

786 signal V_G is a δ-**reliable** estimate if

$$\mathbb{P}\left(V_G \in \mathcal{C}_{\delta,\theta}\right) \geq \frac{1}{2}. \tag{D2}$$

Proposition B.1 applied with $V = V_G$ gives us a sufficient condition for δ-reliability of V_G:

Corollary D.1 *Suppose that the estimate V_G is unbiased. If $trace(\Sigma_G) \leq |\mu_G|^2 \sin^2(\delta)/2$,*

then V_G is δ-reliable.

To make use of this criteria and to compare the reliability of V_G with the reliability of

V_{RL}, we need to compute μ_G and Σ_G. We now explain how these can be calculated for the

ratiometric model.

The state G_j of the protein at position Y_j was read from a particular receptor at some previous time – we say that this protein is **linked** to that particular receptor. Define random variables $\{L_j\}_{j=1}^{N_G}$ which indicate the receptor to which the jth protein is linked: $L_j = k$ means that the jth protein (at Y_j) is linked to the kth receptor (which is at position X_k). Possibly multiple G proteins can be linked to the same receptor, but they may or may not read the same state from that receptor (because they interacted with that receptor at different times, and the receptor state is changing over time).

Let $\eta_i(y)$ denote the conditional density of Y_j, given $L_j = i$. This η_i is depends on the geometric arrangement of all receptors $\{X_k\}_{k=1}^{N_R}$, and we denote this dependence by $\eta_i([X], y)$. Although this quantity is impossible to compute exactly/explicitly, we should think of this as being concentrated in a neighborhood of X_i, as G proteins are most likely to be linked to nearby receptors. Thus,

$$\mathbb{P}(Y_j \in A \mid L_j = i) = \int_A \eta_i([X], y) \, dy, \qquad A \subset S.$$

Given $L_j = i$, the jth G protein takes its state from the ith receptor at some time T_j. Motivated by the analysis above, we make the simplifying assumption that this time (the age of that protein) is exponentially distributed with mean τ, $T_j \sim \text{Exponential}(1/\tau)$, where the parameter τ is given by (D1). We also assume that the ages are independent of G protein positions.

Proposition D.1 *Assume the ratiometric model for G protein dynamics. Given fixed receptor locations $\{X_i\}_{i=1}^{N_R}$, the mean of V_G is*

$$\mu_G = \mathbb{E}[V_G] = N_G \sum_{i=1}^{N_R} p_R(X_i) \int_S \eta_i([X], y) y \, dy \, \mathbb{P}(L_1 = i).$$

The covariance matrix $\Sigma_G = \mathbb{E}[(V_G - \langle V_G \rangle)(V_G - \langle V_G \rangle)^T]$ of V_G is

$$\Sigma_G = N_G M_1 + \frac{(N_G^2 - N_G)}{N_R} M_2 \tag{D3}$$

where

$$M_1 = \sum_{i=1}^{N_R} p_R(X_i) \int_S \eta_i([X], y) yy^T \, dy \, \mathbb{P}(L_1 = i)$$

$$- \sum_{m=1}^{N_R} \sum_{i=1}^{N_R} p_R(X_m) p_R(X_i) \int_S \int_S \eta_m([X], z) \eta_i([X], y) yz^T \, dy \, dz \, \mathbb{P}(L_1 = i) \mathbb{P}(L_2 = m)$$

(D4)

(D5)

and

$$M_2 = N_R \sum_{i=1}^{N_R} \frac{p_R(X_i)(1 - p_R(X_i))}{1 + \tau \lambda(X_i)} \int_S \int_S \eta_i([X], z) \eta_i([X], y) yz^T \, dy \, dz \, \mathbb{P}(L_1 = L_2 = i). \quad \text{(D6)}$$

The quantity

$$\int_S \eta_i([X], y) y \, dy$$

in the expression for μ_G is the mean location of a G protein linked to the i^{th} receptor (at location X_i). Observe that the covariance matrix Σ_G is the sum of two terms. The second term, M_2, results from correlations caused by G proteins possibly interacting with the same receptor. This M_2 is the term where time-scales play a role, through $\tau \lambda(X_i)$. When $\tau \lambda$ is large, meaning that receptor dynamics are fast relative to G protein ages, then M_2 may be negligible.

From Proposition D.1 we can identify two parameter regimes corresponding to different behavior of the random fluctuation of V_G. The covariance of $\frac{1}{N_G} V_G$ is

$$\text{Cov}(V_G/N_G) = (N_G)^{-2} \Sigma_G = \frac{1}{N_G} M_1 + \frac{(1 - 1/N_G)}{N_R} M_2.$$

Recall that the matrix M_2 depends on τ and λ. The quantity $\lambda(x) = k_{\text{on}} C(x) + k_{\text{off}}$ depends on x. Nevertheless, when the gradient is shallow, we have $\lambda \approx \lambda_0 = k_{\text{on}} C_0 + k_{\text{off}}$. Thus, Proposition D.1 shows that the dimensionless parameter $\lambda_0 \tau = (k_{\text{on}} C_0 + k_{\text{off}}) \tau$, which is ratio of the G protein age τ with the receptor turnover time λ_0^{-1}, plays an important role. (The product $\lambda_0 \tau$ is analogous to a Damköhler number.) This analysis suggests two qualitatively different regimes for the behavior of the covariance of V_G:

- **The G-limited regime.** If

$$N_G \ll (1 + \lambda_0 \tau) N_R = N_R + \frac{\lambda_0 L^2}{D_G} \ln(\delta/r_*)$$

then we have

$$\text{Cov}(\frac{1}{N_G}V_G) \approx \frac{1}{N_G}M_1.$$

In this regime, the covariance is approximately that of N_G proteins that are sampling independent receptors. Either N_R is large or $\lambda_0\tau$ is large, so that the receptor states sampled by the G proteins are approximately independent. Consequently, increasing N_R or $\lambda_0\tau$ further does not decrease the covariance as significantly as would an increase in N_G. In other words, small N_G is the main limitation for reliability of V_G in this regime. Note: it is possible that $N_R < N_G < (1 + \lambda_0\tau)N_R$. In other words, even if $N_R < N_G$, the time scales may be such that the covariance of V_G is relatively insensitive to changes in N_R because $(1 + \lambda_0\tau)N_R \gg N_G$. This is consistent with our observation that the estimate V_G^r may be robust even at low N_R, as illustrated in Figure 2 and 3 of the main text. In particular, in this regime the noise-to-signal ratio for V_G^r may be smaller than $\text{trace}(\mathcal{I}_R^{-1})$ from (B6), which is the theoretical lower bound that applies to V_C and to any unbiased estimate based on instantaneous receptor states at a single fixed time.

- **The Receptor-limited regime.** If

$$N_G \gg (1 + \lambda_0\tau)N_R = N_R + \frac{\lambda_0 L^2}{D_G}\ln(\delta/\epsilon),$$

then we have

$$\text{Cov}(\frac{1}{N_G}V_G) \approx \frac{1}{N_R}M_2.$$

In this regime, there are typically many G proteins linked to a given receptor. Variation of the estimate V_G is dominated by fluctuation in the receptor states (which are passed on to the G protein), and there is a relatively high level of correlation in the receptor states recorded by the G proteins linked to the same receptor. Increasing N_G further does not decrease the covariance as significantly as would an increase in either N_R or $\lambda_0\tau$. In other words, even though N_G is large, N_R is the main limitation for reliability of V_G in this regime.

In view of these calculations, we define the quantity

$$\tilde{N}_R = (1 + \lambda_0\tau)N_R = N_R + \frac{\lambda_0 L^2}{D_G}\ln(\delta/\epsilon).$$

We call this the **effective receptor number** – this number represents an effective number of receptor states made available through historical sampling. The quantity $\tilde{N}_R$ should not be thought of as something that applies to a particular cell. Instead, $\tilde{N}_R$ can be regarded as a property of the ratiometric model, a model whose statistical properties are completely determined by parameters N_R, N_G, D_G, L, r_*, k_{off}, K_d and the ligand concentration $C(x)$.

Considering how these two regimes depend on the parameters, we make the following observations:

- Regarding D_G: Faster diffusion of G protein (D_G large) reduces τ. This is because when the diffusion of G protein is fast, the proteins typically interact with receptors very frequently, and their age is lower, so their states are more correlated with other G proteins that also hit the same receptor in the very recent past (the states of linked G proteins are likely to coincide). In the hypothetical extreme $D_G \to \infty$, the age of a G protein goes to zero so that the G proteins are recording only information from the present state of the receptors. In this case, $\tilde{N}_R \approx N_R$. Conversely, slower diffusion (D_G small) increases τ and the effective receptor number $\tilde{N}_R$. This is because with slow diffusion of G protein, their ages tend to be larger and further apart, so even the states of linked G proteins (i.e. G proteins that are linked to the same receptor) are approximately independent.

- Regarding λ_0: Faster receptor dynamics $\lambda_0 \gg 1$ increases $\tilde{N}_R$. This is because the states of a given receptor $R(t_1), \ldots, R(t_n)$ at some times $t_1 < t_2 < \cdots < t_n$ (that do not depend on λ_0) will be approximately independent when λ_0 is large. Thus, if multiple G proteins obtained their state by last interacting with that same receptor, then those proteins are encoding approximately independent state values. Conversely, slow receptor dynamics ($\lambda_0 \ll 1$) decreases $\tilde{N}_R$. A key point here is that changing the receptor turnover time λ_0^{-1} has no effect on the distribution of G protein ages.

- Regarding r_*: decreasing the interaction radius r_* will tend to increase G protein ages relative to the receptor turnover time, since it is more difficult for a protein to reach a receptor. This leads to greater independence in the states encoded by the G protein ensemble.

Thus far we have reasoned under the assumption that ligand concentration $C(x)$ is not

far from uniform:

$$\lambda(x) = k_{\text{on}}C(x) + k_{\text{off}} \approx \lambda_0 = k_{\text{on}}C_0 + k_{\text{off}} = k_{\text{off}}\left(C_0/K_d + 1\right).$$

With non-uniform ligand concentration, the receptor turnover time depends on position. So, in principle, the G protein on front end of the cell (where ligand concentration $C(\theta)$ is highest) should encode a higher level of independence than on the back end, because λ is larger on the front end. However, in the scenarios we are considering the ratio $(C(\theta) + K_d)/(C(-\theta) + K_d))$, with θ denoting the gradient direction, is not very large. So, the difference in this effect between front and back may not be significant.

3. Comparing V_G^r with V_{RL} when N_R is sufficiently large

These expressions for the mean μ_G and covariance Σ_G of V_G^r are complicated. In particular, explicitly computing the probability of linking to a particular receptor is not feasible, as it depends on the geometric configuration of all receptors. Nevertheless, we can gain further insight by making a simplifying assumption that N_R is large and the receptors are evenly distributed over S. This will allow us to simplify μ_G and Σ_G and compare them with μ_R and Σ_R, the mean and covariance of V_{RL}. In particular, we will show that in certain parameter regimes, V_G^r may be a more reliable estimate of θ than V_{RL}. This analysis is consistent with the results of our numerical simulations.

Under the assumption that N_R is large and the receptor locations are well-spaced over S, we expect that a G protein placed uniformly at random on S is equally likely to link with any receptor:

$$\mathbb{P}(L_1 = i) \approx 1/N_R, \quad \forall i = 1, \ldots, N_R. \tag{D7}$$

Furthermore, under this assumption we expect that the conditional density $y \mapsto \eta_i([X], y)$ is concentrated in a small neighborhood of X_i. Thus, we approximate

$$\int_S \eta_i([X], y)y \, dy \approx X_i, \qquad \int_S \eta_i([X], y)yy^T \, dy \approx X_i X_i^T, \qquad \forall i = 1, \ldots, N_R. \tag{D8}$$

Using the approximations (D7) and (D8) we have the following approximations of the mean and covariance of V_G^r. Recall from (B20), that V_{RL} satisfies

$$\mathbb{E}[V_{RL}] = N_R\mu_1, \qquad \text{Cov}(V_{RL}) = N_R\Sigma_{RL}^1,$$

where

$$\mu_1 = \int_S p_R(x)x \, dx, \qquad \Sigma^1_{RL} = \frac{1}{|S|} \int_S p_R(x)xx^T \, dx - \mu_1(\mu_1)^T.$$

Proposition D.2 *Using the approximations* (D7) *and* (D8) *and assuming* N_R *is large, the mean* μ_G *and covariance* Σ_G *of* V^r_G *are*

$$\mu_G = \mathbb{E}[V^r_G] \approx N_G \mu_1$$

and

$$\Sigma_G = Cov(V^r_G) \approx N_G \Sigma^1_{RL} + \frac{(N^2_G - N_G)}{N_R} \Sigma_\tau,$$

where

$$\Sigma_\tau = \frac{1}{|S|} \int_S \frac{p_R(x)(1 - p_R(x))}{1 + \tau\lambda(x)} xx^T \, dx.$$

Using this, we can derive the following remarkable consequence about reliability of V^r_G in the G-limited regime:

Corollary D.2 *When* $N_G > N_R$ *and* $\tau\lambda$ *is large, the estimate* V^r_G *of* θ *may be* δ-*reliable, even while* V_{RL} *is not* δ-*reliable.*

Proof: Using these approximations for μ_G and Σ_G, the condition given in Corollary D.1 for δ-reliability of V^r_G is (approximately):

$$\frac{1}{N_G}\text{trace}(\Sigma^1_{RL}) + \frac{(1 - 1/N_G)}{N_R}\text{trace}(\Sigma_\tau) \leq |\mu_1| \sin^2(\delta)/2.$$

In particular, this holds if

$$\frac{1}{N_G}\text{trace}(\Sigma^1_{RL}) + \frac{1}{N_R}\text{trace}(\Sigma_\tau) \leq |\mu_1| \sin^2(\delta)/2. \tag{D9}$$

On the other hand, the δ-reliability condition for V_{RL}, from Corollary B.1 is

$$\frac{1}{N_R}\text{trace}(\Sigma^1_{RL}) \leq |\mu_1| \sin^2(\delta)/2. \tag{D10}$$

The matrix Σ_τ depends on $\tau\lambda$, but Σ^1_{RL} does not. For any N_R, we may take δ small enough so that V_{RL} is **not** δ-reliable. In particular, (D10) does not hold if δ is small. For this same value of N_R and δ, however, we may take $N_G > N_R$, and then $\lambda\tau$ large enough so that the inequality (D9) holds, implying that V^r_G is δ-reliable, even though V_{RL} is not. In particular, it would be necessary for $\text{trace}(\Sigma_\tau) \leq (1 - N_R/N_G)\text{trace}(\Sigma^1_{RL})$.

 4. Derivation of Proposition D.1

The mean of V_G is:

$$\mathbb{E}[V_G] = \sum_{j=1}^{N_G} \mathbb{E}[G_j Y_j] = N_G \mathbb{E}[G_1 Y_1] = N_G \sum_{i=1}^{N_R} \mathbb{E}[G_1 Y_1 \mid L_1 = i]\mathbb{P}(L_1 = i)$$

$$= N_G \sum_{i=1}^{N_R} \mathbb{E}[R_1]\mathbb{E}[Y_1 \mid L_1 = i]\mathbb{P}(L_1 = i)$$

$$= N_G \sum_{i=1}^{N_R} p_R(X_i) \int_S \eta_i([X], y) y \, dy \mathbb{P}(L_1 = i). \quad \text{(D11)}$$

The covariance matrix $\Sigma_G = \mathbb{E}[(V_G - \langle V_G \rangle)(V_G - \langle V_G \rangle)^T]$ is:

$$\Sigma_G = N_G \mathbb{E}[(G_1 Y_1 - \langle G_1 Y_1 \rangle)(G_1 Y_1 - \langle G_1 Y_1 \rangle)^T]$$

$$+ (N_G^2 - N_G)\mathbb{E}[(G_1 Y_1 - \langle G_1 Y_1 \rangle)(G_2 Y_2 - \langle G_2 Y_2 \rangle)^T]. \quad \text{(D12)}$$

The first term is:

$$\mathbb{E}[(G_1 Y_1 - \langle G_1 Y_1 \rangle)(G_1 Y_1 - \langle G_1 Y_1 \rangle)^T] = \mathbb{E}[(G_1 Y_1)(G_1 Y_1)^T] - \mathbb{E}[(G_1 Y_1)]\mathbb{E}[(G_1 Y_1)^T] \quad \text{(D13)}$$

and

$$\mathbb{E}[(G_1 Y_1)(G_1 Y_1)^T] = \sum_{i=1}^{N_R} \mathbb{E}[(G_1 Y_1)(G_1 Y_1)^T \mid L_1 = i]\mathbb{P}(L_1 = i)$$

$$= \sum_{i=1}^{N_R} \mathbb{E}[R_i^2]\mathbb{E}[Y_1 Y_1^T \mid L_1 = i]\mathbb{P}(L_1 = i)$$

$$= \sum_{i=1}^{N_R} p_R(X_i) \int_S \eta_i([X], y) y y^T \, dy \mathbb{P}(L_1 = i). \quad \text{(D14)}$$

Next,

$$\mathbb{E}[(G_1 Y_1 - \langle G_1 Y_1 \rangle)(G_2 Y_2 - \langle G_2 Y_2 \rangle)^T] = \mathbb{E}[(G_1 Y_1)(G_2 Y_2)^T] - \mathbb{E}[(G_1 Y_1)]\mathbb{E}[(G_2 Y_2)^T]. \quad \text{(D15)}$$

and

$$\mathbb{E}[(G_1 Y_1)(G_2 Y_2)^T] = \sum_{m=1}^{N_R}\sum_{i=1}^{N_R} \mathbb{E}[(G_1 Y_1)(G_2 Y_2)^T \mid L_1 = i, L_2 = m]\mathbb{P}(L_1 = i, L_2 = m)$$

$$= \sum_{m=1}^{N_R}\sum_{i=1}^{N_R} \mathbb{E}[R_i R_m]\mathbb{E}[Y_1 Y_2^T \mid L_1 = i, L_2 = m]\mathbb{P}(L_1 = i)\mathbb{P}(L_2 = m).$$

$$\text{(D16)}$$

The case $i \neq m$ means that the two G proteins are linked to a different receptor (whose states are independent). In this case, $\mathbb{E}[R_i R_m] = \mathbb{E}[R_i]\mathbb{E}[R_m] = p_R(X_i)p_R(X_m)$. However, the case $i = m$ means that both G proteins are linked to the same receptor. By the calculation above (with $n = 2$), we have

$$\mathbb{E}[R_i(t_j)R_i(t_\ell)] = p_R(X_i)^2 + \frac{p_R(X_i)(1 - p_R(X_i))}{1 + \lambda(X_i)\tau}.$$

Therefore,

$$\mathbb{E}[(G_1Y_1)(G_2Y_2)^T] = \sum_{i=1}^{N_R} \left(p_R(X_i)^2 + \frac{p_R(X_i)(1 - p_R(X_i))}{1 + \lambda(X_i)\tau} \right) \mathbb{E}[Y_1Y_2^T \mid L_1 = L_2 = i]\mathbb{P}(L_1 = L_2 = i)$$
$$+ 2\sum_{i<m}^{N_R} p_R(X_i)p_R(X_m)\mathbb{E}[Y_1Y_2^T \mid L_1 = i, L_2 = m]\mathbb{P}(L_1 = i)\mathbb{P}(L_2 = m).$$

(D17)

Putting this all together, we obtain

$$\mathbb{E}[(G_1Y_1 - \langle G_1Y_1 \rangle)(G_2Y_2 - \langle G_2Y_2 \rangle)^T] = \sum_{i=1}^{N_R} \left(\frac{p_R(X_i)(1 - p_R(X_i))}{1 + \lambda(X_i)\tau} \right) \mathbb{E}[Y_1Y_2^T \mid L_1 = L_2 = i]\mathbb{P}(L_1 = L_2 = i)$$

(D18)

and the covariance matrix $\Sigma_G = \mathbb{E}[(V_G - \langle V_G \rangle)(V_G - \langle V_G \rangle)^T]$ is:

$$\Sigma_G = N_G \sum_{i=1}^{N_R} p_R(X_i) \int_S \eta_i([X], y)yy^T \, dy\mathbb{P}(L_1 = i)$$
$$- N_G \sum_{m=1}^{N_R}\sum_{i=1}^{N_R} p_R(X_m)p_R(X_i) \int_S\int_S \eta_m([X], z)\eta_i([X], y)yz^T \, dy \, dz\mathbb{P}(L_1 = i)\mathbb{P}(L_2 = m)$$
$$+ (N_G^2 - N_G) \sum_{i=1}^{N_R} \left(\frac{p_R(X_i)(1 - p_R(X_i))}{1 + \lambda(X_i)\tau} \right) \mathbb{E}[Y_1Y_2^T \mid L_1 = L_2 = i]\mathbb{P}(L_1 = L_2 = i).$$

(D19)

By definition of η_i,

$$\mathbb{E}[Y_1Y_2^T \mid L_1 = L_2 = i] = \int_S\int_S \eta_i([X], z)\eta_i([X], y)yz^T \, dy \, dz.$$

5. Derivation of Proposition D.2

For any continuous function f on the sphere S, when N_R is large enough we have

$$\frac{1}{N_R} \sum_{i=1}^{N_R} f(X_i) \approx \frac{1}{|S|} \int_S f(x) \, dx, \tag{D20}$$

with high probability. This is justified by the law of large numbers, since the receptor positions $\{X_i\}$ are independent. Using this and the approximations (D7) and (D8), we have

$$\mu_G = N_G \sum_{i=1}^{N_R} p_R(X_i) \int_S \eta_i([X], y) y \, dy \, \mathbb{P}(L_1 = i)$$

$$\approx N_G \left(\frac{1}{N_R} \sum_{i=1}^{N_R} p_R(X_i) X_i \right) \approx N_G \frac{1}{|S|} \int_S p_R(x) x \, dx, \tag{D21}$$

and similarly,

$$M_1 \approx \frac{1}{N_R} \sum_{i=1}^{N_R} p_R(X_i) X_i X_i^T - \left(\frac{1}{N_R} \sum_{i=1}^{N_R} p_R(X_i) X_i \right) \left(\frac{1}{N_R} \sum_{m=1}^{N_R} p_R(X_m) X_m \right)$$

$$\approx \frac{1}{|S|} \int_S p_R(x) x x^T \, dx - \left(\frac{1}{|S|} \int_S p_R(x) x \, dx \right) \left(\frac{1}{|S|} \int_S p_R(y) y^T \, dy \right). \tag{D22}$$

Since L_1 and L_2 are independent, $\mathbb{P}(L_1 = L_2 = i) = \mathbb{P}(L_1 = i)\mathbb{P}(L_2 = i) \approx (N_R)^{-2}$. Therefore, using the same reasoning as for M_1, we have

$$M_2 \approx \frac{1}{N_R} \sum_{i=1}^{N_R} \frac{p_R(X_i)(1 - p_R(X_i))}{1 + \tau\lambda(X_i)} X_i X_i^T \approx \frac{1}{|S|} \int_S \frac{p_R(x)(1 - p_R(x))}{1 + \tau\lambda(x)} x x^T \, dx.$$

Appendix E: Local concentration noise versus direction noise

Here is a simple illustration of the phenomenon that in the ratiometric model the gradient of active G protein may be stronger than in the classical model even while the local noise in active G protein may be greater for the ratiometric model. Thus, high-variability in local concentration of active G protein is not necessarily an indicator of poor direction estimation. This is because direction estimation involves a comparison across multiple different regions of the cell surface.

Imagine a greatly simplified, one-dimensional model where the cell has a front and a back: at the front there are N_R receptors and at the back there are N_R receptors. Let $\epsilon \in (0, 1/2)$ be a small parameter representing a small gradient of ligand concentration (toward the front end). Let us suppose that R_f^+ and R_b^+ are independent random variables with distributions

$$R_f^+ \sim \text{Binomial}\left(N_R, \frac{1}{2} + \epsilon\right), \qquad R_b^+ \sim \text{Binomial}\left(N_R, \frac{1}{2} - \epsilon\right).$$

These R_f^+ and R_b^+ represent the number of active receptors at the front and back of the cell, respectively. This statistical model is equivalent to saying that the N_R proteins at the front

end have independent states, each being active with probability $\frac{1}{2} + \epsilon$. Similarly, at the back end, the probability of a receptor being active is $\frac{1}{2} - \epsilon$. Then define

$$R_f^- = N_R - R_f^+, \qquad R_b^- = N_R - R_b^+,$$

which represent the number of inactive receptor at the front and back of the cell, respectively. The mean receptor levels are

$$\mathbb{E}[R_f^{\pm}] = N_R \left(\frac{1}{2} \pm \epsilon \right), \qquad \mathbb{E}[R_b^{\pm}] = N_R \left(\frac{1}{2} \mp \epsilon \right)$$

and the variances are

$$\mathrm{Var}(R_f^+) = \mathrm{Var}(R_f^-) = \mathrm{Var}(R_b^+) = \mathrm{Var}(R_b^-) = N_R \left(\frac{1}{2} + \epsilon \right) \left(\frac{1}{2} - \epsilon \right).$$

So, with $\epsilon > 0$, the receptor gradient has a small bias toward the front end:

$$\mathbb{E}[R_f^+ - R_b^+] = 2N_R \epsilon > 0.$$

In addition to receptor, there are G proteins at the front and back. Suppose there are N_G proteins at the front and N_G proteins at the back. Random variables G_f^+ and G_b^+ will be the number of active G proteins at the front and back, respectively. Given R_f^+ and R_f^-, G proteins at the front end will be active with probability that depends on R_f^+ and R_f^-; what distinguishes the ratiometric from classical models is how this probability depends on R_f^+ and R_f^-. In the ratiometric model, the steady-state probability that a G protein at the front end is active is

$$\frac{R_f^+}{R_f^+ + R_f^-} = \frac{R_f^+}{N_R}.$$

(Rate constants are normalized to one.) In the classical model, the inactivation rate is taken to be constant and uniform across the cell (same at the front and back). Therefore, in the classical model, the steady state probability that a G protein at the front end is active has the form

$$\frac{R_f^+/N_R}{R_f^+/N_R + c}$$

for some constant $c > 0$. Similar statements hold at the back end of the cell, with R_b^+ replacing R_f^+. In either case, classical or ratiometic, the probability that a G protein is active, is a function of the local fraction of active receptor.

In view of this, we take G_f^+ and G_b^+ to be independent random variables having the following distribution: given R_f^+ and R_b^+,

$$G_f^+ \sim \text{Binomial}\left(N_G, f\left(\frac{R_f^+}{N_R}\right)\right), \qquad G_b^+ \sim \text{Binomial}\left(N_G, f\left(\frac{R_b^+}{N_R}\right)\right).$$

where

$$f(r) = \begin{cases} r, & \text{for the ratiometric model,} \\ \frac{r}{r+c}, & \text{for the classical model.} \end{cases}$$

This statistical model is equivalent to saying that the N_G proteins at the front end have independent states, each being active with probability $f\left(\frac{R_f^+}{N_R}\right)$. And similarly for the proteins at the back end.

To compare the two models, we choose the constant c so that the mean total activation rate of G protein is the same. The mean total activation is

$$\mathbb{E}[G_f^+ + G_b^+] = N_G\mathbb{E}[f(R_f^+/N_R)] + N_G\mathbb{E}[f(R_b^+/N_R)].$$

For the ratiometric model, the function f is linear ($f(r) = r$), so this can be computed explicitly:

$$\mathbb{E}[G_f^+ + G_b^+] = N_G\mathbb{E}[R_f^+/N_R] + N_G\mathbb{E}[R_b^+/N_R] = N_G.$$

Therefore, the matching condition requires that the constant c (in the classical model) be chosen so that

$$\mathbb{E}\left[\frac{R_f^+/N_R}{R_f^+/N_R + c}\right] + \mathbb{E}\left[\frac{R_b^+/N_R}{R_b^+/N_R + c}\right] = 1. \tag{E1}$$

This relation for c cannot be solved explicitly. Nevertheless, we claim that when N_R is large and ϵ is small, $c \approx 1/2$. To see this, note that when N_R is large, the distributions of R_f/N_R and R_b/N_R are very concentrated around $\frac{1}{2} + \epsilon$ and $\frac{1}{2} - \epsilon$, respectively. Therefore, when $\epsilon > 0$ is small, (E1) implies that

$$\frac{\frac{1}{2} + \epsilon}{\frac{1}{2} + \epsilon + c} + \frac{\frac{1}{2} - \epsilon}{\frac{1}{2} - \epsilon + c} \approx 1,$$

which implies that $c \approx 1/2$ for small ϵ.

In the case of the classical model, f is nonlinear, and this makes it impossible to calculate explicitly the mean and variance of G_f^+ and G_b^+ in that case. However, as we explained above,

when N_R is large and ϵ is small, the constant c is approximately $1/2$, and the function f may be approximated well by its linear expansion at the value $r = 1/2$:

$$\frac{r}{r + 1/2} \approx \frac{1}{2} + \frac{1}{2}\left(r - \frac{1}{2}\right)$$

This motivates the following modification, which makes calculation explicit in both models: Given R_f^+ and R_b^+, let

$$G_f^+ \sim \text{Binomial}\left(N_G, \ell\left(\frac{R_f^+}{N_R}\right)\right), \qquad G_b^+ \sim \text{Binomial}\left(N_G, \ell\left(\frac{R_b^+}{N_R}\right)\right).$$

where, instead of f defined above, we use the function

$$\ell(r) = \begin{cases} r, & \text{for the ratiometric model,} \\ \frac{1}{2} + \frac{1}{2}\left(r - \frac{1}{2}\right), & \text{for the classical model.} \end{cases} \tag{E2}$$

This ℓ is linear in both cases, and it has the form

$$\ell(r) = \frac{1}{2} + \alpha\left(r - \frac{1}{2}\right), \qquad \alpha = \begin{cases} 1, & \text{for the ratiometric model,} \\ \frac{1}{2}, & \text{for the classical model.} \end{cases}$$

With this modification, the means and variances of G_f^+ and G_b^+ are computable in both models. We compute:

$$\mathbb{E}[G_f^+] = N_G\mathbb{E}\left[\ell\left(\frac{R_f^+}{N_R}\right)\right] = N_G\left(\frac{1}{2} + \alpha\left(\mathbb{E}[R_f^+/N_R] - \frac{1}{2}\right)\right) = N_G\left(\frac{1}{2} + \alpha\epsilon\right), \tag{E3}$$

$$\mathbb{E}[G_b^+] = N_G\mathbb{E}\left[\ell\left(\frac{R_b^+}{N_R}\right)\right] = N_G\left(\frac{1}{2} + \alpha\left(\mathbb{E}[R_b^+/N_R] - \frac{1}{2}\right)\right) = N_G\left(\frac{1}{2} - \alpha\epsilon\right). \tag{E4}$$

In particular, the total activation level is the same for all α:

$$\mathbb{E}[G_f^+ + G_b^+] = N_G, \tag{E5}$$

while the mean front-to-back gradient in active G protein is

$$\mathbb{E}[G_f^+ - G_b^+] = 2N_G\alpha\epsilon = \alpha(N_G/N_R)\mathbb{E}[R_f^+ - R_b^+]. \tag{E6}$$

Recall that the ratiometric case corresponds exactly to $\alpha = 1$, while the classical case corresponds to $\alpha = 1/2$, so the mean gradient is twice as large in the ratiometric case.

Now, let us consider how the variances depend on α, especially the cases $\alpha = 1$ (ratiometric) and $\alpha = 1/2$ (classical). The variance in active G protein, at both the front and back end, can be computed explicitly:

$$\mathrm{Var}(G_f^+) = N_G \mathbb{E}\left[\ell\left(\frac{R_f^+}{N_R}\right)\right]\left(1 - \mathbb{E}\left[\ell\left(\frac{R_f^+}{N_R}\right)\right]\right) + (N_G^2 - N_G)\mathrm{Var}\left(\ell\left(\frac{R_f^+}{N_R}\right)\right) \quad \text{(E7)}$$

$$= N_G\left(\frac{1}{2} + \alpha\epsilon\right)\left(\frac{1}{2} - \alpha\epsilon\right) + (N_G^2 - N_G)\frac{\alpha^2}{N_R}\left(\frac{1}{2} + \alpha\epsilon\right)\left(\frac{1}{2} - \alpha\epsilon\right). \quad \text{(E8)}$$

So, the coefficient of variation of G_f^+ is

$$\mathrm{CV}^2(G_f^+) = \frac{\mathrm{Var}(G_f^+)}{\mathbb{E}[G_f^+]^2} = \frac{N_G\left(\frac{1}{2} + \alpha\epsilon\right)\left(\frac{1}{2} - \alpha\epsilon\right) + (N_G^2 - N_G)\frac{\alpha^2}{N_R}\left(\frac{1}{2} + \alpha\epsilon\right)\left(\frac{1}{2} - \alpha\epsilon\right)}{\left(N_G\left(\frac{1}{2} + \alpha\epsilon\right)\right)^2} \quad \text{(E9)}$$

$$= \left(\frac{1}{N_G} + \left(1 - \frac{1}{N_G}\right)\frac{\alpha^2}{N_R}\right)\frac{\left(\frac{1}{2} - \alpha\epsilon\right)}{\left(\frac{1}{2} + \alpha\epsilon\right)}. \quad \text{(E10)}$$

A similar compuation shows that at the back end we have

$$\mathrm{CV}^2(G_b^+) = \left(\frac{1}{N_G} + \left(1 - \frac{1}{N_G}\right)\frac{\alpha^2}{N_R}\right)\frac{\left(\frac{1}{2} + \alpha\epsilon\right)}{\left(\frac{1}{2} - \alpha\epsilon\right)}. \quad \text{(E11)}$$

For small ϵ, this is approximately

$$\mathrm{CV}^2(G_f^+) \approx \frac{1}{N_G} + \left(1 - \frac{1}{N_G}\right)\frac{\alpha^2}{N_R}, \qquad \mathrm{CV}^2(G_b^+) \approx \frac{1}{N_G} + \left(1 - \frac{1}{N_G}\right)\frac{\alpha^2}{N_R}.$$

In particular, these quantities increase with α for $\alpha \in [0, 1]$. On the other hand, the mean front-to-back gradient

$$\mathbb{E}[G_f^+ - G_b^+] = 2N_G\alpha\epsilon$$

also increases with α. This shows that the directional signal gets stronger with larger α, even though the local noise levels (measured by coefficients of variation $\mathrm{CV}^2(G_f^+)$ and $\mathrm{CV}^2(G_b^+)$) are larger with larger α. In particular, this illustrates the phenomenon that with the ratiometric model ($\alpha = 1$), the local noise in G_f^+ and G_b^+ may be greater than in the classical model ($\alpha = 1/2$), even while the front-to-back gradient in G protein is stronger for the ratiometric model.

This phenomenon can be seen more clearly from examining the joint distributions of $(R_f^+/N_R, G_f^+/N_G)$ and $(R_b^+/N_R, G_b^+/N_G)$ for the two models, see Figure 11. In Figure 12, we plot histrograms representing the marginal distributions of G_f^+/N_G and G_b^+/N_G for the two models. The marginal distributions of G_f^+/N_G and G_b^+/N_G have higher variance for the

ratiometric model, than for the classical model. Figure 13 shows a histrogram representing the distribution of active G protein gradient $(G_f^+ - G_b^+)/N_G$ for both models. We observe that the distribution of this gradient is biased to the right (front end) for the ratiometric model, as explained in the analysis above. From Figure 11, we see how the different activation functions for the ratiometric vs. classical models leads to a bias in the G protein gradient from front to back, producing stronger (steeper) gradient in the ratiometric case, even while the variance in local G protein activation is higher with the ratiometric model.

Appendix F: Classical model

Here we explain why the classical model (nonratiometric) performs more poorly than the ratiometric model. Recall that V_G^r and V_G^c refer to the vector (B4) under the ratiometric and classical models, respectively. We have seen that the ratiometric model V_G^r performs particularly well compared to V_{RL} and V_C when the time scales are such that the G proteins are approximately independent, and $N_G > N_R$. In principle, the classical model estimate V_G^c might also perform well in this scenario, yet our simulations show that this is not the case. An important point derived above is that for the ratiometric model, with N_R sufficiently large, the mean resultant vector is

$$\mathbb{E}[V_G^r] = \frac{N_G}{|S|} \int_S p_G^r(y) y \, dy \approx \frac{N_G}{|S|} \int_S p_R(y) y \, dy$$

where $p_G^r(y)$ is the probability that a G protein at location y is active (under the ratiometric model); $p_R(y)$ is the probability that a receptor at y is active. In particular, the mean vector $\mathbb{E}[V_G^r]$ does not depend on $\lambda_0 \tau_{\text{dif}}$, where $\tau_{\text{dif}} = \tau$ is the diffusive encounter time scale from (D1), and λ_0^{-1} is the time scale of receptor switching. For the classical model, there is an additional time scale $\tau_{\text{off}} = 1/k_i'$, where k_i' is the rate at which G proteins deactivate randomly in the classical model. For the classical model, the mean vector $\mathbb{E}[V_G^c]$ depends on all of these three time scales τ_{off}, τ_{dif}, and λ_0^{-1} in a way that makes the classical model less effective at direction estimation than the ratiometric model.

For the classical model, let $p_G^c(y)$ denote the probability that a G protein at position y is active. The probability p_G^c cannot be computed explicitly, but we analyze an approximate model that has similar features and gives insight into the way that p_G^c depends on the time scales. As we will show, the gradient of p_G^c may be much flatter than that of p_R or p_G^r. In

the classical model, the state $G(t)$ of a G protein at time t and position $Y(t) = y$ is active if and only if that G protein encountered an active receptor (possibly multiple) at some time between time $t - T$ and t, where the random time T has the Exponential distribution with mean τ_{off}. Otherwise, that protein is inactive. The number of receptors that a G protein encounters before deactivation is random and it depends in a complicated way on the receptor configuration, but it is reasonable to suppose that random number N of distinct encounters has mean $\mathbb{E}[N] = \tau_{\text{off}}/\tau_{\text{dif}}$, where

$$\tau_{\text{dif}} = \frac{L^2}{N_R D_G} \ln(\delta/r_*)$$

is the diffusive encounter time scale (see (D1)). Indeed, looking backward in time we may assume that the encounters between a G protein and receptors occur as a Poisson arrival process at rate $1/\tau_{\text{dif}}$ (times between encounters are approximately exponentially distributed with mean τ_{dif}). Thus, N has the distribution of A_T, where A_t is such a Poisson arrival process and $T \sim \text{Exponential}(1/\tau_{\text{off}})$ is independent of the process A. This implies that the number N of distinct receptor encounters has the (shifted) Geometric distribution:

$$\mathbb{P}(N = n) = \int_0^\infty \mathbb{P}(A_t = n \mid T = t) \frac{1}{\tau_{\text{off}}} e^{-t/\tau_{\text{off}}} \, dt$$
$$= \left(\frac{\tau_{\text{dif}}}{\tau_{\text{dif}} + \tau_{\text{off}}} \right) \left(\frac{\tau_{\text{off}}}{\tau_{\text{dif}} + \tau_{\text{off}}} \right)^n, \qquad n = 0, 1, 2, 3, \ldots \tag{F1}$$

with mean $\mathbb{E}[N] = \tau_{\text{off}}/\tau_{\text{dif}}$. This distribution is in close agreement with our simulations; see Figure 14 where we show numerical computation of the empirical distribution of distinct receptors encountered by a G protein before its state resets (in the classical model).

An important consequence of this is that the probability that a G protein encounters zero receptors between resetting events is:

$$\mathbb{P}(N = 0) = \frac{1}{1 + \tau_{\text{off}}/\tau_{\text{dif}}} = \frac{1}{1 + \mathbb{E}[N]}.$$

Such G proteins carry zero information about the receptor states, and thus zero information about the pheromone gradient or local concentration. This is an important difference between the classical and ratiometric models: in the classical model, at a given time t, some fraction (approximately $\frac{1}{1+\tau_{\text{off}}/\tau_{\text{dif}}}$) of proteins carry zero information about the local pheromone concentration because they have not encountered any receptor (active or inactive) since the most recent state resetting event; in the ratiometric model, however, every G

protein carries some information about the local pheromone concentration, via the state of the most recently encountered receptor. Note that $\mathbb{P}(N = 0)$ does not depend in any way on the pheromone concentration; $\mathbb{E}[N]$ depends on the density of receptors, but not on their states.

The presence of "uninformative" G proteins (in the classical model) carrying no information about the pheromone gradient leads to what we call a flattening of the gradient of p_G^c relative to p_R. Assuming encounters are nearly instantaneous, the probability that a given encounter at location x activates the G protein is approximately $p_R(x)$. Assuming that the N encounters are local, then the probability that a G protein at x (under classical model) will be active is then

$$p_G^c(x) = 1 - \mathbb{E}[(1 - p_R(x))^N] \tag{F2}$$

$$= 1 - \sum_{k=0}^{\infty}(1 - p_R(x))^k q(1-q)^k, \qquad q = \frac{\tau_{\text{dif}}}{\tau_{\text{dif}} + \tau_{\text{off}}} \tag{F3}$$

$$= \frac{p_R(x)(1-q)}{q + p_R(x)(1-q)} \tag{F4}$$

$$= \frac{p_R(x)\mathbb{E}[N]}{1 + p_R(x)\mathbb{E}[N]}. \tag{F5}$$

Thus the relationship between p_G^c and p_R depends on the mean number of receptor encounters $\mathbb{E}[N]$. In Figure 15 we plot p_G^c versus p_R for different values of $\mathbb{E}[N]$. When $\tau_{\text{off}}/\tau_{\text{diff}} = \mathbb{E}[N] < < 1$, most G proteins will be inactive, and (F5) shows that

$$\mathbb{E}[N] \ll 1 \qquad \implies \qquad p_G^c(x) \approx 0, \quad \text{for all } x. \tag{F6}$$

However, when $\tau_{\text{off}}/\tau_{\text{diff}} = \mathbb{E}[N] \gg 1$ and p_R not too small, there is saturation – most G proteins will be active, and (F5) shows that

$$\mathbb{E}[N] \gg 1 \qquad \implies \qquad p_G^c(x) \approx 1, \quad \text{for all } x. \tag{F7}$$

The gradient of p_G^c is

$$\nabla p_G^c(x) = \frac{\mathbb{E}[N]}{(1 + p_R(x)\mathbb{E}[N])^2}\nabla p_R(x). \tag{F8}$$

The prefactor in front of ∇p_R can be larger or smaller than 1. We have

$$\frac{\mathbb{E}[N]}{(1 + p_R(x)\mathbb{E}[N])^2} \geq 1$$

if only if $p_R(x) < \frac{1}{\sqrt{\mathbb{E}[N]}} - \frac{1}{\mathbb{E}[N]}$. In particular, if $p_R(x) > \frac{1}{\sqrt{\mathbb{E}[N]}} - \frac{1}{\mathbb{E}[N]}$, then there will necessarily be a flattening effect whereby the prefactor in (F8) is less than 1. Our earlier analysis shows that for shallow gradients, the NSR is smallest when $C_0 \approx K_d$, which implies p_R is near 1/2. In this regime, p_G^c will be flatter than p_R, as shown in Figure 15. In particular, p_G^c is flatter than p_G^r, since for the ratiometric model $p_G^r \approx p_R$ regardless of the time scales. The flattening of p_G^c versus p_R and p_G^r implies less accuracy (higher NSR) for the classical method, since the NSR scales inversely with the gradient squared (i.e. as $1/b^2$ in (B12).)

Appendix G: Captions for Movies

Movie S1. Temporal dynamics of receptor and G-protein vector orientations during Classical gradient sensing. **Top left:** Full simulation showing all receptor and G-protein positions; receptors are color-coded by activation state. **Top middle:** Vector orientations for active (yellow), inactive (gray), and combined (blue) receptor populations. **Top right:** Vectors for the combined receptor (blue) and G-protein (green) populations. The black horizontal line denotes the true gradient direction. **Bottom:** Time traces showing the angle between the pheromone gradient and the receptor (green) and G-protein (cyan) vectors. NG = 2500, NR = 375, $D_m = 0.002$ $\mu m^2/s$, $k_{gi} = 0.0083$ s^{-1}, gradient = 8 to 4 nM. https://youtu.be/beVWFQ6Jzdg

Movie S2. Temporal dynamics of receptor and G-protein vector orientations during Classical gradient sensing. **Top left:** Full simulation showing all receptor and G-protein positions; receptors are color-coded by activation state. **Top middle:** Vector orientations for active (yellow), inactive (gray), and combined (blue) receptor populations. **Top right:** Vectors for the combined receptor (blue) and G-protein (green) populations. The black horizontal line denotes the true gradient direction. **Bottom:** Time traces showing the angle between the pheromone gradient and the receptor (green) and G-protein (cyan) vectors. NG = 2500, NR = 1500, $D_m = 0.002$ $\mu m^2/s$, $k_{gi} = 0.0357$ s^{-1}, gradient = 8 to 4 nM. https://youtu.be/nvEN7Sysycc

Movie S3. Temporal dynamics of receptor and G-protein vector orientations during Classical gradient sensing. **Top left:** Full simulation showing all receptor and G-protein positions; receptors are color-coded by activation state. **Top middle:** Vector orientations for active (yellow), inactive (gray), and combined (blue) receptor populations. **Top right:**

Vectors for the combined receptor (blue) and G-protein (green) populations. The black horizontal line denotes the true gradient direction. **Bottom:** Time traces showing the angle between the pheromone gradient and the receptor (green) and G-protein (cyan) vectors. NG = 2500, NR = 6000, D_m = 0.002 $\mu m^2/s$, k_{gi} = 0.1687 s^{-1}, gradient = 8 to 4 nM. https://youtu.be/egZNPnUeXfM

Movie S4. Temporal dynamics of receptor and G-protein vector orientations during Ratiometric gradient sensing. **Top left:** Full simulation showing all receptor and G-protein positions; receptors are color-coded by activation state. **Top middle:** Vector orientations for active (yellow), inactive (gray), and combined (blue) receptor populations. **Top right:** Vectors for the combined receptor (blue) and G-protein (green) populations. The black horizontal line denotes the true gradient direction. **Bottom:** Time traces showing the angle between the pheromone gradient and the receptor (green) and G-protein (cyan) vectors. NG = 2500, NR = 375, D_m = 0.002 $\mu m^2/s$, gradient = 8 to 4 nM. https://youtu.be/B3Dg8SMIhEE

Movie S5. Temporal dynamics of receptor and G-protein vector orientations during Ratiometric gradient sensing. **Top left:** Full simulation showing all receptor and G-protein positions; receptors are color-coded by activation state. **Top middle:** Vector orientations for active (yellow), inactive (gray), and combined (blue) receptor populations. **Top right:** Vectors for the combined receptor (blue) and G-protein (green) populations. The black horizontal line denotes the true gradient direction. **Bottom:** Time traces showing the angle between the pheromone gradient and the receptor (green) and G-protein (cyan) vectors. NG = 2500, NR = 1500, D_m = 0.002 $\mu m^2/s$, gradient = 8 to 4 nM. https://youtu.be/9SNJQRN9yAQ

Movie S6. Temporal dynamics of receptor and G-protein vector orientations during Ratiometric gradient sensing. **Top left:** Full simulation showing all receptor and G-protein positions; receptors are color-coded by activation state. **Top middle:** Vector orientations for active (yellow), inactive (gray), and combined (blue) receptor populations. **Top right:** Vectors for the combined receptor (blue) and G-protein (green) populations. The black horizontal line denotes the true gradient direction. **Bottom:** Time traces showing the angle between the pheromone gradient and the receptor (green) and G-protein (cyan) vectors. NG = 2500, NR = 6000, D_m = 0.002 $\mu m^2/s$, gradient = 8 to 4 nM. https://youtu.be/

[1] Wilhelm Engelmann. Neue methode zur untersuchung der sauerstoffausscheidung pflanzlicher und thierischer organismen. *Archiv für die gesamte Physiologie des Menschen und der Tiere*, 25(1):285–292, 1881.

[2] Gerhart Drews. Contributions of Theodor Wilhelm Engelmann on phototaxis, chemotaxis, and photosynthesis. *Photosynthesis Research*, 83(1):25–34, January 2005.

[3] Wilhelm Pfeffer. *Locomotorische Richtungsbewegungen durch chemische Reize:(Aus den" Untersuchungen aus dem botanischen Institut zu Tübingen Bd. I. Heft 3 p. 363-482)*. W. Engelmann, 1884.

[4] Hans Molisch. Zur physiologie des pollens, mit besonderer rücksicht auf die chemotropischen bewegungen der pollenschläuche. *Sitzungsberichte*, 102:423–448, 1893.

[5] Bengt Lidforss. Untersuchungen über die reizbewegungen der pollenschläuche. 1. der chemotropismus. *Zeitschrift für Botanik*, 1(7):443–496, 1909.

[6] Herbert Spencer Jennings. *Contributions to the study of the behavior of lower organisms*. Number 16. Carnegie institution of Washington, 1904.

[7] John Tyler Bonner and LJ Savage. Evidence for the formation of cell aggregates by chemotaxis in the development of the slime mold dictyostelium discoideum. *Journal of Experimental Zoology*, 106(1):1–26, 1947.

[8] Theo M Konijn, JG Van De Meene, John T Bonner, and David S Barkley. The acrasin activity of adenosine-3', 5'-cyclic phosphate. *Proceedings of the National Academy of Sciences*, 58(3):1152–1154, 1967.

[9] J E Segall. Polarization of yeast cells in spatial gradients of alpha mating factor. *Proceedings of the National Academy of Sciences*, 90(18):8332–8336, September 1993. Publisher: Proceedings of the National Academy of Sciences.

[10] Debraj Ghose, Katherine Jacobs, Samuel Ramirez, Timothy Elston, and Daniel Lew. Chemotactic movement of a polarity site enables yeast cells to find their mates. *Proceedings of the National Academy of Sciences*, 118(22), June 2021. Publisher: National Academy of Sciences Section: Biological Sciences.

[11] Jayme M. Dyer, Natasha S. Savage, Meng Jin, Trevin R. Zyla, Timothy C. Elston, and

Daniel J. Lew. Tracking shallow chemical gradients by actin-driven wandering of the polarization site. *Current Biology*, 23(1):32–41, 2013.

[12] Karen M Snetselaar, Michael Bölker, and Regine Kahmann. Ustilago maydismating hyphae orient their growth toward pheromone sources. *Fungal genetics and biology*, 20(4):299–312, 1996.

[13] Arthur Henry Reginald Buller. *Researches on Fungi*, volume 2. Longmans, Green and Company, 1922.

[14] Anton Bary. *Comparative morphology and biology of the fungi, mycetozoa and bacteria*. Clarendon press, 1887.

[15] Ilya Metchnikoff. Untersuchungen über die mesodermalen phagocyten einiger wirbeltiere. *Biol Zentralbl*, 3:560–5, 1883.

[16] Theodor Leber. *Die Entstehung der Entzündung und die Wirkung der entzündungerregenden Schädlichkeiten: nach vorzugsweise am Auge angestellten Untersuchungen*. Wilhelm Engelmann, Leipzig, 1891.

[17] Frank R Lillie. The production of sperm iso-agglutinins by ova. *Science*, 36(929):527–530, 1912.

[18] Richard L Miller. Chemotaxis during fertilization in the hydroid campanularia. *Journal of Experimental Zoology*, 162(1):23–44, 1966.

[19] Marc Spehr, Gunter Gisselmann, Alexandra Poplawski, Jeffrey A Riffell, Christian H Wetzel, Richard K Zimmer, and Hanns Hatt. Identification of a testicular odorant receptor mediating human sperm chemotaxis. *Science*, 299(5615):2054–2058, 2003.

[20] Ross Granville Harrison. The outgrowth of the nerve fiber as a mode of protoplasmic movement. *Journal of Experimental Zoology*, 9(4):787–846, 1910.

[21] Timothy E Kennedy, Tito Serafini, JoséR De La Torre, and Marc Tessier-Lavigne. Netrins are diffusible chemotropic factors for commissural axons in the embryonic spinal cord. *Cell*, 78(3):425–435, 1994.

[22] Hans Spemann and Hilde Mangold. über induktion von embryonalanlagen durch implantation artfremder organisatoren. *Archiv für mikroskopische Anatomie und Entwicklungsmechanik*, 100:599–638, 1924.

[23] Kenshi Yoshida, Teruhisa Ozaki, Kenichiro Ushijima, and Hideo Hayashi. Studies on the mechanisms of invasion in cancer. i. isolation and purification of a factor chemotactic for

cancer cells. *International Journal of Cancer*, 6(1):123–132, 1970.

[24] Maryse Bailly, Lin Yan, George M Whitesides, John S Condeelis, and Jeffrey E Segall. Reg-
ulation of protrusion shape and adhesion to the substratum during chemotactic responses of
mammalian carcinoma cells. *Experimental cell research*, 241(2):285–299, 1998.

[25] H Hayashi, K Yoshida, T Ozaki, and K Ushijima. Chemotactic factor associated with invasion
of cancer cells. *Nature*, 226(5241):174–175, 1970.

[26] Julius Adler. Chemoreceptors in bacteria: Studies of chemotaxis reveal systems that detect
attractants independently of their metabolism. *Science*, 166(3913):1588–1597, 1969.

[27] Randall L. Johnson and Peter N. Devreotes. The cyclic amp receptor in dictyostelium dis-
coideum. i. sequence of events following receptor activation. *The Journal of Cell Biology*,
88(3):734–740, 1981.

[28] H. C. Berg and E. M. Purcell. Physics of chemoreception. *Biophysical Journal*, 20(2):193–219,
November 1977.

[29] William Bialek and Sima Setayeshgar. Physical limits to biochemical signaling. *Proceedings*
of the National Academy of Sciences of the United States of America, 102(29):10040–10045,
July 2005.

[30] Kazunari Kaizu, Wiet de Ronde, Joris Paijmans, Koichi Takahashi, Filipe Tostevin, and
Pieter Rein ten Wolde. The Berg-Purcell limit revisited. *Biophysical Journal*, 106(4):976–985,
February 2014.

[31] Robert G. Endres and Ned S. Wingreen. Accuracy of direct gradient sensing by cell-surface
receptors. *Progress in Biophysics and Molecular Biology*, 100(1-3):33–39, October 2009.

[32] Bo Hu, Wen Chen, Wouter-Jan Rappel, and Herbert Levine. Physical limits on cellular sensing
of spatial gradients. *Physical review letters*, 105(4):048104, 2010.

[33] Nicholas T. Henderson, Michael Pablo, Debraj Ghose, Manuella R. Clark-Cotton, Trevin R.
Zyla, James Nolen, Timothy C. Elston, and Daniel J. Lew. Ratiometric GPCR signaling
enables directional sensing in yeast. *PLoS biology*, 17(10):e3000484, October 2019.

[34] Alan Bush, Gustavo Vasen, Andreas Constantinou, Paula Dunayevich, Inés Lucía Patop,
Matías Blaustein, and Alejandro Colman-Lerner. Yeast gpcr signaling reflects the fraction of
occupied receptors, not the number. *Molecular systems biology*, 12(12):898, 2016.

[35] Vinal Lakhani and Timothy C. Elston. Testing the limits of gradient sensing. *PLoS compu-*
tational biology, 13(2):e1005386, February 2017.

[36] Masataka Yanagawa, Michio Hiroshima, Yuichi Togashi, Mitsuhiro Abe, Takahiro Yamashita, Yoshinori Shichida, Masayuki Murata, Masahiro Ueda, and Yasushi Sako. Single-molecule diffusion-based estimation of ligand effects on G protein–coupled receptors. *Science Signaling*, 11(548):eaao1917, September 2018.

[37] Harrie Hendriks. A Cramér-Rao type lower bound for estimators with values in a manifold. *Journal of Multivariate Analysis*, 38(2):245–261, August 1991.

[38] Debraj Ghose, Timothy Elston, and Daniel Lew. Orientation of Cell Polarity by Chemical Gradients. *Annual Review of Biophysics*, 51(1):null, 2022. _eprint: https://doi.org/10.1146/annurev-biophys-110821-071250.

[39] Khem Raj Ghusinga, Roger D Jones, Alan M Jones, and Timothy C Elston. Molecular switch architecture determines response properties of signaling pathways. *Proceedings of the National Academy of Sciences*, 118(11):e2013401118, 2021.

[40] Sally H Zigmond. Ability of polymorphonuclear leukocytes to orient in gradients of chemotactic factors. *Journal of Cell Biology*, 75(2):606–616, 1977.

[41] G Gerisch and U Wick. Intracellular oscillations and release of cyclic amp from dictyostelium cells. *Biochemical and biophysical research communications*, 65(1):364–370, 1975.

[42] Daniel R. Ballon, Paul L. Flanary, Douglas P. Gladue, James B. Konopka, Henrik G. Dohlman, and Jeremy Thorner. DEP-domain-mediated regulation of GPCR signaling responses. *Cell*, 126(6):1079–1093, September 2006.

[43] Christopher G Alvaro and Jeremy Thorner. Heterotrimeric g protein-coupled receptor signaling in yeast mating pheromone response. *Journal of Biological Chemistry*, 291(15):7788–7795, 2016.

[44] Anita Emmerstorfer-Augustin, Christoph M Augustin, Shadi Shams, and Jeremy Thorner. Tracking yeast pheromone receptor ste2 endocytosis using fluorogen-activating protein tagging. *Molecular biology of the cell*, 29(22):2720–2736, 2018.

[45] Christopher DC Allen, K Mark Ansel, Caroline Low, Robin Lesley, Hirokazu Tamamura, Nobutaka Fujii, and Jason G Cyster. Germinal center dark and light zone organization is mediated by cxcr4 and cxcr5. *Nature immunology*, 5(9):943–952, 2004.

[46] Abdelhak Belmadani, Phuong B Tran, Dongjun Ren, Stavroula Assimacopoulos, Elizabeth A Grove, and Richard J Miller. The chemokine stromal cell-derived factor-1 regulates the migration of sensory neuron progenitors. *Journal of Neuroscience*, 25(16):3995–4003, 2005.

[47] Michele K Dougherty, Jürgen Müller, Daniel A Ritt, Ming Zhou, Xiao Zhen Zhou, Terry D Copeland, Thomas P Conrads, Timothy D Veenstra, Kun Ping Lu, and Deborah K Morrison. Regulation of raf-1 by direct feedback phosphorylation. *Molecular cell*, 17(2):215–224, 2005.

[48] Piro Lito, Christine A Pratilas, Eric W Joseph, Madhavi Tadi, Ensar Halilovic, Matthew Zubrowski, Alan Huang, Wai Lin Wong, Margaret K Callahan, Taha Merghoub, et al. Relief of profound feedback inhibition of mitogenic signaling by raf inhibitors attenuates their activity in brafv600e melanomas. *Cancer cell*, 22(5):668–682, 2012.

[49] Ty M. Thomson, Kirsten R. Benjamin, Alan Bush, Tracy Love, David Pincus, Olga Resnekov, Roy C. Yu, Assaf Gordon, Alejandro Colman-Lerner, Drew Endy, and Roger Brent. Scaffold-mediated negative feedback regulates a yeast mapk pathway. *Molecular Systems Biology*, 7:460, 2011.

[50] Kanti V. Mardia and Peter E. Jupp. *Directional Statistics*. Wiley Series in Probability and Statistics. John Wiley & Sons, Chichester, 2000.

[51] S.T. Smith. Covariance, subspace, and intrinsic Crame/spl acute/r-Rao bounds. *IEEE Transactions on Signal Processing*, 53(5):1610–1630, May 2005. Conference Name: IEEE Transactions on Signal Processing.

[52] David Holcman and Zeev Schuss. Time scale of diffusion in molecular and cellular biology. *Journal of Physics A: Mathematical and Theoretical*, 47(17):173001, 2014.

[53] C Chevalier, O Bénichou, B Meyer, and R Voituriez. First-passage quantities of Brownian motion in a bounded domain with multiple targets: a unified approach. *Journal of Physics A: Mathematical and Theoretical*, 44(2):025002, December 2010.

FIG. 1. **Estimating gradient direction using information from both active and inactive receptors.** (**A**) Schematic depicting a spherical cell with receptors randomly distributed on its surface. Each receptor can be active (ligand-bound, yellow) or inactive (unbound, gray), and the gradient direction is inferred by summing receptor vectors and normalizing the sums to obtain unit vectors $\bar{V}_{RL}$, $\bar{V}_R$, and $\bar{V}_C$. Note that $\bar{V}_R$ is flipped to obtain $-\bar{V}_R$ to estimate gradient direction. (**B**) Cumulative distribution functions (CDFs) of angular error for the estimates $\bar{V}_{RL}$, $-\bar{V}_R$, $\bar{V}_C$, for three different receptor numbers ($N_R = 375, 1500, 6000$), where $K_d = 6nM$ and the gradient is centered around $C_0 = K_d$ and ranges from $0.66\,K_d$ to $1.33\,K_d$ (i.e., $b/L = 0.33$). The dashed curve indicates the angular error for an estimate chosen uniformly at random over the cell. (**C**) Noise-to-signal ratio (NSR) comparing active-receptor sensing (yellow), inactive-receptor sensing (gray), and the combined-receptor estimate (blue) to the theoretical lower bound (dark green). Higher receptor counts reduce NSR, with the combined-receptor approach consistently outperforming either active- or inactive-only sensing. (**D**) Cumulative distribution functions (CDFs) of the angular error for the estimates $\bar{V}_{RL}$, $-\bar{V}_R$, $\bar{V}_C$ at fixed $N_R = 375$ but varying the gradient's offset relative to K_d: $C_0/K_d \in \{1/3, 1, 3\}$. As the mean concentration shifts, active and inactive receptors each capture different portions of the ligand profile, and combining their information again improves direction inference.

FIG. 2. **Ratiometric gradient sensing outperforms classical gradient sensing across a range of receptor numbers** N_R. (**A**) Schematic of reactions that constitute classical sensing. (**B**) Schematic of reactions that constitute ratiometric sensing. (**C**) 5 sample traces each of direction derived from G protein distribution over time for ratiometric and classical sensing strategies. $N_R = 375$, $N_G = 2500$, $D_G = 0.002 \mu m^2 s^{-1}$, $k_{on} = 0.0167 s^{-1} nM^{-1}$, $k_{off} = 0.1 s^{-1}$, $K_d = 6 nM$. (**D**) Cumulative distribution functions of angular error comparing gradient sensing between ratiometric ($\bar{V}_G^r$) and classical ($\bar{V}_G^c$) scenarios. $N_R = [375, 3000, 6000]$ using the parameters in (C). (**E**) Noise to signal ratio of direction derived from G protein distribution compared between classical and ratiometric sensing strategies. $N_R = [375, 1500, 3000, 6000]$ using the parameters in (C). For classical sensing, the G protein off-rate was assigned according to $N_R \mapsto k_{go}$ as $[375, 1500, 3000, 6000] \mapsto [0.0083, 0.0357, 0.0769, 0.1687] s^{-1}$.

FIG. 3. **Ratiometric sensing allows G proteins to outperform instantaneous theoretical limit.** (**A**) Temporal tracks of V_C and V_G^r for $N_R = 6000$, $N_G = 2500$, $D_G = 0.002 \mu m^2 s^{-1}$, $k_{on} = 0.0167 s^{-1} nM^{-1}$, $k_{off} = 0.1 s^{-1}$, $K_d = 6nM$. (**B**) Temporal tracks of V_C and V_G^r for $N_R = 375$ using parameters in (A). (**C**) Cumulative distribution functions of angular error comparing gradient sensing efficiency of active G protein in the ratiometric scenario to combined receptor. $N_R = [375, 3000, 6000]$ using parameters in (A). (**D**) Comparison of noise-to-signal ratio with theoretical Cramér-Rao bound which applies to the estimate V_C. For low values of N_R, the noise-to-signal ratio for V_G^r is below this theoretical bound, suggesting that V_G^r utilizes more information than is present in receptor states at a fixed time. $N_R = [188, 375, 750, 1500, 3000, 6000]$ using parameters in (A)

FIG. 4. **Ratiometric sensing enables G proteins to robustly generate a local average of past receptor states.** (**A**) Noise-to-signal ratio for V_G^r, V_C, and the theoretical lower bound as a function of the number of G proteins. $N_G = [181, 625, 1250, 2500, 5000]$, $N_R = 375$, $D_G = 0.002\mu m^2 s^{-1}$, $k_{on} = 0.0167 s^{-1} nM^{-1}$, $k_{off} = 0.1 s^{-1}$, $K_d = 6nM$. (**B**) Cartoon of a single receptor surrounded by G proteins that have recently inherited the receptor's state. For ratiometric signaling, G proteins can inherit state from ON or OFF receptors. For classical signaling, G proteins inherit state from ON receptors, but switch off at basal rate k_{gi}.(**C**) Gradient sensing accuracy (NSR) as a function of receptor switching rate (left: $k_{off} = [0.005, 0.01, 0.05, 0.1]\ s^{-1}$) and G protein diffusion constant (right: $D_G = [0.001, 0.002, 0.004, 0.008]\ \mu m^2 s^{-1}$). $N_G = 2500$, $N_R = 375$. (**D**) Left: Temporal traces of the active G protein fraction for ratiometric and classical signaling at three positions along a gradient corresponding to different mean ligand concentrations: back ($C = 1\ nM$, $p_R = 0.14$), middle ($C = 6\ nM$, $p_R = 0.5$), and front ($C = 18\ nM$, $p_R = 0.75$). The classical model is tuned to match ratiometric at $p_R = 0.5$. Right: Histograms of the active G protein fraction showing that the classical model deviates from p_R at the front and back, while the ratiometric model remains unbiased across all three regions. (**E**) Sweeping through different values of k_{gi} in the classical model shows optimal performance that does not cross instantaneous theoretical limits. $N_G = 2500$, $N_R = 375$. $k_{gi} = [0.0032, 0.0064, 0.0080, 0.0128, 0.0512, 0.1024]$

FIG. 5. **Adaptation to changing gradients.** Default simulation parameters: $N_R = 375$, $N_G = 2500$, $k_{off} = 0.1$ s^{-1}, $K_d = 6$ nM, and $D_G = 0.002$ μm^{2}s^{-1}. (**A**) Cartoon demonstrating simulation where the true gradient direction (black arrow) is flipped by 180°. The G protein direction vector V_G (green arrow) eventually re-orients to match the new gradient. (**B**) Cartoon demonstrating how adaptation time of gradient estimate by G proteins is calculated. Top: The environmental gradient direction is switched at $t = 300$ s from 0° to 180°. Bottom: The direction of the averaged trace derived from 1000 G protein vectors (V_G^r) lags before reaching the new steady state. The time to adapt is defined as the interval required to cross 90° toward the new directional equilibrium. (**C–E**) Parameter sensitivity of adaptation speed and sensing accuracy. Blue curves (left y-axis) indicate time to adapt; green curves (right y-axis) indicate the steady-state noise-to-signal ratio (NSR). (**C**) Sweeping receptor off-rate (k_{off}) from 0.005 to 0.1 s^{-1} shows that faster receptor kinetics improve both adaptation speed and directional accuracy. (**D**) Sweeping the number of receptors (N_R) from 375 to 6000 shows that higher receptor density significantly reduces the time to adapt while maintaining a relatively robust NSR. (**E**) Sweeping G protein diffusion (D_G) from 10^{-3} to 10^{-2} μm^{2}s^{-1} shows that faster diffusion accelerates adaptation but leads to a higher NSR, highlighting a fundamental trade-off between response speed and spatial precision.

FIG. 6. Replicating results in comparison of receptor signals using Smoldyn. **A)** Angular distribution of receptors with the same pheromone gradient centered around K_d but under different receptor number. Corresponds to Fig. 1B. $n = 10$ cells. **B)** Angular distribution of receptors with the same receptor number but under different heights of pheromone gradients. Corresponds to Fig. 1D. $n = 50$ cells.

FIG. 7. Replicating results in comparison between ratiometric and classical active G protein signal and combined receptor signal using Smoldyn. **A)** Angular distributions comparing gradient sensing between ratiometric and classical active G proteins. Corresponds to Fig. 2D. $n = 10$ cells. **B)** Angular distributions comparing gradient sensing between ratiometric active G proteins and combined receptors. Corresponds to Fig. 3C. $n = 10$ cells.

FIG. 8. The figure illustrates the cone $\mathcal{C}_{\delta,\theta}$. The ball centered at μ has radius $r = |\mu|\sin(\delta)$. The condition (B17) guarantees that the standard deviation of V is less than $r/\sqrt{2}$, so that V lies in the ball with probability at least $1/2$.

FIG. 9. This figure illustrates how G proteins record the historical state of a receptor. The states of G proteins G_1, G_2, and G_3 at time t were obtained through interaction with the same receptor, but at different earlier times. The protein G_1 records the receptor state from time $t_1 < t$; we say at that time t the age of G_1 is $t - t_1$. Protein G_2 records the receptor state from time $t_2 < t$; at time t the age of G_2 is $t - t_2$. Even though each protein last interacted with the same receptor, the states they record at time t may differ: G_1 and G_3 are inactive, while G_2 is active.

FIG. 10. This figure illustrates the most recent receptor-interactions among a collection of five G proteins. The states of G proteins G_1 and G_2 at time t were obtained through interaction with the same receptor R_1. Proteins G_1 and G_2 are linked to receptor R_1. However, proteins G_3, G_4, G_5 are linked to receptor R_2, since their most recent receptor interaction was with receptor R_2.

FIG. 11. Empirical joint distribution of $(R_f^+/N_R, G_f^+/N_G)$ and $(R_b^+/N_R, G_b^+/N_G)$, the active receptor and active G protein fractions at front and back of cell. Front of cell: ratiometric (yellow) and classical (green). Back of cell: ratiometric (blue) and classical (dark green). $N_R = N_G = 1000$. $\epsilon = 0.1$. The red line is the (linear) activation function $\ell(r) = r$, which applies to the ratiometric case. The blue and black lines are the activation functions $r/(r + 1/2)$ and its linearization $\frac{1}{2} + \frac{1}{2}(r - \frac{1}{2})$, which applies to the classical model (see equation (E2)).

FIG. 12. Distributions (histograms) of G_f^+/N_G and G_b^+/N_G, the active G protein fraction at the front and back of cell. Front of cell: ratiometric (yellow) and classical (green). Back of cell: ratiometric (blue) and classical (dark green). These correspond to marginal distributions from the joint distribution plot above, Figure 11. $N_R = N_G = 1000$. $\epsilon = 0.1$

FIG. 13. Distributions (histograms) of $(G_f^+ - G_b^+)/N_G$, the active G protein gradient from front to back of cell, for Ratiometric (blue) and Classical (yellow).

FIG. 14. **Blue:** Empirical distribution of N, the number of distinct receptor encounters by a G protein since its most recent state-reset time. $N_R = 375$, $D_G = 0.002(\mu m)^2/sec$, $k_{go} = 0.0083(sec)^{-1}$. The mean number of encounters is $\mathbb{E}[N] = 1.28$. Fraction of proteins not encountering a receptor: 0.42. Mean time to first encounter: 79.9. 20,000 proteins were simulated, for a single realization of the receptor configuration. **Yellow:** For comparison, the Geometric distribution given by (F1).

FIG. 15. Graph of p_G^c versus p_r, for different values of $\mathbb{E}[N] = 2^k - 1$, $k = 0, \ldots, 4$. For comparison, the straight line has slope 1.