Genomic Patterns of Parallel Divergence Across Demographically Heterogeneous

Stickleback Populations in Eastern Canada

*Alan García-Elfring, ¶Antoine Paccard, and *Rowan D. H. Barrett

*Department of Biology, McGill University, Montreal, QC, Canada

¶McGill University Genome Center, McGill University, Montreal, QC, Canada

Corresponding authors: rowan.barrett@mcgill.ca, alan.garcia-elfring@mail.mcgill.ca

Alan García-Elfring ORCID 0000-0002-2093-4041

Antoine Paccard ORCID 0000-0003-0738-5540

Rowan D.H. Barrett ORCID 0000-0003-3044-2531

Significance statement

Threespine stickleback are a model system for studying parallel evolution, yet most genomic research has focused on Pacific and European populations. By examining previously understudied Atlantic Canadian populations, we identify genomic patterns consistent with parallel divergence across markedly heterogeneous demographic contexts, including populations shaped by strong drift as well as gene flow. We detect differentiation near dopamine receptor genes, pointing to a potential role for behavioural and hormonal pathways. Together, these results highlight the role of demographic context in shaping patterns of genomic differentiation associated with freshwater colonization.

Abstract

The threespine stickleback (*Gasterosteus aculeatus*) is a key model in evolutionary

genetics, particularly for studies of parallel evolution, yet most genomic insights derive

from populations on the west coast of North America and in Europe. Here, we use

restriction site-associated DNA sequencing (RAD-seq) of pooled samples to examine

genomic differentiation between marine and freshwater stickleback populations from

Atlantic Canada. Our analyses reveal substantial heterogeneity in the extent and

genomic distribution of marine–freshwater differentiation, with some freshwater

populations showing strong divergence consistent with long-term isolation and drift, and

others exhibiting patterns consistent with ongoing gene flow and admixture. Despite this

demographic variation, we identify genomic regions that are repeatedly differentiated

between marine and freshwater habitats, including loci near dopamine receptor genes

(*Drd4a* and *Drd2l*). Gene ontology analyses of candidate regions show enrichment for

functions related to nervous system development and dopamine receptor activity.

Together, these results indicate that freshwater-associated genomic differentiation in

Atlantic Canadian stickleback occurs across contrasting demographic contexts and

suggest a potential role for neurological and behavioural pathways in adaptation to

freshwater environments.

Introduction

The genetic basis of adaptation underpins the mechanisms of species diversification. A classic system for exploring these processes is the threespine stickleback (*Gasterosteus aculeatus*), native to the oceans of the Northern Hemisphere and an important species in ecological and evolutionary studies over the last century (Bertin 1925; Heuts 1947a, b; Moodie and Reimchen 1976; Nagel and Schluter 1998; Marques et al. 2018; Sanderson et al. 2026). Following the last glacial maximum (~21 kya), retreating glaciers created new freshwater lakes and rivers into which marine stickleback populations dispersed and became isolated. Many of the nascent freshwater populations evolved similar low-plated morph phenotypes, particularly in western North America (Baumgartner and Bell 1984; Colosimo et al. 2005; Hohenlohe et al. 2010). Alleles that confer higher fitness in freshwater environments often persist at low frequencies in marine populations. Bidirectional gene flow is thought to maintain this standing genetic variation, facilitating the repeated colonization of freshwater habitats and the independent evolution of similar phenotypes (the "transporter hypothesis"; Schluter and Conte 2009). As a result, the threespine stickleback is widely regarded as a model system for studying parallel evolution.

However, even in the early stages of establishing stickleback as a model species, researchers noted that the striking parallel evolution observed among freshwater populations might be a geographically restricted phenomenon (Hagen and Moodie 1982), largely confined to western North America (eastern Pacific, e.g., Colosimo et al. 2005) and parts of Europe (e.g., Klepaker 1995). For instance, considering the armour phenotype, in several regions such as eastern North America, Northern Europe, the

Baltic Sea, and eastern Asia, the completely plated morph—typical of marine

environments—also predominates in local freshwater lakes (Münzing 1963; Penczak

1965; Hagen and Moodie 1982; Mäkinen et al. 2008; Raeymaekers et al. 2014;

Ferchaud and Hansen 2016; Yamasaki et al. 2019). This limited phenotypic divergence

between marine and freshwater populations has also been observed at the genetic level

using allozymes (Rafiński et al. 1989). A large-scale genomic study by Fang et al.

(2020a) confirmed that high levels of marine–freshwater (M-FW) genomic differentiation

are primarily restricted to the eastern Pacific, whereas Atlantic populations show

relatively low differentiation. In contrast to the strong divergence seen in British

Columbia populations, many freshwater populations in Eastern Canada are dominated

by the fully plated ecotype (Hagen and Moodie 1982; Haines 2023), although some

exhibit distinct morphological variation (Scott et al. 2023). Despite these observations,

genomic studies in this region remain limited (but see Fang et al. 2020a).

One potential explanation for the reduced extent of parallel evolution and M-FW

genomic differentiation outside the eastern Pacific is the stochastic loss of freshwater-

adapted alleles. Threespine stickleback originated in the Pacific Ocean, where they

have persisted for approximately 26 million years (Matschiner et al. 2011; Betancur-R et

al. 2015). It was only much more recently, during the late Pleistocene (36.9–346.5

thousand years ago), that they expanded their range into the western Pacific and the

Atlantic basin (Orti et al. 1994; Fang et al. 2018; Fang et al. 2020b). As a result of this

range expansion, Atlantic populations exhibit lower genetic diversity compared to their

Pacific counterparts (Fang et al. 2020a). Notably, most Atlantic populations examined to

date have been from Europe, leaving other regions, such as Atlantic Canada, relatively understudied.

Here, we investigate marine and freshwater threespine stickleback from eastern Canada, specifically from the provinces of Nova Scotia and Newfoundland, where fully plated phenotypes have been observed in freshwater habitats (Hagen and Moodie 1982; Haines 2023). Using genome-wide SNP data, we characterize M-FW genomic differentiation and test for signatures of parallel natural selection. Our goal is to gain insight into the evolutionary history and genetic basis of freshwater adaptation in stickleback populations from eastern North America.

Results

Genomic diversity, population structure, and differentiation among populations

We sampled 30 adult threespine stickleback from each of nine populations along the eastern Canadian Atlantic coast (N = 270 total), including four marine and five freshwater sites in Nova Scotia and Newfoundland (Figure 1; Table S1). Genome-wide estimates of nucleotide diversity (π), Watterson's θ, and Tajima's D varied among populations and habitats (Figure 2; Table S2). Marine populations like Antigonish Landing consistently showed high levels of genetic diversity ($\pi \approx 0.31$–0.33) and positive Tajima's D values, consistent with substantial standing variation. Freshwater populations exhibited greater heterogeneity. Blue Pond showed the lowest π and θ values and a near-zero to slightly negative Tajima's D, indicating low genetic diversity consistent with strong drift/founder effects, followed by Lake Ainslie and Black River

(Table S2). In contrast, Pomquet Lake and Pinchgut Lake showed among the highest π

and Tajima's D values observed, indicating an excess of variants at intermediate

frequencies and potential gene flow.

To gain insight into population structure, we performed a principal component

analysis (PCA) on allele frequencies (Figure 2, PCA). Marine and freshwater

populations segregated into three main groups along PC1 (14.2% of variance

explained) and PC2 (13.1%). Freshwater populations characterized by lower genetic

diversity and lower Tajima's D—Blue Pond, Lake Ainslie, and Black River—clustered

together. In contrast, Pomquet Lake and Pinchgut Lake, freshwater populations with

higher genetic diversity and Tajima's D (putative gene flow from the ocean), clustered

together with several marine populations (Porper Pond, Cook's Brook, and Humber

Arm), which extend further along PC1. The marine population Antigonish Landing

formed a distinct cluster.

Hierarchical clustering using the WPGMA method grouped populations into clades

that overall corroborate the PCA (Figure 2 phylogram): Lake Ainslie and Black River

formed one sister group, with Blue Pond appearing as a more genetically divergent

freshwater population. Pomquet Lake and Pinchgut Lake cluster together in a clade

nested among the marine populations with which they overlap in the PCA, showing low

bootstrap support at internal branches, suggesting weak genetic structure consistent

with high connectivity and ongoing gene flow. To this group, Antigonish Landing forms a

sister group with strong bootstrap support (100%), indicating genetic differentiation from

the putatively admixed freshwater and marine clade.

We next examined population structure by visualizing median genome-wide F_{ST}

across all 36 pairwise comparisons based on 18,582 SNPs as a heatmap (Figure 3;

Table S3). Relative divergence varied widely, ranging from 0.0118 to 0.0824. Substantial

genomic heterogeneity was observed among both marine–freshwater (M–FW) and

freshwater–freshwater (FW–FW) comparisons (Figure 4; Figures S1, S2). Notably, both

the lowest and highest levels of genomic differentiation occurred among freshwater

populations: Pomquet Lake (Nova Scotia) and Pinchgut Lake (Newfoundland) were the

least differentiated, whereas Lake Ainslie (Nova Scotia) and Blue Pond (Newfoundland)

were the most differentiated.

Comparisons of relative (F_{ST}) and absolute (d_{XY}) divergence across 20 kb windows

revealed a decoupling between these two measures (Figure 5). Across key population

pairs (e.g., Figure 4), windows with high F_{ST} did not show correspondingly elevated d_{XY},

indicating that extreme allele frequency divergence is not driven by increased sequence

divergence between populations. Instead, high F_{ST} values occur in windows with low d_{XY},

a pattern expected in populations that have experienced genetic drift and possess lower

genetic diversity.

SNPs showing repeated divergence between marine and freshwater populations

Excluding comparisons between populations from the same habitat type (M-M and FW-

FW), we obtained 25,236 SNPs across the 20 possible M-FW comparisons within and

across provinces. We used this SNP set to conduct an exploratory analysis to identify

genomic regions showing repeated differentiation between marine and freshwater

populations, with the aim of generating candidate loci potentially involved in parallel

freshwater adaptation (Figure 6). We identified 2,100 SNPs as F_{ST} outliers in two or

more M-FW comparisons within Nova Scotia (19 SNPs across all six comparisons). In

Newfoundland, we identified 1,306 SNPs as F_{ST} outliers in two or more comparisons.

Comparing interprovincial differentiation among 10 M-FW population comparisons

revealed 3,422 SNPs as F_{ST} outliers in at least two comparisons, with three SNPs

appearing in nine of the 10 comparisons (Figure 6). In the top 5% of the F_{ST} distribution,

we identified 279 SNPs (referred to as 'F_{ST} outliers'), which were outliers in at least 6/20

M-FW comparisons, representing 1.1% of the SNP dataset.

We found that 180 of 279 F_{ST} outliers map within 5 kb of 133 genes (Table S4), with

68 genes containing at least one outlier (Table S5). Notably, two SNPs (T>C at VI:

16449327 and C>T at XX: 14009648) were in the top 5% of F_{ST} values in 19/20 M-FW

comparisons (6/6 in Nova Scotia, 4/4 in Newfoundland, and 9/10 interprovincial

comparisons; Figure 6). The latter SNP is located 920 bp downstream from a dopamine

receptor gene, *Drd2l* (XX: 14002115-14008728). It is worth noting that these markers

show lower coverage in some populations, which is known to bias levels of F_{ST}. (VI:

16449327 coverage range: 5-21X; XX:14009648 coverage range: 5-35X). However,

although both alleles (T and C) were present in marine populations at both loci

(VI:16449327 T:C = 11:39; XX:14009648 T:C = 33:63), the C allele was almost entirely

absent from freshwater populations. Across freshwater habitats, T:C coverage ratios

were 38:2 at the chromosome VI locus and 55:1 at the chromosome XX locus. Notably,

the only freshwater population in which the C ("marine") allele was detected was

Pomquet Lake.

To formally assess significance, we applied a Fisher's Exact Test (FET), which

identified 18 SNPs with significant allele frequency differences (FDR < 0.10) across the

10 marine–freshwater comparisons, each located within 5 kb of different genes (Table

S6). Three of these genes also overlap with FST outlier genes (*dmap1*, *hnrnpm*, and

drd2l). The armour-plating locus (IV:12811481) was not sampled in our RAD-seq

dataset, with the closest SNP mapping 72.3 Kb downstream (IV:12883820).

Gene Ontology

Genes within 5 kb of FST outliers were enriched for biological processes related to

adenylate cyclase-inhibiting dopamine receptor signalling and nervous system

development (Table S7). Enriched molecular functions included neurotrophin binding

and dopamine neurotransmitter receptor activity (Table S8).

Discussion

In this study, we investigated marine and freshwater populations of threespine

stickleback along Canada's east coast to gain insights into their evolutionary history and

the potential genetic basis of freshwater adaptation. Contrary to expectations of low

genome-wide differentiation between marine and freshwater populations in the western

Atlantic (Fang et al. 2020a), we found that differentiation among M-FW populations

varied widely along the east coast of North America. This suggests a complex

evolutionary history, likely involving multiple colonization events, distinct glacial refugia,

or ongoing gene flow. Evidence of parallel evolution was observed as repeated allele

frequency differences at SNP loci across several M-FW population comparisons.

Notably, we identified a consistent signal of differentiation at one SNP near a dopamine

receptor previously found to be differentially expressed between marine and freshwater

populations (Di-Poi et al. 2016). Our results suggest that, even in the likely absence of

key alleles for reduced armour in eastern Canadian populations, behavioural or

endocrine adaptations may have facilitated freshwater persistence.

Heterogeneous marine-freshwater differentiation in Eastern Canada

Current evidence suggests that threespine stickleback colonized the Atlantic during the

late Pleistocene, as recently as ~36.9 thousand years ago (Fang et al. 2018; Fang et al.

2020b). Repeated bottlenecks during this expansion may have led to the stochastic loss

of some freshwater-adaptive alleles, such as the low-plated Eda allele (Fang et al.

2020a). Loci under selection in freshwater are expected to differentiate rapidly following

colonization. Thus, the low M-FW differentiation observed in many Atlantic populations

may reflect a lack of such adaptive alleles in their standing genetic variation, compared

to Pacific populations.

Here, we show that while genome-wide M-FW differentiation is indeed low in some

eastern North American comparisons, other comparisons show levels of differentiation

comparable to those seen on the Pacific coast (Figure 4, Figure S2). Genome-wide π

and θ were consistently higher in marine populations ($\pi \approx 0.31$–0.33) than in freshwater

populations, which exhibited greater heterogeneity. Freshwater populations with low π

and θ, such as Blue Pond, Lake Ainslie, and Black River, showed near-zero or slightly

negative Tajima's D, consistent with reduced genetic diversity and neutral evolution, whereas Pomquet Lake and Pinchgut Lake showed higher π and positive Tajima's D, suggesting an excess of intermediate-frequency variants possibly due to ongoing gene flow. This inference was supported by PCA, where Pomquet Lake and Pinchgut Lake clustered with marine populations rather than with other freshwater populations, like Lake Ainslie and Black River, indicating likely gene flow between these freshwater populations and nearby marine sources.

The highest levels of differentiation were observed between different freshwater populations. For example, two lakes in Newfoundland—Pinchgut Lake and Blue Pond—are only 5 km apart but differ substantially in their genetic divergence from marine populations (Figure 4). Blue Pond is a small, rain-fed lake without input or output streams, making gene flow from neighbouring populations unlikely. Such barriers to gene flow likely promoted divergence through genetic drift. Comparisons of F_{ST} and absolute divergence (d_{XY}) revealed a decoupling of relative and absolute divergence (Figure 5), with high F_{ST} windows showing low d_{XY}, consistent with drift in low-diversity freshwater populations.

Two recent preprints provide important context for interpreting the structure of the marine populations included in our study. Samuk et al. (2025) and Sumarli et al. (2025) analyzed RADseq and whole-genome data, respectively, from several of the same Nova Scotia marine sites sampled here. They found two genetically distinct marine ecotypes — "white" and "common" sticklebacks — co-occur in sympatry across the region, with ongoing gene flow and very low but detectable genomic divergence (FST ≈ 0.01). The common marine stickleback in Nova Scotia could be separated into two genetic

clusters, "mainland" and "Bras d'Or". Despite this shallow divergence, the white ecotype exhibits consistent differences in morphology, reproductive traits, and behaviour, including smaller body size, distinctive coloration, spine length, altered reproductive investment, divergent nesting substrate and location, modified parental care strategies, and distinct courtship displays (Samuk et al. 2025). In contrast, no trophic or ecological differentiation between ecotypes was observed.

Two of our marine sampling sites were also studied by Samuk et al. (2025). In their investigation, Antigonish Landing (western Nova Scotia) consisted almost entirely of individuals from the "common mainland" genetic cluster, whereas Porper Pond (eastern Nova Scotia) was composed almost entirely of individuals belonging to the white ecotype genetic cluster. Fish with ancestry belonging to the common mainland, common Bras d'Or, and white ecotype predominate in western, northern, and eastern Nova Scotia, respectively. We did not observe any obvious phenotypic differentiation among our marine populations sampled in Nova Scotia and Newfoundland, most notably, none exhibited the distinct colouration of the white stickleback ecotype. Given the low divergence and recent estimated split time (~1,000 years), white and common sticklebacks share the vast majority of their genetic variation. Thus, even if a few undetected white sticklebacks individuals were included in our sample, their allele-frequency contribution would be unlikely to account for the localized, parallel allele-frequency shifts observed near dopamine receptor genes.

Our population structure analyses are concordant with the structure reported by Samuk et al. (2025). Principal component analysis (PCA) clearly shows Antigonish Landing and Porper Pond as distinct genetic clusters. Notably, Porper Pond clusters

closely with freshwater populations Pomquet Lake and Pinchgut Lake, consistent with

ongoing gene flow or admixture. One possible interpretation is that white sticklebacks

represent a hybrid population (Marques et al. 2019). It is known that freshwater

stickleback are smaller than marine (Haglund et al. 1990; Aguirre et al. 2022), can

display lighter colouration (Miller et al. 2007;), and show an array of different behaviours

(McPhail and Hay 1983; Di-Poi et al. 2014). However, neither Samuk et al. (2025) nor

Sumarli et al. (2025) included freshwater populations in their analyses, limiting direct

inference about the contribution of freshwater–marine connectivity to this structure.

Multiple colonization events in eastern Canada, similar to the patterns observed in

the Japanese Archipelago (Kakioka et al. 2020) and Europe (Mäkinen et al. 2008) could

have led to heterogeneity in genetic distances among freshwater populations. However,

if freshwater populations had originated from separate sources or refugia (e.g., Mäkinen

and Merilä 2008), they would have accumulated sequence divergence over time,

resulting in high d_{XY}. Instead, the low d_{XY} coupled with high F_{ST} indicates recent

population divergence with minimal absolute sequence differentiation, consistent with

colonization from a single or shared marine source followed by rapid drift-driven

divergence in isolated freshwater lakes. Pinchgut Lake is connected to other lakes

through Georges Lake, with a stream that extends to the ocean, while Pomquet Lake is

a few kilometres (<4 km) from the ocean and likely experiences ongoing gene flow via

Taylor Creek. In contrast, Blue Pond is a small, rain-fed lake without streams. This

geographic variation in connectivity explains the heterogeneous population structure

better than multiple independent colonization events from divergent sources.

Repeated differentiation near dopamine receptor genes

Adaptation to similar environments via re-use of standing genetic variation is considered a primary mechanism underlying parallel evolution (Schluter and Conte 2009; Schlötterer 2023). The low-plated Eda allele, though present at low frequencies in marine populations, confers a fitness advantage in freshwater (Marchinko and Schluter 2007; Barrett et al. 2008; Le Rouzic et al. 2011). Its widespread loss in the Atlantic gene pool may explain the absence of parallel evolution for the low-plated phenotype in eastern Canadian lakes. However, *Eda* is not the only gene involved in freshwater adaptation; genes involved in osmoregulation, such as Na^+/K^+-ATPase and $Na^+/K^+/2Cl^-$ cotransporters, have been repeatedly implicated across global stickleback populations (Deagle et al. 2013; Divino et al. 2016; McCairns and Bernatchez 2010; Garcia-Elfring et al. 2021).

Consistent with previous observations and the predicted loss of the freshwater Eda allele, we did not find signs of molecular evolution near the Eda locus in our populations (Hagen and Moodie 1982; Fang et al. 2020a), although our nearest marker was more than 70 Kb away. Despite overall low differentiation, we identified signals of parallel evolution at 1.1% of SNPs, which showed repeated marine–freshwater divergence both within and across Nova Scotia and Newfoundland. Notably, this included SNPs located near dopamine receptor genes *Drd4a* and *Drd2l*, and candidate genes were enriched for functions related to dopamine receptor activity and nervous system development. Strikingly, the 'C' alleles of two parallel outlier SNPs were almost entirely missing in freshwater populations but remained polymorphic in marine populations, consistent with selection favouring one allele, acting against the other, or both. We provide primers for

each locus that were designed *in silico* (Untergasser et al. 2012); these primers have

not yet been experimentally tested (Table S9).

While individual variation in endocrine function remains less studied than other

physiological traits in evolutionary biology, it holds strong potential for linking genetics,

physiology, and ecology (Williams 2008). In zebrafish (*Danio rerio*), expression levels of

dopamine receptors *Drd2c* and *Drd3* have been associated with individual aggression,

particularly in dominant versus subordinate males (Filby et al. 2010). In stickleback,

exposure to predators has been shown to reduce *Drd4a* expression (Abbey-Lee et al.

2018), and marine and freshwater populations exhibit marked differences in behaviours

such as sociability, aggression, and activity (Di-Poi et al. 2014). At the population level,

Di-Poi et al (2016) showed evolutionary changes in physiological regulatory networks

often target receptor genes—such as *Drd2*—rather than their ligands or

neurotransmitters. In a separate study using stickleback from a Nova Scotia lineage

(Blouw and Hagen 1990), *Drd2* expression was associated with male behavioural trade-

offs between courtship and territorial defence (Barbasch et al. 2023).

The dopaminergic system also plays a key role in osmoregulation through inhibition

of prolactin, a hormone critical for ion retention and water balance in freshwater

environments (Liu et al. 2006; Yamamoto and Vernier 2011; Mancera and McCormick

2019). Prolactin mRNA levels correlate with local ion concentrations in natural

populations of black-chinned tilapia (*Sarotherodon melanotheron*) adapted to different

salinities (Tine et al. 2007). In the euryhaline fish *Scatophagus argus*, renal

dopaminergic signalling via *Drd1* inhibits Na^+/K^+-ATPase under hypo-osmotic stress (Su

et al. 2016). In stickleback, prolactin is associated with freshwater migration and salinity

tolerance (Lam and Hoar 1967; Ishikawa et al. 2016; Pavlova et al. 2020; Taugbøl et al.

2022). Taken together with the population genetic patterns described above, these

results highlight dopaminergic signalling as a plausible axis of freshwater adaptation in

eastern Canadian stickleback. However, because our dataset is pooled RAD-seq and

we cannot estimate local LD decay, SNP-to-gene proximity should be treated as

suggestive, and these dopamine receptor loci are best viewed as candidates motivating

follow-up work. Future studies combining whole-genome data with functional assays

and expression profiling will be required to test whether variation at specific loci

contributes to adaptation in these systems.

Conclusion

Our study revealed heterogeneous genomic differentiation between marine and

freshwater threespine stickleback populations in eastern Canada. Some lakes, in

particular Blue Pond, appear to represent drifted "island" populations with limited gene

flow, whereas others, like Pomquet Lake, maintain higher diversity through ongoing

connectivity with marine populations. Despite this difference in demographic history, we

detected evidence of parallel evolution near dopamine receptor genes, where specific

alleles are largely absent in freshwater but polymorphic in marine populations. Our

results suggest that parallel evolution in eastern Canada may involve selection on

behavioural (or osmoregulatory) phenotypes, shaped by both drift in isolated

populations and on-going selection in connected populations, though further work is

needed to test this hypothesis.

Methods

Field sampling and DNA extraction

We collected 30 adult threespine stickleback (> 30 mm standard length) from nine

populations in eastern (Atlantic) Canada, comprising four marine and five freshwater

populations. Fish were sampled using minnow traps and beach seines during June and

July 2014 in Nova Scotia and Newfoundland, Canada (Figure 1; Table S1). These

samples form the basis of the RAD-seq dataset analyzed in this study.

In Nova Scotia, freshwater populations were sampled from two lakes (Pomquet Lake

and Lake Ainslie) and one stream (Black River), while marine samples were obtained

from Antigonish Landing and Porper Pond. In Newfoundland, freshwater populations

were sampled from Pinchgut Lake and Blue Pond, and marine populations from Cooks

Brook and Humber Arm (Figure 1; Table S1). Genomic DNA was extracted from all

individuals using a standard phenol–chloroform protocol.

DNA sequencing of pooled samples

DNA concentration was quantified using a Picogreen® ds DNA assay (Thermo Fisher

Scientific, Waltham, USA) and an Infinite® 200 Nanoquant (Tecan Group Ltd.

Männedorf, Switzerland). DNA concentrations were normalized across individuals and

pooled by sampling location, resulting in nine pooled samples, each consisting of 30

individuals.

RAD-seq libraries were prepared following the protocols of Paccard et al. (2020).

Briefly, pooled samples were digested with NlaIII and MluCI, ligated to unique barcoded

adapters, size-selected for ~400 bp fragments, enriched for biotin-tagged fragments

using streptavidin beads, and PCR-amplified. Cleanup steps were performed using

AMPure XP beads. Libraries were sent for sequencing at the McGill University and

Genome Quebec Innovation Center (Montreal, Canada) on one lane of Illumina

HiSeq2500 platform, generating 125bp paired-end reads.

Read processing and variant mapping

RAD-seq data was demultiplexed into pool-specific read sets using the

process_radtags module of Stacks (Catchen et al. 2013), based on the barcode

sequences listed in Table S1. We used the *trim-fastq.pl* script of *Popoolation* (Kofler et

al. 2011a) to process raw reads, filtering based on read quality (--quality-threshold 20)

and length (--min-length 50). We then mapped the processed reads to the stickleback

reference genome (BROAD S1) with the program *Bowtie2* (Langmead and Salzberg

2012), using the --end-to-end mapping option. We used *SAMtools* (Li et al. 2009) to

convert the output SAM files to BAM format and subsequently removed reads with

mapping quality below 20 (samtools view -q 20). We generated an mpileup file

(samtools mpileup -B) and converted the mpileup file to the synchronized (sync) format

using *Popoolation2* (Kofler et al. 2011b), which was used for downstream analyses.

Population genetic parameters, population structure, and allele frequency differentiation

Population genetic summary statistics

We estimated population genetic parameters, including nucleotide diversity (π), Watterson's θ, and Tajima's D, using the *variance-sliding.pl* script from PoPoolation2 (Kofler et al. 2011b).

Allele frequency estimation and PCA

To visualize population structure, we performed a principal component analysis (PCA) based on SNP-specific allele frequencies at loci shared across populations, which primarily reflect shared (ancestral) standing variation rather than population-specific rare variants. For each population, allele frequencies were obtained from cleaned and sorted BAM files which were processed using bcftools to generate downstream VCFs. Genome-wide variant calling was performed with bcftools mpileup and bcftools call. From these VCFs, sites were subset to a predefined panel of 18,582 SNP positions shared across populations (datasets defined below).

Allele depths (AD) for reference and alternate alleles were obtained at each SNP using bcftools query, and allele frequencies were computed as the fraction of reads supporting the alternate allele over the total read depth. Population-specific allele frequency files were then combined into a single matrix with SNPs as rows and populations as columns. Before performing principal component analysis (PCA), missing allele frequencies—caused when reads were present but alternate alleles were not confidently called by bcftools in the VCF AD field for certain SNP–population

combinations—were imputed using the mean allele frequency for that SNP across

populations. Unlike bcftools, Popoolation2 does not require confident genotype calls,

only read counts from the sync file. SNPs with zero variance across populations were

excluded. PCA was performed in R using *prcomp* on the centered and scaled allele

frequency matrix, treating populations as observations. Population clustering was

visualized using the first two principal components, with populations labeled and

coloured according to habitat type (marine vs. freshwater).

We estimated relative genetic differentiation among populations using F_{ST}, as defined

by Hartl and Clark (1997), with the *fst-sliding.pl* script from PoPoolation2 (Kofler et al.

2011b). We visualized genetic relationships among marine and freshwater populations

by constructing a WPGMA phylogram based on median of these F_{ST} values. A distance

matrix was generated from genome-wide SNP data, and clustering was performed in R

using the hclust(method = "mcquitty") function. The resulting dendrogram was

visualized as an unrooted phylogram using phangorn::plotBS(), with branch lengths

scaled to genetic distance. Bootstrap support values were calculated from 10,000

replicates, and only values ≥10% are shown. The tree was explicitly unrooted to avoid

assumptions about ancestral directionality.

Low sequencing coverage can bias F_{ST} estimates through sampling variance in allele

frequencies. We therefore additionally assessed allele frequency differences using

Fisher's Exact Test (FET) implemented in *fisher-test.pl of* Popoolation2*.* Analyses were

performed using the following parameters: --min-count 2, --min-coverage 5, --max-

coverage 500, --min-covered-fraction 0, --window-size 1, --step-size 1, --pool-size

30:30:30:30:30:30:30:30:30, and --suppress-noninformative. Only genomic regions

assembled at the chromosome level were analyzed; unplaced scaffolds were excluded.

Also, because isolated populations experiencing strong genetic drift can exhibit elevated

F_{ST} values as a consequence of reduced within-population diversity, we calculated

absolute sequence divergence (d_{XY}). Estimates of d_{XY}, which is equivalent to between-

population nucleotide diversity, were obtained using grenedalf (Czech et al. 2024), a

framework designed, like Popoolation2, for Pool-seq data.

We next sought to identify variants putatively associated with parallel selection. For

each marine–freshwater (M–FW) population comparison, we identified SNPs in the

upper 5% of the F_{ST} distribution (e.g. Yan et al. 2025). This threshold was applied

heuristically to retain a broad range of allele frequency differences across population

pairs with heterogeneous demographic histories.To identify loci with consistent

differentiation (i.e., putative parallelism), we defined three sets of SNPs in the top 5%

across at least two M-FW comparisons: within Nova Scotia, within Newfoundland, and

across both provinces. SNPs shared among all three sets were designated as "F_{ST}

outliers."

To complement the F_{ST} approach, we performed a genome-wide FET scan across 10

intraprovincial M-FW comparisons to identify SNPs showing consistent allele frequency

shifts. To control for multiple testing, raw p-values were converted to q-values using the

qvalue package (Storey et al. 2003), and SNPs with a false discovery rate (FDR) below

10% were retained. This FDR threshold reflects the expected proportion of false

positives among significant results.

Identification of candidate genes and analysis of molecular function

Outlier SNPs are expected to be located within genes potentially under parallel

selection or, more commonly, be located near them, given that we used a reduced

representation approach. To identify such genes, we applied a custom bash script to

map outlier loci to or near protein-coding genes in the reference genome. Our search

was limited to 14,252 protein-coding gene annotations, filtered by the attributes

"ID=gene" and "biotype=protein_coding." Given that our data represent a reduced

portion of the genome, the causal mutation is unlikely to be directly sampled and is

more likely in linkage disequilibrium (LD) with a nearby SNP. Because our dataset is

pooled RAD-seq, we did not estimate local LD decay. To account for this, we examined

genes either associated with outlier loci or located within a heuristically chosen window

of 5 kb around them (e.g., Nichols et al. 2016; Chen et al. 2018). To investigate the traits

putatively under selection, we analyzed the candidate genes for enrichment of

molecular functions using ShinyGO 0.77 (Ge et al. 2020), which links gene lists to

functional categories, such as gene ontology (GO), based on annotation databases

from Ensembl and STRING-db. We assessed enrichment for GO terms in the

categories of biological process and molecular function, using a false discovery rate

(FDR) threshold of 10% and reporting the 20 most significant results. We compared our

list of candidate genes to a background set of 14,252 genes from which outliers were

sampled, using the 'best matching species' option for further analysis.

SNP and windowed datasets

Our analyses used two SNP datasets and one dataset based on 20 kb windows. The

first SNP dataset included 18,582 variants (average minimum coverage across

populations: 56.4X) shared across all nine populations and present in all 36 pairwise

population comparisons, including within- and between-habitat types (M-M, FW-FW, and

M-FW). This dataset was used to estimate population genetic parameters (π, θ, D),

calculate F_{ST} across all 36 comparisons (e.g., Manhattan and heat plots), perform PCA,

and conduct FET scans. For analyses of marine–freshwater parallelism, which excluded

M-M and FW-FW comparisons, we used a second dataset of 25,236 SNPs (average

minimum coverage among populations: 54.5X) informative in the 20 relevant pairwise

comparisons.

When examining absolute divergence, we employed a 20 kb windowed approach,

calculating F_{ST} with Popoolation2 and d_{XY} with grenedalf. We used Popoolation2 to set

the parameters. Because reduced-representation data results in many 20 Kb windows

with zero coverage, we retained informative windows by setting the --min-covered-

fraction parameter (the minimum fraction of a window with coverage between 5-200X

across all populations) to 0.0001, based on the median and mean distances between

variants in the 18,582-SNP dataset (Mean distance: 21,417.65 bp; Median distance:

3,977.00 bp; Std deviation: 40,189.06 bp; Max distance: 1,385,936 bp). Only windows

containing at least three SNPs were included in the analysis. The 0.0001 threshold

ensured that all 20 Kb windows have at least some SNPs retained, while the ≥3-SNP

filter adds statistical robustness by averaging each window's estimate over multiple data

points. This filtering resulted in a total of 7,160 F_{ST} windows from Popoolation2 (average

minimum coverage at each SNP in the window: 24.5X). We subsetted the same 7,160

windows from grenedalf to directly compare F_{ST} to d_{XY} estimates.

Acknowledgments

This work was supported by a Discovery Grant from the Natural Sciences and

Engineering Research Council of Canada (NSERC) awarded to R.B., as well as funding

from the Canada Research Chairs (CRC) program. We thank Mathilde Salamon for her

assistance and valuable contributions throughout the project as well as Dieta Hanson

and the Mullaney family for their assistance with field sampling.

Data availability

All raw sequencing data generated in this study have been deposited in the NCBI

Sequence Read Archive (SRA) under a BioProject accession PRJNA1406969.

References

Abbey-Lee, R.N., Uhrig, E.J., Zidar, J., Favati, A., Almberg, J., Dahlbom, J., Winberg,

S. and Løvlie, H., 2018. The influence of rearing on behavior, brain monoamines,

and gene expression in three-spined sticklebacks. Brain, behavior and evolution,

91(4), pp.201-213.

Aguirre, W.E., Reid, K., Rivera, J., Heins, D.C., Veeramah, K.R. and Bell, M.A., 2022.

Freshwater colonization, adaptation, and genomic divergence in threespine

stickleback. Integrative and Comparative Biology, 62(2), pp.388-405.

Barbasch, T.A., Behrens, C., McLain, M., Arredondo, E. and Bell, A.M., 2023. A

distinct neurogenomic response to a trade-off between social challenge and

opportunity in male sticklebacks (Gasterosteus aculeatus). Biology Letters, 19(11),

p.20230253.

Baumgartner, J.V. and Bell, M.A., 1984. Lateral plate morph variation in California

populations of the threespine stickleback, Gasterosteus aculeatus. Evolution,

pp.665-674.

Barrett, R.D., Rogers, S.M. and Schluter, D., 2008. Natural selection on a major

armor gene in threespine stickleback. Science, 322(5899), pp.255-257.

Bertin, L., 1925. Recherches bionomiques, biométriques et systématiques sur les

épinoches gastérostéidés. Am. Inst. Oceanograph. N. S. Z. Frac. 1: 1-204.

Betancur-R, R., Ortí, G. and Pyron, R.A., 2015. Fossil-based comparative analyses

reveal ancient marine ancestry erased by extinction in ray-finned fishes. Ecology

Letters, 18(5), pp.441-450.

Blouw, D.M. and Hagen, D.W., 1990. Breeding ecology and evidence of reproductive

isolation of a widespread stickleback fish (Gasterosteidae) in Nova Scotia,

Canada. Biological Journal of the Linnean Society, 39(3), pp.195-217.

Catchen, J., Hohenlohe, P.A., Bassham, S., Amores, A. and Cresko, W.A., 2013.

Stacks: an analysis tool set for population genomics. Molecular ecology, 22(11),

pp.3124-3140.

Chen, Z., Farrell, A.P., Matala, A., Hoffman, N. and Narum, S.R., 2018. Physiological

and genomic signatures of evolutionary thermal adaptation in redband trout from

extreme climates. Evolutionary applications, 11(9), pp.1686-1699.

Colosimo, P.F., Hosemann, K.E., Balabhadra, S., Villarreal, G., Dickson, M.,

Grimwood, J., Schmutz, J., Myers, R.M., Schluter, D. and Kingsley, D.M., 2005.

Widespread parallel evolution in sticklebacks by repeated fixation of ectodysplasin

alleles. science, 307(5717), pp.1928-1933.

Czech, L., Spence, J.P. and Expósito-Alonso, M., 2024. grenedalf: population genetic

statistics for the next generation of pool sequencing. Bioinformatics, 40(8),

p.btae508.

Deagle, B.E., Jones, F.C., Absher, D.M., Kingsley, D.M. and Reimchen, T.E., 2013.

Phylogeography and adaptation genetics of stickleback from the H'aida G'waii

archipelago revealed using genome   wide single nucleotide polymorphism

genotyping. Molecular ecology, 22(7), pp.1917-1932.

Di-Poi, C., Lacasse, J., Rogers, S.M. and Aubin-Horth, N., 2014. Extensive

behavioural divergence following colonisation of the freshwater environment in

threespine sticklebacks. PloS one, 9(6), p.e98980.

Di-Poi, C., Bélanger, D., Amyot, M., Rogers, S. and Aubin-Horth, N., 2016. Receptors

rather than signals change in expression in four physiological regulatory networks

during evolutionary divergence in threespine stickleback. Molecular ecology,

25(14), pp.3416-3427.

Divino, J.N., Monette, M.Y., McCormick, S.D., Yancey, P.H., Flannery, K.G., Bell,

M.A., Rollins, J.L., von Hippel, F.A. and Schultz, E.T., 2016. Osmoregulatory

physiology and rapid evolution of salinity tolerance in threespine stickleback

recently introduced to fresh water. Evolutionary ecology research, 17(2), pp.179-

201.

Efron, B., 2004. Large-scale simultaneous hypothesis testing: the choice of a null

hypothesis. Journal of the American Statistical Association, 99(465), pp.96-104.

Fang, B., Merilä, J., Ribeiro, F., Alexandre, C.M. and Momigliano, P., 2018. Worldwide

phylogeny of three-spined sticklebacks. Molecular phylogenetics and evolution,

127, pp.613-625.

Fang, B., Kemppainen, P., Momigliano, P., Feng, X. and Merilä, J., 2020a. On the

causes of geographically heterogeneous parallel evolution in sticklebacks. Nature

Ecology & Evolution, 4(8), pp.1105-1115.

Fang, B., Merilä, J., Matschiner, M. and Momigliano, P., 2020b. Estimating

uncertainty in divergence times among three-spined stickleback clades using the

multispecies coalescent. Molecular phylogenetics and evolution, 142, p.106646.

Feder, A.F., Petrov, D.A. and Bergland, A.O., 2012. LDx: estimation of linkage

disequilibrium from high-throughput pooled resequencing data. PloS one, 7(11),

p.e48588.

Filby, A.L., Paull, G.C., Hickmore, T.F. and Tyler, C.R., 2010. Unravelling the

neurophysiological basis of aggression in a fish model. BMC genomics, 11(1),

pp.1-17.

Ferchaud, A.L. and Hansen, M.M., 2016. The impact of selection, gene flow and

demographic history on heterogeneous genomic divergence: Three spine

sticklebacks in divergent environments. Molecular Ecology, 25(1), pp.238-259.

Garcia Elfring, A., Paccard, A., Thurman, T.J., Wasserman, B.A., Palkovacs, E.P.,

Hendry, A.P. and Barrett, R.D., 2021. Using seasonal genomic changes to

understand historical adaptation to new environments: Parallel selection on

stickleback in highly variable estuaries. Molecular Ecology, 30(9), pp.2054-2064.

Ge, S.X., Jung, D. and Yao, R., 2020. ShinyGO: a graphical gene-set enrichment tool

for animals and plants. Bioinformatics, 36(8), pp.2628-2629.

Hagen, D.W. and Moodie, G.E.E., 1982. Polymorphism for plate morphs in

Gasterosteus aculeatus on the east coast of Canada and an hypothesis for their

global distribution. Canadian Journal of Zoology, 60(5), pp.1032-1042.

Haglund, T.R., Buth, D.G. and Blouw, D.M., 1990. Allozyme variation and the

recognition of the "white stickleback". Biochemical systematics and ecology, 18(7-

8), pp.559-563.

Haines, G.E., 2023. Intraspecific diversity of threespine stickleback (Gasterosteus

aculeatus) populations in eastern Canada. Environmental Biology of Fishes,

106(5), pp.1177-1194.

Hartl, D.L., Clark, A.G. and Clark, A.G., 1997. Principles of population genetics (Vol.

116). Sunderland, MA: Sinauer associates.

Heuts, M.J., 1947a. The Phenotypical Variability of Gasterosteus Aculeatus (L.)

Populations in Belgium.... NV Standaard-boekhandel.

Heuts, M.J., 1947b. Experimental studies on adaptive evolution in Gasterosteus

aculeatus L. Evolution, pp.89-102.

Hohenlohe, P.A., Bassham, S., Etter, P.D., Stiffler, N., Johnson, E.A. and Cresko,

W.A., 2010. Population genomics of parallel adaptation in threespine stickleback

using sequenced RAD tags. PLoS genet, 6(2), p.e1000862.

Ishikawa, A., Kusakabe, M., Kume, M. and Kitano, J., 2016. Comparison of

freshwater tolerance during spawning migration between two sympatric Japanese

marine threespine stickleback species. Evolutionary Ecology Research, 17(4),

pp.525-534.

Kakioka, R., Mori, S., Kokita, T., Hosoki, T.K., Nagano, A.J., Ishikawa, A., Kume, M.,

Toyoda, A. and Kitano, J., 2020. Multiple waves of freshwater colonization of the

three-spined stickleback in the Japanese Archipelago. BMC evolutionary biology,

20(1), p.143.

Klepaker, T., 1995. Postglacial evolution in lateral plate morphs in Norwegian

freshwater populations of the threespine stickleback (Gasterosteus aculeatus).

Canadian Journal of Zoology, 73(5), pp.898-906.

Kofler, R., Orozco-terWengel, P., De Maio, N., Pandey, R.V., Nolte, V., Futschik, A.,

Kosiol, C. and Schlötterer, C., 2011. PoPoolation: a toolbox for population genetic

analysis of next generation sequencing data from pooled individuals. PloS one, 6(1), p.e15925.

Kofler, R., Pandey, R.V. and Schlötterer, C., 2011. PoPoolation2: identifying differentiation between populations using sequencing of pooled DNA samples (Pool-Seq). Bioinformatics, 27(24), pp.3435-3436.

Kolaczkowski, B., Kern, A.D., Holloway, A.K. and Begun, D.J., 2011. Genomic differentiation between temperate and tropical Australian populations of Drosophila melanogaster. Genetics, 187(1), pp.245-260.

Lam, T.J. and Hoar, W.S., 1967. Seasonal effects of prolactin on freshwater osmoregulation of the marine form (trachurus) of the stickleback Gasterosteus aculeatus. Canadian Journal of Zoology, 45(4), pp.509-516.

Langmead, B. and Salzberg, S.L., 2012. Fast gapped-read alignment with Bowtie 2. Nature methods, 9(4), p.357.

Le Rouzic, A., Østbye, K., Klepaker, T.O., Hansen, T.F., Bernatchez, L., Schluter, D. and Vøllestad, L.A., 2011. Strong and consistent natural selection associated with armour reduction in sticklebacks. Molecular ecology, 20(12), pp.2483-2493.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G. and Durbin, R., 2009. The sequence alignment/map format and SAMtools. Bioinformatics, 25(16), pp.2078-2079.

Liu, N.A., Liu, Q., Wawrowsky, K., Yang, Z., Lin, S. and Melmed, S., 2006. Prolactin receptor signaling mediates the osmotic response of embryonic zebrafish lactotrophs. Molecular Endocrinology, 20(4), pp.871-880.

Mäkinen, H.S. and Merilä, J., 2008. Mitochondrial DNA phylogeography of the three-

spined stickleback (Gasterosteus aculeatus) in Europe—evidence for multiple

glacial refugia. Molecular Phylogenetics and Evolution, 46(1), pp.167-182.

Mäkinen, H.S., Cano, J.M. and Merilä, J., 2008. Identifying footprints of directional

and balancing selection in marine and freshwater three spined stickleback

(Gasterosteus aculeatus) populations. Molecular Ecology, 17(15), pp.3565-3582.

Mancera, J.M. and McCormick, S.D., 2019. Role of prolactin, growth hormone,

insulin-like growth factor I and cortisol in teleost osmoregulation. In Fish

osmoregulation (pp. 497-515).

Marchinko, K.B. and Schluter, D., 2007. Parallel evolution by correlated response:

lateral plate reduction in threespine stickleback. Evolution: International Journal of

Organic Evolution, 61(5), pp.1084-1090.

Marques, D.A., Jones, F.C., Di Palma, F., Kingsley, D.M. and Reimchen, T.E., 2018.

Experimental evidence for rapid genomic adaptation to a new niche in an adaptive

radiation. Nature ecology & evolution, 2(7), pp.1128-1138.

Marques, D.A., Meier, J.I. and Seehausen, O., 2019. A combinatorial view on

speciation and adaptive radiation. Trends in ecology & evolution, 34(6), pp.531-

544.

Matschiner, M., Hanel, R. and Salzburger, W., 2011. On the origin and trigger of the

notothenioid adaptive radiation. PLoS one, 6(4), p.e18911.

McCairns, R.S. and Bernatchez, L., 2010. Adaptive divergence between freshwater

and marine sticklebacks: insights into the role of phenotypic plasticity from an

integrated analysis of candidate gene expression. Evolution: International Journal of Organic Evolution, 64(4), pp.1029-1047.

McPhail, J.D. and Hay, D.E., 1983. Differences in male courtship in freshwater and marine sticklebacks (Gasterosteus aculeatus). Canadian Journal of Zoology, 61(2), pp.292-297.

Moodie, G.E.E. and Reimchen, T.E., 1976. Phenetic variation and habitat differences in Gasterosteus populations of the Queen Charlotte Islands. Systematic Zoology, 25(1), pp.49-61.

Münzing, J., 1963. The evolution of variation and distributional patterns in European populations of the three-spined stickleback, Gasterosteus aculeatus. Evolution, pp.320-332.

Nagel, L. and Schluter, D., 1998. Body size, natural selection, and speciation in sticklebacks. Evolution, 52(1), pp.209-218.

Nichols, K.M., Kozfkay, C.C. and Narum, S.R., 2016. Genomic signatures among Oncorhynchus nerka ecotypes to inform conservation and management of endangered Sockeye Salmon. Evolutionary Applications, 9(10), pp.1285-1300.

Orti, G., Bell, M.A., Reimchen, T.E. and Meyer, A., 1994. Global survey of mitochondrial DNA sequences in the threespine stickleback: evidence for recent migrations. Evolution, 48(3), pp.608-622.

Paccard, A., Hanson, D., Stuart, Y.E., von Hippel, F.A., Kalbe, M., Klepaker, T., Skúlason, S., Kristjánsson, B.K., Bolnick, D.I., Hendry, A.P. and Barrett, R.D., 2020. Repeatability of adaptive radiation depends on spatial scale: regional versus

global replicates of stickleback in lake versus stream habitats. Journal of Heredity,

111(1), pp.43-56.

Pavlova, N.S., Neretina, T.V. and Smirnova, O.V., 2020. Dynamics of prolactin axis

genes in the brain of male and female three-spined stickleback Gasterosteus

aculeatus (Gasterostaidae) during short-term freshwater adaptation. Journal of

Ichthyology, 60(2), pp.299-304.

Penczak, T., 1965. Morphological variation of the stickleback (Gasterosteus

aculeatus L.) in Poland. Zoologica Poloniae, 15, pp.3-49.

Rafiński, J., Bańgbura, J. and Przybylski, M., 1989. Genetic differentiation of

freshwater and marine sticklebacks,(Gasterosteus aculeatus) of Eastern Europe.

Journal of Zoological Systematics and Evolutionary Research, 27(1), pp.33-43.

Raeymaekers, J.A., Konijnendijk, N., Larmuseau, M.H., Hellemans, B., De Meester,

L. and Volckaert, F.A., 2014. A gene with major phenotypic effects as a target for

selection vs. homogenizing gene flow. Molecular Ecology, 23(1), pp.162-181.

Samuk, K., Visty, H. and Schluter, D., 2025. Genetic divergence in the absence of

strong ecological differences between coexisting white and common Atlantic

marine sticklebacks. bioRxiv, pp.2025-09.

Sanderson, S., Eckert, L., Barrett, R.D., Reimchen, T.E. and Hendry, A.P., 2026.

Contemporary Evolution of an At-Risk Stickleback Population During a Severe

Drought. Evolutionary Applications, 19(1), p.e70189.

Schlötterer, C., 2023. How predictable is adaptation from standing genetic variation?

Experimental evolution in Drosophila highlights the central role of redundancy and

linkage disequilibrium. Philosophical Transactions of the Royal Society B, 378(1877), p.20220046.

Schluter, D. and Conte, G.L., 2009. Genetics and ecological speciation. Proceedings of the National Academy of Sciences, 106(Supplement 1), pp.9955-9962.

Scott, R.J., Haines, G.E., Biedak, N.R. and Baker, J.A., 2023. Variation in morphology among populations of threespine stickleback (Gasterosteus aculeatus) from western Newfoundland, Canada. Environmental Biology of Fishes, 106(9), pp.1889-1905.

Storey, J.D. and Tibshirani, R., 2003. Statistical significance for genomewide studies. Proceedings of the National Academy of Sciences, 100(16), pp.9440-9445.

Su, M., Mu, X., Gui, L., Zhang, P., Zhou, J., Ma, J. and Zhang, J., 2016. Dopamine regulates renal osmoregulation during hyposaline stress via DRD1 in the spotted scat (Scatophagus argus). Scientific Reports, 6(1), p.37535.

Sumarli, A., Behrens, C., Lucas, G., Olufemi, M.J., Bentzen, P., Bell, A.M., Chain, F.J. and Samuk, K., 2025. Strong but diffuse genetic divergence underlies differentiation in an incipient species of marine stickleback. bioRxiv.

Taugbøl, A., Solbakken, M.H., Jakobsen, K.S. and Vøllestad, L.A., 2022. Salinity induced transcriptome profiles in marine and freshwater threespine stickleback after an abrupt 6 hour exposure. Ecology and Evolution, 12(10), p.e9395.

Tine, M., De Lorgeril, J., Panfili, J., Diop, K., Bonhomme, F. and Durand, J.D., 2007. Growth hormone and Prolactin-1 gene transcription in natural populations of the black-chinned tilapia Sarotherodon melanotheron acclimatised to different

salinities. Comparative Biochemistry and Physiology Part B: Biochemistry and
Molecular Biology, 147(3), pp.541-549.

Untergasser, A., Cutcutache, I., Koressaar, T., Ye, J., Faircloth, B.C., Remm, M. and
Rozen, S.G., 2012. Primer3—new capabilities and interfaces. Nucleic acids
research, 40(15), pp.e115-e115.

Williams, T.D., 2008. Individual variation in endocrine systems: moving beyond the
'tyranny of the Golden Mean'. Philosophical Transactions of the Royal Society B:
Biological Sciences, 363(1497), pp.1687-1698.

Yamamoto, K. and Vernier, P., 2011. The evolution of dopamine systems in
chordates. Frontiers in neuroanatomy, 5, p.21.

Yamasaki, Y.Y., Mori, S., Kokita, T. and Kitano, J., 2019. Armour plate diversity in
Japanese freshwater threespine stickleback (Gasterosteus aculeatus).
Evolutionary Ecology Research, 20(1), pp.51-67.

Yan, X., Arakawa, N., Widayati, K.A., Purba, L.H.P.S., Fahri, F., Suryobroto, B., Terai,
Y. and Imai, H., 2025. Exome analysis reveals species divergence in TYR and
identifies species genetic markers in five endemic Macaca species on Sulawesi
Island. BMC Ecology and Evolution, 25(1), p.66.

Figure 1. Sampling localities of threespine stickleback populations in eastern Canada.
The left panel shows eastern North America, with sampled regions highlighted by black rectangles. Insets display collection sites in Newfoundland (upper right) and Nova Scotia (lower right). Freshwater populations are shown as black circles and marine populations as blue triangles.

Figure 2. Population structure and summary population genetic statistics in Eastern Canadian threespine stickleback.

(Top left) Principal component analysis (PCA) based on genome-wide SNP allele frequencies showing differentiation between marine (M) and freshwater (FW) populations; PC1 and PC2 explain 14.17% and 13.1% of the total variance, respectively. Points are coloured by habitat. (Top right) WPGMA phylogram constructed from genome-wide SNP allele-frequency distances illustrating relationships among populations; bootstrap support values are shown at nodes. (Bottom left) Nucleotide diversity (π) estimated for each population, summarizing within-population genetic variation. (Bottom right) Tajima's D values by population, reflecting deviations in the allele frequency spectrum relative to neutral demographic equilibrium. Marine populations are indicated by (M) and freshwater populations by (FW).

Figure 3. Heatmap of median pairwise F$_{ST}$ among populations. Freshwater populations are shown in black and marine populations in blue.

Figure 4. Manhattan plots of F$_{ST}$ values highlighting heterogeneity among threespine stickleback populations from Eastern Canada. Freshwater populations are labelled in black and marine in blue.

Figure 5. Comparison of absolute (d$_{XY}$) and relative (F$_{ST}$) divergence among populations experiencing heterogeneous demographic histories (as inferred by estimates of Tajima's D).

Figure 6. Overlap of F$_{ST}$ outlier SNPs (top 5%) from marine–freshwater (M–FW) comparisons within and between regions. Bars show the number of SNPs shared among increasing numbers of comparisons, and numbers above each bar indicate the total SNPs detected in that overlap category.

Figure S1. Heterogeneity of M-FW differentiation in Eastern Canada (Newfoundland and Nova Scotia).

Figure S2. Manhattan plots of genome-wide differentiation across marine-marine, marine-freshwater, and freshwater-freshwater population comparisons within, and across, Nova Scotia and Newfoundland.