Greater Expression of DNA Repair Pathways in Sharks vs. Rays/Skates Based on Transcriptomic Analyses

Colton R. Simmons
   a. Nova Southeastern University Dr. Kiran C. Patel College of Osteopathic Medicine, AutoNation Institute for Breast Cancer Research and Care
   b. 3200 S University Dr, Fort Lauderdale, FL 33328, United States
   c. cs2911@nova.edu
Schaefer L. Grant (Corresponding Author)
   a. Johns Hopkins University
   b. 3400 N. Charles Street, Baltimore, MD 21218, United States
   c. Permanent address: 3200 S University Dr, Fort Lauderdale, FL 33328, United States
   d. schaefer.grant16@gmail.com
Lucia Llorente Ruiz
   a. Nova Southeastern University, Fort Lauderdale, FL
   b. 3200 S University Dr, Fort Lauderdale, FL 33328, United States
   c. lullorui@hotmail.com
David W. Kerstetter
   a. Halmos College of Natural Science and Oceanography, Department of Marine and Environmental Science, Nova Southeastern University
   b. 8000 N Ocean Dr, Dania Beach, FL 33004, United States
   c. kerstett@nova.edu
Manasi Pimpley
   a. Nova Southeastern University
   b. 3200 S University Dr, Fort Lauderdale, FL 33328, United States
   c. manasirpimple@gmail.com
Jean J. Latimer
   a. Nova Southeastern University, Barry and Judy College of Pharmacy, Department of Pharmaceutical Sciences, AutoNation Institute for Breast Cancer Research and Care
   b. 3200 S University Dr, Fort Lauderdale, FL 33328, United States
   c. latimer.jean@gmail.com

Abstract

Elasmobranchs are an understudied taxon of cartilaginous fishes. DNA repair studies
have been performed in very few elasmobranchs. Because DNA repair maintains the integrity
of the genetic code, it is important for the survival of elasmobranchs in increasingly polluted
oceans. Oil spills, for example, have been shown to cause DNA adducts in marine animals.
We hypothesized that four elasmobranch species would show differential DNA repair
expression. Dermal tissue was harvested from nurse sharks (*Ginglymostoma cirratum*), spiny
dogfish (also considered to be sharks) (*Squalus acanthias*), yellow stingrays (*Urobatis
jamaicensis*), little skates (*Leucoraja erinacea*), and RNA was isolated. RNA sequencing was
performed using the holocephalan Australian ghostshark (*Callorhincus milii*) reference genome.
ANOVA KEGG pathway analysis revealed that RNA from the sharks manifested significantly
higher expression than those of rays/skates in 4/5 major DNA repair pathways (Base Excision,
Nucleotide Excision, Mismatch Repair, and Homologous Recombination). Each of the four
pathways of DNA repair manifested differential expression of pathway-specific genes (*mpg,
polL, parp4, polD2, xpa, gtf2h3, gtf2h5, ercc8, ercc4, cul4b, rad51D, rad51C, blm, ssbp1, top3a,
rad51, xrcc3, mre11a, brip1, rad54b,* etc.). One gene*, polD2,* was consistently elevated in the
sharks vs. rays/skates in all four pathways. A subunit of *polD*, has been proposed to contribute
to the spreading and amplification of hypermutations in sharks by generating higher diversity of
the T cell receptor repertoire. With increased expression of four major DNA repair pathways,
sharks may be more successful than rays/skates in remediating DNA damage and surviving the
genotoxic effects of increasingly polluted oceans and possibly eluding cancer.

Introduction

Maintaining DNA integrity is critical for preventing diseases such as cancer. Assessing
DNA repair and the potential for genetic damage is essential for identifying environmental
threats to ocean life and informing conservation efforts. Pharmaceuticals, agrochemicals, and
environmental shifts due to climate change all present threats to marine life that involve DNA
damage and repair. Although saltwater studies have previously assessed crustaceans, bivalves
and bony fishes (Clade *Osteichthyes*), fewer DNA damage and repair related studies have been
performed on cartilaginous fishes (Clade *Chondrichthyes*), which includes Subclass Holocephali
(chimaeras) and Subclass Elasmobranchii (sharks, rays, skates) (**Fig. 1**) (Alves, 2022;
Consales and Marsili, 2021, Marques, 2025).
All the major DNA repair pathways have been found in bony fishes within Infraclass
Teleostei (Kienzler et al., 2013). Additionally, many teleosts have an additional repair pathway
for UV damage known as photoenzymatic repair (*per*), which is absent in placental mammals.
None of the five major DNA repair pathways have previously been demonstrated in
elasmobranchs, illustrating the novelty of this study.
Elasmobranch fishes have ecological relevance because of their relatively long life-
spans and broad geographic ranges, with an additional role as a food source to humans.
Ecologically, elasmobranchs generally occupy high trophic roles in many marine ecosystems
and exert top-down pressures in local food webs. As predators, elasmobranchs also have
associated vulnerability to the bioaccumulation of chemical contaminants.
Because of overfishing and environmental contamination, over a third of elasmobranch
species are threatened (Dulvy et al., 2021), especially those with coastal, sedentary lifestyles or
those that are directly in contact with the oceanic sediment. Chemical contamination
assessments of elasmobranchs involve accumulation of metals and organic pollutants known to
cause DNA alterations and DNA damage (Alves, 2022; Chynel, 2021). Global climate change
has caused variations in oceanic salinity and acidification, complications that can contribute to
DNA alterations (Alva, 2017; Alves, 2022). The research on DNA damage in ocean species has
primarily addressed these risks in bony fishes (Wirgin and Walman, 1998; Simoniello et al.,
2009; Canedo et al., 2021).

Little is known about the pollution tolerance of elasmobranchs. Reviews like Gelsleichter
& Walker (2010) and Bezerra et al., (2019) focus mostly on contaminant levels rather than DNA
repair mechanisms to remediate specific forms of DNA damage. Other reviews like Tiktak et al.,
(2020) focus on pollutants from a human consumption perspective. Marra et al., (2019) found
molecular sequences associated with DNA repair mechanisms in the great white shark
genomes (*Carcharodon carcharias*) but did not assess relative RNA expression or functionality.
However, many elasmobranchs spend time in coastal, likely more polluted waters for at least
reproductive purposes, such as nursery grounds for pups and egg cases (Heupel et al., 2018).
90

Fig. 1. Evolution of cartilaginous fishes from the Devonian to the present to show time scales of each phylogeny divergence. The width of the spindles in each sub-figure are proportional to the number of families as a rough estimate of diversity. On the right are visual representations of each of the cartilaginous Chondrichthyan fish species used in the analysis, corresponding to different phylogenetic groups. MA indicates millions of years ago. Evolution diagrams from Epipelagic under Creative Commons license; black fish images are public domain obtained through PhyloPic.org.

91
The present study provides novel information on DNA repair pathways (**Fig. 2**) for four
coastal elasmobranchs. Chondrichthyan fishes (combined Holocephalan chimaeras and
elasmobranch sharks, skates, and rays) diverged from bony fishes – including the
sarcopterygian fishes (and Tetrapoda) during the Ordovician Period (485-443 mybp [millions of
years before the present]) (**Fig. 1**). The fish sampled in this study are shown in black in **Fig. 1**
as well as the species used for the reference genome. Batoids are one of the more recent
groups to appear in Neoselachii, appearing at the end of the Jurassic Period (145 mybp). Note
that these eras are only approximate, since most fossils of elasmobranchs are from teeth,
dentine-covered placoid scales, and other hard structures (e.g., dorsal spines on spiny dogfish),
which makes specimen-based, physical character phylogenies difficult. However, these broad
phylogenetic differences between Galeomorphs and Squaliforms are consistent throughout
several analyses (Maisey et al., 2004; Naylor et al., 2012), especially as distinct from Batoidea
(Aschliman et al., 2012). Others have repeatedly confirmed the differences between Rajids
(skates) and Myliobatiformes (rays) (Villalobas-Segura et al., 2022).
106

Methods

Elasmobranch sampling

All the work in this study was covered by NSU IACUC protocol 2017-DK-01 to DWK. Nurse
sharks (*Ginglymostoma cirratum*, n=3) and yellow stingrays (*Urobatis jamaicensis*, n=3) were
hand captured near Fort Lauderdale, Florida, while spiny dogfish (*Squalus acanthias*, n=3) and

little skates (*Leucoraja erinacea*, n=3) were captured by commercial fishing vessels near
Chatham, Massachusetts. Specimens were brought onto the fishing vessel or dock, and then
sterile techniques were performed on the live animal to remove a small sample of skin. The
tissue sample was then placed into a sterile, RNAse-free Eppendorf tube and snap-frozen in a
dry ice and ethanol mixture. Samples were subsequently stored at -80°C.
117

RNA Isolation and Sequencing
Tissue was pulse homogenized on ice for 20-30 seconds in the presence of RNAse
inhibitor in a fume hood. RNA was isolated using the Qiagen RNAeasy Kit, and RNA integrity
and concentration were determined using an Agilent TapeStation. RNA samples were then
subjected to library preparation using Illumina's TrueSeq Stranded Total RNA library generation
kit and sequenced on a 2x150 base-pairs paired-end run using the NextSeq 500 High Output
(300-cycle; 400 million read flow cell). The data were delivered as Fastq files, which were
analyzed using Partek Flow software (Partek Inc., MO).
126

Data Analysis
Alignment of exonic reads was performed using BWA-MEM (Burrows-Wheeler Aligner)
in Partek Flow software using the Australian ghost shark (*Callorhincus milii*) reference genome.
Aligned reads were then quantified using the Partek Annotation Model E/M to produce gene
counts. Gene set ANOVA (FDR value of $\leq$0.05), an unsupervised form of data analysis, was
used to determine statistically significant pathway differences between the sharks vs.
rays/skates. Since four major DNA repair pathways were among the top 50 significant
pathways, we report those here.
RNA sequencing results were based on three biologic RNA samples from each species
of elasmobranch. Four species in total were used (2 sharks and 2 rays/skates) and 3
individuals per species were included. Data were expressed as the mean ± standard error for
each group. An FDR value of $\leq$0.05 was used to determine significance in ANOVA pathway
analyses for KEGG pathways which included pathways of DNA repair (**Table 1** last column).
Unpaired, one-tailed Student's t tests using a p value of ≤ 0.05 were performed on
individual genes in each DNA repair pathway using Microsoft Excel (version 16.63) and
GraphPad (version 9.3.1). Finally, normalization was performed using counts per million reads.
143
144

Fig. 2. The five major pathways of DNA repair are summarized.

145
Results
147
There are multiple types of DNA damages (base damage, bulky adducts and single-
stranded crosslinks, double stranded breaks (blunt-ended or sticky ended) and mismatched
bases after replication. These damages are resolved by five distinct pathways of DNA repair
(**Fig. 2**). Base Excision Repair (BER) and Nucleotide Excision Repair (NER) remediate
abnormal geometry in the DNA helix generated from single strand damages. BER repairs only
the base that is damaged using evolutionarily conserved and specific glycosylases. NER
resolves damages involving alkylating agents (common as pollutants) and single-stranded
crosslinks by removing the area around the damage and resynthesizing a long patch (27-30 bp).
Double stranded helix damage is resolved either by homologous recombination (HR) or non-
homologous end joining (NHEJ). HR uses a sister chromatid as the template for slow, high
fidelity repair, whereas NHEJ is rapid and error prone in comparison.
An examination of the top 50 out of a total of 164 significant KEGG pathways
differentially expressed between sharks vs ray/skates included four of the five major DNA repair
pathways (**Table 1**). These include BER (**Fig. 2, Table 1** and individual genes shown in **Fig. 3**),
NER (**Fig. 2, Table 1** and individual genes shown in **Fig. 4**), HR (**Fig. 2**, **Table 1** and individual
genes shown in **Fig. 5**) and MMR (**Fig. 2**, **Table 1** and individual genes shown in **Fig. 6**) using
Gene Set ANOVA analyses. The Fanconi Anemia pathway was among the top 50 (**Table 1,**
and individual genes shown in **Fig. 7**) and is shown here because it is a disease state that
includes components of double stranded and single stranded DNA repair pathways. It also
includes related pathways such as cell cycle. The only DNA repair pathway not shown to be
differentially regulated by sharks vs. rays/ skates was NHEJ. Given the critical importance of
DNA repair mechanisms in the cell, it is remarkable that 4/5 pathways of repair show the same
trend (higher in sharks).
We subsequently performed an analysis of pathway specific genes using unpaired
Student's t tests identifying the genes that differentially regulated ($p \leq 0.05$)(**Figs. 3-7**). In
addition, genes common to multiple pathways of DNA repair were identified (*ercc4, rad51*,
pold2). The polymerase delta2 subunit of *polD* was common to multiple pathways (BER, NER,
HR and MMR) and significantly higher in sharks vs. rays/ skates. Polymerase delta is
indispensable for leading and lagging strand synthesis in replication. *polD2* does not encode
the catalytic subunit.
There were also pathway specific genes in each of the four pathways showing higher
expression in sharks vs. ray/skates, so the pathway statistical trends identified in the ANOVA
analysis were not driven by a small number of overlapping genes.

181
182

Gene Set Description	Fold Change	P Value	FDR Step Up
Homologous Recombination	5.42	3.43E-7	6.24E-6
Mismatch Repair	11.08	4.35E-6	4.75E-5
Nucleotide Excision Repair	4.76	1.01E-6	1.38E-5
Base Excision Repair	8.88	1.41E-4	9.65E-4
Fanconi Anemia	1.78	3.13E-5	2.6E-4

Table 1. Unsupervised KEGG pathway analysis. DNA repair related pathways that show differential
expression (for the entire pathway) for sharks vs. rays/skates using Gene Set ANOVA . 4/5 major DNA
repair pathways were among the top 50/164) significant pathways; also included was the Fanconi Anemia
pathway which overlap several major DNA repair pathways. P values and FDR (False Discovery Rate)
values were evaluated for these pathways, as a whole. Pathways are ranked using the more stringent
FDR step up criterion. NHEJ is the only major pathway of DNA repair that is not differentially expressed

between these groups. Fanconi anemia is a disease-related pathway. It includes aspects of NER
(Nucleotide Excision Repair) and of HR (Homologous Recombination).
191
192

Fig. 3. RNA expression via RNA sequencing for genes involved in the **Base Excision Repair Pathway**. Significantly different steady state RNA expression was indicated with an asterisk (*) p≤ 0.05. 4/18 genes were significantly higher in sharks vs. rays/skates: *mpg, polL, parp4* and *polD2*. *polD2* is the major polymerase involved in replication and repair.

193
Base Excision Repair is a pre-replicative repair pathway that repairs small base lesions
caused by alkylation, oxidation and deamination. For Base Excision Repair four genes were
significantly higher in sharks vs. rays/skates: *mpg, polL parp4,* and *polD2*. *polD* is the major
polymerase involved in replication and repair. N-methylpurine-DNA glycosylase (*mpg*) initiates
BER by removing a large variety of deaminated, alkylated and lipid-peroxidation induced purine
adducts. At least 11 distinct glycosylases are known in BER in mammals (Krokan and Bjoras,
2013).
201

Fig. 4. RNA expression for genes involved in the **Nucleotide Excision Repair Pathway**. Significantly different steady state RNA expression was indicated with an asterisk (*) $p \leq 0.05$. 6/29 genes were significantly higher in sharks vs. rays/skates: *xpa, gtf2h3, gtf2h5, ercc8, ercc4, cul4b,* and *polD2.* 1/29 genes was significantly higher in rays/skates compared with sharks: *csa/ercc8* which is involved in transcription coupled NER.

NER is a pre-replicative DNA repair pathway which causes the replication fork to pause
so that the repairosome can remediate the damage before DNA is replicated, avoiding mutation
in the progeny strands. Six genes were significantly higher in sharks vs. rays/skates: *xpa,*
gtf2h3, gtf2h5, ercc4, cul4b, and *polD2.* The *gtf2h* gene products are involved in DNA
unwinding, recruitment of downstream DNA repair factors and verification of bulky lesions on
DNA. These proteins are part of the transcriptional machinery as well as transcription coupled
NER (Wu et al., 2008). *xpa* in concert with damaged DNA initiates repair by binding to
damaged sites and stimulates the 5'-3' helicase activity of *xpd* and the DNA translocase activity
of *xpb* (Fadda, 2015). The product of the *ercc4/xpf* gene interacts with *ercc1* which plays a
crucial role in DNA repair and recombination (Krasikova et al., 2021). The *ercc8/csa* gene
product is involved in transcription-coupled NER as opposed to global genomic NER (Saijo,
2013). This gene is expressed at higher RNA levels in rays/skates vs. sharks.
215

Fig. 5. RNA expression for genes involved in the **Homologous Recombination Repair Pathway**. Significantly different steady state RNA expression was indicated with an asterisk (*) $p \leq 0.05$. 11/27 genes show significantly higher shark vs. ray/skate expression: *rad51D, rad51C, blm, ssbp1, top3a, rad51, xrcc3, mre11a, brip1, rad54b,* and *pold2*.

216
The pathway of DNA repair that did not show differential RNA expression between these
groups was NHEJ (data not shown). NHEJ exists in zebrafish so the finding that NHEJ is not
differentially expressed in elasmobranchs cannot be explained by lack of homology. NHEJ and
HR both repair double stranded breaks, however NHEJ is a faster, more efficient pathway
directly ligating the broken ends of DNA in double strand breaks, although often with some loss
of information. NHEJ is inherently more mutation generating than HR (Stinson and Loparo,
2021).
In the HR pathway eleven genes showed significantly greater expression in sharks vs.
rays/skates (*rad51d*, *rad51c*, *blm*, *ssbp1*, *top3A*, *rad51 xrcc3*, *mre11a*, *brip1*, *rad54b*, *polD2*).
rad51c and *rad51d* are tumor suppressor genes in mammals with known mutations associated
with breast and ovarian cancers. The *xrcc3* gene product, a tumor suppressor gene, is part of
the *rad51* paralog protein complex (Uhrig et al., 2024). The *top3A* gene product (topoisomerase
3 alpha) resolves Holliday junctions (replication forks) and is implicated in Bloom's syndrome-
like disorder when mutated (Martin et al., 2018).

Fig. 6. RNA expression for genes involved in the **Mismatch Repair Pathway** which is a post-replicative repair pathway. Significantly different steady state RNA expression was indicated with an asterisk (*) $p \leq 0.05$. 4/15 genes show significantly higher RNA expression in sharks vs. rays/ skates: *ssbp1, mlh1, rfc3,* and *polD2*.

Mismatch Repair is a post replicative repair process. In Mismatch Repair, four genes
show significantly higher expression in sharks vs. rays/skates (*ssbp1*, *mlh1*, *rfc3*, *polD2*). The
replication protein C factor (*rfc*) plays an important role in MMR, acting as a clamp loader,
loading *pcna* onto DNA which is essential for the *mut* endonuclease activity on the mismatched
nucleotide (Skibbens et al., 2005). *mlh1* is part of a protein complex that detects and binds to
mismatches. Mutation in this gene is associated with colorectal cancer (Pećina-Šlaus et al.,
2020). *mlh1* promoter methylation is now considered to be the primary epigenetic cause of
MMR deficiency, leading to microsatellite instability-high sporadic cancers (Rico-Mendes et al.,
2025). Genes that overlap in two or more pathways include *polD2, rfc, ercc4, Rad51,* and
ssbp1.

Fig. 7. RNA expression for genes involved in the **Fanconi Anemia Pathway.** Significantly different steady state RNA expression was indicated with an asterisk (*) $p \leq 0.05$. 12/27 genes show significantly higher RNA expression in sharks vs. rays/skates: *ube2T, rad51C, blm, top3A, rad51, mlh1, rmi1, ercc4, brip1, atr, fancd2, rev3l*. Two genes (*her9*, *telo2*) show the opposite pattern with sharks being lower than rays/skates.

243
The Fanconi Anemia (FA) pathway is included in this analysis although it is not a
pathway of DNA repair but is instead a disease state, that includes aspects of double strand and
single strand repair as well as replication (Moreno et al., 2021). In addition, the FA pathway
also reveals functions that are related to other aspects of genomic instability like cell cycle
regulation. The *atr* gene encodes a protein kinase, a monitor of DNA damage that is involved in
cell cycle checkpoint regulation upstream of DNA repair pathways. The 12 genes expressed at
higher levels in shark vs. ray/skate are: *ube2T*, *rad51C*, *blm*, *top3A*, *rad51*, *mlh1*, *rmi1*, *ercc4*,
brip1, *atr*, *fancd2*, *rev3L*. The gene *ercc4* is part of the NER pathway. The *mlh1* gene product
is part of mismatch repair. The gene product of *top3A* (topoisomerase gene) relieves
supercoiling upstream of replication forks and is included in the replication pathway. The *rad51*
and *rad51c* genes are part of the HR pathway (Uhrig et al., 2024). The *brip1 gene* encodes a
DNA helicase that interacts with the *brca1* gene (Wang et al., 2023).
256
Discussion
The study of wild species that reflect the unusual evolutionary success represented by
the shark is important for our understanding of fundamental biological processes like DNA
repair. The implications for the avoidance of cancer in increasingly polluted oceans are also
important from the dual perspectives of increased immunity and DNA repair.

Numerous freshwater fishes have been studied for functional DNA repair (reviewed by
Keinzler, et al., 2013), including rainbow trout (*Oncorhynchus mykiss*) and zebrafish (*Danio*
rerio). However, similar DNA repair studies are presently lacking for many marine species,
including marine elasmobranchs. Fewer still have involved examination of the transcriptome for
DNA repair pathways as a whole or assessing the expression of individual genes that are
unique to these pathways and are involved in the genesis of cancer. In addition to pathway
analyses, our study has shown that the gene *polD2* which is differentially expressed in all of
these pathways, is higher in sharks than rays/skates.
Despite a close phylogenetic relationship between the four species of elasmobranchs,
we see a clear distinction in gene expression patterns for 4/5 major pathways of DNA repair with
the two sharks being significantly higher in gene expression. The related pathways of DNA
replication and p53 signaling show the same trend (data not shown). Although DNA repair and
replication are related, one does not drive the other; in other words, high replication does not
necessarily correlate with high repair (Latimer et al., 1996; Latimer et al., 2003).
Most DNA repair genes are tumor suppressor genes. Many of the genes that show
differential gene expression in this study with higher expression in sharks vs. rays/skates are
mutated in human cancers (e.g. *brip1* (Suszynska et al 2020)*, top3a* (Broberg et al 2009)*, xpa*
(Feltes and Bonatto, 2015)*, rad51, atr, mlh1* (Song et al 2015)*, ercc4* (Bogliolo et al 2013)*, polD*
(Gola et al 2023; Mendiratta et al., 2021). *mlh1* has particular importance as an epigenetic
regulator of microsatellite instability (when downregulated) and is used as a biomarker to
determine if a cancer will be high mutational burden and therefore a candidate for
immunotherapy (Rico-Mendez et al., 2025). One could speculate that the expression of these
genes may be correlated with a relative lower rate of cancers in sharks (Sendell-Price et al.,
2023) since they are all expressed at higher steady stage RNA levels in the sharks vs
rays/skates. With greater study of wild species this may become apparent.
In our previous work in cancer, we have found a strong correlation between the
transcriptomic patterns, protein expression and functional repair for NER (Latimer et al., 2010)
particularly when assessed for the entire pathway of repair. NER enzymes are expressed at
low levels compared to many other classes of genes (Christmann and Kaina, 2013). Rather
than finding a few genes differentially expressed we have found 4/5 DNA repair pathways as a
whole, are in the top 50 most significantly regulated pathways out of 164. The remarkable
findings of the present study suggest that sharks could be more successful in surviving polluted
waters containing genotoxic contaminants than rays/skates based on 4/5 significantly different
pathways of DNA repair (Everaarts and Sarkar, 1996). One can envision these repair pathways
being advantageous in the context of oil spills, for example. Carcinogens (specifically alkylating
agents) are present in oil and petrochemicals (Rodrigues et al., 2010). Chemicals used to
disperse oil, such as Corexit 9527A which contains 2-Butoxyethanol, permeabilize cell
membranes thus creating greater nuclear exposure from these genotoxic carcinogens (Hoflack,
1997).
An alternative aspect of DNA repair involves somatic hypermutation which has been
found in sharks in T cell specific genes *tcr* gamma and delta variable regions (Diaz et al., 1998;
Zhu and Hsu, 2010). This hypermutation is believed to play a role in increasing the diversity of
the T cell receptor repertoire, allowing sharks to respond to a wider variety of pathogens. This
hypermutation is a function of certain DNA repair related genes such as error prone DNA
polymerases (E, epsilon or D, delta) (Diaz et al., 1998; Zhu and Hu, 2010). Another potential
ramification of increased T cell receptor diversity and increased DNA repair in general, is the
ability to elude cancer.
Although hypermutation is documented in sharks in cells of the immune system, this
study used the skin of all four species of elasmobranchs. Immune cells could have been

present in the skin. NER is known to be tissue specific in mammals (Latimer et al., 1996;
Latimer et al., 2003; Latimer et al., 2008; Latimer et al., 2015) and remediates two types of UV
based damages. HR is also known to be tissue specific in mammals, being more active in stem
cells and actively proliferating cells (Tichy et al., 2010; Bishop and Schiestl, 2002). Tissue
specific DNA repair has not yet been explored in elasmobranchs but could be a opportunity for
future research, particularly between tissues of sensitivity (e.g., gill filaments) or known sites of
toxin accumulation (e.g., liver).
Sharks have existed on this planet 500 million years whereas *H. sapiens* have existed
300,000 years. The study of sharks and other elasmobranchs is essential for our understanding
of biochemical processes related to genomic stability like DNA repair.

References

Alava, J.J., Cheung, W.W., Ross, P.S. and Sumaila, U.R. (2017). Climate change–contaminant interactions in marine food webs: Toward a conceptual framework. *Global Change Biology 23*, 3984–4001. doi:10.1111/gcb.13667

Alves, L.M.F., Lemos, M.F.L., Cabral, H. and Novais, S.C. (2022). Elasmobranchs as bioindicators of pollution in the marine environment. *Marine Pollution Bulletin 176*, 113418. doi:10.1016/j.marpolbul.2022.113418

Aschliman, N.C., Heithaus, M.R., Musick, J.A., Carrier, J.C., McEachran, J.D. and Claeson, K.M. (2012). Phylogeny of batoidea. *Biology of Sharks and Their Relatives* (2nd edition)*, 73–112. doi:10.1201/b11867-10

Bezerra, M.F., Lacerda, L.D. and Lai, C.-T. (2019). Trace metals and persistent organic pollutants contamination in batoids (Chondrichthyes: Batoidea): A systematic review. *Environmental Pollution 248*, 684–695. doi:10.1016/j.envpol.2019.02.070

Bishop, A.J. and Schiestl, R.H. (2002). Homologous recombination and its role in carcinogenesis. *BioMed Research International 2*, 75–85. doi:10.1155/s1110724302204052

Bogliolo, M., Schuster, B., Stoepker, C., Derkunt, B., Su, Y., Raams, A., Trujillo, J.P., Minguillón, J., Ramírez, M.J., Pujol, R., Casado, J.A., Baños, R., Rio, P., Knies, K., Zúñiga, S., Benítez, J., Bueren, J.A., Jaspers, N.G., Schärer, O.D., de Winter, J.P., Schindler, D., and Surrallés, J. (2013) Mutations in ERCC4, encoding the DNA-repair endonuclease XPF, cause Fanconi anemia. *Am J Hum Genet: 92*(5),800-806. doi: 10.1016/j.ajhg.2013.04.002

Broberg, K., Huynh, E., Schläwicke Engström, K., Björk, J., Albin, M., Ingvar, C., Olsson, H. and Höglund, M. (2009). Association between polymorphisms in RMI1, TOP3A, and BLM and risk of cancer, a case-control study. *BMC Cancer, 9*,140. doi:10.1186/1471-2407-9-140.

Canedo, A., de Jesus, L. W. O., Bailão, E. F. L. C., and Rocha, T. L. (2021). Micronucleus test and nuclear abnormality assay in zebrafish (Danio rerio): Past, present, and future trends. *Environmental pollution, 290*, 118019. https://doi.org/10.1016/j.envpol.2021.118019

Christmann, M., & Kaina, B. (2013). Transcriptional regulation of human DNA repair genes following genotoxic stress: trigger mechanisms, inducible responses and genotoxic adaptation. *Nucleic acids research*, *41*(18), 8403–8420. https://doi.org/10.1093/nar/gkt635

Chynel, M., Munschy, C., Bely, N., Héas-Moisan, K., Pollono, C. and Jaquemet, S. (2021). Legacy and emerging organic contaminants in two sympatric shark species from Reunion Island (Southwest Indian Ocean): Levels, profiles and maternal transfer. *Science of The Total Environment 751*, 141807. doi:10.1016/j.scitotenv.2020.141807

Consales, G., Marsili, L. (2021). Assessment of the conservation status of Chondrichthyans: Underestimation of the pollution threat. *The European Zoological Journal 88*, 165–180. doi:10.1080/24750263.2020.1858981

Cooke, S.J., Bergman, J.N., Madliger, C.L., Cramp, R.L., Beardall, J., Burness, G., Clark, T.D., Dantzer, B., de la Barrera, E., Fangue, N.A., Franklin, C.E., Fuller, A., Hawkes, L.A., Hultine, K.R., Hunt, K.E., Love, O.P., MacMillan, H.A., Mandelman, J.W., Mark, F.C., Martin, L.B., Newman, A.E., Nicotra, A.B., Raby, G.D., Robinson, S.A., Ropert-Coudert, Y., Rummer, J.L.,

373   Seebacher, F., Todgham, A.E., Tomlinson, S. and Chown, S.L.(2021). One hundred research
374   questions in conservation physiology for generating actionable evidence to inform conservation
375   policy and Practice. *Conservation Physiology 9*(1), doi:10.1093/conphys/coab009
376

377   Diaz, M., Greenberg, A. S. and Flajnik, M. F. (1998). Somatic hypermutation of the new antigen
378   receptor gene (NAR) in the nurse shark does not generate the repertoire: possible role in
379   antigen-driven reactions in the absence of germinal centers. *Proceedings of the National*
380   *Academy of Sciences*, *95*(24), 14343-14348. doi: 10.1073/pnas.95.24.14343
381

Dulvy, N.K., Pacoureau, N., Rigby, C.L., Pollom, R.A., Jabado, R.W., Ebert, D.A., Finucci, B.,
Pollock, C.M., Cheok, J., Derrick, D.H., Herman, K.B., Sherman, C.S., VanderWright, W.J.,
Lawson, J.M., Walls, R.H.L., Carlson, J.K., Charvet, P., Bineesh, K.K., Fernando, D., Ralph,
G.M., Matsushiba, J.H., Hilton-Taylor, C., Fordham, S.V., Simpfendorfer, C.A., 2021.
Overfishing drives over one-third of all sharks and rays toward a global extinction crisis. *Current*
Biology 31, 5118–5119. doi:10.1016/j.cub.2021.11.008
388

389   Everaarts, J.M. and Sarkar, A. (1996). DNA damage as a biomarker of marine pollution: Strand
390   breaks in Seastars (Asterias Rubens) from the North Sea. *Water Science and Technology 34*,
391   157–162. doi:10.2166/wst.1996.0616
392

393   Fadda E. (2015). Role of the XPA protein in the NER pathway: A perspective on the function of
394   structural disorder in macromolecular assembly. *Comput Struct Biotechnol J.14*,78-85. doi:
395   10.1016/j.csbj.2015.11.007.
396

397   Feltes, B.C. and Bonatto, D. (2014). Overview of xeroderma pigmentosum proteins architecture,
398   mutations and post-translational modifications. *Mutat Res Rev Mutat Res.* doi:
399   10.1016/j.mrrev.2014.12.002.
400

401   Gelsleichter, J. and Walker, C. (2010). Pollutant exposure and effects in sharks and their
402   relatives. *Marine Biology* 491–537. doi:10.1201/9781420080483-c12
403

404   Gola, M., Stefaniak, P., Godlewski, J., Jereczek-Fossa, B.A. and Starzyńska, A. (2023).
405   Prospects of POLD1 in human cancers: A Review. *Cancers (Basel) 15*(6),1905.
406   doi:10.3390/cancers15061905.
407

408   Heupel, M.R., Kanno, S., Martins, A.P. and Simpfendorfer, C.A. (2019). Advances in
409   understanding the roles and benefits of nursery areas for elasmobranch populations. *Marine*
410   *and Freshwater Research 70*(7), 897-907. doi:10.1071/mf18081
411

412   Hoflack, J. (1997). Alteration in methyl-methanesulfonate-induced poly(adp-ribosyl)ation by 2-
413   butoxyethanol in Syrian hamster embryo cells. *Carcinogenesis 18*, 2333–2338.
414   doi:10.1093/carcin/18.12.2333
415

416   Kienzler, A., Bony, S. and Devaux, A. (2013). DNA repair activity in fish and interest in
417   ecotoxicology: A Review. *Aquatic Toxicology* 134-135, 47–56.
418   doi:10.1016/j.aquatox.2013.03.005
419

420   Krasikova, Y., Rechkunova, N. and Lavrik, O. (2021). Nucleotide excision repair: From
421   molecular defects to neurological abnormalities. *International Journal of Molecular Sciences 22*,
422   6220. doi:10.3390/ijms22126220
423

424   Krokan, H.E. and Bjoras, M. (2013). Base excision repair. *Cold Spring Harbor Perspectives in*
425   *Biology 5*(4), doi:10.1101/cshperspect.a012583
426
427   Latimer, J.J., Hultner, M.L., Cleaver, J.E. and Pedersen, R.A. (1996). Elevated DNA
428   excision repair capacity in the extraembryonic mesoderm of the mid-gestation mouse embryo.
429   *Experimental Cell Research 228*, 19–28. doi: 10.1006/excr.1996.0294
430
431   Latimer, J.J., Nazir, T., Flowers, L.C., Forlenza, M.J., Beaudry-Rodgers, Kelly, C.M.,
432   Conte, J.A., Shestak, K., Kanbour-Shakir, A., and Grant, S.G. (2003). Unique tissue-specific
433   level of DNA nucleotide excision repair in primary human mammary epithelial cultures.
434   *Experimental Cell Res 291*,111-121. doi: 10.1016/s0014-4827(03)00368-9
435
436   Latimer, J.J., Johnson, J.M., Miles, T.D., Dimsdale, J.M., Edwards, R.P., Kelley, J.L.,
437   and Grant, S.G. (2008). Cell-type-specific level of DNA nucleotide excision repair in primary
438   human mammary and ovarian epithelial cell cultures. *Cell and Tissue Research 333*, 461–467.
439   doi:10.1007/s00441-008-0645-1
440
441   Latimer, J.J., Johnson, J.M., Kelly, C.M., Miles, T.D., Beaudry-Rodgers, K.A., Lalanne, N.A.,
442   Vogel, V.G., Kanbour-Shakir, A., Kelley, J.L., Johnson, R.R. and Grant, S.G. (2010). Nucleotide
443   excision repair deficiency is intrinsic in sporadic stage I breast cancer. *Proceedings of the*
444   *National Academy of Sciences 107*, 21725–21730. doi:10.1073/pnas.0914772107
445
446   Latimer, J.J., Makekwana, V.J., Pabon-Padin, Y.R., Pimpley, M.R. and Grant, S.G. (2015).
447   Regulation and dysregulation of mammalian nucleotide excision repair: a pathway to
448   nongermline breast carcinogenesis. *Photochem. Photobiol. 91*(2), 493-500. doi:
449   10.1111/php.12387
450
451   Maisey, J.G., Naylor, J.G. and Ward, D.J. (2004). Mesozoic elasmobranchs, neoselachian
452   phylogeny and the rise of modern elasmobranch diversity. *Mesozoic Fishes 3: Systematics,*
453   *Palaeoenvironments and Biodiversity 3*, 17–56.
454
455   Marques, M.B., Luvizotto-Santos, R. and Hauser-Davis, R.A. (2025). Genetic damage in
456   elasmobranchs: A Review. *Environmental Toxicology and Pharmacology 113*, 104607.
457   doi:10.1016/j.etap.2024.104607
458
Marra, N.J., Stanhope, M.J., Jue, N.K., Wang, M., Sun, Q., Pavinski Bitar, P., Richards, V.P.,
Komissarov, A., Rayko, M., Kliver, S., Stanhope, B.J., Winkler, C., O'Brien, S.J., Antunes, A.,
Jorgensen, S. and Shivji, M.S. (2019). White shark genome reveals ancient elasmobranch
adaptations associated with wound healing and the maintenance of Genome Stability.
Proceedings of the National Academy of Sciences 116, 4446–4455.
doi:10.1073/pnas.1819778116
465
Martin, C.-A., Sarlós, K., Logan, C.V., Thakur, R.S., Parry, D.A., Bizard, A.H., Leitch, A., Cleal,
L., Ali, N.S., Al-Owain, M.A., Allen, W., Altmüller, J., Aza-Carmona, M., Barakat, B.A.Y.,
Barraza-García, J., Begtrup, A., Bogliolo, M., Cho, M.T., Cruz-Rojo, J., Dhahrabi, H.A., Elcioglu,
N.H., Gorman, G.S., Jobling, R., Kesterton, I., Kishita, Y., Kohda, M., Le Quesne Stabej, P.,
Malallah, A.J., Nürnberg, P., Ohtake, A., Okazaki, Y., Pujol, R., Ramirez, M.J., Revah-Politi, A.,
Shimura, M., Stevens, P., Taylor, R.W., Turner, L., Williams, H., Wilson, C., Yigit, G., Zahavich,
L., Alkuraya, F.S., Surralles, J., Iglesias, A., Murayama, K., Wollnik, B., Dattani, M., Heath, K.E.,
Hickson, I.D. and Jackson, A.P. (2018). Mutations in TOP3A cause a bloom syndrome-like

disorder. *The American Journal of Human Genetics 103*, 221–231.
doi:10.1016/j.ajhg.2018.07.001
476
477   Mendiratta, G., Ke, E., Aziz, M., Liarakos, D., Tong, M. and Stites, E.C. (2021). Cancer gene
478   mutation frequencies for the U.S. population. *Nature Communications 12*, doi:10.1038/s41467-
479   021-26213-y
480
481   Moreno, O., Paredes, A., Suarez-Obando, F. and Rojas, A., (2021). An update on Fanconi
482   anemia: Clinical, cytogenetic and molecular approaches. *Biomedical Reports 15*.
483   doi:10.3892/br.2021.1450
484
485   Naylor, G.J., Caira, J.N., Jensen, K., Rosana, K.A., Straube, N. and Lakner, C. (2012).
486   Elasmobranch Phylogeny: A mitochondrial estimate based on 595 species. *Biology of Sharks*
487   *and Their Relatives* (2nd edition). doi:10.1201/b11867-9
488
489   Pećina-Šlaus, N., Kafka, A., Salamon, I. and Bukovac, A. (2020). Mismatch repair pathway,
490   genome stability and cancer. *Frontiers in Molecular Biosciences 7*,
491   doi:10.3389/fmolb.2020.00122
492
493   Pratt, J. H. L., Pratt, T. C., Knotek, R. J., Carrier, J. C. and Whitney, N.M. (2022). Long-term use
494   of a shark breeding ground: Three decades of mating site fidelity in the nurse shark,
495   Ginglymostoma cirratum. *PLoS One 17*(10), e0275323. doi: 10.1371/ journal.pone.0275323
496
497   Rico-Méndez, M. A., Trujillo-Rojas, M. A., Ayala-Madrigal, M. L., Hernández-Sandoval, J. A.,
498   González-Mercado, A., Gutiérrez-Angulo, M., Romero-Quintana, J. G., Valenzuela-Pérez, J. A.,
499   Ramírez-Ramírez, R., Flores-López, B. A., & Moreno-Ortiz, J. M. (2025). *MLH1* methylation
500   status and microsatellite instability in patients with colorectal cancer. *Genes*: *16*(2), 182.
501   https://doi.org/10.3390/genes16020182
502
503   Rodrigues, F.P., Angeli, J.P., Mantovani, M.S., Guedes, C.L. and Jordão, B.Q., (2010).
504   Genotoxic evaluation of an industrial effluent from an oil refinery using plant and animal
505   bioassays. *Genetics and Molecular Biology 33*, 169–175. doi:10.1590/s1415-
506   47572010005000006
507
508   Saijo M. (2013). The role of Cockayne syndrome group A (CSA) protein in transcription-coupled
509   nucleotide excision repair. *Mech Ageing Dev 134*(5-6),196-201. doi:
510   10.1016/j.mad.2013.03.008.
511
512   Sendell-Price, A.T., Tulenko, F.J., Pettersson, M., Kang, D., Montandon, M., Winkler, S., Kulb,
513   K., Naylor, G.P., Phillippy, A., Fedrigo, O., Mountcastle, J., Balacco, J.R., Dutra, A., Dale, R.E.,
514   Haase, B., Jarvis, E.D., Myers, G., Burgess, S.M., Currie, P.D., Andersson, L. and Schartl, M.
515   (2023) Low mutation rate in epaulette sharks is consistent with a slow rate of evolution in
516   sharks. *Nat Commun 19*;14(1), 6628. doi:10.1038/s41467-023-42238-x.
517
518   Simoniello, M.F., Gigena, F., Poletta, G., Loteste, A., Kleinsorge, E., Campana, M., Scagnetti, J.
519   and Parma, M.J. (2009). Alkaline comet assay for genotoxic effect detection in Neotropical Fish
520   prochilodus lineatus (Pisces, Curimatidae). *Bulletin of Environmental Contamination and*
521   *Toxicology 83*, 155–158. doi:10.1007/s00128-009-9771-z
522

523 Skibbens R. V. (2005). Unzipped and loaded: the role of DNA helicases and RFC clamp-loading
524 complexes in sister chromatid cohesion. *The Journal of cell biology*, *169*(6), 841–846.
525 https://doi.org/10.1083/jcb.200503129
526
Song, H., Dicks, E., Ramus. S.J., Tyrer, J.P., Intermaggio, M.P., Hayward, J., Edlund, C.K.,
Conti, D., Harrington, P., Fraser, L., Philpott, S., Anderson, C., Rosenthal. A., Gentry-Maharaj,
A., Bowtell, D.D., Alsop, K., Cicek, M.S., Cunningham, J.M., Fridley, B.L., Alsop, J., Jimenez-
Linan, M., Høgdall, E., Høgdall, C.K., Jensen, A., Kjaer, S.K., Lubiński, J., Huzarski, T.,
Jakubowska, A., Gronwald, J., Poblete, S., Lele, S., Sucheston-Campbell, L., Moysich, K.B.,
Odunsi, K., Goode, E.L., Menon, U., Jacobs, I.J., Gayther, S.A., Pharoah, P.D. Contribution of
Germline Mutations in the RAD51B, RAD51C, and RAD51D Genes to Ovarian Cancer in the
Population. *J Clin Oncol. 33*(26), 2901-2907. doi: 10.1200/JCO.2015.61.2408.
535
536 Stinson, B.M. and Loparo, J.J. (2021). Repair of DNA double-strand breaks by the
537 nonhomologous end joining pathway. *Annual Review of Biochemistry 90*, 137–164.
538 doi:10.1146/annurev-biochem-080320-110356
539
540 Suszynska, M., Ratajska and Kozlowski, P. (2020). BRIP1, RAD51C, and RAD51D mutations
541 are associated with high susceptibility to ovarian cancer: mutation prevalence and precise risk
542 estimates based on a pooled analysis of ~30,000 cases. *J Ovarian Res.13*(1): 50. doi:
543 10.1186/s13048-020-00654-3.
544
545 Tichy, E.D., Pillai, R., Deng, L., Liang, L., Tischfield, J., Schwemberger, S.J., Babcock, G.F.,
546 Stambrook, P.J., (2010). Mouse embryonic stem cells, but not somatic cells, predominantly use
547 homologous recombination to repair double-strand DNA breaks. *Stem Cells and Development
548 19*, 1699–1711. doi:10.1089/scd.2010.0058
549
550 Tiktak, G.P., Butcher, D., Lawrence, P.J., Norrey, J., Bradley, L., Shaw, K., Preziosi, R. and
551 Megson, D., (2020). Are concentrations of pollutants in sharks, rays and skates
552 (Elasmobranchii) a cause for concern? A systematic review. *Marine Pollution Bulletin 160*,
553 111701. doi:10.1016/j.marpolbul.2020.111701
554
555 Uhrig, M.E., Sharma, N., Maxwell, P., Gomez, J., Selemenakis, P., Mazin, A.V. and Wiese, C.,
556 (2024). Disparate requirements for RAD54L in replication fork reversal. *Nucleic Acids Research
557 52*, 12390–12404. doi:10.1093/nar/gkae828
558
559 Villalobos-Segura, E., Marramà, G., Carnevale, G., Claeson, K.M., Underwood, C.J., Naylor,
560 G.J., and Kriwet, J.,(2022). The phylogeny of Rays and skates (Chondrichthyes:
561 Elasmobranchii) based on morphological characters revisited. *Diversity 14*, 456.
562 doi:10.3390/d14060456
563
564 Wang, R., Zhang, J., Cui, X., Wang, S., Chen, T., Niu, Y., Du, X., Kong, J., Wang, L. and Jiang,
565 Y. (2023). Multimolecular characteristics and role of BRCA1 interacting protein C-terminal
566 helicase 1 (BRIP1) in human tumors: A pan-cancer analysis. *World Journal of Surgical
567 Oncology 21*(1), doi:10.1186/s12957-022-02877-8
568
569 Wu, W., Liu H., Lei, R., Chen, D., Zhang, S., Lv, J., Wang, Y., Fan. W., Qian, J., Jin, G., Ma, H.,
570 Miao, R., Hu. Z., Wang, H., Jin, L., Wei, Q., Shen, H., Huang, W., Lu, D. (2009) Genetic variants
571 in GTF2H1 and risk of lung cancer: a case-control analysis in a Chinese population. *Lung
572 Cancer 63*(2),180-186. doi: 10.1016/j.lungcan.2008.05.014.
573

Wirgin, I. and Waldman, J.R. (1998). Altered gene expression and genetic damage in North
American fish populations. *Mutation Research/Fundamental and Molecular Mechanisms of*
Mutagenesis 399, 193–219. doi:10.1016/s0027-5107(97)00256-x

Zhu, C. and Hsu, E. (2010). Error-prone DNA repair activity during somatic hypermutation in
shark B lymphocytes *J.Immunol.185* (9), 5336-5347. doi: 10.4049/jimmunol. 1000779

Acknowledgements
Funding was provided by the NSU President's Faculty Research Grant Program (award to JJL
and DWK) and the AutoNation Institute for Breast Cancer Research and Care (JJL). We
appreciate the vessel time by Captains Doug Feeney (F/V *Noah*) and Dave Gelfman (F/V *Horse*
Mackerel), plus the field assistance by Jacqueline Reuder (NSU). Many thanks to Stephen
Zimberg and Carol Murphy for their critical reading of this work. Special acknowledgement goes
to Jennifer Kalil for her work on figure preparation. This paper is dedicated to the memory of
Professor Arnold Miller for instilling a love of science in all who knew him.