Title Page:

Arcuate Fasciculus: Evolutionary Convergence in Marmosets and Humans

Yufan Wang,[1,2,3] Luqi Cheng,[4] Deying Li,[1,2,3] Yuheng Lu,[5,6] William D. Hopkins,[7] Chet C. Sherwood,[8] Ting Xu,[9] Cirong Liu,[10,11,12] George Paxinos,[13] Tianzi Jiang,[1,2,3,14] Congying Chu,[1,2] Lingzhong Fan[1,2,3,15,16,*]

[1]Beijing Key Laboratory of Brainnetome and Brain-Computer Interface, Institute of Automation, Chinese Academy of Sciences, Beijing 100190, China

[2]Brainnetome Center, Institute of Automation, Chinese Academy of Sciences, Beijing 100190, China

[3]School of Artificial Intelligence, University of Chinese Academy of Sciences, Beijing 100049, China

[4]School of Life and Environmental Sciences, Guilin University of Electronic Technology, Guilin 541004, China

[5]School of Biomedical Engineering, Tsinghua University, Beijing 100084, China

[6]Tsinghua Laboratory of Brain and Intelligence, Tsinghua University, Beijing 100084, China

[7]Department of Comparative Medicine, University of Texas MD Anderson Cancer Center, Bastrop, TX 78602, USA

[8]Department of Anthropology and Center for the Advanced Study of Human Paleobiology, The George Washington University, Washington 20052, USA

[9]Center for the Integrative Developmental Neuroscience, Child Mind Institute, New York, NY 10022, USA

[10]Institute of Neuroscience, CAS Center for Excellence in Brain Science and Intelligence Technology, Chinese Academy of Sciences, Shanghai 200031, China

[11]University of Chinese Academy of Sciences, Beijing 100049, China

[12]Key Laboratory of Genetic Evolution & Animal Models, Chinese Academy of Sciences, Shanghai 200031, China

[13]Neuroscience Research Australia and The University of New South Wales, Sydney NSW 2031,

Australia

[14]Xiaoxiang Institute for Brain Health and Yongzhou Central Hospital, Yongzhou 425000, China

[15]School of Life Sciences and Health, University of Health and Rehabilitation Sciences, Qingdao

266000, China

[16]Shandong Key Lab of Complex Medical Intelligence and Aging, Binzhou Medical University,

Yantai 264003, China

*Corresponding author: Lingzhong Fan. Email: lingzhong.fan@ia.ac.cn

Abstract

The marmoset is a highly vocal platyrrhine monkey that shares key anatomical and functional features with humans, providing a unique opportunity to illuminate the phylogenetic origins of diverging connectivity profiles and their transformations throughout evolution. Although the similarity of vocalization features between humans has been reported, whether marmosets possess an arcuate fasciculus homolog is not known. In this study, we delineated the white matter tracts in marmosets, establishing homologies with those observed in other primates, including macaques, chimpanzees, and humans. The presence of an arcuate fasciculus homolog in marmosets was confirmed by tracer and ultra-high-resolution diffusion magnetic resonance imaging datasets. Using a connectivity blueprint approach, we compared cortical connectivity patterns across these species and found the arcuate fasciculus in marmosets terminates in the ventral frontal cortex, with a similarity to humans that exceeds that observed in macaques, something corroborated by quantitative analyses after transforming all brains into a common space. To explore arcuate fasciculus' support for species-specific vocalizations, the activation patterns of vocal communications in marmosets and humans were associated with arcuate fasciculus connectivity. Collectively, our findings suggest that a dorsal pathway, which emerged early in marmoset evolution, has evolved convergently with humans, despite their distant phylogenetic kindship.

Introduction

The common marmoset (*Callithrix jacchus*) is a highly vocal platyrrhine monkey that shares certain features of vocalization with humans, such as pitch perception, vocalization control, and antiphonal calling [1-3]. The species is increasingly recognized as valuable in neuroscience and biomedical research [4,5], suitable for comparative analyses to elucidate the origins of evolutionary adaptations in brain structure and function along the phylogenetic tree [6,7]. The marmoset serves as a promising comparative reference for understanding human brain evolution because of a small, less furrowed brain that facilitates neuroimaging and electrophysiological recording [8], along with its well-known sophisticated and distinctive social life characterized by cooperative family groups, strong pair bonds, shared parental care, and vocal communication to maintain group cohesion [9,10].

Recent studies have reported the capacity of increased voluntary control of vocalizations in nonhuman primates [11,12], suggesting a potential neuroanatomical precursor of language shared with humans [13]. This is supported by the presence of a cortical dorsal-ventral dual stream architecture in several primate species, including marmosets. As a main component of the dorsal stream, the arcuate fasciculus (af) has been confirmed in our closest relatives, such as great apes and macaque monkeys, through non-invasive mapping of brain connections using diffusion MRI (dMRI) [14,15]. Current views emphasize the implication of the arcuate fasciculus for human speech and language processing [16], with its evolutionary adaptations underpinning the complex cognitive functions observed in humans compared to nonhuman primates [14,17-19]. However, the arcuate fasciculus homolog in marmosets, potentially linking the posterior temporal and ventral frontal regions dorsally, and its comparative connectivity patterns across species require further clarification.

In this study, we identified an arcuate fasciculus homolog in marmosets, consistent with evidence in other primates [20-22]. The presence of an af homolog in marmosets was confirmed using tracer and ultra-high resolution dMRI scanning data. To comprehensively explore cortical connection patterns across species, we reconstructed major white matter tracts in marmosets in a similar manner and discovered that the arcuate fasciculus connectivity in the marmoset ventral frontal cortex resembles human patterns more closely than those of macaques. The connectivity

pattern of the af across species was associated with species-specific vocal communication

activation patterns. We quantitatively examined these by transformed them into a common space

by functional alignment, revealing a human-unique extension into the temporal cortex and a

remarkable similarity between humans and marmosets in the anterior connection of the af. Our

findings highlight species-specific cortical connectivity and suggest convergent evolution of the

arcuate fasciculus connections in marmosets and humans, despite an early divergence in the

phylogenetic tree. The logic of our analysis is outlined in Figure 1, providing an overview of the

study's methodology.

Figure 1. Analysis pipeline. (A) Given the similar vocalization features shared between

marmosets and humans, we ask, do marmosets have an arcuate fasciculus? **(B)** Tracer and

ultra-high-resolution dMRI data were used to reveal the existence of af in marmosets and disambiguation with slf. The arcuate fasciculus homolog in marmosets was reconstructed in a manner that corresponds with those available for other primates. **(C)** The connectivity blueprints were constructed for marmosets, macaques, chimpanzees, and humans. This common space approach was further used to examine regions with significant evolutionary changes. **(D)** Connectivity pattern of af across species were associated with the vocal perception and production, and further transformed into a common space for quantitative investigation of their uniqueness.

Results

Marmosets have an arcuate fasciculus

Controversial is the existence in the marmoset of the homologue of the arcuate fasciculus (af), a dorsal pathway interconnecting the posterior temporal cortex and ventral frontal cortex (Figure 2A). We collected both retrograde and anterograde tracer data in the marmoset brain published previously, specifically in Brain/MINDS Marmoset Connectivity Resource (BMCR) [23,24] and Marmoset Brain Connectivity Atlas (MBCA) [25,26]. The available retrograde tracing results showed there are connections between area 45 (A45) and the belt auditory cortex, specifically the auditory cortex caudomedial area (AuCM), and temporoparietal transitional area (Tpt) (Figure 2B). Furthermore, the anterograde neural tracer injected in A45 presented a clear dorsal pathway above the insula (Figure 2C, left and middle) connecting with AuCM and Tpt (Figure 2C, right).

Inspired by the results from tracer experiments, we also investigate this dorsal connection using an ultra-high resolution dMRI dataset [27]. We set A45 in the ventral frontal cortex and the AuCM as the seed and target for diffusion tractography, and vice versa. The same procedure was conducted for A45 and the Tpt. The averaged tracking probability maps showed a dominant dorsal pathway in both hemispheres (Figure 2D), providing evidence of the existence of an af homolog in marmosets.

Figure 2. Both tracer and dMRI tracking showed a dorsal pathway between vPFC and pSTG. (A) Do marmosets have an arcuate fasciculus, which dorsally interconnects vPFC and pSTG? **(B)** Both retrograde and anterograde tracer data in the marmoset brain were collected. The retrograde tracing experiments showed connections between A45 and AuCM and Tpt. **(C)** The anterograde tracer injected in the A45 presented a clear dorsal pathway above the insula, connecting the AuCM and Tpt. **(D)** A dorsal pathway that connects A45 and Tpt/AuCM was clearly observed in the high-resolution dMRI image of marmosets, and this pathway is stronger than the ventral counterpart. A45, area 45; AuCM, auditory cortex caudomedial area; Tpt, temporoparietal transitional area; AC, auditory cortex; af, arcuate fasciculus; lf, lateral fissure; sts, superior temporal sulcus.

To reconstruct the arcuate fasciculus in marmosets, we created tractography protocols which

were defined similarly to those previously established for other primates [20-22,28], while adjusting

for the specific neuroanatomical structures of marmosets (Figure S1A). Besides, given that the

arcuate fasciculus and superior longitudinal fasciculus (slf) are often entangled, especially in

post-mortem blunt dissections, the three branches of the superior longitudinal fasciculus were also

considered (Figure S1B-D). In the ultra-high resolution individual diffusion space, we

reconstructed four white matter tracts (Figure 3) and found clearly distinct pathways for each tract

(Figure 3B), suggesting the disambiguation of af from the three different branches of the slf in

marmoset.

Figure 3. Validation of arcuate fasciculus in marmosets using an ultra-high-resolution
dataset. (A) An arcuate fasciculus homolog was identified in the marmoset. **(B)** The arcuate
fasciculus was distinct from the three branches of the superior longitudinal fasciculus, as shown in
the high-resolution dMRI image of marmosets. A45, area 45; AuCM, auditory cortex caudomedial
area; Tpt, temporoparietal transitional area; af, arcuate fasciculus; slf, superior longitudinal
fasciculus; lf, lateral fissure; sts, superior temporal sulcus.

We additionally used high-resolution dMRI datasets from humans, chimpanzees, and

macaques [29-31] and reconstructed the arcuate fasciculus using the previously established

tractography protocols. All four species showed generally consistent arched shapes that

interconnect the posterior temporal and ventral frontal areas, with the most obvious difference

being that the human af extends to the middle and inferior temporal cortex (Figure 4). Given the

differing brain sizes across species, we resampled the diffusion data according to their brain sizes

and scanning resolution, and reconstructed the af similarly. The results showed that the af was also

found in the resampled data (Figure S2). These findings collectively support strongly the existence

of an arcuate fasciculus homolog in marmosets.

Figure 4. Validation of arcuate fasciculus in four species using ultra-high-resolution datasets.
The arcuate fasciculus of four species were reconstructed using comparable tractography protocols.
All four species showed generally consistent arched shapes that interconnect posterior temporal
and ventral frontal areas, with the most obvious difference being that human af extends to middle
and inferior temporal cortex. cs, central sulcus; lf, lateral fissure; sts, superior temporal sulcus; as,
arcuate sulcus; ots, occipitotemporal sulcus; ips, intraparietal sulcus; ahr, anterior ramus of lateral
fissure; vPFC, ventral prefrontal cortex; AC, auditory cortex; af, arcuate fasciculus; slf, superior
longitudinal fasciculus; unc, uncinate fasciculus; cg, cingulum bundle; cc, corpus callosum; ic,
internal capsule; ac, anterior commissure; ec, external capsule; ex, extreme capsule; fx, fornix;
mdlf, middle longitudinal fasciculus; ilf, inferior longitudinal fasciculus; pcg, parahippocampal
cingulum; ifof, inferior fronto-occipital fasciculus.

Convergent connectivity in marmoset and human arcuate fasciculus

To provide a comprehensive depiction of the connectional changes across cortex between the four

species, we developed protocols to reconstruct other major white matter tracts in a manner

consistent with those available for other primates (Figure 5A, Figure S1). The tracts were

described using the terminology previously employed for comparable species in the

supplementary results. The robustness and generalizability of the tractography protocols were

tested using an independent marmoset dataset (Brain/MINDS marmoset dataset) with a variety of

data quality and acquisition parameters (Figure S3) [32].

Figure 5. Marmoset white matter tracts and tract-to-region connectivity pattern of af. (A)
Horizontal, coronal, and sagittal maximal intensity projections of the population percentage tract
atlases are shown in the center (display range = 30% - 100% of population coverage) with the 3D
visualization displayed in the periphery. **(B)** The connectivity pattern of the arcuate fasciculus in
marmosets, macaques, chimpanzees, and humans. af, arcuate fasciculus; slf1, superior longitudinal
fasciculus I; slf2, superior longitudinal fasciculus II; slf3, superior longitudinal fasciculus III; mdlf,
middle longitudinal fasciculus; ilf, inferior longitudinal fasciculus; ifof, inferior fronto-occipital
fasciculus; unc, uncinate fasciculus; fa, frontal aslant tract; vof, vertical occipital fasciculus; cbd,
cingulum bundle: dorsal; cbp, cingulum bundle: peri-genual; cbt, cingulum bundle: temporal; fx,
fornix; ac, anterior commissure; fma, forceps major; fmi, forceps minor; mcp, middle cerebellar
peduncle; cst, corticospinal tract; ar, acoustic radiation; or, optic radiation; atr, anterior thalamic
radiation; str, superior thalamic radiation; ptr, posterior thalamic radiation.

To investigate the cortical connectivity patterns of the arcuate fasciculus, we counted the

number of times the af tractogram that reached the gray/white matter interface of each cortical

region, obtaining a projection map of af connectivity for four species. All four species showed

connections with the ventral frontal and premotor regions, inferior parietal regions, and superior

temporal regions, but the af only extended into the middle temporal regions in humans (Figure 5B).

Specifically, the af of marmosets connected to A45, proisocortical motor region (ProM), ventral

area 6 (A6V), parietal area PF/PFG (PF/PFGb), Tpt, auditory belt area medial (BeltM), and other

regions (Figure 5B, top).

We next calculated the "connectivity blueprint" by multiplying the unwrapped white matter

tract matrix by the whole-brain connectivity matrix [21], whose rows represented each subregion's

connectivity distribution pattern. We leveraged the connectivity blueprints for humans,

chimpanzees, macaques, and marmosets to compare the connectional changes in the arcuate

fasciculus across species. A modified dissimilarity measure used to quantify the difference

between two probability distributions, the symmetric Kullback-Leibler (KL) divergence [21], was

calculated to measure the dissimilarity of the regional connectivity blueprints of humans and each

of three nonhuman primates between each subregion in the corresponding Brainnetome atlas. The

minimum divergence for each subregion of the Human Brainnetome Atlas (HumanBNA) [33] was

denoted as the connectivity divergence for that region, resulting in a connectivity divergence map

(Figure 6A). Higher values on the divergence map indicate that a region in humans had a

connectivity pattern that was more dissimilar to the corresponding regions in the other species, i.e.,

it might not be represented in this close phylogenetic relative [34].

As shown in the left panel of Figure 6A and Figure S4A, the regions with the most different

connectivity patterns between humans and marmosets were mainly found in the anterior and

middle cingulate cortex and the dorsolateral prefrontal cortex (PFC). The regions with the most

different connectivity patterns between humans and macaques were in the dorsal and medial

prefrontal cortex and posterior inferior parietal lobule (IPL) (Figure 6A, middle; Figure S4B). In

contrast, the regions with the most different connectivity patterns between humans and

chimpanzees were located in the middle and posterior temporal lobe, posterior IPL, anterior

precuneus (Pcun), insula, and inferior frontal gyrus (IFG) (Figure 6A, right; Figure S4C).

217
Figure 6. Connectivity divergence across marmosets, macaques, chimpanzees, and humans.
(A) From left to right: the connectivity divergence map between humans and marmosets,
macaques, and chimpanzees, respectively. Humans present unique connectivity profiles in the
middle temporal regions, lateral frontal cortex, posterior parietal cortex, and anterior cingulate
cortex compared with the other three nonhuman primates. **(B)** Connectivity strength of af with
subregions with high connectivity divergence. For example, the connection between A45c and af
was weak in chimpanzees and macaques, but stronger in marmosets and strongest in humans. **(C)**
The connectivity changes after "virtually lesioning" the white matter tracts connected with A45c,
including af, slf3, unc, and atr. The connectivity divergences between species changed differently
across tracts, with the af leading to various changes across cortical regions compared to other three
tracts. Focusing on the A45c, the KL divergence between humans and marmosets showed the
smallest change, while the macaques and chimpanzees exhibited larger changes. A45c, caudal area
45; A22c, caudal area 22; A21c, caudal area 21; af, arcuate fasciculus; slf3, superior longitudinal
fasciculus III; unc, uncinate fasciculus; atr, anterior thalamic radiation; ifs, inferior frontal sulcus;

prcs, precentral sulcus; lf, lateral fissure; ahr, anterior ramus of lateral fissure. *n.s.*, non-significant,

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

Meanwhile, minimum divergence indicates which subregion in the nonhuman primate brain

showed the most similar connectivity pattern to that of humans, i.e., a "homologous" subregion

defined by connectivity. This method was validated using cytoarchitectonic data to ensure

cross-species correspondence [35,36] (Figure S5). For example, when examining human caudal area

45 (A45c), the KL divergence between its connectivity blueprint and that of each marmoset

subregion was calculated. The minimum divergence was observed in area 45 in marmosets (Figure

S5A), which shared similar laminar features with human area 45, including larger pyramidal cells

in layer III and a well-defined granular layer IV (Figure S5B) [37,38], suggesting cytoarchitectonic

homology between the two species in this region.

We focused on brain regions connected to the arcuate fasciculus. Interestingly, we found that

the connection between the af and ventral PFC (A45c) was stronger in marmosets than in

macaques (Figure 6B, a; $t_{\text{human-chimp}} = 5.17$, $p_{\text{corrected}} < 0.001$; $t_{\text{human-macaque}} = 5.08$, $p_{\text{corrected}} < 0.001$;

$t_{\text{human-marmoset}} = 4.19$, $p_{\text{corrected}} < 0.01$; $t_{\text{marmoset-chimp}} = 0.76$, $p = 0.45$; $t_{\text{marmoset-macaque}} = 4.85$, $p_{\text{corrected}} <$

0.01). This suggests the connectivity of marmosets along the dorsal pathway is more similar to

that in humans than in macaques, consistent with previous findings [1]. The af extended and curved

into the temporal lobe (caudal area 22 [A22c]) and even caudodorsally to the caudal area 21 (A21c)

(a caudal area in the middle temporal regions) resulting in human-unique connectivity of the af

compared to the other three nonhuman primates (Figure 6B, b & c; A22c: $t_{\text{human-chimp}} = 7.68$,

$p_{\text{corrected}} < 0.001$; $t_{\text{human-macaque}} = 3.90$, $p_{\text{corrected}} < 0.05$; $t_{\text{human-marmoset}} = 6.62$, $p_{\text{corrected}} < 0.001$; A21c:

$t_{\text{human-chimp}} = 13.52$, $p_{\text{corrected}} < 0.001$; $t_{\text{human-macaque}} = 5.70$, $p_{\text{corrected}} < 0.001$; $t_{\text{human-marmoset}} = 9.81$,

$p_{\text{corrected}} < 0.001$). This conclusion was supported using another macaque and marmoset datasets

(Figure S6).

As mentioned in the previous findings, the dorsal connection with the ventral PFC was more

similar between humans and marmosets, a platyrrhine monkey that split from their last shared

common ancestor much earlier than macaques and chimpanzees. This may suggest the marmosets

af has continued to evolve along the phylogenetic lineage, becoming more similar to humans

through convergent evolution under socio-environmental conditions that favor similarities in

function. We further explored how connectivity would change after "virtually lesioning" the af in the four species (Figure 6C, top left). The connectivity divergences between species changed differently (Kolmogorov-Smirnov test: marmoset: $p = 0.969$; macaque: $p = 0.007$; chimpanzee: $p = 0.216$; Figure 6C, bottom left). Focusing on A45c, the KL divergence between humans and marmosets showed the smallest change ($\Delta_{\mathrm{marmoset}} = -0.02$), while macaques and chimpanzees showed higher changes ($\Delta_{\mathrm{macaque}} = -0.51$ and $\Delta_{\mathrm{chimp}} = -0.37$; Figure 6C). These differences were not obvious when knocking out other white matter tracts, including the third branch of superior longitudinal fasciculus (slf3) ($\Delta_{\mathrm{marmoset}} = -0.03$; $\Delta_{\mathrm{macaque}} = -0.10$, $\Delta_{\mathrm{chimp}} = -0.19$), uncinate fasciculus (unc) ($\Delta_{\mathrm{marmoset}} = -0.10$; $\Delta_{\mathrm{macaque}} = -0.01$, $\Delta_{\mathrm{chimp}} = -0.05$), and anterior thalamic radiation (atr) ($\Delta_{\mathrm{marmoset}} = -0.13$; $\Delta_{\mathrm{macaque}} = -0.14$, $\Delta_{\mathrm{chimp}} = -0.08$; Figure 6C, right). This suggests the specificity and involvement of af projections in the ventral PFC compared to other connected tracts. Meanwhile, these findings provide additional evidence for the higher similarity of the af connection with the ventral PFC between humans and marmosets.

Finally, given the shared features of vocalization between humans and marmosets, we sought to assess the involvement of af in vocal communication in humans and marmosets. The human af is well known for its role in speech and language processing, something supported by the overlap between part of its cortical connection with the functional activation of "speech" obtained from the NeuroSynth database [39] (Figure 7A; see also the "speech perception" and "speech production" in Figure S7). Specifically, auditory cortex and more posterior superior temporal cortex were activated during speech perception, in contrast, inferior frontal gyrus, ventral premotor and motor cortex were activated during speech production.

Figure 7. Comparison of af connectivity patterns across species. **(A)** Activation patterns of human speech and the connectivity patterns of af. **(B)** The transformed activation patterns in marmoset cortex. **(C)** Activation patterns of marmoset vocal communications. **(D)** The af connectivity patterns transformed from macaques and marmosets using the functional alignment techniques. **(E)** Binarized af connectivity patterns of the three species and quantitative Dice coefficients and tract extension ratios across these maps. **(F)** Weighted correlation maps of the actual human map and the transformed macaque and marmoset maps. The left panel showed the distribution of weighted correlation values in the A45c.

We transformed these activation maps to marmoset cortex based on functional-alignment techniques [40,41] (Figure 7B), which showed overlap with the connectivity pattern of af in marmosets, including A6V, and etc. Although whole-brain imaging of marmosets during vocal communication are difficult to obtain, previous fMRI and electrophysiological experiments provide important evidence of the involvement of these cortical regions in vocalization perception [42] and also natural vocal communication [43] (Figure 7C).

We further quantitatively evaluate across species the similarity of connectivity patterns of af by transforming them into a common space using the same functional-alignment techniques (Figure 7D). Note the chimpanzees were not considered here because their functional data were unavailable. We first calculated the Dice coefficient to assess the overlap similarity between the

actual and transformed tract connection map, and a "tract extension ratio" to assess how much of

the human tract projections extend into areas where other species do not [18]. It can be observed that

marmosets showed the highest tract connection overlap with humans compared to macaques (left:

$Dice_{\text{macaque}} = 0.196$, $Dice_{\text{marmoset}} = 0.255$; right: $Dice_{\text{macaque}} = 0.227$, $Dice_{\text{marmoset}} = 0.271$; Figure

7E). The tract extension ratios for the three species showed a similar degree, with all values

greater than 1, indicating the af in humans extended further, into more brain areas, during

evolution (left: $Ratio_{\text{macaque}} = 3.457$, $Ratio_{\text{marmoset}} = 2.251$; right: $Ratio_{\text{macaque}} = 2.627$, $Ratio_{\text{marmoset}}$

$= 2.089$; Figure 7E).

To further localize where the af showed similar connections across species, we calculated the

weighted local correlation map between the actual and transformed tract connection map, where

higher values indicated brain areas with both the tract of human and the other species showing a

termination. Focusing on the ventral PFC (A45c), the results showed that marmosets exhibited

higher similarity with humans than macaques (median correlation in the left hemisphere: $r_{\text{macaque}} =$

0.0107; $r_{\text{marmoset}} = 0.0426$; right: $r_{\text{macaque}} = 0.0097$; $r_{\text{marmoset}} = 0.0119$; Figure 7F).

Discussion

In this study, we identified an arcuate fasciculus homolog in marmoset brains that corresponds with those in other primates. We also found that marmosets exhibited a more similar connection pattern of the arcuate fasciculus with humans in the ventral frontal cortex, something reconfirmed by quantitative analyses after transformation into a common space. Our results suggest the presence of an earlier-emerged dorsal pathway in marmosets that convergently evolved aspects of its connectivity with humans.

Previous studies have largely focused on the Catarrhini, neglecting potential evolutionary changes in the Platyrhini after their ancestors split. By leveraging species-equivalent tractography protocols, which are informed by anatomical insights into reconstructing the arcuate fasciculus in human, chimpanzee, and macaque brains [20], we identified a dorsal fiber pathway that connects the posterior temporal and ventral frontal regions, traversing through the inferior parietal and ventral premotor areas, while also being distinct from the three branches of the superior longitudinal fasciculus, suggesting the presence of the long suspected arcuate fasciculus homolog in marmosets. Additionally, tract-tracing experiments showed that area 45 in the ventral frontal cortex receives monosynaptic auditory input from the belt auditory cortex [25,26] and also sends information to the superior temporal cortex and posterior parietal cortex [23,24]. These dMRI and tracer observations on the putative arcuate fasciculus might push back the emergence of this dorsal pathway further than the split from a common ancestor with marmosets, approximately 35 million years ago [14,44]. The present data provide further evidence for the auditory dorsal pathway among primate species, a pathway previously observed in tractography following seeding in auditory regions [1,27], supporting species-specific vocalizations during social behaviors.

Most interestingly, unlike the weak connection to the ventral frontal cortex in macaques, marmosets showed a stronger connection, which was more similar to that of humans, suggesting the dorsal pathway in marmosets may have continued to evolve independently in closer alignment with humans than with macaques after the marmoset lineage diverged [1]. The similarity in the ventral frontal cortex between humans and marmosets may help explain the similarity in the perception of vocalizations in marmosets and speech in humans [45]. Production and use of

vocalizations in primates show similar spectral characteristics and neural circuits with human

speech [46], particularly the prefrontal and premotor cortex in marmosets, which has been found to

be associated with the production of social communication signals across the marmoset vocal

repertoire [47]. Another complex process associated with both vocal perception and production,

previously unreported in nonhuman primates—modification of vocal production according to the

acoustic environment, i.e., voluntary vocal control—was observed in marmosets, a talent that

allows them to avoid acoustic interference and maintains effective communication [48,49].

Our data raise the question of whether these changes in the arcuate fasciculus are unique to

marmosets or are shared among other platyrrhines, because no studies have yet examined this

feature in other platyrrhine species, such as capuchin monkeys, howler monkeys, or others. It is

also unknown whether these species share an ancestral trait which undergoes parallel evolutionary

modifications independently. This calls for broader comparative studies to map af connectivity

across more primate species even with different vocalization patterns and to integrate genetic, and

developmental perspectives.

Comparable vocal communication complexity may have exerted similar selective pressure on

brain connectivity in humans and marmosets, both species having evolved in environments

requiring adaptability and intricate social interactions, which might have driven the enhancement

of auditory-motor integration. The highly cooperative social structures favored stronger

sensorimotor integration for real-time social coordination or even predictive social processing [50-52],

reinforcing the af-like pathways beyond vocal communications.

In contrast to the findings in the ventral frontal cortex, the relatively high connectivity

differences in the arcuate fasciculus within the middle and posterior temporal cortex, along with

observed hemispheric differences between species, underscore a potential human-specific

adaptation [53,54]. These adaptations may have played a role in the evolution of language in the

human lineage, particularly in enabling the development of more complex linguistic abilities, such

as speech processing [54,55]. Despite these differences, the vocal behavior among primate species

provides a window into the foundational elements of human speech, suggesting an early

evolutionary origin of some of its components. This behavior was further refined in human

language to enhance social communication [13].

In addition to its relevance to speech, this study provided a detailed marmoset white matter

tract atlas comparable to those of other primate species, facilitating cross species investigations of

similarities and differences in fiber pathways. As an important primate model, the common

marmoset shares key brain organizational features with larger primates, making it feasible to

identify marmoset white matter tracts based on comparative evidence from other primates [27,56].

Although previous research has presented a detailed delineation of white matter pathways using

ultra-high-resolution dMRI datasets [27], that mapping was limited to a single *ex vivo* subject,

reducing its generalizability to larger cohorts and its utility for direct quantitative comparison with

other species. The literature is now complemented by the robust and reproducible tractography

protocols we used, protocols grounded in prior anatomical knowledge and registered to individual

space to reconstruct white matter tracts consistently across subjects while accounting for

individual variability. The established population-based marmoset white matter tract atlas will

serve as a resource for comparative research.

In conclusion, we employed dMRI to map marmoset white matter tracts, identifying an

arcuate fasciculus homolog and its convergent similarities with humans in the ventral frontal

cortex. This provided an opportunity to extend comparative research to more species and reveal

general organizational principles of primate brains. Future exploration that also take mammals and

birds into consideration could offer new insights into how intelligence evolved from divergent

neural structures [57]. Although this study preliminarily investigated the functional association of af

connectivity, future comparative analyses combining electrophysiological recordings, behavioral,

or lesion studies could provide direct evidences linking af connectivity to vocal or cognitive

functions, supporting the investigation of cognitive specializations in humans and nonhuman

primates.

Methods

Data acquisition and preprocessing

For an intuitive reading and comparison across datasets, the scanning parameters of the diffusion dataset used in current study, including spatial resolution and diffusion directions, are summarized in Table S1 (ultra-high-resolution datasets) and S2 (lower-resolution datasets).

Marmoset dataset 1: Marmoset Brain Mapping (MBM)

Marmoset (*Callithrix jacchus*; $n = 24$, 17 males; mean age, 4.17±1.88 y; age range, 1.88-9.00 y) MRI scans were obtained from a data archive of the Marmoset Brain Mapping Project (https://marmosetbrainmapping.org/) [58]. The experimental procedures followed policies established by the US Public Health Service Policy on Humane Care and Use of Laboratory Animals. All procedures were approved by the Animal Care and Use Committee (ACUC) of the National Institute of Neurological Disorders and Stroke, National Institutes of Health.

All the marmosets underwent a 3–4-week acclimatization protocol, as previously described [58,59]. After completing the training, they were properly acclimated to lying in the sphinx position in an MRI-compatible cradle with their heads comfortably restrained with 3D-printed anatomically conforming helmets.

The marmosets were scanned in a 7T/30 cm horizontal MRI (Bruker, Billerica, USA) equipped with a 15 cm customized gradient set capable of 450 mT/m gradient strength (Resonance Research Inc., Billerica, USA) and an 8-channel phased-array RF coil custom-built for marmosets [60]. T2-weighted structural images were collected at a $0.25 \times 0.25 \times 0.5$ mm resolution (TR = 6000 ms, TE = 9 ms, flip angle = 90°, FOV = 28×36 mm, matrix size = 112×144, 38 axis slices, slice thickness = 0.5 mm). Multishell diffusion MRI data were collected using a 2D diffusion-weighted spin-echo EPI sequence with the following parameters: TR = 5.1 s, TE = 38 ms, number of segments = 88, FOV = 36×28 mm, matrix size = 72×56, slice thickness = 0.5 mm, a total of 400 DWI images for two-phase encodings (blip-up and blip-down) and each had 3 b values (8 $b = 0$, 64 $b = 1000$ s/mm^2, and 128 $b = 2000$ s/mm^2). The multishell gradient sampling scheme was generated using the Q-shell sampling method.

The preprocessing of the *in vivo* diffusion MRI dataset incorporated denoising, correcting for

eddy currents- and EPI-induced distortions using pairs of diffusion data acquired with opposite

phase encoding (blip-up and blip-down), and merging the preprocessed pairs into one dataset. The

nonlinear spatial registration from the individual space to the DTI template of MBMv3 space [61]

was carried out using ANTs, which were then used to transform the landmarks defined on the

template into individual space for tractography. Voxel-wise estimates of the fiber orientation

distribution were calculated using bedpostx [62], allowing for three fiber orientations for the dataset.

Marmoset dataset 2: Brain/MINDS Marmoset Brain MRI Dataset (Brain/MINDS)

Another cohort of adult marmosets was obtained from the Brain/MINDS Marmoset Brain MRI

Dataset (https://dataportal.brainminds.jp/marmoset-mri-na216) [32,63] by retaining only the adult

ones. Our study included 110 adult marmosets with both structural and *in vivo* diffusion-weighted

images (50 males; mean age, 4.38±2.47 y; age range, 1.73-10.02 y). All procedures were approved

by the Animal Experiment Committees of the RIKEN Center for Brain Science (CBS) and

conducted following the Guidelines for Conducting Animal Experiments of RIKEN CBS.

The marmosets were scanned in a 9.4-T BioSpec 94/30 unit (Bruker Optik GmbH, Ettlingen,

Germany) and a transmit and receive coil with an inner diameter of 86 mm. T1-weighted

structural images were collected using a magnetization-prepared rapid gradient echo (MP-RAGE)

with the following parameters: TR = 6000 ms, TE = 2 ms, flip angle = 12°, number of averages =

1, inversion time = 1600 ms, voxel size = $0.27 \times 0.27 \times 0.54$ mm. T2-weighted structural images

were collected using rapid acquisition with relaxation enhancement (RARE), with the following

parameters: TR = 4000 ms, TE = 22 ms, RARE factor = 4, flip angle = 90°, number of averages =

1, voxel size = $0.27 \times 0.27 \times 0.54$ mm. Multishell diffusion MRI data were collected using a

spin-echo EPI sequence with the following parameters: TR = 3 s, TE = 25.6 ms, number of

segments = 6, voxel size = $0.35 \times 0.35 \times 0.7$ mm, a total of 94 DWI images (4 b = 0, 30 b = 1000

s/mm^2, and 60 b = 3000 s/mm^2). The preprocessing of the dataset was the same with the MBM

dataset.

Marmoset dataset 3: High-resolution marmoset data

A formalin-fixed brain sample from one male common marmoset (4.5 y) with

ultra-high-resolution MRI scanning was used [27]. Details of procedures before MRI scanning can

be found in previous publications [27]. The brain sample was scanned on a 7T/30-cm horizontal bore

MRI spectrometer (Bruker Biospin) with a 30-mm inner diameter quadrature millipede coil (ExtendMR). Multishell diffusion MRI data were collected using a 3D diffusion-weighted multi-shot spin-echo EPI sequence with the following parameters: TR = 200 ms, TE = 29 ms, number of averages = 1, number of segments = 88, FOV = 38 × 29.76 × 29.76 mm, matrix size = 474 × 372 × 372, resolution = 80 μm isotropic, a total of 204 DWI images with four b values (8 b = 0, 6 b =30 s/mm^2, 64 b = 2400 s/mm^2, and 126 b = 4800 s/mm^2). The diffusion MRI data of ex $vivo$ brain samples had been preprocessed using the DIFFPREP of TORTOISE as in the previous study [27].

Human dataset 1: Human Connectome Project (HCP)

Data from 40 healthy human adults (17 males) were selected. These were the 40 subjects that were utilized in the construction of the Human Brainnetome Atlas [33] as well as the ones available from the S1200 subjects release of the Human Connectome Project (HCP) database (http://www.humanconnectome.org/) [64]. All the scans and data from the individuals included in the study had passed the HCP quality control and assurance standards.

The scanning procedures and acquisition parameters were detailed in previous publications [65]. In brief, T1w images were acquired with a 3D MPRAGE sequence on a Siemens 3T Skyra scanner equipped with a 32-channel head coil with the following parameters: TR = 2400 ms, TE = 2.14 ms, flip angle = 8°, FOV = 224×320 mm^2, voxel size = 0.7 mm isotropic. DWI images were acquired using single-shot 2D spin-echo multiband echo planar imaging on a Siemens 3 Tesla Skyra system (TR = 5520 ms, TE = 89.5 ms, flip angle = 78°, FOV = 210×180 mm). These consisted of three shells (b-values = 1000, 2000, and 3000 s/mm^2) with 90 diffusion directions isotropically distributed among each shell and six b = 0 acquisitions within each shell at a spatial resolution of 1.25 mm isotropic voxels.

The diffusion-weighted images were preprocessed using FSL [66]. Following preprocessing, voxel-wise estimates of the fiber orientation distribution were calculated using bedpostx [62], allowing for three fiber orientations for the human dataset.

Human dataset 2: High-resolution human data

A whole-brain in $vivo$ diffusion MRI dataset acquired at 760 μm isotropic resolution was used [30]. The data were acquired on the MGH-USC 3T Connectom scanner equipped with high-strength

gradient ($G_{max} = 300$ mT/m), a custom-built 64-channel phased-array coil and personalized
motion-robust head stabilizer, using the recently developed SNR-efficient simultaneous multi-slab
imaging technique termed gSlider-SMS. Advanced parallel imaging reconstruction with
ghost-reduction algorithm (Dual-Polarity GRAPPA) and reversed phase-encoding acquisition
were also performed to mitigate ghosting artifacts and image distortion. The data was totally
acquired at 1260 q-space points including 420 directions at $b = 1000$ s/mm^2 and 840 directions
at $b = 2500$ s/mm^2 across the 9 sessions, and was employed by an optimized preprocessing
pipeline.

Chimpanzee dataset 1: National Chimpanzee Brain Resource (NCBR)

We obtained 46 adult chimpanzees (18 males) with T1w and DWI data from the National
Chimpanzee Brain Resource (NCBR, http://www.chimpanzeebrain.org/). The data were acquired
using previously described procedures at the Emory National Primate Research Center (ENPRC)
on a 3T MRI scanner under propofol anesthesia (10 mg/kg/h) [67]. All procedures followed
protocols approved by ENPRC and the Emory University Institutional Animal Care and Use
Committee (IACUC, approval no. YER-2001206). All data were obtained before the 2015
implementation of U.S. Fish and Wildlife Service and National Institutes of Health regulations
governing research with chimpanzees. All chimpanzee scans were completed by the end of 2012;
no new data were acquired for this study.

T1w images were collected at an $0.7 \times 0.7 \times 1$ mm resolution. DWI images were acquired
using a single-shot spin-echo echo-planar sequence for 60 diffusion directions (b = 1000 s/mm^2,
TR = 5900 ms; TE = 86 ms; 41 slices; 1.8 mm isotropic resolution). DWI images with
phase-encoding directions (left-right) of opposite polarity were acquired to correct susceptibility
distortion. Five b = 0 s/mm^2 images were also acquired with matching imaging parameters for
each repeat of a set of DWI images.

The diffusion-weighted images were preprocessed similarly using FSL [66]. Following
preprocessing, voxel-wise estimates of the fiber orientation distribution were calculated using
bedpostx [62], allowing for two fiber orientations for the chimpanzee dataset due to the *b*-value in
the diffusion data.

Chimpanzee Dataset 2: High-resolution chimpanzee data

A whole-brain chimpanzee dMRI dataset was acquired at 500 µm isotropic voxel size on a preclinical Bruker Biospec 94/30 MRI system at 9.4T (Paravision 6.0.1), using a $G_{max} = 300$ mT/m gradient system and a 154-mm transmit-receive quadrature radiofrequency coil (Bruker BioSpin). [29]. Details of procedures before MRI scanning can be found in previous publications [29]. Diffusion MRI data were collected using a segmented 3D EPI spin-echo sequence with the following parameters: TR = 1,000 ms, TE = 58.9 ms, matrix size = $240 \times 192 \times 144$, no partial Fourier, no parallel acceleration, EPI segmentation factor of 32 and EPI-Readout-BW of 400 kHz, 55 DWI images with $b = 5000$ s/mm^2. Three interspersed $b = 0$ images without diffusion-weighting and an additional $b = 0$ volume were acquired with reversed-phase encoding direction for correction.

Macaque dataset 1: TheVirtualBrain (tvb)

Data from eight adult rhesus macaque monkeys (all males) were obtained from TheVirtualBrain [68]. All surgical and experimental procedures were approved by the Animal Use Subcommittee of the University of Western Ontario Council on Animal Care (AUP no. 2008–125) and followed the Canadian Council of Animal Care guidelines. Surgical preparation and anesthesia as well as imaging acquisition protocols have been previously described [68]. Images were acquired using a 7T Siemens MAGNETOM head scanner. Two diffusion-weighted scans were acquired for each animal with each scan having an opposite phase encoding in the superior-inferior direction at 1 mm isotropic resolution, allowing for the correction of susceptibility-related distortion. The data were acquired with 2D EPI diffusion for five animals, and a multiband EPI diffusion sequence was used for the remaining three animals. In all cases, the data were acquired with $b = 1000$ s/mm^2, 64 directions, and 24 slices. A 3D T1w image was also collected for each animal (128 slices, resolution: 0.5 mm isotropic).

The diffusion-weighted images were preprocessed similarly using FSL [66]. Following preprocessing, voxel-wise estimates of the fiber orientation distribution were calculated using bedpostx [62], allowing for two fiber orientations for the macaque dataset due to the b-value in the diffusion data.

Macaque dataset 2: UC-Davis macaque data (ucd)

Data from nineteen adult rhesus macaque monkeys (all females) were available from PRIME-DE

(https://fcon_1000.projects.nitrc.org/indi/indiPRIME.html) [69]. The neuroimaging experiments and

associated procedures were performed at the California National Primate Research Center under

protocols approved by the University of California, Davis Institutional Animal Care and Use

Committee [70]. Images were acquired on a 3T Siemens Skyra scanner with four-channel clamshell

coil. Two diffusion-weighted scans were acquired for each animal with opposite phase encoding,

allowing for the correction of susceptibility-related distortion. The data were acquired with $b =$

1600 s/mm^2, 60 directions. T1w images were collected for each animal with resolution at 0.3 mm

isotropically. The preprocessing of the dataset was the same with the macaque tvb dataset.

Macaque dataset 3: High-resolution macaque data

An ultra-high resolution postmortem macaque dMRI dataset was acquired at 200 μm isotropic

voxel size on a 7 Tesla small animal MRI system equipped with 650 mT/m Resonance Research

gradient coils [31]. Details of procedures before MRI scanning can be found in previous publications

[31]. Diffusion MRI data were collected using a 3D spin-echo pulse sequence with the following

parameters: TR = 100 ms, TE = 32.1 ms, matrix size = 384 × 288 × 256, 30 DWI images

with $b = 4000$ s/mm^2 and a $b = 0$ volume were acquired.

Tractography protocols

Tractography protocols of white matter tracts

The protocols to reconstruct white matter tracts were defined as similarly as possible to those

previously defined for other primates [20-22,28] albeit adjusted for the neuroanatomical structures of

marmosets. All protocols included a midsagittal section as an exclusion mask for isolating the

tracts within one of the two hemispheres, except for the anterior commissure, forceps major,

forceps minor, and middle cerebellar peduncle, which link brain regions between the left and right

hemispheres. We here only provided descriptions of arcuate fasciculus and superior longitudinal

fasciculus, while the those of other major white matter tracts could be found in supplementary

methods.

The marmoset arcuate fasciculus was reconstructed with a seed in the PF/PFG, a posterior

target axially posterior to the terminus of the lateral fissure, and an anterior target at the level of

the ventral premotor cortex, posterior to the ventral prefrontal cortex.

For the three branches of the superior longitudinal fasciculus, a coronal mask in the white

matter underneath the primary motor cortex and primary sensorimotor cortex was used as the seed

along with the two target masks. Anteriorly, target masks for the first, second, and third branches

of the slf were coronal sections through the BA8/BA9, BA9/BA46, and BA45/BA47, respectively.

Posteriorly, target masks for the first, second, and third branches of the slf were coronal sections

through the PE, IP/PG, and PF/PFG, respectively. For each subcomponent, the seed was

positioned in alignment with the posterior target while moved anteriorly into the primary

motor/sensorimotor regions. The exclusion mask consisted of an axial mask underneath the

parietal cortex, an axial mask excluding the subcortical cortex, and a coronal mask preventing

ventral longitudinal tracts.

Tractography to reconstruct white matter tracts

Probabilistic diffusion tractography was performed using FSL's probtrackx2 [62]. The seed, target,

exclusion, and stop masks, which were defined in volumetric standard space, were warped to

individual space for tractography. The tractography was run in two directions (seed to target and

target to seed) with the following parameters: each seed voxel was sampled 10,000 times, with a

curvature threshold of 0.2, and a number of steps of 3200. The step length was set to one fourth of

voxel size according to each dataset. The resulting tractograms were normalized by dividing by

the waypoint number and warped back to template space. The normalized tractograms of tracts

were thresholded at 0.005 for individuals and then averaged to create a group-level tractogram for

each tract. The tracts were reconstructed in MNI152 standard space for humans, Yerkes29 space

for chimpanzees, F99 space for macaques, and MBMv3 space for marmosets.

Construction of connectivity patterns of tracts

We counted the number of times the tractogram of white matter tract that reached the gray/white

matter interface of each cortical region, obtaining a projection map of the tract for each species.

These projection maps were normalized by the size of seed mask size (in voxels) for each subject

of each species to allow valid cross-species comparisons.

Tractography between regions of interest

To be consistent with the tracer results, we selected cortical regions based on Paxinos atlas of

marmosets [35]. For each pair of regions of interest, i.e., A45 and AuCM, and A45 and Tpt, we

conducted diffusion tractography by using one region as the seed and the other as the target, and

vice versa, to generate two sets of tracking probability maps. The two probability maps were

averaged into a single map to represent the final map of the connection. Dorsal and ventral

waypoints were defined separately to compare the relative strength of the pathways based on the

tracking results.

Connectional comparison across four species

Construction of connectivity blueprints

We performed probabilistic tractography using FSL's probtrackx2 [62] to map the whole-brain

connectivity pattern. Specifically, the white surface was set as a seed region tracking to the rest of

the brain with the ventricles removed. The pial surface was used as a stop mask to prevent

streamlines from crossing sulci. Each vertex was sampled 10,000 times (10,000 trackings) based

on the orientation probability model for each voxel with a curvature threshold of 0.2, and a

number of steps of 3200. The step length was set to one fourth of voxel size according to each

dataset. This resulted in a (*whole-surface vertices*) $\times$ (*whole-brain voxels*) matrix for further

analysis.

The homologous white matter tracts of four species were reconstructed using XTRACT

following protocols [20,22]. Tractography of the white matter tracts was then vectorized to a (*tracts*)

$\times$ (*whole-brain voxels*) matrix and was multiplied by the whole-brain connectivity matrix, deriving

a (*tracts*) $\times$ (*whole-surface vertices*) matrix, i.e., the connectivity blueprint [21]. The matrix rows

showed each vertex's connectivity distribution pattern.

We further averaged the connectivity profiles of the vertices within subregions of atlas of

each species. Specifically, we averaged the connectivity profiles of the vertices within each

subregion of the Human Brainnetome Atlas (HumanBNA) [33], Chimpanzee Brainnetome Atlas

(ChimpBNA) [34], Macaque Brainnetome Atlas (MacBNA) [71], and Marmoset Brain Mapping Atlas

(MBM) [72] to form the regional connectivity blueprints across subjects for humans, chimpanzees,

macaques, and marmosets respectively. Note that the middle cerebellar peduncle (mcp) does not

project to the cortex. We disregarded that tract when constructing the blueprints. So, the final

tracts = 44.

Connectivity divergence between species

We used connectivity blueprints built for each species to explore the connectivity divergence across species [21,34]. We calculated the symmetric Kullback Leibler (KL) divergence for each pair of subregions in the MBM and HumanBNA and searched for the minimal value for each subregion in the HumanBNA. Specifically, for an example subregion in the HumanBNA, a connectivity blueprint represents how it was connected to each white matter tract. Using the KL divergence measure, this connectivity profile was compared with that of all the marmoset subregions. The subregion with the connectivity blueprint most similar to the human one, i.e., with the smallest KL divergence, was picked, and this KL divergence value was assigned to the human subregion. A higher value on this cross-species divergence map means that the subregion has a more dissimilar connectivity profile that is absent in marmosets. The same procedure was conducted between humans and chimpanzees and humans and macaques.

Comparison of connectivity strength of af

We compared individual connectivity strength of af with subregions across four species. Specifically, we focused on caudal area 45 (A45c), caudal area 22 (A22c), and caudal area 21 (A21c). Connectivity probability was compared between species using two-sample t test and significance was determined after correcting for multiple comparisons using Bonferroni correction.

"Virtually lesioning" the tracts in connectivity blueprints

To examine the effect of specific white matter tract on the connectivity divergence, we proposed a "virtually lesioning" approach by setting the connectivity probability values of the tract to zero and recalculating the minimum KL divergence. The difference of the connectivity divergences between species were examined by Kolmogorov-Smirnov test.

Functional alignment and tract connection comparison

Functional alignment across species

The cross-species functional alignment among marmosets, macaques, and humans were used [40,41]. Specifically, the homologous landmarks in three species were identified for subsequently extracting matched functional components using the joint-embedding approach, which were input

into multimodal surface matching (MSM) [73]. The macaque-human transformation was used as an intermediate step in constructing the cortical transformation between marmosets and humans. Chimpanzees were not considered here because their functional data was unavailable.

Alignment of functional activation

To investigate the association between functional activation by speech and the connectivity patterns of af. We selected terms including "speech", "speech perception", and "speech production" from the NeuroSynth database [39] (https://neurosynth.org/) and projected onto surface. The activation maps were transformed into marmoset surface for comparison with the connectivity patterns of af in marmosets.

Comparison of connectivity patterns of af in a common space

We compared connectivity patterns of af across species in a common human space. We transformed connectivity patterns of marmoset and macaque to human space. We calculated Dice coefficients and tract extension ratio to quantify the amount of overlap in the tract maps [18]. The Dice coefficient was computed for the binarized actual human tract map and the map transformed from the other species. The tract extension ratio was calculated as the ratio of the number of vertices covered by the human tract map to the number of vertices covered by both the human and the other tract map.

To intuitively and quantitatively compare tract connections between humans and nonhuman primates, we calculated weighted local correlation maps of the human map and the maps transformed from other species. Specifically, the local correlation map was computed using a sliding window around every vertex on the sphere, which was adjusted to up-weight the brain areas where the tract is represented on the surface. Higher values in the weighted correlation map indicated brain areas where both the tract of human and the other species show a termination.

Data availability

The tractography protocols in standard space for producing marmoset white matter tracts and their surface projection data are available online (https://github.com/FANLabCASIA/MarmosetWM). The tractography protocols for humans, chimpanzees, and macaques are available at https://git.fmrib.ox.ac.uk/rmars/chimpanzee-tractography-protocols.

The MBM dataset and the high-resolution single marmoset subject data are available at https://marmosetbrainmapping.org/data.html. The Brain/MINDS dataset is available at https://dataportal.brainminds.jp/marmoset-mri-na216. The human data are available from the Human Connectome Project (https://db.humanconnectome.org/). The high-resolution single human subject data is available at https://doi.org/10.5061/dryad.nzs7h44q2. The chimpanzee data are available at the National Chimpanzee Brain Resource (http://www.chimpanzeebrain.org/). The high-resolution single chimpanzee subject data is available from the Evolution of Brain Connectivity Project (https://doi.org/10.17617/3.O5XSI9). The TVB macaque data are available at https://doi.org/10.18112/openneuro.ds001875.v1.0.3. The UC-Davis macaque data are available at https://fcon_1000.projects.nitrc.org/indi/PRIME/ucdavis.html. The high-resolution single macaque subject data is from a previously shared resource. The Brain/MINDS Marmoset Connectivity Resource is available at https://dataportal.brainminds.jp/marmoset-tracer-injection. The Marmoset Brain Connectivity Atlas is available at https://analytics.marmosetbrain.org/graph.html. The human and marmoset Nissl-stained sections are from a previously shared resource.

Code availability

The HCP-Pipeline can be found at https://github.com/Washington-University/HCPpipelines, and the NHP-HCP-Pipeline can be found at https://github.com/Washington-University/NHPPipelines. The neuroimaging preprocessing tools used are freely available, including FreeSurfer v6.0 (http://surfer.nmr.mgh.harvard.edu/) and FSL v6.0.7 (https://fsl.fmrib.ox.ac.uk/fsl/fslwiki). The brain maps were presented using MRIcroGL v20 (https://www.nitrc.org/projects/mricrogl) and Workbench v1.5.0 (https://www.humanconnectome.org/software/connectome-workbench). The tracts were visualized using ITK-SNAP 4.0.1 (http://www.itksnap.org/) and Paraview 5.11.0

706   (https://www.paraview.org/).

Acknowledgments

We thank Kadharbatcha Saleem for valuable discussion. This work was partially supported by

STI2030-Major Projects (Grant No. 2021ZD0200203), the Natural Science Foundation of China

(Grant Nos. 82072099, 82202253, 62250058), the China Postdoctoral Science Foundation

(2022M722915, 2024M761725), and Guangxi Science and Technology Base and Talent Special

Project (Grant No. AD22035125). Data were provided in part by the National Chimpanzee Brain

Resource (supported by NIH NS092988, NIH HG011641, NIH AG067419, NIH AG087945, NSF

EF-2021785, and NSF DRL-2219759), the Duke Center for In Vivo Microscopy NIH/NIBIB (P41

EB015897), Human Connectome Project from WU-Minn Consortium (Principal Investigators:

David Van Essen and Kamil Ugurbil; 1U54MH091657) funded by the 16 NIH Institutes and

Centers that support the NIH Blueprint for Neuroscience Research and by the McDonnell Center

for Systems Neuroscience at Washington University. The authors appreciate the English language

and editing assistance of Rhoda E. and Edmund F. Perozzi, PhDs.

Competing interests

723   Authors declare no competing interests.

References

Zhang, Y., Shen, S. X., Bibic, A. & Wang, X. Evolutionary continuity and divergence of auditory dorsal and ventral pathways in primates revealed by ultra-high field diffusion MRI. *Proc Natl Acad Sci U S A* **121**, e2313831121, doi:10.1073/pnas.2313831121 (2024).

Song, X., Osmanski, M. S., Guo, Y. & Wang, X. Complex pitch perception mechanisms are shared by humans and a New World monkey. *Proc Natl Acad Sci U S A* **113**, 781-786, doi:10.1073/pnas.1516120113 (2016).

Choi, J. Y., Takahashi, D. Y. & Ghazanfar, A. A. Cooperative vocal control in marmoset monkeys via vocal feedback. *J Neurophysiol* **114**, 274-283, doi:10.1152/jn.00228.2015 (2015).

Okano, H. Current Status of and Perspectives on the Application of Marmosets in Neurobiology. *Annu Rev Neurosci* **44**, 27-48, doi:10.1146/annurev-neuro-030520-101844 (2021).

Okano, H., Hikishima, K., Iriki, A. & Sasaki, E. The common marmoset as a novel animal model system for biomedical and neuroscience research applications. *Semin Fetal Neonatal Med* **17**, 336-340, doi:10.1016/j.siny.2012.07.002 (2012).

Tendler, B. C. *et al.* The Digital Brain Bank, an open access platform for post-mortem imaging datasets. *Elife* **11**, doi:10.7554/eLife.73153 (2022).

Bryant, K. L. *et al.* Diffusion MRI data, sulcal anatomy, and tractography for eight species from the Primate Brain Bank. *Brain Struct Funct* **226**, 2497-2509, doi:10.1007/s00429-021-02268-x (2021).

Schaeffer, D. J., Liu, C., Silva, A. C. & Everling, S. Magnetic Resonance Imaging of Marmoset Monkeys. *ILAR J* **61**, 274-285, doi:10.1093/ilar/ilaa029 (2020).

Samandra, R., Haque, Z. Z., Rosa, M. G. P. & Mansouri, F. A. The marmoset as a model for investigating the neural basis of social cognition in health and disease. *Neurosci Biobehav Rev* **138**, 104692, doi:10.1016/j.neubiorev.2022.104692 (2022).

Miller, C. T. *et al.* Marmosets: A Neuroscientific Model of Human Social Behavior. *Neuron* **90**, 219-233, doi:10.1016/j.neuron.2016.03.018 (2016).

Sinnott, J. M., Stebbins, W. C. & Moody, D. B. Regulation of voice amplitude by the monkey. *J Acoust Soc Am* **58**, 412-414, doi:10.1121/1.380685 (1975).

Brumm, H., Voss, K., Kollmer, I. & Todt, D. Acoustic communication in noise: regulation of call characteristics in a New World monkey. *J Exp Biol* **207**, 443-448, doi:10.1242/jeb.00768 (2004).

Hage, S. R. Language evolution in primates. *Science* **385**, 713-714, doi:10.1126/science.ado4341 (2024).

Balezeau, F. *et al.* Primate auditory prototype in the evolution of the arcuate fasciculus. *Nat Neurosci* **23**, 611-614, doi:10.1038/s41593-020-0623-9 (2020).

Rilling, J. K. *et al.* The evolution of the arcuate fasciculus revealed with comparative DTI. *Nat Neurosci* **11**, 426-428, doi:10.1038/nn2072 (2008).

Becker, Y., Loh, K. K., Coulon, O. & Meguerditchian, A. The Arcuate Fasciculus and language origins: Disentangling existing conceptions that influence evolutionary accounts. *Neurosci Biobehav Rev* **134**, 104490, doi:10.1016/j.neubiorev.2021.12.013 (2022).

Ardesch, D. J. *et al.* Evolutionary expansion of connectivity between multimodal association areas in the human brain compared with chimpanzees. *Proc Natl Acad Sci U S A* **116**, 7101-7106, doi:10.1073/pnas.1818512116 (2019).

Eichert, N. *et al.* Cross-species cortical alignment identifies different types of anatomical

reorganization in the primate temporal lobe. *Elife* **9**, doi:10.7554/eLife.53232 (2020).

19   Roumazeilles, L. *et al.* Longitudinal connections and the organization of the temporal cortex in

macaques, great apes, and humans. *PLoS Biol* **18**, e3000810, doi:10.1371/journal.pbio.3000810 (2020).

20   Bryant, K. L., Li, L., Eichert, N. & Mars, R. B. A comprehensive atlas of white matter tracts in the

chimpanzee. *PLoS Biol* **18**, e3000971, doi:10.1371/journal.pbio.3000971 (2020).

21   Mars, R. B. *et al.* Whole brain comparative anatomy using connectivity blueprints. *Elife* **7**,

doi:10.7554/eLife.35237 (2018).

22   Warrington, S. *et al.* XTRACT - Standardised protocols for automated tractography in the human

and macaque brain. *Neuroimage* **217**, 116923, doi:10.1016/j.neuroimage.2020.116923 (2020).

23   Watakabe, A. *et al.* Local and long-distance organization of prefrontal cortex circuits in the

marmoset brain. *Neuron* **111**, 2258-2273 e2210, doi:10.1016/j.neuron.2023.04.028 (2023).

24   Skibbe, H. *et al.* The Brain/MINDS Marmoset Connectivity Resource: An open-access platform

for cellular-level tracing and tractography in the primate brain. *PLoS Biol* **21**, e3002158,

doi:10.1371/journal.pbio.3002158 (2023).

25   Majka, P. *et al.* Towards a comprehensive atlas of cortical connections in a primate brain:

Mapping tracer injection studies of the common marmoset into a reference digital template. *J Comp*

Neurol **524**, 2161-2181, doi:10.1002/cne.24023 (2016).

26   Majka, P. *et al.* Open access resource for cellular-resolution analyses of corticocortical

connectivity in the marmoset monkey. *Nat Commun* **11**, 1133, doi:10.1038/s41467-020-14858-0

(2020).

27   Liu, C. *et al.* A resource for the detailed 3D mapping of white matter pathways in the marmoset

brain. *Nat Neurosci* **23**, 271-280, doi:10.1038/s41593-019-0575-0 (2020).

28   Bryant, K. L. *et al.* A map of white matter tracts in a lesser ape, the lar gibbon. *Brain Struct Funct*,

doi:10.1007/s00429-023-02709-9 (2023).

29   Eichner, C. *et al.* Detailed mapping of the complex fiber structure and white matter pathways of

the chimpanzee brain. *Nat Methods* **21**, 1122-1130, doi:10.1038/s41592-024-02270-1 (2024).

30   Wang, F. *et al.* In vivo human whole-brain Connectom diffusion MRI dataset at 760 microm

isotropic resolution. *Sci Data* **8**, 122, doi:10.1038/s41597-021-00904-z (2021).

31   Calabrese, E. *et al.* Investigating the tradeoffs between spatial resolution and diffusion sampling

for brain mapping with diffusion tractography: time well spent? *Hum Brain Mapp* **35**, 5667-5685,

doi:10.1002/hbm.22578 (2014).

32   Hata, J. *et al.* Multi-modal brain magnetic resonance imaging database covering marmosets with a

wide age range. *Sci Data* **10**, 221, doi:10.1038/s41597-023-02121-2 (2023).

33   Fan, L. *et al.* The Human Brainnetome Atlas: A New Brain Atlas Based on Connectional

Architecture. *Cereb Cortex* **26**, 3508-3526, doi:10.1093/cercor/bhw157 (2016).

34   Wang, Y. *et al.* The Chimpanzee Brainnetome Atlas reveals distinct connectivity and gene

expression profiles relative to humans. *Innovation (Camb)* **6**, 100755, doi:10.1016/j.xinn.2024.100755

(2025).

35   Paxinos, G., Watson, C., Petrides, M., Rosa, M. G. & Tokuno, H. *The Marmoset Brain in*

Stereotaxic Coordinates.   (Elsevier Academic Press, 2012).

36   Mai, J. K., Majtanik, M. & Paxinos, G. *Atlas of the human brain*.   (Academic press, 2015).

37   Amunts, K. *et al.* Broca's region revisited: cytoarchitecture and intersubject variability. *J Comp*

Neurol **412**,   319-341,   doi:10.1002/(sici)1096-9861(19990920)412:2<319::aid-cne10>3.0.co;2-7

(1999).

38   Palomero-Gallagher, N. & Zilles, K. Differences in cytoarchitecture of Broca's region between
human, ape and macaque brains. *Cortex* **118**, 132-153, doi:10.1016/j.cortex.2018.09.008 (2019).

39   Yarkoni, T., Poldrack, R. A., Nichols, T. E., Van Essen, D. C. & Wager, T. D. Large-scale
automated synthesis of human functional neuroimaging data. *Nat Methods* **8**, 665-670,
doi:10.1038/nmeth.1635 (2011).

40   Xu, T. *et al.* Cross-species functional alignment reveals evolutionary hierarchy within the
connectome. *Neuroimage* **223**, 117346, doi:10.1016/j.neuroimage.2020.117346 (2020).

41   Xu, T. *et al.* in *The Organization for Human Brain Mapping (OHBM) 2023 Annual Meeting*
(Montreal, Canada, 2023).

42   Dureux, A., Zanini, A. & Everling, S. Mapping of facial and vocal processing in common
marmosets with ultra-high field fMRI. *Commun Biol* **7**, 317, doi:10.1038/s42003-024-06002-1 (2024).

43   Miller, C. T., Thomas, A. W., Nummela, S. U. & de la Mothe, L. A. Responses of primate frontal
cortex neurons during natural vocal communication. *J Neurophysiol* **114**, 1158-1171,
doi:10.1152/jn.01003.2014 (2015).

44   Van Essen, D. C. *et al.* Cerebral cortical folding, parcellation, and connectivity in humans,
nonhuman primates, and mice. *Proc Natl Acad Sci U S A* **116**, 26173-26180,
doi:10.1073/pnas.1902299116 (2019).

45   Osmanski, M. S. & Wang, X. Perceptual specializations for processing species-specific
vocalizations in the common marmoset (Callithrix jacchus). *Proc Natl Acad Sci U S A* **120**,
e2221756120, doi:10.1073/pnas.2221756120 (2023).

46   Cheney, D. L. & Seyfarth, R. M. Flexible usage and social function in primate vocalizations. *Proc*
Natl Acad Sci U S A **115**, 1974-1979, doi:10.1073/pnas.1717572115 (2018).

47   Zhao, L. & Wang, X. Frontal cortex activity during the production of diverse social
communication calls in marmoset monkeys. *Nat Commun* **14**, 6634, doi:10.1038/s41467-023-42052-5
(2023).

48   Roy, S., Miller, C. T., Gottsch, D. & Wang, X. Vocal control by the common marmoset in the
presence of interfering noise. *J Exp Biol* **214**, 3619-3629, doi:10.1242/jeb.056101 (2011).

49   Oren, G. *et al.* Vocal labeling of others by nonhuman primates. *Science* **385**, 996-1003,
doi:10.1126/science.adp3757 (2024).

50   Jiang, Y. *et al.* Constructing the hierarchy of predictive auditory sequences in the marmoset brain.
Elife **11**, doi:10.7554/eLife.74653 (2022).

51   Clery, J. C., Hori, Y., Schaeffer, D. J., Menon, R. S. & Everling, S. Neural network of social
interaction observation in marmosets. *Elife* **10**, doi:10.7554/eLife.65012 (2021).

52   Dureux, A., Zanini, A., Selvanayagam, J., Menon, R. S. & Everling, S. Gaze patterns and brain
activations in humans and marmosets in the Frith-Happe theory-of-mind animation task. *Elife* **12**,
doi:10.7554/eLife.86327 (2023).

53   Bryant, K. L. *et al.* Connectivity profile and function of uniquely human cortical areas. *bioRxiv*,
doi:10.1101/2024.06.20.599486 (2024).

54   Sierpowska, J. *et al.* Comparing human and chimpanzee temporal lobe neuroanatomy reveals
modifications to human language hubs beyond the frontotemporal arcuate fasciculus. *Proc Natl Acad*
Sci U S A **119**, e2118295119, doi:10.1073/pnas.2118295119 (2022).

55   Braunsdorf, M. *et al.* Does the temporal cortex make us human? A review of structural and

functional diversity of the primate temporal lobe. *Neurosci Biobehav Rev* **131**, 400-410,
doi:10.1016/j.neubiorev.2021.08.032 (2021).

56 Schmahmann, J. D. & Pandya, D. N. *Fiber Pathways of the Brain*. (Oxford University Press,
2006).

57 Gattoni, G. & Tosches, M. A. Constrained roads to complex brains. *Science* **387**, 716-717,
doi:10.1126/science.adv2609 (2025).

58 Tian, X. *et al.* An integrated resource for functional and structural connectivity of the marmoset
brain. *Nat Commun* **13**, 7416, doi:10.1038/s41467-022-35197-2 (2022).

59 Silva, A. C. *et al.* Longitudinal functional magnetic resonance imaging in animal models. *Methods*
Mol Biol **711**, 281-302, doi:10.1007/978-1-61737-992-5_14 (2011).

60 Papoti, D. *et al.* Design and implementation of embedded 8-channel receive-only arrays for
whole-brain MRI and fMRI of conscious awake marmosets. *Magn Reson Med* **78**, 387-398,
doi:10.1002/mrm.26339 (2017).

61 Liu, C. *et al.* Marmoset Brain Mapping V3: Population multi-modal standard volumetric and
surface-based templates. *Neuroimage* **226**, 117620, doi:10.1016/j.neuroimage.2020.117620 (2021).

62 Behrens, T. E., Berg, H. J., Jbabdi, S., Rushworth, M. F. & Woolrich, M. W. Probabilistic
diffusion tractography with multiple fibre orientations: What can we gain? *Neuroimage* **34**, 144-155,
doi:10.1016/j.neuroimage.2006.09.018 (2007).

63 Hata, J., Nakae, K., Yoshimaru, D. & Okano, H. Brain/MINDS Marmoset Brain MRI Dataset
NA216 and eNA91: (DataID: 4624). doi:10.24475/bminds.mri.thj.4624.

64 Van Essen, D. C. *et al.* The WU-Minn Human Connectome Project: an overview. *Neuroimage* **80**,
62-79, doi:10.1016/j.neuroimage.2013.05.041 (2013).

65 Glasser, M. F. *et al.* The minimal preprocessing pipelines for the Human Connectome Project.
Neuroimage **80**, 105-124, doi:10.1016/j.neuroimage.2013.04.127 (2013).

66 Jenkinson, M., Beckmann, C. F., Behrens, T. E., Woolrich, M. W. & Smith, S. M. Fsl. *Neuroimage*
62, 782-790, doi:10.1016/j.neuroimage.2011.09.015 (2012).

67 Chen, X. *et al.* Brain aging in humans, chimpanzees (Pan troglodytes), and rhesus macaques
(Macaca mulatta): magnetic resonance imaging studies of macro- and microstructural changes.
Neurobiol Aging **34**, 2248-2260, doi:10.1016/j.neurobiolaging.2013.03.028 (2013).

68 Shen, K. *et al.* A macaque connectome for large-scale network simulations in TheVirtualBrain. *Sci*
Data **6**, 123, doi:10.1038/s41597-019-0129-z (2019).

69 Milham, M. P. *et al.* An Open Resource for Non-human Primate Imaging. *Neuron* **100**, 61-74 e62,
doi:10.1016/j.neuron.2018.08.039 (2018).

70 Baxter, M. G., Santistevan, A. C., Bliss-Moreau, E. & Morrison, J. H. Timing of cyclic estradiol
treatment differentially affects cognition in aged female rhesus monkeys. *Behav Neurosci* **132**, 213-223,
doi:10.1037/bne0000259 (2018).

71 Lu, Y. *et al.* Macaque Brainnetome Atlas: A multifaceted brain map with parcellation, connection,
and histology. *Sci Bull (Beijing)*, doi:10.1016/j.scib.2024.03.031 (2024).

72 Liu, C. *et al.* A digital 3D atlas of the marmoset brain based on multi-modal MRI. *Neuroimage*
169, 106-116, doi:10.1016/j.neuroimage.2017.12.004 (2018).

73 Robinson, E. C. *et al.* Multimodal surface matching with higher-order smoothness constraints.
Neuroimage **167**, 453-465, doi:10.1016/j.neuroimage.2017.10.037 (2018).