Evaluation of nanopore sequencing for increasing accessibility of eDNA studies

in biodiverse countries

Daniel Gygax[1,2,3] *, Sabina Ramirez[1], Moses Chibesa[4], Twakundine Simpamba[5], Michael

Riffel[1], Tom Riffel[6], Geofrey Zulu[6], Amrita Srivathsan[7], Reindert Nijland[8], Lara Urban[2,3,9]*

1Faunomics, Rifcon GmbH, Hirschberg, Germany

2Technical University of Munich, School of Life Sciences, Freising, Germany

3Helmholtz Munich, Neuherberg, Germany

4Department of Zoology and Aquatic Sciences, The Copperbelt University, Kitwe, Zambia

5Department of National Parks and Wildlife, Chilanga, Zambia

6Nsanga Conservation, Zambia

7Center for Integrative Biodiversity Discovery, Museum für Naturkunde, Berlin, Germany

8Marine Animal Ecology Group, Wageningen University, Wageningen, The Netherlands

9Institute for Food Safety and Hygiene, University of Zürich, Zürich, Switzerland

*Corresponding authors: dgygax90@gmail.com & lara.h.urban@gmail.com

Abstract

Biodiversity loss is a global challenge of the 21st century. Environmental DNA (eDNA)-based

metabarcoding offers a cost- and time-efficient alternative to conventional biodiversity

surveys, enabling detection of rare, cryptic, and elusive species from environmental samples.

However, limited access to genomic technologies restricts the application of eDNA

metabarcoding in highly biodiverse remote regions and low- and middle-income countries

(LMICs). Here, we directly compared the latest portable nanopore sequencing methods with

established Illumina sequencing for vertebrate eDNA metabarcoding of Zambian water

samples. Our results show that due to recent improvements in sequencing chemistry and

optimized basecalling, nanopore sequencing data can recapitulate or even surpass

established protocols, demonstrating the feasibility of *in situ* biodiversity assessments. eDNA-

and camera trap-based species detections had minimal overlap in species detections,

suggesting a complementary rather than substituting application of these biodiversity

monitoring technologies. We finally demonstrate that our entire eDNA workflow can be

successfully implemented in a mobile laboratory under remote field conditions by completing

all steps—from sample collection to data analysis—within the Luambe National Park in

Zambia. This approach has important implications for capacity building in LMICs and for

overcoming limitations associated with sample export.

Keywords

eDNA, portable sequencing, vertebrates, Zambia, metabarcoding

Introduction

Biodiversity loss is one of the major challenges of the 21st century [1], but efforts to prevent biodiversity loss are often hindered by a lack of knowledge of species distributions and dynamics [2,3]. Environmental DNA (eDNA) approaches can provide a cost- and time-efficient alternative to conventional biodiversity monitoring methods by detecting taxa, including rare, cryptic, and elusive species, from metabarcoding their genetic material in environmental samples such as water, soil, or air, and subsequent amplicon sequencing [4-6]. Such eDNA approaches have recently become cheaper and more reliable in monitoring the temporal and spatial distribution and dynamics of species, populations, and communities [7][8][9]. The lack of access to genomic technology, however, hinders the application of metabarcoding approaches in remote areas and low- and middle-income countries (LMICs), which often contain the globally largest amount of biodiversity [10-12]. Simultaneously, international treaties such as the Convention on International Trade in Endangered Species of Wild Fauna and Flora [13] and data sovereignty principles as specified by the Nagoya Protocol on Access and Benefit Sharing [14] and the Global Indigenous Data Alliance [15] often require the generation and usage of such genomic data close to the species' origin.

Long-read nanopore sequencing technology as developed by Oxford Nanopore Technologies (ONT) is the first genomic technology that enables rapid *in situ* sequencing at low upfront investment costs through its portable devices. This technology therefore holds the promise of putting eDNA research in line with international treaties and data sovereignty principles, while saving time and resources otherwise spent on sample transport and exportation, shortening the gap between research and decision-making and potentially empowering local researchers, conservationists, and decision-makers [11,16]. First applications of nanopore sequencing to metabarcoding in general [17-19] and eDNA specifically [20,21] have shown promising results in species but in comparison to the established sequencing platform for metabarcoding, short-read Illumina sequencing, remained affected by the relatively high sequencing error rates. Bludau *et al.* have, however, also shown that this lack in sequencing accuracy can be

compensated by nanopore sequencing's long-read capability, which enabled species-level

resolution in their protistan studies by substantially increasing the targeted amplicon size.

While the error rate of this previously applied nanopore sequencing chemistry ("R9" chemistry)

was estimated at anywhere between 5% and 22% [17,19], recent substantial improvements

in the nanopore design ("R10" chemistry) and basecalling algorithms now allow for highly

accurate nanopore read- and consensus-level accuracy. Doorenspleet *et al.* were the first to

use these accuracy improvements to apply nanopore-based eDNA to successfully monitor

marine vertebrates from aquarium and sea water. Similarly, Bludau *et al*. showed that an

increase of the targeted amplicon size also improves species detection when using the latest

nanopore chemistry; however, fragmentation of the marine vertebrate's DNA in real-world

environments such as the sea inhibits the amplification of long stretches of DNA [22].

Here, we conducted an eDNA study to directly compare the latest improvements in portable

nanopore sequencing with established Illumina sequencing as well as camera trapping for

monitoring terrestrial vertebrate diversity from freshwater samples in Zambia. We additionally

tested the *in situ* implementation of eDNA methods in a mobile laboratory setting by applying

our complete eDNA workflow from water collection to data analysis on site in Luambe National

Park, Zambia. Zambia serves as a good example of a biodiversity-rich LMIC that shares

borders and ecosystems with eight other African countries and is estimated to be home to 242

mammals, 757 bird, 74 amphibian and 156 reptiles [23]. Zambia has further been described

as "a large patchwork of important ecosystems" [25]., with its subtropical savanna ecosystems

being characterized by extreme oscillations of precipitation and water availability during the

wet and dry seasons [24,25]. This seasonality results in an increased reliance on mobility for

both humans and animals to access water [25] and therefore imposes challenges for

monitoring highly mobile vertebrates across large areas. In arid and semi-arid systems where

surface water is often scarce, natural and artificial water holes further act as gathering sites

for mammals and birds [26], making them potentially optimal sources of eDNA for monitoring

vertebrate fauna [27-29].

Materials and Methods

Environmental DNA collection

For sequencing comparisons, we took water samples at five locations in the Luambe and Lukusuzi National Parks within the Luangwa Valley in Eastern Zambia between July 13th and 16th, 2023 (Fig A-F). In Luambe National Park, three water bodies were sampled: an artificial pond near an established tourism facility (CAM: 12.45880° S, 32.14662° E) (Fig 1B) and two permanent natural ponds (P2: 12.48503° S, 32.17606° E; P3: 12.48246° S, 32.19048° E) (Fig 1C-D). In Lukusuzi National Park, samples were collected from two sites along the Lukusuzi River (L1: 12.71192° S, 32.50254° E; L2: 12.71247° S, 32.50226° E) (Fig 1E-F). These locations were chosen because previous camera trap-based monitoring had identified them as hotspots of vertebrate activity (Fig 1A-F). At each water body, three biological replicates (A, B, and C) were collected. For the *in situ* application (see "*In situ* eDNA application"), freshwater samples were collected at three sites of the Mwendi Lagoon (Mw: 12.53318° S, 32.1435° E) in Luambe National Park between June 11th and 16th, 2025 (Fig 1G).

Fig 1. Map and pictures of study sites. A) Map showing the sampling locations within Luambe and Lukusuzi National Parks (Mw: blue dot) ; B) CAM: Artificial pond in Luambe

Camp; C) P2: Natural waterhole 1 in Luambe NP; D) P3: Natural waterhole 2 in Luambe NP; E) L1: Lukuzusi backwater site 1; F) L2: Lukuzusi backwater site 2; G) Mw: Mwendi Lagoon (sampling site for the *in situ* application). Basemap: Esri, National Geographic Style (accessed September 2025) via QGIS.

We collected freshwater samples by using sterile plastic scoops and storing them in sterile Whirl-Pak® bags (Thermo Scientific, Germany) for transport to the camp. At the camp, one liter of water per sample was filtered using 250ml cups with 0.45 µm nitrocellulose filters (Thermo Scientific Nalgene, Germany) and a Vampire Vacuum Pump (Bürkle, Germany) to speed up the process. For samples with high sediment content, multiple nitrocellulose filters were required to filter the entire liter due to clogging; for these samples, final DNA extracts were subsequently pooled across filters per replicate. After filtration, the nitrocellulose filters were first stored in sterile kraft bags and then placed in sterile ziplock bags with silica gel beads. The samples were frozen at -20ºC before and after transportation from Zambia to Germany. For the sequencing comparisons, all molecular work was subsequently conducted at the Helmholtz Center Munich laboratories in Germany under the corresponding research permits from the Zambian Department of National Parks and Wildlife (Reference number: NPW/8/27/1).

Environmental DNA extraction

The nitrocellulose filters were cut in half using a sterile scalpel, sterile gloves, and a FFP2 mask, then stored separately in 1.5 ml Eppendorf tubes at -20°C until further processing; half of a filter was then used for DNA extraction while the other half was kept as a backup. DNA was extracted from all samples using the DNeasy PowerSoil Kit (Qiagen) following the manufacturer's instructions, with the exception that DNA was eluted in only 50 µl of elution buffer in the final step.

For the filter negative control, one liter of sterile Milli-Q® water was filtered. Together with all biological replicates across all sampling sites, this resulted in 16 DNA extracts, whose DNA

concentrations were measured with an Invitrogen™ Qubit™ 3.0 Fluorometer using the High

Sensitivity (HS) Assay.

PCR amplification and metabarcoding

Metabarcoding was performed using two mitochondrial primer sets. The vertebrate-specific

12S rRNA primer pair 12SV05 (forward 5′-TTAGATACCCC ACTATGC-3′; reverse 5′-

TAGAACAGGCT CCTCTAG-3′) amplifies a ~97 bp fragment of the 12S rRNA gene [30].

Mammals were further targeted with the 16S rRNA primer pair 16Smam1 (forward 5′-

CGGTTGGGGTGACCTCGGA-3′) and 16Smam2 (reverse 5′-

GCTGTTATCCCTAGGGTAACT-3′), which amplifies a ~95 bp fragment of the 16S rRNA gene

[31].

Each sample extract was amplified in PCR triplicates, and a filter negative control in two PCR

replicates, plus one PCR control, resulting in overall 48 PCR products. The PCR reactions

were set up in 25 µL volumes consisting of 2 µL DNA template, 12.5 µL Phusion® High-Fidelity

PCR Master Mix, 1.25 µL of each primer, and 8 µL of DNase-free water. For the 12S rRNA

primer pair the cycling parameters were as follows: 95°C for 10 minutes, followed by 40 cycles

of 94°C for 30 s, 60°C for 30 s, and 70°C for 60 s, with a final extension at 72°C for 7 min. For

the 16S rRNA primer pair, the cycling parameters were as follows: 95°C for 10 min, followed

by 40 cycles of 95°C for 12 s, 59°C for 30 s and 70°C for 25 s, with a final extension at 72 °C

for 7 min.

A symmetrical tagged primer approach was used to multiplex the samples during amplification.

Unique 9 bp tags, generated using Barcode Generator

(https://github.com/lcomai/barcode_generator), were added to the 5′ ends of both the forward

and reverse primers for each primer set. This tag length has been shown to be suitable for

multiplexing amplicons for nanopore sequencing [32]. All tags were designed to differ by at

least 3 bp from each other, and the same tag was used on both the forward and reverse

primers of each PCR product.

Following amplification, one replicate PCR product was respectively visualized on 2% agarose

gels with GelRed and Invitrogen™ 100 bp DNA. Amplification was confirmed by the presence

of a band of the expected size, which was the case for all visualized samples (including the

negative controls).

Sequencing library preparation and sequencing

The 48 PCR products were pooled by taking 10 µl of each product and then cleaned using a

2:1 bead-to-DNA ratio with AMPure XP Beads (Beckman Coulter, Germany), followed by

resuspension in 100 µL of DNase-free water. This final pool was separated into two 50 µL

volumes as input for nanopore and Illumina sequencing, respectively.

Nanopore sequencing was performed using the portable MinION Mk1B device. Library

preparation was done using the Ligation Sequencing Kit (SQK-LSK114), following the

manufacturer's Ligation Sequencing Kit Amplicons protocol with the following modification: At

the end of the protocol, 30 µL of elution buffer was used instead of the specified 15 µL. This

was done to simultaneously prepare two libraries, which was possible due to high DNA input

into the library preparation. Both libraries were consecutively sequenced for 16 hours on one

flow cell, separated by a flow cell wash step.

Illumina sequencing was performed by Novogene, GmbH, Germany, using their 2x150 bp

protocol on a NovaSeq 6000.

Data processing and OTU clustering

Nanopore raw data was basecalled using Dorado's (v1.0.2) super high accuracy model (SUP

v5.0.0; https://github.com/nanoporetech/dorado). The resulting FASTQ files were

demultiplexed using OBITools4's obimultiplex command, allowing for a maximum of two errors

in the tags (https://git.metabarcoding.org/obitools/obitools4). Primers were trimmed using

Cutadapt (v4.9) [33]. Subsequent Operational Taxonomic Unit (OTU) processing steps were

performed using VSEARCH (v2.28.1) [33]: We first filtered the reads by setting the VSEARCH

fastq_maxee parameter to thresholds of 1.0: For a 97 bp sequence, a fastq_maxee value of

1.0 corresponds to average per-base quality scores of approximately 20 using the formula: Q

$= -10 \times \log_{10}(\text{fastq\_maxee} \div L)$, where Q is the average per-base Phred quality score and L is

the read length. Therefore, lower fastq_maxee thresholds require higher average per-base

quality scores to keep the expected error per read below the specified limit. The next steps

included dereplication (with singleton removal), chimera removal, and clustering at a 99%

sequence identity threshold; the final OTUs were pooled across respective PCR replicates.

Illumina paired-end data were first merged and then demultiplexed with Obitools 4 allowing for

a maximum of two errors in the tags. Primers were trimmed using Cutadapt (v4.9) [33].

Subsequent OTU processing steps were identical to the OTU processing steps of the

nanopore sequencing data.

Taxonomic assignments

Amphibian, bird, mammalian, and bird species lists reported in Zambia were obtained from

the   SASCAL   portal

(http://data.sasscal.org/metadata/view.php?view=doc_documents&id=3154, 2017). This full

species list (S1 Table) was used as a list of regional species, and to assess our study's

database incompleteness database (S2 Table).

Taxonomic assignment of the OTUs was done using the global alignment function

usearch_global VSEARCH (v2.28.1) [34] against the Midori2 databases Unique Nucleotide

short ribosomal RNA version 266 (MIDORI2_UNIQ_NUC_GB266_srRNA_BLAST) and the

Unique   Nucleotide   long   ribosomal   RNA   version   266

(MIDORI2_UNIQ_NUC_GB266_lrRNA_BLAST), which contain curated non-redundant 12S

and 16S rRNA sequences, respectively [35]. Only the best taxonomic hits with at least 90%

sequence similarity score and 80% query coverage score were taken and then filtered for local

taxa detections based on the list of regional species: Assignments with ≥98% identity to a

reference sequence from Zambia were considered reliable species-level identifications:

matches between 95–98% identity were assigned at the genus level, while those between 90–95% identity were conservatively classified at the family level. We only focused our subsequent analyses on primary non-terrestrial vertebrate classes. Taxa detected in the negative controls were removed from the data.

Camera Trapping

One week prior to eDNA sampling, camera traps (SECACAM Wild-Vision Full HD 5.0) were put up at sampling locations to validate species detections. Cameras, equipped with passive infrared sensors, were set to picture-mode and recorded for 24 hours. The number of deployed camera traps varied from 2 to 5 cameras across sites, depending on the size and accessibility of the water bodies. Cameras were positioned to cover the entire waterbodies with priority on established drinking places by large mammals. Drinking places were determined by visual assessments on mammal tracks, scat, or further signs of presence.

***In situ* eDNA application**

We finally conducted our entire protocol including sample collection, molecular work, and bioinformatic analyses on site in the Nsanga Research Camp within Luambe National Park. Freshwater samples were collected daily from three sites (A–C) spaced ~30 m apart at the Mwendi Lagoon (Mw: 12.53318° S, 32.1435° E) in Luambe National Park from June 11[th] to 16[th]. For the molecular work, we used a mobile laboratory setting assembled within a camping tent (S4 Appendix). For the bioinformatic analyses, we used a high-performing laptop (Razer Blade 15.6″ with 16 GB RAM, an NVIDIA RTX 4070 GPU, and an Intel Core i7 processor). We again used camera trapping in parallel (see "Camera Trapping"), with one camera per sampling site and daily assessments from sampling day 2.

We implemented the following modifications of our nanopore sequencing-based eDNA protocol for the *in situ* approach:

1) To reduce contamination risk, the following protocol steps were modified: Instead of

using 0.45 µm nitrocellulose filters and centralized processing by a vacuum pump in

the camp, Smith and Root™ eDNA Filter Packs were used to filter three liters of

wateron site while minimizing handling. We used PES (Polyethersulfone) filters with a

larger pore size (1.2 µm) to reduce clogging when filtering larger volumes of water.

With this approach it took between 15 min and 25 minutes to filter 3L of water. The

filters were stored in sterile bags and stored at -20ºC until DNA extraction.

2) We used the DNeasyPowerWater Kit (Qiagen) for DNA extraction following the

manufacturer's instructions, with the exception that DNA was eluted in 50 µl of elution

buffer in the final step. We used this kit instead of the DNeasy PowerSoil Kit (Qiagen)

since it allows to process the entire filter in a faster manner. For the filter negative

controls, we used blank filters instead of filtering sterile Milli-Q® water due to lack of

access to such volumes of sterile water. We used one blank filter for each of the three

DNA extractions of six samples. This resulted in 21 DNA extracts, whose DNA

concentrations were measured with an onsite Invitrogen™ Qubit™ 3.0 Fluorometer

using the High Sensitivity (HS) Assay.

3) We incorporated human blockers in the PCR reaction to reduce human contamination.

The PCR reactions changed but the same symmetrical tagging approach as described

previously was used for sample multiplexing. Specifically, the PCR reactions were set

up in 20 µL volumes consisting of 3 µL DNA template, 0.75 U AmpliTaq Gold, 1× Gold

PCR Buffer and 2.5 mM $MgCl_2$ (all reagents from Applied Biosystems); 0.6 µM each of

5′ nucleotide tagged forward and reverse primer; 0.2 mM dNTP mix (Invitrogen);

0.5 mg/mL bovine serum albumin (BSA, Bio Labs), and 3 µM human blocker (5′–3′

TACCCCACTATGCTTAGCCCTAAACCTCAACAGTTAAATC–spacerC3 for the 12S

rRNA primers [36] and 5′–3′ GCGACCTCGGAGCAGAACCC–spacerC3 for the 16S

rRNA primerss [37]). For the 12S rRNA PCR protocol, the number of cycles was

increased from 40 to 45 cycles, and instead of the 60°C we used 59°C. For the 16S

rRNA PCR protocol, instead of the 59 °C we used 51 °C. We confirmed amplification

263   using 2% EX pre-cast agarose gels run with the E-Gel™ Power Snap Electrophoresis

264   (Thermo Fischer Scientific). For the PCR pool clean-up, we further changed the

265   bead:sample ratio from 2:1 to 1.5:1 to reduce primer-dimmer carryover.

266  4) Nanopore sequencing was performed using the portable MinION Mk1D device with

267   better temperature control than the Mk1B device.

268 **Results**

269 Read counts varied widely across samples for both sequencing technologies and primer set.

270 For Illumina 12S samples ranged from 3 to 161,352 reads per sample (mean = 48,459; SD =

271 41,615). For Illumina 16S yields ranged from 448 to 470,489 (mean = 121,653; SD = 89,537).

272 For Nanopore 12S yields spanned 18 to 2,424,297 (mean = 273,710; SD = 543,585). For

273 Nanopore 16S yields ranged from 2,029 to 1,433,579 (mean = 227,434; SD = 309,771).

274 Our database revision revealed that the availability of genetic references for Zambian

275 biodiversity in the NCBI repository was limited across all primary terrestrial vertebrate classes

276 58.14% of occurring amphibian species were covered by 12S rRNA entries, 68.60% by 16S

277 rRNA, but only 6.98% by full mitochondrial references. 42.78% of reptile species were covered

278 by 12S rRNA entries, 64.43% by 16S rRNA, but only 3.09% by full mitochondrial references.

279 28.38% of avian species were covered by 12S rRNA entries, 16.43% by 16S rRNA, and

280 15.64% by full mitochondrial references. Finally, mammals were relatively well represented

281 with 58.80% (12S rRNA), 53.60% (16S rRNA) and 57.60% (full mitochondria), respectively.

282 All four primary terrestrial vertebrate classes were detected by nanopore and Illumina

283 sequencing; as *Xenopus* sp*.* was detected by negative controls, it was excluded from

284 downstream analyses. For both sequencing technologies the highest number of local species

285 detections occurred for mammals (17 and 11 species, respectively; Fig 2). Altogether, 18 taxa

286 were detected by both sequencing technologies while seven taxa were only detected by

287 nanopore, and one taxon only by Illumina sequencing (Fig 2). The two primer sets showed

288 little overlap in amplified taxa, with the 12s rRNA primer set yielding 21 taxa and the 16s rRNA

primers 9 taxa, with only four of them in common. Most taxa were detected at the CAM site

(n=15) followed by P3 and L1 (n=7, respectively), and L2 (n=6) and P2 (n=5). While most sites

showed little overlap in detected taxa, a few common species like *Hippopotamus amphibius*

*(*Hippopotamus*), Chlorocebus pygerythrus* (Vervet monkey) and *Papio cynocephalus* (Yellow

baboon) were detected at nearly all sites (Fig 2).

Taxonomic ranks				Sampling sites				
Class	**Order**	**Family**	**Species**	**CAM**	**P2**	**P3**	**L1**	**L2**
Amphibians	Anura	Ptychadenidae	*Ptychadena anchietae*	▲				
Birds	Charadriiformes	Charadriidae	*Vanellus armatus*		●			
	Anseriformes	Anatidae	*Anas sp.*					●
	Bucerotiformes	Bucorvidae	*Bucorvidae sp.*	●				
	Cuculiformes	Cuculidae	*Cuculidae sp.*	●				
	Accipitriformes	Accipitridae	*Accipitridae sp.*	●				
	Psittaciformes	Psittacidae	*Poicephalus sp.*	●				
	Anseriformes	Anatidae	*Sarkidiornis melanotos*	▲				
Mammals	Artiodactyla	Hippopotamidae	*Hippopotamus amphibius*	●▲	●▲	●▲	●▲	●▲
	Artiodactyla	Bovidae	*Syncerus caffer*		▲			
	Artiodactyla	Bovidae	*Tragelaphus sp.*	●		●		●
	Artiodactyla	Bovidae	*Tragelaphus scriptus*			▲		
	Carnivora	Herpestidae	*Helogale parvula*			●▲	●▲	
	Carnivora	Felidae	*Felidae sp.*	●		●	●	
	Carnivora	Herpestidae	*Herpestes sp.*				●	
	Carnivora	Viverridae	*Viverridae sp.*				●	
	Chiroptera	Vespertilionidae	*Afronycteris nanus*	●▲				
	Chiroptera	Vespertilionidae	*Vespertilionidae sp.*	▲				
	Chiroptera	Rhinolophidae	*Rhinolophus sp.*	●		●		
	Primates	Cercopithecidae	*Chlorocebus pygerythrus*	●▲	●▲	●▲	●▲	
	Primates	Cercopithecidae	*Papio cynocephalus*	▲	▲		▲	▲
	Rodentia	Muridae	*Muridae sp.*		●			
	Rodentia	Gliridae	*Graphiurus kelleni*					●
	Rodentia	Gliridae	*Graphiurus microtis*					●
	Rodentia	Muridae	*Pelomys fallax*	▲				
Reptiles	Testudines	Pelomedusidae	*Pelusios sinuatus*					●

Legend			
Marker	12S	●	
	16S	▲	
	Both	●▲	
Seq. tech	Nanopore		
	Illumina		
	Both		

Fig 2. Primary terrestrial vertebrate taxa detected. Detections are depicted across the five Zambian

sampling sites eDNA primers sets (12S or 16S rRNA, or both) and sequencing technology (nanopore

and Illumina sequencing, or both).

We next compared all our taxa detections between the different sequencing technologies and

camera trapping (Fig 3.). Briefly, camera trapping-based detection was biased towards birds

and mammals with 10 and 9 species detected, respectively, and no detections for amphibians

or reptiles (S3 Table) while eDNA-based methods yielded 25 detections across all four primary

terrestrial vertebrates, out of which 14 were at the species level and 5 at the genus and 7 at

the family level, respectively. We observed little overlap between the taxa detected by camera

traps and eDNA; this overlap increased slightly with taxonomic ranks (Fig 3).

Fig 3. Shared taxa detections between camera trap-based biodiversity monitoring and freshwater-based eDNA-based monitoring using nanopore and Illumina sequencing. Taxonomic A) species, B) genus, and C) family detections. D) Examples of camera trap pictures: *left*: Blacksmith lapwing (*Vanellus armatus*) detected by all three approaches; *middle*: Hippopotamus (*Hippopotamus amphibius*) detected by all three approaches; and *right*: Small-spotted Genet (*Genetta genetta*) detected by camera trapping and by nanopore sequencing on the family level (*Viverridae sp.*).

We finally validated a full mobile laboratory workflow based on nanopore sequencing under field conditions (Methods). Over six consecutive sampling days at three sites of Mwendi Lagoon, we detected ten primary terrestrial vertebrate taxa using eDNA, and seven with camera traps, with four of the taxa detected by both methods (Fig 4A). Within the eDNA detections, nine taxa were recovered with the 12S rRNA primer, and four with the 16S rRNA primer, including three taxa amplified by both primers (Fig 4B). No local species was detected in the negative controls. Taxonomic richness varied across the three sites: Site A yielded the highest richness by eDNA approaches (n = 9), followed by sites B and C (n = 4 and n = 3, respectively); more taxa could be detected by eDNA than by camera trapping at A and C, and *vice versa* at B (Fig 4C).

Fig 4. Number of primary terrestrial vertebrate taxa accumulated over six days detected. A) *in situ* eDNA and camera trapping; B) the two different primers used for *in situ* eDNA (12S and 16S rRNA); and C) by all approaches across the three sampling sites at the Mwendi Lagoon, Zambia.

We found that frequent visitors (e.g*., Loxodonta africana*, *Papio. cynocephalus*) and semi-aquatic taxa (e.g., *H. amphibius*, *Xenopus sp*.) were detected on multiple days and by more than one approach, whereas birds, bats, and rodents were typically recorded on a single day and by only one method. All antelopes—*Aepyceros. melampus, Tragelaphus. scriptus, Kobus. vardonii*, and *Kobus. ellipsiprymnus*—were detected exclusively by camera trapping and were not recovered from eDNA samples (Fig 5).

Class	Order	Species	Method	Day 1	Day 2	Day 3	Day 4	Day 5	Day 6
Amphibians	Anura	*Hyperolius sp.*	eDNA					•	•
			Camera traps	NA					
	Anura	*Xenopus sp.*	eDNA					•	
			Camera traps	NA					
Birds	Galliformes	*Numida meleagris*	eDNA			•			
			Camera traps	NA					
	Pelecaniformes	*Threskiornis aethiopicus*	eDNA						•
			Camera traps	NA					
Mammals	Artiodactyla	*Hippopotamus amphibius*	eDNA	•▲	•▲	•▲	•▲	•▲	•▲
			Camera traps	NA					
	Artiodactyla	*Aepyceros melampus*	eDNA						
			Camera traps	NA					
	Primates	*Papio cynocephalus*	eDNA			•		•	
			Camera traps	NA					
	Chiroptera	*Epomophorus pusillus*	eDNA				•		
			Camera traps	NA					
	Rodentia	*Thallomys paedulcus*	eDNA						▲
			Camera traps	NA					
	Proboscidea	*Loxodonta africana*	eDNA	▲	▲	▲	▲		
			Camera traps	NA					
Amphibians	Anura	*Xenopus sp.*	eDNA	•		•	•	•	•
			Camera traps	NA					
Birds	Galliformes	*Numida meleagris*	eDNA						
			Camera traps	NA					
	Passeriformes	*Vidua macroura*	eDNA				•		
			Camera traps	NA					
Mammals	Artiodactyla	*Hippopotamus amphibius*	eDNA	•▲	•▲	•▲	•▲	▲	•▲
			Camera traps	NA					
	Artiodactyla	*Kobus vardonii*	eDNA						
			Camera traps	NA					
	Artiodactyla	*Tragelaphus scriptus*	eDNA						
			Camera traps	NA					
	Artiodactyla	*Kobus ellipsiprymnus*	eDNA						
			Camera traps	NA					
	Primates	*Papio cynocephalus*	eDNA	•▲	•▲		•▲	•▲	
			Camera traps	NA					
Amphibians	Anura	*Xenopus sp.*	eDNA	•	•	•	•	•	•
			Camera traps	NA					
Mammals	Artiodactyla	*Hippopotamus amphibius*	eDNA	•▲	•▲	•▲	•▲	•▲	•▲
			Camera traps	NA					
	Proboscidea	*Loxodonta africana*	eDNA	•▲	▲	▲			
			Camera traps	NA					

Legend		
Point	A	
	B	
	C	
Marker	12S	•
	16S	▲
	Both	•▲

Fig 5. Primary terrestrial vertebrate taxa detected by *in situ* eDNA and camera trapping. Detections are shown for three sampling point (A, B, C) at the Mwendi Lagoon, Zambia, across six consecutive sampling days (Methods). For eDNA detections, the primers 12S (●) and 16S (▲) rRNA are indicated.

Discussion

While eDNA approaches hold promise for providing a cost- and time-efficient alternative to conventional biodiversity monitoring approaches such as camera trapping, the *in situ* application of eDNA in biodiverse areas and countries has so far been hampered by the reliance of standard protocols on Illumina sequencing technology, which requires expensive machinery that is mostly only available at centralized sequencing facilities or research laboratories [4,5,21]. It is, however, important to conduct eDNA analyses on site—to obtain results without delay, empower local decision-makers, and to fulfill international treaties on

data sovereignty, material export, and benefit sharing when it comes to genetics-based

biodiversity monitoring [14,15].

Here, we established portable nanopore sequencing for eDNA analysis of freshwater samples

from Zambia for the detection of primary terrestrial vertebrate species and then compared

established Illumina and nanopore sequencing-based eDNA results with each other based on

the same DNA extracts and the same targeted marker gene regions. We show that the

improvements in nanopore sequencing accuracy through sequencing chemistry updates

(R10.4.1) and improved data basecalling algorithms (Dorado SUP) lead to mostly comparable

species detections between the two eDNA approaches, with nanopore sequencing even

detecting more local species than Illumina sequencing. Based on these results, we applied a

fully mobile approach based on the portable nanopore genomic technology and cost-effective

sample multiplexing for *in situ* biodiversity monitoring on site in the Luambe National Park in

Zambia, confirming the *in situ* applicability of our eDNA approach.

Due to its long sequencing reads, nanopore sequencing further holds promise for extending

future metabarcoding studies to longer marker gene regions, or entire mitochondrial genomes

[18,21,22,38]. For example, Bludau *et al.* showed that nanopore-sequenced amplicons of the

full-length 18S rRNA gene from soil protistan communities achieved greater taxonomic

classification accuracy down to the species level than short-read Illumina-sequenced

amplicons of the Variable Region 9 of the same gene, which only provided reliable

classification on the phylum level [18]. Similarly, Doorenspleet, *et al.* compared nanopore-

sequenced amplicons of a 2 kb region of fish mitochondrial DNA against the commonly used

MiFish primer pair targeting a ~170 bp region. They showed that while the longer amplicons

allowed for more species assignment in a controlled aquarium condition, the shorter amplicons

performed better in natural environmental settings. This is probably because eDNA is often

highly fragmented due to various environmental factors such as UV radiation, microbial

activity, and enzymatic degradation, resulting in higher persistence time for shorter fragments

[39-41]; such shorter DNA fragments are furthermore likely to amplify during PCR reactions

[42]. When using eDNA to monitor vertebrates, especially primary terrestrial vertebrates from

freshwater samples like presented in our study, the samples' DNA might therefore be too

fragmented to allow for efficient long-read gene sequencing, which remains to be assessed.

The suitability of amplicon length in metabarcoding studies might ultimately depend on

different factors, such as the targeted taxa, the eDNA source, and the amount of genetic

variability contained within the amplified marker. In highly diverse regions (e.g., the Luangwa

Valley) that host several sympatric closely related species, for example among antelopes and

felids, the use of longer amplicons could provide the taxonomic resolution needed to

discriminate among species.

Based on our database exploration, we found that, even for small marker gene regions (such

as the 12S and 16S rRNA regions used in this study), genomic databases remain incomplete,

particularly for megadiverse but understudied ecosystems [42-44]—such as Zambia's

subtropical savannahs ecosystems as assessed in this study. The accurate mapping of

marker gene OTUs to publicly available databases such as the entire NCBI nucleotide or the

MIDORI2 databases strongly depends on the regional taxonomic coverage and can lead to

unassigned sequences or false-positive assignments in the case of lacking references [45-

47]. We therefore emphasize that building or populating genetic databases with regional

genetic diversity is crucial for the successful application of eDNA based monitoring

approaches; until databases are more complete, any eDNA application must be interpreted

while taking the availability of genetic information into account.

We further compared our eDNA approaches with more conventional biodiversity monitoring

achieved through simultaneously applied camera trapping Both our eDNA approaches

showed little overlap with camera-based species detections. Several methodological and

ecological factors could explain the discrepancies. As discussed previously, database

completeness, primer choice, and amplicon resolution can influence which taxa are detected

by eDNA. In addition, the interaction between species and the eDNA source can determine

the likelihood of detection. For example, three recent studies have shown that waterborne

eDNA from African waterholes can successfully reveal mammalian and other vertebrate

communities [27-29]. However, we argue that these methods are likely to perform better for

certain species than others, depending on their ecology. Water-dwelling or drought-intolerant

species, such as many mammal or bird species, are more likely to leave detectable DNA

traces in such waterholes. eDNA persistence in waterholes has further shown to be relatively

short (Farell, 2022), which can contribute to variability in detection rates among species. Our

findings illustrate this ecological bias, with eDNA preferentially detecting species with frequent

or prolonged contact with water, such as hippos, elephants, and amphibians. Species that

interact only briefly with water bodies, such as antelopes and felids, may not shed sufficient

DNA to be consistently detected by eDNA, but could reliably be detected by camera trapping.

Similarly, drought-resistant species or those able to obtain water from food resources may

leave little or no trace in water samples. On the other hand, camera traps might miss species

that cannot be easily detected visually, such as, for example, reptiles, amphibians, and small

mammals such as rodents, bats, and pangolins. Overall, the limited overlap between eDNA

and camera trapping suggests their complementary application and highlights the value of

integrating additional eDNA sources such as air, soil, or blood-feeding insects [48-52].

In summary, we show that nanopore sequencing technology can be used to reliably carry out

eDNA-based biodiversity monitoring in remote locations, including in highly diverse LMICs

such as Zambia. The mobile laboratory and protocols we describe can be easily adapted for

processing other eDNA sources, or for other conservation purposes such as pathogen

surveillance. These efforts can pave the way towards democratization of genomic technology

in general, and eDNA applications specifically. However, further efforts are needed to (i) make

eDNA-based approaches more cost-effective so that capacity can be built for routine

applications in LMICs, particularly for conservation purposes where resources are often

limited, and to (ii) contribute to regional reference database developments to better reflect

global genetic biodiversity.

Supplementary Material

All supplementary material is currently available [here](). The code used for this study is available

at: https://github.com/dmgr90/eDNA-Zambia-1

Data Accessibility Statement

All sequencing data is deposited in the NCBI Sequence Read Archive (SRA; accession

number: PRJNA1195924).

Acknowledgements and Benefit-Sharing Statement

We thank the Department of National Parks and Wildlife for this fruitful collaboration. This

research was carried out with their authorization under the research permit NPW/8/27/1.

References

1. Butchart, S. H. M., Walpole, M., Collen, B., van Strien, A., Scharlemann, J. P. W., Almond, R. E. A., Baillie, J. E. M., Bomhard, B., Brown, C., Bruno, J., Carpenter, K. E., Carr, G. M., Chanson, J., Chenery, A. M., Csirke, J., Davidson, N. C., Dentener, F., Foster, M., Galli, A., … Watson, R. 2010;. Global Biodiversity: Indicators of Recent Declines. Science, 328(5982), 1164–1168. doi: 10.1126/science.118751

2. Kindsvater, H. K., Dulvy, N. K., Horswill, C., Juan-Jordá, M.-J., Mangel, M., & Matthiopoulos, J. 2018;. Overcoming the Data Crisis in Biodiversity Conservation. Trends in Ecology & Evolution, 33(9), 676–688. doi: 10.1016/j.tree.2018.06.00

3. Vié, J.-C., Hilton-Taylor, C., & Stuart, S. N. 2009;. Wildlife in a changing world: An analysis of the 2008 IUCN red list of threatened species. IUCN ; Lynx Edicions

4. Lewis, D. 2019;. Rare bird's detection highlights promise of 'environmental DNA'. Nature, 575(7783), 423–424. doi: 10.1038/d41586-019-03522-

5. Matthias, L., Allison, M. J., Maslovat, C. Y., Hobbs, J., & Helbing, C. C. 2021;. Improving ecological surveys for the detection of cryptic, fossorial snakes using eDNA on and under artificial cover objects. Ecological Indicators, 131, 108187. doi: 10.1016/j.ecolind.2021.10818

6. Mizumoto, H., Mitsuzuka, T., & Araki, H. 2020;. An Environmental DNA Survey on Distribution of an Endangered Salmonid Species, Parahucho perryi, in Hokkaido, Japan. Frontiers in Ecology and Evolution, 8, 569425. doi: 10.3389/fevo.2020.56942

7. Itakura, H., Wakiya, R., Yamamoto, S., Kaifu, K., Sato, T., & Minamoto, T. 2019;. Environmental DNA analysis reveals the spatial distribution, abundance, and biomass of

Japanese eels at the river-basin scale. Aquatic Conservation: Marine and Freshwater
Ecosystems, 29(3), 361–373. doi: 10.1002/aqc.3058

8. Nardi, C. F., Fernández, D. A., Vanella, F. A., & Chalde, T. 2019;. The expansion of exotic
Chinook salmon (Oncorhynchus tshawytscha) in the extreme south of Patagonia: An
environmental DNA approach. Biological Invasions, 21(4), 1415–1425. doi: 10.1007/s10530-
018-1908-8

9. Sales, N. G., Wangensteen, O. S., Carvalho, D. C., Deiner, K., Præbel, K., Coscia, I.,
McDevitt, A. D., & Mariani, S. 2021;. Space-time dynamics in monitoring neotropical fish
communities using eDNA metabarcoding. Science of The Total Environment, 754, 142096.
doi: 10.1016/j.scitotenv.2020.142096

10. Fisher, B., & Christopher, T. 2007;. Poverty and biodiversity: Measuring the overlap of
human poverty and the biodiversity hotspots. Ecological Economics, 62(1), 93–101. doi:
10.1016/j.ecolecon.2006.05.020

11. Urban, L., Perlas, A., Francino, O., Martí-Carreras, J., Muga, B. A., Mwangi, J. W., Boykin
Okalebo, L., Stanton, J. L., Black, A., Waipara, N., Fontsere, C., Eccles, D., Urel, H., Reska,
T., Morales, H. E., Palmada-Flores, M., Marques-Bonet, T., Watsa, M., Libke, Z., ... Van
Oosterhout, C. 2023;. Real-time genomics for One Health. Molecular Systems Biology,
e11686. doi: 10.15252/msb.20231168

12. Watsa, M., Erkenswick, G. A., Pomerantz, A., & Prost, S. 2020;. Portable sequencing as
a teaching tool in conservation and biodiversity research. PLOS Biology, 18(4), e3000667.
doi: 10.1371/journal.pbio.3000667

13. Convention on International Trade in Endangered Species of Wild Fauna and Flora |

CITES. (n.d.). Retrieved December 6, 2024, from https://cites.org/eng/disc/text.php

14. The Nagoya Protocol on Access and Benefit-sharing. (n.d.). Retrieved December 6, 2024,

from https://www.cbd.int/ab

15. CARE Principles—Global Indigenous Data Alliance. (n.d.). Retrieved December 6, 2024,

from   https://www.gida-global.org/care

16. Urban, L., Miller, A. K., Eason, D., Vercoe, D., Shaffer, M., Wilkinson, S. P., Jeunen, G.-

J., Gemmell, N. J., & Digby, A. 2022;. Non-invasive real-time genomic monitoring of the

critically endangered kākāpō [Preprint]. Genomics. doi: 10.1101/2022.11.14.516431

17. Baloğlu, B., Chen, Z., Elbrecht, V., Braukmann, T., MacDonald, S., & Steinke, D. 2021;. A

workflow for accurate metabarcoding using nanopore MinION sequencing. Methods in

Ecology and Evolution, 12(5), 794–804. doi: 10.1111/2041-210X.13561

18. Bludau, D., Sieber, G., Shah, M., Deep, A., Boenigk, J., & Beisser, D. 2025;. Breaking the

Standard: Can Oxford Nanopore Technologies Sequencing Compete With Illumina in

Protistan Amplicon Studies? Environmental DNA, 7(2), e70084. doi: 10.1002/edn3.70084

19. Van Der Reis, A. L., Beckley, L. E., Olivar, M. P., & Jeffs, A. G. 2023;. Nanopore short-

read sequencing: A quick, cost-effective and accurate method for DNA metabarcoding.

Environmental DNA, 5(2), 282–296. doi: 10.1002/edn3.374

20. Egeter, B., Veríssimo, J., Lopes-Lima, M., Chaves, C., Pinto, J., Riccardi, N., Beja, P., &

Fonseca, N. A. 2021;. Metabarcoding with MinION: Speeding up the detection of invasive

aquatic species using environmental DNA and nanopore sequencing. ARPHA Conference

Abstracts, 4, e65036. doi: 10.3897/aca.4.e6503

21. Bista, I., & Lino, A. 2025. Long-read sequencing for biodiversity analyses – a

comprehensive guide. EcoEvoRxiv (preprint), published August 26, 2025.

22. Doorenspleet, K., Jansen, L., Oosterbroek, S., Kamermans, P., Bos, O., Wurz, E., Murk,

A., & Nijland, R. 2025;. The Long and the Short of It: Nanopore-Based eDNA Metabarcoding

of Marine Vertebrates Works; Sensitivity and Species-Level Assignment Depend on Amplicon

Lengths. Molecular Ecology Resources, 25(4), e14079. doi: 10.1111/1755-0998.1407

23. Government of the Republic of Zambia. 2015;. Zambia's second national biodiversity

strategy and action plan (NBSAP-2) 2015–2025. Ministry of Lands, Natural Resources and

Environmental Protection.

24. Critical Ecosystem Partnership Fund (CEPF). Final Project Completion Report: Project

Number SG68954. 2

017. Available from: https://www.cepf.net/sites/default/files/final-report-sg68954.pdf

25. Center for Large Landscape Conservation (CLLC). 2020;. Zambia FOCUS-BRI Report.

Retrieved from https://largelandscapes.org/wp-content/uploads/Zambia-FOCUS-BRI.pdf

26. Thrash, I., Theron, G. K., & du Bothma, J. P. 1995; Dry season herbivore densities around

drinking troughs in the Kruger National Park. *Journal of Arid Environments, 29*(2), 213–219.

doi: 10.1016/S0140-1963(05)80091-6

27. Farrell, M. J., Govender, D., Hajibabaei, M., Van Der Bank, M., & Davies, T. J. 2022;

Environmental DNA as a management tool for tracking artificial waterhole use in savanna

ecosystems. *Biological Conservation, 274,* 109712. doi: 10.1016/j.biocon.2022.109712

28. Seeber, P. A., McEwen, G. K., Löber, U., Förster, D. W., East, M. L., Melzheimer, J., &

Greenwood, A. D. 2019; Terrestrial mammal surveillance using hybridization capture of

environmental DNA from African waterholes. *Molecular Ecology Resources, 19,* 1486–1496.

doi: 10.1111/1755-0998.13069

29. Schenekar, T., Baxter, J., Phukuntsi, M. A., Sedlmayr, I., Weckworth, B., & Mwale, M.

2024; Optimizing waterborne eDNA capture from waterholes in savanna systems under

remote field conditions. *Molecular Ecology Resources, 24*(4), e13942. doi: 10.1111/1755-

0998.13942

30. Riaz, T., Shehzad, W., Viari, A., Pompanon, F., Taberlet, P., & Coissac, E. 2011;.

ecoPrimers: Inference of new DNA barcode markers from whole genome sequence analysis.

Nucleic Acids Research, 39(21), e145–e145. doi: 10.1093/nar/gkr732

31. Taylor, P. G. 1996; Reproducibility of ancient DNA sequences from extinct Pleistocene

fauna. *Molecular Biology and Evolution*. 13(1): 283–285. doi: 10.1093/oxfordjournals.molbev

32. Srivathsan, A., Feng, V., Suárez, D., Emerson, B., & Meier, R. 2023;. Rapid species

discovery and identification with real-time barcoding facilitated by ONTbarcoder 2.0 and

Oxford Nanopore R10.4 [Preprint]. Ecology. doi: 10.1101/2023.06.26.546538

33. Martin, M. 2011; Cutadapt Removes Adapter Sequences from High-Throughput

Sequencing Reads. EMBnet Journal, 17, 10-12

34. Rognes, T., Flouri, T., Nichols, B., Quince, C., & Mahé, F. 2016;. VSEARCH: A versatile

open source tool for metagenomics. PeerJ, 4, e2584. doi: 10.7717/peerj.258

35. Leray, M., Knowlton, N., & Machida, R. J. 2022;. MIDORI2: A collection of quality

controlled, preformatted, and regularly updated reference databases for taxonomic

assignment of eukaryotic mitochondrial sequences. *Ecology and Evolution*, 12(4), e0303. doi:

10.1002/edn3.303

36. Calvignac-Spencer, S., Merkel, K., Kutzner, N., Kühl, H., Boesch, C., Kappeler, P. M.,

Metzger, S., Schubert, G., & Leendertz, F. H. 2013; Carrion fly-derived DNA as a tool for

comprehensive and cost-effective assessment of mammalian biodiversity. *Molecular Ecology*.

22(4): 915–924. doi: 10.1111/mec.12183

37. Vestheim, H., & Jarman, S. N. 2008; Blocking primers to enhance PCR amplification of

rare sequences in mixed samples – a case study on prey DNA in Antarctic krill stomachs.

Frontiers in Zoology. 5: 12. doi: 10.1186/1742-9994-5-12

38. Urban, L., Holzer, A., Baronas, J. J., Hall, M. B., Braeuninger-Weimer, P., Scherm, M. J.,

Kunz, D. J., Perera, S. N., Martin-Herranz, D. E., Tipper, E. T., Salter, S. J., & Stammnitz, M.

R. 2021;. Freshwater monitoring by nanopore sequencing. eLife, 10, e61504. doi:

10.7554/eLife.61504

39. Bista, I., Carvalho, G. R., Tang, M., Walsh, K., Zhou, X., Hajibabaei, M., Shokralla, S.,

Seymour, M., Bradley, D., Liu, S., Christmas, M., & Creer, S. 2018;. Performance of amplicon

and shotgun sequencing for accurate biomass estimation in invertebrate community samples.

Molecular Ecology Resources, 18(5), 1020–1034. doi: 10.1111/1755-0998.1288

40. Deagle, B. E., Jarman, S. N., Coissac, E., Pompanon, F., & Taberlet, P. 2014;. DNA

metabarcoding and the cytochrome c oxidase subunit I marker: Not a perfect match. Biology

Letters, 10(9), 20140562. doi: 10.1098/rsbl.2014.056

41. Jo, T., Murakami, H., Masuda, R., Sakata, M. K., Yamamoto, S., & Minamoto, T. 2017;.

Rapid degradation of longer DNA fragments enables the improved estimation of distribution

and biomass using environmental DNA. Molecular Ecology Resources, 17(6). doi:

10.1111/1755-0998.12685

42 Deiner, K., Bik, H. M., Mächler, E., Seymour, M., Lacoursière-Roussel, A., Altermatt, F.,

Creer, S., Bista, I., Lodge, D. M., Vere, N., Pfrender, M. E., & Bernatchez, L. 2017;.

Environmental DNA metabarcoding: Transforming how we survey animal and plant

communities. Molecular Ecology, 26(21), 5872–5895. doi: 10.1111/mec.14350

43. Li, F., Zhang, Y., Altermatt, F., Zhang, X., Cai, Y., & Yang, Z. 2022;. Gap analysis for DNA-

based biomonitoring of aquatic ecosystems in China. Ecological Indicators, 137, 108732. doi:

10.1016/j.ecolind.2022.10873

44. Marques, V., Milhau, T., Albouy, C., Dejean, T., Manel, S., Mouillot, D., & Juhel, J. 2021;.

GAPeDNA: Assessing and mapping global species gaps in genetic databases for eDNA

metabarcoding. Diversity and Distributions, 27(10), 1880–1892. doi: 10.1111/ddi.13142

45. Supple, M. A., & Shapiro, B. 2018;. Conservation of biodiversity in the genomics era.

Genome Biology, 19(1), 131. doi: 10.1186/s13059-018-1520-3

46. Keck, F., Couton, M., & Altermatt, F. 2023;. Navigating the seven challenges of taxonomic

reference databases in metabarcoding analyses. Molecular Ecology Resources, 23(4), 742–

755. doi: 10.1111/1755-0998.13746

47. Mugnai, F., Costantini, F., Chenuil, A., Leduc, M., Gutiérrez Ortega, J. M., & Meglécz, E.

2023;. Be positive: Customized reference databases and new, local barcodes balance false

taxonomic assignments in metabarcoding studies. PeerJ, 11, e14616. doi: 10.7717/peerj.1461

48 Polling, M., Buij, R., Laros, I., & de Groot, G. A. 2024;. Continuous daily sampling of

airborne eDNA detects all vertebrate species identified by camera traps. Environmental DNA,

First published: 22 August 2024. doi: 10.1002/edn3.591

49. Lynggaard, C., Frøslev, T. G., Johnson, M. S., Olsen, M. T., & Bohmann, K. 2024;.

Airborne environmental DNA captures terrestrial vertebrate diversity in nature. Molecular

Ecology Resources, 24(1), e13840. doi: 10.1111/1755-0998.13840

50. Urban, L., Miller, A. K., Eason, D., Vercoe, D., Shaffer, M., Wilkinson, S. P., Jeunen, G.-

J., Gemmell, N. J., & Digby, A. 2023;. Non-invasive real-time genomic monitoring of the

critically endangered kākāpō. eLife, 12, RP84553. doi: 10.7554/eLife.84553.2

51. Vieira, C. J. S. P., Gyawali, N., Onn, M. B., et al. 2024;. Mosquito bloodmeals can be used

to determine vertebrate diversity, host preference, and pathogen exposure in humans and

wildlife. Scientific Reports, 14, 23203. doi: 10.1038/s41598-024-73820-y

52. Reeves, L. E., Gillett-Kaufman, J. L., Kawahara, A. Y., & Kaufman, P. E. 2018;. Barcoding

blood meals: New vertebrate-specific primer sets for assigning taxonomic identities to host

DNA from mosquito blood meals. PLoS Neglected Tropical Diseases, 12(8), e0006767. doi:

10.1371/journal.pntd.0006767