Title Page

Inverse association between serum vitamin B_{12} level and abundance of potential B_{12}-

producing gut microbes in Indian children

Nisha Chandel1‡, Priyansh Patel1‡, Pramod R. Somvanshi[1,2], Anil Kumar Verma[3], Vivek

Thakur[1*]

[1]Department of Systems & Computational Biology, School of Life Sciences, University of

Hyderabad, India

[2]Asian Centre for Endocrine Pancreas Research, Asian Healthcare Foundation, AIG Hospitals,

Hyderabad, India

[3]ICMR-National Institute of Research in Tribal Health (NIRTH), Jabalpur, India

‡ The authors have contributed equally to the work

* Corresponding author: vivek22@uohyd.ac.in

List of abbreviations

p-adj: Adjusted p-value

CNNS: Comprehensive National Nutrition Survey

IF: Intrinsic factor

MAGs: Metagenome-Assembled Genomes

MCM: Mitochondrial L-Methyl-malonyl-CoA mutase

MetH: Methionine synthase

NRC: Nutrition Rehabilitation Centre

RDI: Recommended dietary intake

Vit. B_{12}: Vitamin B_{12}

Abstract

Background: The human gut microbiome is a natural source of essential micronutrients like B-vitamins, which are utilized by both the host and other community members. The prevalence and abundance of known B-vitamin producers and B-vitamin biosynthesis pathways have already been reported in gut microbiome cohorts of a few countries including India.

Objective: To test whether the presence of B-vitamin producers/biosynthetic pathways translates into serum B-vitamin levels, taking B_{12} as a case example.

Methods: Fecal samples were collected from non-deficient (serum B_{12} level > 210 pg/mL, n=29) and B_{12} deficient (serum B_{12} level < 210 pg/mL, n=30) children from a tribal region of central India. Whole metagenomic DNA was extracted, sequenced, and analyzed for taxonomic profiling and diversity comparisons. Differentially abundant taxa between two groups were identified. The prevalence and abundance of potential B_{12} producers were compared, and their association with serum B_{12} level was established.

Results: A comparison of *within-sample* diversity between the two groups didn't show any difference; however, *between-sample* diversity was significantly less in the B_{12} deficient group. Differential abundance testing also showed different microbiome structure in the B_{12} deficient group, where an increased abundance of B_{12} transporter-carrying *Bacteroides thetaiotaomicron*, a few pathogenic species, and ten known B_{12} producers was observed. Potential B_{12} producers were also significantly prevalent and abundant in the deficient group. Their cumulative abundance was also significantly higher in the deficient group and showed a negative association with serum B_{12} levels.

45 **Conclusion:** A higher abundance of potential B_{12} producers in the deficient group suggested an

46 adaptive mechanism by the gut microbiome to meet the community's B_{12} requirements, by

47 selectively promoting the growth of B_{12} producers.

Introduction

Cobalamin (Vitamin B_{12}) is one of the structurally most complex biomolecules, exclusively synthesized by bacteria and a few archaea [1, 2]. It is needed as a co-factor only for two enzymes in humans, namely, cytosolic Methionine synthase (MetH) and mitochondrial L-Methyl-malonyl-CoA mutase (MCM) [3]. Being an essential micronutrient, it must be acquired by ingestion [1]. Its deficiency impairs DNA synthesis, and causes erythroblast apoptosis, resulting in anemia (reviewed by Koury and Ponka; 2004) [4]. B_{12} depleted diet, malabsorption, and defects in the cellular delivery and uptake system are the causes of deficiency, as reviewed in multiple studies [5-7]. The recommended dietary intake (RDI) of vitamin B_{12} (Vit. B_{12}) ranges from 0.9 to 2.4 micrograms from infants to adults, which can be sourced from an animal-based diet, and fortified foods [8]. Ingestion of such foods brings Vit. B_{12} to the small intestine, where it binds to an intrinsic factor (IF) released from the stomach. This IF-B_{12} complex gets absorbed in the small intestine. The unabsorbed B_{12} is utilized by bacteria in both the small and large intestine (see the review articles) [7, 9].

Biosynthesis of Vit. B_{12} is one of the essential functions of the gut bacterial community [10, 11]. However, approximately 25% of sequenced gut bacterial genomes had the genetic potential to synthesize corrinoids (cobalamin and related cofactors), whereas 80% were predicted to use them [12], indicating that the majority of the community members depend on B_{12} producers and unabsorbed dietary B_{12} for their requirement.

Vit. B_{12} modulates the gut community structure, as it is best known for its role as a cofactor and gene expression regulator in bacteria [13]. Different gut bacterial species require distinct corrinoids, and they obtain specific ones by corrinoid remodeling, which limits their availability

to the host [13, 14]. Several possible competitive and symbiotic interactions within the gut

community indicated the co-evolution of community members, which possibly has also benefited

the vitamin homeostasis of the host [11].

Earlier, we reported the biosynthetic potential of gut microbes for all B vitamins in Indian

cohorts [10] and estimated a slightly higher contribution of such microbes to RDI than a previous

report [11]. On the contrary, a contemporary country-wide nutritional survey (Comprehensive

National Nutrition Survey, CNNS-2018) reported a worrying level of prevalence of Vit. B_{12}

deficiency in the Indian population [15]. This made us wonder if i) a high abundance of B_{12}

producers or B_{12} biosynthesis pathways in the gut gets translated into host serum B_{12} levels, and

ii) serum B_{12} deficiency shows any association with the gut community structure. The current

study aims to answer these questions by analyzing fecal metagenome data from B_{12} deficient

(n=30) and non-deficient (n=29) children from the Balaghat district of Madhya Pradesh, India,

where we assumed that fecal metagenome is a proxy of the gut metagenome and tested the

association of host serum B_{12} level with potential gut B_{12} producers (Figure 1A).

Materials and Methods

Sample Collection

Sampling sites included the Nutrition Rehabilitation Centre (NRC), Anganwadi centres (part of

the Integrated Child Development Services program), and the Government Primary School of

wards 3, 4, and 9 of Baihar village in the Balaghat district of Madhya Pradesh state, India.

Children who hadn't taken antibiotics for the last month and didn't have diarrhoea, constipation,

other gastrointestinal diseases, or any other disease were included in this study. Metadata about

all subjects such as age, gender, BMI, diet, and antibiotics or supplements usage, was gathered using a questionnaire and consent form, which was approved by the Institutional Ethics Committee (UH/IEC/2022/389), University of Hyderabad, India and ICMR-National Institute of Research in Tribal Health, India (Table S1). The basic summary of the data for the two groups is provided in Table 2. The fecal sample collection was done from January 2023 to March 2023, involving children with ages ranging between 2-12 years as follows: serum B_{12} profiling of 158 children was first done at Thyrocare Technologies Ltd. (Mumbai) using Chemi Luminescent Microparticle Intrinsic factor Immunoassay (CMIA; Intra assay (%CV): 5.6, Inter assay (%CV): 6.8, Sensitivity: 125 pg/mL) or Competitive Chemi Luminescent Immuno Assay (CLIA; Intra assay (%CV): 5.0%, Inter assay (%CV): 9.2 %; Sensitivity: 45 pg/ml) methods, and they were classified into non-deficient (>210 pg/ml; n=118) and B_{12} deficient groups (<210 pg/ml; n=40). After obtaining 40 B_{12} deficient cases, further screening of serum B_{12} profiling was stopped as it was an exploratory study. The fecal samples, however, could be obtained only for 30 of them. To make both groups comparable, 29 non-deficient children were randomly selected for fecal sampling. About 30 participants per group is a common starting point in exploratory research due to its ability to balance statistical power with practical feasibility, its suitability for detecting small-to-medium effects in high-dimensional microbiome data, and its alignment with power analysis estimates for achieving a reasonable likelihood of detecting true effects in small-to-medium-sized studies. The fecal samples were collected in sterile stool containers and stored at 0°C within 1 to 5 hours and later were transported in dry ice to miBiome Therapeutics LLP Mumbai, India, for DNA extraction and sequencing (Figure 1A).

Fecal Metagenomic DNA extraction and quantification

Whole metagenomic DNA from all 59 samples was isolated using the QIAamp PowerFecal Pro

DNA Kit according to the manufacturer's instructions. The DNA samples were quantified on

NanoDrop One and Qubit using water and a standard as a control, respectively. DNA quality was

estimated by agarose gel electrophoresis.

Library Preparation and Sequencing

Libraries were prepared using QIAseq® FX DNA library kit from Qiagen. Briefly, 150 ng of

DNA was used for fragmentation and end repair, followed by adapter ligation. Excess adapters

were removed using AMPure XP purification beads, followed by PCR enrichment of adapter-

ligated DNA and clean-up of library products as instructed by the manufacturer. The cleaned

libraries were quantitated on a Qubit fluorometer, and appropriate dilutions were loaded on HS

D1000 screen tape to determine the size range of the fragments and the average library size.

Eventually, paired-end sequencing, with reads of size 150 bases each, was done using the

Illumina *Novaseq6000* platform.

Pre-processing of metagenomic reads

The quality of sequencing data was checked using FastQC v0.12.1 [16] and MultiQC v1.22.1

[17]. Since all reads had an average *Phred*-score > 25, only ambiguous bases and adapters were

removed using *fastp* [18]. Further, host contamination in the filtered reads was removed by

aligning reads to the human genome (GRCh38) using Bowtie2 v2.5.4 [19], and unaligned reads

were extracted using SAMtools v1.20 [20]. The processed reads were used for downstream

analysis (Figure 1B).

Taxonomy profiling and species filtration

The compositional profiling of each sample was done using MetaPhlAn 4.0.6 (Metagenomic

Phylogenetic Analysis) [21]. To further reduce the risk of contamination or alignment artifacts,

species with very low mean relative abundance (<= 0.001%) or prevalence (<= 15%) were

removed (Figure 1B).

Diversity analysis

The filtered compositional profiles were used to measure alpha and beta diversity between the

two groups. The alpha diversity (or within-sample diversity) was measured by using the *vegan*

package [22] using multiple measures such as species richness (*specnumber()* function),

Shannon index (*diversity()*), and Simpson index (*diversity()*). The beta diversity (or between-

sample diversity) analysis was first performed using a dimensionality reduction approach:

Principal Coordinate Analysis (PCoA). It was done using the *cmdscale()* function on Bray-Curtis

distance matrix. Alternatively, the beta diversity analysis was performed using pair-wise distance

measures like Bray-Curtis, Jaccard distance, and Robust Aitchison distances; all were calculated

using the *adonis()* function in the *vegan* package in R.

Vit. B_{12} pathway analysis at the community (or microbiome) level

For Vit. B_{12} biosynthesis pathway analysis, filtered reads of samples from each group were co-

assembled using the *de novo* assembly tool, MEGAHIT (v1.2.9) [23]. The contigs of each co-

assembly were annotated using the Prokka tool (v1.14.5) [24]. Gene abundance in each sample

was estimated by mapping filtered reads from the respective sample to the gene sequences of the

corresponding co-assembly using the Bowtie2 tool (v2.5.4) [19], and the read counts were

estimated by the Expectation-Maximization-based tool, RSEM (v1.3.3) [25] (Figure 1B).

To examine the presence of B_{12} biosynthesis pathway(s) in any sample, the reference genes of

the known B_{12} biosynthesis pathways or subpathways in bacteria were first obtained from

MetaCyc [26] (Table 1). Since the majority of the reference genes were from *Salmonella*

typhimurium genome (RefSeq ID: GCF_000006945.2) [27], the genome annotation using Prokka

tool [24] was used to obtain the protein sequences. The reference genes which were from other

bacteria, namely, *Listeria innocua serovar 6a* (strain ATCC BAA-680 / CLIP 11262),

Escherichia coli (strain K12), Priestia megaterium, Desulfuromonas acetoxidans (strain DSM

684 / 11070), *Eubacterium callanderi*, their protein sequences were obtained from UniProt

database [28], and appended to *S. Typhimurium* protein sequences. The orthologs of the

appended set were identified among translated gene sequences of each co-assembly using the

OrthoFinder tool (v2.5.5) [29] (Figure 1B).

The pathway abundance in each sample was estimated by taking the median value of normalized

abundances of all genes of the pathway. The abundance of any gene was independent of the

genomes to which they belonged.

Vit. B_{12} pathway analysis at Metagenome-Assembled Genomes (MAGs) or species level

The above analysis was based on B_{12} biosynthesis through cooperativity at the community level.

Additionally, the contribution from the individual MAGs with complete pathways was also

considered for comparison between the groups. The contigs from each co-assembly were binned

to obtain MAGs using the CONCOCT tool (v1.1.0) [30]. Their quality was assessed with

CheckM2 tool (v1.0.2) [31], and MAGs with completeness <= 90 % were discarded. Complete

MAGs were annotated using the Prokka tool (v1.14.5) [24], and their taxonomy was predicted

using GTDB toolkit [32]. Orthologs of the B_{12} biosynthesis pathways gene in each MAG were

identified by using the OrthoFinder tool (v2.5.5) [29] (Figure 1B). The mean contig depth

(calculated by CONCOCT) belonging to a particular MAG was used as its abundance. The

proportion of samples having a MAG was termed as its prevalence. Pathways with completeness

<= 80% and present only in 5 MAGs or less than that were eliminated from the downstream

analysis.

Differential abundance testing and other statistical Analyses

All statistical analyses were performed in R (version 4.2.2; www.r-project.org). The difference in

beta diversity was statistically tested using the *adonis()* function in the vegan package, which

uses PERMANOVA. Statistical testing for differentially abundant species was done using a

negative binomial regression model-based *Wald* test implemented in DESeq2 [33] after

controlling for the effect of confounders, namely batch, age, gender, diet, and BMI, which uses

Benjamini-Hochberg procedure (FDR) for multiple hypothesis testing. The adjusted *p-value (p-*

adj) <= 0.05 was considered significant. The known B_{12} producers were inferred by comparing

the differentially abundant species to a comprehensive list from our previous study [10].

All the statistical comparisons between groups were performed using the non-parametric

(*Wilcoxon rank-sum)* test. The proportion of MAGs carrying B_{12} pathways (potential B_{12}

producers) between the two groups was compared using the proportionality test function

(*prop.test()*). The association between serum B_{12} levels and the cumulative abundance of B_{12}

producers was assessed using a multiple linear regression model in R (*lm()*), controlling for

potential confounders such as age, gender, diet, and BMI. All figures were generated using the

ggplot2 package [34] in R.

Results

The prevalence of B_{12} deficiency in the tribal children cohort was higher than the

national/state average for school-age children

Serum B_{12} profiling of 158 children from the Balaghat district of central India showed a

deficiency in ~25% of samples (Supplemental Table 2). Compared to CNNS-2018 survey data,

though the prevalence in the current study had a similar increasing trend with age (10%, 28%,

and 39.3%, in pre-school, school-age and adolescent, respectively), however, the prevalence in

the school-aged group was relatively higher (~28%, n=90) (Supplemental Table 2).

From the total of 158 subjects, two groups of similar size were made (see methods) for further

study: one with a normal B_{12} level (n=29, range: 212-810 pg/ml, median=365 pg/ml) and another

with B_{12} deficiency (n=30, range: 82-198 pg/ml, median=169 pg/ml) (Figure 1A, Supplemental

Figure 1A; Table 2; Supplemental Table 1). Whole metagenome shotgun sequencing of fecal

samples of both the groups resulted in ~ 25 million paired-end reads per sample (~24 million

after quality control) (Supplemental Figure 1B).

Mixed trend for alpha (*within-sample*) diversity in the B_{12} deficient group

Taxonomic profiling results showed the presence of 1225 species (Supplemental Table 3), where

most of the species belonged to phylum Bacillota, Bacteroidota, Actinomycetota, and

Pseudomonadota (Figure 2A). It was also observed that ~20% of the reads remained

unclassified, which may be partly due to the uniqueness of the Indian gut microbiome (Figure

2A). Species present only in a few samples and with low abundance were filtered (see materials

and methods), and the remaining 374 species were considered for downstream analysis. Within-

sample diversity showed no significant difference in the B_{12}-deficient group when estimated

using metrics based on both the richness and evenness, such as the Shannon index ($p = 0.11$) and

Simpson index ($p = 0.10$). However, it was significantly higher when estimated using only the

species richness ($p = 0.05$) (Figure 2B and C; Supplemental Figure 2A).

Significant reduction in beta (*between-sample*) diversity in the B_{12} deficient group indicated

convergence

To examine if the microbiome composition was different in B_{12}-deficient condition, PCoA using

the Bray-Curtis dissimilarity index was done. B_{12} deficient samples clustered together, partially

overlapping with the non-deficient ones along the first principal coordinate, which captured

~24% of the variation (Figure 3A). The higher similarity (i.e., lower distance) among the B_{12}

deficient samples compared to non-deficient ones was statistically confirmed using the

PERMANOVA test ($p = 0.004$) (Figure 3B). Similar trends of lower beta diversity in the B_{12}

deficient group were observed using other distance measures like the Jaccard distance ($p =$

0.005) and the Robust Aitchison distance ($p = 0.07$), further confirming different gut community

structure in B_{12} deficient condition (Figure 3C; Supplemental Figure 2B).

Ten *known* B_{12} producers were among sixty-two differentially more abundant species in the

B_{12} deficient group

To find out differentially abundant species in any of the groups, their abundance was statistically

compared. A total of 157 significantly abundant species, 62 were more abundant in the B_{12}

deficient group (FDR = 0.05, $\log_2$ fold change >=3) (Figure 3D), which included the B_{12} transport system carrying species of genus *Bacteroides,* pathogenic species *Sutterella faecalis, Clostridiaceae unclassified SGB4771,* mucin degrader *Akkermansia muciniphila,* and others, which are listed in Supplemental Table 4 [35 - 38]. The non-deficient group had a higher abundance of lactate metabolizing species of genus *Veillonella, Streptococcus thermophilus,* plant-based diet associate *Prevotella stercoria* (previously known as *P. copri*), a short-chain fatty acid (acetic acid) producer *Eubacterium siraeum,* probiotic potential candidate *Blautia* and others (Supplemental Table 4) [39 - 43]. This indicated gut community perturbation in B_{12} deficient condition, where a higher abundance of at least a couple of pathogens and a lower abundance of a few beneficial bacteria (Figure 3D) was observed. Further, the known B_{12} producers among the differentially abundant species were examined: the B_{12} deficient group had about ten known B_{12} producers, while only six were observed in the non-deficient group (Figure 3D).

Cooperativity-based B_{12} biosynthesis pathway abundance showed no difference in the B_{12} deficient group

Anticipating the incompleteness of the literature information on B_{12} producers, the pathway abundance was further inferred directly from the metagenomic data, and its distribution was compared between the two groups. The pathway abundance in each sample was estimated assuming a cooperative interaction among community members for its biosynthesis. The abundance of all five B_{12} biosynthesis pathways/subpathways (Supplemental Table 5) was found to be statistically similar (*p > 0.05*), indicating no difference between the groups for the given assumption (Figure 4A-C; Supplemental Figure 3A and B).

Potential B_{12} producers were generally more prevalent and abundant in the B_{12} deficient group

An alternative to the cooperativity-based B_{12} biosynthesis assumption would be considering the contribution from the MAGs with a complete B_{12} biosynthesis pathway (i.e., the potential B_{12} producers) for abundance estimation. Out of the total of 324 MAGs in the non-deficient group and 367 in the B_{12} deficient group, 135 and 161 were complete, respectively. While the number of MAGs having a particular B_{12} biosynthesis pathway varied between both groups, their proportion was statistically similar (Table 1). Among five pathways, anaerobic, GDP-I, and Cobinamide salvage were present in more than 5 MAGs, so only these were considered for further analysis. The prevalence of the potential B_{12} producers was significantly higher in the B_{12} deficient group for all three pathways; however, their abundance was significantly higher for two of the pathways, with the anaerobic pathway an exception (*p=0.3*) (Figure 4C–I).

The cumulative abundance of potential B_{12} producers was higher in the B_{12} deficient group

Instead of considering the abundance of potential B_{12} producers for individual pathways, their cumulative abundance in each sample was calculated, and a group-wise comparison was made. The B_{12} deficient group had a much higher cumulative abundance (*p = 7.955e-09*) (Figure 5A). To further establish the association between serum B_{12} level and the cumulative abundance of B_{12} producers, a multiple linear regression analysis was carried out. Results showed a negative association, where with a decline in serum B_{12} levels, the abundance of B_{12} producers significantly increased (*p = <0.001, β_0 = 6.42, β_1 = -0.02, Std. Error = 0.006*) (Figure 5B).

Moreover, applying local regression (locally weighted scatterplot smoothing, LOWESS) revealed a refined trend, suggesting that the cumulative abundance of potential B_{12} producers increased non-linearly as the serum B_{12} levels approached the deficient range (Supplemental Figure 4). The majority of the B_{12} producers belonged to the phylum Bacillota, followed by Bacteroidota. 18.4% and 21.21% of them remained unclassified in B_{12} deficient and non-deficient groups, respectively (Supplemental Table 6, Supplemental Figure 5A and B).

Discussion

To the best of our knowledge, the current study is the first one to directly elucidate the association between serum B_{12} level and gut microbiome at an improved species-level resolution and provide a better understanding than the existing reports. There have been multiple studies in humans and mice to examine the association0 [44-52], but all of them except one were done using 16S rRNA amplicon data and had a resolution at the genus level. The exception was a study on Saudi Arabian females using shotgun metagenome sequencing data; however, their aim was slightly different as they examined the association between Vit. B_{12} status and gut microbiota in the context of obesity [51].

In our study, we observed no significant differences in alpha diversity between the two groups, while a lower beta diversity was noted in the Vit. B_{12} deficient condition. These findings were partially consistent with previous reports. However, the literature evidence on the association between Vit. B_{12} status and gut microbial diversity remains limited. A systematic review on Vit. B_{12} and the gut microbiome summarized that both alpha and beta diversity increased with an increase in Vit. B_{12} levels, as reported in *in-vivo* studies; however, findings from mouse models

and observational human studies were heterogeneous [53]. Two of the mouse studies did not

show any difference in alpha diversity [46,54], while the third reported lower alpha diversity, but

statistical significance was not reported [55]. Similarly, two human studies did not show any

difference [44,52], while one reported an increase in alpha diversity with B_{12} intake [45].

Regarding beta diversity, three studies did not show any difference [46,55,56], while one study

reported different diversity at the genus level in mice [54]. In humans, two studies showed no

difference [44,52], and one showed different beta diversity [45]. Therefore, while majority of the

studies have reported similar findings as observed in the current study, others have observed

differing outcomes, highlighting variability in the existing literature, possibly due to the

difference in sequencing data, sample sizes, experimental variables.

The different microbiome structure was also supported by the differential abundance of a few

species in either group (Supplemental Table 4). Along with the enrichment of pathogenic

species, the deficient group had a higher abundance of a corrinoid transporter-carrying

Bacteroides thetaiotaomicron, and several other potential B_{12} producers. A lower relative

abundance of the genus *Bacteroides* was reported to be associated with high B_{12} intake in

humans [45, 47], and with B_{12} supplementation in mice and in-vitro models [53]. However, this

trend was conflicted by another study, which reported no such change in the abundance [44]. *B.*

thetaiotaomicron, a mucin generalist, has a crucial role in health and disease pathogenesis [57],

and is also known to have multiple copies of corrinoid transporter (BtuFBCD) in its genome,

which gives an advantage to this bacterium in distinguishing B_{12} analogs produced by several

other members [12, 14, 58]. The higher relative abundance of *B. thetaiotaomicron* in B_{12}

deficient groups suggests a scarcity of the B_{12} in the gut. Overall, a higher abundance of genus

Bacteroides is a characteristic feature of a Vit. B_{12}-depleted microenvironment.

Besides, the taxonomic classification of B_{12} producers and their comparison with the available

knowledge of producers in the current study also revealed several new species that have

biosynthetic potential. For instance, MAGs belonging to species *A. muciniphila, Agathobacter*

faecis, Agathobacter rectalis, Alistipes senegalensis, Anaerobutyricum hallii, Veillonella atypica

etc., had B_{12} biosynthesis potential (Supplemental Table 6).

Enrichment of known B_{12} producers and higher prevalence and abundance of potential B_{12}

producers in the deficient group, as well as their negative association with serum B_{12} level,

indicated that host B_{12} level is linked with the gut microbiome. We hypothesize that this

association may reflect a selection pressure on the microbial community to favor those that could

grow by synthesizing their own B_{12} over those solely depending on the cobalamin analogs in the

lumen. An alternate hypothesis, which may however require more supporting evidence, can be a

feedback response possibly originating from the host to the gut microbial community, in a way

that enhances microbial B_{12} biosynthesis, so that it could source a fraction of the microbiota-

derived B_{12} to meet its DRI and thereby partially prevent acute deficiency. If that being the case,

then the gut microbiome would possess buffering ability for homeostasis of some of the essential

micronutrients, wherein certain species respond to the micronutrient deficiency in a threshold-

dependent manner (at subnormal levels) and undergo microbiome remodeling with an increase

in abundance of relevant species for its replenishment. A sustained microbial abundance, along

with Vit. B_{12} deficiency of subnormal levels (~82 pg/ml) indicates a possibility of a threshold-

dependent switch-like response (Supplemental Figure 4) that takes care of severe deficiency;

however, it is insensitive to achieve clinically sufficient Vit. B_{12} levels. Such a sub-optimal

feedback response may be evolutionarily efficient in sparing raw material for *de-novo* Vit. B_{12}

biosynthesis under severe micronutrient deficiency in the long term (Supplemental Figure 6).

In our past work, we highlighted the higher abundance of B-vitamin producers and biosynthesis

pathways in tribal cohorts as a desirable feature and discussed possible links to different diets,

limited access to antibiotics, etc. [10]. Similarly, Das et al. (2019) reported the Chinese

population to have higher B-vitamin biosynthesis pathways compared to the other three

nationalities [59]. Having observed higher B_{12} producers and biosynthesis pathways in the serum

B_{12} deficient group, the previous findings should be re-interpreted based on the hypothesized

feedback model. In both previous reports, the B-vitamin status of subjects was unknown, and the

groups with higher abundance suggest that the higher abundance could be due to dietary

consumption of just adequate or sub-optimal levels of vitamins that are in a range where the

switch of the feedback mechanism is active. This indicates that the vitamin levels are partly

derived from dietary sources and partly from the microbiome in the healthy host.

Limitations and future directions

This study was based on case-control design, and hence cause-effect relationship can't be

established. The results were based on a small sample size involving only children. The profiling

of other biomarkers of Vit. B_{12} status (holotranscobalamin, methylmalonic acid, and

homocysteine) and profiling of biochemical markers to determine the comprehensive health and

nutritional status of an individual was not done, which restricted us from taking into account

other confounding effects in the analysis. To ascertain the proposed hypothesis and causality of

the observed differences, longitudinal/interventional studies required to be done. Further, the

microbial signatures identified in our study could also be experimentally investigated in a

longitudinal study on an animal model by administering B_{12} deficient and sufficient diets.

Conclusion

This is the first study to directly elucidate the association between serum B_{12} level and gut

microbiome at the species or strain level. The gut community in the B_{12}-deficient group was

enriched with mucin-degrading bacteria, a few pathogens, and known B_{12}-producers. With a

higher occurrence of potential B_{12}-producers in the deficient group and their negative association

with serum B_{12} level, we hypothesize that the gut microbiome responds to the deficiency by

selectively promoting the growth of producers, which meets the community's demand for Vit.

B_{12}.

Acknowledgements

We would like to acknowledge the University of Hyderabad's Institution of Eminence (IoE) for

funding our research and the Centre for Modelling, Simulations, and Design (CMSD) for

allowing the use of their High-performance Computer facility. Dr Suyash Shrivastava, and Ms

Prathysha, Scientists/Project staff from ICMR-National Institute of Research in Tribal Health,

Jabalpur, Madhya Pradesh, for preparing the questionnaire; Dr. Devaraj Parasannanavar,

Scientist D, ICMR-National Institute of Health, Hyderabad, for his valuable suggestions for

grouping samples based on clinical data.

Authors' contributions

Designed research: VT, AKV, NC; prepared questionnaire: AKV; collected samples and data:

PP, AKV; performed data analysis: NC, VT; performed statistical analysis: NC, PR; Interpreted

findings: NC, VT, PR; wrote manuscript: NC, PP, VT, PR; All authors have read and approved

the final manuscript.

Data Availability

The raw data is publicly available at NCBI Sequencing Read Archive under Bioproject ID

PRJNA1114651.

Conflicts of Interest

The authors declare that there are no conflicts of interest.

Funding

The project and manpower involved in this research were financially supported by funding from

the University of Hyderabad-Institution of Eminence (grant number: UoH-IoE-RC2-21-023)

from 30 Aug 2021 to 30th Aug 2024.

Reference

1. Roth, JR. Lawrence, JG. Rubenfield, M. Kieffer-Higgins, S. Church, GM.

Characterization of the cobalamin (vitamin B_{12}) biosynthetic genes of Salmonella

typhimurium. Journal of bacteriology. 1993 Jun;175(11):3303-16.

2. Roth, JR. Lawrence, JG. Bobik, TA. Cobalamin (coenzyme B_{12}): synthesis and biological

significance. Annual review of microbiology. 1996 Oct;50(1):137-81.

3. Allen, RH. Stabler, SP. Savagem DG. Lindenbaum, J. Metabolic abnormalities in

cobalamin (vitamin B_{12}) and folate deficiency. The FASEB journal. 1993

Nov;7(14):1344-53.

4. Koury, MJ. Ponka, P. New insights into erythropoiesis: the roles of folate, vitamin B_{12},

and iron. Annu. Rev. Nutr.. 2004 Jul 14;24(1):105-31.

5. Allen, LH. Causes of vitamin B_{12} and folate deficiency. Food Nutr Bull. 2008 Jun;29(2

Suppl):S20-34; discussion S35-7. doi: 10.1177/15648265080292S105. PMID: 18709879.

6. Green, R. Miller, JW. Vitamin B_{12} deficiency. InVitamins and hormones 2022 Jan 1 (Vol.

119, pp. 405-439). Academic Press.

7. Stabler, SP. Allen, RH. Vitamin B_{12} deficiency as a worldwide problem. Annu. Rev.

Nutr.. 2004 Jul 14;24(1):299-326.

8. National Institute of Health; Office of dietary supplements,

https://ods.od.nih.gov/factsheets/VitaminB12-HealthProfessional/ Accessed 30 Oct 2024

9. Rowley, CA. Kendall, MM. To B_{12} or not to B_{12}: Five questions on the role of cobalamin

in host-microbial interactions. PLoS pathogens. 2019 Jan 3;15(1):e1007479.

10. Chandel, N. Somvanshi, PR. Thakur, V. Characterisation of Indian gut microbiome for

B-vitamin production and its comparison with Chinese cohort. British Journal of

Nutrition. 2024 Feb;131(4):686-97.

11. Magnúsdóttir, S. Ravcheev, D. de Crécy-Lagard, V. Thiele, I. Systematic genome

assessment of B-vitamin biosynthesis suggests co-operation among gut microbes.

Frontiers in genetics. 2015 Apr 20;6:148.

12. Degnan, PH. Barry, NA. Mok, KC. Taga, ME. Goodman, AL. Human gut microbes use
multiple transporters to distinguish vitamin B_{12} analogs and compete in the gut. Cell Host
& Microbe. 2014 Jan 15;15(1):47-57.

13. Degnan, PH. Taga, ME. Goodman, AL. Vitamin B_{12} as a modulator of gut microbial
ecology. Cell metabolism. 2014 Nov 4;20(5):769-78.

14. Allen, RH. Stabler, SP. Identification and quantitation of cobalamin and cobalamin
analogues in human feces. The American journal of clinical nutrition. 2008 May
1;87(5):1324-35.

15. Comprehensive   National   Nutrition   Survey   (2016-2018)
https://nhm.gov.in/WriteReadData/l892s/1405796031571201348.pdf Accessed 20 Dec
2024

16. Andrews, S. Babraham Bioinformatics-FastQC a Quality Control Tool for High
Throughput   Sequence   Data.   2010
https://www.bioinformatics.babraham.ac.uk/projects/fastqc (accessed August 2024)

17. Ewels, P. Magnusson, M. Lundin, S. Käller, M. MultiQC: summarize analysis results for
multiple tools and samples in a single report. Bioinformatics. 2016 Oct 1;32(19):3047-8.

18. Chen, S. Zhou, Y. Chen, Y. Gu, J. fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics. 2018 Sep 1;34(17):i884-90.

19. Langmead, B. Salzberg, SL. Fast gapped-read alignment with Bowtie 2. Nature methods.
2012 Apr;9(4):357-9.

20. Li, H. Handsaker, B. Wysoker, A. Fennell, T. Ruan, J. Homer, N. et al. The sequence
alignment/map format and SAMtools. bioinformatics. 2009 Aug 15;25(16):2078-9.

480 21. Blanco-Míguez, A. Beghini, F. Cumbo, F. McIver, LJ. Thompson, KN. Zolfo, M. et al.

481   Extending and improving metagenomic taxonomic profiling with uncharacterized species

482   using MetaPhlAn 4. Nature Biotechnology. 2023 Nov;41(11):1633-44.

483 22. Dixon, P. VEGAN, a package of R functions for community ecology. Journal of

484   vegetation science. 2003 Dec;14(6):927-30.

485 23. Li, D. Luo, R. Liu, CM. Leung, CM. Ting, HF. Sadakane, K. et al. MEGAHIT v1. 0: a

486   fast and scalable metagenome assembler driven by advanced methodologies and

487   community practices. Methods. 2016 Jun 1;102:3-11.

488 24. Seemann, T. Prokka: rapid prokaryotic genome annotation. Bioinformatics. 2014 Jul

489   15;30(14):2068-9.

490 25. Li, B. Dewey, CN. RSEM: accurate transcript quantification from RNA-Seq data with or

491   without a reference genome. BMC bioinformatics. 2011 Dec;12:1-6.

492 26. Caspi, R. Billington, R. Ferrer, L. Foerster, H. Fulcher, CA. Keseler, IM. et al. The

493   MetaCyc database of metabolic pathways and enzymes and the BioCyc collection of

494   pathway/genome databases. Nucleic acids research. 2016 Jan 4;44(D1):D471-80.

495 27. The National Center for Biotechnology Information. https://www.ncbi.nlm.nih.gov/

496   Accessed 01 Aug 2024

497 28. Bairoch, A. Apweiler, R. Wu, CH. Barker, WC. Boeckmann, B. Ferro, S. et al. The

498   universal protein resource (UniProt). Nucleic acids research. 2005 Jan

499   1;33(suppl_1):D154-9.

500 29. Emms, DM. Kelly, S. OrthoFinder: phylogenetic orthology inference for comparative

501   genomics. Genome biology. 2019 Dec;20:1-4.

30. Alneberg, J. Bjarnason, BS. de Bruijn, I. Schirmer, M. Quick, J. Ijaz, UZ. et al.
CONCOCT: clustering contigs on coverage and composition. arXiv preprint
arXiv:1312.4038. 2013 Dec 14.

31. Chklovski, A. Parks, DH. Woodcroft, BJ. Tyson, GW. CheckM2: a rapid, scalable and
accurate tool for assessing microbial genome quality using machine learning. Nature
Methods. 2023 Aug;20(8):1203-12.

32. Chaumeil, PA. Mussig, AJ. Hugenholtz, P. Parks, DH. GTDB-Tk: a toolkit to classify
genomes with the Genome Taxonomy Database.

33. Love, MI. Huber, W. Anders, S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome biology. 2014 Dec;15:1-21.

34. Wexler, AG. Schofield, WB. Degnan, PH. Folta-Stogniew, E. Barry, NA. Goodman, AL.
Human gut Bacteroides capture vitamin B_{12} via cell surface-exposed lipoproteins. Elife.
2018 Sep 18;7:e37138.

35. Hiippala, K. Kainulainen, V. Kalliomäki, M. Arkkila, P. Satokari, R. Mucosal prevalence
and interactions with the epithelium indicate commensalism of Sutterella spp. Frontiers in
microbiology. 2016 Oct 26;7:1706.

36. Muñiz Pedrogo, DA. Chen, J. Hillmann, B. Jeraldo, P. Al-Ghalith, G. Taneja, V. et al. An
increased abundance of Clostridiaceae characterizes arthritis in inflammatory bowel
disease and rheumatoid arthritis: a cross-sectional study. Inflammatory bowel diseases.
2019 Apr 11;25(5):902-13.

37. Derrien, M. Collado, MC. Ben-Amor, K. Salminen, S. de Vos, WM. The Mucin degrader
Akkermansia muciniphila is an abundant resident of the human intestinal tract. Applied
and environmental microbiology. 2008 Mar 1;74(5):1646-8.

38. Scheiman, J. Luber, JM. Chavkin, TA. MacDonald, T. Tung, A. Pham, LD. et al. Meta-
omics analysis of elite athletes identifies a performance-enhancing microbe that functions
via lactate metabolism. Nature medicine. 2019 Jul;25(7):1104-9.

39. Huang, YY. Lu, YH. Liu, XT. Wu, WT. Li, WQ. Lai, SQ. et al. Metabolic properties,
functional characteristics, and practical application of Streptococcus thermophilus. Food
Reviews International. 2024 Feb 17;40(2):792-813.

40. Precup, G. Vodnar, DC. Gut Prevotella as a possible biomarker of diet and its eubiotic
versus dysbiotic roles: a comprehensive literature review. British Journal of Nutrition.
2019 Jul;122(2):131-40.

41. Moore, WC. Johnson, JL. Holdeman, LV. Emendation of Bacteroidaceae and
Butyrivibrio and descriptions of Desulfomonas gen. nov. and ten new species in the
genera Desulfomonas, Butyrivibrio, Eubacterium, Clostridium, and Ruminococcus.
International journal of systematic and evolutionary microbiology. 1976 Apr;26(2):238-
52.

42. Liu, X. Mao, B. Gu, J. Wu, J. Cui, S. Wang, G. et al. Blautia—a new functional genus
with potential probiotic properties?. Gut microbes. 2021 Jan 1;13(1):1875796.

43. Valentini, L. Pinto, A. Bourdel-Marchasson, I. Ostan, R. Brigidi, P. Turroni, S. et al.
Impact of personalized diet and probiotic supplementation on inflammation, nutritional
parameters and intestinal microbiota–The "RISTOMED project": Randomized controlled
trial in healthy older people. Clinical nutrition. 2015 Aug 1;34(4):593-602.

44. Boran, P. Baris, HE. Kepenekli, E. Erzik, C. Soysal, A. Dinh, DM. The impact of vitamin
B_{12} deficiency on infant gut microbiota. European journal of pediatrics. 2020
Mar;179:385-93.

45. Gurwara, S. Ajami, NJ. Jang, A. Hessel, FC. Chen, L. Plew, S. et al. Dietary nutrients
involved in one-carbon metabolism and colonic mucosa-associated gut microbiome in
individuals with an endoscopically normal colon. Nutrients. 2019 Mar 13;11(3):613.

46. Lurz, E. Horne, RG. Määttänen, P. Wu, RY. Botts, SR. Li, B. et al. Vitamin B_{12}
deficiency alters the gut microbiota in a murine model of colitis. Frontiers in nutrition.
2020 Jun 5;7:83.

47. Carrothers, JM. York, MA. Brooker, SL. Lackey, KA. Williams, JE. Shafii, B. et al.
Fecal microbial community structure is stable over time and related to variation in
macronutrient and micronutrient intakes in lactating women. The Journal of nutrition.
2015 Oct 1;145(10):2379-88.

48. Forgie, AJ. Pepin, DM. Ju, T. Tollenaar, S. Sergi, CM. Gruenheid, S. Willing, BP. Over
supplementation with vitamin B_{12} alters microbe-host interactions in the gut leading to
accelerated Citrobacter rodentium colonization and pathogenesis in mice. Microbiome.
2023 Feb 3;11(1):21.

49. Babakobi, MD. Reshef, L. Gihaz, S. Belgorodsky, B. Fishman, A. Bujanover, Y. Effect
of maternal diet and milk lipid composition on the infant gut and maternal milk
microbiomes. Nutrients 2020; 12: 2539 [Internet].

50. Selma-Royo, M. García-Mantrana, I. Calatayud, M. Parra-Llorca, A. Martínez-Costa, C.
Collado, MC. Maternal diet during pregnancy and intestinal markers are associated with
early gut microbiota. European journal of nutrition. 2021 Apr;60:1429-42.

51. Al-Musharaf, S. Aljuraiban, GS. Al-Ajllan, L. Al-Khaldi, N. Aljazairy, EA. Hussain, SD
et al. Vitamin B_{12} status and gut microbiota among Saudi females with obesity. Foods.
2022 Dec 11;11(24):4007.

52. Herman, DR. Rhoades, N. Mercado, J. Argueta, P. Lopez, U. Flores, GE. Dietary habits

of 2-to 9-year-old American children are associated with gut microbiome composition.

Journal of the Academy of Nutrition and Dietetics. 2020 Apr 1;120(4):517-34.

53. Guetterman, HM. Huey, SL. Knight, R Fox, AM. Mehta, S. Finkelstein, JL. Vitamin B-

12 and the gastrointestinal microbiome: a systematic review. Advances in Nutrition. 2022

Mar 1;13(2):530-58.

54. Kelly, CJ. Alexeev, EE. Farb, L. Vickery, TW. Zheng, L. Eric, LC. Kitzenberg, DA.

Battista, KD. Kominsky, DJ. Robertson, CE. Frank, DN. Oral vitamin B_{12} supplement is

delivered to the distal gut, altering the corrinoid profile and selectively depleting

Bacteroides in C57BL/6 mice. Gut Microbes. 2019 Nov 2;10(6):654-62.

55. Zhu, X. Xiang, S. Feng, X. Wang, H. Tian, S. Xu, Y. Shi, L. Yang, L. Li, M. Shen, Y.

Chen, J. Impact of cyanocobalamin and methylcobalamin on inflammatory bowel disease

and the intestinal microbiota composition. Journal of agricultural and food chemistry.

2018 Dec 20;67(3):916-26.

56. Park, S. Kang, S. Kim, DS. Folate and vitamin B-12 deficiencies additively impaire

memory function and disturb the gut microbiota in amyloid-β infused rats. International

journal for vitamin and nutrition research. 2019 Dec 16.

57. Wang, C. Zhao, J. Zhang, H. Lee, YK. Zhai, Q. Chen, W. Roles of intestinal bacteroides

in human health and diseases. Critical reviews in food science and nutrition. 2021 Nov

25;61(21):3518-36.

58. Desai, MS. Seekatz, AM. Koropatkin, NM. Kamada, N. Hickey, CA. Wolter, M. et al. A

dietary fiber-deprived gut microbiota degrades the colonic mucus barrier and enhances

pathogen susceptibility. Cell. 2016 Nov 17;167(5):1339-53.

59. Das, P. Babaei, P. Nielsen, J. Metagenomic analysis of microbe-mediated vitamin

metabolism in the human gut microbiome. BMC genomics. 2019 Dec;20:1-1.

617 **Table 1. List of the B$_{12}$ biosynthesis pathways or sub-pathways examined in this study,**

618 **number of MAGs from each group having one of the pathways, and *p-value* of comparison**

619 **of their proportion between groups.**

Pathway/Subpathway name	Short name	MetaCyc ID	Non-deficient	B$_{12}$ Deficient	*p-value*
Adenosylcobalamin biosynthesis (anaerobic)	anaerobic	PWY-5507	27	46	*0.11*
AdoCbl biosynthesis from adenosylcobinamide-GDP I	GDP-1	PWY-5509	63	85	*0.3505*
AdoCbl biosynthesis from adenosylcobinamide-GDP II	GDP-II	PWY-7975	1	1	*_*
AdoCbl salvage from cobalamin	Cobalamin salvage	PWY-6268	6	5	*1*
Superpathway of AdoCbl salvage from cobinamide	Cobinamide salvage	COBALSYN-PWY	44	67	*0.139*
Total MAGs			**135**	**161**	

627 **Table 2. Summary of the metadata collected from samples.**

	Non-deficient (n=29)	B_{12} deficient (n=30)
Serum B_{12} level (pg/ml)	392.83 (±156.73)	161.63 (±30.80)
Age in years (mean± (sd))	6.4 (±2.6)	7.8 (±2.6)
Weight in kg (mean± (sd))	16.3 (±5.0)	18.8 (±4.8)
BMI (mean±(sd))	13.1 (±2.3)	13.9 (±2.1)
Gender	Female=16, Male=13	Female=21, Male=9
Diet	Nonvegetarian=24, Vegetarian=5	Nonvegetarian=30
Antibiotics	No	One subject took antibiotics 2 months before sampling

629
630

Main Figures

Figure 1. Methodology. **A** sampling and **B** computational analysis. BioRender software was

used to create the figure under an academic license.

Figure 2. Alpha (within-sample) diversity trends. **A** Bar plot showing the relative abundance of

detected phyla in the data. The Box plots represent within-sample diversity between Non-

deficient and B_{12} deficient groups: **B** Richness (*p-value = 0.05*), and **C** Shannon diversity (*p-*

value = 0.11).

Figure 3. Beta (between-sample) diversity trends. **A** Principal coordinate analysis of Bray-curtis

distances, **B** Bray-curtis dissimilarity index (*p-value = 0.004*), **C** Jaccard distance (*p-value =*

0.005), and **D** Volcano plot showing differentially abundant species between non-deficient and

B_{12} deficient groups (*p-adj <= 0.05*). Differentially abundant known B_{12} producers have been

highlighted.

Figure 4. B_{12} Biosynthesis pathway abundance in two groups. **A** anaerobic (*p-value = 0.48*), **B**

GDP-I (*p-value = 0.09*), **C** Cobinamide salvage (*p-value = 0.32*). Abundance (median contig

coverage) and Prevalence (%) of potential B_{12} producers in two groups. **D** anaerobic pathway

abundance (*p-value = 0.303*), **E** prevalence (*p-value = 0.007*), **F** GDP-I pathway abundance

(*p-value = 0.01*), **G** prevalence (*p-value = 1.246e-06*), **H** Cobinamide salvage abundance (*p-

value = 0.03*), **I** prevalence (*p-value = 2.817e-06*).

Figure 5. A Cumulative abundance of potential B_{12} producers between two groups, **B**

Relationship between serum B_{12} level and cumulative abundance of B_{12} producers. The y-axis is

the median contig coverage.

Supplementary Tables

Supplemental Table 1. Metadata of samples selected for metagenomic analysis.

Supplemental Table 2. Age-wise prevalence of B_{12} deficiency in tribal children from Balaghat.

Supplemental Table 3. Taxonomic profile of all samples.

Supplemental Table 4. Differentially abundant species in B_{12} deficient group with respect to the

non-deficient group.

Supplemental Table 5. Genes involved in various pathways and subpathways of cobalamin

biosynthesis.

Supplemental Table 6. Taxonomic classification of vitamin B_{12} pathway carrying MAGs

identified in both the groups.

Supplemental Figure 1. Summary of sampled B_{12} level and fecal microbiome sequencing data.

A Distribution of serum B_{12} levels in Non-deficient and B_{12} deficient groups. **B** Distribution of

average reads per sample before and after filtering.

Supplemental Figure 2. Diversity trends. The Box plots represent diversity between Non-

deficient and B_{12} deficient groups: **A** Simpson diversity (*p-value = 0.10*), and **B** Robust aitchison

(*p-value = 0.07*).

Supplemental Figure 3. B_{12} Biosynthesis pathway abundance in two groups. A GDP-II (*p-*

value = 0.47), **B** Cobalamin salvage (*p-value = 0.84)*

Supplemental Figure 4. Locally weighted scatterplot smoothing (LOWESS) of data in Figure

5B showed that the cumulative abundance of potential B_{12} producers (median contig coverage)

increased non-linearly when the serum B_{12} level lies in the deficient range.

Supplemental Figure 5. Taxonomic classification of potential B_{12} producers at the phylum

level in: A Non-deficient group, **B** B_{12} deficient group.

Supplemental Figure 6. The putative Host-microbiome feedback regulation loop **A.** In case of

sufficient Vit. B_{12} bioavailability, the serum Vit. B_{12} is maintained above the deficiency

threshold, wherein the microbiome shares the Vit. B_{12} from the host with a relatively lesser

contribution towards *de-novo* Vit. B_{12} synthesis due to putative inhibitory effect of serum Vit.

B_{12} levels on microbiome B_{12} synthesis. **B.** In the case of insufficient Vit. B_{12} uptake or

transport (due to lack of intrinsic factor), serum Vit. B_{12} levels fall below a threshold that leads

to microbiome remodeling to enhance the abundance of Vit. B_{12} producers and the inhibitory

effect on Vit. B_{12} synthesis is relieved so that it is sourced from the microbiome origin to stall

the severe Vit. B_{12} deficiency.