1 **Tumor Landscape Analysis:**
2 **An Ecologically Informed Framework to Understand Tumor Microenvironments**
3
4 Luis Cisneros[1,3], Merih Deniz Toruner[2], Martin E. Fernandez-Zapico[1], Carlo C. Maley[3], Ryan M. Carr[1]
5
6 [1]Department of Oncology, Mayo Clinic, Rochester, MN
7 [2]Warren Alpert Medical School, Brown University, Providence, RI
8 [3]Arizona Cancer Evolution Center, Center for Evolution and Medicine, and Biodesign Center for Biocomputing,
9 Security and Society, Arizona State University, Tempe, AZ
10
11 **Running Title:** Tumor Landscape Analysis Using Ecological Statistics
12
13 **Corresponding Author:** Ryan M. Carr, 200 First Street SW**,** Rochester, MN 55905, e-mail:
14 carr.ryan@mayo.edu
15
16 **Conflict of Interest Statement:** The authors declare no conflicts of interest.
17
18 **Word Count:** 6,132
19
Number of Figures and Tables: 6

Abstract

Tumor microenvironments (TMEs) are spatially complex and dynamic systems shaped by evolutionary

pressures, tissue architecture, and cellular interactions. To capture this complexity, we developed the Tumor

Landscape Analysis (TLA) pipeline, a computational framework applying principles from landscape ecology

and spatial statistics to quantitatively characterize tumor spatial heterogeneity. TLA integrates spatially

resolved pathology data, including whole-cell, point-based, and region-level formats, and computes multiscale

metrics to assess cell distributions, neighborhood relationships, and tissue-level organization. The framework

leverages ecological indices such as the Morisita-Horn index, Ripley's H function, and Shannon diversity to

quantify intercellular proximity, spatial clustering, and cellular diversity. It also uses the concept of local

microenvironments (LMEs), data-driven ecological niches defined by local cell-type abundance and spatial

uniformity, enabling unsupervised and reproducible classification of tumor regions. The use of fragmentation

metrics, including patch density, shape complexity, and interspersion, provide further insight into spatial

disorganization and emergent tissue architecture. TLA is agnostic to imaging modality and biological context,

supporting broad applicability across tumor types and sample formats. By translating complex tissue

architectures into interpretable spatial metrics, the pipeline enables integrative analyses that link spatial

ecology to clinical and molecular phenotypes. This approach facilitates a deeper understanding of how spatial

features contribute to tumor progression, therapeutic resistance, and clinical outcomes, offering new

opportunities for precision oncology rooted in spatial systems biology.

Introduction

Cancer is a dynamic disease shaped by genetic and epigenomic variations, as well as environmental pressures driving adaptive changes within the tissue ecosystem. Tumor initiation and progression involve continuously evolving interactions among tumor cells, stromal elements, and non-cellular components of the microenvironment. Landscape ecology offers a compelling framework to interpret this spatial complexity (1). Systemic and local regulators, combined with structural constraints, promote clonal divergence, cell migration, metastasis, and the emergence of therapy-resistant subclones (2,3). Therapeutic interventions act as selective pressures, enhancing clonal diversity and resistance (4,5). Moreover, the coevolution between cancer cells and their microenvironment creates a feedback loop that sustains progression despite treatment (6,7). Viewing cancer through an eco-evolutionary lens not only deepens our understanding of tumor behavior but also opens pathways for risk stratification and treatment strategies leveraging cancer's ecological dynamics.

The functional integrity of tissues depends on the spatial coordination of diverse cell types. Disease disrupts these interactions, compromising function and altering spatial organization—much like habitat fragmentation in ecology, which leads to biodiversity loss and systemic environmental shifts (8). This parallel underscores the value of a landscape ecology approach in cancer biology (9). The rise of spatial biology has driven demand for tools capable of analyzing cellular spatial structures and interactions. Advances in multiplexed imaging now allow high-resolution mapping of tissue complexity, enabling deep insights into the tumor microenvironment (TME) (10–12). In this context, the TME can be viewed as a multicellular ecosystem, where intercellular interactions and spatial heterogeneity critically influence tumor progression and therapeutic resistance (13). Traditional spatial assessments rely on qualitative histopathologic review—insightful but labor-intensive and variable (14). Landscape ecology, by contrast, provides quantitative tools to analyze spatial heterogeneity through concepts such as species mixing, parasitism, predator-prey dynamics, and spatial entropy. These metrics can objectively assess cellular diversity, segregation, and interaction patterns (15), revealing non-random spatial relationships. Ecological models adapted for oncology help elucidate immune-tumor interactions, necrosis, and stromal architecture (16,17).

Increasingly, ecological metrics are used to model tumor heterogeneity and population dynamics. The Morisita-Horn index, for instance, quantifies immune-tumor colocalization with prognostic value in breast cancer (18). Computational approaches have further advanced this paradigm: Pacheco et al. (2014) employed evolutionary game theory to model cell dynamics in monoclonal gammopathy (19), while Andor et al. (2016) used tools like EXPANDS and PyClone to map clonal evolution and its correlation with clinical outcomes (20). These tools help reconstruct tumor evolutionary trajectories, offering insights into progression, resistance, and prognosis. Integrating ecological and computational metrics into oncology thus holds promise for translating spatial complexity into clinical utility.

Emerging imaging platforms enable unprecedented resolution of the tumor landscape. Studies by Heindl et al.

and Keren et al. demonstrate how digital pathology and multiplex imaging reveal single-cell interactions within

the TME (21–23). Three-dimensional reconstructions now illuminate invasion patterns and regional

heterogeneity in greater depth (24), supporting not only visualization but also spatially resolved analysis of

tumor architecture.

Despite these advances, applying ecological frameworks to cancer remains methodologically complex. Tumor

tissues are inherently heterogeneous and dynamic, requiring tools capturing spatial and temporal variation. To

meet this need, we developed a Tumor Landscape Analysis (TLA) pipeline, a computational framework

inspired by landscape ecology and environmental systems modeling. TLA integrates spatial statistics and

ecological metrics to quantify heterogeneity across scales, capturing cellular distributions, interclass

relationships, and microarchitectural features within the TME. The pipeline includes data preprocessing,

computation of landscape metrics, and statistical aggregation of spatial features. These outputs can be

integrated with disease variables to define associations between ecological patterns and clinical phenotypes.

By translating tissue architecture into interpretable spatial data, TLA provides a powerful strategy to decode

tumor ecology, offering novel insights into prognosis, disease progression, and therapeutic response.

Materials and Methods

TLA Pipeline Architecture

The TLA pipeline is a computational framework designed to quantify spatial heterogeneity in tumor tissues by

integrating concepts from landscape ecology and spatial statistics (**Figure 1**). It accommodates diverse data

types, including whole-cell segmentations, cell centroid coordinates, and region-level annotations. The pipeline

analyzes individual cell features, computes neighborhood-level spatial profiles, and evaluates tissue

architecture using fragmentation metrics. The framework identifies local microenvironments (LMEs), which

represent discrete ecological niches based on local cell-type abundance and spatial uniformity. These outputs

are synthesized into interpretable spatial maps and multiscale metrics, which can be integrated with clinical

and molecular data to explore associations between spatial architecture and disease phenotypes. TLA is

designed to be broadly applicable across imaging platforms and cancer types, supporting a systems-level

approach to tumor ecology.

Figure 1. Tumor Landscape Analysis (TLA) Workflow showing the different stages of the analysis pipeline. Left panel: Tissue and Cell classification are performed using computer vision assisted by machine learning, and provides the data analyzed in TLA. Middle panel: The lower level of analysis consists of individual cell characterization. Then point statistics is done for cell coordinate data. Regional analysis includes the characterization of regions and local spatial profiles. Finally, with data provided from tissue classification, or local microenvironment (LME) classification based on local profiles, discrete patches, or niches, are defined and used for estimating fragmentation statistics. Right panel: TLA measures that are estimated at the whole-slide scale (including global spatial statistics and fragmentation statistics) are collected and summarized for patient-level analysis that includes statistical modeling, and survival analysis in combination with clinical variables and outcomes.

Data Types and Segmentation Approaches

Accurate segmentation and classification of tissue features are foundational for spatial analysis

(23,24,29,32,33). Digital pathology uses deep learning methods and multiplex imaging to generate spatially

resolved maps of tissues. These computational methods generally fall into three categories based on the scale

and type of segmentation performed, including the identification and classification of whole-cell images, cell

nuclei locations, and multiple-cell tissue component regions.

Cell-Level Spatial Metrics

Cell-level descriptions enable the generation of summary statistics that characterize cellular properties based

on groupings defined by cell type or spatial criteria, such as membership within specific regions of interest,

tissue compartments, or spatially resolved transcriptomic domains (37). These statistics are essential for

quantifying localized phenotypic features relevant to tissue ecology and tumor biology. By linking individual

cellular characteristics to their spatial context, such analyses facilitate detailed inferences about spatial

dynamics and the relationships between cellular behaviors and tissue-level organization.

When digital pathology provides whole-cell detection data, several types of measurements can be derived

directly from the properties of individual cells. Morphological features, including area, perimeter, centroid,

equivalent radius, and circularity, provide information about cell shape and size. Expression characteristics can

be quantified through multiplex imaging, which yields marker-specific intensity values for each cell (22,23,32).

Additionally, cellular context can be assessed through adjacency analyses that quantify the number and types

of neighboring cells, as well as the edge lengths shared between them.

These cell-level features form the basis for multi-scale modeling of the tumor microenvironment and spatially

informed biological hypotheses. Detailed mathematical derivations for calculating equivalent radius, circularity,

and adjacency edge features are provided in the **Supplementary Methods**.

Global Point-Based Spatial Statistics

Point statistics are employed to characterize spatial features when digital pathology data are provided in the

form of cell coordinates. This analytical approach focuses on understanding the spatial distribution of discrete

events (i.e., cells) by comparing observed patterns to a null hypothesis of Complete Spatial Randomness

(CSR). Under CSR, point events are assumed to occur independently and uniformly across space. Deviations

from CSR generally arise due to two fundamental types of spatial effects, first-order effects and second-order

effects (26).

First-order effects reflect large-scale spatial variations in point intensity driven by environmental influences or

tissue architecture. These are typically manifested in differences in point abundance across the landscape.

Quantification of first-order effects includes summary metrics such as the total number of points, class-specific

counts and densities, and the proportional composition of different point classes.

Second-order effects arise from interactions among individual points and influence the fine-scale structure of

the spatial pattern. These effects are assessed using second-order summary statistics that evaluate spatial

relationships between pairs of points, thereby revealing biotic interactions occurring at local scales. Key

second-order metrics include the Morisita-Horn Index, which measures compositional similarity between spatial

populations; the Colocalization Index, adapted from ecological studies to quantify the spatial overlap between

two point classes (18); the Nearest Neighbor Index, which evaluates proximity between different cell types; and

Ripley's H function, which assesses clustering or dispersion patterns across a range of spatial scales.

Mathematical definitions and derivations for all second-order indices are provided in the **Supplementary**

Methods.

Regional-Level and Local Spatial Profiles

When spatial statistics are calculated locally, the result is a two-dimensional spatial map that captures

variations in tissue architecture across a sample. These localized statistical profiles are particularly valuable for

investigating spatial heterogeneity, enabling the identification of microenvironmental differences that may not

be apparent from global analyses.

Local density maps represent spatial profiles of first-order indices, including local cell densities estimated using

kernel smoothing techniques such as Gaussian or Epanechnikov kernels (38,39). These maps can also

incorporate additional spatial features, such as Euclidean distances from each location to tissue compartment

boundaries or other macroscopic structures of biological relevance.

The Getis-Ord Gi*-score is another powerful local metric that detects statistically enriched or depleted regions

for specific cell types (40–42), highlighting localized clustering or exclusion patterns within the tissue.

Furthermore, second-order point statistics—such as colocalization indices, nearest neighbor distances, and

Ripley's H function—can be adapted to local computation. In our methodology, these measures are estimated

using spatial convolution smoothing to perform the necessary summations across neighborhoods. Specifically,

we employ a Gaussian kernel to compute running values of each statistical index in a moving window across

the tissue. This results in a high-resolution spatial profile in which each pixel is assigned a value representing

the local neighborhood estimate of the index. Detailed mathematical formulations and implementation

strategies for the convolutional approach are provided in the **Supplementary Methods**.

Local Microenvironments (LME) Classification

While the optimal definition of tumor microenvironments should ideally incorporate biological insights including

signaling pathways, expression profiles, and functional states, we propose an agnostic, data-driven framework

to define Local Microenvironment (LME) categories using only cell-type abundances and spatial organization

derived from point-location data.

This approach draws on methods from field ecology, where landscapes are divided into quadrats, small,

spatially defined regions used to quantify local population distribution. To adapt this strategy, we apply a

convolution bandwidth parameter that sets the spatial resolution for assessing each cell's neighborhood. For

each "quadrat", we compute the abundance of each cell type and a mixing index, which quantifies spatial

uniformity across "sub-quadrats" within that neighborhood. Instead of rectangular quadrats, we use circular

regions defined by the Gaussian kernel implemented as the spatial convolution approach.

The mixing index is based on a univariate form of the Morisita-Horn index. It compares the observed cell
distribution across sub-quadrats to a uniform (complete spatial randomness) expectation. A high mixing score
indicates even spatial distribution, while a low score reflects clustering or segregation. Additional mathematical
details of the mixing index computation and bandwidth kernel can be found in **Supplementary Methods**.
194
We define LME categories according to the combination of local cell densities and the mixing score. For each
cell type, distributions of abundance and mixing score values are generated from all quadrat scores across all
samples considered in a training set. This set should span the range of all expected microenvironments
relevant to the study. Each of these distributions is then partitioned into three regions, representing low,
medium, and high values. Because the mixing index is not well defined in the context of low-abundance
distributions, we collapsed the nine original categories into three broader classifications that align with our LME
framework. The criteria for this reclassification are as follows: the "B" (Bare) category denotes regions of low
cell abundance, irrespective of the degree of mixing; the "S" (Segmented) category corresponds to moderate
to high cell abundance with low mixing, indicating clustered distributions; and the "M" (Mixed) category
represents moderate to high abundance with high mixing, indicative of more uniformly dispersed populations.
Notably, due to the sensitivity of the mixing index to cell abundance, cases with medium abundance and
intermediate mixing levels are also categorized as "Mixed" (M), since the index tends to underestimate mixing
in the presence of low point counts.
208
Following this classification, each cell type within a defined quadrat is assigned one of the B, S, or M labels. A
composite LME code is then generated by concatenating the labels across all tracked cell types within that
region. In a system containing three distinct cell types, this results in 27 possible LME combinations (e.g., BBB,
BSM, MMM), each representing a unique local ecological state (**Supplementary Methods**). To generate a
spatially resolved map of the tissue landscape, we apply a convolution-based smoothing function, assigning
each pixel the most probable LME category based on its surrounding microenvironment.
215
216   This approach yields a high-resolution classification of emergent spatial domains that are ecologically and
217   biologically meaningful. It enables spatial stratification of tissue samples, identification of niche-level tumor
218   features, and correlation with clinical phenotypes and treatment outcomes.
219
Landscape Fragmentation and Composition Metrics
Tumor tissue can be conceptualized as an ecological mosaic—a spatial assembly of distinct biological niches
or "patches" that correspond to non-overlapping regions defined by their cellular composition and
microenvironmental context (43–46). To quantify the spatial complexity inherent in these tissue landscapes, we
apply fragmentation analysis, a set of metrics designed to capture both the compositional and configurational
properties of categorical tissue maps.

Composition refers to first-order spatial properties and quantifies the proportion of the tissue landscape occupied by each region type, such as tumor, stroma, or immune-dense areas. This provides insight into the relative abundance of different tissue compartments.

Configuration, by contrast, captures second-order spatial characteristics, describing how patches are spatially arranged, their geometric irregularity, and the degree to which they appear fragmented or aggregated within the overall tissue architecture.

This analytical framework is flexible and can be applied to a variety of spatial data sources, including regional-level spatial statistics, maps derived from Local Microenvironmental Ecology (LME) classification, or outputs from segmentation algorithms. The only requirement is that the input consists of a categorical spatial profile composed of discrete, labeled patches, enabling a reproducible and biologically interpretable analysis of tissue organization.

Landscape metrics used to characterize tissue architecture can be broadly categorized into three major classes: geometrical properties, shape metrics, and diversity metrics.

Geometrical properties describe the fundamental physical attributes of spatial patches, such as area, perimeter, and the total edge length at interfaces between adjacent regions. These metrics also include normalized area-to-perimeter ratios. Together, they provide insight into class dominance, rarity, and the extent of potential interfaces for cell-cell or region-region interactions, which may influence tumor-immune dynamics or stromal boundaries.

Shape metrics assess the morphological complexity of individual patches. These include the perimeter-area ratio, which is sensitive to spatial scale; the shape index, which offers a scale-independent measure of shape complexity; and the fractal dimension, which captures deviations from idealized Euclidean forms and approximates contour irregularity. These measures help to quantify the structural complexity of regions within the tumor microenvironment.

Diversity metrics evaluate the variety and distribution of patch types across the tissue. Examples include the Shannon diversity index, patch density, the proportion of the landscape occupied by each class, and contagion, a measure of the degree of interspersion and adjacency between different patch types. Complete mathematical definitions and formulations for shape index, fractal dimension, and other derived metrics are provided in the **Supplementary Methods**.

Landscape fragmentation metrics can be assessed at three spatial scales: the patch, class, and landscape
levels. At the class level, the patch-specific metrics are aggregated to summarize spatial attributes of each
patch type. Key metrics include the total number of patches, cumulative area, and the proportion of the
landscape that each class occupies. The patch density metric, the number of patches per unit area, serves as
a general index of fragmentation, as higher densities often indicate greater disruption of contiguous regions.
The largest patch index further reflects dominance by indicating the proportion of the landscape occupied by
the most expansive patch of a given class. Similarly, edge density normalizes total boundary length to the
landscape area, quantifying the extent of physical interfaces where cell-type mixing or functional transitions
may occur.

At the landscape level, metrics such as the landscape shape index summarize edge complexity across all
classes, while contagion, also referred to as interspersion, estimates the likelihood that neighboring pixels
belong to the same class, thereby capturing spatial aggregation or dispersion of tissue elements. Lastly, the
Shannon diversity index quantifies overall ecological diversity within the tissue by considering both the richness
(number of distinct classes) and the evenness (distributional balance among them), offering a holistic view of
tumor microenvironment heterogeneity. Metric equations, index formulations, and analytical thresholds are
detailed in the **Supplementary Methods**.

Together, these hierarchical metrics provide a comprehensive lens for interpreting the spatial organization of
tumor tissue. For example, a tissue with high patch density but small average patch area likely exhibits
pronounced fragmentation, with frequent shifts in microenvironmental context. In contrast, a low contagion
score would suggest spatial disorganization or fine-grained intermingling of distinct tissue regions, as might be
seen in poorly differentiated tumors or areas of immune infiltration. Conversely, a high contagion or shape
index might indicate well-demarcated regions or organized compartments, such as fibrotic stroma or tumor
nests. Additionally, the Shannon index serves as a sensitive metric for spatial diversity, where high values
reflect complex ecosystems with balanced tissue class distributions, while low values correspond to
monocultures or highly dominant cell states. By synthesizing these spatial features, fragmentation analysis
enables quantitative assessment of tissue architecture, identification of recurring spatial motifs, and potential
correlation with clinical outcomes such as prognosis or treatment response.

Data and Code Availability
All code for data processing and analysis associated with the current submission is available at the CarrLab
GitHub repository: https://github.com/CarrLab/TLA.
All documentation for usage and future updates will be maintained in this web location.

Results

A foundational approach to understanding spatial organization in biological tissues involves analyzing the distributions of discrete entities —such as cells or events— across space. In ecology, spatial point pattern analyses quantify the effects of population dynamics, niche partitioning, and inter-species interactions, which manifest as first-order effects (based on cell density) and second-order effects (based on spatial proximity) (25,26). These same principles apply to tumor biology, where spatial distributions of cells shape microenvironmental conditions that influence tumor progression, metastasis, treatment resistance, and adaptation.

In cancer, cellular spatial patterns are central to the formation of the TME and frame the evolutionary context of disease development. Local patterns of immune infiltration, stromal organization, and necrosis mirror ecological systems, linking single-cell phenotypes to tissue-level functionality. Cellular organization governs signaling, apoptosis, proliferation, and differentiation, and disruptions in these configurations can drive pathological remodeling (27). Viewing tumors as ecological systems motivates the integration of landscape ecology and spatial statistics into digital pathology pipelines (28–31).

To apply landscape statistics to a tissue, the cells first must be identified and classified into different cell types (generally through digital pathology techniques). Then, landscape ecology statistics can be applied to the tissues at four spatial scales: the cell level, the regional/local level, local microenvironments, and landscape fragmentation.

We present results for examples that illustrate outcomes for the three data formats introduced in the methods section. In each case, we show some typical features quantified using our methodology.

Whole-Cell Detection

First, whole-cell detection methods use highly multiplexed imaging to identify and phenotype individual cells across a tissue section. For example, Keren et al. (2018) applied multiplexed ion beam imaging (MIBI) to analyze triple-negative breast cancer samples using a 36-marker panel (22). A convolutional neural network (CNN) was trained to classify cells based on marker expression, producing cell-level phenotype maps with corresponding spatial coordinates. The output for each sample was a rasterized map of individually segmented cells annotated with phenotypic labels, as shown in **Figure 2**.

Figure 2. Whole-cell Detection Using Multiplexed Ion Beam Imaging of Triple Negative Breast Cancer. A. Example of a raster map of individually identified whole cells, colored by phenotype. The whole sample for this methodology is only 1mm x 1mm. B. Small 200um x 200um section showing details of cell contacts. The difference in sizes between cell types is evident. C. Heatmap representing frequencies of adjacency contacts between cancer cells (C), immune cells (I), Fibroblasts (F) and endothelial cells (E). Adjacencies of immune cells with other immune cells correspond to 45.50% of the interactions in this sample, while cancer cells with cancer cells are 28.86% of them, and cancer cells with immune cells are 22.57%. All the rest of the interaction pairings are rare (<1%) (indicated by blue). D. Diagram showing some of the morphological properties calculated for individual cells, including area A, perimeter P, equivalent radius r and centroid (x, y). E. Distributions of cell area and circularity for each cell type class. The circularity is calculated from the area and perimeter as C= "exp" (1- $P^2/4A\pi$), which equals 1 for a circular shape and goes down to 0 for less circular shapes, elongated or very dendritic. In this sample, cancer cells are indeed typically larger in area and more circular than other cell types.

This data is used to make measurements for basic cell morphology, including area and circularity, as well as

cellular context accounted by the frequencies of adjacencies. More statistics can be collected for this level of

analysis, including other cell morphology features, more sophisticated measures of context that weight

adjacencies by the edge length of each contact, and accounting of expression profiles at single-cell levels,

given that multiplex platforms provide such richness of data. Additionally, centroid coordinates can be

estimated geometrically for each cell and used downstream for point statistics, LME classification, and

fragmentation analysis. However, our examples of these analyses use other data sources mainly because the
MIBI platform is limited in the size of the mapped landscape. These samples are only 1mm across, which
constrains the capacity to detect large-scale tissue features and long-range effects. Also, some of the metrics
implemented in these analyses require segmenting the region of interest (ROI) into smaller quadrats to build
spatial distributions, and each of these quadrats needs to contain an appropriate number of cells to grant
statistical robustness; this is not feasible with such a small field of view. Thus, we limit the MIBI analyses to a
few cell-level statistics.
343
In **Figure 2** we show a sample case. Individually identified cells are mapped out and colored by cell type (A
and detail in B). We show examples of the distributions of cell areas (D) and circularity (E) for this slide, as well
as a heatmap diagram showing the frequencies of adjacencies (C). These results demonstrate that tumor cells
in this TNBC sample are consistently larger and more circular than other cell types. Also, lymphocytes tend to
be in contact preferentially with other lymphocytes (as is evident by the large cluster of lymphocytes in the
lower side of the sample), and tumor cells also preferentially interact with other tumor cells, as well as
lymphocytes as a close second.
351
Cell Location Data
Nuclear segmentation has been used to characterize cell types in hematoxylin and eosin (H&E)-stained
histology. In studies of ductal carcinoma in situ (DCIS), CNN-based classifiers were trained to assign nuclear
centroids to specific cell phenotypes (34,35). These analyses identified three major cell populations: epithelial
cells, immune cells, and other stromal cells. The spatial format for each biopsy comprised a map of labeled
nuclear centroids, with each point denoting the location and phenotype of an individual nucleus (**Figure 3**).
358
359 Many measurements can be derived from a point-statistics analysis. This analysis will be published in much
360 greater detail elsewhere, but here we present key examples to illustrate our methodology and to display the
361 behavior of different point statistics measures. Additionally, this type of data can be used downstream for
362 creating LME categories and fragmentation analysis. To keep this article concise, we limit these results to
363 examples of spatial statistics metrics and the LME construction (**Figure 4**).
364
365 In **Figure 3** we can see examples for each of the point-statistics metrics, a schematic that explains extreme
366 cases for each measure, and centroid overlays of specific real cases that represent each of these extremes.
367

Figure 3. Cell-Level Spatial Metrics in DCIS. A. Examples for extreme cases of the Nearest Neighbor Distance index. The top panel row displays a diagram of a well-mixed situation, with NND close to 0, showing that different cell types are typically uniformly distributed with respect to each other. To the right there is a real data example showing the NND of lymphocytes (blue) around tumor cells (red) averaged over tumor cells is low, indicating that lymphocytes are typically intermixed with tumor cells. The second row shows a diagram of a segregated state, in which the NND between red and blue dots are long. To the right a real data example of this situation is presented, in which the separation between tumor cells (red) and lymphocytes (blue) is evident. Both examples correspond to the NND of lymphocytes around tumor cells, which actual value of the index shown in the lower right corners. B. Examples for the Ripley's H index are shown. This is a metric calculated for a specific scale, d=40um. In this case, the separation being quantified is between features of this specific scale size. Thus, the mixed case, upper left diagram, shows features well mixed together at the resolution scale of the circle. The lower left diagram shows the separation between the point classes at that same scale. In real data examples, in which we can see groups of tumor cells (red) and groups of lymphocytes (blue) intermixed together (upper right) and separation between clusters regardless of the scale (lower right). Actual index values are shown in the inset box. Curves of the value of the index for larger scale values are shown to the far right in both cases, showing that in the first example the index increases slightly with d and then drops after a scale of 200um, while in the lower example it consistently drops, but not a lot. It is generally expected to observe lower values at longer scales as small features get smoothed out at such scales. C. Examples for the Colocalization Index, which quantifies the similarity between two spatial distributions of points. In the upper case, the diagram shows two segregated distributions of points, with a value of the index close to zero, which can be observed in the real example on the right-hand side. Whereas the lower example shows a situation in which the two distributions are like each other, and an index close to one. The real case example for this case is not obvious. Similarly to the RHF index, there is a set scale for the calculation of this index, in this case given by the size of the bins used to construct a discrete spatial distribution (i.e. as a histogram). The scale used for these examples is 40 μm. Even though lymphocytes (blue) are not exactly infiltrated in clusters of tumor cells (red), at the scale of smoothing the regions where there is a high density of tumor cells also have a higher density of lymphocytes. This example is important to indicate a caveat in using this metric as a measure of infiltration.

Nearest-neighbor Distance Index (NND) (**Figure 3A**): the diagrams show that a low value of this index is a

signature of CSR, meaning that the two cell types considered (in these cases they are DCIS cells and

lymphocytes) are well mixed and the frequency of finding a lymphocyte next to a DCIS cell is like the frequency

of finding another DCIS cell. This effect is evident in the detail shown on the right-hand panel, where we can

see that lymphocytes are thoroughly mixed and infiltrated in the clouds of DCIS cells. On the other hand, large

values of NND indicate that the typical distance from DCIS cells to the closest lymphocytes is larger than the

distance to the closest DCIS cell, hence representing a case of segregation of lymphocytes away from DCIS

cells. This can be seen in the right-side detail panel where lymphocytes are excluded from the DCIS cell

clusters.

Ripley's H function (RHF) (**Figure 3B**): in this case again, a low extreme is indicative of CSR. An important

distinction is that the properties conveyed by this metric have a characteristic scale defined by the radius used

to calculate H. A way to understand this metric is similar to the principle for NND, but instead of considering

just the first neighbor, count the ratio of the number of points of each type inside a circle of radius d. Therefore

the "mixing" or "clustering" is defined with respect to this scale. For this particular example H was calculated at

a scale of $d = 40\mu\mathrm{m}$ (a small green circle in the lower left corner shows the dimension of the scale used). In

this sense, cells could be clustered forming features, and this metric could still have a low value if those

features are smaller than the scale d and they, as coarse-graining features instead of the individual cells, are

well mixed and satisfy a CSR condition. We can see this in our example, on the right-hand side we see

clusters of DCIS cells mixed with lymphocytes providing a well-mixed condition at the scale used, though at the

cell scale, this example would show separation between the two populations. In contrast, the other extreme

case clearly shows a separation between the populations indicating a typically large distance from DCIS cells
and groups of lymphocytes.

Colocalization Index (MHI) (**Figure 3C**): for the case of the colocalization index, measured employing the
Morisita-Horn index. The high extreme ($M \sim 1$) is not necessarily a case of CSR, though CSR would certainly
have this result. In a more general sense, what this metric indicates is the similarity between the spatial
distributions. If the spatial distribution of the two cell types is not uniform, but are similar to each other, we will
observe this extreme, as illustrated in the lower panel. The scale of the quadrats used, $750 \times 750 [\mu m^2]$, comes
into play in much the same way as the scale for RHF. As we see in the real data example, lymphocytes follow
the spatial distribution of DCIS cells, suggested by the fact that regions with large concentrations of DCIS cells
also have larger concentrations of lymphocytes, even though at lower scales they are not infiltrated. On the
other hand, the other extreme $M \sim 0$ reflects a condition of total separation of the two populations, at the scale
of the spatial distribution binning, as is clearly shown in the example case.

Region Detection Data
We present region-level results from two studies. The first one is comprised of regional results derived from the
point-statistics analysis in our DCIS example (see previous section). In **Figure 4** we show the spatial profiles
for local measurements of NND, RHI, and MHI, showing how these indices can be estimated in small regions
to represent the variations in space.

Figure 4. Regional-Level and Local Spatial Profiles in DCIS. A. Hot and cold density regions estimated using the Getis-Ord Gi* statistic, for each of the cell phenotypes in the DCIS study. Regions colored red are hot spots where the density of cells is larger than expected according to the total cell density of the sample (i.e. statistical significance using a z-score test). Green indicates regions where the cell density is within the range of expected values. Blue regions (not observed here) indicate cold spots where cell density is lower than expected. B. Local Spatial Profiles for the Colocalization Index are shown for two examples, between lymphocytes and fibroblasts and between cancer cells and lymphocytes. The colocalization index is calculated in the neighborhood around each pixel using a spatial convolution with a Gaussian kernel with a 20um bandwidth. In this way, these profiles represent local variations of these metrics, in this case showing that most of the slide has moderate to low values of colocalization, with some small regions where the distributions of cell types are similar. C. Local Spatial Profiles for the Ripley's H function of fibroblasts around lymphocytes and lymphocytes around tumor cells, calculated locally using a spatial convolution method with a Gaussian kernel with a 20um bandwidth. The extent of the slide shows mostly intermediate values, with some larger values in the periphery of the middle region, where tumor clusters are, indicating that segregation of cell types happens away from those areas. D. Local Spatial Profiles for the Nearest Neighbor Distance index of fibroblasts around lymphocytes and lymphocytes around tumor cells, calculated locally using a spatial convolution method with a Gaussian kernel with a 20um bandwidth. Like panel C, larger values indicate segregation of cell types whereas small values indicate closer proximity between cell types.

For the spatial profile of NND of DCIS cells around lymphocytes, we can see a region of negative values,

whereas in most other regions we see values typically close to zero. A negative value of this metric indicates

that DCIS cells (test cells) are closer to any lymphocytes (reference cells) than other lymphocytes, this

happens when the density of reference cells is small while that of test cells is large, meaning that test cells are

common in the region and reference cells are individually infiltrated and surrounded by test cells. On the other

hand, a positive score indicates clustering of test cells around reference cells, while close to zero values

indicate an even distribution. In this sample, the distributions of DCIS cells and lymphocytes are pretty even

across the tissue landscape. For the local Ripley's H score, we show an example of lymphocytes around other

lymphocytes that show regions of clustering of these types of cells. Notice that the reference cell density is

calculated locally, thus the local values of H reflect clustering within regions (which is a circle

with $r = 0.75\mathrm{mm}$). This profile is thus a display of the intensity of local relative variations of the density of

lymphocytes. Finally, for the colocalization index, we present an example of the overlap of DCIS cells and

lymphocytes, showing that for the most part, the distributions of these cell types are very similar across the

sample, with some variations in regions on the edges of it that display moderate separation of the distribution

patterns, mostly driven by low densities of lymphocytes.

In addition to the three basic point statistic metrics, we also estimated the profile for the Getis-Ord Gi* statistic

for the density of lymphocytes. This is a regional statistic that detects locations in which there is higher, or

lower than expected cell density for a particular cell type. We show specifically the regions that are statistically

significant for the z-statistic, corresponding to hot (red) and cold (blue) regions for cell density, while green

regions indicate expected densities. Interestingly, this sample shows quite a lot of local clustering across the

sample, including a middle region where DCIS cells are concentrated.

Finally, in landscape ecology, ecosystems are often decomposed into spatially distinct biological niches,

referred to as "ecologic phenotypes" or LME, which represent specific combinations of species coexisting

within environmental contexts. Analogously, tumor tissues can be understood as ecological mosaics composed

of functionally distinct regions, each shaped by local cellular composition and spatial configuration. We present

an example of the spatial profile of LME categories defined by the local abundance and mixing scores for each

cell type, as described in the methods (**Figure 5**). We can see that this strategy is capable of differentiating

regions with different population configurations as different types of micro-ecologies. There is a main region

with profiles for all DCIS cells that are both abundant and mixed with segregated lymphocytes (red region) and

a region with segregated DCIS cells and lymphocytes (green area) (**Figure 5B**). This separation is evident

from the point distribution profiles separated by classes (**Figure 5C**).

Figure 5 Local Microenvironments (LME). A. Local values of cell abundance and mixing for all samples in the study are collected and summarized. Distributions of values are used to assign low, medium and high levels, as observed collectively in all samples. B. Example of LME classification based on local values of cell abundance and mixing. C. Quantified frequency of LME subtypes of the example landscape in panel B. (S) Segmented environments are those in which cells are clustered together; they have moderate to high abundance and low mixing. (M) Mixed environments are those where cells are mixed uniformly; they have moderate to high abundance and high mixing. Because the mixing score is sensitive to low abundance, medium abundance levels are considered "mixed" (M) if the mixing is medium, as mixing is biased to lower values with lower abundances. With its scheme, (B, S, M) codes are assigned for each individual class, forming categories like: `BBB, BBS, BBM …. MMM`. Each of these categories represents a basic type of Local Micro-Environment that is consistent across the entire study and can be used to define ecological phenotypes in the TLA.

As an unsupervised and quantitative framework, LME analysis provides a reproducible and scalable method

for partitioning tumor microenvironments using only spatially resolved cell type data. While it is not intended to

replace biologically curated models, this strategy offers a principled way to delineate tumor architecture and

immune dynamics, especially when integrated with molecular profiling or clinical annotation.

All these spatial profiles can be further analyzed using fragmentation analysis tools to measure the

morphological and configuration properties of the "habitats" in the sample. For this, it is necessary to convert

them into discrete maps composed of a set of patches or local habitats. This is already the case for the Getis-

Ord hot/cold region profiles and the LME profiles. In the case of the local statistics profiles, a possibility is to

stratify the continuous values into discrete contour levels and use those as the patches for the fragmentation

analysis. In this sense, if we visualize these profiles as surface landscapes with mountains and valleys, the

fragmentation analysis of contour levels is a way to assess the ruggedness or smoothness of such a surface.

Fragmentation Statistics

Our last example is the regional classification data of PDAC resections. In this case, discrete regions

representing different tissue compartments are identified using machine-learning image processing. The

regional segmentation data does not allow for point-statistics analysis, but it can be processed using our

fragmentation analysis approach. In **Figure 6** we show some examples of fragmentation metrics that were

significant in this study. Here we present extreme example profiles for the Shape Index, the Patch Density, and

Contagion metrics. In this study, these metrics were predictive of the time to recurrence after neoadjuvant

therapy in PDAC. The shape index is a normalized measure of the perimeter-to-area ratio of the patches

conforming to the landscape. A large value of this index indicates that the typical patches have irregular and

complex shapes which can be an indication of fragmentation and suggests a dynamic in which signaling and

other processes of transport or exchange of chemical substrates between niches. We see that the lower

extreme case shows typically small patches with globular and compact shapes, while the higher extreme case

shows elongated and complex-shape patches. The nature of interactions between patches would be radically

different in these two cases. The patch density is simply a measure of the number of patches per unit area.

The extreme cases are clear, showing a case with a moderate number of patches, especially stroma, for the

lower extreme and a very rich and populated landscape for the higher extreme, with many small tumor patches

and a more fragmented stroma. Finally, contagion is a metric that quantifies the level of cohesiveness of the

landscape, expressed as the probability that two randomly picked neighboring locations are of the same class.

In the lower extreme case, we could see that depending on the location in the landscape, the two types of
regions can be very intertwined, especially in the middle of the sample, suggesting that it's generally likely that
two random neighboring locations would be of different classes. The opposite is true for the other extreme
case. In which, though the tumor stroma is very segmented, the stroma is much more prevalent in space and
two random neighboring locations will likely be stroma. It's important to point out that this measure is
normalized for the abundance of each class, but a high value of contagion indicates the likelihood of two
neighbors being of the same class, but it says nothing about the balance of the probabilities of different
classes.

Third, region detection approaches segment larger structural compartments of the tissue. In pancreatic ductal
adenocarcinoma (PDAC), a machine learning model was used to delineate tumor and stromal regions in post-
treatment surgical resection samples (36). This region-level segmentation enabled the generation of spatial
maps that facilitate downstream analyses, including the identification of specific cell types within distinct tissue
compartments and investigations of spatial relationships, such as proximity to the basement membrane or
blood vessels (**Figure 6**).

Figure 6. Landscape Fragmentation and Composition Metrics in PDAC. A. The Contagion Index: this is a metric of diversity and complexity of a landscape mosaic that quantifies the probability that two randomly chosen neighboring locations (pixels) belong to the same patch class, thus measuring the level of connectivity, or conversely fragmentation, of the landscape. These are two case examples showing extreme high (upper case) and extreme low (lower case) of this metric in the PDAC study data, and explanation diagrams (right hand panels). B. The patch density: the number of patches, of any class, per unit area. Two examples and explanation diagrams are shown. C. The shape index: this is a morphology measure that accounts for the relation of perimeter and area that is adjusted for size. Since the perimeter of a patch scales linearly with its length scale, while the area scales as the square of it, this measure is defined as a fourth of the perimeter divided by the square root of the area and is a measure of shape independent of scale. Compact, circular or squared, shapes have low perimeter to area ratio, and thus have low shape index values, while more complex, or dendritic, shapes have longer perimeters for the same area, thus having large shape index. Examples and diagrams of extreme cases are shown.

Discussion

Discussion

Interactions between different cell types in the TME shape disease progression and influence therapeutic
outcomes. Those interactions and the ecology of the TME can be studied through spatial analyses of the
different cell types. While technological advances in spatial biology have enabled increasingly detailed
characterizations of tissue architecture, many current analytic approaches still rely heavily on reductionist
models or narrowly focused metrics. In this study, we present a comprehensive analytical framework, the
Tumor Landscape Analysis (TLA) pipeline, that adapts core principles from landscape ecology to better
capture the complexity of tumor ecosystems. This framework moves beyond conventional spatial biology by
offering a multiscale, quantitative, and ecologically informed view of the TME.

Conventional spatial analyses typically depend on identifying specific cell types and evaluating their
relationships through simple distance metrics, adjacency maps, or nearest-neighbor interactions. These
methods, while informative, often operate in isolation from the broader context of the tissue's spatial
organization. They rarely account for higher-order spatial properties such as fragmentation, aggregation, or
habitat structure—features that are central to ecological systems and, by extension, highly relevant to tumor
biology. The TLA framework addresses these limitations by incorporating a broader spectrum of spatial
statistics, adapted from ecological theory. Metrics such as the Morisita-Horn index, Ripley's H function, and the
Shannon diversity index allow for the characterization of tissue landscapes in terms of both cellular
composition and spatial configuration. Rather than focusing exclusively on pairwise cell interactions or
compartmental delineations, TLA provides a means to assess how cells and regions are distributed, how they
interact across distance scales, and how these patterns may reflect underlying biological processes such as
immune infiltration, stromal remodeling, or clonal expansion.

One of the key innovations introduced here is the concept of Local Microenvironments (LMEs), defined not by
prior biological assumptions but by empirical patterns of cell abundance and spatial uniformity. This
unsupervised classification yields discrete ecological niches within the tissue, offering a reproducible and
interpretable means of partitioning tumor architecture. Importantly, this approach is data-driven yet biologically
intuitive: regions identified as "segmented," "mixed," or "bare" correspond closely with phenomena observed in
pathology, such as lymphocyte clustering, immune-excluded zones, or necrotic cores. In parallel, the
application of fragmentation metrics, long used in landscape ecology to describe habitat loss and
environmental disruption, provides a novel lens through which to assess tumor morphology. Measures such as
patch density, shape complexity, and interspersion offer insight into the degree of spatial disorganization within
a tissue. These attributes are not merely descriptive; they have the potential to serve as spatial biomarkers,
capturing aspects of tumor behavior that may correlate with aggressiveness, treatment resistance, or
recurrence risk. They are likely to change dramatically with treatment.

Perhaps most significantly, TLA allows for integration across data types and spatial scales. Whether analyzing whole-cell segmentations, centroid point maps, or region-level classifications, the framework applies a consistent set of spatial tools that enable comparisons across tissue types, disease states, and imaging platforms. This adaptability is essential for translating spatial biology into clinical practice, where tissue samples vary in quality, modality, and resolution. Looking ahead, the potential applications of this framework are substantial. When applied to longitudinal or post-treatment samples, TLA could facilitate the reconstruction of tumor evolution under therapeutic pressure, revealing how spatial features emerge, shift, or regress. Its integration with molecular or transcriptomic data could yield composite ecological-genomic signatures, providing a richer context for understanding tumor behavior and cell-cell interactions. Moreover, TLA-derived metrics could be incorporated into predictive models or decision-support tools aimed at guiding therapy selection or patient stratification.

Taken together, the Tumor Landscape Analysis framework represents a shift in how we conceptualize and quantify tumor ecosystems. By drawing on principles from landscape ecology, it enables a more holistic understanding of spatial heterogeneity, one that captures not only what cell types are present in the TME, but how they are organized, how they interact, and what patterns emerge from their collective behavior. This perspective is both timely and necessary, as oncology continues to move toward more personalized, spatially informed models of care.

Acknowledgments

This work was also supported by the Gerstner Family Foundation Career Development Award, the Grand Forks Career Development Award, and the Mayo Clinic Center for Clinical and Translational Science (CCaTS) through a Small Grant (UL1TR000135). Ryan Carr gratefully acknowledges this funding support, which was instrumental in the development and execution of this study.

References

1. Urban DL, O'Neill RV, Shugart HH Jr. Landscape Ecology. Bioscience. Oxford University Press (OUP); 1987;37:119–27.

2. Greaves M, Maley CC. Clonal evolution in cancer. Nature. Springer Science and Business Media LLC; 2012;481:306–13.

3. Maley CC, Aktipis A, Graham TA, Sottoriva A, Boddy AM, Janiszewska M, et al. Classifying the evolutionary and ecological features of neoplasms. Nat Rev Cancer. Nature Publishing Group; 2017;17:605–19.

4. Marusyk A, Janiszewska M, Polyak K. Intratumor heterogeneity: The Rosetta stone of therapy resistance. Cancer Cell. Elsevier BV; 2020;37:471–84.

5. Read A. Chapter 23 – ecology, evolution, and the cancer patient. Ecology and Evolution of Cancer. Elsevier Inc.; 2017;255–7.

6. Ibrahim-Hashim A, J. Gillies R, S. Brown J, A. Gatenby R. Coevolution of tumor cells and their microenvironment: "niche construction in cancer." Ecology and Evolution of Cancer. Elsevier; 2017. page 111–7.

7. Angelova M, Mlecnik B, Vasaturo A, Bindea G, Fredriksen T, Lafontaine L, et al. Evolution of metastases in space and time under immune selection. Cell. Elsevier BV; 2018;175:751-765.e16.

8. Saunders D, Hobbs RJ, Margules C. Biological consequences of ecosystem fragmentation: A review. Conservation Biology. [Wiley, Society for Conservation Biology]; 1991;5:18–32.

9. Merlo LMF, Pepper JW, Reid BJ, Maley CC. Cancer as an evolutionary and ecological process. Nat Rev Cancer. Springer Science and Business Media LLC; 2006;6:924–35.

10. Moffitt JR, Lundberg E, Heyn H. The emerging landscape of spatial profiling technologies. Nat Rev Genet. Springer Science and Business Media LLC; 2022;23:741–59.

11. Bressan D, Battistoni G, Hannon GJ. The dawn of spatial omics. Science. American Association for the Advancement of Science; 2023;381:eabq4964.

12. Bull JA, Moore JW, Mulholland EJ, Leedham SJ, Byrne HM. MuSpAn: A Toolbox for Multiscale Spatial Analysis [Internet]. bioRxiv. 2024 [cited 2025 Mar 20]. page 2024.12.06.627195. Available from: https://www.biorxiv.org/content/10.1101/2024.12.06.627195v3.abstract

13. Daoust SP, Fahrig L, Martin AE, Thomas F. From forest and agro-ecosystems to the microecosystems of the human body: what can landscape ecology tell us about tumor growth, metastasis, and treatment options? Evol Appl. Wiley; 2013;6:82–91.

14. Bridge P, Fielding A, Rowntree P, Pullar A. Intraobserver variability: Should we worry? J Med Imaging Radiat Sci. Elsevier BV; 2016;47:217–20.

15. Colombo EH, Martínez-García R, López C, Hernández-García E. Spatial eco-evolutionary feedbacks mediate coexistence in prey-predator systems [Internet]. arXiv [q-bio.PE]. bioRxiv; 2019. Available from: http://arxiv.org/abs/1902.03016

16. Lloyd MC, Rejniak KA, Brown JS, Gatenby RA, Minor ES, Bui MM. Pathology to enhance precision medicine in oncology: lessons from landscape ecology. Adv Anat Pathol. 2015;22:267–72.

17. Yuan Y. Spatial Heterogeneity in the Tumor Microenvironment. Cold Spring Harb Perspect Med. 2016;6:a026583.

18. Maley CC, Koelble K, Natrajan R, Aktipis A, Yuan Y. An ecological measure of immune-cancer colocalization as a prognostic factor for breast cancer. Breast Cancer Res. 2015;17:131.

19. Pacheco JM, Santos FC, Dingli D. The ecology of cancer from an evolutionary game theory perspective. Interface Focus. The Royal Society; 2014;4:20140019.

20. Andor N, Graham TA, Jansen M, Xia LC, Aktipis CA, Petritsch C, et al. Pan-cancer analysis of the extent and consequences of intratumor heterogeneity. Nat Med. Springer Science and Business Media LLC; 2016;22:105–13.

21. Heindl A, Nawaz S, Yuan Y. Mapping spatial heterogeneity in the tumor microenvironment: a new era for digital pathology. Lab Invest. Nature Publishing Group; 2015;95:377–84.

22. Keren L, Bosse M, Marquez D, Angoshtari R, Jain S, Varma S, et al. A structured tumor-immune microenvironment in triple negative breast cancer revealed by multiplexed ion beam imaging. Cell. Elsevier BV; 2018;174:1373-1387.e19.

23. Keren L, Bosse M, Thompson S, Risom T, Vijayaragavan K, McCaffrey E, et al. MIBI-TOF: A multiplexed imaging platform relates cellular phenotypes and tissue structure. Sci Adv. American Association for the Advancement of Science (AAAS); 2019;5:eaax5851.

24. Kiemen AL, Braxton AM, Grahn MP, Han KS, Babu JM, Reichel R, et al. CODA: quantitative 3D reconstruction of large tissues at cellular resolution. Nat Methods. Springer Science and Business Media LLC; 2022;19:1490–9.

25. Jia G, Yu X, Fan D, Jia J. Mechanism underlying the spatial pattern formation of dominant tree species in a natural secondary forest. PLoS One. Public Library of Science (PLoS); 2016;11:e0152596.

26. Ben-Said M. Spatial point-pattern analysis as a powerful tool in identifying pattern-process relationships in plant ecology: an updated review. Ecol Process [Internet]. Springer Science and Business Media LLC; 2021;10. Available from: http://dx.doi.org/10.1186/s13717-021-00314-4

27. Wei C-J, Xu X, Lo CW. Connexins and cell signaling in development and disease. Annu Rev Cell Dev Biol. Annual Reviews; 2004;20:811–38.

28. Nam S, Chong Y, Jung CK, Kwak T-Y, Lee JY, Park J, et al. Introduction to digital pathology and computer-aided pathology. J Pathol Transl Med. The Korean Society of Pathologists and The Korean Society for Cytopathology; 2020;54:125–34.

29. Barisoni L, Lafata KJ, Hewitt SM, Madabhushi A, Balis UGJ. Digital pathology and computational image analysis in nephropathology. Nat Rev Nephrol. Springer Science and Business Media LLC; 2020;16:669–85.

30. Giansanti D. The artificial intelligence in digital pathology and digital radiology: Where are we? Healthcare (Basel). MDPI AG; 2020;9:30.

31. Hanna MG, Ardon O, Reuter VE, Sirintrapun SJ, England C, Klimstra DS, et al. Integrating digital pathology into clinical practice. Mod Pathol. Elsevier BV; 2022;35:152–64.

32. Greenwald NF, Miller G, Moen E, Kong A, Kagel A, Dougherty T, et al. Whole-cell segmentation of tissue images with human-level performance using large-scale data annotation and deep learning. Nat Biotechnol. Springer Science and Business Media LLC; 2022;40:555–65.

33. Aeffner F, Zarella MD, Buchbinder N, Bui MM, Goodman MR, Hartman DJ, et al. Introduction to digital image analysis in whole-slide imaging: A white paper from the digital pathology association. J Pathol Inform. Elsevier BV; 2019;10:9.

34. Sobhani F, Muralidhar S, Hamidinekoo A, Hall AH, King LM, Marks JR, et al. Spatial interplay of tissue
hypoxia and T-cell regulation in ductal carcinoma in situ. NPJ Breast Cancer. Nature Publishing Group;
2022;8:105.

35. Zhang H, AbdulJabbar K, Grunewald T, Akarca AU, Hagos Y, Sobhani F, et al. Self-supervised deep
learning for highly efficient spatial immunophenotyping. EBioMedicine. Elsevier; 2023;95:104769.

36. Diao JA, Wang JK, Chui WF, Mountain V, Gullapally SC, Srinivasan R, et al. Human-interpretable image
features derived from densely mapped cancer pathology slides predict diverse molecular phenotypes. Nat
Commun. Springer Science and Business Media LLC; 2021;12:1613.

37. Sun S, Zhu J, Zhou X. Statistical analysis of spatial expression patterns for spatially resolved
transcriptomic studies. Nat Methods. Springer Science and Business Media LLC; 2020;17:193–200.

38. Rosenblatt M. Remarks on some nonparametric estimates of a density function. Ann Math Stat. Institute of
Mathematical Statistics; 1956;27:832–7.

39. Parzen E. On estimation of a probability density function and mode. Ann Math Stat. Institute of
Mathematical Statistics; 1962;33:1065–76.

40. Songchitruksa P, Zeng X. Getis–Ord spatial statistics to identify hot spots by using incident management
data. Transp Res Rec. SAGE Publications; 2010;2165:42–51.

41. Ord JK, Getis A. Local spatial autocorrelation statistics: Distributional issues and an application. Geogr
Anal. Wiley; 1995;27:286–306.

42. Getis A, Ord JK. The analysis of spatial association by use of distance statistics. Geogr Anal. Wiley;
1992;24:189–206.

43. Mcgarigal, K., Cushman, S., & Ene, E. FRAGSTATS v4: Spatial Pattern Analysis Program for Categorical
Maps. 2012;

44. Hesselbarth MHK, Sciaini M, With KA, Wiegand K, Nowosad J. landscapemetrics: an open-source R tool
to calculate landscape metrics. Ecography (Cop). 2019;42:1648–57.

45. Nowosad J, Stepinski TF. Information theory as a consistent framework for quantification and classification
of landscape patterns. Landsc Ecol. 2019;34:2091–101.

46. Bosch M. PyLandStats: An open-source Pythonic library to compute landscape metrics. PLoS One. Public
Library of Science (PLoS); 2019;14:e0225734.

A Nearest Neighbor Distance

Well Mixed, Random, NND~0

Segregated, NND>0

B Ripley's H Function

Well Mixed, Random, RHF~0

Clustering of 🔴 around 🔵 , RHF>0

C Colocalization Index

CI~0

CI~1

Cancer cells • Lymphocytes • Fibroblasts

NND=0.2355

NND=1.4101

RHF=0.0084

RHF=0.8284

CI=0.0974

CI=0.9031

Distance (mm)

RHF(c,l)

r (μm)

A

Fibroblasts
(μ=123.04, σ=126.96)

Lymphocytes
(μ=122.36, σ=277.14)

DCIS Cells
(μ=239.90, σ=319.35)

Cells per Quadrant

Fibroblasts
(μ=0.75, σ=0.18)

Lymphocytes
(μ=0.60, σ=0.18)

DCIS Cells
(μ=0.74, σ=0.17)

Mixing Index per Quadrant

B

C

LME Categories (tlf)

- MMM
- MMS
- MMB
- MSM
- MSS
- MSB
- MBM
- MBS
- MBB
- SMM
- SMS
- SMB
- SSM
- SSS
- SSB
- SBM
- SBS
- SBB
- BMM
- BMS
- BMB
- BSM
- BSS
- BSB
- BBM
- BBS
- BBB

Cancer **Stroma**

A Contagion Index

Contagion Index High

Congation Index Low

B Patch Density

Patch Density High

Patch Density Low

C Shape Index

Shape Index High

Shape Index Low

Distance (mm)