Manuscript

40
41
42 **Characterization of iGABASnFR2 for in vivo mesoscale imaging of intracortical GABA**
43 **dynamics**
44

45 **Authors:** Edris Rezaei*[1],Setare Tohidi[1], Mojtaba Nazari-Ahangarkolaee[1],Javad Karimi
46 Abadchi[2]
47
48 [1]Canadian Centre for Behavioural Neuroscience, University of Lethbridge; Lethbridge, AB,
49 Canada
50
51 [2]Department of Psychiatry, the Douglas Research Centre, McGill University, Montréal, QC,
52 Canada
53
54 *** Corresponding and first author:**
55
56 Edris Rezaei (edris.rezaei@uleth.ca)
57
58 **Address:**
59 Edris Rezaei, Department of Neuroscience, University of Lethbridge, 4401 University Dr W,
60 Lethbridge, AB T1K 3M4, Canada

80
81
Abstract

While genetically encoded sensors have advanced the study of cortical excitation, tools for large-
scale imaging of inhibition remain limited. Visualizing extracellular GABA dynamics in vivo is
essential for understanding how inhibitory networks shape brain activity across sensory,
behavioral, and pharmacological states. To validate and apply the genetically encoded sensor
iGABASnFR2 for wide-field imaging of extracellular GABA, and to characterize how cortical
inhibition reorganizes across brain states, sensory modalities, and after GABA transporter
blockade. We performed mesoscale imaging in head-fixed C57BL/6 mice systemically
expressing iGABASnFR2. Recordings were conducted under isoflurane anesthesia, during quiet
wakefulness, natural sleep (NREM and REM), and after administration of the GAT-1 inhibitor
Tiagabine. We analyzed both sensory-evoked and spontaneous GABA signals using time-series,
spectral, and seed-pixel correlation analyses. iGABASnFR2 demonstrated strong and modality-
specific GABAergic responses to sensory stimulation, with faster and stronger activation in the
contralateral cortex. Although the general spatial patterns of sensory-evoked GABA responses
were consistent across anesthesia and quiet wakefulness, the amplitude, timing, and spread of
these responses were significantly greater during wakefulness. During spontaneous activity,
cortical GABA levels and connectivity modulated by brain state: GABA amplitude and
interhemispheric synchrony were highest during quiet wakefulness but reduced during NREM
sleep. Tiagabine elevated baseline GABA levels, abolished stimulus-evoked responses, and
enhanced local and long-range inhibitory synchrony. iGABASnFR2 enables reliable, high-
resolution imaging of cortical GABA dynamics in vivo. These results demonstrate that inhibitory
signaling is dynamically structured across brain states and can be pharmacologically modulated.
This tool offers new opportunities to explore the role of inhibition in health and disease at the
mesoscale level.

Keywords: iGABASnFR2; GABA; wide-field imaging; inhibition; spontaneous activity.

114 **Introduction**
115
The nervous system is composed of two basic cell types: neurons and glial cells. Neurons can be
categorized into excitatory and inhibitory types. Excitatory neurons primarily release glutamate
which is the major neurotransmitter in the nervous system to facilitate communication between

neurons across different brain regions[1], while inhibitory neurons mainly release gamma-
aminobutyric acid (GABA) to stabilize neural networks by balancing excitatory activity and
preventing excessive neuronal firing[2]. Maintaining this precise balance between excitation and
inhibition is essential for sensory processing, memory formation, and motor control[3-4]. Through
temporally and spatially precise modulation of neuronal activity, inhibitory signaling contributes
to a wide range of brain functions, including sensory processing, circuit refinement, and the
regulation of oscillatory dynamics[5-9]. Disruptions in GABAergic signaling underlie a variety of
neurological and psychiatric disorders, including epilepsy, schizophrenia, and autism spectrum
disorders, highlighting the clinical importance of understanding GABAergic modulation in both
health and disease[10-16]. Despite the importance of GABA in regulating cortical activity, direct,
real-time visualization of GABA dynamics in vivo remains unknown. The development of
genetically encoded neurotransmitter sensors has significantly expanded the ability to monitor
neural activity across spatial and temporal scales[17]. For instance, the glutamate sensor iGluSnFR
enabled high-resolution, real-time imaging of excitatory neurons, providing insights into cortical
connectivity and sensory-evoked activity in awake and anesthetized animals[18-19]. Building on
advances in sensor engineering, the genetically encoded fluorescent sensor iGABASnFR2 was
developed to detect extracellular GABA dynamics[20]. This sensor permits real-time, in vivo
monitoring of extracellular GABA, providing a powerful tool for visualizing GABA's spatial and
temporal dynamics across large-scale cortical regions. Wide field imaging with iGABASnFR2
enables comprehensive mapping of inhibitory circuits. However, a detailed characterization of
iGABASnFR2 functionality across different brain states, sensory modalities, and
pharmacological conditions is still needed. In this study, we comprehensively characterized
spontaneous and sensory-evoked GABAergic activity across the dorsal cortex using wide-field
imaging of iGABASnFR2 sensor in natural sleep, awake mice, and under 1% isoflurane
anesthesia. Additionally, we assess the effects of Tiagabine, a GABA reuptake inhibitor, to
further characterize sensor performance and demonstrate its sensitivity to pharmacological
manipulation of cortical GABA levels. These results not only validate the robustness and
sensitivity of iGABASnFR2 as a practical tool for monitoring GABAergic neurotransmission in
vivo but also lay a solid foundation for future studies into inhibitory dynamics in health and
disease, including conditions like epilepsy, schizophrenia, and autism spectrum disorders.
149

150 **Material and methods**
151

152 **Animal subjects**
153

The University of Lethbridge animal care committee approved all procedures, which adhered to
the guidelines set forth by the Canadian Council on Animal Care and Use. We used 20 young
adult (6–8 weeks old) C57BL/6 mice (12 males and 8 females). Mice were accommodated in
transparent plastic cages within a 12-hour light-dark cycle, with lights turning on at 7:30 AM and

provided unrestricted food and water access. Room temperature was maintained at 24 ± 2 °C,
and relative humidity was kept between 40% and 50%.
160
161   **Viral constructs**
162
163   The viral constructs originated from the Viral Vector Core of the Canadian Neurophotonics
164   Platform (RRID: SCR_016477). Plasmids encoding iGABASnFR2 and csGFP were obtained
165   from Addgene, and AAV2/PHP.N-CAG-iGABASnFR2 and AAV2/PHP.N-CAG-csGFP viral
166   vectors were subsequently designed, packaged, and purified by the CNP Viral Vector Core at a
167   final concentration of 1.5×10^{13} GC/mL.
168
169   **Retro-orbital injection**
170
For retro-orbital viral injections[21], 4–6-week-old C57BL/6 mice were anesthetized with 3%
isoflurane and maintained at 2–2.5% during the procedure. Metacam (5 mg/kg, subcutaneous)
was administered for analgesia, and body temperature was maintained using a heating blanket.
Topical anesthesia (0.5% proparacaine hydrochloride) was applied to the eye, followed by gentle
pressure to induce mild eye protrusion. A 30-gauge needle was then inserted through the medial
canthus at a 30° angle into the retro-orbital sinus. Mice received 1.4×10^{11} genome copies (GC)
of either AAV2/PHP.N-CAG-iGABASnFR2 or AAV2/PHP.N-CAG-csGFP, using the PHP.N
capsid for widespread CNS expression, as originally described by[22] .
179
180   **Drug Administration**
181
182   Tiagabine hydrochloride (Cat #SML0035, Sigma-Aldrich Canada) was purchased from Sigma-
183   Aldrich. Tiagabine was dissolved in sterile saline to achieve a concentration of 5 mg/ml. For
184   intraperitoneal (IP) injection, mice received Tiagabine at a dosage of 10 mg/kg body weight.
185   Control animals were administered an equivalent volume of sterile saline.
186
187   **Surgical procedure and post-operative Care**
188
C57BL/6 mice received buprenorphine (0.05–0.1 mg/kg, subcutaneously) approximately 30
minutes before surgery, followed by anesthesia with isoflurane (1–2% in oxygen) delivered via a
nose cone. After shaving and sterilizing the scalp, lidocaine (0.5%, 5 mg/mL, subcutaneously)
was administered at the incision site for local anesthesia (0.04–0.08 mL for mice weighing 25–55
g). A midline incision was made to expose the skull, and the overlying skin was carefully
removed to avoid damaging the underlying bone. A custom-designed head plate was affixed to
the skull using C&B Metabond Quick Base (Parkell, NY, USA) mixed with C&B Metabond
Clear L-Powder (3 g, Product Code S399, Parkell, Japan). A sterile 12-mm circular glass
coverslip (Carolina Biological Supply, Cat. No. 633005) was placed on the skull surface and

sealed in place with the same adhesive. For hippocampal recordings, a bipolar electrode made of
two twisted 50 μm Teflon-coated stainless-steel wires (A-M Systems) was slowly inserted
through a craniotomy at a 57° angle relative to vertical. Electrode placement was guided by
continuous monitoring of signal quality using both visual and auditory feedback. After, the
electrode was secured to the skull using Krazy Glue, followed by dental cement. An
electromyography (EMG) electrode was also implanted into the neck muscles to monitor muscle
activity. Following surgery, animals were housed individually in temperature-controlled
recovery cages and received subcutaneous injections of Baytril (enrofloxacin, 10 mg/kg),
meloxicam (5 mg/kg), and 1 mL of warm sterile saline. These injections were administered once
every 24 hours for three days postoperatively, in accordance with institutional guidelines. After
this recovery period, animals were monitored twice daily for the remainder of the experiment.
209
210
211 **iGABASnFR2 imaging under anesthesia**
212
Following, the animals were anesthetized with isoflurane (2.5% for induction, followed by 1%
for maintenance). The depth of anesthesia was confirmed by assessing reflexes. Once adequately
anesthetized, each mouse was positioned in a head-stage, and the head was securely head-fixed.
A homoeothermic blanket was utilized to maintain their body temperature, and isoflurane was
administered via a nosepiece. The isoflurane concentration was adjusted to 1% to initiate the
imaging procedure. Images were acquired using a microscope consisting of a front- to-front pair
of video lenses with a field of view measuring 8.6×8.6 mm. The camera's focal plane was
positioned 0.5–1 mm (about 0.04) below the cortical surface. A 12-bit charge-coupled device
(CCD) camera (1M60 Pantera Dalsa, Waterloo, ON) and an EPIX E8 frame grabber with XCAP
3.8 imaging software (EPIX, Inc., Buffalo Grove, IL) were used to capture images at a frame rate
of 80 Hz. These imaging parameters have been employed in previous studies[23-25]. Carefully
designed data collection protocols support the robustness of our findings. Sequential illumination
was achieved using alternating blue and green LEDs[26]. The timing of LED alternation is
illustrated in Supplementary Figure 1. Blue light (473 nm, filtered through a 467–499 nm
bandpass) was used to excite the iGABASnFR2 indicator, and green light (530 nm, filtered
through a 527/42 nm bandpass) was used for intrinsic signal imaging of blood volume. A
bandpass emission filter (shown in Figure 1) was positioned in front of the CCD camera to
enable selective detection of either fluorescence or reflectance signals. Blue and green LEDs
were synchronized and alternated on a frame-by-frame basis using TTL triggering, resulting in
interleaved acquisition of fluorescence and reflectance images at 40 Hz per channel.
Additionally, images of reflectance, crucial for blood artifact corrections, were evaluated within
the current pipeline[27-29]. Anesthetized iGABASnFR2 imaging of spontaneous activity was
conducted without sensory stimulation for 15- minute sessions.
236
Sensory stimulation

We captured the iGABASnFR2 signal in reaction to varied peripheral simulations while utilizing
urethane anesthesia, following the methodology outlined in previous studies. Sensory stimuli
were employed to map the functional areas of the hind-limb somatosensory, forelimb
somatosensory, auditory, visual, and barrel cortices[23,30]. Sensory stimuli were applied to map
cortical regions corresponding to forelimb, hindlimb, whisker, visual, and auditory modalities.
For forelimb and hindlimb stimulation, a piezoelectric bending actuator delivered a single 300-
ms tap via a square pulse directly to the skin of one forelimb or hindlimb. Whisker stimulation
targeted the C2 whisker, which was attached to a piezoelectric actuator (Q220-A4-203YB, Piezo
Systems, Inc., Woburn, MA) and deflected using a single 300-ms square pulse . Visual stimuli
consisted of a single 20-ms pulse of 435 nm light (LED), delivered at a fixed distance and height
relative to the right eye. For each sensory modality, 40 stimulus presentations were delivered
with a 10-second interstimulus interval to calculate average cortical responses. The timing of
stimulus delivery is illustrated in Supplementary Figure 1.

Habituation

After the 7-day recovery period from surgery, mice gradually habituated to head restraint in the
recording environment. Initially, each mouse was placed individually on the recording platform
along with Cheerios cereal, allowing them to explore freely and become comfortable. Mice were
progressively acclimated to eating the cereal while head-fixed, beginning with 5-minute sessions
and increasing by 5 minutes each day until reaching 60 minutes.

iGABASnFR2 imaging during wakefulness

Following habituation, wakefulness recordings began. Each animal underwent recording sessions
every three days, completing three to four sessions per mouse. Recordings were consistently
performed at the same time of day to reduce variability and stress. Between sessions, mice were
returned to their home cages for rest and recovery before the next recording. After finishing all
wakefulness sessions, mice proceeded to the sleep recording phase.

iGABASnFR2 imaging during sleep

To optimize conditions for natural sleep under head restraint, mice were transferred from their
colony housing to a separate room at noon the day before recording. Sleep was restricted for six
hours by gentle stimulation (using a cotton-tip stick) whenever signs of drowsiness were
observed. Following 6 hours of sleep deprivation, mice were placed overnight in large, enriched
cages containing a running wheel, Cheerios, and a water container to promote exploration and
natural sleep. The next morning (~9:00 AM), animals were transferred to the imaging platform
for sleep recordings. Afterward, they were returned to their home cages for at least three days of
recovery before any further recordings. This sleep deprivation protocol is commonly used to

induce moderate but physiologically meaningful sleep pressure[31], which is known to trigger a
homeostatic increase in slow-wave activity (SWA) during subsequent NREM sleep.

Preprocessing
280

Image stacks were first de-interleaved to separate the GABA-sensitive fluorescence signal (blue
channel) and the hemodynamic reflectance (green channel) signal. The correct channel
assignment was verified by computing pixel-wise correlations between the first frame of each
stack and reference images corresponding to each illumination wavelength. To quantify
extracellular GABA dynamics, the relative fluorescence change ($\Delta F/F$) was calculated on a per-
pixel basis using a 2-second pre-stimulus window to define baseline fluorescence (F_0). The
$\Delta F/F$ signal was defined as:
288

289 $$\Delta F/F(t) = (F(t) - F_0) / F_0$$
290

291 where $F(t)$ is the fluorescence intensity at time t, and F_0 is the baseline fluorescence. To correct
292 for hemodynamic artifacts[27-29], a pixel-wise linear regression was applied, in which a scaled
293 version of the reflectance signal $R(t)$ was subtracted from the fluorescence trace:
294

295 $$F_{corrected}(t) = F(t) - \alpha \cdot R(t)$$
296

The corrected fluorescence signal was then normalized to the baseline ($\Delta F/F$) and bandpass
filtered between 31–74 Hz using a second-order Butterworth filter to remove low-frequency drift
and high-frequency noise. Trial-averaged $\Delta F/F$ responses were generated to assess sensory-
evoked activity. All preprocessed data—including corrected $\Delta F/F$, raw fluorescence, and
reflectance signals—were saved in float32 format.

ROI-based fluorescence analysis
Following preprocessing, imaging data from specific regions of interest (ROIs)—defined as 3×3-
pixel areas (~40,401 μm^2) centered around anatomical coordinates corresponding to stimulation
sites)— were extracted. Baseline correction was conducted by subtracting the mean fluorescence
signals calculated from a 1-second pre-stimulus period from the post-stimulus fluorescence
signals. The signals were further filtered to eliminate slow baseline drifts using a high-pass filter
(>0.1 Hz) and to reduce high-frequency noise using a low-pass filter (<5 Hz). From the filtered
signals, several key parameters were derived, including peak amplitude (maximum $\Delta F/F_0$
within a 1-second post-stimulus interval), time-to-peak, and decay time (duration for the
fluorescence to fall to 50% of peak amplitude). For visualization, the corrected $\Delta F/F_0$ signals
were plotted with indicators marking peak and decay times, facilitating clear interpretation of
response dynamics. Mean responses across trials were computed separately for each sensory
modality, with variability assessed by plotting the standard error of the mean (SEM) as shaded
regions surrounding the mean trace.

Motion detection and exclusion

Motion artifacts were identified and excluded from analyses using electromyography (EMG) signals recorded simultaneously with imaging data. EMG recordings were smoothed using a median filter and squared to enhance the detection of muscle activity periods. An activity threshold was established at the 95th percentile of the processed EMG signal to identify movement onset and offset events. These EMG-detected motion periods were temporally aligned with imaging frames via synchronized camera clock signals. Frames coinciding with detected movements were subsequently removed from analysis, ensuring the seed pixel correlation analysis reflected only stationary periods free of motion-related artifacts.

Seed-pixel correlation analysis

Seed pixel correlation analysis was performed to evaluate functional connectivity based on spontaneous GABA activity from sleep and anesthetized mice. preprocessed data were spatially registered to the Allen Brain Atlas, enabling anatomical alignment and inter-subject comparisons. Regions of interest (ROIs)—specifically the barrel cortex (BC), visual cortex (VC), hindlimb (HL), and forelimb (FL)—were defined using anatomical coordinates derived from the atlas. Within each ROI, seed pixels were selected to serve as reference points for correlation-based connectivity analysis.

Motion signal extraction, alignment, and sleep scoring

Behavioral videos were used to monitor animal movement during imaging. Motion signals were extracted using FaceMap[32], which computes frame-to-frame pixel intensity changes in user-defined regions of interest (ROIs). Five ROIs—nose, whisker pad, ear, shoulder, and trunk—were selected to capture both facial and body movements. To synchronize video with neural data, camera frame pulses were recorded on analog channels during acquisition. These were used to align motion traces with electrophysiological recordings (LFP and EMG, sampled at 2 kHz). Traces were interpolated or trimmed as needed, z-scored, and averaged across ROIs to generate composite facial and body motion signals. This enabled detection of both gross movement and small twitches, such as those occurring during REM sleep, which may not be captured by EMG alone. With motion aligned, vigilance states were classified as wakefulness, NREM, or REM sleep using combined behavioral and physiological features. Wakefulness was marked by visible movement and high EMG activity. NREM sleep was defined by low EMG, a low hippocampal theta-to-delta power ratio, and the presence of Large Irregular Activity (LIA) in the LFP. REM sleep was characterized by minimal EMG activity, a high theta-to-delta ratio, and continuous hippocampal theta. In head-fixed recordings, pupil constriction served as an additional marker of

355   sleep onset[33-35]. This multimodal approach enabled robust, accurate classification of sleep stages
356   across different experimental conditions.
357
358   **Statistical analysis**
359
360   All data processing and analyses were performed using custom scripts written in MATLAB
361   R2024a.
362
363   **Results**
364
365   **Experimental workflow, imaging setup, and cortical expression of iGABASnFR2**
366
To characterize extracellular GABA dynamics, we first injected AAV2/PHP.N-CAG-
iGABASnFR2 and AAV2/PHP.N-CAG-csGFP systemically via retro-orbital injection (**Figure
1A**). Imaging was performed under isoflurane anesthesia, during quiet wakefulness, NREM and
REM sleep. The imaging setup (**Figure 1B**) utilized a bilateral cranial window and a CCD-based
wide-field microscope that captured fluorescence across an 8.6 × 8.6 mm field of view.
Sequential blue (470 nm) and green (530 nm) LED illumination enabled alternating frame
acquisition and hemodynamic correction[27-29]. To validate sensor expression, we conducted a
histological analysis (**Figure 1C**). A robust expression of iGABASnFR2 was revealed in cortical
and hippocampal regions. Coronal brain sections (**Figure 1Cii**) demonstrated consistent
expression across animals. To assess inter-animal variability and regional expression strength,
fluorescence intensity profiles were extracted using ImageJ from defined cortical areas (**Figure
1Ciii**), and to further quantify and visualize overall trends, these profiles were averaged across
animals, with the mean ± SEM shown (**Figure 1Civ**) confirming uniform sensor expression
suitable for quantitative cortical imaging.
381
382
383   **iGABASnFR2 reveals modality- and hemisphere-specific cortical inhibition under**
384   **anesthesia**
385
Previous studies on sensory processing have primarily focused on excitatory neuronal responses,
often using calcium or glutamate indicators to map stimulus-evoked activity across the cortex
[19,23,36]. However, much less is known about how sensory stimuli engage inhibitory networks,
particularly at the mesoscale. Inhibitory interneurons play a crucial role in shaping sensory
responses, modulating cortical excitability, and controlling the timing and precision of neural
coding[37-38]. To examine the spatiotemporal dynamics of GABA activity across the cortex, we
delivered contralateral whisker, hindlimb, forelimb, and visual stimulation under 1% isoflurane
anesthesia and recorded GABA activity using the iGABASnFR2 sensor. As shown in Figure 2A,
each sensory modality shows a localized increase in GABAergic fluorescence within the

corresponding primary sensory cortex, with response onsets occurring approximately 75–150 ms
after stimulus. All sensory modalities elicited more robust and spatially localized GABAergic
activity. Temporal response profiles (**Figure 2B**) demonstrated stronger GABAergic activation
in the contralateral hemisphere relative to the ipsilateral side across all sensory modalities.
Specifically, the contralateral visual cortex exhibited the highest peak amplitude, followed by
hindlimb, whisker, and forelimb cortices while ipsilateral responses were generally weaker and
slower. To quantify these differences, we extracted response features including decay time, time
to peak, and peak amplitude across all sensory regions (**Figure 2C**). A two-way ANOVA
revealed significant main effects of sensory region on all three measures of inhibitory response
dynamics. Time to peak ($p = 0.0092$), decay time ($p = 0.0006$), and peak amplitude ($p = 0.0011$)
all varied significantly across sensory regions, indicating region-specific characteristics of
GABAergic inhibition. Laterality (ipsilateral vs. contralateral) had a significant main effect only
on peak amplitude ($p < 0.0001$), with contralateral responses consistently showing higher
amplitudes. No significant effects of laterality were observed for time to peak ($p = 0.35$) or decay
time ($p = 0.77$). Furthermore, there were no significant interactions between region and laterality
for any of the three metrics (time to peak: $p = 0.22$; decay: $p = 0.83$; peak amplitude: $p = 0.78$),
suggesting that hemispheric differences in GABAergic inhibition were consistent across
modalities and not dependent on specific sensory regions. To confirm that these delayed signals
were specific to GABAergic dynamics and not artifacts of hemodynamics or sensor excitation,
we used csGFP-expressing mice under identical imaging conditions as a negative control. As
shown in Supplementary **Figure 2**, csGFP mice showed no significant sensory-evoked or
spontaneous fluorescence changes, confirming that the iGABASnFR2 signals reflect GABAergic
activity. Auditory stimulation under anesthesia evoked clear iGABASnFR2 responses in the
auditory cortex (Supplementary Figure 3), further validating sensor specificity. To assess cortical
coordination during sensory processing, we performed seed-pixel correlation analysis across 10
anatomically defined cortical regions. This revealed structured and modality-specific inhibitory
networks, with the strongest intrahemispheric connectivity observed contralateral to the stimulus
(Supplementary Figure 4). Across sensory modalities—including whisker (**Video 1**), hindlimb
(**Video 2**), forelimb (**Video 3**), and visual (**Video 4**) stimulation—sensory input activates
thalamocortical projections targeting layer 4 of the primary sensory cortices, leading to early
excitatory responses[39-40]. These are primarily mediated by excitatory neurons and refined by fast
feedforward inhibition from parvalbumin (PV)- expressing interneurons[41-42]. Subsequently, a
delayed GABAergic response emerges, largely driven by somatostatin (SST)-expressing
interneurons providing feedback inhibition to modulate dendritic activity and maintain cortical
stability[9,43,44]. This response is consistently stronger and earlier in the contralateral hemisphere,
which receives direct thalamic input, whereas the ipsilateral hemisphere exhibits a weaker and
more delayed GABAergic response, likely due to slower callosal transmission and reduced
excitatory drive[45-46]. Together, these results demonstrate that sensory-evoked GABA dynamics
follow a consistent temporal structure across modalities and hemispheres and confirm that

iGABASnFR2 reliably detects extracellular GABA responses with spatiotemporal precision
across cortical regions, establishing its utility for mesoscale mapping of inhibitory dynamics.
436
437 **Sensory-evoked and spontaneous GABA activity in quiet wakefulness resembles**
438 **anesthesia-induced patterns**
439
Cortical brain states vary across behavioral conditions, shaping spontaneous activity and sensory
processing. During active behavior, such as locomotion or whisking, cortical activity becomes
desynchronized, and inhibition is modulated to refine sensory gain[47-49]. In contrast, during quiet
wakefulness and under light anesthesia, neuronal activity is dominated by slow, synchronized
fluctuations that reflect reduced arousal and a shift toward global inhibitory tone[43,50]. While
previous studies have shown that GABAergic interneurons contribute significantly to these state-
dependent dynamics[36,51], it remains unclear whether the spatiotemporal profile of extracellular
GABA during quiet wakefulness resembles that observed under anesthesia. To examine whether
GABAergic responses to sensory stimulation and spontaneous activity in quiet awake mice
exhibit spatiotemporal dynamics similar to those observed under anesthesia, we performed wide-
field imaging of the cortex using the iGABASnFR2 sensor. Head-fixed mice were imaged in
both quite awake and anesthetized states, the latter induced by 1% isoflurane. EMG recordings
from a neck muscle electrode, along with video monitoring of body and whisker movement,
were used to classify behavioral state (**Figure 3A**). Power spectral analysis of EMG signals
confirmed a reduction in muscle tone under anesthesia compared to the quiet awake state. We
recorded cortical GABA signals evoked by contralateral visual or whisker stimulation, averaging
responses across 40 trials for each condition (**Figure 3B**). Both anesthetized and quiet awake
states showed robust stimulus-evoked increases in extracellular GABA in primary sensory areas,
including the primary visual cortex (VISp), anterior visual area (VISa), barrel cortex (BC), and
primary motor cortex (M1). In Figure 3Bi, which shows cortical GABA responses to visual
stimulation, the quiet awake state is characterized by an earlier onset and more widespread
GABA release across the visual cortex. In contrast, under anesthesia, the GABA signal appears
later and is more spatially restricted. Similarly, in Figure 3Bii, following whisker stimulation, the
quiet awake state shows strong and broad activation of the contralateral barrel cortex (BC) and
associated motor areas (M1). This response is both faster and more spatially extensive than in the
anesthetized state. To assess how GABAergic responses vary across brain states, we analyzed
their temporal dynamics during visual and whisker stimulation (**Figure 3C**). Responses were
stronger and more spatially distinct during quiet wakefulness than under anesthesia—VISp >
VISa for visual input and BC > M1 for whisker input. Quantitative analysis (Figure 3D)
confirmed significantly higher peak amplitudes and longer decay times in the awake state.Time
to peak was generally shorter in the awake state, suggesting faster inhibitory onset, though most
differences were not statistically significant. However, BC in wakefulness responded
significantly faster than M1 under anesthesia. Decay times were longer under anesthesia,
especially in BC, suggesting more prolonged inhibition when cortical activity is suppressed.

These results highlight the influence of both brain state and region on the strength and timing of
GABAergic responses.
476

Mesoscale imaging of cortical GABA dynamics during natural sleep and wakefulness
478

While many studies have explored cortical dynamics across sleep and wake states using
electrophysiological methods and excitatory activity sensors[52-54], the ability to track inhibitory
signaling at mesoscale resolution across natural brain states remains limited. To further assess
iGABASnFR2 performance across different brain states, we examined cortical GABA dynamics
during quiet wakefulness, non-rapid eye movement (NREM) sleep, and rapid eye movement
(REM) sleep. Mesoscale iGABASnFR2 imaging was combined with simultaneous hippocampal
LFP and EMG recordings in head-fixed mice. Animals were allowed to transition naturally
between vigilance states while cortical GABA levels were monitored (Figure 4A–C). GABA
signals were highest during wakefulness and reduced during NREM sleep (Figure 4D). Spectral
analysis showed a reduction in low-frequency GABA fluctuations during REM compared to both
wakefulness and NREM (Figure 4E), suggesting diminished slow GABA oscillations during
REM sleep. To further assess spatial coordination of cortical GABA activity, we computed
pairwise correlation maps across the cortex during each brain state over multiple days. As shown
in (**Figure 4F**), Cortical GABA signals exhibited strong bilateral synchrony during quiet
wakefulness, which was markedly reduced during NREM sleep and only moderately diminished
during REM. This pattern is evident in interhemispheric correlation heatmaps (Figure 4F), where
NREM shows the most substantial decrease in bilateral synchrony, while REM correlations
remain relatively higher. Quantitative analysis confirmed that mean interhemispheric correlation
values were lower during NREM sleep compared to both wakefulness and REM (Figure 4G).
During transitions from REM to wakefulness, we observed a sharp increase in cortical GABA
levels (Figure 4H). This transition was accompanied by a broad re-engagement of cortical
GABAergic activity across multiple regions (**Figure 4I**), suggesting rapid reinstatement of
GABA response upon arousal from REM sleep. A detailed overview of the motion-based and
electrophysiological features used for behavioral state classification is provided in
Supplementary Figure 5, which illustrates the temporal alignment of whisker pad and nose
motion, EMG power, body motion, theta-to-delta ratio, and hippocampal LFP signals across
NREM and REM sleep transitions. Together, these findings demonstrate that iGABASnFR2
reliably captures spontaneous, brain state-dependent fluctuations in extracellular GABA with
high temporal and spatial resolution. It supports its utility for long-term mesoscale imaging of
cortical inhibition under natural physiological conditions.
509

510 **Intracortical long-range GABAergic correlations revealed by seed-pixel analysis across**
511 **brain states**
512

To assess the organization of spontaneous extracellular GABA dynamics in the cortex, we used seed-pixel correlation mapping of iGABASnFR2 fluorescence to investigate intracortical long-range connectivity across different brain states. By placing seed pixels in primary sensory regions, we generated correlation maps of spontaneous GABA dynamics across awake, NREM, and REM sleep states (Figure 5A). In the awake state, these maps revealed strong bilateral synchrony and widespread long-range connectivity between distant cortical regions. These spatial patterns of functional connectivity closely resemble those previously observed with excitatory signals using iGluSnFR and Ca²□ imaging, suggesting that spontaneous GABA activity also reflects underlying anatomical connectivity and shared network drive[24,25,30]. During NREM sleep, seed-pixel correlations were reduced, indicating a decoupling of large-scale inhibitory networks consistent with cortical slow-wave activity. In contrast, REM sleep preserved many of the bilateral and local connections seen in wakefulness, although with moderate reductions in correlation strength. We further quantified interhemispheric connectivity using pairwise correlation matrices of bilateral cortical regions (**Figure 5B**). These patterns, consistently captured using iGABASnFR2, highlight the sensor's sensitivity to state-dependent fluctuations in extracellular GABA and its utility for mapping mesoscale inhibitory networks in vivo.

Tiagabine elevates baseline GABA levels but dampens sensory-evoked responses and reorganizes cortical inhibitory connectivity

To examine how pharmacological inhibition of GABA reuptake influences cortical GABA dynamics, we administered Tiagabine—a selective GAT-1 inhibitor[55]—under 1% isoflurane anesthesia and monitored extracellular GABA levels using iGABASnFR2. Mice were head-fixed throughout the experiment. We first recorded visually evoked GABA responses (~6 minutes), followed by a ~15-minute baseline period of spontaneous activity. While the mouse remained head-fixed, Tiagabine was injected intraperitoneally. Spontaneous activity was then recorded for another ~15 minutes before delivering a second round of visual stimulation to assess post-Tiagabine responses. Before Tiagabine injection, visual stimulation evoked robust and spatially localized increases in GABA signals within the contralateral primary visual cortex (Figure 6Ai). Following Tiagabine administration, the same sensory stimulus failed to evoke any detectable response (Figure 6Aii). This complete loss of evoked GABA activity is also evident in the time-series traces (Figure 6Aiii). These recordings were acquired at 150 Hz using the blue fluorescence channel under continuous (non-strobing) illumination.

To better understand this loss of evoked responsiveness, we analyzed spontaneous GABA activity using dual-wavelength strobing imaging (40 Hz blue, 40 Hz green). In the primary visual cortex , the raw blue fluorescence signal, averaged across 6 animals, showed a sustained elevation[56-57]in baseline GABA levels following Tiagabine injection (Figure 6Bi), consistent with extracellular GABA accumulation due to GAT-1 blockade. Reflectance signal from the same region also increased post-injection (Figure 6Bii), confirming overall signal elevation.

Quantitative analysis revealed that Tiagabine significantly increased the duration, frequency, and
peak amplitude of spontaneous GABA transients across animals (Figure 6Biii), reflecting
heightened inhibitory tone.
To assess how elevated GABA levels affect network organization, we performed seed-pixel
correlation analysis (0.1–5 Hz) of spontaneous iGABASnFR2 activity. Following Tiagabine
injection, seed regions in sensory cortices showed increased local synchrony and expanded
spatial correlation patterns (Figure 6C). Interhemispheric correlation matrices also revealed
stronger bilateral connectivity, particularly between homologous sensory areas (Figure 6D).
These results further validate the ability of iGABASnFR2 to detect changes in extracellular
GABA levels under both baseline and stimulated conditions, highlighting its sensitivity to
dynamic alterations in cortical inhibition and its potential utility in pharmacological studies
targeting GABAergic signaling.
565
566 **Discussion**
567
We used mesoscale imaging with the genetically encoded GABA sensor iGABASnFR2 to map
the spatiotemporal dynamics of extracellular GABA across the intact mouse cortex in vivo. By
characterizing extracellular GABA dynamics in both sensory-evoked and spontaneous across
brain states—including anesthesia, quiet wakefulness, NREM, and REM sleep—we provide, for
the first time, a comprehensive mesoscale imaging of GABA dynamics in the cortex. our results
establish iGABASnFR2 as a valuable tool for investigating GABAergic tone and state-dependent
fluctuations in extracellular GABA, complementing previous studies that used iGluSnFR and
voltage-sensitive dyes to map excitatory signaling and depolarization across large- scale cortical
networks[19,23]. The robust and state-dependent changes in extracellular GABA we observed—
during sensory stimulation, spontaneous activity, and pharmacological manipulation—
demonstrate that inhibitory tone is dynamically regulated across brain states and cortical regions.
579
580 **Interpreting iGABASnFR2 signals**
581
iGABASnFR2 fluorescence changes are likely shaped by a combination of GABA release from
interneurons, diffusion through the extracellular space, and clearance by GABA transporters,
particularly GAT-1[58-60]. The timing of sensory-evoked responses—characterized by a delayed
onset (~75 ms) and prolonged signal lasting several hundred milliseconds—is consistent with the
dynamics of SST-positive interneuron-mediated inhibition, which is known to follow the rapid
activation of PV+ interneurons[41,43]. The prolonged decay phase of iGABASnFR2 signals
(**Figure 2C**) suggests that extracellular GABA may persist longer, potentially contributing to
modulatory or volume transmission effects[61].
590
591 **Cortical GABA responses to sensory input are conserved across brain states**
592

Sensory-evoked results revealed robust, modality-specific patterns of extracellular GABA
activation across the cortex, reflecting both localized responses and broader, network-level
dynamics (**Figure 2A**). Each stimulus modality selectively activated its corresponding primary
sensory area (**Figure 2B**), with activity spreading from the focal point of stimulation to more
distal, functionally connected cortical regions. This widespread response pattern suggests that
inhibitory activity can propagate across large-scale networks, challenging the traditional view of
inhibition as strictly local.

In our study, contralateral responses were consistently stronger and faster than ipsilateral
responses, likely reflecting direct thalamocortical projections to the primary sensory cortex. In
contrast, the delayed and weaker ipsilateral responses are consistent with slower callosal
transmission and interhemispheric integration (Figure 2B–C), as reported in previous studies of
cortical sensory processing[45]. In addition, despite differences in global cortical state and arousal,
sensory-evoked GABAergic responses were consistently observed across brain states, suggesting
a preserved functional role of inhibition, even as response strength and timing are modulated
(Figure 3B–D). This robustness underscores the reliability of iGABASnFR2 for monitoring
inhibitory dynamics and reflects the essential role of inhibition in stabilizing excitation and
regulating cortical gain, as also supported by prior calcium imaging studies in awake animals [47].

Spontaneous GABA dynamics reveal state-dependent connectivity

In our study, interhemispheric GABA synchrony was lowest during NREM sleep (Figure 4D-G),
suggesting that inhibition in this state becomes locally structured and functionally decoupled
across hemispheres. REM sleep (Figure 4D-G) showed partial recovery of interhemispheric
GABA synchrony, suggesting a reorganization of inhibitory networks distinct from both
wakefulness and NREM. This contrasts with vascular-based findings; for instance, a study
[62]reported strong bilateral hemodynamic coherence during both NREM and REM, reflecting
broader, nonspecific signals. However, they also noted that neurovascular signals integrate
contributions from neurons, astrocytes, and metabolism, potentially decoupling them from
electrophysiological activity. Our direct measurement of extracellular GABA provides a distinct
view, revealing that strong vascular synchrony during NREM does not imply coordinated
inhibitory signaling. Indeed, the low GABA synchrony we observe during NREM contrasts with
the high bilateral [HbT] coherence reported by Turner and colleagues [62], highlighting a
dissociation between vascular and inhibitory network dynamics.

Similar patterns have been observed in BOLD fMRI studies, which report reduced large-scale
connectivity during NREM and partial restoration during REM[63-65]. These converging lines of
evidence highlight arousal state as a key factor in shaping both vascular and neural network
dynamics and emphasize the central, yet state-specific, role of inhibition in large-scale brain
coordination during sleep.

Functional inhibitory architecture aligns with cortical structural organization

Seed-pixel correlation results revealed structured long-range inhibitory connectivity (**Figure**
5A). In the awake state, we observed strong bilateral synchrony between homologous sensory
areas and strong local correlations within modalities (**Figure 5Ai, 5Bi**). These maps are similar
to those derived from excitatory indicators, including iGluSnFR and VSD[19,23,66], supporting the
view that inhibitory and excitatory networks are functionally integrated, but exhibit distinct
temporal dynamics[42]. In NREM sleep, long-range inhibitory connectivity weakened, suggesting
a shift toward localized processing (**Figures 5Aii, 5Bii**) while REM sleep partially restored
(**Figures 5Aiii, 5Biii**). State-dependent changes highlight the flexibility of cortical inhibitory
networks, which adapt dynamically to behavioral states, promoting broad coordination during
wakefulness and preserving localized stability during sleep[67].

Tiagabine elevates extracellular GABA and disrupts sensory-evoked responses via altered
cortical synchrony

Our data show that Tiagabine disrupts sensory-evoked GABAergic responses by elevating
extracellular GABA and altering cortical network dynamics. Before injection, visual stimulation
produced robust, spatially confined increases in GABA sensor signals in the visual cortex
(Figure 6A). After Tiagabine administration, however, the same stimulus no longer evoked
detectable GABA responses—despite a sustained increase in baseline fluorescence (Figure 6Bi),
confirming that the sensor remained active. Although not used for quantification, an increase in
green reflectance signal (Figure 6Bii) suggested broader changes in cortical vascular or
metabolic state. The absence of evoked activity in the presence of elevated GABA points to a
nonlinear relationship between inhibitory tone and sensory responsiveness. This may involve
excessive tonic inhibition, desynchronized interneuron networks, or saturation of GABAergic
circuits. In our study, Tiagabine-induced increases in extracellular GABA were clearly detected
by iGABASnFR2, with fluorescence peaking around 10 minutes post-injection—consistent with
prior microdialysis reports[68]. This elevation likely enhanced tonic inhibition via extrasynaptic
GABA_A receptors[69], reducing neuronal excitability and suppressing phasic, stimulus-driven
responses. This mechanism helps explain the loss of stimulus-locked GABA transients. Similar
effects have been observed in rodent EEG studies, where Tiagabine increases low-frequency
synchronization and, at higher doses, induces hypersynchronous activity resembling absence
seizures—likely a result of widespread tonic inhibition[70]. Our results align with these findings
and demonstrate that iGABASnFR2 effectively detects sustained changes in extracellular
GABA. However, the loss of evoked transients following Tiagabine suggests the sensor may
reach a functional ceiling under conditions of high ambient GABA, limiting its dynamic range.
Though Tiagabine does not directly cause transporter reversal, elevated intracellular GABA and
transporter saturation could still favor nonvesicular GABA release[67], further amplifying tonic
signaling. Together, these findings highlight both the strengths and limitations of iGABASnFR2:
it is sensitive to pharmacologically induced increases in extracellular GABA but may
underestimate fast, transient responses when tonic levels are elevated. This underscores the

importance of considering inhibitory tone and network state when interpreting GABA imaging
data.
673
674 **Conclusion**
675

Our findings establish iGABASnFR2 as a robust sensor for widefield imaging of extracellular
GABA in the cortex, revealing both strong stimulus-locked responses and state-dependent
dynamics. Our experiments show that inhibition is not merely local or reactive but is organized
into large-scale motifs that shift with arousal state and can be pharmacologically tuned. For a
long time, efforts to understand how excitation and inhibition interact across the cortex have
been limited by the lack of tools that can monitor both processes in real time—especially in
awake, behaving animals. That's now changing. Genetically encoded sensors like
iGABASnFR2, along with glutamate and calcium indicators for excitatory activity, offer a new
window into the excitation-inhibition (E/I) balance with spatial, temporal, and cell-type
specificity. Importantly, the development of red-shifted glutamate sensors, such as R-iGluSnFR
[71], now makes it possible to image GABA and glutamate simultaneously, opening the door to
more complete views of circuit function in vivo.This capability will be essential for
understanding how the E/I balance is regulated during normal brain function—and how it
becomes disrupted in neurological and psychiatric disorders. Many neurological and psychiatric
disorders—such as epilepsy, autism, and schizophrenia—involve disruptions in E/I balance, yet
direct measurements of these dynamics in vivo have remained elusive. Simultaneous imaging of
glutamate and GABA now enables researchers to observe how excitation and inhibition interact
across space and time, how this interaction is modulated by behavioral state, neuromodulators, or
genetic risk factors, and how it breaks down in pathological conditions. These insights are
critical for developing targeted interventions that restore E/I balance and stabilize network
function in disease.

Acknowledgments

We thank Di Shao and the Animal Welfare Committee at the University of Lethbridge for their
support in enabling the animal experiments. We also thank Majid H. Mohajerani for providing
access to laboratory resources and equipment during the data collection phase of this project. We
thank Nicholas J Michelson and Tim H. Murphy for performing the initial experiment.
702
703 **Funding**
704
705 The author(s) declare that financial support was received for the research, authorship, and/or
706 publication of this article. This work was supported by Alberta Innovate and Natural Sciences
707 and Engineering Research Council of Canada (grant 390930), awarded to Majid H. Mohajerani.
708

Author contributions

E.R. conceptualized the study. E.R. designed and conducted the main experiments. M.N. contributed to additional experimental procedures. J.K.A. implemented the strobing imaging system. E.R. and S.T. analyzed the data. E.R. prepared all the figures, wrote, and revised the manuscript. All authors have read and agreed to the submitted version of the manuscript.

Supplementary material

Supplementary material and methods are available.

Disclosures

The authors declare that there are no financial interests, commercial affiliations, or other potential conflicts of interest that could have influenced the objectivity of this research or the writing of this paper.

The code and data Availability

The code and all data analyzed and used to produce the main findings of this study have been deposited on Dryad. Source data files are available for Figures 2, 3, 5, and 6.

References

[1]Zhou, Y., & Danbolt, N. C. (2014). Glutamate as a neurotransmitter in the healthy brain. *Journal of Neural Transmission*, *121*(8), 799–817. https://doi.org/10.1007/s00702-014-1180-8

[2]Froemke, R. C. (2015). Plasticity of cortical excitatory-inhibitory balance. *Annual Review of Neuroscience*, *38*, 195–219. https://doi.org/10.1146/annurev-neuro-071714-034002

[3]Tatti, R., Haley, M. S., Swanson, O. K., Tselha, T., & Maffei, A. (2017). Neurophysiology and regulation of the balance between excitation and inhibition in neocortical circuits. *Biological Psychiatry, 81*(10), 821–831. https://doi.org/10.1016/j.biopsych.2016.09.017

[4]Yuste, R. (2005). Origin and classification of neocortical interneurons. *Neuron, 48*(4), 524–527. https://doi.org/10.1016/j.neuron.2005.11.012

[5]Markram, H., Toledo-Rodriguez, M., Wang, Y., Gupta, A., Silberberg, G., & Wu, C. (2004).

Interneurons of the neocortical inhibitory system. *Nature Reviews Neuroscience, 5*(10), 793–807.

https://doi.org/10.1038/nrn1519

[6]Tremblay, R., Lee, S., & Rudy, B. (2016). GABAergic interneurons in the neocortex: From

cellular properties to circuits. *Neuron, 91*(2), 260–292.

https://doi.org/10.1016/j.neuron.2016.06.033

[7]Kepecs, A., & Fishell, G. (2014). Interneuron cell types are fit to function. *Nature, 505*(7483),

318–326. https://doi.org/10.1038/nature12983

[8]Ferguson, B. R., & Gao, W.-J. (2018). PV interneurons: Critical regulators of E/I balance for

prefrontal cortex-dependent behavior and psychiatric disorders. *Frontiers in Neural Circuits, 12*,

Article 37. https://doi.org/10.3389/fncir.2018.00037

[9]Urban-Ciecko, J., & Barth, A. L. (2016). Somatostatin-expressing neurons in cortical networks.

Nature Reviews Neuroscience, 17(7), 401–409. https://doi.org/10.1038/nrn.2016.53

[10]De Lanerolle, N. C., Kim, J. H., Robbins, R. J., & Spencer, D. D. (1989). Hippocampal

interneuron loss and plasticity in human temporal lobe epilepsy. *Brain Research, 495*(2), 387–

395. https://doi.org/10.1016/0006-8993(89)90234-5

[11]Lewis, D. A., & Moghaddam, B. (2006). Cognitive dysfunction in schizophrenia: Convergence

of γ-aminobutyric acid and glutamate alterations. *Archives of Neurology, 63*(10), 1372–1376.

https://doi.org/10.1001/archneur.63.10.1372

[12]Curley, A. A., & Lewis, D. A. (2012). Cortical basket cell dysfunction in schizophrenia. *The*

Journal of Physiology, 590(Pt 4), 715–724. https://doi.org/10.1113/jphysiol.2011.224659

[13]Marín, O. (2012). Interneuron dysfunction in psychiatric disorders. *Nature Reviews*

Neuroscience, 13(2), 107–120. https://doi.org/10.1038/nrn3155

[14]Rubenstein, J. L. R., & Merzenich, M. M. (2003). Model of autism: Increased ratio of

excitation/inhibition in key neural systems. *Genes, Brain and Behavior, 2*(5), 255–267.

https://doi.org/10.1034/j.1601-183x.2003.00037.x

[15]Ben-Ari, Y. (2006). Seizures beget seizures: The quest for GABA as a key player. *Critical*

Reviews in Neurobiology, 18(1–2), 135–144. https://doi.org/10.1615/critrevneurobiol.v18.i1-

2.140

[16]Gonzalez-Burgos, G., & Lewis, D. A. (2008). GABA neurons and the mechanisms of network

oscillations: Implications for understanding cortical dysfunction in schizophrenia. *Schizophrenia*

Bulletin, 34(5), 944–961. https://doi.org/10.1093/schbul/sbn070

[17]Marvin, J. S., Borghuis, B. G., Tian, L., Cichon, J., Harnett, M. T., Akerboom, J., Gordus, A.,

Renninger, S. L., Chen, T.-W., Bargmann, C. I., Orger, M. B., Schreiter, E. R., Demb, J. B., Gan,

W.-B., Hires, S. A., & Looger, L. L. (2013). An optimized fluorescent probe for visualizing

glutamate neurotransmission. *Nature Methods, 10*(2), 162–170.

https://doi.org/10.1038/nmeth.2333

[18]Marvin, J. S., Scholl, B., Wilson, D. E., Podgorski, K., Kazemipour, A., Müller, J. A., Schoch,

S., Urra Quiroz, F. J., Rebola, N., Bao, H., Little, J. P., Tkachuk, A. N., Cai, E., Hantman, A. W.,

Wang, S. S.-H., DePiero, V. J., Borghuis, B. G., Chapman, E. R., Dietrich, D., DiGregorio, D.

A., Fitzpatrick, D., & Looger, L. L. (2018). Stability, affinity, and chromatic variants of the

glutamate sensor iGluSnFR. *Nature Methods, 15*(11), 936–939. https://doi.org/10.1038/s41592-

018-0171-3

[19]Xie, Y., Chan, A. W., McGirr, A., Xue, S., Xiao, D., Zeng, H., & Murphy, T. H. (2016).

Resolution of high-frequency mesoscale intracortical maps using the genetically encoded

glutamate sensor iGluSnFR. *The Journal of Neuroscience, 36*(4), 1261–1272.

https://doi.org/10.1523/JNEUROSCI.2744-15.2016

[20]Marvin, J. S., Shimoda, Y., Magloire, V., Leite, M., Kawashima, T., Jensen, T. P., Kolb, I.,

Knott, E. L., Novak, O., Podgorski, K., Leidenheimer, N. J., Rusakov, D. A., Ahrens, M. B.,

Kullmann, D. M., & Looger, L. L. (2019). A genetically encoded fluorescent sensor for in vivo

imaging of GABA. *Nature Methods, 16*(8), 763–770. https://doi.org/10.1038/s41592-019-0471-2

[21]Yardeni, T., Eckhaus, M., Morris, H. D., Huizing, M., & Hoogstraten-Miller, S. (2011). Retro-

orbital injections in mice. *Lab Animal, 40*(5), 155–160. https://doi.org/10.1038/laban0511-155

Yuste, R. (2005). Origin and classification of neocortical interneurons. *Neuron, 48*(4), 524–527.

https://doi.org/10.1016/j.neuron.2005.11.012

[22]Chan, K. Y., Jang, M. J., Yoo, B. B., Greenbaum, A., Ravi, N., Wu, W.-L., Sánchez-Guardado,

L., Lois, C., Mazmanian, S. K., Deverman, B. E., & Gradinaru, V. (2017). Engineered AAVs for

efficient noninvasive gene delivery to the central and peripheral nervous systems. *Nature*

Neuroscience, 20(8), 1172–1179. https://doi.org/10.1038/nn.4593

[23]Mohajerani, M. H., Chan, A. W., Mohsenvand, M., LeDue, J., Liu, R., McVea, D. A., Boyd, J.

D., Wang, Y. T., Reimers, M., & Murphy, T. H. (2013). Spontaneous cortical activity alternates

between motifs defined by regional axonal projections. *Nature Neuroscience, 16*(10), 1426–

1435. https://doi.org/10.1038/nn.3499

[24]Vanni, M. P., & Murphy, T. H. (2014). Mesoscale transcranial spontaneous activity mapping in

GCaMP3 transgenic mice reveals extensive reciprocal connections between areas of

somatomotor cortex. *Journal of Neuroscience, 34*(48), 15931–15946.

https://doi.org/10.1523/JNEUROSCI.1818-14.2014

[25]Silasi, G., Xiao, D., Vanni, M. P., Chen, A. C. N., & Murphy, T. H. (2016). Intact skull chronic

windows for mesoscopic wide-field imaging in awake mice. *Journal of Neuroscience Methods,*

267, 141–149. https://doi.org/10.1016/j.jneumeth.2016.04.012

[26]Xiao, D., Forys, B. J., Vanni, M. P., & Murphy, T. H. (2021). MesoNet allows automated

scaling and segmentation of mouse mesoscale cortical maps using machine learning. *Nature*

Communications, 12(1), 5992. https://doi.org/10.1038/s41467-021-26255-2

[27]Kramer, R. H., & Levitan, H. (1979). Cerebral cortical microfluorometry at isosbestic

wavelengths for correction of vascular artifact. *Science, 205*(4407), 693–696.

https://doi.org/10.1126/science.223243

[28]Ma, Y., Shaik, M. A., Kim, S. H., Kozberg, M. G., Thibodeaux, D. N., Zhao, H. T., Yu, H., &

Hillman, E. M. C. (2016). Wide-field optical mapping of neural activity and brain

haemodynamics: Considerations and novel approaches. *Philosophical Transactions of the Royal*

Society B: Biological Sciences, 371(1705), 20150360. https://doi.org/10.1098/rstb.2015.0360

[29]Scott, B. B., Thiberge, S. Y., Guo, C., Brody, C. D., Karpova, A. Y., & Tank, D. W. (2018).

Imaging cortical dynamics in GCaMP transgenic rats with a head-mounted widefield

macroscope. *Neuron, 100*(5), 1045–1058.e5. https://doi.org/10.1016/j.neuron.2018.09.050

[30]Afrashteh, N., Inayat, S., Bermudez-Contreras, E., Luczak, A., McNaughton, B. L., &

Mohajerani, M. H. (2021). Spatiotemporal structure of sensory-evoked and spontaneous activity

revealed by mesoscale imaging in anesthetized and awake mice. *Cell Reports, 37*(10), 110081.

https://doi.org/10.1016/j.celrep.2021.110081

[31]Vyazovskiy, V. V., Cirelli, C., Pfister-Genskow, M., Faraguna, U., & Tononi, G. (2008).

Molecular and electrophysiological evidence for net synaptic potentiation in wake and

depression in sleep. *Nature Neuroscience, 11*(2), 200–208. https://doi.org/10.1038/nn2035

[32]Syeda, A., Zhong, L., Tung, R., Long, W., Pachitariu, M., & Stringer, C. (2024). Facemap: A

framework for modeling neural activity based on orofacial tracking. *Nature Neuroscience, 27*(1),

187–195. https://doi.org/10.1038/s41593-023-01490-6

[33]Yüzgeç, Ö., Prsa, M., Zimmermann, R., & Huber, D. (2018). Pupil size coupling to cortical

states protects the stability of deep sleep via parasympathetic modulation. *Current Biology,*

28(3), 392–400.e3. https://doi.org/10.1016/j.cub.2017.12.049

[34]Karimi Abadchi, J., Nazari-Ahangarkolaee, M., Gattas, S., Bermudez-Contreras, E., Luczak,

A., McNaughton, B. L., & Mohajerani, M. H. (2020)**.** Spatiotemporal patterns of neocortical

activity around hippocampal sharp-wave ripples. *eLife, 9*, e51972.

https://doi.org/10.7554/eLife.51972

[35]Turner, K. L., Gheres, K. W., & Drew, P. J. (2023). Relating pupil diameter and blinking to

cortical activity and hemodynamics across arousal states. *Journal of Neuroscience, 43*(6), 949–

964. https://doi.org/10.1523/JNEUROSCI.1244-22.2022

[36]Chen, Q., Cichon, J., Wang, W., Qiu, L., Lee, S.-J. R., Campbell, N. R., Destefino, N., Goard,

M. J., Fu, Z., Yasuda, R., Looger, L. L., Arenkiel, B. R., Gan, W.-B., & Feng, G. (2012).

Imaging neural activity using Thy1-GCaMP transgenic mice. *Neuron, 76*(2), 297–308.

https://doi.org/10.1016/j.neuron.2012.07.011

[37]Atallah, B. V., Bruns, W., Carandini, M., & Scanziani, M. (2012). Parvalbumin-expressing

interneurons linearly transform cortical responses to visual stimuli. *Neuron, 73*(1), 159–170.

https://doi.org/10.1016/j.neuron.2011.12.013

[38]Wilson, N. R., Runyan, C. A., Wang, F. L., & Sur, M. (2012). Division and subtraction by

distinct cortical inhibitory networks in vivo. *Nature, 488*, 343–348.

https://doi.org/10.1038/nature11347

[39]Petersen, C. C. H. (2007). The functional organization of the barrel cortex. *Neuron*, 56(2), 339–

355. https://doi.org/10.1016/j.neuron.2007.09.017

[40]Ridder, W. H., & Nusinowitz, S. (2006). The visual evoked potential in the mouse—Origins

and response characteristics. *Vision Research*, 46(6–7), 902–913.

https://doi.org/10.1016/j.visres.2005.09.006

863 [41]Gabernet, L., Jadhav, S. P., Feldman, D. E., Carandini, M., & Scanziani, M. (2005).

864 Somatosensory integration controlled by dynamic thalamocortical feed-forward inhibition.

865 *Neuron*, 48(2), 315–327. https://doi.org/10.1016/j.neuron.2005.09.022

866 [42]Isaacson, J. S., & Scanziani, M. (2011). How inhibition shapes cortical activity. *Neuron*, 72(2),

867 231–243. https://doi.org/10.1016/j.neuron.2011.09.027

868 [43]Muñoz, W., Tremblay, R., Levenstein, D., & Rudy, B. (2017). Layer-specific modulation of

869 neocortical dendritic inhibition during active wakefulness. *Science*, 355(6328), 954–959.

870 https://doi.org/10.1126/science.aag2599

871 [44]Gentet, L. J., Kremer, Y., Taniguchi, H., Huang, Z. J., & Petersen, C. C. H. (2012). Unique

872 functional properties of somatostatin-expressing GABAergic neurons in mouse barrel cortex.

873 *Nature Neuroscience*, 15(4), 607–612. https://doi.org/10.1038/nn.3051

874 [45]Ferezou, I., Haiss, F., Gentet, L. J., Aronoff, R., Weber, B., & Petersen, C. C. H. (2007).

875 Spatiotemporal dynamics of cortical sensorimotor integration in behaving mice. *Neuron*, 56(5),

876 907–923. https://doi.org/10.1016/j.neuron.2007.10.007

877 [46]Minamisawa, G., Kwon, S. E., Chevee, M., Brown, S. P., & O'Connor, D. H. (2018). A non-

878 canonical feedback circuit for rapid interactions between somatosensory cortices. *Cell Reports*,

879 23(9), 2718–2731. https://doi.org/10.1016/j.celrep.2018.04.113

880 [47]Pakan, J. M. P., Lowe, S. C., Dylda, E., Keemink, S. W., Currie, S. P., Coutts, C. A., &

881 Rochefort, N. L. (2016). Behavioral-state modulation of inhibition is context-dependent and cell

882 type specific in mouse visual cortex. *eLife, 5*, e14985. https://doi.org/10.7554/eLife.14985

883 [48]Gentet, L. J., Avermann, M., Matyas, F., Staiger, J. F., & Petersen, C. C. H. (2010). Membrane

884 potential dynamics of GABAergic neurons in the barrel cortex of behaving mice. *Neuron, 65*(3),

885 422–435. https://doi.org/10.1016/j.neuron.2010.01.006

886 [49]Keller, G. B., & Mrsic-Flogel, T. D. (2018). Predictive processing: A canonical cortical

887 computation. *Neuron, 100*(2), 424–435. https://doi.org/10.1016/j.neuron.2018.10.003

888 [50]McGinley, M. J., Vinck, M., Reimer, J., Batista-Brito, R., Zagha, E., Cadwell, C. R., Tolias, A.

889 S., Cardin, J. A., & McCormick, D. A. (2015). Waking state: Rapid variations modulate neural

890 and behavioral responses. *Neuron, 87*(6), 1143–1161.

891 https://doi.org/10.1016/j.neuron.2015.09.012

[51]Pfeffer, C. K., Xue, M., He, M., Huang, Z. J., & Scanziani, M. (2013). Inhibition of inhibition

in visual cortex: The logic of connections between molecularly distinct interneurons. *Nature*

Neuroscience, 16(8), 1068–1076. https://doi.org/10.1038/nn.3446

[52]Dash, M. B., Douglas, C. L., Vyazovskiy, V. V., Cirelli, C., & Tononi, G. (2009). Long-term

homeostasis of extracellular glutamate in the rat cerebral cortex across sleep and waking states.

Journal of Neuroscience, 29(3), 620–629. https://doi.org/10.1523/JNEUROSCI.5486-08.2009

[53]Nazari, M., Karimi Abadchi, J., Naghizadeh, M., Bermudez-Contreras, E. J., McNaughton, B.

L., Tatsuno, M., & Mohajerani, M. H. (2023). Regional variation in cholinergic terminal activity

determines the non-uniform occurrence of cortical slow waves during REM sleep in mice. *Cell*

Reports, 42(5), 112450. https://doi.org/10.1016/j.celrep.2023.112450

[54]Karimi Abadchi, J., Rezaei, Z., Knöpfel, T., McNaughton, B. L., & Mohajerani, M. H. (2023).

Inhibition is a prevalent mode of activity in the neocortex around awake hippocampal ripples in

mice. *eLife, 12*, e79513. https://doi.org/10.7554/eLife.79513

[55]Dalby, N. O. (2000). GABA-level increasing and anticonvulsant effects of three different

GABA uptake inhibitors. *Neuropharmacology, 39*(12), 2399–2407.

https://doi.org/10.1016/s0028-3908(00)00075-7

[56]Fink-Jensen, A., Suzdak, P. D., Swedberg, M. D., Judge, M. E., Hansen, L., & Nielsen, P. G.

(1992). The gamma-aminobutyric acid (GABA) uptake inhibitor, tiagabine, increases

extracellular brain levels of GABA in awake rats. *European Journal of Pharmacology, 220*(2–3),

197–201. https://doi.org/10.1016/0014-2999(92)90748-s

[57]Wu, Y., Wang, W., & Richerson, G. B. (2001). GABA transaminase inhibition induces

spontaneous and enhances depolarization-evoked GABA efflux via reversal of the GABA

transporter. *Journal of Neuroscience, 21*(8), 2630–2639.

https://doi.org/10.1523/JNEUROSCI.21-08-02630.2001

[58]Scimemi, A. (2014). Plasticity of GABA transporters: An unconventional route to shape

inhibitory synaptic transmission. *Frontiers in Cellular Neuroscience, 8*, 128.

https://doi.org/10.3389/fncel.2014.00128

[59]Overstreet, L. S., & Westbrook, G. L. (2003). Synapse density regulates independence at

unitary inhibitory synapses. *Journal of Neuroscience, 23*(7), 2618–2626.

https://doi.org/10.1523/JNEUROSCI.23-07-02618.2003

[60]Glykys, J., & Mody, I. (2007). The main source of ambient GABA responsible for tonic

inhibition in the mouse hippocampus. *The Journal of Physiology, 582*(Pt 3), 1163–1178.

https://doi.org/10.1113/jphysiol.2007.134460

[61]Farrant, M., & Nusser, Z. (2005). Variations on an inhibitory theme: Phasic and tonic

activation of GABA(A) receptors. *Nature Reviews Neuroscience, 6*(3), 215–229.

https://doi.org/10.1038/nrn1625

[62]Turner, K. L., Gheres, K. W., Proctor, E. A., & Drew, P. J. (2020). Neurovascular coupling and

bilateral connectivity during NREM and REM sleep. *eLife, 9*, e62071.

https://doi.org/10.7554/eLife.62071

[63]Horovitz, S. G., Braun, A. R., Carr, W. S., Picchioni, D., Balkin, T. J., Fukunaga, M., & Duyn,

J. H. (2009). Decoupling of the brain's default mode network during deep sleep. *Proceedings of*

the National Academy of Sciences of the United States of America, 106(27), 11376–11381.

https://doi.org/10.1073/pnas.0901435106

[64]Larson-Prior, L. J., Zempel, J. M., Nolan, T. S., Prior, F. W., Snyder, A. Z., & Raichle, M. E.

(2009). Cortical network functional connectivity in the descent to sleep. *Proceedings of the*

National Academy of Sciences of the United States of America, 106(11), 4489–4494.

https://doi.org/10.1073/pnas.0900924106

[65]Tagliazucchi, E., & Laufs, H. (2014). Decoding wakefulness levels from typical fMRI resting-

state data reveals reliable drifts between wakefulness and sleep. *Neuron, 82*(3), 695–708.

https://doi.org/10.1016/j.neuron.2014.03.020

[66]Mohajerani, M. H., McVea, D. A., Fingas, M., & Murphy, T. H. (2010). Mirrored bilateral

slow-wave cortical activity within local circuits revealed by fast bihemispheric voltage-sensitive

dye imaging in anesthetized and awake mice. *Journal of Neuroscience, 30*(10), 3745–3751.

https://doi.org/10.1523/JNEUROSCI.4423-09.2010

[67]Niethard, N., Burgalossi, A., & Born, J. (2017). Plasticity during sleep is linked to specific
regulation of cortical circuit activity. *Frontiers in Neural Circuits, 11*, 65.
https://doi.org/10.3389/fncir.2017.00065

[68]Johansson, T., Fredholm, B. B., & Ahlberg, D. (1992). In vivo evidence that tiagabine
increases extracellular GABA concentrations in the brain by blocking GABA uptake. *European*
Journal of Pharmacology, 213(1), 127–132. https://doi.org/10.1016/0014-2999(92)90083-5

[69]Semyanov, A., Walker, M. C., Kullmann, D. M., & Silver, R. A. (2004). Tonically active GABA$_A$ receptors: modulating gain and maintaining the tone. *Trends in Neurosciences*, 27(5), 262–269. https://doi.org/10.1016/j.tins.2004.03.005

[70]Lancel, M., Faulhaber, J., & Deisz, R. A. (2009). Effect of the GABA uptake inhibitor tiagabine on sleep and EEG power spectra in the rat. *British Journal of Pharmacology*, 136(2), 204–212. https://doi.org/10.1038/sj.bjp.0701769

[71]Wu, J., Abdelfattah, A. S., Zhou, H., Ruangkittisakul, A., Qian, Y., Ballanyi, K., & Campbell, R. E. (2018). Genetically encoded glutamate indicators with altered color and topology. *ACS Chemical Biology, 13*(7), 1832–1837. https://doi.org/10.1021/acschembio.7b01085

Biography

Edris Rezaei is a Ph.D. candidate in the Canadian Centre for Behavioural Neuroscience at the University of Lethbridge, Canada. He is interested in how inhibition shapes neural activity, particularly its role in cognition and neurological disorders.

Setare Tohidi completed her master's degree at the University of Lethbridge. She is now working as a Data Scientist (Junior Associate) at Synechron in Canada.

Mojtaba Nazari□Ahangarkolaee completed her master's degree at the University of Lethbridge, affiliated. He is now working as a Data Scientist at the University of Lethbridge.

J. Karimi Abadchi is currently a postdoctoral researcher at the Douglas Research Centre in Montreal. He was a Ph.D. student at the University of Lethbridge.

Figure captions

Figure 1. Schematic of experimental workflow, imaging setup, and an expression of iGABASnFR2. A. Experimental timeline. AAV2/PHP.N-CAG-iGABASnFR2 or the control virus AAV2/PHP.N-CAG-csGFP was systemically administered via retro-orbital injection into 6- week-old C57BL/6 mice. Four weeks after viral injection, the animals underwent cranial window surgery. Following a 7-day recovery period, animals gradually habituated to the head-fixation setup over the course of another week. Sensory-evoked imaging was then conducted under 1% isoflurane anesthesia. After this session, longitudinal recordings during quiet wakefulness and natural sleep were carried out for up to six weeks. Mice were subsequently euthanized, and brains were perfused for histological analysis. B. imaging setup. (i) Illustration of the cranial window implantation over the skull following scalp removal. (ii) representative

image showing the cortical imaging area, indicated by the dashed white line. (iii) Schematic of the wide-field imaging setup: a blue LED (470 nm) was used for excitation, and signals were captured using a CCD camera at 530 nm emission. (iv) Map of the bilateral craniotomy showing targeted cortical regions based on the Allen Mouse Brain Atlas reference. C. expression of iGABASnFR2. (i) Schematic of coronal sectioning locations (a–f) along the anterior-posterior axis. (ii) Coronal sections (a–f) show strong iGABASnFR2 expression in the cortex and hippocampus. (iii) The region marked by a white dashed rectangle in panel C(iid) was used to extract fluorescence profiles across animals (n=4), highlighting unique expression and inter-animal expression consistency. (iv) The average profile with SEM as a shaded region.

Figure 2. Sensory-evoked GABAergic responses in the neocortex measured by iGABASnFR2 imaging. A. Montages of the wide bilateral craniotomy, with bregma marked by a white circle. Cortical GABAergic activation patterns are shown in a mouse anesthetized with isoflurane (1%) following: (i) whisker stimulation (300 ms), (ii) hindlimb stimulation (300 ms), (iii) forelimb stimulation (300 ms), and (iv) visual stimulation (20 ms) of the eye using an LED. Sensory-evoked extracellular GABA signals were detected using the iGABASnFR2 sensor. Activation is observed within 50–375 ms post-primary sensory cortex activation. Responses represent an average of 40 trials. The second image in the second row indicates anterior (A), posterior (P), medial (M), and lateral (L) directions. B. Time series of sensory-evoked GABA responses. The time series of GABA responses for each sensory stimulation was measured from the respective primary sensory regions. Contralateral responses are shown in black, and ipsilateral responses are shown in red. Data are presented as mean ± SEM, with responses extracted from 3×3-pixel regions of interest (ROI) (~40,401 μm²), n = number of animals. C. Summary of sensory-evoked GABA response features. Decay time (ms), Peak amplitude ($\Delta F/F$) and Time to peak (ms) for contralateral and ipsilateral responses. Data are shown as mean ± SEM, with contralateral responses in blue and ipsilateral responses in red. Statistical Analysis (Two-Way ANOVA): A two-way ANOVA revealed a significant effect of Region on decay time (p = 0.0479), peak amplitude (p = 0.0048), and time to peak (p < 0.0001), indicating sensory-region-dependent differences in GABA responses. Laterality (contralateral vs. ipsilateral) significantly affected peak amplitude (p = 0.0076) but not decay time (p = 0.9874) or time to peak (p = 0.5535). A significant Region × Laterality interaction was found for time to peak (p = 0.0055), indicating lateralization effects vary across sensory regions.

Figure 3. Sensory-evoked and spontaneous GABA activity in quiet wakefulness resembles patterns observed under anesthesia. (A) Individual example of neck muscle EMG (>1 Hz) from a head-restrained mouse under isoflurane (1%) anesthesia (top trace) and quite awake (bottom trace). The power spectra of the EMG signals differ between states of anesthesia (blue) and wakefulness (red). (B) Representative cortical GABA signals are taken from the iGABASnFR2 sensor in response to contralateral visual or whisker stimulation during anesthesia or in quiet awake states. The images represent an average of 40 trials of stimulation (C) Quantification of GABA signals in response to sensory stimulation under anesthesia and quiet wakefulness. Plots show averaged responses from 3×3-pixel ROIs (~0.04 mm²) within VISp (blue) and VISa (red) for visual stimulation and BC (blue) and M1 (red) for whisker stimulation (Ciii, Civ). Shaded areas represent SEM. (D) Statistical comparison of peak amplitude, time to

1028 peak, and decay time of GABA responses across states. *P < 0.05, **P < 0.01, one-way
1029 ANOVA. Error bars indicate SEM.
1030
Figure 4. Combined electrophysiological recording and mesoscale iGABASnFR2 imaging
of GABA activity during wakefulness and sleep. Spatiotemporal dynamics of GABA activity
over a 1-s period during wakefulness, NREM, and REM sleep. Scale bar, 2 mm. (B)
Representative traces of GABA activity (RSC), hippocampal LFP, and EMG power during
wakefulness, NREM, and REM sleep in a head-fixed mouse. Baseline fluorescence (F□) was
calculated as the mean signal over the recording session. (C) Expanded view of GABA activity,
LFP, and EMG power corresponding to the time windows in (B). (D) Group mean normalized
GABA signal across wakefulness, NREM, and REM sleep (n = 5 mice, Kruskal-Wallis test with
Nemenyi post hoc correction; $P = 0.003$ overall, all pairwise comparisons significant at $P < 0.05$)
(E) Spectral power of GABA signal in retrosplenial cortex (RSC) across wakefulness, NREM,
and REM sleep. (F) Cortical GABA activity correlation maps across quiet wakefulness, NREM,
and REM sleep over two recording days. **(G)** Mean interhemispheric correlation of cortical
GABA activity across wakefulness, NREM, and REM sleep (n = 3 mice). Mean correlation
between units significantly increased from quiet wakefulness to NREM and REM sleep across
all mice ($p□<□0.05$, repeated measures ANOVA with Bonferroni-corrected paired t-tests). **(H)**
Simultaneous recordings of cortical GABA signals, hippocampal LFP spectrogram, and EMG
power during a REM-to-wakefulness transition. **(I)** Time-lapse montage showing cortical GABA
dynamics during the REM-to-wake transition shown in (H). Scale bar: 2 mm.
1049
Figure 5. Brain state–dependent patterns of extracellular GABA dynamics measured by
iGABASnFR2. (A) Seed-pixel correlation maps (0.1–5Hz) of spontaneous iGABASnFR2
fluorescence across brain states. (i) Awake, (ii) NREM sleep, and (iii) REM sleep. Correlation
maps are shown for seed regions in the barrel cortex (BC), forelimb cortex (FL), hindlimb cortex
(HL), and visual cortex (VC). (B) Interhemispheric correlation matrices of spontaneous
extracellular GABA signals across brain states. (i) Awake, (ii) NREM sleep, and (iii) REM sleep.
Matrices display Pearson correlation coefficients between bilateral cortical regions based on
iGABASnFR2 fluorescence. Regions include MOp – Primary motor area, MOs – Secondary
motor area, SSp_bfd – Primary somatosensory area (barrel field), SSp_ll – Lower limb, SSp_ul –
Upper limb, VISam – Anteromedial visual area, VISp – Primary visual area, VISpm –
Posteromedial visual area, RSPagl – Agranular retrosplenial area, VISa – Anterior visual area.
Widespread interhemispheric connectivity is observed during wakefulness and REM sleep, with
reduced correlations during NREM sleep. Blue crosses indicate the location of the seed pixel
used for correlation mapping. number of animals=5.
1064
Figure 6. Effect of Tiagabine administration on extracellular GABA activity in mice
expressing iGABASnFR2. (A) Sensory-evoked GABA responses recorded with the blue
channel (iGABASnFR2) at 150 Hz under continuous illumination (no strobing). **(i)**
Representative montages of visually evoked GABAergic activity before Tiagabine
administration. **(ii)** The same stimulation after Tiagabine injection. **(iii)** Time series of
contralateral visually evoked iGABASnFR2 signals from the primary visual cortex (VISp).
Traces represent mean ± SEM from 5 animals, extracted from 3×3-pixel ROIs (~0.04 mm²). **(B)**
Tiagabine-induced changes in spontaneous GABA activity in VISp, recorded using dual-channel
strobing acquisition (80 Hz total: 40 Hz blue, 40 Hz green). **(i)** Blue fluorescence signal

(iGABASnFR2) from VISp, averaged across 6 animals. Traces show signal before (red) and
after (black) Tiagabine injection. The red trace shows a photobleaching trend; the black trace
shows increased fluorescence post-injection. **(ii)** Reflectance signal from the same VISp region,
averaged across 6 animals. **(iii)** Quantification of spontaneous iGABASnFR2 transients in VISp
before and after Tiagabine injection. Box plots show duration (left), frequency (middle), and
peak amplitude (right) of fluorescence transients. Data represents mean ± SEM across 6 animals
(n = 6). Tiagabine significantly increased the duration, frequency, and amplitude of spontaneous
GABAergic transients compared to baseline. Statistical comparisons were made using paired t-
tests; all changes were significant ($p < 0.05$). **(iv)** Schematic of cortical regions of interest
(ROIs) used for seed-pixel correlation analysis: barrel cortex (orange), forelimb (red), hind limb
(green), and visual cortex (brown). **(C)** Functional connectivity of spontaneous iGABASnFR2
activity, recorded with dual-channel strobing (40 Hz blue, 40 Hz green). **(i)** Seed-pixel
correlation maps (0.1–5 Hz) before and after Tiagabine injection. Blue asterisks indicate seed
pixel locations in the selected ROIs. **(ii)** Interhemispheric correlation matrices of spontaneous
iGABASnFR2 activity before and after Tiagabine injection ($n = 5$ mice), showing enhanced
bilateral cortical connectivity post-injection. Regions: MOp – Primary motor area, MOs –
Secondary motor area, SSp_bfd – Primary somatosensory area (barrel field), SSp_ll – Lower
limb, SSp_ul – Upper limb, VISam – Anteromedial visual area, VISp – Primary visual area,
VISpm – Posteromedial visual area, RSPagl – Agranular retrosplenial area, VISa – Anterior
visual area. The interior is at the top of all maps. Blue crosses indicate the location of the seed
pixel used for correlation mapping.

1095

1096

A

C57BL/6 mice
(6 weeks-old)

viral injection — 4 weeks — sensory evoked activity anesthetized 1% — 1 week — perfusion
2 week — surgery & habituation — 6 weeks — sleep and wakefulness recording

B

(i) Glass window, Head fixing bolt, Skull

(ii) A L + R P

(iii) Set-up for iGABASnFR2 imaging — CCD, 530 nm filter, 470 nm LED, 530 nm LED

(iv)

MOSR - Right Secondary Motor Cortex
MOpR - Right Primary Motor Cortex
SSp_mR - Right Primary Somatosensory Cortex (Mouth)
SSp_ulR - Right Primary Somatosensory Cortex (Upper Limb)
SSp_llR - Right Primary Somatosensory Cortex (Lower Limb)
SSp_trR - Right Primary Somatosensory Cortex (Trunk)
SSp_nuR - Right Primary Somatosensory Cortex (Nucleus)
SSp_nR - Right Primary Somatosensory Cortex (Nose)
SSp_bfdR - Right Primary Somatosensory Cortex (Barrel Field)
SPaglR - Right Anterolateral Part of Retrosplenial Cortex (Granular)
VISaR - Right Anterior Visual Cortex
VISamR - Right Anteromedial Visual Cortex
VISpmR - Right Posteromedial Visual Cortex
VISpR - Right Primary Visual Cortex
VISrlR - Right Rostrolateral Visual Cortex
VISaIR - Right Anterolateral Visual Cortex
VISR - Right Visual Cortex
VISIR - Right Secondary Visual Cortex (Intermediate)
RSPdR - Right Retrosplenial Cortex (Dorsal)

C

(i) a b c d e f

(ii) a b c d e f

(iii) Mouse 1, Mouse 2, Mouse 3, Mouse 4 — iGABASnFR2 Intensity vs Distance (µm)

(iv) average — SEM, Mean — Distance (µm)

A

Anesthetized-1% Isoflurane

i Stim — Whisker stim

ii Stim — Hindlimb stim

iii Stim — Forelimb stim

iv Stim LED — Visual stim

0 ms 75 ms 150 ms 225 ms 300 ms 375 ms 450 ms 525 ms 600 ms 675 ms

2 mm

B

Whisker stimulus, n=19 — Black: Ipsilateral response, Red: Contralateral response

Hindlimb stimulus, n=9

Whisker stimulus, n=19

Hindlimb stimulus, n=9

Visual stimulus, n=13

Forelimb stimulus, n=6

Visual stimulus, n=13

Forelimb stimulus, n=6

C

Time to peak (s) — BC, FL, HL, VC

Decay (s) — BC, FL, HL, VC

Peak amplitude $\Delta F/F_0$ (%) — BC, FL, HL, VC

Black: Ipsilateral response, Red: Contralateral response

A EMG (anesthetized)

1 s   0.1 mV

EMG (quiet awake)

EMG power spectrum

Bi Visual stimulation (300 ms)

Anesthetized $\Delta F/F_0$ (%)   0.2

Awake $\Delta F/F_0$ (%)   0.6

2 mm   0ms → π → 25 ms   50 ms   75 ms   100 ms   125 ms   150 ms

ii Whisker stimulation (300 ms)

Anesthetized $\Delta F/F_0$ (%)   0.4

Awake $\Delta F/F_0$ (%)   1.5

2 mm   0ms → π → 25 ms   50 ms   75 ms   100 ms   125 ms   150 ms

C iGABASnFR2 $\Delta F/F_0$ (%)

Quiet awake   Visual stimulus   VISp VISa n=5

Anesthetized   Visual stimulus   n=13

Quiet awake   Whisker stimulus   BC M1 n=5

Anesthetized   Whisker stimulus   n=19

Time (s)

D

Peak amplitude $\Delta F/F_0$ (%)   ** * ***   Anest (n=13) Quiet awake (n=5)   VISp VISa

Time to peak (ms)   Anest (n=13) Quiet awake (n=5)

Decay (ms)   * *   Anest (n=13) Quiet awake (n=5)   VISp VISa

Peak amplitude $\Delta F/F_0$ (%)   ** ** **   Anest (n=19) Quiet awake (n=5)

Time to peak (ms)   *   Anest (n=19) Quiet awake (n=5)

Decay (ms)   * *   Anest (n=19) Quiet awake (n=5)   BC M1

A

i

Anesthetized-1% Isoflurane

ΔF/V₀ (%)

300 ms tone pip at 12 kHz

Auditory stim

| 0s | 75ms | 150ms | 225ms | 300ms | 375ms | 450ms | 525ms | 600ms | 675ms |

ii

AUDpR - Right Primary Auditory Cortex (Posterior)

iii