Title: Hippocampal ripples initiate cortical dimensionality expansion for memory retrieval

Casper Kerrén[1*], Sebastian Michelmann[2] & Christian F Doeller[1,3]

Affiliations:

[1] *Max Planck Institute for Human Cognitive and Brain Sciences; Leipzig, Germany*

[2] *Department of Psychology, New York University; New York City, New York*

[3] *Kavli Institute for Systems Neuroscience, Centre for Neural Computation, Egil and Pauline Braathen and Fred Kavli Centre for Cortical Microcircuits, Jebsen Centre for Alzheimer's Disease, NTNU Norwegian University of Science and Technology; Trondheim, Norway*

*Corresponding author. Email: kerren@cbs.mpg.de

Abstract

How are past experiences reconstructed from memory? Learning is thought to compress external inputs into low-dimensional hippocampal representations, later expanded into high-dimensional cortical activity during recall. Hippocampal ripples—brief high-frequency bursts linked to retrieval—may initiate this expansion. Analysing intracranial EEG data from patients with pharmaco-resistant epilepsy during an episodic memory task, we found that cortical dimensionality increased following ripple events during correct, but not incorrect, retrieval. This expansion correlated with faster reaction times and reinstatement of the target association. Crucially, hippocampal theta and cortical gamma phase–amplitude coupling emerged after ripples but before cortical expansion, suggesting a mechanism for ripple-driven communication. Ripple events also marked the separation of task-relevant variables in cortical state space, revealing how hippocampal output reshapes the geometry of memory representations to support successful recall.

Main Text

Episodic memory allows us to store detailed records of past experiences and consciously reconstruct those experiences at later points in time (*1*). Like any computational system though, the human brain operates with finite resources (*2*). To cope with these constraints, efficient encoding and retrieval of memories is thought to rely on compression and expansion of neural representations (*3–6*). During encoding, environmental information flows through cortical and subcortical regions to the hippocampus, where memories are initially stored (*7*, *8*). During retrieval, internal or external cues trigger the hippocampus to detect matches with stored traces (*9*). A partial match initiates pattern completion, leading to memory reactivation and reconstruction in cortical networks (*10*, *11*). Yet, how hippocampal pattern completion gives rise to cortical reinstatement remains poorly understood.

Recent studies have begun to conceptualise this process as shifts in the geometric relationships of points in neural state space, a framework that has found broader applications in studies of decision-making and working memory (*12–19*). We hypothesise that memory retrieval might likewise involve a transformation in dimensionality, such that low-dimensional hippocampal representations are expanded into a higher-dimensional cortical state, allowing mnemonic information to be decoded for successful recall (*5*, *10*, *20*, *21*). Preliminary evidence supports this hypothesis, showing a shift from semantic to perceptual representations along the ventral visual stream during retrieval (*22–26*). However, whether and how the hippocampus might initiate this dimensionality expansion is unclear.

One potential means to drive cortical representational expansion is through hippocampal ripples. Ripples are known to coordinate the transfer of compressed representations and changes of brain-wide functional connectivity during offline periods in rodents (*27–31*). In humans, an increase of hippocampal ripple rates precedes episodic memory recall (*32–37*) and neocortical reinstatement of previously encoded memories consistently follows hippocampal ripple events (*32–35*). In parallel, theoretical accounts and empirical evidence have highlighted cross-frequency interactions—particularly theta-gamma-phase–amplitude coupling (TG-PAC) —as a mechanism for coordinating long-range communication (*38–44*). While TG-PAC has been widely studied during mnemonic processing particularly within hippocampus (*45–53*), its role during retrieval, and specifically in mediating ripple-initiated cortical transformations, remains largely unexplored.

We hypothesise that the reconstruction of information in cortical regions is supported by a ripple-based mechanism, where compressed representations are expanded via PAC-based

connectivity between hippocampus and cortex. To test this hypothesis, we analysed
intracranial data from 12 patients with pharmaco-resistant epilepsy as they performed an
associative memory task (Fig. 1, A and B, fig. S1). We asked whether ripple events could
trigger both an increase in cortical representational dimensionality and TG-PAC between
hippocampus and cortex—linking local hippocampal dynamics to global cortical
transformations during successful episodic retrieval (Fig. 1C).

Fig. 1. Paradigm, implantation scheme, hypothesis
(A) During encoding, participants associated a noun either with one of two colours (*left*) or one of two
scenes (*right*; alternating across runs). During retrieval, participants were prompted with the noun
alongside response options, yielding either successful associative memory (AM+) or unsuccessful
associative memory (AM-; combining incorrect and 'don't know' responses). (**B**) Hippocampal (*top*)
and extra-hippocampal (*bottom*) iEEG contacts included in the analyses, highlighted in different
colours for each participant. (**C**) Schematic depiction of the hypothesis: During encoding, the
representation of an event is stored in hippocampus as a lower-dimensional pointer to the higher-
dimensional representation in cortical areas. During retrieval, hippocampal ripples initiate a process
that initiates a hippocampal-cortical connectivity through theta-gamma phase-amplitude coupling (TG-
PAC), leading to the low-dimensional representation being expanded in cortical areas, which is linked
to behavioural memory performance.
20

21 **Behavioural results**
22 The experiment yielded a balanced amount of successful and unsuccessful associative
23 memory trials (AM+: M = 49.95%, SE = 5.12%; AM-: M = 50.05%, SE = 5.12%; t(1,11) = -.01,

1   p = .992). Response latencies for AM+ trials were faster (M = 1.90 sec, SE = .12 sec) than

2   for AM- trials (M = 2.06 sec, SE = .12 sec; t(1,11) = -2.68, p = .021).

Greater ripple-density for successful retrieval

On average, 787.33 hippocampal ripples (SE = 145.97) per participant were detected during

all retrieval trials, after excluding false positives, with a spectral mean peak at 89.17 Hz (SE

= .57 Hz) (Fig. 2A; fig. S1 for participant-specific ripple plots). Previous research has shown

that ripple density increases prior to successful free recall performance (e.g., (*32*)). To

assess whether this finding extends to associative recognition as employed here, we

extracted the ripple density for each trial, i.e., the number of ripples normalised by the trial's

reaction time, and averaged across trials in each condition. Ripple density was higher during

AM+ (M = .24 Hz, SE = .03 Hz) compared to AM- (M = .19 Hz, SE = .02 Hz), t(11) = 3.58, p

< .01 (Fig. 2B).

Fig. 2. Hippocampal ripple density increases during successful memory retrieval

(**A**) Grand average hippocampal ripple for all participants, aligned to ripple peak at time 0, with

spectral grand average as inset, showing a peak at 89 Hz. (**B**) Ripple density significantly differed

between AM+ and AM-. Circles represent participants. (**C**) Ripple density dynamics across time. *Left*:

aligned to retrieval cue onset. *Right*: aligned to response time. Horizontal black lines denote

significant differences between AM+ and AM- (p < .05, corrected for multiple comparisons across

time).

Additionally, we examined whether the increased ripple rate occurred at specific time points

during the successful retrieval and observed that ripple density was increased 800-1400ms

post cue (for cue-aligned data; Fig. 2C, left) and leading up to the retrieval response (for

response aligned data; Fig. 2C, right). These findings are consistent with previously reported

latencies of episodic memory processes (*54*).

Neocortical reinstatement is linked to hippocampal ripple events

Building on these findings, we next examined the relationship between hippocampal ripple

events and the reinstatement of information from encoding. Specifically, we assessed

whether cortical patterns present during encoding re-emerged around ripple events during

retrieval.

Our results confirm that ripple-locked memory reinstatement underlies successful

associative recognition memory (AM+). The successful decoding of memory content was

closely yoked to ripple peaks in our data and continued until the end of the analysis epoch

(see figure S2 for various control analysis confirming that reinstatement was specific to the

ripple latency of the current trial). This extends previous findings of the role of hippocampal

ripples in the replay of encoding activity during free recall of verbal material (*34*) and

suggests a generalised role of ripples in the reinstatement of information-rich cortical

representations.

Fig. 3. Target memory decoding and dimensionality transformation are locked to ripple events.

(**A**) Stimulus-locked encoding x ripple-locked retrieval target decoding accuracies. Results show significant increase of cortical target decodability for successful vs. unsuccessful retrieval following hippocampal ripple events (p < .05, corrected for multiple comparisons). (**B**) Estimating dimensionality for cortical contacts yielded significantly higher dimensionality for AM+ (blue) versus AM- (red) following ripple events (time 0 on x-axis). Inset: The increase in dimensionality was significantly negatively correlated with reaction time. (**C**) Linear mixed-effects modelling (Epoch Half: pre and post ripple, Trial Type: AM+ and AM-) of cortical dimensionality revealed a significant interaction, indicating a greater increase in dimensionality post-ripple for AM+. Asterisks denote significance compared to 0 for AM+ and the interaction of pre and post ripple time-window and AM+ and AM-. (**D**) dPC1, 2, and 4 revealed significant decoding accuracy of target identity in the time window of dimensionality expansion (highlighted in green bar). Significant intervals of decodability are highlighted with horizontal lines. Euclidean distance between clusters increased around ripple onset with a peak 700ms post ripple-events (dashed line). (**E**) Plotting dPCs 1, 2 and 4, showed overlapping representations of experimental variables before ripples (left), whereas their separability increased after ripples (right).

Hippocampal ripple-induced dimensionality expansion increases the separability of cortical representations

These findings suggest that hippocampal ripples initiate a cortical "decompression" of memory representations, allowing latent information to be reconstructed during retrieval (*5*, *10*, *20*, *21*) (Fig. 1C). To formally test this, we first examined changes in representational geometry using principal component analysis (PCA), applied to iEEG data time-locked to ripple events.

We found that cortical dimensionality significantly increased following ripples during AM+ trials, compared to AM– trials, with effects emerging 470–840 ms post-ripple (two clusters; both p<.05). Dimensionality estimates were robust across parameters (fig. S3, A and B) and strongly correlated with the original signal ($r_{spearman}$ .87, p < .001; fig. S3C). Linear mixed-effects modelling confirmed higher dimensionality for AM+ trials overall (Estimate = .09, SE = .03, t(19,988) = 2.82, p = .005), and for post- vs. pre-ripple epochs (Estimate = .14, SE = .03, t(19,988) = 4.39, p < .001), with a significant interaction indicating greater ripple-induced expansion for remembered events (Estimate = -.09, SE = .02, t(19,988) = -4.16, p < .001; Fig. 3C). Paired samples t-tests supported this effect, showing increased dimensionality following ripples in AM+ trials (before: M = 2.43, SE = .05; after: M = 2.50, SE = 0.06; t(11) = 2.22, p = .049) , but no change in AM– trials (before: M = 2.44, SE = 0.06; after: M = 2.41, SE = 0.06;), t(11) = -0.45, p = .66).

Crucially, higher post-ripple dimensionality during AM+ trials was associated with faster

response times ($r_{spearman}$ = -.669, p = .02; Inset Fig. 3B), and stronger cortical reinstatement

($r_{spearman}$ = .656, p = .02), linking representational expansion to both behavioural and neural

markers of successful retrieval. Control analyses confirmed that these effects were specific

to ripple events and not observed in surrogate data (fig. S4), reinforcing the functional role

of ripple-triggered dimensionality expansion.

While PCA preserves the geometry of neural activity while maximising variance, and LDA

enhances interpretability by maximising class separation, each has limitations. PCA mixes

task-related variance and does not isolate the contribution of experimental variables,

whereas LDA distorts the original geometry of the neural state space in service of

classification. Demixed PCA (dPCA; (55)), addresses both limitations by simultaneously

reducing population activity and disentangling it into components aligned with specific task

parameters. As illustrated in Kobak et al. (55), dPCA separates latent structure in a way that

retains both class-specific information and the original representational geometry, making it

particularly well-suited for characterising the organisation and temporal evolution of memory-

related cortical states. Using this approach, we could examine how ripple events affected the

organisation of cortical states, not just their dimensionality (55).

We reconstructed neural activity using components reflecting target association (blue, red,

indoor, outdoor), memory performance (AM+ and AM-), their interaction (target association x

memory performance), and a condition-independent component reflecting time. According to

our hypothesis, cortical representations of these variables would separate during the period

of dimensionality expansion (~400–800 ms post-ripple; highlighted in green in Figure 3D),

thereby enhancing decodability in high-dimensional space.

We first confirmed that our reconstruction accurately captured the data. Specifically, the first

50 dPCs accounted for as much variance as our estimated total signal (see Supplementary

Materials, and figure S5A). To assess whether these demixed components carried task-

relevant information, we applied cross-validated decoding, which tests whether the structure

revealed by dPCA generalises to unseen data, providing a direct readout of when and how

experimental variables are represented in neural population activity (55).

Among all components, the interaction between target association and memory performance

(dPC1) explained most variance (fig. S5B), capturing retrieval-related dynamics. This

component showed high decoding accuracy across almost the entire epoch and peaked

shortly after ripple events. The second component (dPC2), which captured target

association-specific brain dynamics, showed significant decoding accuracy both around ripple events and during dimensionality expansion (Fig. 4A and fig. S5D). Finally, the fourth component (dPC4), linked to memory performance, significantly distinguished AM+ from AM- trials, with effects emerging shortly after ripple events and during dimensionality expansion (Fig. 4A and fig. S5F). For full output, see figure S5.

These findings confirm that dPCA effectively disentangled experimental variables in a lower-dimensional space. To quantify this representational reorganisation, we applied k-means clustering to the dPCA components and assessed the Euclidean distance between clusters. To evaluate clustering quality, we used the Silhouette score, where higher scores indicate stronger clustering. Using 200 ms time bins around ripple events, we observed increased cluster separation beginning at ripple onset and peaking ~700 ms later (Fig. 3D, dashed line). To further quantify this effect, we plotted the state-space organisation of these components during dimensionality expansion (400-800ms post-ripple events) and compared it to corresponding pre-ripple period (400-800ms pre-ripple). Before ripple events, we identified six clusters (silhouette score = .83), whereas eight clusters emerged after ripple events (silhouette score = .90) (Fig. 3E). This was accompanied by a significant increase in Euclidean distance between clusters (pre-ripple: M = 4.06, SE = .044; post-ripple: M = 4.78, SE = .048; t(319) = 5.04, p < .01). A similar pattern emerged when using dPC 1, 3, and 4 (fig. S5H, and Supplementary materials). In both visualisations, we observed a clear separation of all experimental variables following ripple events, a pattern that was less distinct before ripple onset.

Theta-gamma phase-amplitude coupling coordinates information flow between hippocampus and cortex

Our results demonstrate that hippocampal ripples initiate a cortical expansion during episodic memory retrieval in which the separation between experimental variables increases. But what mechanisms enable this transformation of information? A candidate mechanism is the coupling between hippocampal theta and cortical gamma rhythms (*38, 39, 44, 56–58*). Despite theoretical accounts emphasising its importance, there is currently little direct evidence in humans that theta–gamma-phase-amplitude-coupling (TG-PAC) supports long-range communication during episodic memory retrieval.

One likely reason for this gap is the considerable variability in retrieval timing across trials, which poses a challenge for aligning neural events consistently. Our findings suggest that hippocampal ripples can be leveraged to pinpoint the exact moments of reinstatement. Thus, we hypothesised that communication between the hippocampus and cortex would occur

shortly after ripple onset, but prior to the cortical expansion of memory representations, and

would be critical for reconstructing the memory trace. To test this, we examined TG-PAC in

the time window surrounding ripple events.

Fig. 5. Phase-amplitude coupling following ripple events are related to dimensionality

expansion.

(A) On average gamma oscillations peaked at 60 Hz, whereas theta on average peaked at 6 Hz. **(B)**

Significant TG-PAC around peak frequencies for respective frequency (0 on x and y axes). **(C)**. Top:

A time-resolved TG-PAC commencing around ripple events and persisted for approximately 400ms

after the events. Bottom: Plotting the time-resolved TG-PAC averaged across the y-axis in top panel.

(D) A positive correlation between TG-PAC and cortical dimensionality expansion, such that stronger

TG-PAC was related increased dimensionality expansion. **(F)** A cross-correlation showed that TG-

PAC leads dimensionality expansion with a peak at approximately 260ms.

We identified peak theta and gamma frequencies in hippocampal and cortical signals after

removing the 1/f background using the FOOOF algorithm (*59*). Time–frequency data were

then aligned to these peaks (±2 Hz for theta, ±10 Hz for gamma), and phase–amplitude

coupling was computed. Empirical PAC values were compared to a null distribution

generated by shuffling trials 250 times per participant and time point. This allowed us to

assess TG-PAC in narrow-band activity centred on peak frequencies, time-locked to ripple

events (−1 to +1 s). The average peak theta and gamma frequencies were 6.08 Hz

(SE = 0.69 Hz) and 59.58 Hz (SE = 4.75 Hz), respectively (Fig. 4A). This revealed

significant coupling between hippocampal theta phase and cortical gamma amplitude around

the identified peaks ($p_{cluster}$<.05; Fig. 4B).

To examine the timing of TG-PAC relative to ripple events, we conducted a time-resolved

analysis using sliding windows. This revealed that coupling emerged shortly after ripple

onset and persisted for ~300 ms ($p_{clusters}$ < .05; Fig. 4C). Notably, TG-PAC strength in this

post-ripple window correlated with ripple-related increases in cortical dimensionality (r = .59,

p = .04; Fig. 4D), linking cross-frequency coupling to representational transformation during

retrieval.

We hypothesised that TG-PAC would follow hippocampal ripples but precede cortical

dimensionality expansion. To test this, we performed a cross-correlation analysis against a

null distribution of 1000 phase-shuffled surrogates. The analysis revealed a significant

positive lag, peaking at ~260 ms ($p_{cluster}$ = .02; Fig. 4E), suggesting that TG-PAC may play

a mechanistic role in triggering the cortical expansion of memory representations following

ripple events.

Discussion

Episodic memory is thought to optimise neural processing by compressing information during encoding and expanding it during retrieval (*3–6*). In particular, the hippocampus plays a central role in this dynamic, matching retrieval cues with stored memory traces and triggering pattern completion to reconstruct the full memory in the cortex (*9–11*). Yet, the neural mechanism through which hippocampal pattern completion gives rise to cortical memory reinstatement remains poorly understood.

We hypothesised that memory retrieval involves a transformation in neural dimensionality. Specifically, that low-dimensional representations in the hippocampus expand into high-dimensional cortical activity patterns. This shift would allow higher order areas to decode rich mnemonic content (*5, 10, 20, 21*). Leveraging the spatial and temporal resolution of human intracranial EEG (iEEG) while participants performed an associative memory task, we demonstrate that that successful memory retrieval was associated with stronger reactivation of encoding-related cortical patterns (Fig. 3A). Notably, this was accompanied by an increase in neural dimensionality (Fig. 3, B–C), which was linked to both faster response times (Fig. 3B) and greater reinstatement strength—suggesting a functional role for dimensional expansion in supporting retrieval.

Our findings position hippocampal ripples as key triggers of cortical reinstatement during episodic memory retrieval. While prior work has linked ripples to hippocampal–cortical communication and memory-related activity in both humans and non-human animals (*29, 30, 32–35, 60–63*), we show that ripples are temporally aligned with both increased cortical dimensionality and reinstatement of encoded information. This extends previous literature by providing direct evidence that ripples mark a transition from compressed to high-dimensional cortical states supporting the reconstruction of specific mnemonic content. Moreover, our results suggest that ripple-related reflect a broader cortical mechanism for memory retrieval.

In our study, ripple-triggered cortical reinstatement occurred reliably during associative retrieval and was tightly time-locked to the onset of ripple events (Fig. 2A, fig. S2). Ripple density was also higher during correct trials, particularly in the pre-response period, suggesting that that ripples functionally drive cortical memory reactivation. Crucially, this reinstatement was accompanied by an increase in neural dimensionality. The finding puts forward a novel account of ripple events that do not simply trigger the reactivation of static memory traces, but instead facilitate an expansion of compressed representations into complex, high-dimensional cortical states. Such expansion may reflect the recruitment of a

broader set of cortical areas and the unfolding of richer, more differentiated representational patterns across the network, which then enables flexible coding of specific mnemonic content.

Our findings are also in line with recent work on the time-scales of hippocampal–cortical interactions during memory retrieval. Michelmann and colleagues (*64*) reported that hippocampal activity precedes cortical reinstatement by ~740 ms during predictive recall under naturalistic conditions, while in another study, the same authors found that hippocampal–cortical information flow precedes cortical state transitions by 500–700 ms during memory search (*65*). In line with these findings, we observed a ~400–800 ms lag from hippocampal ripples to cortical dimensionality expansion and a maximal separability of task-related variables at 700 ms post-ripples. The duration of this effect and its extended temporal window are evidence that hippocampal-driven reactivation unfolds over many synaptic transmission points to realise the expansion of a memory-trace. I.e., hippocampal outputs may initiate distributed cortical reorganisation and enable the dynamic reconstruction of mnemonic content across time and brain regions (*66, 67*).

To characterise the representational changes that accompany this process, we used demixed principal component analysis (dPCA). Task-relevant variables, such as memory accuracy and target identity, became more separable following ripple events (Fig. 3E), with this separability peaking precisely at the time of dimensionality expansion (Fig. 3D). The neural state space also became more complex, as indicated by an increased number of optimal clusters and greater Euclidean distances between them. These results suggest that ripples support the cortical "decompression" of memory content, allowing for finer-grained separation of mnemonic features.

In our data, the increase in representational structure did not differentiate between correctly remembered and forgotten associations (Fig. 3E, right). This suggests that the ripple-mediated reorganisation of cortical dynamics alone is not diagnostic of retrieval success and other elements can cause memory processes to fail. This mirrors findings in non-human primates, where structured neural representations were observed even in error trials during working memory tasks. For example, Rigotti and colleagues found that although dimensionality collapsed in incorrect trials, task-relevant information could still be decoded— indicating an underlying structure (*68*). Similarly, in our data, incorrect trials showed lower overall dimensionality (Fig. 3B), but retained structured representations of task variables (Fig. 3E). These findings suggest that absolute separability is not the sole determinant of retrieval

success; rather, it is the richness or expressiveness of the neural state—as assessed via

PCA (Fig. 3B)—that better predicts retrieval speed and accuracy.

To our knowledge, there is currently no direct evidence in humans for hippocampal theta–

cortical gamma phase–amplitude coupling (TG-PAC) during episodic memory retrieval. Here,

we show that TG-PAC peaks shortly after ripple events (Fig. 4C) and precedes an increase

in cortical dimensionality by ~260 ms (Fig. 4E), suggesting it may act as a bridge between

the initiation and reconstruction of retrieved events. This extends previous findings of TG-

PAC during encoding and retrieval in the hippocampus (*45*, *47*, *58*, *69*), during spatial

memory retrieval (*70*), and across distributed cortical areas during autobiographical recall

(*71*).

While prior studies have linked slow and fast gamma to encoding and retrieval, respectively

(*44*, *72*, *73*), we found no clear frequency split—participants showed peaks in both bands.

These differences may reflect region-specific gamma dynamics outside the medial temporal

lobe.

Our findings extend recent work linking neural dimensionality to flexible coding in decision-

making and navigation (e.g., (13)), showing that episodic retrieval also involves dynamic,

high-dimensional cortical states (5). Events with greater dimensionality during encoding are

more likely to be remembered (74), raising the possibility that rich encoding geometry

facilitates later reconstruction. Future work should test whether dimensionality at encoding

predicts reinstatement fidelity.

We propose a framework in which ripples trigger a shift from compressed hippocampal

codes to expanded cortical states. This transition may be mediated by ripple-induced

synchrony (75), followed by cortical desynchronisation in alpha/beta bands (76), which has

been linked to improved memory fidelity and recall (77, 78). Such desynchronisation may

support low-synchrony, high-dimensional representations that promote selective decoding

while reducing interference (68, 79–82). Understanding how ripple-induced synchrony

interacts with cortical desynchronisation will be an important direction for future research on

the temporal dynamics of memory reinstatement.

Acknowledgement

The authors would like to thank Juergen Fell and Bernhard Staresina for sharing their data.

Funding

Max Planck Society (C.K. and C.F.D.)

Deutsche Forschungsgemeinschaft project 437219953 (S.M.)

Author contributions

Conceptualization: C.K.

Methodology: C.K. and S.M.

Investigation: C.K.

Visualization: C.K.

Funding acquisition: S.M. and C.F.D.

Project administration: C.F.D.

Supervision: S.M. and C.F.D.

Writing – original draft: C.K., S.M. and C.F.D.

Writing – review & editing: C.K., S.M. and C.F.D.

Competing interests

Authors declare that they have no competing interests.

Data and materials availability

The data and code that support the conclusions of this study are available upon reasonable request from C.K. (kerren@cbs.mpg.de) and are also accessible online at GitHub.

Supplementary Materials

Materials and Methods

Figs. S1 to S5

Tables S1

References (*45-46, 55, 59, 83*–94)

References

1. E. Tulving, *Elements of Episodic Memory* (Oxford University Press, 1983).

2. L. de Almeida, M. Idiart, J. E. Lisman, Memory retrieval time and memory capacity of the CA3 network: Role of gamma frequency oscillations. *Learn. Mem.* 14, 795–806 (2007).

3. C. J. Bates, R. A. Jacobs, Efficient data compression in perception and perceptual memory. *Psychological Review* 127, 891–917 (2020).

4. H. B. Barlow, "Possible Principles Underlying the Transformations of Sensory Messages" in *Sensory Communication*, W. A. Rosenblith, Ed. (The MIT Press, 2012; https://academic.oup.com/mit-press-scholarship-online/book/20714/chapter/180090664), pp. 216–234.

5. C. Kerrén, D. Reznik, C. F. Doeller, B. J. Griffiths, Exploring the role of dimensionality transformation in episodic memory. *Trends in Cognitive Sciences*, doi: 10.1016/j.tics.2025.01.007 (2025).

6. D. Reznik, P. Majka, M. G. P. Rosa, M. P. Witter, C. F. Doeller, Phylogeny of neocortical-hippocampal projections provides insight in the nature of human memory. bioRxiv [Preprint] (2024). https://doi.org/10.1101/2024.05.09.593130.

7. H. Eichenbaum, T. Otto, N. J. Cohen, Two functional components of the hippocampal memory system. *Behavioral and Brain Sciences* 17, 449–517 (1994).

8. D. O. Hebb, *The Organization of Behavior; a Neuropsychological Theory* (Wiley, Oxford, England, 1949) *The organization of behavior; a neuropsychological theory*.

9. K. Duncan, N. Ketz, S. J. Inati, L. Davachi, Evidence for area CA1 as a match/mismatch detector: A high☐resolution fMRI study of the human hippocampus. *Hippocampus* 22, 389–398 (2012).

10. J. L. McClelland, B. L. McNaughton, R. C. O'Reilly, Why there are complementary learning systems in the hippocampus and neocortex: insights from the successes and failures of connectionist models of learning and memory. *Psychol Rev* 102, 419–457 (1995).

11. H. Eichenbaum, N. J. Cohen, *From Conditioning to Conscious Recollection: Memory Systems of the Brain* (Oxford University Press, 2004; https://doi.org/10.1093/acprof:oso/9780195178043.001.0001).

12. C. K. Machens, R. Romo, C. D. Brody, Functional, but not anatomical, separation of "what" and "when" in prefrontal cortex. *J Neurosci* 30, 350–360 (2010).

13. A. Kikumoto, A. Bhandari, K. Shibata, D. Badre, A transient high-dimensional geometry affords stable conjunctive subspaces for efficient action selection. *Nat Commun* 15, 8513 (2024).

14. R. Bartolo, R. C. Saunders, A. R. Mitz, B. B. Averbeck, Dimensionality, information and learning in prefrontal cortex. *PLOS Computational Biology* 16, e1007514 (2020).

15. V. Mante, D. Sussillo, K. V. Shenoy, W. T. Newsome, Context-dependent computation by recurrent dynamics in prefrontal cortex. *Nature* 503, 78–84 (2013).

16. C. D. Harvey, P. Coen, D. W. Tank, Choice-specific sequences in parietal cortex during a virtual-navigation decision task. *Nature* 484, 62–68 (2012).

17. S. Hedayati, R. E. O'Donnell, B. Wyble, A model of working memory for latent representations. *Nat Hum Behav* 6, 709–719 (2022).

18. J. D. Murray, J. Jaramillo, X.-J. Wang, Working Memory and Decision-Making in a Frontoparietal Circuit Model. *J. Neurosci.* 37, 12167–12186 (2017).

19. S. L. Brincat, M. Siegel, C. von Nicolai, E. K. Miller, Gradual progression from sensory to task-related processing in cerebral cortex. *Proceedings of the National Academy of Sciences* 115, E7202–E7211 (2018).

20. E. Spens, N. Burgess, A generative model of memory construction and consolidation. *Nat Hum Behav*, doi: 10.1038/s41562-023-01799-z (2024).

21. M. Jazayeri, S. Ostojic, Interpreting neural computations by examining intrinsic and embedding dimensionality of neural activity. *Current Opinion in Neurobiology* 70, 113–120 (2021).

22. J. Linde-Domingo, M. S. Treder, C. Kerrén, M. Wimber, Evidence that neural information flow is reversed between object perception and object reconstruction from memory. *Nat Commun* 10, 179 (2019).

23. J. Lifanov, J. Linde-Domingo, M. Wimber, Feature-specific reaction times reveal a semanticisation of memories over time and with repeated remembering. *Nat Commun* 12, 3177 (2021).

24. S. Mirjalili, P. Powell, J. Strunk, T. James, A. Duarte, Context Memory Encoding and Retrieval Temporal Dynamics are Modulated by Attention across the Adult Lifespan. *eNeuro* 8, ENEURO.0387-20.2020 (2021).

25. C. Kerrén, Y. Zhao, B. J. Griffiths, A reduction in self-reported confidence accompanies the recall of memories distorted by prototypes. *Commun Psychol* 2, 1–11 (2024).

26. C. Kerrén, J. Linde-Domingo, B. Spitzer, Prioritization of semantic over visuo-perceptual aspects in multi-item working memory. bioRxiv [Preprint] (2022). https://doi.org/10.1101/2022.06.29.498168.

27. D. Khodagholy, J. N. Gelinas, G. Buzsáki, Learning-enhanced coupling between ripple oscillations in association cortices and hippocampus. *Science* 358, 369–372 (2017).

28. S. P. Jadhav, G. Rothschild, D. K. Roumis, L. M. Frank, Coordinated Excitation and Inhibition of Prefrontal Ensembles during Awake Hippocampal Sharp-Wave Ripple Events. *Neuron* 90, 113–127 (2016).

29. J. Karimi Abadchi, M. Nazari-Ahangarkolaee, S. Gattas, E. Bermudez-Contreras, A. Luczak, B. L. McNaughton, M. H. Mohajerani, Spatiotemporal patterns of neocortical activity around hippocampal sharp-wave ripples. *Elife* 9, e51972 (2020).

30. N. K. Logothetis, O. Eschenko, Y. Murayama, M. Augath, T. Steudel, H. C. Evrard, M. Besserve, A. Oeltermann, Hippocampal-cortical interaction during periods of subcortical silence. *Nature* 491, 547–553 (2012).

31. H. R. Joo, L. M. Frank, The hippocampal sharp wave–ripple in memory retrieval for immediate use and consolidation. *Nat Rev Neurosci* 19, 744–757 (2018).

32. Y. Norman, E. M. Yeagle, S. Khuvis, M. Harel, A. D. Mehta, R. Malach, Hippocampal sharp-wave ripples linked to visual episodic recollection in humans. *Science* 365, eaax1030 (2019).

33. Y. Norman, O. Raccah, S. Liu, J. Parvizi, R. Malach, Hippocampal ripples and their coordinated dialogue with the default mode network during recent and remote recollection. *Neuron* 109, 2767-2780.e5 (2021).

34. A. P. Vaz, S. K. Inati, N. Brunel, K. A. Zaghloul, Coupled ripple oscillations between the medial temporal lobe and neocortex retrieve human memory. *Science* 363, 975–978 (2019).

35. A. P. Vaz, J. H. Wittig, S. K. Inati, K. A. Zaghloul, Replay of cortical spiking sequences during human memory retrieval. *Science* 367, 1131–1134 (2020).

36. S. Henin, A. Shankar, H. Borges, A. Flinker, W. Doyle, D. Friedman, O. Devinsky, G. Buzsáki, A. Liu, Spatiotemporal dynamics between interictal epileptiform discharges and ripples during associative memory processing. *Brain* 144, 1590–1602 (2021).

37. J. J. Sakon, M. J. Kahana, Hippocampal ripples signal contextually mediated episodic recall. *Proc Natl Acad Sci U S A* 119, e2201657119 (2022).

38. E. Nyhus, T. Curran, Functional role of gamma and theta oscillations in episodic memory. *Neurosci Biobehav Rev* 34, 1023–1035 (2010).

39. P. Fries, Rhythms for Cognition: Communication through Coherence. *Neuron* 88, 220–235 (2015).

40. R. T. Canolty, R. T. Knight, The functional role of cross-frequency coupling. *Trends Cogn Sci* 14, 506–515 (2010).

41. A. Sirota, S. Montgomery, S. Fujisawa, Y. Isomura, M. Zugaro, G. Buzsáki, Entrainment of Neocortical Neurons and Gamma Oscillations by the Hippocampal Theta Rhythm. *Neuron* 60, 683–697 (2008).

42. A. Hyafil, A.-L. Giraud, L. Fontolan, B. Gutkin, Neural Cross-Frequency Coupling: Connecting Architectures, Mechanisms, and Functions. *Trends in Neurosciences* 38, 725–740 (2015).

43. J. Fell, N. Axmacher, The role of phase synchronization in memory processes. *Nat Rev Neurosci* 12, 105–118 (2011).

44. L. L. Colgin, Theta-gamma coupling in the entorhinal-hippocampal system. *Curr Opin Neurobiol* 31, 45–50 (2015).

45. B. J. Griffiths, M. C. Martín-Buro, B. P. Staresina, S. Hanslmayr, Disentangling neocortical alpha/beta and hippocampal theta/gamma oscillations in human episodic memory formation. *NeuroImage* 242, 118454 (2021).

46. R. T. Canolty, E. Edwards, S. S. Dalal, M. Soltani, S. S. Nagarajan, H. E. Kirsch, M. S. Berger, N. M. Barbaro, R. T. Knight, High Gamma Power Is Phase-Locked to Theta Oscillations in Human Neocortex. *Science* 313, 1626–1628 (2006).

47. L. Saint Amour di Chanaz, A. Pérez-Bellido, X. Wu, D. Lozano-Soldevilla, D. Pacheco-Estefan, K. Lehongre, E. Conde-Blanco, P. Roldan, C. Adam, V. Lambrecq, V. Frazzini, A. Donaire, M. Carreño, V. Navarro, A. Valero-Cabré, L. Fuentemilla, Gamma amplitude is coupled to opposed hippocampal theta-phase states during the encoding and retrieval of episodic memories in humans. *Current Biology* 33, 1836-1843.e6 (2023).

48. E. L. Newman, S. N. Gillet, J. R. Climer, M. E. Hasselmo, Cholinergic Blockade Reduces Theta-Gamma Phase Amplitude Coupling and Speed Modulation of Theta Frequency Consistent with Behavioral Effects on Encoding. J Neurosci 33, 19635–19646 (2013).

49. J. Daume, J. Kamiński, A. G. P. Schjetnan, Y. Salimpour, U. Khan, M. Kyzar, C. M. Reed, W. S. Anderson, T. A. Valiante, A. N. Mamelak, U. Rutishauser, Control of working memory by phase–amplitude coupling of human hippocampal neurons. *Nature* 629, 393–401 (2024).

50. F. Mormann, J. Fell, N. Axmacher, B. Weber, K. Lehnertz, C. E. Elger, G. Fernández, Phase/amplitude reset and theta–gamma interaction in the human medial temporal lobe during a continuous word recognition memory task. *Hippocampus* 15, 890–900 (2005).

51. A. Fernández-Ruiz, A. Oliva, G. A. Nagy, A. P. Maurer, A. Berényi, G. Buzsáki, Entorhinal-CA3 Dual-Input Control of Spike Timing in the Hippocampus by Theta-Gamma Coupling. *Neuron* 93, 1213-1226.e5 (2017).

52. A. B. L. Tort, R. W. Komorowski, J. R. Manns, N. J. Kopell, H. Eichenbaum, Theta-gamma coupling increases during the learning of item-context associations. *Proc Natl Acad Sci U S A* 106, 20942–20947 (2009).

53. N. Axmacher, M. M. Henseler, O. Jensen, I. Weinreich, C. E. Elger, J. Fell, Cross-frequency coupling supports multi-item working memory in the human hippocampus. *Proc Natl Acad Sci U S A* 107, 3228–3233 (2010).

54. B. P. Staresina, M. Wimber, A Neural Chronometry of Memory Recall. *Trends Cogn Sci* 23, 1071–1085 (2019).

55. D. Kobak, W. Brendel, C. Constantinidis, C. E. Feierstein, A. Kepecs, Z. F. Mainen, X.-L. Qi, R. Romo, N. Uchida, C. K. Machens, Demixed principal component analysis of neural population data. *Elife* 5, e10989 (2016).

56. G. Buzsáki, A. Draguhn, Neuronal oscillations in cortical networks. *Science* 304, 1926–1929 (2004).

57. M. Tamura, T. J. Spellman, A. M. Rosen, J. A. Gogos, J. A. Gordon, Hippocampal-prefrontal theta-gamma coupling during performance of a spatial working memory task. *Nat Commun* 8, 2182 (2017).

58. B. Lega, J. Burke, J. Jacobs, M. J. Kahana, Slow-Theta-to-Gamma Phase–Amplitude Coupling in Human Hippocampus Supports the Formation of New Episodic Memories. *Cereb. Cortex* 26, 268–278 (2016).

59. T. Donoghue, M. Haller, E. J. Peterson, P. Varma, P. Sebastian, R. Gao, T. Noto, A. H. Lara, J. D. Wallis, R. T. Knight, A. Shestyuk, B. Voytek, Parameterizing neural power spectra into periodic and aperiodic components. *Nat Neurosci* 23, 1655–1665 (2020).

60. R. F. Helfrich, J. D. Lendner, B. A. Mander, H. Guillen, M. Paff, L. Mnatsakanyan, S. Vadera, M. P. Walker, J. J. Lin, R. T. Knight, Bidirectional prefrontal-hippocampal dynamics organize information transfer during sleep in humans. *Nat Commun* 10, 3572 (2019).

61. B. P. Staresina, J. Niediek, V. Borger, R. Surges, F. Mormann, How coupled slow oscillations, spindles and ripples coordinate neuronal processing and communication during human sleep. *Nat Neurosci* 26, 1429–1437 (2023).

62. Z. Xiao, X. Wang, J. Zhang, J. Ou, L. He, Y. Qu, X. Hu, T. Behrens, Y. Liu, Human hippocampal ripples predict the alignment of experience to a grid-like schema. bioRxiv [Preprint] (2025). https://doi.org/10.1101/2025.01.08.632069.

63. R. Kaplan, M. H. Adhikari, R. Hindriks, D. Mantini, Y. Murayama, N. K. Logothetis, G. Deco, Hippocampal Sharp-Wave Ripples Influence Selective Activation of the Default Mode Network. *Curr Biol* 26, 686–691 (2016).

64. S. Michelmann, A. R. Price, B. Aubrey, C. K. Strauss, W. K. Doyle, D. Friedman, P. C. Dugan, O. Devinsky, S. Devore, A. Flinker, U. Hasson, K. A. Norman, Moment-by-moment tracking of naturalistic learning and its underlying hippocampo-cortical interactions. *Nat Commun* 12, 5394 (2021).

65. S. Michelmann, P. Dugan, W. Doyle, D. Friedman, L. Melloni, C. K. Strauss, S. Devore, A. Flinker, O. Devinsky, U. Hasson, K. A. Norman, Fast-timescale hippocampal processes bridge between slowly unfurling neocortical states during memory search. [Preprint] (2025). https://doi.org/10.1101/2025.02.11.637471.

66. D. Reznik, R. Trampel, N. Weiskopf, M. P. Witter, C. F. Doeller, Dissociating distinct cortical networks associated with subregions of the human medial temporal lobe using precision neuroimaging. *Neuron* 111, 2756-2772.e7 (2023).

67. D. Reznik, D. S. Margulies, M. P. Witter, C. F. Doeller, Evidence for convergence of distributed cortical processing in band-like functional zones in human entorhinal cortex. *Current Biology* 34, 5457-5469.e2 (2024).

68. M. Rigotti, O. Barak, M. R. Warden, X.-J. Wang, N. D. Daw, E. K. Miller, S. Fusi, The importance of mixed selectivity in complex cognitive tasks. *Nature* 497, 585–590 (2013).

69. A. C. Heusser, D. Poeppel, Y. Ezzyat, L. Davachi, Episodic sequence memory is supported by a theta-gamma phase code. Nat Neurosci 19, 1374–1380 (2016).

70. U. Vivekananda, D. Bush, J. A. Bisby, S. Baxendale, R. Rodionov, B. Diehl, F. A. Chowdhury, A. W. McEvoy, A. Miserocchi, M. C. Walker, N. Burgess, Theta power and theta-gamma coupling support long-term spatial memory retrieval. *Hippocampus* 31, 213–220 (2021).

71. N. Roehri, L. Bréchet, M. Seeber, A. Pascual-Leone, C. M. Michel, Phase-Amplitude Coupling and Phase Synchronization Between Medial Temporal, Frontal and

Posterior Brain Regions Support Episodic Autobiographical Memory Recall. *Brain Topogr* 35, 191–206 (2022).

72. L. L. Colgin, T. Denninger, M. Fyhn, T. Hafting, T. Bonnevie, O. Jensen, M.-B. Moser, E. I. Moser, Frequency of gamma oscillations routes flow of information in the hippocampus. *Nature* 462, 353–357 (2009).

73. B. J. Griffiths, G. Parish, F. Roux, S. Michelmann, M. van der Plas, L. D. Kolibius, R. Chelvarajah, D. T. Rollings, V. Sawlani, H. Hamer, S. Gollwitzer, G. Kreiselmeyer, B. Staresina, M. Wimber, S. Hanslmayr, Directional coupling of slow and fast hippocampal gamma with neocortical alpha/beta oscillations in human episodic memory. *Proc Natl Acad Sci U S A* 116, 21834–21842 (2019).

74. J. Sheng, L. Zhang, C. Liu, J. Liu, J. Feng, Y. Zhou, H. Hu, G. Xue, Higher-dimensional neural representations predict better episodic memory. Sci Adv 8, eabm3829 (2022).

75. G. Buzsáki, L. W. Leung, C. H. Vanderwolf, Cellular bases of hippocampal EEG in the behaving rat. *Brain Res* 287, 139–171 (1983).

76. G. Parish, S. Hanslmayr, H. Bowman, "The Sync/deSync model: How a synchronized hippocampus and a de-synchronized neocortex code memories" (preprint, Neuroscience, 2017); https://doi.org/10.1101/185231.

77. B. J. Griffiths, S. D. Mayhew, K. J. Mullinger, J. Jorge, I. Charest, M. Wimber, S. Hanslmayr, Alpha/beta power decreases track the fidelity of stimulus-specific information. *eLife* 8, e49562 (2019).

78. M. C. Martín-Buro, M. Wimber, R. N. Henson, B. P. Staresina, Alpha Rhythms Reveal When and Where Item and Associative Memories Are Retrieved. *J. Neurosci.* 40, 2510–2518 (2020).

79. N. A. Cayco-Gajic, C. Clopath, R. A. Silver, Sparse synaptic connectivity is required for decorrelation and pattern separation in feedforward networks. *Nat Commun* 8, 1116 (2017).

80. F. Lanore, N. A. Cayco-Gajic, H. Gurnani, D. Coyle, R. A. Silver, Cerebellar granule cell axons support high dimensional representations. *Nature neuroscience* 24, 1142 (2021).

81. C. Higgins, Y. Liu, D. Vidaurre, Z. Kurth-Nelson, R. Dolan, T. Behrens, M. Woolrich, Replay bursts in humans coincide with activation of the default mode and parietal alpha networks. *Neuron* 109, 882-893.e7 (2021).

82. S. Michelmann, B. P. Staresina, H. Bowman, S. Hanslmayr, Speed of time-compressed forward replay flexibly changes in human episodic memory. *Nat Hum Behav* 3, 143–154 (2019).