Chromosome-scale genome assembly of the European common cuttlefish *Sepia officinalis*

Simone Rencken[1,2] https://orcid.org/0009-0003-9341-2898, Georgi Tushev[1] https://orcid.org/0000-0002-3340-9422, David Hain[1,3] https://orcid.org/0000-0002-8979-7938, Elena Ciirdaeva[1] https://orcid.org/0009-0000-7360-7725, Oleg Simakov[4] https://orcid.org/0000-0002-3585-4511, Gilles Laurent[1] https://orcid.org/0000-0002-2296-114X

[1] Max Planck Institute for Brain Research, Frankfurt am Main, Germany
[2] Radboud University, Donders Institute for Brain, Cognition and Behaviour, Nijmegen, Netherlands
[3] Faculty of Biological Sciences, Goethe University, Frankfurt am Main, Germany
[4] Department of Neuroscience and Developmental Biology, University of Vienna, Vienna, Austria

Abstract

Coleoid cephalopods, a subclass of mollusks, exhibit remarkable adaptations, including the largest brains among invertebrates, camera-like eyes, and a distinctive embryonic development. They possess an advanced behavioral repertoire including dynamic camouflage.

The common cuttlefish *Sepia officinalis* has served as a key model organism in various research fields, spanning biophysics, neurobiology, behavior, evolution, ecology and biomechanics. More recently, it has become a model to investigate the neural mechanisms underlying cephalopod camouflage, using quantitative behavioral approaches alongside molecular techniques to characterize the identity, evolution and development of neuronal cell types.

Despite significant interest in this animal, a high-quality, annotated genome of its species is still lacking. To address this, we sequenced and assembled a chromosome-scale genome for *S. officinalis*. The final assembly spans 5.68 billion base pairs and comprises 47 repeat-rich chromosomes. Gene linkage analysis confirms the existence of 47 chromosomes, revealing clear homologies with related species such as *Euprymna scolopes* and *Doryteuthis pealeii*. Our work includes a comprehensive gene annotation and full-length transcript predictions that should be helpful for further evolutionary and single-cell expression studies. This genome provides a valuable resource for future research on the evolution, brain organization, information processing, development, and behavior in this important clade.

Introduction

Coleoid cephalopods (octopus, squid, cuttlefish) are a highly derived group of mollusks, characterized by the largest nervous systems among all invertebrates (ca. 500 million neurons in an adult octopus[1,2], ca. 200 million neurons in adult cuttlefish) and specializations with a great historical importance for neuroscience (e.g., "giant axons"[3] and "giant synapses"[4–6]). These animals exhibit very sophisticated behaviors such as dynamic camouflage[7–13], learning[14–16], social communication[17–19] and hunting[20–22] as well as two-stage sleep[23–25]. Because of these characteristics, cephalopods have been the focus of many fundamental studies by biologists, biophysicists and physiologists over the past century[26–35]. Thanks to advances in sequencing technologies, coleoid cephalopods are now also emerging as animal models in molecular neurobiology, evolution and genomics[36]. Recent studies in particular examined cephalopod biology from the perspectives of single-cell gene expression[37–40], genome topology and gene regulation[41].

Despite the technical progress, the genomes of cephalopods remain challenging to assemble because of their large sizes and high repeat fractions, creating a bottleneck for further molecular studies. While the genomes of *Octopodiformes* (Octopus, Eledone, Argonauta) are often comparable in size to that of humans (around 3 Gigabases (Gb)[42,43]) or possibly even smaller[44], the typical genomes of *Decapodiformes* (squids and cuttlefish) are often twice as large, reaching 6 Gb[45,46]. The biggest contributing factors to this genome expansion are transposable elements (TEs), with different TE classes differentiating squid and cuttlefish genomes from those of Octopodiformes[47]. Besides such differences in genome sizes, karyotypes in coleoid cephalopods are also unusual when compared to those of other invertebrates, including mollusks: for example, haploid chromosome numbers are around 30 in octopuses and around 46 in squids and cuttlefish[46,48], while they are between 8 and 19 in non-cephalopod mollusks[49–56]. Karyotype reconstruction and validation techniques in animals with such large genomes are also not as developed as they are for more familiar invertebrate and vertebrate species.

The common cuttlefish *Sepia officinalis* has been used as a model organism for studies in neural control and development[57,58], behavior[7,15], evolution[59], environmental studies[60,61] and biomechanics[62,63]. Found in the Mediterranean, the English Channel and the European Atlantic, it can produce large egg clutches containing hundreds of embryos, several times throughout its semelparous reproductive cycle[64,65]. More recently, this species has emerged as an important model to study the neural basis of cephalopod camouflage[66,67] building on

70 extensive behavioral studies[7–13] and neuronal cell type characterization, evolution and
71 development[68–70].
72

Despite a widespread interest in this animal, including the recent sequencing of several
cephalopod species as part of the Aquatic Symbiosis Genomics project[71], a high-quality,
annotated genomic resource for this species is still lacking. We describe here the
sequencing, assembly, and annotation of the *Sepia officinalis* genome. Our analysis
provides evidence for the existence of 47 chromosomes, with clear homologies to the
chromosomes of related species such as *Euprymna scolopes* and *Doryteuthis pealeii*. We
predict full-length transcripts and provide functional gene annotation that should inform
future evolutionary and transcriptomic studies. Taken together, this genome should be an
important resource for scientists interested in genome evolution, brain organization,
information processing, development and behavior.

83 Results

84 We sequenced the genome of a 6-month-old *Sepia officinalis* individual (probably male, see
85 below), reared from eggs collected in the Portuguese Atlantic. The DNA was extracted from
86 brain tissue and sequenced using long-read (PacBio HiFi) and chromatin conformation (Hi-
87 C) methods (Figure 1A,B).

88 Genome size and heterozygosity

89 The haploid genome size of *S. officinalis* was estimated to be ~5.14 Gb based on k-mer
90 estimation from the long-read data, with a high repeat content of 54 %[72,73]. The
91 heterozygosity rate was estimated to be 1.03 %, which is higher than in octopus genomes,
92 yet moderate among marine invertebrates[42,43,46].
93

94 The size of the scaffolded assembly was 5.68 Gb (Figure 1C), thus about 10% greater than
95 our initial GenomeScope estimate. Indeed, the sizes of most published metazoan genome
96 assemblies deviate by more than 10% from estimates, with proportionally greater deviations
97 as assembly sizes increase[74].
98

99 Completeness of the genome was assessed using BUSCO[75] with *metazoa_odb10*. It was
100 95.2% [single copy: 93.5%, duplicated: 1.7%], with 3.2% fragmented and 1.6% missing
101 BUSCOs. The BUSCO score with *mollusca_odb10* for the final assembly was 77.2%

completeness [single copy: 75.9%, duplicated: 1.3%] with 4.3% fragmented and 18.5% missing BUSCOs.

Karyotype

The assembly was generated from PacBio HiFi reads and Hi-C reads with ca. 23-fold and 11-fold coverage, respectively. Scaffolding was performed using Hi-C reads, placing 95.8 % of bases in 47 scaffolds. The Hi-C heatmap in Figure 1D shows the chromatin contacts in clusters corresponding to these 47 scaffolds. It is consistent with the proposal that they each represent a single chromosome.

To increase confidence in our assembled *Sepia officinalis* karyotype, we used several scaffolding programs and manually curated the scaffolds using two independent approaches. The first used HapHiC[76], a tool based on Hi-C data that utilizes allele information from primary assembly programs, and scaffolds assemblies with the constraint of an expected number of chromosomes. The resulting Hi-C contact maps with 46, 47, 48, 49 and 50 chromosomes (Supplementary Figure 1) show support for 47 chromosomes; an input of 46 scaffolds resulted in one clear chromosome merger (blue arrowhead); higher input numbers resulted in false chromosome splits (black arrowheads, 48, 49 and 50 scaffolds).

Second, we performed manual curation of the scaffolding result obtained with YAHS[77] using JBAT[78]. These two approaches, conducted independently and both based on Hi-C signal, converged on 47 chromosomal scaffolds.

Karyotype comparisons with other Decapods

Previous karyotyping studies in decapod cephalopods based on cytogenetic techniques provide widely varying estimates of chromosome numbers, reflecting the difficulty of resolving large chromosome numbers *in situ*. For example, 1n=46 chromosomes has been reported for two species of cuttlefish (*Sepia esculenta* and *Sepia lycidas*) and three loliginid squids[79]; 1n=34 chromosomes has been reported for *Sepia arabica*[80] and 1n=24 chromosomes in *Sepia pharaonis*[81]. In *Sepia officinalis*, a karyotype of 1n=52 has even been described from testis samples[82]. More recently, several cephalopod genomes have been scaffolded at chromosome-scale resolution, resolving karyotypes *in silico*. From those recent studies, a haploid karyotype of 46 chromosomes seems to be common among *Decapodiformes*[46] (Figure 2A).

Comparison with another chromosome-scale *Sepia officinalis* assembly

A chromosome-scale assembly for *Sepia officinalis* was released recently by the Wellcome Sanger Institute's Darwin Tree of Life project[83] (DToL, GCA_964300435.1). That genome was assembled from a male individual using PacBio Sequel II and Arima2 Hi-C data, with a final assembly size of 5.8 Gb. In that study, the haploid chromosome number was estimated to be 49. We scaffolded their assembly using our Dovetail Omni-C data, and the resulting contact map again supports 47 chromosomes (Supplementary Figure 2A), consistent with our results. We next aligned the two *S. officinalis* assemblies using minimap2[84] to assess the differences between them (Supplementary Figure 2B). The "Sanger" assembly appears to split three chromosomes of our assembly (chromosomes 1, 5 and 42) and to merge two (chromosomes 40 + 43). In conclusion, available evidence supports a haploid karyotype of 47 chromosomes in *Sepia officinalis*.

Comparisons with other chromosome-scale *Decapodiformes* genomes

To further investigate the possible reasons why our estimate of *Sepia officinalis's* chromosome numbers deviates from those in other studies, we compared our assembly with the chromosome-scale genomes of two other decapod cephalopods, *Euprymna scolopes*[41] (a *sepiolidae*) and *Doryteuthis pealeii*[46] (a *loliginidae*), both described as having 46 chromosomes. Using orthogroups to perform linkage analysis, we could detect a clear chromosome homology between *Doryteuthis* and *Sepia*, and to a lesser extent between *Euprymna* and *Sepia* (Figure 2B). Dotplots showing detailed pairwise synteny comparisons are shown in Supplementary Figure 3 (*Sepia* to *Doryteuthis*) and Supplementary Figure 4 (*Sepia* to *Euprymna*). The inferred species tree as part of this analysis places *Euprymna* as sister to *Sepia* and *Doryteuthis*, matching recent phylogenetic analyses[50,85]. However, this tree was constructed without the inclusion of outgroup taxa, and therefore lacks a reliable root.

This comparison revealed that chromosomes 40 and 43 in *Sepia* are merged into one in *Euprymna* and in *Dorytheutis* (Figure 2C), as they were in the DToL *Sepia officinalis* assembly (Supplementary Figure 2B). Our scaffolding approach showed no contact between chromosomes 40 and 43 in our *Sepia* Hi-C data, providing support for these chromosomes being separate in *Sepia officinalis*.

We also compared our assembly to a chromosome-scale *Sepia esculenta* assembly (1n=46, GCA_964036315.1), and recently annotated by genome liftover from *Sepia pharaonis*[86]. That assembly, constructed from a female individual, showed a low read coverage for a

putative sex chromosome in a ZZ/Z0 sex determination system[87]. Our syntenic comparison

indicated a strong homology between the putative Z chromosome of *Sepia esculenta* and

chromosome 46 of *Sepia officinalis* (Supplementary Figure 5A). Because we did not record

the sex of the *Sepia officinalis* animal used for genome sequencing and assembly, we tried

to infer the sex by replicating the analysis used to identify the putative Z chromosome in

Sepia esculenta[87]. For this, we aligned short-reads (Illumina) from the same *Sepia officinalis*

individual to the assembly and examined the normalized read coverage for each

chromosome (Supplementary Figure 5B). In contrast to the low coverage observed in the

female *Sepia esculenta* assembly (Supplementary Figure 5C), we observed no decrease in

read coverage for chromosome 46, suggesting that, if the alleged sex chromosome identity

is correct, then our material came from a male animal.

Genome repeat landscape

After creating a custom repeat library for *Sepia officinalis* using RepeatModeler[88], we

masked the genome using RepeatMasker[89], resulting in 71.17% masked bases. The

categories of repeats are shown in Figure 3A. Most repeats (39.65% of total bp) were not

characterized and presumably represent ancient repeats that diverged beyond recognition[90].

Retroelements constituted the largest characterized repeat category (17.32%) followed by

DNA transposons (5.92%); 5.83% were annotated as simple repeats. As observed in other

Decapodiformes, the *Sepia* genome contained almost no short interspersed nuclear

elements (SINEs), supporting the hypothesis that the SINE expansion observed in

octopuses[47] occurred independently in their lineage.

Gene modeling and annotation

The genome was further annotated using BRAKER[75,91–106] combining short- and long-read

RNA-seq data and publicly available protein data from multiple molluscan species

(*Doryteuthis pealeii*[46], *Euprymna scolopes*[107], *Octopus bimaculoides*[46], *Octopus vulgaris*[43],

Nautilus pompilius[108], and *Pecten maximus*[51]). A total of 18,663 gene models and 23,768

proteins were annotated.

The gene annotation was evaluated using BUSCO[75] v5.5.0 in protein mode, showing high

completeness of the annotation (metazoa_odb10: C:98.2%[S:77.3%,D:20.9%], F:1.0%,

M:0.8%, n:954 and mollusca_odb10: C:81.5%[S:59.6%,D:21.9%], F:0.9%, M:17.6%,

n:5295). We further checked the completeness and consistency of our gene models using

OMArk[109], and compared them to genome annotations for two other cuttlefish species, *Sepia*

pharaonis and *Sepia lycidas* (Figure 3B). The completeness is assessed by comparing the

annotated genes to conserved orthologs (present in 80% of extant species) in a given

taxonomic clade, here the superorder *Lophotrochozoa*. The *S. officinalis* annotation misses

181 out of 2373 genes, which is higher than *S. lycidas* (96 genes) but much lower than *S.*

pharaonis (458 genes).

In a consistency assessment, where the presence of known gene families from the lineage is

evaluated (Figure 3C), *S. officinalis* contains low proportions of taxonomically inconsistent or

fragmented proteins (ca. 13%) similar to *S. lycidas* (9%), giving confidence in the annotation.

In contrast, more than 50% of *S. pharaonis* proteins are labeled as inconsistent or

fragmented, which could indicate annotation errors[109].

Overall, the annotations contain different numbers of predicted proteins, likely caused by

differences in the annotation method and reference data used. *S. officinalis* contains the

least proteins (23,768) compared to *S. lycidas* (35,949) and *S. pharaonis* (53,515). The *S.*

officinalis annotation thus represents a conservative yet high-quality gene set. In the future,

this annotation could be enhanced by incorporating more long-read RNA sequencing data,

for example.

Lastly, we assigned orthology information to the *S. officinalis* proteome from InterPro[110] and

eggNOG-mapper[111] to aid the interpretability of the resource for future transcriptomic or

proteomic studies. Overall, 89% of proteins (21,204 out of 23,768) received an annotation

from InterPro from at least one of their databases. 59% of proteins (14,126 out of 23,768)

were annotated by eggNOG-mapper, reflecting the more stringent orthology filters and

prioritization of full-length matches implemented in the program[112]. Similar to previous

studies of other cephalopod genomes[42], we observed a high number of C2H2 zinc finger

transcription factors (1,538 proteins, InterPro annotation) and protocadherins (339 proteins,

InterPro), suggesting a conserved expansion of these gene families among cephalopods.

Further analyses should reveal more expanded gene families in *S. officinalis* that underlie

species-specific adaptations.

232 Discussion

This study presents a high-quality, chromosome-scale genome assembly and annotation for

the common cuttlefish *Sepia officinalis*, providing an additional important resource for

comparative genomics, neurobiology, and evolutionary studies within cephalopods and

mollusks. The assembly was derived from a combination of PacBio HiFi long reads, Dovetail

Omni-C chromatin conformation capture, and multiple rounds of scaffolding and manual
curation, resulting in a genome size of 5.68 Gb and a haploid karyotype of 47 chromosomes.
This resource represents a significant step forward for cephalopod genomics, especially
within the *Decapodiformes*, where large and repetitive genomes have hindered previous
efforts.

Chromosome Number and Karyotypic Variation

Our results strongly support a haploid chromosome number of 47 in *S. officinalis*, a
conclusion reinforced by two independent scaffolding approaches (YAHS[77] and HapHiC[76])
and supported by Hi-C contact maps. This karyotype contrasts with the 46 chromosomes
reported in other decapod cephalopods, such as *Euprymna scolopes* and *Doryteuthis*
pealeii[46]. Syntenic comparisons revealed that chromosomes 40 and 43 in *S. officinalis*
correspond to a single chromosome in these other species. The current Hi-C data for *Sepia*
officinalis provides clear evidence that chromosomes 40 and 43 are distinct, with no
interaction contact. Because these two chromosomes are merged also in *S. esculenta*, it is
plausible that this difference results from a *S. officinalis*-specific fission. However, our results
show how difficult it still is to assemble large genomes with high karyotype numbers.
Therefore, validation and comparisons with other *Sepia* species' genomes will be important.
This, in turn, will help identify chromosomal fusion-with-mixing events that are clade specific.
Early branching order in *Decapodiformes* has been notoriously unstable[50,85,113–116]; thus,
such rare and irreversible phylogenetic characters could be useful in further phylogenetic
analysis of this clade[48,117].

We also compared our results to the recently published *S. officinalis* genome from the
Darwin Tree of Life project (DToL)[83], which proposed 49 chromosomes. The discrepancy is
attributable to one merger in our assembly and three apparent chromosome splits in the
DToL assembly. While these differences could stem from technical choices such as different
Hi-C methods (Arima v2 vs. Dovetail Omni-C) and sequencing depth (83-fold vs. 11-fold
coverage), our assembly is strongly supported by Hi-C contact data, especially in regions
where the DToL assembly splits chromosomes that appear contiguous in our data.
Interestingly, the DToL assembly merges the chromosomes 40 and 43 - supporting a 46-
chromosome model as ancestral state. However, this is not supported by re-scaffolding with
the current Omni-C dataset, which again shows them as separate. This discrepancy might
reflect technical differences, but it could also be hinting at population-level variability in
chromosomal architecture; that is, some individuals may carry a fusion while others do not.
Intraspecific chromosome number variation is rare but not unprecedented; for instance,

chromosomal polymorphism has been described in the butterfly *Leptidea sinapis* across

different populations[118,119]. Notably, the two *S. officinalis* individuals used for sequencing

originated from different regions - the Portuguese Atlantic (this study) and the French

Mediterranean (DToL project), raising the possibility of geographic variation in chromosome

structure. Future population-level analyses will be critical to determine whether *S. officinalis*

exhibits karyotypic polymorphism.

Genome Size and Repeat Landscape

The size of the genome of *S. officinalis* is estimated at 5.68 Gb - comparable to other

Decapodiformes and roughly double the size of typical *Octopodiformes* genomes. Repeat

annotation revealed that over 71% of the genome is repetitive, dominated by unclassified

elements and retrotransposons. The near absence of SINEs reinforces the idea that their

expansion in octopuses was a lineage-specific event[47]. These results underscore the

complexity and evolutionary dynamism of the cephalopod genome, shaped by waves of

transposon activity that likely played a role in the evolution of novel traits[41,46].

Gene Annotation and Comparative Assessment

We annotated 18,663 gene models and 23,768 proteins using a combination of short- and

long-read RNA-seq data and molluscan protein references. Compared with other annotated

cuttlefish genomes (*S. lycidas*, *S. pharaonis*), our annotation is conservative in gene number

but stands out in terms of completeness and taxonomic consistency. OMArk-based

evaluations confirmed that our gene models have low levels of fragmentation and

contamination, and high lineage-specific coherence. Striking differences across species in

the number of predicted genes, duplications, and missing orthologs likely reflect variation in

the annotation pipelines and reference data rather than true biological divergence. Still, our

dataset provides a high-confidence foundation that can be refined with additional

transcriptomic data from long-read techniques and diverse tissues.

As in other coleoid genomes, we observed notable expansions of gene families such as

C2H2 zinc finger transcription factors and protocadherins, genes implicated in neural

development and plasticity[42,45]. These conserved expansions across cephalopods point to

genomic underpinnings of their cognitive complexity and behavioral sophistication.

Conclusions and Outlook

This study delivers a foundational genomic resource for *Sepia officinalis*, a model for studying neural control, behavior, and adaptive camouflage. The chromosome-scale assembly and annotation open the door to in-depth studies of gene regulation, neural circuit development, and genome evolution across coleoid cephalopods. Moreover, the identification of possible population-level karyotype variation highlights the importance of incorporating broader sampling strategies in future genomic studies.

As we move toward a more complete picture of cephalopod genome evolution, the integration of chromosomal synteny, regulatory architecture, and transcriptomic diversity across species will be especially important. The *S. officinalis* genome represents an important step on this path, enabling high-resolution comparative and functional analyses of one of the most enigmatic and evolutionarily successful invertebrate lineages.

Materials & Methods

Animal husbandry

All research and animal care procedures were carried out following the institutional guidelines that comply with national and international laws and policies (DIRECTIVE 2010/63/EU; German animal welfare act; FELASA guidelines). European cuttlefish *Sepia officinalis* were hatched from eggs collected in the Portuguese Atlantic and reared in a seawater system, at 20°C. The closed system contains 4,000l of artificial seawater (ASW; Instant Ocean) with a salinity of 33‰ and pH of 8–8.5. Water quality was tested daily and adjusted as required. Trace elements and amino acids were supplied weekly. Marine LED lights above each tank provided a 12/12-h light/dark cycle with gradual on- and off-sets at 07:00 and 19:00. The animals were fed live food (either *Hemimysis* spp. or small *Palaemonetes* spp.) *ad libitum* twice per day. The animals were housed together in 120-l glass tanks with a constant water through-flow resulting in five complete water exchanges per hour. Enrichment consisted of natural fine-grained sand substrate, seaweed (*Caulerpa prolifera*), rocks of different sizes and various natural and man-made three-dimensional objects.

Tissue preparation

Animals underwent terminal anesthesia to prevent animal suffering or distress, following the Guidelines for the Care and Welfare of Cephalopods in Research[120,121]. Animals were

transferred to a bucket containing ethanol 2% (v/v) in ASW. After 5 minutes, animals were
gently probed for simple avoidance reflexes, and the ethanol concentration in the ASW was
gradually increased to a maximum of 5%. Sufficient depth of anesthesia was reached when
the animal no longer reacted even to stronger stimuli (touching and pinching with tweezers),
and no reaction was observed when a hand was moved in the visual area and the cornea
was touched[122]. In this deep anesthesia, the animals were decapitated and the tissue was
rapidly transferred into ice-cold, calcium-free Ringer solution (460 mM NaCl, 10 mM KCl, 51
mM $MgCl_2$, 10 mM glucose, 2 mM Glutamine, 10 mM HEPES, pH 7.4) bubbled with oxygen.
Brain and body tissue was dissected under a stereoscope and flash-frozen in liquid nitrogen
before storage at -80ºC until further use.
341
342   Unless stated otherwise, sequencing libraries were prepared from the same individual (6-
343   month-old adult *Sepia officinalis*, F1 from eggs collected in Portugal).

Long-read Whole Genome Library Preparation and Sequencing

The long-read library was prepared and sequenced at the MPI for Plant Research in
Cologne, Germany. Genomic DNA was extracted from flash-frozen brain tissue with the
MagAttract HMW DNA kit (Qiagen, Cat. no. 67563) and the sequencing library was prepared
with the SMRTbell express template prep kit 2.0 (PacBio, Cat. no. 100-938-900). The library
was sequenced on 5 SMRT cells on the PacBio Sequel II with the Sequel II binding kit 2.2
(102-089-000), Sequel II sequencing kit 2.0 (101-820-200) and SMRT Cell 8M tray (101-
389-001).

Long-read RNA library preparation and sequencing

RNA was isolated from 5-10 mg of flash-frozen tissue (brain, mantle/epidermis,
arm/tentacle). First, tissue samples were homogenized in TRIzol using a homogenizer
(VDI12/S12N-5S, VWR, Germany). RNA was isolated and DNAseI treated according to the
Direct-zol RNA Mini prep kit from ZymoResearch (R2050). RNA was quantified by
spectrophotometry (Nanodrop, Thermo Scientific, USA) and its quality was assessed with a
RNA 6000 Nanochip on Bioanalyzer (Agilent Technologies, Germany).
The Iso-Seq libraries were prepared and sequenced at the Sequencing facility of the MPI for
Plant Research, using a method targeting the 5' cap and poly-A tail of mRNAs, described in
detail in[123]. Briefly, mRNA was pooled from the 8 tissue samples and cDNA was synthesized
using the TeloPrime Full-Length cDNA Amplification Kit V2 (Lexogen, Cat Nr: 013.08 or
013.24). The sequencing library was prepared with the SMRTbell express template prep kit
2.0 (PacBio, Cat. no. 100-938-900). The pooled library was sequenced on 1 SMRT cell on

the PacBio Sequel II system using the Sequel II binding kit 2.1 (101-843-000), Sequel II

sequencing kit 2.0 (101-820-200) and SMRT Cell 8M tray (101-389-001).

Further, a separate library was prepared from optic lobe cDNA and sequenced on a second

SMRT cell using the same reagents described above.

Omni-C Library Preparation and Sequencing

An Omni-C library was prepared and sequenced at the MPI for Plant Research in Cologne,

Germany. The library was prepared from brain tissue using the DovetailTM Omni-C™ Kit

(Dovetail Genomics, Cat. No. 21005) according to the manufacturer's protocol. Briefly,

chromatin was fixed in place in the nucleus. Fixed chromatin was digested with DNase I then

extracted. Chromatin ends were repaired and ligated to a biotinylated bridge adapter

followed by proximity ligation of adapter-containing ends. After proximity ligation, crosslinks

were reversed and the DNA was purified from proteins. Purified DNA was treated to remove

biotin that was not internal to ligated fragments. The sequencing library was generated using

Illumina-compatible adapters. Biotin-containing fragments were isolated using streptavidin

beads before PCR enrichment of the library. The library was sequenced on an Illumina

NextSeq 2000 platform to generate 400 million 2 x 150 bp read pairs.

Short-read DNA sequencing

High molecular weight genomic DNA was isolated from brain tissue (optic lobe) using NEB

Monarch gDNA Purification Kit (T3010S). First, 10 mg of flash-frozen tissue were submerged

in 100 µl of tissue lysis buffer, cut into small pieces and transferred into a 1.5 ml tube

containing an additional 100 µl of lysis buffer. Then, 3 µl of 20 U/µl Proteinase K was added

to the tissue sample and incubated for 2 h at 56°C in a thermal mixer with agitation until

tissue pieces were completely dissolved. After incubation, tissue lysate was purified from the

debris by centrifugation for 3 min at maximal speed (Eppendorf, Germany). Clean

supernatant was transferred to a new 1.5 ml tube and incubated with 3 µl RNase A for 5 min

at 56°C with agitation at full speed. After RNase A incubation tissue lysate was column

purified and eluted with 80 µl of elution buffer according to the manufacturer protocol. The

purity of gDNA was assessed using A260/280 and A260/230 absorbance ratios measured

by NanoDrop spectrophotometer (ThermoFisher Scientific). The integrity of the obtained

gDNA was checked using 0,75% agarose gel electrophoresis by loading 100 ng of gDNA

sample and using Hind III digested Lambda DNA (NEB#3012) as a marker.

DNA sequencing libraries were prepared from 500 ng of high MW gDNA using Illumina DNA

PCR-Free Tagmentation Library Prep Kit (20041794) and UD Indexes Set A (20026121).

Obtained dual-indexed single-stranded libraries were quantified using the Qubit single-

stranded DNA (ssDNA) assay kit (Q10212, ThermoFisher Scientific). 1000 pM of the final

library were run on P3 reagents and Illumina NextSeq2000 using Illumina DNA PCR-Free

Sequencing and Indexing primer (20041797) with 151 bp paired-end reads.

Short-read RNA library preparation and sequencing

For short-read RNA sequencing, tissue from another animal (8-month-old adult, F0 from

eggs collected in Normandie, France) was used. RNA was isolated from 5 mg of flash-frozen

tissues (brain, skin and retina). First, tissue samples were homogenized in TRIzol using a

homogenizer (VDI12/S12N-5S, VWR, Germany) following DNAseI treatment and column

purification according to the Direct-zol RNA Micro prep kit from ZymoResearch (R2062). The

RNA integrity and quantity were measured with the Qubit fluorometer (Invitrogen, Q33216)

and the 2100 Bioanalyzer (Agilent Technologies, Germany).

RNA-seq libraries were prepared from 300 ng of total RNA, using the Illumina TruSeq

stranded mRNA library prep kit (Cat:20020594) and IDT for Illumina xGen UDI-UMI Adapters

(Cat:10005903). Libraries were sequenced on an Illumina NextSeq500 (Mid output, 300 cyc)

and NextSeq2000 (P3, 300 cyc) with 145 bp paired-end reads.

Nuclear genome assembly

Before assembly, PacBio HiFi reads were subjected to additional trimming to remove

residual adapter sequences using NCBI's VecScreen tool. Genome size was estimated

using GenomeScope[72] from Illumina short reads and PacBio HiFi data. The primary

assembly was generated with hifiasm[124] using a combination of HiFi and Hi-C reads.

Scaffolding was performed iteratively with YAHS[77] on the phased haplotype 1. We adjusted

the run parameters for scaffolding resolutions (-r

1000,2000,5000,10000,20000,30000,50000,70000,100000,150000,200000,250000,300000,

350000,400000,500000,700000,1000000,2000000,5000000,10000000,15000000,20000000

,30000000,40000000,50000000,60000000,70000000,80000000,100000000,110000000,120

000000,150000000,170000000,200000000,500000000), repetitions per resolution (-R3),

minimum read mapping quality (-q1) and telomeric sequence (--telo-motif "TTAGGG") from

the only experimentally determined telomere motif for cephalopods (*Amphioctopus*

areolatus[125], accessed via TeloBase[126]) to optimize the tool's performance for a large,

fragmented assembly. Finally, the assembly was manually curated using JBAT[78] to place

residual scaffolds into chromosome scaffolds. Different versions of the assembly were

evaluated based on HiC coverage, mRNA alignment and completeness based on BUSCO[75]

v5.5.0 (*metazoa_odb10* and *mollusca_odb10*).

Mitochondrial genome assembly

To assemble the mitochondrial genome, we aligned a published *S. officinalis* mitochondrial genome[127] (NC_007895.1) to the PacBio Hi-Fi and Omni-C reads using minimap2[84]. The hits were extracted using seqtk[128] subseq and assembled using hifiasm[124]. The resulting contigs contained multiple repeats of the circular mitochondrial genome, and were aligned back to NC_007895.1 to extract the final sequence.

Nuclear genome annotation

Repetitive elements in the genome were softmasked using RepeatMasker v4.1.7-p1[89] (with rmblast v2.14.1+) after creating a custom repeat library with RepeatModeler v.2.0.6[88].

Gene models were created using BRAKER3[75,91–106], run via the Docker container, on the softmasked genome using both RNA-seq and protein data. We used long-read Iso-Seq data and short-read Illumina RNA-seq data generated for this study (see above). For protein data, publicly available peptide annotations for *Doryteuthis pealeii*[46], *Euprymna scolopes*[107], *Octopus bimaculoides*[46], *Octopus vulgaris*[43], *Nautilus pompilius*[108], *Pecten maximus*[51] were used for training, as well as a previously generated proteome for *Sepia officinalis* from StringTie gene models (this study). RNA-seq data was input into BRAKER3 using the —bam option, protein files were specified with the–prot_seq option. The BUSCO completeness of the resulting gene set was maximized using TSEBRA within BRAKER on the BUSCO lineage *metazoa_odb10*.

For the initial protein set used as input for BRAKER, the short and long RNA reads were aligned to the genome using *minimap2*[84]. Transcript models were predicted with *StringTie* v3.0.0[129] using the *--conservative* and *--mix* options. The resulting GTF files were combined using the transcript merge mode, resulting in a set of non-redundant transcripts. Finally, TransDecoder[130] v5.7.0 was run with default parameters to translate coding regions in the transcripts.

The final gene annotations for *S. officinalis* were assessed for completeness using OMArk v.0.3.0[109] with the ancestral clade *Lophotrochozoa* and without including splice information, accessed via their webserver. Proteins were annotated with orthology information using InterProScan[110] v5.73-104 including lookup of annotations and GO terms (options -iprlookup -goterms). Further orthology information was added using the eggNOG-mapper v2.1.12[111] webserver with the eggNOG v5.0 database and default parameters.

Whole genome alignment and synteny analysis

For whole genome alignments, the assembly produced for *Sepia officinalis* in this study and the published assembly from the Darwin Tree of Life project (GCA_964300435.1) were aligned using *minimap2*[84] and visualized with a custom script in R v4.4.2[131].

For synteny analyses, peptide and gtf files were downloaded for *Doryteuthis pealeii* and *Euprymna scolopes.* Annotation files for *Sepia esculenta* were recently generated by liftover annotation from *Sepia pharaonis*[87] and downloaded from Zenodo[86]. Synteny analyses between all chromosomes of the compared species were performed using the R package GENESPACE v.1.2.3[132]. *S. officinalis* was used as the reference species for all synteny maps.

Coverage analysis

Read coverage across chromosomes was analyzed as described recently[87]. Short reads were aligned using STAR v2.7.11b[133] to our chromosome-scale assembly. For *Sepia esculenta*, short reads (ERR12945500) and assembly (GCA_964036315.1) were downloaded from NCBI. The sequencing depth was calculated using mosdepth[134] by a window size of 500,000 bp and normalized to the median coverage of the first chromosome.

Data availability

The genome assembly and raw sequencing reads can be found at the BioProject PRJNA1091451 on NCBI. The code for the genome assembly and annotation is available at https://gitlab.mpcdf.mpg.de/mpibr/laur/cuttlefishomics/soffgenome. Genome annotation files are deposited at https://public.brain.mpg.de/Laurent/sepoff/annotation/.

Author contributions

O.S., and G.L. conceived of the study design. S.R., O.S. and G.L. wrote the first draft of the manuscript. D.H. collected tissue for sequencing. E.C. extracted DNA and RNA from tissue and performed Illumina sequencing. G.T. performed genome assembly. G.T. and S.R. manually curated and annotated the genome. S.R., G.T., and O.S. performed genomic analyses. S.R. created the figures. S.R., O.S. and G.L. coordinated and led the project. G.L. acquired funding for this study.

496   All authors contributed to the review and editing of the manuscript.

Acknowledgments

We thank Bruno Huettel for PacBio and Omni-C sequencing. We thank Xitong Liang,

Theodosia Woo and Mathieu Renard for help with tissue dissection. We thank Darrin

Schultz, Dalila Destanović and Thea Rogers for helpful discussions and advice on manual

curation and gene modeling.

This work was funded by the Max Planck Society (GL) and the European Research Council

(GL; ERC grant CAMOUFLAGE, 101141501). O.S. was supported by the ERC's Horizon

2020: European Union Research and Innovation Programme, grant No. 945026.

All authors declare no conflict of interest. Views and opinions expressed are those of the

authors only and do not necessarily reflect those of the European Union or the European

Research Council Executive Agency. Neither the European Union nor the granting authority

can be held responsible for them.

Figures

Figure 1: *Sepia officinalis* assembly statistics and quality control. A) Specimen of *S.*
officinalis (credit: Stephan Junek, MPI Brain Research). **B)** Overview of the assembly
workflow. Genome size was estimated using GenomeScope[72,73]. The primary assembly was
generated from long DNA reads (PacBio Sequel II) and HiC reads (Dovetail OmniC) with
hifiasm[124]. Assembly was scaffolded with YAHS[77]. **C)** Snail plot of chromosome-scale *S.*
officinalis assembly generated using blobtools2[135] showing scaffold statistics, base
composition and BUSCO[75] completeness against the metazoa_odb10 database. **D)** HiC
heatmap showing 47 chromosome-scale scaffolds with very few sequences in unplaced
scaffolds. Heatmap was generated using juicebox[78], 0-7039 observed counts (balanced) are
shown.

Figure 2: Syntenic comparison of three decapod species. A) Taxonomy of selected
cephalopod species showing their genome size and (haploid) chromosome numbers.
Taxonomy information downloaded from NCBI taxonomy browser, divergence times
from[85,136]. **B)** Synteny relationship between chromosomes of *E. scolopes*[46] (top), *D. pealeii*[46]
(middle) and *S. officinalis* (bottom). Riparian plot was generated using orthogroups located
on chromosomes of the species. *Euprymna* chromosomes 45 and 46 are not shown
because they contain too few orthogroups. **C)** Synteny plot of *Sepia* chromosomes 40
(magenta) and 43 (darkblue), that are joined in the other species and cause the different
haploid chromosome number in *Sepia*. Species icons are from biorender.com.

Figure 3: Genome annotation for *Sepia officinalis*. A) Annotation of repeat landscape of
the *S. officinalis* genome, annotated using RepeatModeler[88]. Full repeat landscape is shown
on the left, annotated repeats (excluding unclassified or simple repeats) are shown on the
right. **B-C)** Quality control of gene annotation and comparison to two other cuttlefish species
using OMArk[109]. Results shown for *Sepia lycidas* (GCA_963932145.1, Ensembl Genebuild),
Sepia officinalis (BRAKER, this study) and *Sepia pharaonis*[137] (BRAKER). *Lophotrochozoa*
was used as the ancestral clade. **B)** Completeness assessed by the presence of genes
conserved in the clade, classified as *single* or multiple copies (*duplicated*), or *missing*. **C)**
Consistency assessed by the proportion of proteins placed in the correct lineage
(*consistent*); placement in incorrect lineages randomly (*inconsistent*) or to specific species
(*contamination*), or no placement in known gene families (*unknown*).

Supplementary Figure 1: HapHiC scaffolding for different numbers of expected chromosomes show 47 chromosomes as most supported. Hi-C contact maps from HapHiC[76] are shown for 46, 47, 48, 49 and 50 expected chromosome scaffolds. Assembled chromosomes are shown as blue boxes, HiC signal indicating a false (unsupported) merger is shown by cyan arrow, false splits are shown by black arrows.

Supplementary Figure 2: Comparison of *Sepia officinalis* chromosome-scale
assemblies. A) HiC heatmap of GCA_964300435.1 scaffolded against Omni-C data
collected from a different individual (this study). Assembled chromosomes are shown as blue
boxes, HiC signal indicating a false merge is shown by cyan arrow, false splits are shown by
black arrows. **B)** Whole-genome alignment of both *S. officinalis* assemblies (this study: rows,
GCA_964300435.1: columns).

Supplementary Figure 3: Syntenic relationship between *S. officinalis* and *D. pealeii* chromosomes. Dot plot showing finer-resolution syntenic anchor hits (perfectly collinear blast hits within the same orthogroup). Genes are ordered along the chromosomes, gridlines are shown every 1000 genes.

Supplementary Figure 4: Syntenic relationship between *S. officinalis* and *E. scolopes* chromosomes. Dot plot showing finer-resolution syntenic anchor hits (perfectly collinear blast hits within the same orthogroup). Genes are ordered along the chromosomes, gridlines are shown every 1000 genes. *E. scolopes* chromosomes 45 and 46 are not shown because they contain too few orthogroups.

**Supplementary Figure 5: Syntenic comparison of four decapod species hints at a
cephalopod sex chromosome. A)** Riparian plot showing synteny relationships of
chromosomes from four decapod species, generated using orthogroups. *Euprymna*
chromosomes 45 and 46 are not shown because they contain too few orthogroups.
Chromosome split in *S. officinalis* compared to other species is shown in purple, putative sex
chromosome as identified recently[87] is shown in cyan. **B)** Normalized coverage of
sequencing data in *S. officinalis* chromosomes. **C)** Normalized coverage of short reads to
female *S. esculenta* genome, reproduced from[87]. Decrease in read coverage for
chromosome 46 visible, the putative Z sex chromosome. Read depth was calculated from
Illumina gDNA reads in windows of 500,000 bp and normalized to the median coverage of
chromosome 1. Box plots showing median divergence (box dividing line), interquartile range
(box), and 1.5 times the interquartile range (whiskers). Putative Z chromosome is highlighted
in cyan.

References

1. Giuditta, A., Libonati, M., Packard, A., and Prozzo, N. (1971). Nuclear counts in the brain lobes of octopus vulgaris as a function of body size. Brain Res. *25*, 55–62.

2. Young, J. (1963). The number and sizes of nerve cells in Octopus. In Proceedings of the Zoological Society of London (Wiley Online Library), pp. 229–254.

3. Hodgkin, A.L., Huxley, A.F., and Katz, B. (1952). Measurement of current-voltage relations in the membrane of the giant axon of Loligo. J. Physiol. *116*, 424–448. https://doi.org/10.1113/jphysiol.1952.sp004716.

4. Llinás, R., Steinberg, I.Z., and Walton, K. (1981). Presynaptic calcium currents in squid giant synapse. Biophys. J. *33*, 289–321. https://doi.org/10.1016/S0006-3495(81)84898-9.

5. Llinás, R., Steinberg, I.Z., and Walton, K. (1981). Relationship between presynaptic calcium current and postsynaptic potential in squid giant synapse. Biophys. J. *33*, 323–351. https://doi.org/10.1016/S0006-3495(81)84899-0.

6. Llinas, R.R. (1984). The Squid Giant Synapse. In Current Topics in Membranes and Transport The Squid Axon., A. Kleinzeller, ed. (Academic Press), pp. 519–546. https://doi.org/10.1016/S0070-2161(08)60483-9.

7. Hanlon, R.T., and Messenger, J.B. (2018). Cephalopod Behaviour 2nd ed. (Cambridge University Press) https://doi.org/10.1017/9780511843600.

8. Josef, N., Berenshtein, I., Rousseau, M., Scata, G., Fiorito, G., and Shashar, N. (2017). Size Matters: Observed and Modeled Camouflage Response of European Cuttlefish (Sepia officinalis) to Different Substrate Patch Sizes during Movement. Front. Physiol. *7*.

9. Osorio, D., Ménager, F., Tyler, C.W., and Darmaillacq, A.-S. (2022). Multi-level control of adaptive camouflage by European cuttlefish. Curr. Biol. *32*, 2556-2562.e2. https://doi.org/10.1016/j.cub.2022.04.030.

10. Marshall, N.J., and Messenger, J.B. (1996). Colour-blind camouflage. Nature *382*, 408–409. https://doi.org/10.1038/382408b0.

11. Zylinski, S., Osorio, D., and Shohet, A. j (2009). Perception of edges and visual texture in the camouflage of the common cuttlefish, Sepia officinalis. Philos. Trans. R. Soc. B

Biol. Sci. *364*, 439–448. https://doi.org/10.1098/rstb.2008.0264.

12. Zylinski, S., Darmaillacq, A.-S., and Shashar, N. (2012). Visual interpolation for contour
completion by the European cuttlefish (Sepia officinalis) and its use in dynamic
camouflage. Proc. R. Soc. B Biol. Sci. *279*, 2386–2390.
https://doi.org/10.1098/rspb.2012.0026.

13. Zylinski, S., Osorio, D., and Shohet, A.J. (2009). Cuttlefish camouflage: context-
dependent body pattern use during motion. Proc. R. Soc. B Biol. Sci. *276*, 3963–3969.
https://doi.org/10.1098/rspb.2009.1083.

14. Jozet-Alves, C., Bertin, M., and Clayton, N.S. (2013). Evidence of episodic-like memory
in cuttlefish. Curr. Biol. *23*, R1033–R1035. https://doi.org/10.1016/j.cub.2013.10.021.

15. Schnell, A.K., Boeckle, M., Rivera, M., Clayton, N.S., and Hanlon, R.T. (2021). Cuttlefish
exert self-control in a delay of gratification task. Proc. R. Soc. B Biol. Sci. *288*,
20203161. https://doi.org/10.1098/rspb.2020.3161.

16. Boycott, B.B., and Young, J.Z. (1955). A memory system in Octopus vulgaris Lamarck.
Proc. R. Soc. Lond. Ser. B - Biol. Sci. *143*, 449–480.
https://doi.org/10.1098/rspb.1955.0024.

17. Mather, J.A. (2006). Behaviour Development: A Cephalopod Perspective. Int. J. Comp.
Psychol. *19*. https://doi.org/10.46867/ijcp.2006.19.01.02.

18. Hall, K., and Hanlon, R. (2002). Principal features of the mating system of a large
spawning aggregation of the giant Australian cuttlefish Sepia apama (Mollusca:
Cephalopoda). Mar. Biol. *140*, 533–545.

19. Norman, M.D., Finn, J., and Tregenza, T. (1999). Female impersonation as an
alternative reproductive strategy in giant cuttlefish. Proc. R. Soc. Lond. B Biol. Sci. *266*,
1347–1349. https://doi.org/10.1098/rspb.1999.0786.

20. Sampaio, E., Seco, M.C., Rosa, R., and Gingins, S. (2021). Octopuses punch fishes
during collaborative interspecific hunting events. Ecology *102*, e03266.
https://doi.org/10.1002/ecy.3266.

21. Sampaio, E., Sridhar, V.H., Francisco, F.A., Nagy, M., Sacchi, A., Strandburg-Peshkin,
A., Nührenberg, P., Rosa, R., Couzin, I.D., and Gingins, S. (2024). Multidimensional
social influence drives leadership and composition-dependent success in octopus–fish

hunting groups. Nat. Ecol. Evol. *8*, 2072–2084. https://doi.org/10.1038/s41559-024-

02525-2.

22. Feord, R.C., Sumner, M.E., Pusdekar, S., Kalra, L., Gonzalez-Bellido, P.T., and Wardill,

T.J. (2020). Cuttlefish use stereopsis to strike at prey. Sci. Adv. *6*, eaay6036.

https://doi.org/10.1126/sciadv.aay6036.

23. Medeiros, S.L. de S., Paiva, M.M.M. de, Lopes, P.H., Blanco, W., Lima, F.D. de,

Oliveira, J.B.C. de, Medeiros, I.G., Sequerra, E.B., de Souza, S., Leite, T.S., et al.

(2021). Cyclic alternation of quiet and active sleep states in the octopus. iScience *24*,

102223. https://doi.org/10.1016/j.isci.2021.102223.

24. Iglesias, T.L., Boal, J.G., Frank, M.G., Zeil, J., and Hanlon, R.T. (2019). Cyclic nature of

the REM sleep-like state in the cuttlefish Sepia officinalis. J. Exp. Biol. *222*, jeb174862.

https://doi.org/10.1242/jeb.174862.

25. Pophale, A., Shimizu, K., Mano, T., Iglesias, T.L., Martin, K., Hiroi, M., Asada, K.,

Andaluz, P.G., Van Dinh, T.T., Meshulam, L., et al. (2023). Wake-like skin patterning

and neural activity during octopus sleep. Nature *619*, 129–134.

https://doi.org/10.1038/s41586-023-06203-4.

26. Young, J.Z. (1974). The central nervous system of Loligo I. The optic lobe. Philos. Trans.

R. Soc. Lond. B Biol. Sci. *267*, 263–302.

27. Young, J.Z. (1962). The optic lobes of Octopus vulgaris. Philos. Trans. R. Soc. Lond. B.

Biol. Sci. *245*, 19–58. https://doi.org/10.1098/rstb.1962.0005.

28. Young, J. (1960). The Visual System of Octopus:(1) Regularities in the Retina and Optic

Lobes of Octopus in Relation to Form Discrimination. Nature *186*, 836–839.

29. Young, J.Z. (1991). Computation in the Learning System of Cephalopods. Biol. Bull. *180*,

200–208. https://doi.org/10.2307/1542389.

30. Young, J.Z. (1963). Light-and Dark-Adaptation in the Eyes of Some Cephalopods. Proc.

Zool. Soc. Lond. *140*, 255–272. https://doi.org/10.1111/j.1469-7998.1963.tb01863.x.

31. Young, J.Z. (1962). The retina of cephalopods and its degeneration after optic nerve

section. Philos. Trans. R. Soc. Lond. B. Biol. Sci. *245*, 1–18.

32. Nixon, M., and Young, J.Z. (2003). The brains and lives of cephalopods (Oxford

University Press).

33. Case, N.M., Gray, E., and Young, J. (1972). Ultrastructure and synaptic relations in the
optic lobe of the brain of Eledone and Octopus. J. Ultrastruct. Res. *39*, 115–123.

34. Dilly, P., Gray, E.G., and Young, J.Z. (1963). Electron microscopy of optic nerves and
optic lobes of Octopus and Eledone. Proc. R. Soc. Lond. B Biol. Sci. *158*, 446–456.

35. Stephens, P., and Young, J.Z. (1969). The glio-vascular system of cephalopods. Philos.
Trans. R. Soc. Lond. B. Biol. Sci. *255*, 1–12.

36. Baden, T., Briseño, J., Coffing, G., Cohen-Bodénès, S., Courtney, A., Dickerson, D.,
Dölen, G., Fiorito, G., Gestal, C., Gustafson, T., et al. (2023). Cephalopod-omics:
Emerging Fields and Technologies in Cephalopod Biology. Integr. Comp. Biol. *63*, 1226–
1239. https://doi.org/10.1093/icb/icad087.

37. Styfhals, R., Zolotarov, G., Hulselmans, G., Spanier, K.I., Poovathingal, S., Elagoz, A.M.,
De Winter, S., Deryckere, A., Rajewsky, N., Ponte, G., et al. (2022). Cell type diversity in
a developing octopus brain. Nat. Commun. *13*, 7392. https://doi.org/10.1038/s41467-
022-35198-1.

38. Songco-Casey, J.O., Coffing, G.C., Piscopo, D.M., Pungor, J.R., Kern, A.D., Miller, A.C.,
and Niell, C.M. (2022). Cell types and molecular architecture of the Octopus
bimaculoides visual system. Curr. Biol. *0*. https://doi.org/10.1016/j.cub.2022.10.015.

39. Duruz, J., Sprecher, M., Kaldun, J.C., Al-Soudy, A.-S., Lischer, H.E., van Geest, G.,
Nicholson, P., Bruggmann, R., and Sprecher, S.G. (2023). Molecular characterization of
cell types in the squid Loligo vulgaris. eLife *12*, e80670.
https://doi.org/10.7554/eLife.80670.

40. Gavriouchkina, D., Tan, Y., Ziadi-Künzli, F., Hasegawa, Y., Piovani, L., Zhang, L.,
Sugimoto, C., Luscombe, N., Marlétaz, F., and Rokhsar, D.S. (2022). A single-cell atlas
of bobtail squid visual and nervous system highlights molecular principles of convergent
evolution. bioRxiv, 2022.05.26.490366. https://doi.org/10.1101/2022.05.26.490366.

41. Schmidbaur, H., Kawaguchi, A., Clarence, T., Fu, X., Hoang, O.P., Zimmermann, B.,
Ritschard, E.A., Weissenbacher, A., Foster, J.S., Nyholm, S.V., et al. (2022). Emergence
of novel cephalopod gene regulation and expression through large-scale genome
reorganization. Nat. Commun. *13*, 2172. https://doi.org/10.1038/s41467-022-29694-7.

42. Albertin, C.B., Simakov, O., Mitros, T., Wang, Z.Y., Pungor, J.R., Edsinger-Gonzales, E.,
Brenner, S., Ragsdale, C.W., and Rokhsar, D.S. (2015). The octopus genome and the

evolution of cephalopod neural and morphological novelties. Nature *524*, 220–224. https://doi.org/10.1038/nature14668.

43. Destanović, D., Schultz, D.T., Styfhals, R., Cruz, F., Gómez-Garrido, J., Gut, M., Gut, I., Fiorito, G., Simakov, O., Alioto, T.S., et al. (2023). A chromosome-level reference genome for the common octopus, Octopus vulgaris (Cuvier, 1797). G3 GenesGenomesGenetics, jkad220. https://doi.org/10.1093/g3journal/jkad220.

44. Yoshida, M., Hirota, K., Imoto, J., Okuno, M., Tanaka, H., Kajitani, R., Toyoda, A., Itoh, T., Ikeo, K., Sasaki, T., et al. (2022). Gene Recruitments and Dismissals in the Argonaut Genome Provide Insights into Pelagic Lifestyle Adaptation and Shell-like Eggcase Reacquisition. Genome Biol. Evol. *14*, evac140. https://doi.org/10.1093/gbe/evac140.

45. Belcaid, M., Casaburi, G., McAnulty, S.J., Schmidbaur, H., Suria, A.M., Moriano-Gutierrez, S., Pankey, M.S., Oakley, T.H., Kremer, N., Koch, E.J., et al. (2019). Symbiotic organs shaped by distinct modes of genome evolution in cephalopods. Proc. Natl. Acad. Sci. *116*, 3030–3035. https://doi.org/10.1073/pnas.1817322116.

46. Albertin, C.B., Medina-Ruiz, S., Mitros, T., Schmidbaur, H., Sanchez, G., Wang, Z.Y., Grimwood, J., Rosenthal, J.J.C., Ragsdale, C.W., Simakov, O., et al. (2022). Genome and transcriptome mechanisms driving cephalopod evolution. Nat. Commun. *13*. https://doi.org/10.1038/s41467-022-29748-w.

47. Marino, A., Kizenko, A., Wong, W.Y., Ghiselli, F., and Simakov, O. (2022). Repeat Age Decomposition Informs an Ancient Set of Repeats Associated With Coleoid Cephalopod Divergence. Front. Genet. *13*.

48. Simakov, O., Bredeson, J., Berkoff, K., Marletaz, F., Mitros, T., Schultz, D.T., O'Connell, B.L., Dear, P., Martinez, D.E., Steele, R.E., et al. (2022). Deeply conserved synteny and the evolution of metazoan chromosomes. Sci. Adv. *8*, eabi5884. https://doi.org/10.1126/sciadv.abi5884.

49. Zou, Y., Fu, J., Liang, Y., Luo, X., Shen, M., Huang, M., Chen, Y., You, W., and Ke, C. (2024). Chromosome-level genome assembly of the ivory shell Babylonia areolata. Sci. Data *11*, 1201. https://doi.org/10.1038/s41597-024-04001-9.

50. Chen, Z., Baeza, J.A., Chen, C., Gonzalez, M.T., González, V.L., Greve, C., Kocot, K.M., Arbizu, P.M., Moles, J., Schell, T., et al. (2025). A genome-based phylogeny for Mollusca is concordant with fossils and morphology. Science *387*, 1001–1007. https://doi.org/10.1126/science.ads0215.

743   51. Zeng, Q., Liu, J., Wang, C., Wang, H., Zhang, L., Hu, J., Bao, L., and Wang, S. (2021).
744   High-quality reannotation of the king scallop genome reveals no 'gene-rich' feature and
745   evolution of toxin resistance. Comput. Struct. Biotechnol. J. *19*, 4954–4960.
746   https://doi.org/10.1016/j.csbj.2021.08.038.

747   52. Sun, S., Han, X., Han, Z., and Liu, Q. (2024). Chromosomal-scale genome assembly
748   and annotation of the land slug (Meghimatium bilineatum). Sci. Data *11*, 35.
749   https://doi.org/10.1038/s41597-023-02893-7.

750   53. Männer, L., Schell, T., Spies, J., Galià-Camps, C., Baranski, D., Ben Hamadou, A.,
751   Gerheim, C., Neveling, K., Helfrich, E.J.N., and Greve, C. (2024). Chromosome-level
752   genome assembly of the sacoglossan sea slug Elysia timida (Risso, 1818). BMC
753   Genomics *25*, 941. https://doi.org/10.1186/s12864-024-10829-7.

754   54. Ma, B., Jin, W., Fu, H., Sun, B., Yang, S., Ma, X., Wen, H., Wu, X., Wang, H., and Cao,
755   X. (2023). A High-Quality Chromosome-Level Genome Assembly of a Snail
756   Cipangopaludina cathayensis (Gastropoda: Viviparidae). Genes *14*.
757   https://doi.org/10.3390/genes14071365.

758   55. Liu, Z., Huang, Y., Chen, H., Liu, C., Wang, M., Bian, C., Wang, L., and Song, L. (2023).
759   Chromosome-level genome assembly of the deep-sea snail Phymorhynchus
760   buccinoides provides insights into the adaptation to the cold seep habitat. BMC
761   Genomics *24*, 679. https://doi.org/10.1186/s12864-023-09760-0.

762   56. Peñaloza, C., Gutierrez, A.P., Eöry, L., Wang, S., Guo, X., Archibald, A.L., Bean, T.P.,
763   and Houston, R.D. (2021). A chromosome-level genome assembly for the Pacific oyster
764   Crassostrea gigas. GigaScience *10*, giab020.
765   https://doi.org/10.1093/gigascience/giab020.

766   57. Arias-Montecino, A., Sykes, A., Álvarez-Hernán, G., de Mera-Rodríguez, J.A., Calle-
767   Guisado, V., Martín-Partido, G., Rodríguez-León, J., and Francisco-Morcillo, J. (2024).
768   Histological and scanning electron microscope observations on the developing retina of
769   the cuttlefish (Sepia officinalis Linnaeus, 1758). Tissue Cell *88*, 102417.
770   https://doi.org/10.1016/j.tice.2024.102417.

771   58. O'Brien, C.E., Mezrai, N., Darmaillacq, A.-S., and Dickel, L. (2017). Behavioral
772   development in embryonic and early juvenile cuttlefish (Sepia officinalis). Dev.
773   Psychobiol. *59*, 145–160. https://doi.org/10.1002/dev.21476.

59. Bellingham, J., Morris, A.G., and Hunt, D.M. (1998). The Rhodopsin Gene of the

Cuttlefish Sepia Officinalis: Sequence and Spectral Tuning. J. Exp. Biol. *201*, 2299–
2306. https://doi.org/10.1242/jeb.201.15.2299.

60. Chemello, G., Faraoni, V., Notarstefano, V., Maradonna, F., Carnevali, O., and
Gioacchini, G. (2022). First Evidence of Microplastics in the Yolk and Embryos of
Common Cuttlefish (Sepia officinalis) from the Central Adriatic Sea: Evaluation of
Embryo and Hatchling Structural Integrity and Development. Anim. Open Access J.
MDPI *13*. https://doi.org/10.3390/ani13010095.

61. Court, M., Macau, M., Ranucci, M., Marquês, T., Repolho, T., Lopes, V.M., Rosa, R.,
and Paula, J.R. (2024). Oxygen loss compromises growth and cognition of cuttlefish
newborns. Proc. Biol. Sci. *291*, 20241291. https://doi.org/10.1098/rspb.2024.1291.

62. Gladman, N.W., and Askew, G.N. (2023). The hydrodynamics of jet propulsion
swimming in hatchling and juvenile European common cuttlefish, Sepia officinalis. J.
Exp. Biol. *226*. https://doi.org/10.1242/jeb.246225.

63. Yang, T., Jia, Z., Chen, H., Deng, Z., Liu, W., Chen, L., and Li, L. (2020). Mechanical
design of the highly porous cuttlebone: A bioceramic hard buoyancy tank for cuttlefish.
Proc. Natl. Acad. Sci. U. S. A. *117*, 23450–23459.
https://doi.org/10.1073/pnas.2009531117.

64. Von Boletzky, S. (1987). Fecundity variation in relation to intermittent or chronic
spawning in the cuttlefish, Sepia officinalis L. (Mollusca, Cephalopoda). Bull. Mar. Sci.
40.

65. Laptikhovsky, V., Salman, A., Onsoy, B., and Katagan, T. (2003). Fecundity of the
common cuttlefish, Sepia officinalis L. (Cephalopoda, Sepiidae): A new look at an old
problem. Sci. Mar. *67*, 279–284.

66. Reiter, S., Hülsdunk, P., Woo, T., Lauterbach, M.A., Eberle, J.S., Akay, L.A., Longo, A.,
Meier-Credo, J., Kretschmer, F., Langer, J.D., et al. (2018). Elucidating the control and
development of skin patterning in cuttlefish. Nature *562*, 361–366.
https://doi.org/10.1038/s41586-018-0591-3.

67. Woo, T., Liang, X., Evans, D.A., Fernandez, O., Kretschmer, F., Reiter, S., and Laurent,
G. (2023). The dynamics of pattern matching in camouflaging cuttlefish. Nature *619*,
122–128. https://doi.org/10.1038/s41586-023-06259-2.

68. Imarazene, B., Andouche, A., Bassaglia, Y., Lopez, P.-J., and Bonnaud-Ponticelli, L.

(2017). Eye Development in Sepia officinalis Embryo: What the Uncommon Gene

Expression Profiles Tell Us about Eye Evolution. Front. Physiol. *8*, 613.

https://doi.org/10.3389/fphys.2017.00613.

69. Cocci, P., Mosconi, G., and Palermo, F.A. (2023). Effect of polycyclic aromatic

hydrocarbons on homeobox gene expression during embryonic development of

cuttlefish, Sepia officinalis. Chemosphere *325*, 138315.

https://doi.org/10.1016/j.chemosphere.2023.138315.

70. Andouche, A., Bassaglia, Y., Baratte, S., and Bonnaud, L. (2013). Reflectin genes and

development of iridophore patterns in Sepia officinalis embryos (Mollusca,

Cephalopoda). Dev. Dyn. *242*, 560–571. https://doi.org/10.1002/dvdy.23938.

71. McKenna, V., Archibald, J., Beinart, R., Dawson, M., Hentschel, U., Keeling, P., Lopez,

J., Martin-Duran, J., Petersen, J., Sigwart, J., et al. (2024). The Aquatic Symbiosis

Genomics Project: probing the evolution of symbiosis across the Tree of Life [version 2;

peer review: 1 approved, 1 approved with reservations]. Wellcome Open Res. *6*.

https://doi.org/10.12688/wellcomeopenres.17222.2.

72. Ranallo-Benavidez, T.R., Jaron, K.S., and Schatz, M.C. (2020). GenomeScope 2.0 and

Smudgeplot for reference-free profiling of polyploid genomes. Nat. Commun. *11*, 1432.

https://doi.org/10.1038/s41467-020-14998-3.

73. Vurture, G.W., Sedlazeck, F.J., Nattestad, M., Underwood, C.J., Fang, H., Gurtowski, J.,

and Schatz, M.C. (2017). GenomeScope: fast reference-free genome profiling from short

reads. Bioinformatics *33*, 2202–2204. https://doi.org/10.1093/bioinformatics/btx153.

74. Hjelmen, C.E. (2024). Genome size and chromosome number are critical metrics for

accurate genome assembly assessment in Eukaryota. Genetics *227*, iyae099.

https://doi.org/10.1093/genetics/iyae099.

75. Simão, F.A., Waterhouse, R.M., Ioannidis, P., Kriventseva, E.V., and Zdobnov, E.M.

(2015). BUSCO: assessing genome assembly and annotation completeness with single-

copy orthologs. Bioinformatics *31*, 3210–3212.

https://doi.org/10.1093/bioinformatics/btv351.

76. Zeng, X., Yi, Z., Zhang, X., Du, Y., Li, Y., Zhou, Z., Chen, S., Zhao, H., Yang, S., Wang,

Y., et al. (2024). Chromosome-level scaffolding of haplotype-resolved assemblies using

Hi-C data without reference genomes. Nat. Plants *10*, 1184–1200.

https://doi.org/10.1038/s41477-024-01755-3.

77. Zhou, C., McCarthy, S.A., and Durbin, R. (2023). YaHS: yet another Hi-C scaffolding
tool. Bioinformatics *39*, btac808. https://doi.org/10.1093/bioinformatics/btac808.

78. Dudchenko, O., Shamim, M.S., Batra, S.S., Durand, N.C., Musial, N.T., Mostofa, R.,
Pham, M., Hilaire, B.G.S., Yao, W., Stamenova, E., et al. (2018). The Juicebox
Assembly Tools module facilitates de novo assembly of mammalian genomes with
chromosome-length scaffolds for under $1000. Preprint at bioRxiv,
https://doi.org/10.1101/254797 https://doi.org/10.1101/254797.

79. Gao, Y.M., and Natsukari, Y. (1990). Karyological studies on seven cephalopods. Venus
Jpn. J. Malacol. *49*, 126–145.

80. Jazayeri, A., Papan, F., Motamedi, H., and Mahmoudi, S. (2011). Karyological
investigation of Persian Gulf cuttle fish (sepia arabica) in the coasts of Khuzestan
province. Life Sci. J. *8*.

81. Ebrahimi Pour, M. (2009). The Study of Persian Gulf Cuttlefish (Sepia pharaonis)
Chromosome Via Incubation of Blood Cells. Appl. Biol. *22*, 1–8.
https://doi.org/10.22051/jab.2009.3307.

82. Vitturi, R., Rasotto, M.B., and Farinella-Ferruzza, N. (1982). The chromosomes of 16
molluscan species. Ital. J. Zool. *49*, 61–71.

83. The Darwin Tree of Life Project Consortium (2022). Sequence locally, think globally: The
Darwin Tree of Life Project. Proc. Natl. Acad. Sci. *119*, e2115642118.
https://doi.org/10.1073/pnas.2115642118.

84. Li, H. (2018). Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics *34*,
3094–3100. https://doi.org/10.1093/bioinformatics/bty191.

85. López-Córdova, D.A., Avaria-Llautureo, J., Ulloa, P.M., Braid, H.E., Revell, L.J., Fuchs,
D., and Ibáñez, C.M. (2022). Mesozoic origin of coleoid cephalopods and their abrupt
shifts of diversification patterns. Mol. Phylogenet. Evol. *166*, 107331.
https://doi.org/10.1016/j.ympev.2021.107331.

86. Coffing, G., TITTES, S., Small, S., Songco-Casey, J.O., Pungor, J., Piscopo, D., Miller,
A., Niell, C., and Kern, A. (2024). Data for: Cephalopod Sex Determination and its
Ancient Evolutionary Origin. (Zenodo). https://doi.org/10.5281/zenodo.14010217
https://doi.org/10.5281/zenodo.14010217.

87. Coffing, G.C., Tittes, S., Small, S.T., Songco-Casey, J.O., Piscopo, D.M., Pungor, J.R.,
Miller, A.C., Niell, C.M., and Kern, A.D. (2025). Cephalopod sex determination and its
ancient evolutionary origin. Curr. Biol. *35*, 931-939.e4.
https://doi.org/10.1016/j.cub.2025.01.005.

88. Flynn, J.M., Hubley, R., Goubert, C., Rosen, J., Clark, A.G., Feschotte, C., and Smit,
A.F. (2020). RepeatModeler2 for automated genomic discovery of transposable element
families. Proc. Natl. Acad. Sci. *117*, 9451–9457.
https://doi.org/10.1073/pnas.1921046117.

89. Smit, A., Hubley, R., and Green, P. RepeatMasker http://www.repeatmasker.org.

90. Feschotte, C. (2008). The contribution of transposable elements to the evolution of
regulatory networks. Nat. Rev. Genet. *9*, 397–405. https://doi.org/10.1038/nrg2337.

91. Hoff, K.J., Lomsadze, A., Borodovsky, M., and Stanke, M. (2019). Whole-Genome
Annotation with BRAKER. Methods Mol. Biol. Clifton NJ *1962*, 65–95.
https://doi.org/10.1007/978-1-4939-9173-0_5.

92. Brůna, T., Hoff, K.J., Lomsadze, A., Stanke, M., and Borodovsky, M. (2021). BRAKER2:
automatic eukaryotic genome annotation with GeneMark-EP+ and AUGUSTUS
supported by a protein database. NAR Genomics Bioinforma. *3*, lqaa108.
https://doi.org/10.1093/nargab/lqaa108.

93. Brůna, T., Gabriel, L., and Hoff, K.J. (2024). Navigating Eukaryotic Genome Annotation
Pipelines: A Route Map to BRAKER, Galba, and TSEBRA. Preprint at arXiv,
https://doi.org/10.48550/arXiv.2403.19416 https://doi.org/10.48550/arXiv.2403.19416.

94. Gabriel, L., Hoff, K.J., Brůna, T., Borodovsky, M., and Stanke, M. (2021). TSEBRA:
transcript selector for BRAKER. BMC Bioinformatics *22*, 566.
https://doi.org/10.1186/s12859-021-04482-0.

95. Hoff, K.J., Lange, S., Lomsadze, A., Borodovsky, M., and Stanke, M. (2016). BRAKER1:
Unsupervised RNA-Seq-Based Genome Annotation with GeneMark-ET and
AUGUSTUS. Bioinforma. Oxf. Engl. *32*, 767–769.
https://doi.org/10.1093/bioinformatics/btv661.

96. Stanke, M., Schöffmann, O., Morgenstern, B., and Waack, S. (2006). Gene prediction in
eukaryotes with a generalized hidden Markov model that uses hints from external
sources. BMC Bioinformatics *7*, 62. https://doi.org/10.1186/1471-2105-7-62.

97. Stanke, M., Diekhans, M., Baertsch, R., and Haussler, D. (2008). Using native and
syntenically mapped cDNA alignments to improve de novo gene finding. Bioinformatics
24, 637–644. https://doi.org/10.1093/bioinformatics/btn013.

98. Li, H. (2023). Protein-to-genome alignment with miniprot. Bioinformatics *39*, btad014.
https://doi.org/10.1093/bioinformatics/btad014.

99. Iwata, H., and Gotoh, O. (2012). Benchmarking spliced alignment programs including
Spaln2, an extended version of Spaln that incorporates additional species-specific
features. Nucleic Acids Res. *40*, e161. https://doi.org/10.1093/nar/gks708.

100.   Gotoh, O. (2008). A space-efficient and accurate method for mapping and aligning
cDNA sequences onto genomic sequence. Nucleic Acids Res. *36*, 2630–2638.
https://doi.org/10.1093/nar/gkn105.

101.   Buchfink, B., Xie, C., and Huson, D.H. (2015). Fast and sensitive protein alignment
using DIAMOND. Nat. Methods *12*, 59–60. https://doi.org/10.1038/nmeth.3176.

102.   Kovaka, S., Zimin, A.V., Pertea, G.M., Razaghi, R., Salzberg, S.L., and Pertea, M.
(2019). Transcriptome assembly from long-read RNA-seq alignments with StringTie2.
Genome Biol. *20*, 278. https://doi.org/10.1186/s13059-019-1910-1.

103.   Brůna, T., Lomsadze, A., and Borodovsky, M. (2024). GeneMark-ETP significantly
improves the accuracy of automatic annotation of large eukaryotic genomes. Genome
Res. *34*, 757–768. https://doi.org/10.1101/gr.278373.123.

104.   Huang, N., and Li, H. (2023). compleasm: a faster and more accurate
reimplementation of BUSCO. Bioinformatics *39*, btad595.
https://doi.org/10.1093/bioinformatics/btad595.

105.   Pertea, G., and Pertea, M. (2020). GFF Utilities: GffRead and GffCompare. Preprint
at F1000Research, https://doi.org/10.12688/f1000research.23297.2
https://doi.org/10.12688/f1000research.23297.2.

106.   Gabriel, L., Brůna, T., Hoff, K.J., Ebel, M., Lomsadze, A., Borodovsky, M., and
Stanke, M. (2024). BRAKER3: Fully automated genome annotation using RNA-seq and
protein evidence with GeneMark-ETP, AUGUSTUS, and TSEBRA. Genome Res. *34*,
769–777. https://doi.org/10.1101/gr.278090.123.

107.   Rogers, T.F., Yalçin, G., Briseno, J., Vijayan, N., Nyholm, S.V., and Simakov, O.

(2024). Gene modelling and annotation for the Hawaiian bobtail squid, Euprymna
scolopes. Sci. Data *11*, 40. https://doi.org/10.1038/s41597-023-02903-8.

108.   Zhang, Y., Mao, F., Mu, H., Huang, M., Bao, Y., Wang, L., Wong, N.-K., Xiao, S., Dai,
H., Xiang, Z., et al. (2021). The genome of Nautilus pompilius illuminates eye evolution
and biomineralization. Nat. Ecol. Evol. *5*, 927–938. https://doi.org/10.1038/s41559-021-
01448-6.

109.   Nevers, Y., Rossier, V., Train, C.M., Altenhoff, A., Dessimoz, C., and Glover, N.
(2022). Multifaceted quality assessment of gene repertoire annotation with OMArk.
bioRxiv, 2022.11.25.517970. https://doi.org/10.1101/2022.11.25.517970.

110.   Blum, M., Andreeva, A., Florentino, L.C., Chuguransky, S.R., Grego, T., Hobbs, E.,
Pinto, B.L., Orr, A., Paysan-Lafosse, T., Ponamareva, I., et al. (2025). InterPro: the
protein sequence classification resource in 2025. Nucleic Acids Res. *53*, D444–D456.
https://doi.org/10.1093/nar/gkae1082.

111.   Cantalapiedra, C.P., Hernández-Plaza, A., Letunic, I., Bork, P., and Huerta-Cepas, J.
(2021). eggNOG-mapper v2: Functional Annotation, Orthology Assignments, and
Domain Prediction at the Metagenomic Scale. Mol. Biol. Evol. *38*, 5825–5829.
https://doi.org/10.1093/molbev/msab293.

112.   Huerta-Cepas, J., Forslund, K., Coelho, L.P., Szklarczyk, D., Jensen, L.J., von
Mering, C., and Bork, P. (2017). Fast Genome-Wide Functional Annotation through
Orthology Assignment by eggNOG-Mapper. Mol. Biol. Evol. *34*, 2115–2122.
https://doi.org/10.1093/molbev/msx148.

113.   Tanner, A.R., Fuchs, D., Winkelmann, I.E., Gilbert, M.T.P., Pankey, M.S., Ribeiro,
Â.M., Kocot, K.M., Halanych, K.M., Oakley, T.H., da Fonseca, R.R., et al. (2017).
Molecular clocks indicate turnover and diversification of modern coleoid cephalopods
during the Mesozoic Marine Revolution. Proc. R. Soc. B Biol. Sci. *284*, 20162818.
https://doi.org/10.1098/rspb.2016.2818.

114.   Anderson, F.E., and Lindgren, A.R. (2021). Phylogenomic analyses recover a clade
of large-bodied decapodiform cephalopods. Mol. Phylogenet. Evol. *156*, 107038.
https://doi.org/10.1016/j.ympev.2020.107038.

115.   Lindgren, A.R., Pankey, M.S., Hochberg, F.G., and Oakley, T.H. (2012). A multi-gene
phylogeny of Cephalopoda supports convergent morphological evolution in association
with multiple habitat shifts in the marine environment. BMC Evol. Biol. *12*, 129.

https://doi.org/10.1186/1471-2148-12-129.

116.   Sanchez, G., Setiamarga, D.H.E., Tuanapaya, S., Tongtherm, K., Winkelmann, I.E.,
Schmidbaur, H., Umino, T., Albertin, C., Allcock, L., Perales-Raya, C., et al. (2018).
Genus-level phylogeny of cephalopods using molecular markers: current status and
problematic areas. PeerJ *6*, e4331. https://doi.org/10.7717/peerj.4331.

117.   Schultz, D.T., Haddock, S.H.D., Bredeson, J.V., Green, R.E., Simakov, O., and
Rokhsar, D.S. (2023). Ancient gene linkages support ctenophores as sister to other
animals. Nature *618*, 110–117. https://doi.org/10.1038/s41586-023-05936-6.

118.   Lukhtanov, V.A., Dincă, V., Talavera, G., and Vila, R. (2011). Unprecedented within-
species chromosome number cline in the Wood White butterfly Leptidea sinapis and its
significance for karyotype evolution and speciation. BMC Evol. Biol. *11*, 109.
https://doi.org/10.1186/1471-2148-11-109.

119.   Lukhtanov, V.A., Dincă, V., Friberg, M., Šíchová, J., Olofsson, M., Vila, R., Marec, F.,
and Wiklund, C. (2018). Versatility of multivalent orientation, inverted meiosis, and
rescued fitness in holocentric chromosomal hybrids. Proc. Natl. Acad. Sci. *115*, E9610–
E9619. https://doi.org/10.1073/pnas.1802610115.

120.   Ponte, G., Roumbedakis, K., COST Action FA1301, Galligioni, V., Dickel, L.,
Bellanger, C., Pereira, J., Vidal, E.A., Grigoriou, P., Alleva, E., et al. (2023). General and
species-specific recommendations for minimal requirements for the use of cephalopods
in scientific research. Lab. Anim. *57*, 26–39.
https://doi.org/10.1177/00236772221111261.

121.   Fiorito, G., Affuso, A., Basil, J., Cole, A., de Girolamo, P., D'Angelo, L., Dickel, L.,
Gestal, C., Grasso, F., Kuba, M., et al. (2015). Guidelines for the Care and Welfare of
Cephalopods in Research –A consensus based on an initiative by CephRes, FELASA
and the Boyd Group. Lab. Anim. *49*, 1–90. https://doi.org/10.1177/0023677215580006.

122.   Andrews, P.L.R., Darmaillacq, A.-S., Dennison, N., Gleadall, I.G., Hawkins, P.,
Messenger, J.B., Osorio, D., Smith, V.J., and Smith, J.A. (2013). The identification and
management of pain, suffering and distress in cephalopods, including anaesthesia,
analgesia and humane killing. J. Exp. Mar. Biol. Ecol. *447*, 46–64.
https://doi.org/10.1016/j.jembe.2013.02.010.

123.   Cartolano, M., Huettel, B., Hartwig, B., Reinhardt, R., and Schneeberger, K. (2016).
cDNA Library Enrichment of Full Length Transcripts for SMRT Long Read Sequencing.

PLOS ONE *11*, e0157779. https://doi.org/10.1371/journal.pone.0157779.

124.   Cheng, H., Concepcion, G.T., Feng, X., Zhang, H., and Li, H. (2021). Haplotype-
resolved de novo assembly using phased assembly graphs with hifiasm. Nat. Methods
18, 170–175. https://doi.org/10.1038/s41592-020-01056-5.

125.   Adachi, K., Ohnishi, K., Kuramochi, T., Yoshinaga, T., and Okumura, S.-I. (2014).
Molecular cytogenetic study in Octopus (Amphioctopus) areolatus from Japan. Fish. Sci.
80, 445–450. https://doi.org/10.1007/s12562-014-0703-4.

126.   Lyčka, M., Bubeník, M., Závodník, M., Peska, V., Fajkus, P., Demko, M., Fajkus, J.,
and Fojtová, M. (2024). TeloBase: a community-curated database of telomere
sequences across the tree of life. Nucleic Acids Res. *52*, D311–D321.
https://doi.org/10.1093/nar/gkad672.

127.   Akasaki, T., Nikaido, M., Tsuchiya, K., Segawa, S., Hasegawa, M., and Okada, N.
(2006). Extensive mitochondrial gene arrangements in coleoid Cephalopoda and their
phylogenetic implications. Mol. Phylogenet. Evol. *38*, 648–658.
https://doi.org/10.1016/j.ympev.2005.10.018.

128.   Li, H. (2013). Seqtk: a fast and lightweight tool for processing FASTA or FASTQ
sequences.

129.   Shumate, A., Wong, B., Pertea, G., and Pertea, M. (2022). Improved transcriptome
assembly using a hybrid of long and short reads with StringTie. PLOS Comput. Biol. *18*,
e1009730. https://doi.org/10.1371/journal.pcbi.1009730.

130.   Haas, B.J. GitHub - TransDecoder/TransDecoder: TransDecoder source.
https://github.com/TransDecoder/TransDecoder.

131.   R Core Team (2024). R: A Language and Environment for Statistical Computing (R
Foundation for Statistical Computing).

132.   Lovell, J.T., Sreedasyam, A., Schranz, M.E., Wilson, M., Carlson, J.W., Harkess, A.,
Emms, D., Goodstein, D.M., and Schmutz, J. (2022). GENESPACE tracks regions of
interest and gene copy number variation across multiple genomes. eLife *11*, e78526.
https://doi.org/10.7554/eLife.78526.

133.   Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P.,
Chaisson, M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner.

Bioinforma. Oxf. Engl. *29*, 15–21. https://doi.org/10.1093/bioinformatics/bts635.

134.   Pedersen, B.S., and Quinlan, A.R. (2018). Mosdepth: quick coverage calculation for
genomes and exomes. Bioinformatics *34*, 867–868.
https://doi.org/10.1093/bioinformatics/btx699.

135.   Challis, R., Richards, E., Rajan, J., Cochrane, G., and Blaxter, M. (2020). BlobToolKit
– Interactive Quality Assessment of Genome Assemblies. G3 GenesGenomesGenetics
10, 1361–1374. https://doi.org/10.1534/g3.119.400908.

136.   Kröger, B., Vinther, J., and Fuchs, D. (2011). Cephalopod origin and evolution: A
congruent picture emerging from fossils, development and molecules. BioEssays *33*,
602–613. https://doi.org/10.1002/bies.201100001.

137.   Song, W., Li, R., Zhao, Y., Migaud, H., Wang, C., and Bekaert, M. (2021). Pharaoh
Cuttlefish, Sepia pharaonis, Genome Reveals Unique Reflectin Camouflage Gene Set.
Front. Mar. Sci. *8*.