Title: Increased rates of hybridization in swordtail fish are associated with water pollution

Authors:

Benjamin M Moran[1,2,3], Wilson F Ramírez-Duarte[2,4,5], Daniel L Powell[1,2,6], Terrance TL Yang[1],

Theresa R Gunn[1,2], Gaston I Jofre-Rodríguez[2,7], Cheyenne Y Payne[8,9], Erik NK Iverson[10],

Gabriel E Preising[1,2], Shreya M Banerjee[3], Alexandra E Donny[1,11], Rhea Sood[1], John J

Baczenas[1], Gabriela María Vázquez Adame[2], Carla Gutiérrez-Rodríguez[12], Chelsea M

Rochman[4], Molly Schumer[1,2,13]*, Gil G Rosenthal[2,14]*

Affiliations:

[1]Department of Biology, Stanford University

[2]Centro de Investigaciones Científicas de las Huastecas "Aguazarca", A.C.

[3] Present Address: Department of Evolution and Ecology, University of California at Davis

[4]Department of Ecology & Evolutionary Biology, University of Toronto

[5]Present Address: Department of Environmental & Physical Sciences, Concordia University of

Edmonton

[6] Department of Biological Sciences, Louisiana State University

[7]Department of Biology, Virginia Commonwealth University

[8]Department of Ecology and Evolutionary Biology, University of California, Santa Cruz

[9]Southwest Fisheries Science Center, National Marine Fisheries Service, National Oceanic and

Atmospheric Administration

[10]Department of Integrative Biology, University of Texas at Austin

[11]Department of Genome Sciences, University of Washington

[12] Red de Biología Evolutiva, Instituto de Ecología A.C

[13]Freeman Hrabowski Fellow, Howard Hughes Medical Institute

[14]Dipartimento di Biologia, Università di Padova

***co-supervised this work

Running Head: Anthropogenic hybridization in swordtails

Keywords: hybridization, hybrid populations, anthropogenic change, water pollution, mate

choice, freshwater fish, swordtail fish

Abstract

Biodiversity loss can occur when disturbance compromises the reproductive barriers

between species, causing them to collapse into a single population through hybridization. Recent

research has documented an increasing number of cases where anthropogenic environmental

disturbance is associated with new hybrid populations, but few studies have been able to

quantitatively compare potential environmental drivers and test possible mechanisms connecting

interspecific hybridization to anthropogenic disturbance. Here, we combine genomic and

chemical surveys to explore the loss of reproductive isolation between the sister species

Xiphophorus malinche and *X. birchmanni*, fishes whose riverine habitat in the Sierra Madre

Oriental of Mexico is increasingly impacted by human-mediated disturbance. By sequencing

whole genomes of thousands of fish, we characterize the landscape of hybridization between

these species in four distinct rivers. Ancestry structure varied dramatically across rivers, ranging

from stable coexistence to clinal hybrid zones, hinting that dynamics of hybridization in this

system may be environmentally dependent. In one river, we discovered that a sudden transition

in patterns of hybridization coincides with its passage through an urbanized area, with upstream

sites showing distinct ancestry clusters and downstream sites showing a swarm of hybrids with

variable ancestry. Water chemistry measurements show that water quality changes significantly

over this area, including in parameters known to disrupt fish olfaction and mating. We

hypothesize that the hybrid swarm downstream of the town is driven by disruption of olfaction

that impacts mating preferences. We test this by sequencing mothers and embryos from sites

downstream of the urbanized area, and find reduced assortative mating, in contrast to strong

assortative mating in sites upstream. We document alterations of the olfactory epithelium at the

same site which are consistent with chemical irritation. Taken together, our work illuminates the

mechanisms linking anthropogenic disturbance to the breakdown of reproductive isolation in

hybridizing species.

Introduction

Biologists have known for decades that hybridization can be associated with environmental disturbance (1, 2). Disturbed habitats bring together species which do not coexist under other circumstances, create novels environments which may provide better habitat for hybrids than those of undisturbed habitats, and may alter the interactions between species that previously coexisted without interbreeding (1–3). While natural changes to the environment may have been the major source of disturbance historically, human impacts on the environment (e.g. land conversion, pollution, resource extraction) may play an analogous role in promoting hybridization in recent decades (4–7). As human impacts on the environment become ubiquitous (8), the connection between disturbance and hybridization would suggest the number of hybrid zones should be growing as well. In fact, the rate of hybridization appears to be increasing across a range of taxa, largely due to anthropogenic impacts on the environment (3).

In investigating the connection between disturbance and hybridization, early researchers hypothesized that hybrids appear in disturbed areas because the barriers that prevent mating between closely related species can break down in novel environmental contexts (1). Such "premating" barriers often arise early in the divergence of two lineages (9, 10) and thus may be especially important in the origin and preservation of biodiversity (11–13). The link between disturbance and hybridization has been supported by empirical work that shows that premating barriers, such as a behavioral preference to mate assortatively by species, are highly context-dependent (14–18), sensitive to environmental conditions (19, 20), and may disappear in the face of disturbance that impacts mechanisms of mate discrimination (6, 21, 22). Human modification

of the environment can interfere with mechanisms used to identify conspecifics, either by

directly affecting signal production or transmission or by abolishing conspecific preferences in

the choosy sex (6, 23, 24). In aquatic species, in particular, a variety of chemical pollutants have

been shown to ablate or even reverse conspecific mate preferences (6, 23–27).

If assortative mating is a major reproductive barrier between species (or genetically

distinct subpopulations), weakening this barrier can lead to a sudden collapse of population

structure and phenotypic divergence (28–32), often referred to as a "hybrid swarm". Since mate

discrimination is typically weaker in hybrids than in parental species (21), this can lead to

irreversible phenotypic and genetic homogenization through hybridization (i.e. "speciation

reversal" or "genetic swamping"). Although hybridization may sometimes lead to the formation

of new species (33), hybrid speciation without a change in chromosome number appears to be

relatively rare (34), and the immediate consequence of the formation of hybrid swarms is more

often the loss of genetic differentiation and functional biodiversity (7, 35). Moreover, early-

generation hybrids often experience inviability or sterility due to intrinsic hybrid

incompatibilities (36, 37) between the parental genomes (although we note that hybridization can

also facilitate adaptation; (38, 39)). Thus, in addition to its direct effects of homogenizing

divergent species, hybridization may also decrease overall population size and fecundity.

The swordtail fishes of the genus *Xiphophorus* provide a compelling system in which to

study the barriers to gene flow in natural populations and their breakdown following

anthropogenic disturbance. As model systems for hybridization and mate choice, *Xiphophorus*

species have been subject to numerous studies showing that mating preferences can be altered by

environmental and social context (15, 16, 21, 23). Swordtails are found in the eastern slopes of

the Sierra Madre Oriental and Gulf coastal plain of Mexico, a region experiencing growing

human population density and agricultural land conversion since the 20th century (40, 41). This

change in land cover has impacted water quality in the region, while also threatening the area's

exceptional biodiversity (42, 43). In particular, the sister species *X. malinche* and *X. birchmanni*

overlap in the montane rivers of Hidalgo, Mexico, where steep slopes create frequent barriers to

migration (waterfalls, cascades, and subterranean river reaches) and coincide with dramatic

ecological variation over short geographic distances. *X. malinche* occupy cool headwater streams

while *X. birchmanni* are found in warmer lower-elevation rivers, creating the potential for

ecological selection against hybrids (44, 45). In addition, early-generation hybrids between these

species suffer from severe genetic incompatibilities (46–48). However, some hybrids are viable

and fertile, and natural hybrid zones can be found in multiple independent stream reaches (49,

50). Researchers first reported hybridization between *X. malinche* and *X. birchmanni* in 1997,

whereas hybrids were not detected when sampling this site a decade earlier (51, 52). Meanwhile,

the period of 1980-1995 saw a 37% increase in population in the state of Hidalgo (53).

Independent genetic approaches have estimated that these hybrid zones are on the order of ~100

generations old (54), consistent with hybridization beginning in the late 1900s. The recent onset

of hybridization begs the question of what triggered it, and whether it was tied to the concurrent

socio-ecological transformation of the region.

Though we lack ecological data from these rivers prior to the beginning of hybridization,

spatial patterns in present day populations may provide insights into the drivers of hybridization.

The genetic structure in most hybrid populations is best described as a hybrid swarm lacking

population structure (49). One exception to this pattern among described populations is a river

reach just upstream of the town of Calnali, Hidalgo. There, genetic studies have described two

subpopulations of *malinche*-like and *birchmanni*-like admixed fish that exist in sympatry (Fig. 1;

(55). Sequencing over multiple years showed that this bimodal population structure has been

stable over time, and is best explained by strong assortative mating between individuals of the

same ancestry group (55). However, recent sampling of a population downstream in the same

river showed that population structure disappears just downstream of the town of Calnali (46),

raising the possibility that assortative mating breaks down after exposure to a point pollution

source.

Here, we use whole-genome sequencing of 2620 fish and comprehensive surveys of

water chemistry to study the relationship between environmental conditions and patterns of

hybridization across multiple *X. malinche* × *X. birchmanni* hybrid zones in Hidalgo, Mexico. We

survey hybrid ancestry in four rivers and compare hybrid population structure across

demographically independent hybrid zones. We find that assortative mating between ancestry

clusters in the Río Calnali breaks down at the precise point where anthropogenic inputs

drastically change water chemistry, leading to a sudden change in population structure. Based on

comprehensive water chemistry measurements, we identify multiple pollutants which spike at the

point of population structure collapse. We find that this pattern of population structure collapse is

unique among rivers surveyed, with other river systems showing little change in population

structure over broad geographic distances. Using land use classification of satellite imagery, we

confirm that these changes in water chemistry downstream on the Río Calnali are coupled with a

uniquely high density of built infrastructure in close proximity to the river. Given the known

importance of olfactory communication in swordtail mate choice, we test for evidence of damage

to olfactory rosettes as a result of natural or lab exposure to pollutants. We find evidence of

greater histological damage in polluted sites compared to less disturbed sites. This work thus

establishes a quantitative link between environment, phenotype, and genomics in an

anthropogenic hybrid zone, deepening our understanding of human impacts on biodiversity.

Materials and Methods

Study area and sample collection

We collected fin clips from 44 sites across four stream basins which together compose the Río Atlapexco drainage of Hidalgo, Mexico (see Table S1 for full list with GPS coordinates) for low-coverage whole genome sequencing. Fish were collected using baited minnow traps and anesthetized in a buffered solution of 100 mg/mL MS-222 dissolved in stream water (Stanford APLAC protocol #33071). Fish were allowed to recover after fin clips were collected, then were returned to the pool from which they had been sampled. Tissue was stored in 95% ethanol for DNA extraction. A subset of female fish from the Calnali Low population (Fig. 1) were euthanized and stored with their embryos in 95% ethanol for analysis of assortative mating (see below). In addition, 30 fish each from the Aguazarca and Calnali Low populations were decapitated following euthanasia, with whole heads preserved in 10% neutral buffered formalin for olfactory histology (see below).

DNA extraction and coverage whole-genome sequencing

DNA was extracted from preserved fin clips using the Agilent DNAdvance extraction kit (Beckman Coulter, Brea, California). The protocol for extractions followed the manufacturer's instructions for the tissue extraction version of the protocol except that we used half-reactions. Extracted DNA was quantified on a BioTek Cytation5 (Agilent, Santa Clara, CA) microplate reader, and diluted to 10 ng/µl for library preparation. We used a previously described

tagmentation-based protocol to prepare DNA for sequencing (44). Libraries were dual-indexed

via amplification (12 PCR cycles) and purified using 18% SPRI beads. Libraries were quantified

on a Qubit Fluorometer (Thermo Scientific, Wilmington, DE) and library size distribution was

visualized on an Agilent 4200 Tapestation (Agilent, Santa Clara, CA). Libraries were sequenced

at Weill Cornell Medical College, Harvard Bauer Core, and Admera Health Services (South

Plainfield, NJ).

Local ancestry inference

Hybrids between *X. birchmanni* and *X. malinche* harbor ancestry from both species that

varies locally along the genome and globally between individuals and populations. We used an

approach developed by our group for accurate local ancestry inference in *X. malinche × X.*

birchmanni hybrids that relies on a hidden Markov model and known ancestry informative sites

that distinguish *X. birchmanni* and *X. malinche* to infer local ancestry (*ancestryinfer*; (56)). Both

simulation studies (56) and analyses of populations of known ancestry have indicated that this

approach has a low error rate (~0.1% per ancestry informative site (47, 57)). Likewise, inference

of local ancestry is robust to a number of technical challenges including parameter

misspecification (56).

We inferred local ancestry using *ancestryinfer* from whole genome sequence data for all

individuals for which we collected more than 300,000 reads from the sequencing approach

described above. This coverage cutoff was based on previous simulation studies (56) and

resulted in 12 to 702 individuals per site post-filtering (median 32; Table S1). *Ancestryinfer*

outputs posterior probabilities of ancestry for all informative sites along the genome. In the case

of *X. malinche* × *X. birchmanni* hybrids, this results in estimates of ancestry at 729,167 ancestry

informative sites across 24 chromosomes. Since we were primarily interested in accurately

estimating global admixture proportion for individuals along a spatial cline, we converted

posterior probabilities to 'hard calls' using a threshold of 0.9. For example, if an individual had

greater than 0.9 posterior probability for *X. malinche* ancestry, we converted that site to an *X.*

malinche state. If a site did not have a posterior probability ≥0.9 for any ancestry state, we

converted the site to missing data.

We used these hard calls to estimate the proportion of the genome derived from each

parental species. We estimated this proportion using two approaches. First, we calculated the

ratio of the number of alleles called for the *X. malinche* state out of the total number of alleles

called (treating heterozygous sites as 1 allele for each parental species). We also used a spatially

aware approach, given that there is variation in the distance between ancestry informative sites.

To do so, we calculated the proportion of the genome covered by high-confidence *X. malinche* or

X. birchmanni ancestry tracts, excluding regions that were not confidently called from the

denominator. In practice, this approach generated nearly identical results (Supp. Info 1; Fig. S1).

With these results in hand, we visualized the distribution of admixture proportion in

populations along the Río Calnali, Río Pochula, Río Huazalingo, and Río Conzintla (Fig. 2). As

previous results had indicated the presence of bimodal ancestry structure in upstream populations

of the Río Calnali, we also calculated Hartigan's dip statistic (58), which quantifies departures

from a unimodal distribution, on samples from each collection site. Since we used this test across

many sites, we applied a Bonferroni correction for the number of tests. Samples were collected

between 2006 and 2024, with the highest sampling density between 2014 and 2022 (Table S2).

Given the broad time period over which samples were collected, we also evaluated whether

ancestry structure was stable at a given site across the study period (Supp. Info 2; Fig. S2; (55).

Water chemistry surveys

We chose a subset of 32 sampling sites that captured the full range of variation in

ancestry structure present in each river and collected multimodal water quality measurements

from these sites between February 2022 and September 2024 (Fig. 1). Due to changes in

equipment availability over time, sample sizes for individual water quality measurements varied

between 0 and 13 per location and measure; for all measures and sample sizes per location, see

Table S3. We measured temperature, conductivity, dissolved oxygen, and fluorescent dissolved

organic matter (fDOM) using YSI EXO 2 and EXO 1s multiparameter sondes (Xylem Inc.).

Turbidity was measured in nephelometric turbidity units (NTUs) with an Orion AQ4500

Turbidity Meter (Thermo Fisher). Using a YSI 9500 photometer, we applied colorimetric assays

to measure pH, total alkalinity, total hardness, calcium hardness, and concentrations of sulfate,

sulfite, sulfide, phosphate, ammonia, nitrite, and nitrate. Sonde measurements, turbidity

measurement, and colorimetry tests were performed at the site of sampling to ensure the accurate

measurement of unstable chemicals, while further measurements were performed on collected

water samples. For measurements of dissolved organic carbon (DOC) and total nitrogen, we

stored samples in glass vials capped with PTFE/silicone septa, which were combusted at 500 °C

for 1 hour and rinsed ten times with ultrapure water (Milli-Q® Merck) prior to sample collection.

Water samples were processed on-site by syringe-filtering through a 0.45 μm polyethersulfone

(PES) membrane, acidifying to pH 2-3 using concentrated hydrochloric acid, and capping with

no air bubbles remaining in the vial. DOC and N measurements were carried out on a Shimadzu

TOC-L analyzer using the TC/IC and TN protocols, respectively, at the Environmental

Engineering Laboratory of the National Autonomous University of Mexico (UNAM) and the

Environmental Measurements Facility at Stanford University. Total and dissolved metals were

measured from samples collected in 50 mL Falcon tubes triple-rinsed with sample water, with

samples for dissolved metals filtered through 0.45 μm PES and all samples acidified to a pH of

~2 using trace metal grade nitric acid. Inductively coupled plasma mass spectrometry (ICP-MS)

detection of 39 ions was carried out following a modified version of the APHA3030B/6020A

methods by ALS Environmental (Waterloo, ON; see Table S4 for full list of ions and QA/QC).

Given that many metals in the filtered samples were below the lower detection limits of ICP-MS,

we used ICP-MS measurements of total concentration from unfiltered samples in all downstream

statistical analyses.

Geospatial Analyses

To complement our measurements of water chemistry, we measured land use in each

river catchment as a proxy for anthropogenic disturbance. Briefly, we used Google Earth Engine

to access Sentinel-2 multispectral imagery from the month of May in the years 2020-2023 (59),

removed images with > 20% cloud cover, masked pixels flagged as clouds, then took the median

intensity for each band and pixel to construct a composite image with bands 11 (1610 nm, short-

wave infrared, 20 m^2 resolution), 8 (842 nm, near-infrared, 10 m^2), and 2 (490 nm, blue, 10 m^2).

This spectral combination provides better contrast between built infrastructure, herbaceous, and

woody vegetation than the visible spectrum (60). All further analyses were performed in ArcGIS

Pro (version 3.3.1) with the Spatial Analyst and Image Analyst extensions. We segmented the

three-band raster file using the Segmentation tool, with a spectral detail of 18, a spatial detail of

19, and a minimum size per object of 20 pixels. Next, we selected four land use types for

classification: water, forest (continuous tree cover, excluding plantation-style agriculture), non-

forest vegetation (grassland, pasture, or cultivated crops, including plantation-style orchard), and

developed land (buildings, mines, and roads). We included water surfaces (primarily to prevent

their misclassification as other land use types), forest as the primary undisturbed land cover in

the area, and non-forest vegetation as an additional form of human disturbance beyond hardened

infrastructure. Using 3 m^2-resolution imagery of the study area in March 2022 downloaded from

Planet Labs (61), we identified 60 polygons of each land class to serve as training samples.

Given the short time frame between the capture of our Sentinel and Planet Labs image sets, we

assumed that there would be negligible change in the proportion of land classes. We then used

the Classification tool to perform object-based land use classification using a support vector

machine algorithm with maximum 500 samples per class. We visually inspected the resulting

classification for pixels misclassified as developed land, and removed these pixels from the

dataset prior to further analyses (0.8% of all pixels).

To delineate catchments for each site, we downloaded a digital elevation model (DEM) of

the state of Hidalgo with 15 m^2 resolution from the Mexican National Institute for Statistics and

Geography (62). Starting with this DEM, we calculated flow direction and flow accumulation,

snapped each site to the nearest pour point pixel, and calculated the land area which drained into

the river upstream of a given site (Fig. S3), using the Flow Direction, Flow Accumulation, Snap

Pour Points, and Watershed tools, respectively. To identify the land area with greatest potential

for runoff into waterways (i.e. greatest influence on water chemistry), we used the Raster

Calculator tool. We converted all cells which received flow from at least 3000 uphill cells into a

polyline object representing the relevant waterways in our study area (Fig. S3). We then used the

Buffer tool to identify the land area within 500 m, 100 m, and 50 m of these waterways, and used

the Clip Raster tool to identify the subset of classified land use pixels within each of these

buffers. Using the Zonal Histogram tool, we summed the pixel counts of each land use category

in each site's subcatchment, then calculated the proportion of the total pixels per subcatchment

composed of each land use type. Finally, we calculated cumulative land use within a site's

watershed by summing the land use categories within its own site-specific subcatchment and

within the subcatchments of all upstream sites.

Statistical analysis of environmental data

We explored overall differences in water chemistry between sites and drainages using

Principal Component Analysis (PCA). Given differing sample sizes of ICP-MS measurement

relative to other measures, we conducted two separate PCAs, one for ICP-MS metals

measurement ($n = 66$ measurements) and one for all other data ($n = 58$). For water chemistry

parameters other than metals, we included only measurements which were consistently collected

throughout the length of the field study, including dissolved oxygen (mg/L), fDOM (quinine

sulfate units), conductivity (μS/cm), total hardness (mg/L $CaCO_3$ equivalent), total alkalinity

(mg/L $CaCO_3$ equivalent), pH, nitrite (mg/L N), ammonia (mg/L N), phosphate (mg/L PO_4),

sulfate (mg/L SO_4), and turbidity (NTU). Of the 39 ions measured in the metals measurements,

we retained 30 which had more than one measurement above the lower detection limits of ICP-

MS. For the purposes of this analysis, metals measurements below the detection limit were set to

halfway between zero and the lower detection limit. For example, for cadmium, the detection

limit was 5 ng/L and we set values below the detection limit to 2.5 ng/L. While this approach has

some limitations (63, 64) compared to alternative approaches using imputation (65), we could

not apply imputation approaches to our data given the possibility of systematic differences in

concentrations of many of the metals between rivers. However, we explored the potential

impacts of this analysis choice on our conclusions by repeating PCAs with the measurements

below the detection limit set to the detection limit or zero, and found our results were

qualitatively unchanged (Fig. S4).

We also used the PCA analysis to qualitatively compare patterns within and across rivers

and identify chemicals of interest for further statistical analysis. Specifically, we chose chemicals

which had negative loadings on both PC1 and PC2 in either the non-metal or metal PCA because

we found that this approach clearly separated sites downstream on the Río Calnali from the

others (Fig. 4). From this set of parameters, we next investigated which had evidence for

affecting fish neurobiology, olfaction, or mate choice in the literature. These parameters included

DOM (23), turbidity (66), nitrite (67), ammonia (68, 69), copper (70, 71), lead (72), cadmium

(73, 74), zinc (75), and aluminum (76). In the analysis of ICP-MS metals data, where the effects

of samples below the detection limit prevented the use of parametric tests, we used Kruskal-

Wallis tests to identify differences in concentration within and between drainages. For all other

variables, we used ANOVA and Tukey's HSD tests to confirm differences between the

downstream Calnali sites and the other groups, testing for differences in the log-transformed

response variables as a function of drainage, sampling month, and their interaction. We also used

the same approach to explore differences in water chemistry between watersheds (Supp. Info 3;

Fig. S5-6) and over different years across the sampling period (Supp. Info 4; Fig. S7-9).

Finally, we tested whether broad-scale variation in water chemistry was associated with

variation in land use along the rivers. We created a combined dataset of all parameters described

above, calculated the site-wise average of each water quality parameter across all measurements,

and then constructed a PCA ($n = 30$ sites). We found that the variation in the PCA largely

mirrored our prior analyses on subsets of the data discussed above (Fig. S10). We then conducted

two separate linear regressions with the top two principal components of water chemistry

variation as response variables, each as a function of five explanatory variables: the site-specific

and cumulative proportion of developed land upstream of the site, the site-specific and

cumulative proportion of non-forest vegetation upstream of the site, and the total watershed area.

We repeated our analyses considering only land use within 50 m of the closest point of the river,

and also evaluated the effects of using different approaches to account for detection limits in the

ICP-MS measurements (setting measurements to the detection limit, zero, or a random uniform

variable drawn between zero and the detection limit; Table S10-12).

Analysis of ancestry assortative mating in Calnali Low

Previous work has shown that the vast majority of individuals above the town of Calnali

on the Río Calnali fall into one of two ancestry clusters: a *X. birchmanni* biased cluster that

derived ~75% of its genome from *X. birchmanni* and a *X. malinche* biased cluster that derives

>90% of its genome from *X. malinche*. These distinct ancestry clusters are maintained by strong

ancestry-assortative mating (55). In this study, we identify the Calnali Low population as one of

the first sampling sites where this genetic structure breaks down along the Río Calnali. As a

result, we were interested in evaluating levels of ancestry assortative mating in this population.

To do so, we analyzed mother-embryo samples from the Calnali Low population,

including previously published samples collected in 2020 (47), as well as newly sequenced

samples from a 2017 collection. Because swordtails are livebearing fish which exhibit multiple

paternity, when we collect pregnant *X. malinche* × *X. birchmanni* females we also collect ~10

offspring from an average of three fathers. Although we cannot directly assay the paternal

genotypes, since the maternal genotype can be directly determined, we can use the ancestry of

the offspring to infer the ancestry of the male the female mated with, since offspring will inherit

approximately 50% of their genome from each parent. Thus, the difference between the mother's

genome-wide ancestry and the genome-wide ancestry of her offspring is a useful proxy for

paternal ancestry.

We have previously used this approach to quantify ancestry assortative mating in

upstream populations on the Río Calnali (55), and repeat this approach here. Specifically, we ask

about the observed difference between maternal and offspring ancestry genome-wide in the

Calnali Low population, compared to expectations under a scenario where females mate at

random with respect to male ancestry. To generate a distribution of offspring ancestry expected

under no assortative mating, we performed simulations. For each focal female in our dataset ($n =$

74), we randomly sampled a mate from our full dataset of 702 adults from the Calnali Low

population. Next, we calculated the mean of the maternal and simulated paternal ancestry. We

added noise to this average ancestry by sampling from a random normal distribution with the

mean set to the variance in admixture proportion among full siblings observed in our data (Supp.

Info 6). We repeated this procedure until we had simulated offspring for all of the focal females.

We generated 1,000 replicate simulations and compared the observed and simulated differences

in mother-offspring admixture proportions in the Calnali Low population.

Inferring the strength of assortative mating from observed maternal-offspring ancestry

distributions

393   We found evidence that the observed mother-offspring ancestry patterns in the Calnali

Low population differed subtly from patterns expected if mating were truly random with respect

to ancestry (Fig. 5B). However, this difference from random mating was much weaker than

observed in upstream populations on the Río Calnali. To quantitatively compare the strength of

ancestry-assortative mating consistent with the observed data, we used the same approach we

previously used to quantify the strength of ancestry assortative mating in the Aguazarca hybrid

population (55).

400   Briefly, we used an approximate Bayesian computation approach to evaluate what levels

of ancestry-assortative mating were consistent with our observed data. We drew a probability that

a female would accept a mate with an ancestry proportion that differed from hers by more than

5% (see Supp. Info 6 for simulations using different thresholds) from a random uniform

distribution ranging from 0 (perfect assortative mating) to 1 (random mating). We refer to this

value as the strength of assortative mating. For each female we then randomly drew a mate from

the population ancestry distribution. If the mate fell within the ancestry threshold, the mate was

accepted. If it did not, we used a random binomial weighted by the strength of assortative mating

to determine if the mate was accepted. If the mate was rejected, we continued this procedure

until the female was matched with a mate, then simulated offspring as described above. This was

repeated until all females were matched with a mate. We repeated this simulation procedure for a

total of 100,000 simulations.

We used a simple rejection sampling approach to select simulations that closely matched

the empirical data. For each simulation, we calculated the mean and variance of the difference in

ancestry between mothers and their offspring. If both summary statistics fell within 5% of the

observed value for the Calnali Low population (mean 6.6%, variance 0.46%), we accepted the

simulation. This allows us to estimate the range of values of assortative mating consistent with

our data. We plotted the posterior distribution of accepted assortative mating strengths and

identified the maximum a posteriori (MAP) estimate. We repeated this procedure with other

thresholds of ancestry proportion differences (see Supp. Info 6; Fig. S11). To evaluate whether

results were affected by our approach in modeling assortative mating, we also explored an

alternative approach modelling assortative mating as a continuous beta function (Supp. Info 7;

Fig S12).

Histology in wild-caught fish

Like their relatives (77), *X. malinche × birchmanni* hybrids have plate-like olfactory

rosettes with beds of sensory epithelia surrounded by ridges of nonsensory tissue (Fig. S13).

Sensory epithelial beds contain a combination of ciliated and microvillous olfactory neurons,

which collect and transmit chemical information to the olfactory bulb, and ciliated sustentacular

cells, which increase flow over the beds to heighten olfactory sensitivity (78, 79). The

intervening nonsensory ridges are composed largely of connective tissue and goblet cells, the

latter of which are responsible for producing the glycoprotein-based mucus which physically

shields the epithelium from irritants (77, 80). Based on reports from the literature (70, 80–83),

we quantified the loss of cilia from the sensory epithelial surface and the proliferation of goblet

cells in the nonsensory epithelium and treated these two measures as a proxy for olfactory

disruption.

To understand whether polluted water was directly impacting the sensory periphery of

fish in *X. malinche* × *birchmanni* populations, we collected a total of 30 fish each from the

Aguazarca and Calnali Low populations in May (n = 10 per site) and September (n = 20) 2022.

Fishes were euthanized by MS-222 overdose and decapitation, after which heads were fixed in

10% buffered formalin for at least 48 hours. Fixed crania were dissected to remove the gills and

lower jaw, decalcified for 8 hours in 1X PBS with 10% formic acid, then dehydrated for 20

minutes each in 30%, 50%, and 70% ethanol before long-term storage in 70% ethanol. Samples

were submitted to Histo-Tec Laboratories (Hayward, CA) for xylene clearing, paraffin

embedding, sectioning, and staining. Horizontal sections were taken at 5 µm intervals with a

rotary microtome after removing 75-150 µm of tissue from the dorsal surface, with deeper

sectioning needed to reach the olfactory epithelia in larger individuals. One slide per individual

was prepared with hematoxylin and eosin (H&E) staining to observe gross morphological

changes (i.e. presence/absence of ciliary on olfactory epithelium). Samples from one collection

(May 2022) were also subject to Alcian Blue / Periodic Acid-Schiff stain for mucoproteins to

identify mucus-producing goblet cells, with eosin as a counterstain. Individuals for whom

olfactory rosettes were not present in the Histo-Tec slides (3 from Aguazarca and 1 from Calnali

Low for H&E slides, 1 each for AB/PAS slides) were excluded from further analysis.

A blinded observer examined the slides at 1000X magnification using oil immersion. The

observer defined the bounds of each sensory epithelial bed, and estimated the proportion of the

surface area of each bed on which cilia were visible to the nearest 10%. Further, we scanned and

imaged each slide at 40x magnification via a Biotek Cytation5 microplate reader, utilizing the in-

place ruler on the image to calculate the length of each bed in ImageJ. We normalized ciliated

proportions by bed length to calculate the total proportion of the sensory epithelium with visible

cilia in each individual, then used a Mann-Whitney U test to test for differences in ciliated

surface proportion between fish from Aguazarca and Calnali Low. Kolmogorov-Smirnov tests

indicated that the ciliated surface proportions were not significantly different between the two

sampling months ($P > 0.05$), and that other potentially confounding technical factors (number of

sensory beds, individual bed length, and total bed length) also did not differ significantly

between the two populations or sampling months ($P > 0.05$). To estimate the density of goblet

cells, a blinded observer counted all cells positive for AB/PAS staining within a bed, and we

tested for a significant difference between populations in the number of cells per individual,

corrected for total rosette length, using Mann-Whitney U tests. We also performed chemical

treatment trials in laboratory settings to explore whether we could induce similar patterns

through short-term chemical exposure (Supp. Info 8; Fig. S14-15).

Results

Population structure across rivers

Ancestry structure varied markedly among the four rivers sampled for this study. One

river, the Pochula, showed a relatively smooth cline in ancestry from pure *X. malinche* at high

elevations to *X. birchmanni*-like hybrids at lower elevations. The other three rivers varied in the

degree of population structure, with at least one site in the Calnali, Huazalingo, and Conzintla

showing evidence of significant departures from unimodality based on the distribution of

individual ancestry proportions (Fig. 2; Hartigan's Dip Test *P < 0.0001*). Examination of the

variation in individual ancestry in these populations supported the presence of two

subpopulations with distinct genome-wide ancestry.

Though we identify subpopulations of different ancestry in each river, we infer that the

drivers of these patterns differ dramatically. On the Río Huazalingo, the presence of two

subpopulations was only detected at a site directly downstream of a waterfall separating pure *X.*

malinche from admixed *X. birchmanni*. At this site, pure *X. malinche* individuals were sometimes

detected downstream of the waterfall, but this pattern varied between sampling seasons, hinting

that the *X. malinche* individuals may not persist in this population (Supp. Info 2; Fig. S2). In the

Río Conzintla, we observed sympatry between nearly-pure *X. malinche* and *X. birchmanni*; no

recent-generation hybrids were detected at this site. Downstream of these sympatric sites, the *X.*

malinche cluster disappears and leaves a single *X. birchmanni* ancestry cluster (Fig. 2; Table S5).

The ancestry fraction in this *X. birchmanni* cluster was stable across all sites upstream of a point

where the river passed underground (Kolmogorov-Smirnov Test $P > 0.05$), suggesting that there

is little ongoing gene flow between species in this drainage.

By contrast, in the Río Calnali, we observe a dramatic collapse in population structure

centered on the town of Calnali (Fig. 2). Sites upstream of the town, spanning 3.15 river

kilometers, show strong departure from unimodality in their ancestry distributions ($D = 0.12 -$

0.21, $P < 0.00001$ for all sites; Table S5). *X. malinche*-like and *X. birchmanni*-like subpopulations

co-exist in sympatry across these sites, but sites downstream of the town plaza show evidence of

a unimodal distribution ($D = 0.01 - 0.07$, $P > 0.18$ across all sites; Table S5) with intermediate

mean ancestry and increased variance in individual ancestry proportion. Intriguingly, average *X.*

malinche ancestry fraction increased downstream of the town of Calnali (Fig. 2), in contrast to

expectations given the geographical distribution of *X. malinche* populations in other rivers. To

further investigate this pattern, we collected samples from tributaries flowing into this section of

the Calnali, and concluded that migration of individuals with high *X. malinche* ancestry from

tributaries into the main river is a plausible explanation for this observed shift in ancestry (Supp.

Info 9; Fig. S18).

Land use across drainages

Land use quantification showed that human impact on the Río Calnali watershed was

indeed higher than other rivers. The reaches where the river passed through the town of Calnali

had the highest proportion of developed land in the study area, with unparalleled levels of

urbanization compared to surrounding areas. This increased level of urbanization was centered

on the location of genetic population structure collapse in the town of Calnali (Fig. 3). When

calculating the cumulative watershed land input of a given point, the downstream sites on the Río

Calnali had the highest proportions of developed land in the study area (Fig 3B), especially when

considering land within a short distance of the river (Fig. 3E; Fig. S19H). In contrast, we

observed no similar association of cleared land (vegetation, with the exclusion of forest) with

population structure on the Río Calnali for any given stream reach or at the watershed scale.

When compared to other rivers, though, the Río Calnali did show higher levels of cumulative

cleared area within a short distance of the river (Fig. S20F, S20H). Overall, our findings suggest

that disruption of forest is ubiquitous throughout the Atlapexco drainage, including in rivers with

low levels of hybridization. By contrast, the level of built infrastructure is especially high in the

Calnali subcatchment.

Water chemistry variation aligns with breakdown of population structure

Water chemistry varied dramatically both between rivers and between sites on the same

river. Principal component analysis demonstrated that the majority of variation in water quality

could be summarized by two axes: one that largely separates the sites by river (PC1), and another

separating the sites upstream of the town of Calnali from the other rivers and those downstream

of Calnali (PC2; Fig. 4). Sites downstream of the town of Calnali were associated with increased

dissolved organic matter, turbidity, ammonia, and nitrite, as well as reduced dissolved oxygen, all

consistent with increased anthropogenic impacts (Fig. 4). Statistical tests indicated that levels of

turbidity, DOM, nitrite, and ammonia were all significantly elevated in the sites downstream on

the Río Calnali relative to the upstream sites and to other rivers (Fig. 4, Supp. Fig S16; Table S7-

S8). Across rivers, there was variation in total conductivity, alkalinity, hardness, and pH, with sites in the Río Conzintla having the highest values of each of these metrics and sites along the Río Calnali having the lowest (Fig. 4; Fig. S22).

High concentrations of a range of metals have been previously linked to breakdowns in neurological or olfactory function in fish (27). Given the potential relationship between such breakdown and changes in ancestry assortative mating, we were interested in whether any metals were elevated in waters downstream of the town of Calnali, but not in other sites or across rivers. A principal component analysis showed that concentrations of many dissolved ions, measured by ICP-MS, were highest in the Conzintla and lowest in the Calnali (Fig. 4; Fig. S24). However, the transition from the sites upstream to downstream of the town of Calnali was also associated with a dramatic increase in the concentration of numerous ions, especially major industrial metals (e.g. iron, aluminum, copper, lead, manganese; Fig. 4). Multiple metals significantly increased in concentration downstream of the town of Calnali when individually tested, including copper, lead, cadmium, aluminum, and zinc (Kruskal Wallis $P < 0.001$; Fig. 4; Fig. S23). On the other hand, other ions with known effects on fish neurobiology or olfaction showed similar or higher concentration in the Conzintla than in the Calnali downstream sites, including manganese, arsenic, and calcium (Supp. Info 3; Fig. S24). Though we lack sufficient sampling to formally analyze these effects, we also found some evidence for temporal fluctuations in metal concentrations across rivers (Supp. Info 4; Fig. S7-9).

Finally, to quantify whether changes in water chemistry parameters were connected to land use change, we independently modelled the first two principal components of water chemistry variation as a function of both the site-wise and cumulative area of developed and cleared land, accounting for cumulative watershed area as a covariate. We found that PC1 was

strongly correlated with the cumulative area of both developed ($P < 0.001$) and cleared land ($P <$

0.001), while PC2 was strongly associated with the proportion of developed ($P = 0.032$) and

cleared ($P = 0.002$) land immediately upstream of a site. We found that limiting our analysis to

land in close proximity to the river increased the relative importance of developed land as a

predictor of both PC1 and PC2. In general, our results were not sensitive to various approaches

to analyzing ICP-MS values below the detection limit (Table S9-S12). We thus concluded that

the proportion of built infrastructure in the watershed was the strongest predictor of water

chemistry variation, with the most dramatic impacts in the Río Calnali downstream of the town

of Calnali.

Strength of ancestry-assortative mating in the Calnali Low and Aguazarca hybrid populations

In the upstream hybrid population on the Río Calnali, the Aguazarca population, we

previously documented bimodal ancestry structure (55). This strong ancestry structure appears to

be driven by nearly complete assortative mating by ancestry type in this population (55). By

contrast, the Calnali Low population does not show evidence of bimodal ancestry structure,

although variance in ancestry is high, potentially indicative of recent admixture between the *X.*

birchmanni and *X. malinche* ancestry clusters (Fig. 2). We used paired mother-embryo

sequencing to evaluate whether mating patterns differed from expectations under random mating

with respect to ancestry. Intriguingly, despite the dramatic breakdown in population structure, we

found that mothers and offspring were less different in ancestry than expected under simulations

of truly random mating (Fig. 5A). However, the average difference in mother-offspring ancestry

was significantly greater in the Calnali Low hybrid population (0.0655) than the prior estimate

from the Aguazarca population (0.022; *P < 0.0001* by simulation), hinting that assortative mating

by ancestry is substantially weaker in the Calnali Low hybrid population.

Given this result, we sought to quantify the strength of ancestry assortative mating in the

Calnali Low population and compare it to that observed in the upstream Aguazarca population.

We used an approximate Bayesian computation approach to ask what strength of ancestry

assortative mating was consistent with the average difference in ancestry between mothers and

offspring that we observe in the Calnali Low hybrid population (see Methods). Based on

accepted simulations, we recovered a well-resolved posterior distribution of the strength of

ancestry assortative mating, and estimated the maximum a posteriori (MAP) to be 0.65 (Fig. 5B).

This is substantially weaker than estimates from the Aguazarca population, in which estimates of

assortative mating by ancestry approach 1 (55). Thus, our results suggest that weaker assortative

mating contributes to the breakdown in population structure in the Calnali Low hybrid

population.

Olfactory disruption as a function of water chemistry

Changes in olfaction can affect mate discrimination in *Xiphophorus (23, 26)*, and we

found that fish in heavily impacted sites on the Río Calnali have been exposed to high levels of a

range of compounds that may impact olfactory function. We therefore evaluated evidence for

physical damage of the olfactory rosettes of wild caught fish from different sites on the Río

Calnali, and found that fish collected from the upstream site, Aguazarca, and the downstream

site, Calnali Low, showed significant changes in olfactory morphology. H&E staining allowed us

to observe the motile and sensory cilia which may be lost in the face of chemical exposure (70,

81, 82). We found marginally lower proportions of ciliated surface area on the olfactory

epithelium at the Calnali Low site (Mann-Whitney Test P-value = 0.08; Fig. 5C). Likewise,

AB/PAS staining clearly distinguished goblet cells through the deep blue staining of the

protective mucoproteins they produce, and revealed significantly higher densities of goblet cells

at Calnali Low (Mann-Whitney Test P-value = 0.002; Fig. 5D-F), consistent with a

compensatory response to shield the olfactory epithelium from toxicants.

Discussion

A more nuanced view of the genetic and environmental landscape of hybrid zones will allow biologists to move past a binary picture of hybridization's presence or absence, and begin to pinpoint the specific changes and mechanisms by which anthropogenic disturbance can drive hybridization. By surveying genome-wide ancestry distributions at dozens of sites where *X. birchmanni* and *X. malinche* coexist, we found that the patterns of hybridization varied drastically within and between rivers. We detected some degree of gene flow between *X. malinche* and *X. birchmanni* in all four rivers surveyed, but the amount and spatial distribution of admixture differed across each river (Fig. 2, Supp. Info 9; Fig. S16-17). Thus, while there is some potential for hybridization whenever related species overlap, the implications for biodiversity are highly variable.

At one end of the spectrum of genetic homogenization, the Río Conzintla showed two slightly admixed ancestry clusters which co-occur at multiple sites across 3.7 river kilometers, resulting in a "bimodal" distribution of genome-wide ancestry at these sites. Within these sites, individuals derived <15% of their genome-wide ancestry from the other species (mean ± SD "minor" parent ancestry of 5.6 ± 3.1% at Xochicoatlán, 10.6 ± 4.5% at Mixtla). While this level of admixture is evolutionarily significant, its functional consequences are likely limited by genome-wide selection against introgression (49), and populations with similar admixture rates have been designated as "conservation populations" in other cases of anthropogenic hybridization (84). The stability of within-cluster admixture proportions across the entire length of the Río Conzintla, combined with our observation of only 1 recent hybrid out of 239 individuals, suggests that gene flow between clusters in this river is extremely limited. The stable

hybrid zone in this stream thus serves as a contrast to other rivers, and comparisons in water

quality may illuminate the potential drivers of breakdown in assortative mating in other contexts.

In contrast, the Huazalingo and Pochula rivers showed no sites with bimodal ancestry

structure, but rather a sharp transition in genome-wide ancestry moving downstream. These

rivers appear more similar to classic clinal hybrid zones, where ancestry patterns are shaped by

the countervailing effects of gene flow and selection against hybrids (85). While the lack of any

structured populations in these rivers prevented us from assessing the effects of environmental

variation on rates of hybridization, we found that overall water chemistry in these rivers was

quite similar to the Río Conzintla. We speculate that anthropogenic impacts may have played a

role in the onset of hybridization in these rivers, but it is difficult to test such hypotheses in

contemporary hybrid populations, since the weakening of mate preference in highly admixed

individuals means that such clines could continue to exist even after environmental triggers

disappear (86–88). These rivers nonetheless can provide important insights into the role of

migration and ecological selection in structuring *X. malinche × X. birchmanni* hybrid zones

(Supp. Info 9).

Lastly, the Río Calnali showed a dramatic change in ancestry structure from upstream to

downstream. We observed strong bimodality in genome-wide ancestry at sites upstream of the

town of Calnali, with individuals falling into one of two subpopulations, a *X. malinche*-like

ancestry cluster and a *X. birchmanni*-like cluster. These results align with prior work

documenting the presence of strong assortative mating by ancestry within *X. malinche* and *X.*

birchmanni-like hybrid clusters at the Aguazarca site on the Río Calnali (Supp. Info 6) (55).

Within a span of less than two river kilometers where the river traverses the town of Calnali, this

bimodal population structure collapses into a hybrid swarm, with the range of observed

admixture proportions spanning the continuum between the upstream clusters (Fig. 2). The Río

Calnali thus provides a powerful opportunity to study the environmental determinants of

reproductive isolation through comparisons within and between rivers.

We took advantage of this remarkable system by measuring a large array of chemical

parameters across rivers in parallel with our genomic surveys. In doing so, we identified

substantial chemical variation both between rivers of the Atlapexco drainage and between sites

upstream and downstream of the town of Calnali (Fig. 4). PCA highlighted river-specific water

chemistry signatures, which we speculate may be driven by geology and the impacts of mining in

the Conzintla drainage (see Supp. Info 3; Fig. S5). However, we also observe an independent

chemical change among sites on the Río Calnali above and below the town of Calnali, while the

other rivers were relatively stable in water chemistry along their lengths (Fig. 4). By quantifying

land use across each drainage, we confirmed that the Calnali drainage is indeed unique both in

the scale of built infrastructure relative to catchment area, as well as the proximity of

development to the river (Fig. 3, Fig. S19). The clearing of forests for pasture and cropland could

also contribute to the chemical disturbances we observe, but may be a less likely explanation

given the relatively small differences in agricultural land areas within and between drainages

(Fig. S20), as well as the low-intensity agricultural practices typical of the area (89, 90). Both

agricultural and urban development have been linked to some of the changes in water chemistry

that we observe in the downstream Río Calnali (91–95). In particular, many metals whose

concentrations change along the course of the Calnali have been previously associated with both

domestic wastewater and runoff from roads (93).

Among the many compounds which varied between rivers and disturbance regimes, a

substantial number have been previously shown to affect olfaction in fishes (dissolved organic

matter, copper, aluminum, zinc (23, 96, 97)) or fish neurology more generally (ammonia, cobalt,

and cadmium (68, 74, 98)). Many of these parameters were specifically elevated in the Río

Calnali, and especially so in the areas downstream of the town. Thus, they are particularly

promising candidates for the drivers of ancestry structure breakdown. On the other hand, we

found that other ions with olfactory or neurological effects in the ecotoxicology literature

(manganese, arsenic, calcium (26, 99–101)) were most extreme in the Conzintla, where

reproductive isolation is maintained (Fig. S22, Fig. S24), suggesting that variation in these

specific parameters is not a primary driver of hybridization in *Xiphophorus*. Variation in pH

between streams may also play a role, as low pH increases metal bioavailability (Fig. S22; (96)).

Having established differences in both the degree of population structure and in water

chemistry between sites in the Río Calnali, we next explored the loss of olfaction-based

assortative mate preferences as a biological mechanism which might connect the two. Mother-

embryo sequencing and simulations demonstrate that assortative mating is substantially

weakened at downstream sites on the Río Calnali, but surprisingly, it is not entirely absent (Fig.

5B). Ongoing assortative mating could be driven by downstream migration, with mothers either

mating prior to migration, or retaining some ability to mate assortatively even after arriving in

highly polluted waters. Alternatively, weak assortative mating at the population scale could be

produced by differential vulnerability of various signal modalities to disturbance (e.g. visual vs.

olfactory cues, (102, 103)) or fluctuations in the degree of sensory disturbance fish experience as

a result of variation in environmental conditions (Supp. Info 4). Regardless of the mechanism,

this observation supports theoretical predictions that even a partial breakdown in assortative

mating may result in the emergence of hybrid swarms (28).

Finally, our histological sampling of fishes in the upstream and downstream Calnali shows changes in cellular abundance and morphology consistent with olfactory damage. The loss of cilia in the sensory epithelium has previously been observed after exposure to heavy metals and anthropogenic hydrocarbons (78, 104), and could directly reduce olfactory sensitivity by disrupting water flow through the nares (105, 106) or the function of olfactory sensory neurons (107). Likewise, an increased number of goblet cells has been linked to inflammation secondary to nasal irritation in other systems, and may be a protective response to noxious chemical exposure (80, 83). These histological analyses represent the first evidence of higher olfactory damage in a wild hybrid swarm relative to nearby structured populations. Whether these changes themselves weaken assortative mating, or are correlated with other causative changes (e.g. transport and accumulation of heavy metals in the brain's olfactory bulb (108)) is an exciting direction for future work. While our results support the hypothesis that the breakdown of assortative mating along the Río Calnali is mediated at least in part by olfactory dysfunction, our short-term laboratory exposures of *X. birchmanni* to ecologically relevant levels of ammonia, copper, humic acid, or their combination did not reproduce the effect (Fig. S15). The specific chemical concentrations and exposure durations necessary to disrupt pre-mating isolation thus remains a promising area for further study.

Taken as a whole, our study presents an integrative window into the environmental and organismal drivers of hybridization between *X. birchmanni* and *X. malinche*. By showing that dramatic water chemistry differences are tightly linked to changes in population structure, we confirm the relevance of prior behavioral work showing the potential for changes in water chemistry to disrupt assortative mating in these species (23, 26). Though many studies have documented a connection between environmental conditions and hybridization in freshwater

fishes (3, 27), our analyses of assortative mating and olfactory histology provide a deeper

understanding of the proximal mechanisms by which water chemistry changes might disrupt

reproductive isolation. This multidimensional approach to the study of hybrid zones sheds light

on the routes by which anthropogenic change can impact biodiversity on small spatiotemporal

scales. As human impacts on the environment continue to expand, a greater understanding of the

drivers of hybridization will help to mitigate human impacts on biodiversity.

Figure 1. Study area and sampling sites. **A)** Overview of Río Atlapexco basin, highlighted in orange within the inset Mexican state map, whose area is indicated in yellow on the gray outline of Mexico. Rivers are shown as blue lines, the four major rivers studied are labelled, sites where fish were collected are yellow circles, and sites where water chemistry data were collected are orange diamonds. The black box encloses the area displayed in panel **B**. **B)** Sites sampled on Río

Calnali. Thick gray lines show major roadways, and thin black lines show streets in the town of

Calnali. Images show representative male *Xiphophorus* fishes collected from two populations on

the Río Calnali (AGZC = Aguazarca, CALL = Calnali Low). These sites were also the locations

used for sample collection for histology. In the Río Calnali, *X. malinche*-like fish are found in the

headwaters, *X. malinche*-like and *X. birchmanni*-like fishes are found upstream of the town of

Calnali, and fish with intermediate genotypes and phenotypes are found downstream of town.

Figure 2. Genetic ancestry results from *X. malinche* × *X. birchmanni* hybrids in the Río

Atlapexco Drainage. Histograms show ancestry distributions at sampling locations along four

rivers in the Río Atlapexco drainage. Sites are arranged with upstream sites on the left side of the

X-axis, and downstream sites on the right. Y-axis corresponds to the proportion of genome-wide

ancestry derived from *X. malinche* among individuals in that populations (1 = pure *X. malinche*,

0 = pure *X. birchmanni*), and the width of bars indicates the frequency of individuals in that

ancestry bin. Building icon on the Río Calnali transect denotes the sites located within the town

of Calnali. X-axis text shows site numbers, as in Fig. 1. Bolded X-axis numbers indicate sites

with significant deviations from a unimodal ancestry distribution, either across all sampling

timepoints (asterisk) or at a subset of sampling times (dagger, Supp. Info. 2).

Figure 3. Results of land use classification analysis in the Río Atlapexco basin. **A**) Map of

classification results from 2020-2023 hyperspectral imagery. 10 m² pixels were classified as

developed (buildings, mines, and roads), cleared land (grassland, pasture, or cultivated crops)

forest (continuous tree cover, excluding plantation-style agriculture), or water by an SVM

algorithm, with training polygons identified from 3 m^2-resolution 2022 imagery from Planet

Labs (61). River polygons are shown as blue lines over classified pixels. **B-E)** Percent of land

classified as developed within different drainages of the Río Atlapexco basin. Points show the

proportion of pixels in each site's subcatchment that were classified as developed. Left column

(**B** and **D**) shows the site-wise proportions, or the subcatchment of the section of river between a

site and the closest upstream site. Right column (**C** and **E**) shows the cumulative subcatchment

of all sections of the river upstream of a given site. Top row (**B** and **C**) shows calculations that

include pixels of any distance from a stream, while bottom row (**D** and **E**) shows calculations

limited to pixels within 50 m of a stream (defined as pixels with accumulated flow > 3000 cells,

depicted with river polygons in **A**). Color denotes the drainage that the site occurs in, and lines

connect sites in the same river based on their order. The line corresponding to the Río Calnali

branches because multiple headwater tributaries were sampled.

Figure 4. Water chemistry data from four rivers in the Río Atlapexco drainage. **A-B)** Principal

component analysis (PCA) biplot for **A)** parameters measured using the sonde, turbidimeter,

colorimeter, and **B)** parameters measured using ICP-MS. Points represent sampling events, and

color and shape denote drainage, with sites on the Río Calnali upstream and downstream of the

town of Calnali labelled separately. Labelled arrows represent variable loadings onto first two

principal components, with highly correlated variables showing similar vectors. **C-E)** Plots of the

change in water chemistry over the course of the river, focusing on parameters that were elevated

in sites downstream of the town of Calnali. These include **C)** fluorescent dissolved organic

matter, **D)** ammonia, and **E)** total copper. The building icon on the X-axis denotes the sites

located within the town of Calnali on the Río Calnali transect. X-axis indicates distance

downstream from the first site sampled on a river, points indicate individual measurements, and

lines show a loess-spline fit of change over the river course. Note that the Y-axis is log-scaled in

787   the case of ammonia measurements (**D**). Translucent points below the gray dashed line in **E)**

788   represent measured values below the detection limit of the ICP-MS assay. Points below this limit

789   are plotted at half the detection threshold, with jittering to prevent overlap and represent sample

790   size, but do not represent point estimates of the concentration.

Figure 5. Mating patterns and histological phenotypes in fishes upstream and downstream of the

town of Calnali. **A)** Results of mother-embryo sequencing analysis based on individuals

collected at the Calnali Low site, downstream of the town of Calnali. X-axis shows the

proportion of each mother's genome derived from *X. malinche*, and Y-axis shows the difference

in genome-wide *X. malinche* ancestry between each mother and embryo. Mean maternal-

offspring ancestry difference was significantly but modestly lower in real data (black) than in

simulated data modeling random mating (blue; *P < 0.001* by simulation). **B)** Inferred strength of

assortative mating from observed maternal-offspring ancestry distributions. Blue bars depict the

simulated probability of assortative mating in accepted simulations, with 0 representing random

mating. Dashed line denotes maximum a posteriori estimate ($P_{assortative} = 0.65$). **C-D)** Results of

olfactory histology of fishes from sites upstream (Aguazarca) and downstream (Calnali Low) of

the town of Calnali. **C)** Proportion of each fish's sensory epithelium with visible cilia, visualized

with H&E staining. **D)** Number of mucus-producing goblet cells per μm of olfactory rosette

surface area, visualized by staining with AB/PAS. *P*-values indicate results of Mann-Whitney *U*

Test for difference between sites. **E-F)** Representative AB/PAS stains of olfactory rosettes of *X.*

malinche × birchmanni hybrids from **E)** Aguazarca, and **F)** Calnali Low. Mucoproteins are

stained blue by AB/PAS, eosin counterstain appears as pink. Arrowheads indicate locations of

goblet cells.

Acknowledgements

The authors thank members of the Schumer and Rosenthal laboratories for helpful discussion and/or feedback on earlier versions of this manuscript. We thank the Federal Government of Mexico for permission to collect fish (Permiso de Pesca de Fomento no. PPF/DGOPA-064/20). Stanford University and the Stanford Research Computing Center provided computational support for this project. We thank Dra. Susana Saval Bohórquez of the UNAM Environmental Engineering Laboratory and Dr. Guangchao Li and Douglas Turner of the Stanford Environmental Measurements Facility for technical and experimental support. This work was supported by a Knight–Hennessy Scholars fellowship and NSF GRFP 2019273798 to B.M.M., a CONACyT fellowship to GIJ, NSF Awards 1723266, 1755327 and 1354172 to GGR, and Human Frontiers in Science Grant RGY0081 to MS & CMR.

References

1. E. Mayr, *Systematics and the Origin of Species: from the Viewpoint of a Zoologist* (Columbia

University Press, 1942).

2. E. Anderson, Hybridization of the habitat. *Evolution* 1–9 (1948).

3. K. C. Grabenstein, S. A. Taylor, Breaking Barriers: Causes, Consequences, and Experimental

Utility of Human-Mediated Hybridization. *Trends in Ecology & Evolution* **33**, 198–212

(2018).

4. R. Ubsdell, Studies on variation and evolution in Centaurium erythraea Rafn and C. littorale

(D. Turner) Gilmour in the British Isles. 3. Breeding systems, floral biology and general

discussion. *Watsonia* **12**, 225–232 (1979).

5. C. Brochmann, Hybridization and distribution of Argyranthemum coronopifolium

(Asteraceae – Anthemideae) in the Canary Islands. *Nordic Journal of Botany* **4**, 729–736

(1984).

6. O. Seehausen, J. J. M. van Alphen, F. Witte, Cichlid Fish Diversity Threatened by

Eutrophication That Curbs Sexual Selection. *Science* **277**, 1808–1811 (1997).

7. P. Vonlanthen, *et al.*, Eutrophication causes speciation reversal in whitefish adaptive

radiations. *Nature* **482**, 357–362 (2012).

8. E. C. Ellis, Anthropogenic transformation of the terrestrial biosphere. *Philosophical

Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*

369, 1010–1035 (2011).

9.   J. A. Coyne, H. A. Orr, *Speciation* (Sinauer Associates, 2004).

10. D. A. Turissini, J. A. McGirr, S. S. Patel, J. R. David, D. R. Matute, The Rate of Evolution of

Postmating-Prezygotic Reproductive Isolation in Drosophila. *Molecular Biology and*

Evolution **35**, 312–334 (2018).

11. D. I. Bolnick, T. J. Near, Tempo of Hybrid Inviability in Centrarchid Fishes (ᵀᴱᴸeostei:

Centrarchidae). *Evolution* **59**, 1754–1767 (2005).

12. C. E. Wagner, L. J. Harmon, O. Seehausen, Ecological opportunity and sexual selection

together predict adaptive radiation. *Nature* **487**, 366–369 (2012).

13. O. Puebla, E. Bermingham, F. Guichard, Pairing dynamics and the origin of species.

Proceedings of the Royal Society of London B: Biological Sciences **279**, 1085–1092 (2012).

14. J.-G. J. Godin, S. E. Briggs, Female mate choice under predation risk in the guppy. *Animal*

Behaviour **51**, 117–130 (1996).

15. P. M. Willis, M. J. Ryan, G. G. Rosenthal, Encounter rates with conspecific males influence

female mate choice in a naturally hybridizing fish. *Behavioral Ecology* **22**, 1234–1240

(2011).

16. M. N. Verzijden, Z. W. Culumber, G. G. Rosenthal, Opposite effects of learning cause

asymmetric mate preferences in hybridizing species. *Behavioral Ecology* **23**, 1133–1139

(2012).

17. J. Hunt, R. Brooks, M. D. Jennions, Female Mate Choice as a Condition-Dependent Life-

History Trait. *The American Naturalist* **166**, 79–92 (2005).

18. A. E. Hausmann, *et al.*, Light environment influences mating behaviours during the early

stages of divergence in tropical butterflies. *Proceedings of the Royal Society B: Biological*

Sciences **288**, 20210157 (2021).

19. K. S. Pfennig, Facultative mate choice drives adaptive hybridization. *Science* **318**, 965–967

(2007).

20. G. G. Rosenthal, D. Stuart-Fox, others, "Environmental disturbance and animal

communication" in *Behavioural Responses to a Changing World: Mechanisms and*

Consequences, (Oxford University Press, 2015), pp. 16–31.

21. G. G. Rosenthal, Individual mating decisions and hybridization. *Journal of Evolutionary*

Biology **26**, 252–255 (2013).

22. B. Seifert, J. Kulmuni, P. Pamilo, Independent hybrid populations of Formica polyctena

Xrufa wood ants (Hymenoptera: Formicidae) abound under conditions of forest

fragmentation. *Evol Ecol* **24**, 1219–1237 (2010).

23. H. S. Fisher, B. B. M. Wong, G. G. Rosenthal, Alteration of the chemical environment

disrupts communication in a freshwater fish. *Proc R Soc B* **273**, 1187–1193 (2006).

24. J. L. Ward, M. J. Blum, Exposure to an environmental estrogen breaks down sexual isolation

between native and invasive species. *Evolutionary Applications* **5**, 901–912 (2012).

25. P. C. Hubbard, E. N. Barata, A. V. M. Canario, Possible disruption of pheromonal

communication by humic acid in the goldfish, Carassius auratus. *Aquatic Toxicology* **60**,

169–183 (2002).

26. D. L. Powell, A. D. Rose, G. G. Rosenthal, A widely-used pollutant causes reversal of

conspecific mate preference in a freshwater fish. [Preprint] (2022). Available at:

https://www.biorxiv.org/content/10.1101/2022.09.07.507014v1 [Accessed 2 September

2023].

27. W. F. Ramirez-Duarte, *et al.*, Hybridization in the Anthropocene – how pollution and climate

change disrupt mate selection in freshwater fish. *Biological Reviews* **0** (2024).

28. R. T. Gilman, J. E. Behm, Hybridization, Species Collapse, and Species Reemergence After

Disturbance to Premating Mechanisms of Reproductive Isolation. *Evolution* **65**, 2592–2605

(2011).

29. D. J. Hasselman, *et al.*, Human disturbance causes the formation of a hybrid swarm between

two naturally sympatric fish species. *Molecular Ecology* **23**, 1137–1152 (2014).

30. T. B. Malek, J. W. Boughman, I. Dworkin, C. L. Peichel, Admixture mapping of male nuptial

colour and body shape in a recently formed hybrid population of threespine stickleback.

Molecular Ecology **21**, 5265–5279 (2012).

31. B. Alric, *et al.*, Local human pressures influence gene flow in a hybridizing Daphnia species

complex. *Journal of Evolutionary Biology* **29**, 720–735 (2016).

32. T. van Hengstum, *et al.*, Human-induced hybridization among congeneric endemic plants on

Tenerife, Canary Islands. *Plant Syst Evol* **298**, 1119–1131 (2012).

33. J. I. Meier, *et al.*, Ancient hybridization fuels rapid cichlid fish adaptive radiations. *Nature*

Communications **8**, 14363 (2017).

34. S. A. Taylor, E. L. Larson, Insights from genomes into the evolutionary importance and
prevalence of hybridization in nature. *Nat Ecol Evol* **3**, 170–177 (2019).

35. R. J. Abbott, N. H. Barton, J. M. Good, Genomics of hybridization and its evolutionary
consequences. *Molecular Ecology* **25**, 2325–2332 (2016).

36. M. Schumer, R. Cui, G. G. Rosenthal, P. Andolfatto, Reproductive Isolation of Hybrid
Populations Driven by Genetic Incompatibilities. *PLoS Genet* **11**, e1005041 (2015).

37. C. Veller, N. B. Edelman, P. Muralidhar, M. A. Nowak, Recombination and selection against
introgressed DNA. *Evolution* **77**, 1131–1144 (2023).

38. C. Chen, K. S. Pfennig, Female toads engaging in adaptive hybridization prefer high-quality
heterospecifics as mates. *Science* **367**, 1377–1379 (2020).

39. N. B. Edelman, J. Mallet, Prevalence and Adaptive Impact of Introgression. *Annu Rev Genet*
55, 265–283 (2021).

40. M. Ribeiro Palacios, Land use change in Southern Huasteca, Mexico; drivers and
consequences for livelihood and ecosystem services. (2012).

41. H. R. Hernández, M. A. Robledo, J. R. A. Rivera, J. F. Martinez, Spatial Configuration of
Land-Use/Land-Cover in the Pujal-Coy Project Area, Huasteca Potosina Region, Mexico.
Ambio **37**, 381–389 (2008).

42. M. L. Martínez, *et al.*, Effects of land use change on biodiversity and ecosystem services in
tropical montane cloud forests of Mexico. *Forest Ecology and management* **258**, 1856–
1863 (2009).

43. L. Chapa-Vargas, K. Monzalvo-Santos, Natural protected areas of San Luis Potosí, Mexico:

ecological representativeness, risks, and conservation implications across scales.

International Journal of Geographical Information Science **26**, 1625–1641 (2012).

44. C. Payne, *et al.*, Genomic insights into variation in thermotolerance between hybridizing

swordtail fishes. *Mol Ecol* (2022). https://doi.org/10.1111/mec.16489.

45. Z. W. Culumber, D. B. Shepard, S. W. Coleman, G. G. Rosenthal, M. Tobler, Physiological

adaptation along environmental gradients and replicated hybrid zone structure in swordtails

(Teleostei: Xiphophorus). *Journal of Evolutionary Biology* **25**, 1800–1814 (2012).

46. D. L. Powell, *et al.*, Natural hybridization reveals incompatible alleles that cause melanoma

in swordtail fish. *Science* **368**, 731–736 (2020).

47. B. M. Moran, *et al.*, A lethal mitonuclear incompatibility in complex I of natural hybrids.

Nature **626**, 119–127 (2024).

48. M. Schumer, Y. Brandvain, Determining epistatic selection in admixed populations. *Mol Ecol*

25, 2577–2591 (2016).

49. M. Schumer, *et al.*, Natural selection interacts with recombination to shape the evolution of

hybrid genomes. *Science* **360**, 656–660 (2018).

50. Z. W. Culumber, *et al.*, Replicated hybrid zones of *Xiphophorus* swordtails along an

elevational gradient. *Mol Ecol* **20**, 342–356 (2011).

51. M. Rauchenberger, I. K. D. Kallman, Monophyly and Geography of the Rio P'anuco Basin

Swordtails (Genus Xiphophorus) with Descriptions of Four New Species. *AMERICAN*

MUSEUM NOVITATES 44 (1990).

52. G. G. Rosenthal, *et al.*, Dissolution of Sexual Signal Complexes in a Hybrid Zone between

the Swordtails Xiphophorus birchmanni and Xiphophorus malinche (Poeciliidae). *Copeia*

2003, 299–307 (2003).

53. INEGI, México en cifras. (1998). Available at:

https://www.inegi.org.mx/app/areasgeograficas/ [Accessed 6 April 2025].

54. M. Schumer, *et al.*, High-resolution mapping reveals hundreds of genetic incompatibilities in

hybridizing fish species. *eLife* **3**, e02535 (2014).

55. M. Schumer, *et al.*, Assortative mating and persistent reproductive isolation in hybrids. *Proc*

Natl Acad Sci USA **114**, 10936–10941 (2017).

56. M. Schumer, D. L. Powell, R. Corbett-Detig, Versatile simulations of admixture and accurate

local ancestry inference with mixnmatch and ancestryinfer. *Molecular Ecology Resources*

20, 1141–1151 (2020).

57. S. M. Banerjee, *et al.*, Complex hybridization between deeply diverged fish species in a

disturbed ecosystem. *Evolution* **77**, 995–1005 (2023).

58. J. A. Hartigan, P. M. Hartigan, The Dip Test of Unimodality. *The Annals of Statistics* **13**, 70–

84 (1985).

59. European Union/ESA/Copernicus, Sentinel-2 MSI: MultiSpectral Instrument, Level-2A.

(2020).

60. J. Transon, R. D'Andrimont, A. Maugnard, P. Defourny, Survey of Hyperspectral Earth

Observation Applications from Space in the Sentinel-2 Context. *Remote Sensing* **10**, 157

(2018).

61. Planet Labs, Planetscope Scene, March 29 2022, Instrument PS2.SD, Strip 5529184. (2022).

62. INEGI, Continuo de elevaciones mexicano y modelos digitales de elevación v3. (2013).

63. A. Kamal, K. Wichuk, D. McCartney, K. Londry, C. Felske, Effect of Treated Wood on

Biosolids Composting. *Compost Science & Utilization* **25**, 178–193 (2017).

64. K. J. Pieper, *et al.*, Evaluating Water Lead Levels During the Flint Water Crisis. *Environ. Sci.*

Technol. **52**, 8124–8132 (2018).

65. D. R. Helsel, *Nondetects and data analysis. Statistics for censored environmental data.*

(2005).

66. O. Seehausen, *et al.*, Speciation through sensory drive in cichlid fish. *Nature* **455**, 620–626

(2008).

67. J.-H. Kim, S. K. Kim, Y. B. Hur, Toxic effects of waterborne nitrite exposure on antioxidant

responses, acetylcholinesterase inhibition, and immune responses in olive flounders,

Paralichthys olivaceus, reared in bio-floc and seawater. *Fish & Shellfish Immunology* **97**,

581–586 (2020).

68. Y. K. Ip, S. F. Chew, D. J. Randall, "Ammonia toxicity, tolerance, and excretion" in *Fish*

Physiology, Nitrogen Excretion., (Academic Press, 2001), pp. 109–148.

69. D. J. Randall, T. K. N. Tsui, Ammonia toxicity in fish. *Marine Pollution Bulletin* **45**, 17–23

(2002).

70. J. A. Hansen, J. D. Rose, R. A. Jenkins, K. G. Gerow, H. L. Bergman, Chinook salmon

(Oncorhynchus tshawytscha) and rainbow trout (Oncorhynchus mykiss) exposed to copper:

Neurophysiological and histological effects on the olfactory system. *Environmental*

Toxicology and Chemistry **18**, 1979–1991 (1999).

71. W. A. Dew, A. Azizishirazi, G. G. Pyle, Contaminant-specific targeting of olfactory sensory

neuron classes: Connecting neuron class impairment with behavioural deficits.

Chemosphere **112**, 519–525 (2014).

72. J.-W. Lee, *et al.*, Toxic effects of lead exposure on bioaccumulation, oxidative stress,

neurotoxicity, and immune responses in fish: A review. *Environmental Toxicology and*

Pharmacology **68**, 101–108 (2019).

73. C. R. Williams, E. P. Gallagher, Effects of cadmium on olfactory mediated behaviors and

molecular biomarkers in coho salmon (Oncorhynchus kisutch). *Aquatic Toxicology* **140–**

141, 295–302 (2013).

74. K. Heffern, K. Tierney, E. P. Gallagher, Comparative effects of cadmium, zinc, arsenic and

chromium on olfactory-mediated neurobehavior and gene expression in larval zebrafish

(*Danio rerio*). *Aquatic Toxicology* **201**, 83–90 (2018).

75. J. T. Hentig, C. A. Byrd-Jacobs, Exposure to zinc sulfate results in differential effects on

olfactory sensory neuron subtypes in adult zebrafish. *International journal of molecular*

sciences **17**, 1445 (2016).

76. D. A. Klaprat, S. Brown, T. J. Hara, The effect of low pH and aluminum on the olfactory

organ of rainbow trout, Salmo gairdneri. *Environmental Biology of Fishes* **22**, 69–77

(1988).

77. E. Zeiske, R. Melinkat, H. Breucker, J. Kux, Ultrastructural studies on the epithelia of the

olfactory organ of cyprinodonts (teleostei, cyprinodontoidea). *Cell Tissue Res.* **172**, 245–

267 (1976).

78. P. A. Triana-Garcia, "Studies on The Olfactory Biology and Behavioral Responses to Copper

Exposure of The Native Indicator Species Delta Smelt (Hypomesus transpacificus)," UC

Davis. (2021).

79. P. A. Triana-Garcia, G. A. Nevitt, J. B. Pesavento, S. J. Teh, Gross morphology, histology,

and ultrastructure of the olfactory rosette of a critically endangered indicator species, the

Delta Smelt, Hypomesus transpacificus. *J Comp Physiol A Neuroethol Sens Neural Behav*

Physiol **207**, 597–616 (2021).

80. T. Shimizu, "Mucus, Goblet Cell, Submucosal Gland" in *Nasal Physiology and*

Pathophysiology of Nasal Disorders, T. M. Önerci, Ed. (Springer, 2013), pp. 1–14.

81. M. Lazzari, S. Bettini, L. Milani, M. G. Maurizii, V. Franceschini, Differential response of

olfactory sensory neuron populations to copper ion exposure in zebrafish. *Aquatic*

Toxicology **183**, 54–62 (2017).

82. K. E. Klima, F. M. Applehans, Copper exposure and the degeneration of olfactory receptors in rainbow trout (Oncorhynchus mykiss). *Chemical Speciation & Bioavailability* **2**, 149–154 (1990).

83. D. Saucier, L. Astic, Morpho-functional alterations in the olfactory system of rainbow trout (*Oncorhynchus mykiss*) and possible acclimation in response to long-lasting exposure to low copper levels. *Comparative Biochemistry and Physiology Part A: Physiology* **112**, 273–284 (1995).

84. Fish and Wildlife Service, US Department of the Interior, Endangered and Threatened Wildlife and Plants; 90-Day Finding on a Petition To List the Colorado River Cutthroat Trout. *Federal Register* **69**, 21151–21158 (2004).

85. N. H. Barton, K. S. Gale, "Genetic Analysis of Hybrid Zones" in *Hybrid Zones and the Evolutionary Process*, p. 33.

86. B.-J. Velthuis, W. Yang, T. Van Opijnen, J. H. Werren, Genetics of female mate discrimination of heterospecific males in *Nasonia* (Hymenoptera, Pteromalidae). *Animal Behaviour* **69**, 1107–1120 (2005).

87. O. Svensson, *et al.*, The genetics of mate preferences in hybrids between two young and sympatric Lake Victoria cichlid species. *Proceedings of the Royal Society B: Biological Sciences* **284**, 20162332 (2017).

88. E. M. Schmidt, K. S. Pfennig, Hybrid female mate choice as a species isolating mechanism: environment matters. *Journal of Evolutionary Biology* **29**, 865–869 (2016).

89. R. M. Paulette, Uno de los rostros de la migración interna. Marginación y pobreza en la
sierra hidalguense. *Sociológica México* 157–191 (2015).

90. S. Cruz Hernández, *et al.*, Saberes tradicionales locales y el cambio climático global. *Revista*
mexicana de ciencias agrícolas **11**, 1917–1928 (2020).

91. Y. Xia, C. Ti, D. She, X. Yan, Linking river nutrient concentrations to land use and rainfall in
a paddy agriculture–urban area gradient watershed in southeast China. *Science of The Total*
Environment **566–567**, 1094–1105 (2016).

92. E. Apeagyei, M. S. Bank, J. D. Spengler, Distribution of heavy metals in road dust along an
urban-rural gradient in Massachusetts. *Atmospheric Environment* **45**, 2310–2323 (2011).

93. J. J. Sansalone, J. P. Hird, F. K. Cartledge, M. E. Tittlebaum, Event-Based Stormwater
Quality and Quantity Loadings from Elevated Urban Infrastructure Affected by
Transportation. *Water Environment Research* **77**, 348–365 (2005).

94. S. Yu, *et al.*, Anthropogenic land uses elevate metal levels in stream water in an urbanizing
watershed. *Science of The Total Environment* **488–489**, 61–69 (2014).

95. V. Paul, M. S. Sankar, S. Vattikuti, P. Dash, Z. Arslan, Pollution assessment and land use land
cover influence on trace metal distribution in sediments from five aquatic systems in
southern USA. *Chemosphere* **263**, 128243 (2021).

96. D. H. Baldwin, C. P. Tatara, N. L. Scholz, Copper-induced olfactory toxicity in salmon and
steelhead: Extrapolation across species and rearing environments. *Aquatic Toxicology* **101**,
295–297 (2011).

97. A. E. Smith, "The influence of sub-lethal zinc exposure on olfaction of alarm cues and associated alarm behaviours in juvenile coho salmon," University of British Columbia. (2023).

98. L. Sonnack, *et al.*, Effects of metal exposure on motor neuron development, neuromasts and the escape response of zebrafish embryos. *Neurotoxicology and Teratology* **50**, 33–42 (2015).

99. H. Tjälve, C. Mejàre, K. Borg-Neczak, Uptake and Transport of Manganese in Primary and Secondary Olfactory Neurones in Pike. *Pharmacology & Toxicology* **77**, 23–31 (1995).

100. Y. Rodríguez-Agudelo, *et al.*, Motor alterations associated with exposure to manganese in the environment in Mexico. *Science of The Total Environment* **368**, 542–556 (2006).

101. D. B. Szymkowicz, *et al.*, Exposure to arsenic during embryogenesis impairs olfactory sensory neuron differentiation and function into adulthood. *Toxicology* **420**, 73–84 (2019).

102. J. Lunt, D. L. Smee, Turbidity interferes with foraging success of visual but not chemosensory predators. *PeerJ* **3**, e1212 (2015).

103. S. M. Leahy, M. I. McCormick, M. D. Mitchell, M. C. O. Ferrari, To fear or to feed: the effects of turbidity on perception of risk by a marine fish. *Biology Letters* **7**, 811–813 (2011).

104. S. J. Song, B. Park, K. Jo, C.-S. Kim, Damage to Olfactory Organs of Adult Zebrafish Induced by Diesel Particulate Matter. *International Journal of Molecular Sciences* **23**, 407 (2022).

105. I. Reiten, *et al.*, Motile-Cilia-Mediated Flow Improves Sensitivity and Temporal Resolution

of Olfactory Computations. *Current Biology* **27**, 166–174 (2017).

106. J. P. L. Cox, Hydrodynamic aspects of fish olfaction. *Journal of The Royal Society Interface*

5, 575–593 (2008).

107. D. P. McEwen, P. M. Jenkins, J. R. Martens, "Chapter 12 Olfactory Cilia: Our Direct

Neuronal Connection to the External World" in *Current Topics in Developmental Biology*,

Ciliary Function in Mammalian Development., (Academic Press, 2008), pp. 333–370.

108. F. W. Sunderman, Nasal Toxicity, Carcinogenicity, and Olfactory Uptake of Metals. *Ann*

Clin Lab Sci **31**, 3–24 (2001).

1088

1089