Plasticity, not genetics, shapes individual responses to thermal stress in non-native populations of the European green crab (*Carcinus maenas*)

Article type: Research article
Running title: Green crab thermal tolerance and plasticity

Yaamini R. Venkataraman[1]*, Julia C. Kelso[1,2], Catlin Payne[1,3], Heidi L. Freitas[1,3], Jasmine Kohler[1,4], Carolyn K. Tepolt[1]

[1]Biology Department, Woods Hole Oceanographic Institution, Woods Hole, MA, USA
[2]Department of Genetics, Harvard Medical School, Boston, MA, USA
[3]Department of Biology, University of Massachusetts, Boston, Boston, MA, USA
[4]Cape Cod Community College, West Barnstable, MA, USA

* = corresponding author, yaamini.venkataraman@gmail.com

Abstract

Temperature is a major driver of individual performance in ectotherms, with this impact depending on stressor intensity and duration. Differences in individual response across temperature, time, and populations are shaped by the interplay between evolutionary adaptation and phenotypic plasticity. Some populations are able to thrive in novel and changing environments despite limited genetic diversity, raising the question of how plasticity and adaptation interact after significant genetic diversity loss. The European green crab (*Carcinus maenas*) is a textbook example of this phenomenon: invasive populations boast a broad thermal tolerance and exceptional thermal flexibility even after repeated genetic bottlenecks. Despite this loss of diversity overall, prior work has found a strong population-level association between variation at a specific extended genomic region (supergene), cold tolerance, and sea surface temperature. We conducted a series of three experiments using righting response to characterize sublethal thermal tolerance and plasticity in introduced green crab populations, then determined if these factors were associated with supergene genotype for individual adult crabs. Crabs showed signs of stress after exposure to a 30⁰C heat shock in one experiment. Interestingly, a second experiment exposing *C. maenas* to repeated 24-hour heat shocks showed that prior heat shock conferred beneficial plasticity during a subsequent event. The third experiment examined cold acclimation over multiple timepoints up to 94 hours. At 5⁰C, certain crabs exhibited an acclimatory response where righting slowed dramatically at first, and then gradually sped up after a longer period of cold exposure. Several crabs failed to right at 1.5⁰C, which could be indicative of dormancy employed to reduce energy consumption in colder conditions. There were no significant relationships between individual plasticity and supergene genotype in any experiment. Linking population-level genetic associations with individual-level physiology is complex, and reflects the impact of environmental conditions such as temperature throughout life history in shaping adult phenotype. Our results highlight the robust thermal tolerance and plasticity that adult green crabs maintain despite a substantial reduction in genetic diversity, and underscore the importance of probing population-level genotype-phenotype associations at the individual level.

Keywords: European green crab, acute stress tolerance, thermal plasticity, genotype, heat shock, heat priming, cold acclimation

Introduction

Temperature is a primary driver of physiology in marine invertebrates (Somero,

Lockwood and Tomanek 2017), and substantial effort has focused on quantifying its impacts on

these species (Yao and Somero 2014). In some circumstances, acute thermal stress may elicit

a pronounced stress response, such as higher mortality, which then subsides after extended

periods of stress exposure (Zou *et al.* 2023). Alternatively, short-term exposure can be less

stressful due to acclimatory potential not present in response to longer-term stressors (Madeira

et al. 2018; Morley *et al.* 2019). Populations within a species can also exhibit distinct responses

to thermal stress due local adaptation or thermal history (Sanford and Kelly 2011; Fields, Cox

and Karch 2012; Falfushynska, Phan and Sokolova 2016). These different responses across

temperature, time, and populations can be attributed to various mechanisms, including shifts in

metabolic pathways, changes in lipid membrane composition, and differential gene expression

(Pörtner 2002; Somero 2010, 2012). Understanding these mechanisms may aid in predicting

organismal responses to more frequent thermal extremes (Calvin *et al.* 2023) and uncover why

certain species and populations thrive in inherently stressful conditions (Somero 2010).

Intraspecific variation in physiology often arises from some combination of evolutionary

adaptation and phenotypic plasticity (Fox *et al.* 2019). Evolutionary adaptation depends on

genetic differences within a species that alter fitness in a given environment (Fox *et al.* 2019).

For example, genetic differences over five decades in *Daphnia magna* were associated with

changes in critical thermal maxima (Cuenca Cambronero *et al.* 2018). Phenotypic plasticity, or

the ability for a single genotype to produce more than one phenotype, also plays a critical role in

shaping an individual organism's thermal tolerance (Hendry 2016; Fox *et al.* 2019). Thermal

priming is a common plastic response in many species including marine invertebrates, in which

prior exposure to heat stress confers beneficial plasticity in response to future shocks (Martell

2022; Glass *et al.* 2023; Abbas *et al.* 2024). Repeated heat shocks boosted thermal tolerance of

the abalone *Haliotis tuberculata* with no long-term detrimental effects (Abbas *et al.* 2024), while

prior exposure to elevated temperatures improved *Nematostella vectensis* growth, development,

and acute heat tolerance (Glass *et al.* 2023). Populations in variable environments often evolve

greater capacity for plasticity (Handelsman *et al.* 2013; Landy *et al.* 2020). However, acclimatory

plasticity may not fully mediate negative impacts of thermal stress (Gunderson and Stillman

2015), and tradeoffs between tolerance and plasticity can produce thermally-tolerant genotypes

with lower plasticity (Barley *et al.* 2021; Griffiths *et al.* 2024). For example, experimental

evolution of the copepod *Acartia tonsa* produced genotypes with a 2-5ºC increase in LD50 over

40 generations, but with decreases in plasticity of over 60% (Sasaki and Dam 2021).

Understanding the interplay of these mechanisms may uncover why certain species thrive in

inherently stressful conditions (Somero 2010).

If changes in thermal physiology are driven by selection and plasticity, then does

plasticity alone explain why some populations boast broad environmental tolerances with limited

genetic diversity? Several introduced species thrive in novel environments even after substantial

losses of genetic variation (Roman and Darling 2007; Estoup *et al.* 2016). However, even highly

bottlenecked populations often retain some genetic diversity, including (and perhaps especially)

at adaptive markers (Swindell and Bouzat 2005; Oliver and Piertney 2012; Teixeira and Huber

2021). Population genomic research has identified specific genetic markers correlated with

various environmental parameters across populations in many introduced species (Schwander,

Libbrecht and Keller 2014; Zhang *et al.* 2023; Ma *et al.* 2024), but it remains a challenge to link

these population-level association studies with individual-level physiological mechanisms

(Riginos *et al.* 2016; Lind and Lotterhos 2024). Some studies have successfully identified

genome-phenome linkages at various levels of biological hierarchy. For example, genetic

markers associated with warmer and more arid conditions are positively correlated with

Anopheles gambiae larval thermal tolerance (Rocca *et al.* 2009), and flies (*Drosophila*

suboscura) with putatively warm-adapted genotypes had higher abundance of heat shock

proteins than flies with cold-adapted genotypes (Calabria *et al.* 2012). Individual survival within

one population of adult dwarf surf clams *Mulina lateralis* at 30ºC for 17 days was associated

with variation in genes in the ETHR/EHF signaling pathway (Wang *et al.* 2024). Additionally,

these genes were more highly expressed in heat-tolerant clams, further illustrating the

importance of these population-level genetic markers to individual thermal tolerance (Wang *et*

al. 2024).

The European green crab (*Carcinus maenas*) provides the perfect opportunity to study

these interactions in genetically depauperate yet highly plastic populations. Like many non-

indigenous species, *C. maenas* boasts broad thermal tolerance and plasticity (Davidson,

Jennions and Nicotra 2011; Zerebecki and Sorte 2011; Sorte *et al.* 2013), with upper thermal

limits higher and lower thermal limits lower than other temperate crustaceans (Tepolt and

Somero 2014; Tepolt 2024). Their evolution in variable estuarine environments spanning 4-23ºC

in the native range may have shaped this wide thermal optima range (Compton, Leathwick and

Inglis 2010; Jost, Podolski and Frederich 2012). A variety of mechanisms may underpin this

broad thermal tolerance, including "dumping" hemocyanin-bound oxygen to extend upper

thermal limits (Jost, Podolski and Frederich 2012) and altering hemolymph magnesium content

in colder temperatures to reduce the blockage of calcium channels that otherwise would impair

movement (Frederich *et al.* 2000; Wittmann *et al.* 2010). Seasonal differences in thermal limits

(Cuculescu, Hyde and Bowler 1998; Hopkin *et al.* 2006) and rapid homeoviscous adaptation

(Chapelle 1978, 1986; El Babili *et al.* 1997) observed in *C. maenas* underscore the importance

of phenotypic plasticity to the species' success across broad thermal gradients.

Green crabs in invasive North American populations are characterized by substantial

losses of genome-wide genetic diversity, but still maintain broad thermal plasticity (Tepolt and

Palumbi 2015) (see **Appendix S1** for a detailed *C. maenas* invasion history). North American *C.*

maenas inhabit a broader thermal range than in their native range (Compton, Leathwick and

Inglis 2010), and are thriving in sub-arctic waters in Newfoundland (Kelley, de Rivera and

Buckley 2013; Best, McKenzie and Couturier 2017). While introduced *C. maenas* lost

substantial genetic diversity, the variation remaining in these populations may underpin thermal

plasticity. Population genomics of *C. maenas* revealed a group of single nucleotide

polymorphisms (SNPs) in strong linkage disequilibrium that were significantly associated with

thermal tolerance on a population level; these SNPs likely represent an extended region of

reduced recombination, or supergene (Tepolt and Palumbi 2020). Crab populations with higher

heart rates at 0ºC had a higher frequency of the putatively "cold-adapted" supergene allele

(Tepolt and Palumbi 2020), while populations experiencing warmer average sea surface

temperatures had a higher frequency of the putatively "warm-adapted" supergene allele (Tepolt

and Palumbi 2020). In a separate study, this supergene represented nearly all of the genomic

variation associated with sea surface temperature in northeast Pacific populations (Tepolt *et al.*

2022). Among the dozens of single-nucleotide differences between the two main supergene

alleles are two nonsynonymous SNPs in *hypoxia-inducible factor 1-alpha* (Tepolt and Palumbi

2020): as a key regulator of hypoxia, it may also be essential in marine invertebrate responses

to temperature and offers a potential functional link between supergene genotype and thermal

phenotype (Pörtner, Langenbuch and Michaelidis 2005; Pörtner, Bock and Mark 2017). While

the genomic data suggest a strong link between genomic adaptation and population-level

thermal tolerance, it is unclear if or how these differences translate to individual thermal

plasticity. Supergene genotype had no impact on individual response to acute cold stress in

adult *C. maenas* from several northwest Atlantic populations (Coyle *et al.* 2019); however, the

crabs were only exposed to 4.5ºC for a maximum of six hours. Given the complexities of linking

population-level associations with individual physiology (Gagnaire and Gaggiotti 2016; Riginos

et al. 2016; Tam *et al.* 2019; Lind and Lotterhos 2024), investigating the role of the supergene at

the individual level may help elucidate the mechanisms that shape thermal tolerance in the

species.

149   We conducted three experiments to investigate individual thermal tolerance and

150 plasticity in adult crabs in response to thermal extremes, then tested whether the observed

151 responses were linked to the temperature-associated supergene (**Figure 1**). We first compared

152 responses to acute heat shock in crabs from Washington and Maine ("Experiment 1"), and

153 hypothesized that crabs from Maine, which experience higher summer temperatures, would

154 display a higher heat tolerance than their Washington counterparts. To understand plasticity in

155 response to heat stress, we exposed crabs from Washington to repeated acute heat shocks

156 ("Experiment 2"). We predicted that "priming" from previous heat shock exposure would confer

157 beneficial plasticity in response to subsequent acute heat stress. Finally, we characterized cold

158 tolerance and plasticity for crabs from Massachusetts and Washington over multiple timepoints

159 across 94 hours ("Experiment 3"), to explore the extent and time scale of plasticity at low

160 temperatures. We expected that crabs would slow upon initial cold exposure, but might exhibit

161 evidence of thermal acclimation over longer periods. We also expected crabs from MA to have a

162 higher cold tolerance than WA crabs due to lower winter temperatures in the northwest Atlantic

163 than in the northeast Pacific. Across all three experiments, we tested whether and how

164 supergene genotype influenced thermal tolerance and plasticity. We expected crabs with at

165 least one copy of the putatively warm-adapted allele to be more robust to heat stress.

166 Conversely, we hypothesized that the putatively cold-adapted allele would improve response to

167 short- and medium-term cold acclimation. Our results show how investigating physiological

168 response at the individual organism level can elucidate how plasticity and genetic background

169 shape thermal tolerance in genetically depauperate but highly successful populations.

170 **Methods**

171 *General experimental setup*

A series of experiments was conducted to assess the impact of short- (~24 hours) to

medium-term (94 hours) heat and cold stress on introduced *C. maenas* populations in both the

northwest Atlantic and northeast Pacific. Experimental crabs were individually labeled using

waterproof paper and superglue, and the propus and dactyl segments from the third walking leg

on the right side (dorsal view) were preserved in 95% ethanol for subsequent genotyping. These

joints were taken as the third walking leg has no significant role in crab righting behavior

(Young, Peck and Matheson 2006). If the crab did not have a third walking leg on the right side,

the third walking leg was taken from the left.

Crabs were placed in 21 cm x 26 cm x 41 cm (22.386 L) tanks filled with 800 micron

filtered raw seawater from Great Harbor, MA, USA (Experiments 1 and 3) or 50 micron filtered

raw seawater from Vineyard Sound, MA, USA (Experiment 2), and equipped with an air stone

and filter (**Appendix S2**). Crabs were distributed to balance sex, carapace width, and

integument color among tanks. Crabs with orange-red or red integument color were not included

in experiments; these colors indicate an extended time in intermolt and "red" green crabs have

less robust tolerance to a range of physiological stressors (Styrishave, Rewitz and Andersen

2004). Remaining crabs had integument colors at the beginning of the experiment ranging from

blue to yellow-orange. Tanks were checked for mortalities, cleaned, and fed daily (**Appendix**

S2). Ammonia concentrations were checked every third day and filters were replaced every five

days (**Appendix S2**). A 12-hour:12-hour light:dark cycle was maintained for all tanks (Tepolt

and Somero 2014).

Crabs were initially kept in acclimation conditions for at least five days (see details

below). Water temperature was maintained during the acclimation period using either a flowing

water bath or environmental chamber manipulation, and heated or cooled using a 300-Watt

Deluxe Titanium Heating Tubes (Finnex, USA) with an external temperature controller (bayite,

China) or environmental chamber, respectively (**Appendix S2**). Water temperature in each tank

was recorded every 15 minutes using two HOBO Data Loggers (Onset, USA), one at the water

surface and one at the bottom of the tank. A non-parametric Kruskal-Wallis test

(`kruskal.test` from R `stats` package (R Core Team 2024) was used to confirm that

temperature treatments were statistically different from each other and that replicate tanks were

not different from each other.

Measuring thermal tolerance with time-to-right

Time-to-right (TTR) was used in all experiments to assess sublethal thermal tolerance.

TTR was defined as the number of seconds *C. maenas* spent on righting behavior (defined in

(Young, Peck and Matheson 2006)), including "hesitation time" after crabs are placed dorsal

side up but before moving their fifth pereopod pair. Individual crabs were placed with their dorsal

side on the bottom surface of the tank and given 90 seconds to right. If a crab did not right

within 90 seconds, it was manually righted and recorded as "not righting" for that trial. Crabs

were rested for 10 seconds before repeating the TTR trial twice for a total of three TTR

measurements per crab per trial. Readings were always conducted by the same person within

an experiment to reduce error. To control for other factors that may influence righting response,

crabs were weighed, measured, sexed, and assessed for integument color and missing legs

after each TTR trial.

Supergene genotyping

Crabs were genotyped at a single-nucleotide polymorphism (SNP) in the *structural*

maintenance of chromosomes protein 3 (SMC) gene to understand how individual variation at

the putative supergene impacts thermal tolerance. This SNP is part of a group of SNPs in very

strong linkage disequilibrium in the putative supergene, and is diagnostic of supergene allele

(Tepolt & Palumbi 2020). DNA was extracted from ~4 mm^2 of leg joint tissue using the DNeasy

Blood & Tissue Kit (QIAGEN, Valencia, CA, USA) following manufacturer's instructions.

Incubation with the ATL Lysis Buffer and Proteinase K occurred for at least two hours. The final

eluate was pipetted back onto the spin column, incubated, and re-eluted to increase DNA yield.

Genotyping was done with one of two equivalent methods. For Experiment 2, a 126 bp

region of the SMC gene containing a C/T single nucleotide polymorphism (SNP) diagnostic for

supergene allele was amplified using the SMC F and R primers and methods specified in Coyle

et al. (2019) (**Table S1**). Amplified product was sent to Sequegen (Worcester, MA, USA) for

Sanger sequencing. Resultant chromatograms were edited and aligned using Sequencher

v.5.4.6 (Gene Codes, Ann Arbor, MI, USA). Samples from Experiments 1 and 3 were genotyped

using a restriction digest assay. A 690 bp region of the SMC gene containing the same

diagnostic C/T SNP was amplified using SMC_long F and R primers (**Table S1**). Amplified

product was digested with 0.5 µL of AluI enzyme (Promega, USA or ThermoFisher, USA) at

37ºC for one hour. The recognition site includes a CG dinucleotide at 88 bp in the PCR product,

meaning that digestion only occurs for C alleles There is an additional non-target cut site at 477

bp that is digested for all genotypes. Digested product was run out on a 1.5% agarose gel with

GelRed® Nucleic Acid Gel Stain (Biotium, Inc, Fremont, CA, USA) at 90 V for at least 65

minutes then imaged. A 477 bp band indicated TT homozygotes, a 389 bp band indicated CC

homozygotes, and both bands indicated CT heterozygotes (**Figure S1**). Both CC and CT

genotypes had also an 88 bp band. All genotypes had a 213 bp band due to digestion at the

non-target site (**Figure S1**). The C allele is found in the cold-adapted supergene allele, while the

T allele is found in the warm-adapted supergene allele.

Experiment 1: Responses to acute heat shock across populations

This experiment gauged *C. maenas* response to an acute heat shock in WA and ME

populations (**Appendix S3**). The WA and ME crabs were placed in tanks with 15-17 crabs each,

separated by site. All crabs were acclimated at room temperature (~22-24ºC) for at least five

days. After the acclimation period, crabs were exposed to a heat ramp of ~1ºC/hour to 30ºC

over 6 hours, then remained at 30⁰C for 22 hours (**Figure S3**). TTR was assessed before and at

the end of the 30⁰C heat shock. Prior to analysis, the three replicate TTR measurements for

each individual crab at each time point were averaged and log-transformed. Trials where the

crab did not right itself within 90 seconds were excluded from averaging. After experiments were

concluded, crabs were sacrificed and dissected to identify and count trematode parasite cysts in

the hepatopancreas tissue per Blakeslee et al. (2015), with the modification that all

hepatopancreas tissue was examined on 12.5 cm x 7.5 cm squash plates. Parasite intensity

was calculated by dividing metacercarial cyst count by crab carapace width. Since *C. maenas*

populations in the northwest Atlantic are parasitized while counterparts in the northeast Pacific

are not, parasite intensity was quantified to control for any extraneous factors impacting righting

when comparing ME and WA crab responses.

Linear mixed effects models (`lmer` from R `lme4` package; (Bates *et al.* 2015)) were

used in three steps to evaluate how various factors impacted log-transformed average righting

response time with crab ID used as a random effect to correct for repeated measurements of

the same animals. First, a null model was constructed with crab sex, integument color, carapace

width, weight, whether or not a crab was missing a fifth pereopod ("swimmer"), and parasite

intensity (ME only) as explanatory variables to understand how these factors explained variation

in righting response. An Analysis of Variance (ANOVA; `anova` from R `stats` package (R Core

Team 2024)) was used to perform a likelihood ratio test and compare the null model with a

model testing the influence of a specific factor (*ie.* only carapace width or parasite infection

intensity). Second, any significant variables identified by ANOVA tests were included in a new

model along with timepoint. Timepoint was modeled as a categorical variable because it directly

corresponded with heat shock (hour 0 = no heat shock, hour 22 = heat shock). Significant

factors (*ie.* timepoint, any significant demographic variables) identified by ANOVA tests were

then included in a third and final null model. This "no genotype" null model was compared to

three separate models containing significant factors identified by the previous two null models

and either genotype, presence of the C allele, or presence of the T allele. A final set of ANOVA

tests were used to discern if these genotype-specific factors significantly impacted log-

transformed average righting response. For all models, *P*-values were corrected for multiple

comparisons using a Bonferroni correction. Factors were considered significant when adjusted p

< 0.05. Normality and homoscedasticity were verified visually. Pairwise tests were conducted

with `emmeans` v.1.11.0 for any significant factor (Lenth *et al.* 2018).

Experiment 2: The role of "heat priming" in shaping thermal tolerance

The goal of this experiment was to understand how repeated heat shocks impacted *C.*

maenas thermal plasticity. After the acclimation period (**Appendix S4**), two tanks (treatment

group) were exposed to a ~30ºC heat shock for 20 hours, while the other two tanks (control

group) were maintained at 15ºC. After the initial heat shock, all tanks were maintained at 15ºC

for 48 hours. Both the control and treatment groups were then exposed to a second heat shock

of ~30°C for 24 hours.

Righting response was measured for all crabs after the acclimation period (day 0), at the

end of the first heat shock (day 1), and at the end of the second heat shock (day 4).

Measurements taken for crabs experiencing heat shocks were conducted at 30ºC. A three-step

series of linear mixed effects models was used to assess the impact of sex, integument color,

carapace width, weight, a missing swimmer, treatment, day, the interaction of treatment and

day, genotype, presence of the C allele, and presence of the T allele on log-transformed

average TTR, with crab ID used as a random effect to correct for repeated measurements of the

same animals. (see *Experiment 1* for specific methods). For this experiment, day was modeled

as a categorical predictor, as it was a proxy for heat shock timing. Pairwise tests were

conducted with `emmeans` v.1.11.0 for any significant factor (Lenth *et al.* 2018). Oxygen

consumption was also measured for a subset of crabs in this experiment (see **Appendix S4** for

details).

Experiment 3: Changing cold tolerance across populations over time

This experiment investigated how short- and medium-term exposure to cold

temperatures influenced thermal tolerance and plasticity in green crabs from MA and WA over

the course of 94 hours (**Appendix S5**). At hours 0, 4, 22, 28, 46, and 94, righting response was

assessed for at least seven crabs per tank. A series of linear mixed effects models was used to

assess the impact of sex, integument color, carapace width, weight, a missing swimmer,

treatment, time, the interaction of treatment and time, genotype, presence of the C allele, and

presence of the T allele on log-transformed average TTR, with crab ID used as a random effect

to correct for repeated measurements of the same animals (see *Experiment 1* for specific

methods). For this experiment, timepoint was modeled as a continuous predictor. Pairwise tests

were conducted with `emmeans` v.1.11.0 for any significant factor (Lenth *et al.* 2018). To obtain

pairwise differences between different timepoints, timepoint was re-estimated as a categorical

variable.

A different set of models was used to evaluate how various factors impacted failure to

right within the 90 second cutoff. For these models, TTR for each crab was coded as a

"success" (righting within 90 seconds) or "failure" (no righting within 90 seconds). Binomial linear

mixed effects models with crab ID as a random effect were used to understand how various

factors impacted binary righting response in three steps. First, a null model was constructed

with crab sex, integument color, carapace width, weight, and whether or not a crab was missing

a swimmer as explanatory variables to understand how these factors explained variation in

binary righting response. An Analysis of Variance (ANOVA; `anova` from R `stats` package (R

Core Team 2024)) was used to perform a likelihood ratio test and compare the null model with a

model testing the influence of a specific factor (*ie.* only carapace width or weight). Second, any

significant variables identified by ANOVA tests were included in a new model along with

temperature, timepoint, and their interaction. Significant factors (*ie.* timepoint, any significant

demographic variables) identified by ANOVA tests were then included in a third and final null

model. This "no genotype" null model was compared to three separate models containing

significant factors identified by the previous two null models, and either genotype, presence of

the C allele, or presence of the T allele. A final set of ANOVA tests were used to discern if these

genotype-specific factors significantly impacted log-transformed average righting response.

Factors were considered significant at p < 0.05 after Bonferroni correction.

Sample size decreased over the course of the experiment due to 5-6 crabs per

treatment (1-2 crabs per tank) being randomly selected for dissection at each timepoint for

future analyses. To account for this, a sensitivity analysis was used to confirm the validity of the

average time-to-right and failure-to-right modeling results (see **Appendix S5** for details).

To compare responses to temperature between populations, separate linear mixed

effects models were created to evaluate the impact of population and significant variables from

the population-specific models on average time to right, or on failure to right. Generalized linear

models (`glm` from R `stats` package (R Core Team 2024)) were also used to compare the

effect of population on failure to right at each timepoint for crabs in the colder temperature

treatment. All *P*-values were corrected for multiple comparisons using a Bonferroni correction.

Results

Experiment 1: Responses to acute heat shock across populations

Righting response was evaluated before and after a 22 hour heat shock to understand

the factors that impact short-term heat tolerance. Timepoint had a significant impact on righting

response in both populations (**Figure 2A**; WA comparison: $\chi^2 = 20.85$, *P*-value < 0.0001; ME

comparison: $\chi^2 = 19.17$, *P*-value < 0.0001). Crabs from both WA and ME had significantly

slower TTR after heat shock (mean$_{WA, 22\ hours}$ = 3.22 s ± 4.22 s; mean$_{ME, 22\ hours}$ = 2.22 s ± 2.15 s)

than before (mean$_{WA, 0\ hours}$ = 1.16 s ± 0.26 s; mean$_{ME, 0\ hours}$ = 1.23 s ± 1.15 s). Neither genotype

(χ^2 = 0.48, *P*-value = 1), presence of the C allele (χ^2 = 3.02 x 10^{-4}, *P*-value = 1), nor presence of

the T allele (χ^2 = 3.24, *P*-value = 1) had a significant impact on average TTR in WA *C. maenas*

(**Figure 2B**). A similar trend was identified in the ME population, which only had CT and CC

crabs (**Figure S2B**). After heat shock, CT crabs had a slightly faster righting response (mean$_{ME,}$

$_{22\ hours,\ CT}$ = 1.46 s ± 0.20 s) than CC crabs (mean$_{ME,\ 22\ hours,\ CC}$ = 2.67 s ± 2.62 s), but this

difference was not significant after multiple test correction (genotype: χ^2 = 5.65, P-value = 0.12)

(**Figure 2B**). Righting response was not impacted by sex, integument color, carapace width,

weight, or whether or not a crab was missing a fifth pereopod in this or any other experiment

(**Table S3**). Parasite infection intensity did not impact righting response in ME crabs (**Table S3**).

Crab ID explained 4.11% and 21.69% of the variation in final WA and ME models (timepoint

only), respectively.

Experiment 2: The role of "heat priming" in shaping thermal tolerance

The righting responses of *C. maenas* that previously experienced heat stress were

compared to the responses of crabs naïve to stress in order to understand how repeated

exposures impact thermal plasticity (**Figure S5**). We note that there were small temperature

differences between tanks for the second heat shock (**Table S2**; **Figure S5**). Because of this

experimental difference, tank was used as a random effect in subsequent analyses to account

for significant temperature differences.

Average TTR for an individual crab was significantly impacted by treatment (χ^2 = 20.52,

P-value = 0.0004), day (χ^2 = 24.16, *P*-value = 0.0008), and their interaction (χ^2 = 15.69, *P*-value

= 0.004) (**Figure 3A**). These trends were driven by differences in TTR between control and

treatment crabs at the end of the experiment. On day 4, *C. maenas* exposed to a prior heat

shock had faster and less variable righting times (mean$_{treatment}$ = 1.18 seconds ± 0.62 seconds)

than their counterparts that did not experience a prior heat shock (mean$_{control}$ = 4.78 s ± 4.98 s)

(**Figure 3A**; t$_{control-treatment}$ = 4.48, *P*-value$_{control-treatment}$ = 0.001). Within-treatment comparisons

revealed that crabs that did not experience a prior heat shock had significantly slower righting

response on day 4 than they did on days 0 or 1 (**Figure 3A**; $t_{day0-day4}$ = -4.02, P-value$_{day0-day4}$ =

0.0005; $t_{day1-day4}$ = -5.06, P-value$_{day1-day4}$ < 0.0001).

Supergene genotype was not associated with thermal response and priming. Genotype

(χ^2 = 3.52, P-value = 1), presence of the C allele (χ^2 = 1.67, P-value = 1), and presence of the T

allele (χ^2 = 2.73, P-value = 1) had no significant impact on average TTR. Although these trends

were not statistically significant, crabs with at least one T allele had slightly slower TTR than

their counterparts without a T allele, irrespective of prior heat shock exposure (**Figure 3B**;

mean$_{treatment,day\ 4,\ T}$ = 1.13 s ± 0.68 s, mean$_{treatment,day\ 4,\ no\ T}$ = 0.80 s ± 0.06 s, mean$_{control,\ day\ 4,\ T}$ =

6.17 s ± 6.01 s, mean$_{control,\ day\ 4,\ no\ T}$ = 2.83 s ± 2.44 s). Crab ID and tank accounted for 23.41% or

0% of the variance in TTR in the final model (treatment, time, and their interaction only),

respectively.

Experiment 3: Changing cold tolerance across populations over time

Crabs from MA and WA were ramped down to target temperatures of 1.5ºC or 5ºC and

held for several days to understand the factors that govern short- and medium-term cold

tolerance (**Figure S7**). All tanks reached experimental temperatures by hour 22 of the 94 hour

experiment. Actual temperatures experienced were slightly higher than the targets (**Table S2**).

Differences in temperature conditions between these two experiments were likely due to the

additional tanks used for WA and higher ambient temperatures in July 2024 versus June 2024

impacting cooling efficiency of the environmental chamber.

Righting response of MA crabs was significantly impacted by temperature (χ^2 = 77.69, P-

value < 0.0001), time (χ^2 = 161.78, P-value < 0.0001), and their interaction (χ^2 = 51.81, P-value

< 0.0001) (**Figure 4A**). Crabs at 1.5ºC had significantly slower TTR than crabs in 5ºC at all

timepoints except for hour 0, when all crabs were at their common 15ºC acclimation

temperature (**Table S4**). Within each treatment, there were also significant differences in TTR

between timepoints (**Figure 4A; Table S5**). Supergene genotype ($\chi^2 = 0.61$, *P*-value = 1),

presence of the C allele ($\chi^2 = 0.38$, *P*-value = 1), or presence of the T allele ($\chi^2 = 0.19$, *P*-value

= 1) did not significantly impact average TTR in MA *C. maenas* (**Figure 4B**). Crab ID explained

10.99% of the variance in the final model (temperature, time, and their interaction only).

Throughout the experiment, some MA *C. maenas* failed to right at 1.5ºC, but none failed

to right at 5ºC. Shortly after first reaching 1.5ºC at hour 22, 90% of crabs failed to right, while

64.3% of the remaining crabs failed to right at the end of the experiment (**Figure 4C**). Failure to

right was significantly associated with temperature ($\chi^2 = 56.05$, *P*-value < 0.0001) and time ($\chi^2 =$

64.72, *P*-value < 0.0001), but not their interaction ($\chi^2 = 1.07 \times 10^{-11}$, *P*-value = 1), genotype ($\chi^2 =$

3.31, *P*-value = 1), nor the presence of the C allele ($\chi^2 = 1.99 \times 10^{-11}$, *P*-value = 1). The majority

of crabs that failed to right had at least one T allele (77.8% of crabs at 22 hours, 66.7% of crabs

remaining at 94 hours) (**Figure 4D**). However, the impact of the T allele on failure to right in the

MA population was not significant after multiple test correction ($\chi^2 = 3.31$, *P*-value = 0.82).

Failure to right was not impacted by sex, integument color, carapace width, weight, or whether

or not a crab was missing a fifth pereopod in any population (**Table S7**). There were no

differences between the final model and the most parsimonious model derived from the

sensitivity test for either MA model (**Table S8**).

Crabs from WA responded to 1.5ºC and 5ºC in a similar fashion to their MA

counterparts. Temperature ($\chi^2 = 113.80$, *P*-value < 0.0001), time ($\chi^2 = 287.84$, *P*-value <

0.0001), and their interaction ($\chi^2 = 65.95$, *P*-value < 0.0001) significantly impacted average

righting response in WA *C. maenas* (**Figure 5A**). Righting response in 1.5ºC was significantly

slower than TTR in 5ºC for all timepoints except hour 0, prior to the start of the cold ramp (**Table**

S5). Within a treatment, there were also significant differences in righting response over time

(**Table S6**). There was no significant impact of genotype ($\chi^2 = 1.06$, *P*-value = 1), presence of

the C allele ($\chi^2 = 0.40$, *P*-value = 1), or presence of the T allele ($\chi^2 = 0.54$, *P*-value = 1) on

average TTR in WA crabs (**Figure 5B**). Righting response was not impacted by sex, integument

color, carapace width, weight, or whether or not a crab was missing a fifth pereopod (**Table S3**).

Crab ID explained 5.32% of variance in the final model (temperature, time, and their interaction

only). There were no differences between the final model and the most parsimonious model

derived from the sensitivity test (**Table S6**).

In comparison to the MA population, a smaller proportion of WA *C. maenas* failed to right

at 1.5°C: 34.5% WA crabs failed to right at hour 22 and 40% of the remaining crabs failed to

right at hour 94 (**Figure 5C**). Failure to right in WA crabs was related to temperature ($x^2 = 27.10$,

P-value < 0.0001) and time ($x^2 = 23.46$, *P*-value < 0.0001), but not their interaction ($x^2 = 0$, *P*-

value = 1). There was also no impact of genotype ($x^2 = 2.64$, *P*-value = 1), presence of the C

allele ($x^2 = 2.60$, *P*-value = 1), or presence of the T allele ($x^2 = 0.06$, *P*-value = 1) on failure to

right in WA *C. maenas*. Time did not have a significant impact in the most parsimonious model

derived for the sensitivity test, and was therefore removed from the final model for the full

dataset (**Table S8**). Temperature was still significant in the revised model ($x^2 = 26.57$, *P*-value <

0.0001).

Population (MA or WA) did not have a significant impact on average TTR ($x^2 = 0.47$, *P*-

value = 0.49) or overall failure to right ($x^2 = 0$, *P*-value = 1). Population-specific impacts on

failure to right at 1.5°C were also examined for crabs at each timepoint. There was no significant

impact of population at any timepoint (*P*-value > 0.05 for all comparisons).

Discussion

We examined tolerance and phenotypic plasticity in response to heat and cold

exposures in a highly-successful species in two regions where it has been introduced and lost

significant genetic diversity relative to its native range. We found that acute exposure to sub-

lethal high temperature elicits a stress response which could be ameliorated by repeated sub-

lethal stress exposure. Exposure to cold temperatures led to slower righting or failed righting.

However, some crabs were able to recover, suggesting longer-term plasticity at cold

temperatures. Interestingly, a genomic region strongly associated with temperature and thermal

physiology at the population level showed no relationship with thermal physiology at the

individual level in our experiments. The maintenance of broad thermal tolerance and plasticity in

righting response in adult crabs with no apparent influence of supergene genotype suggests

that adult tolerance is largely reliant on outcomes of selection during early life history in addition

to robust phenotypic plasticity.

Green crabs display adaptive plasticity at sub-lethal temperatures

Our work investigating *C. maenas* response to heat shock adds to the body of work

highlighting the broad thermal tolerance of this species outside of its native range. We expected

that *C. maenas* would speed up righting response during the heat shock because metabolic

rates increase for ectotherms at warmer temperatures (Somero, Lockwood and Tomanek 2017).

Instead, we found that crabs from ME and WA slowed TTR after heat shock. This contrasts with

Jost et al. (2012), who report TTR starting to slow at 34ºC for ME *C. maenas*, with no decrease

at 32ºC. Methodological differences may be responsible for these contrasting results. While our

study held crabs at 30ºC for ~22 hours after a ~1ºC/hour heat ramp, Jost et al. (2012) used a

6ºC/hour increase in temperature. The faster temperature used in Jost et al. (2012) may

activate different acute response pathways, leading to a different physiological outcome.

Alternatively, the slower TTR observed in our study could indicate that thermal thresholds may

be reduced with prolonged exposure to an elevated temperature. Interestingly, crabs in the

treatment group experiencing their first 30ºC heat stress on day 1 in Experiment 2 did not

exhibit stress-related slowing of righting response, while crabs in the control group experiencing

a heat shock on day 4 did significantly slow their righting response. This unexpected result

could be the product of individual variation in stress response within our sampled population:

crab ID contributed to ~20% of variance in righting response, which was the highest out of any

experiment. Future work should repeat this experiment with a larger sample size and a range of

heat shock timing and duration to clarify the impact of initial heat stress on physiology.

The response of WA crabs to repeated heat stress demonstrates that negative impacts

of acute sub-lethal stress may be ameliorated by adaptive plasticity. This adaptive plasticity may

manifest in different ways, such as production of molecular chaperones like heat shock proteins

(HSPs) and transcription factors (Somero, Lockwood and Tomanek 2017). After the heat shock

subsides, these heat shock-responsive molecules and regulatory pathways may remain active

in the organism, preparing it for similar conditions within a certain timeframe. While not widely

explored in crustaceans, beneficial priming (*ex.* heat hardening or hormetic priming) has been

documented in arthropods such as *Drosophila melanogaster* (Sejerkilde, Sørensen and

Loeschcke 2003; Sørensen *et al.* 2008; Moghadam *et al.* 2019), as well as a variety of marine

taxa, including seagrasses (Pazzaglia *et al.* 2021), anemones (Glass *et al.* 2023), corals (Martell

2022), shellfish (Abbas *et al.* 2024), and sticklebacks (Spence-Jones *et al.* 2025). Studies have

also shown that priming may be a contributing factor to increased thermal tolerance in non-

indigenous species. Repeated heat stress increased thermal tolerance in *Semimytilus algosus*

and *Mytilus edulis* mussels (Lenz *et al.* 2018) and priming improved heat tolerance in non-native

Bactrocera dorsalis and *Bactrocera correcta* flies (Gu *et al.* 2019). The conditions under which

beneficial plasticity manifests should be evaluated in future experiments by altering the intensity

of heat shocks as well as the time in between the first and second heat shocks.

Righting response is plastic after short- and medium-term cold exposure

Albeit less frequently studied than heat tolerance, cold tolerance often plays an

important role in dictating species distributions (Wasson *et al.* 2025). Organisms that are able to

tolerate cooler conditions may be able to effectively colonize northern latitudes with limited

competition (Walther *et al.* 2009; Christiansen *et al.* 2015; Dijkstra, Westerman and Harris

2017). Our work demonstrates adult green crabs can maintain motor function at 5ºC — with

some maintaining motor function at 1.5ºC — which may allow them a competitive advantage in

colder conditions. These findings are in line with previous research that highlights the extensive

cold tolerance of adult *C. maenas*. For example, *C. maenas* successfully reproduce in sub-arctic

conditions in Newfoundland (Best, McKenzie and Couturier 2017), and can survive at 2ºC for at

least two months (Rivers, McKenzie and McGaw 2025), and below 5ºC for over four months

(Kelley, de Rivera and Buckley 2013). In prior work in a UK population in the native range, some

crabs failed to right at 0ºC, while crabs that maintained movement had a ~5 second average

righting response at 0ºC (Young, Peck and Matheson 2006). We observed an average righting

response of 33.09 s ± 22.58 s (MA) or 33.59 s ± 22.29 s (WA) at 1.5ºC. These apparent

differences in cold tolerance between crab populations may be due in part to genetic or

environmental differences. However, it is likely that methodology plays a role. We included

hesitation time, or the time before moving the fifth pereopod pair after crabs are placed dorsal

side up, as part of the righting response, while Young et al. (2006) does not, which likely

increased our measurements relative to theirs. Nevertheless, it is clear that *C. maenas* has a

robust cold tolerance, which may have facilitated northern range expansion into Newfoundland

(Blakeslee *et al.* 2010) and ongoing spread in Alaska (NOAA 2022).

In addition, *C. maenas* displayed some ability to acclimate to chronic cold stress. At 5ºC,

righting response for several MA and WA crabs sped up at the end of the experiment relative to

earlier timepoints (**Figure S8**). We also observed evidence of acclimation to 1.5ºC: while many

crabs failed to right when initially exposed to 1.5ºC, some individuals "recovered" and were able

to right at a later timepoint (**Figure S8**). This recovery suggests that failure to right is thermally

plastic, and may be related to dormancy in this species. Historical work suggests that when

temperature is ≤5ºC *C. maenas* enter torpor (Berrill 1982; Young, Peck and Matheson 2006), an

energy conservation mechanism typically employed in colder conditions (Pörtner and Playle

1998). Recent work suggests that crabs enter a dormant state (metabolic slowing with limited

movement) which may facilitate energy conservation while also allowing for opportunistic

feeding and locomotive activity (*ex.* burying for predator avoidance) (Rivers, McKenzie and

McGaw 2025). The use of dormancy is also supported by prior research which found a steady

decrease in heart rate with no cardiac breakpoint when cooling crabs from 15ºC to 0ºC,

suggesting that *C. maenas* has sub-zero cold tolerance (Tepolt and Somero 2014). Adult

locomotive plasticity in cold temperatures may be facilitated by additional mechanisms such as

regulation of hemolymph magnesium to prevent calcium channel blockage and restore

locomotor activity (Frederich *et al.* 2000; Aronson *et al.* 2015); increasing the number of

polyunsaturated lipids present in muscle tissue to provide membrane fluidity (Chapelle 1978);

and downregulating the cell cycle regulator *cyclin D1* to reduce cell proliferation (Kelley, de

Rivera and Buckley 2013). In addition to exploring the mechanisms that govern adult dormancy

in cold conditions, attention should be paid to cold tolerance mechanisms throughout life history.

Larval development and survival in *C. maenas* are compromised when temperatures are below

10ºC (Nagaraj 1993; Hines *et al.* 2004; deRivera *et al.* 2007), and modeling work suggests that

cold temperatures have played a key in limiting larval survival and spread in the northeast

Pacific (Du *et al.* 2024). Additional manipulative experiments with larval crabs may clarify

differences in cold tolerance associated with ontogeny.

We explored the relationship between supergene and cold acclimation at the individual

level in part because of a significant population-level association between supergene allele and

cold tolerance in prior work (Tepolt and Palumbi 2020). Surprisingly, in our study the cold-

adapted supergene did not significantly impact TTR at 1.5ºC or 5ºC. This is consistent with

Coyle et al. (2019), who found that failure to right after acute exposure to 4.5ºC was associated

with mitochondrial haplotype, not supergene genotype, in male crabs from Gulf of Maine

populations. Even though supergene distributions were similar between MA and WA

populations, the MA population had a higher proportion of crabs that failed to right during the

experiment. While these differences among populations may be a product of the slightly

different experimental temperatures for MA and WA crabs, it is unlikely that this ~0.5ºC

temperature difference is solely responsible. The switch from an active to dormant state in *C.*

maenas occurs between 4-6ºC (Young, Peck and Matheson 2006; Rivers, McKenzie and

McGaw 2025), which is well above the coldest temperature used in this experiment. While

mitochondrial diversity present in the northwest Atlantic could interact with nuclear diversity to

shape thermal tolerance, to our knowledge there is limited mitochondrial haplotype diversity

within the MA population (Darling *et al.* 2014; Jeffery *et al.* 2017; Lehnert *et al.* 2018). Instead,

we suggest that MA crabs utilize physiological mechanisms that elicit a more pronounced

dormancy than WA crabs as a response to colder winters in the northwest Atlantic (**Figure 1**).

Even though we acclimated all crabs to the same conditions prior to experimentation, there is

evidence that distal environmental history can impact physiological response through within-

and across-generation legacy effects in marine invertebrates (Donelan *et al.* 2020; Ashey and

Rivest 2021; Fernández *et al.* 2025). Future work should account for legacy effects, and

examine responses of crabs during the winter season to ascertain if population differences are a

result of environmental history or lack of seasonal acclimation.

Conclusion

Our work highlights how thermal plasticity contributes to the broad thermal tolerance of

successful non-indigenous species, even after a substantial loss of genetic diversity. We found

that crabs exhibit potentially beneficial plasticity in response to sub-lethal heat stress.

Additionally, crabs display a robust cold tolerance and the potential to acclimate to cold

temperatures over the course of 94 hours. Although previous work demonstrates a strong

association between supergene allele frequency and both cold tolerance and sea surface

temperature at the population level (Tepolt and Palumbi 2020) This suggests that, rather than

being causally linked, environmental conditions like temperature likely shape both supergene

distribution and phenotypic plasticity throughout life history, including during larval dispersal and

settlement. In order to clarify the relationship between population- and individual-level tolerance,

future work should compare if and how the supergene influences thermal tolerance in planktonic

larvae, newly-settled juveniles, and adult crabs.

While we didn't see significant relationships between genotype and individual righting

response, it is possible that supergene genotype plays a role at a different level of biological

organization. Since the association was established using heart rate at $0^{\circ}C$ (Tepolt and Palumbi

2020), it may be useful to examine the relationship between genotype and heart rate plasticity in

individual crabs. Supergene genotype could also impact molecular pathways without changes in

whole-organism physiology. For example, a study in coral larvae detected metabolic

reprogramming under elevated temperature without a decrease in survival (Huffmyer *et al.*

2024), and American lobster (*Homarus americanus*) exposure to ocean acidification resulted in

broad metabolic reprogramming that was not associated with changes to resting metabolism

(Noisette *et al.* 2021). Future work should examine if the warm- and cold-adapted supergene

alleles are associated with differences in metabolic pathways and gene expression that allow for

physiological homeostasis.

Author contributions

YRV, JCK, and CP conceived, designed, and carried out all experiments with assistance from

HLF and input from CKT. YRV, JCK, and JK conducted molecular assays. JCK, CP, and HLF

conducted preliminary data analysis. YRV refined all data analysis. YRV wrote the initial draft

with input from JCK and CP; and YRV and CKT revised and edited this manuscript. All authors

reviewed and approved the final manuscript.

Acknowledgements

We are grateful to Chelsey Buffington and Brian Turner at Washington Fish and Wildlife and

Tait Nygaard and Isabelle See at Quahog Bay Conservancy for providing crabs used in this

experiment, and the PEP and CC-CREW interns that volunteered with data collection. We also

appreciate two anonymous reviewers for their constructive feedback on an earlier draft.

Funding

This work was supported by the NSF Postdoctoral Research Fellowships in Biology Program

under Grant No. 2209018 to YRV. JCK was supported by NSF REU OCE-2150401. CP was

supported by the Woods Hole Partnership Education Program (PEP). HLF was supported by

NSF GEOPATHS grant ICER-2023192 to the Community College Research Experiences at

Woods Hole Oceanographic Institution (CC-CREW) program. JK was supported by the Woods

Hole Oceanographic Institution Academic Programs Office Funds through the Blue Economy

program. Additional project support, including support of CKT, was provided by NSF OCE-

1850996.

Conflict of Interest Statement

The authors have no conflicts of interest to report.

Data availability

All raw data, scripts, results, and supplementary material are available at

https://github.com/yaaminiv/cold-green-crab.

References

Abbas ASA, Collins M, Ellis R *et al.* Heat hardening improves thermal tolerance in abalone, without the trade-offs associated with chronic heat exposure. *J Therm Biol* 2024;**124**:103963.

Aronson RB, Frederich M, Price R *et al.* Prospects for the return of shell-crushing crabs to Antarctica. *J Biogeogr* 2015;**42**:1–7.

Ashey J, Rivest EB. Effect of environmental history on the physiology and acute stress response of the eastern oyster *Crassostrea virginica*. *Mar Ecol Prog Ser* 2021;**674**:115–30.

Barley JM, Cheng BS, Sasaki M *et al.* Limited plasticity in thermally tolerant ectotherm populations: evidence for a trade-off. *Proc Biol Sci* 2021;**288**:20210765.

Bates D, Mächler M, Bolker B *et al.* Fitting linear mixed-effects models Usinglme4. *J Stat Softw* 2015;**67**, DOI: 10.18637/jss.v067.i01.

Berrill M. The life cycle of the Green crab *Carcinus maenas* at the northern end of its range. *J Crustacean Biol* 1982;**2**:31–9.

Best K, McKenzie C, Couturier C. Reproductive biology of an invasive population of European green crab, *Carcinus maenas*, in Placentia Bay, Newfoundland. *Manag Biol Invasion* 2017;**8**:247–55.

Birk MA. *respirometry: Tools for Conducting and Analyzing Respirometry Experiments.*, 2023.

Blakeslee AMH, Keogh CL, Fowler AE *et al.* Assessing the effects of trematode infection on invasive green crabs in eastern north america. *PLoS One* 2015;**10**:e0128674.

Blakeslee AMH, McKenzie CH, Darling JA *et al.* A hitchhiker's guide to the Maritimes: anthropogenic transport facilitates long-distance dispersal of an invasive marine crab to Newfoundland. *Divers Distrib* 2010;**16**:879–91.

Calabria G, Dolgova O, Rego C *et al.* Hsp70 protein levels and thermotolerance in *Drosophila subobscura*: a reassessment of the thermal co-adaptation hypothesis: Testing for thermal co-adaptation in *Drosophila*. *J Evol Biol* 2012;**25**:691–700.

Calvin K, Dasgupta D, Krinner G *et al. IPCC, 2023: Climate Change 2023: Synthesis Report. Contribution of Working Groups I, II and III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [core Writing Team, H. Lee and J. Romero (eds.)]. IPCC, Geneva, Switzerland.* Arias P, Bustamante M, Elgizouli I, et al. (eds.). Intergovernmental Panel on Climate Change (IPCC), 2023.

Carlton JT, Cohen AN. Episodic global dispersal in shallow water marine organisms: the case history of the European shore crabs *Carcinus maenas* and *C. aestuarii. J Biogeogr* 2003;**30**:1809–20.

Chapelle S. The influence of acclimation temperature on the fatty acid composition of an aquatic Crustacean (*Carcinus maenas*). *J Exp Zool* 1978;**204**:337–46.

Chapelle S. Aspects of phospholipid metabolism in crustaceans as related to changes in

environmental temperatures and salinities. *Comp Biochem Physiol B* 1986;**84**:423–39.

Christiansen JS, Sparboe M, Saether B-S *et al.* Thermal behaviour and the prospect spread of
an invasive benthic top predator onto the Euro-Arctic shelves. *Divers Distrib* 2015;**21**:1004–
13.

Compton TJ, Leathwick JR, Inglis GJ. Thermogeography predicts the potential global range of
the invasive European green crab (*Carcinus maenas*). *Divers Distrib* 2010;**16**:243–55.

Coyle AE, Voss ER, Tepolt CK *et al.* Mitochondrial genotype influences the response to cold
stress in the European green crab, *Carcinus maenas*. *J Exp Biol* 2019;**222**:jeb203521.

Cuculescu M, Hyde D, Bowler K. Thermal tolerance of two species of marine crab, *Cancer*
pagurus and *Carcinus maenas*. *J Therm Biol* 1998;**23**:107–10.

Cuenca Cambronero M, Beasley J, Kissane S *et al.* Evolution of thermal tolerance in
multifarious environments. *Mol Ecol* 2018;**27**:4529–41.

Darling JA, Bagley MJ, Roman J *et al.* Genetic patterns across multiple introductions of the
globally invasive crab genus *Carcinus*. *Mol Ecol* 2008;**17**:4992–5007.

Darling JA, Tsai Y-HE, Blakeslee AMH *et al.* Are genes faster than crabs? Mitochondrial
introgression exceeds larval dispersal during population expansion of the invasive crab
Carcinus maenas. *R Soc Open Sci* 2014;**1**:140202.

Davidson AM, Jennions M, Nicotra AB. Do invasive species show higher phenotypic plasticity
than native species and, if so, is it adaptive? A meta-analysis. *Ecol Lett* 2011;**14**:419–31.

deRivera CE, Hitchcock NG, Teck SJ *et al.* Larval development rate predicts range expansion of
an introduced crab. *Mar Biol* 2007;**150**:1275–88.

Dijkstra JA, Westerman EL, Harris LG. Elevated seasonal temperatures eliminate thermal
barriers of reproduction of a dominant invasive species: A community state change for
northern communities? *Divers Distrib* 2017;**23**:1182–92.

Donelan SC, Hellmann JK, Bell AM *et al.* Transgenerational plasticity in human-altered
environments. *Trends Ecol Evol* 2020;**35**:115–24.

Du J, Tepolt CK, Grason EW *et al.* Dispersal pathways of European green crab larvae into and
throughout the eastern Salish Sea. *Prog Oceanogr* 2024;**223**:103245.

El Babili M, Bodennec J, Carsol MA *et al.* Effects of temperature and intracellular pH on the
sphingomyelin metabolism in the gills of crab, *Carcinus maenas*. *Comp Biochem Physiol B*
Biochem Mol Biol 1997;**117**:125–33.

Estoup A, Ravigné V, Hufbauer R *et al.* Is there a genetic paradox of biological invasion? *Annu*
Rev Ecol Evol Syst 2016;**47**:51–72.

Falfushynska HI, Phan T, Sokolova IM. Long-term acclimation to different thermal regimes
affects molecular responses to heat stress in a freshwater clam *Corbicula fluminea*. *Sci*
Rep 2016;**6**:39476.

Fernández F, Guzmán-Rivas F, Urzúa Á *et al.* Maternal investment of kelp crabs from central

Chile is associated with the level of environmental fluctuations and predictability. *Mar Ecol*
Prog Ser 2025;**755**:79–93.

Fields PA, Cox KM, Karch KR. Latitudinal variation in protein expression after heat stress in the
salt marsh mussel *Geukensia demissa*. *Integr Comp Biol* 2012;**52**:636–47.

Fox RJ, Donelson JM, Schunter C *et al.* Beyond buying time: the role of plasticity in phenotypic
adaptation to rapid environmental change. *Philos Trans R Soc Lond B Biol Sci*
2019;**374**:20180174.

Frederich M, Sartoris FJ, Arntz WE *et al.* Haemolymph Mg(2+) regulation in decapod
crustaceans: physiological correlates and ecological consequences in polar areas. *J Exp*
Biol 2000;**203**:1383–93.

Gagnaire P-A, Gaggiotti OE. Detecting polygenic selection in marine populations by combining
population genomics and quantitative genetics approaches. *Curr Zool* 2016;**62**:603–16.

Glass BH, Jones KG, Ye AC *et al.* Acute heat priming promotes short-term climate resilience of
early life stages in a model sea anemone. *PeerJ* 2023;**11**:e16574.

Griffiths JS, Sasaki M, Neylan IP *et al.* The potential for experimental evolution to uncover trade-
offs associated with anthropogenic and climate change adaptation. *Glob Chang Biol*
2024;**30**:e17584.

Gunderson AR, Stillman JH. Plasticity in thermal tolerance has limited potential to buffer
ectotherms from global warming. *Proc Biol Sci* 2015;**282**:20150401.

Gu X, Zhao Y, Su Y *et al.* A transcriptional and functional analysis of heat hardening in two
invasive fruit fly species, *Bactrocera dorsalis* and *Bactrocera correcta*. *Evol Appl*
2019;**12**:1147–63.

Handelsman CA, Broder ED, Dalton CM *et al.* Predator-induced phenotypic plasticity in
metabolism and rate of growth: rapid adaptation to a novel environment. *Integr Comp Biol*
2013;**53**:975–88.

Hendry AP. Key Questions on the Role of Phenotypic Plasticity in Eco-Evolutionary Dynamics. *J*
Hered 2016;**107**:25–41.

Hines A, Ruiz GM, Hitchcock NG *et al.* Projecting Range Expansion of Invasive European
Green Crabs (*Carcinus maenas*) to Alaska: Temperature and Salinity Tolerance of Larvae.
2004.

Hopkin RS, Qari S, Bowler K *et al.* Seasonal thermal tolerance in marine *Crustacea*. *J Exp Mar*
Bio Ecol 2006;**331**:74–81.

Huang B, Liu C, Banzon V *et al.* Improvements of the Daily Optimum Interpolation Sea Surface
Temperature (DOISST) version 2.1. *J Clim* 2021;**34**:2923–39.

Huffmyer AS, Ashey J, Strand E *et al.* Coral larvae increase nitrogen assimilation to stabilize
algal symbiosis and combat bleaching under increased temperature. *PLoS Biol*
2024;**22**:e3002875.

Jeffery NW, DiBacco C, Wringe BF *et al.* Genomic evidence of hybridization between two

independent invasions of European green crab (*Carcinus maenas*) in the Northwest
Atlantic. *Heredity (Edinb)* 2017;**119**:154–65.

Jost JA, Podolski SM, Frederich M. Enhancing thermal tolerance by eliminating the pejus range:
a comparative study with three decapod crustaceans. *Mar Ecol Prog Ser* 2012;**444**:263–74.

Kelley AL, de Rivera CE, Buckley BA. Cold tolerance of the invasive *Carcinus maenas* in the
east Pacific: molecular mechanisms and implications for range expansion in a changing
climate. *Biol Invasions* 2013;**15**:2299–309.

Landy JA, Oschmann A, Munch SB *et al.* Ancestral genetic variation in phenotypic plasticity
underlies rapid evolutionary changes in resurrected populations of waterfleas. *Proc Natl*
Acad Sci U S A 2020;**117**:32535–44.

Lehnert SJ, DiBacco C, Jeffery NW *et al.* Temporal dynamics of genetic clines of invasive
European green crab (*Carcinus maenas*) in eastern North America. *Evol Appl*
2018;**11**:1656–70.

Lenth R, Singmann H, Love J *et al.* Emmeans: Estimated marginal means, aka least-squares
means. *R package version* 2018;**1**:3.

Lenz M, Ahmed Y, Canning-Clode J *et al.* Heat challenges can enhance population tolerance to
thermal stress in mussels: a potential mechanism by which ship transport can increase
species invasiveness. *Biol Invasions* 2018;**20**:3107–22.

Lind BM, Lotterhos KE. The limits of predicting maladaptation to future environments with
genomic data. *Genomics* 2024.

Madeira C, Mendonça V, Flores AAV *et al.* High thermal tolerance does not protect from chronic
warming – A multiple end-point approach using a tropical gastropod, *Stramonita*
haemastoma. *Ecol Indic* 2018;**91**:626–35.

Ma L-J, Cao L-J, Chen J-C *et al.* Rapid and repeated climate adaptation involving chromosome
inversions following invasion of an insect. *Mol Biol Evol* 2024;**41**:msae044.

Martell HA. Thermal priming and bleaching hormesis in the staghorn coral, *Acropora cervicornis*
(Lamarck 1812). *J Exp Mar Bio Ecol* 2022:151820.

Moghadam NN, Ketola T, Pertoldi C *et al.* Heat hardening capacity in *Drosophila melanogaster*
is life stage-specific and juveniles show the highest plasticity. *Biol Lett* 2019;**15**:20180628.

Morley SA, Peck LS, Sunday JM *et al.* Physiological acclimation and persistence of ectothermic
species under extreme heat events. *Glob Ecol Biogeogr* 2019;**28**:1018–37.

Nagaraj M. Combined effects of temperature and salinity on the zoeal development of the green
crab, *Carcinus maenas* (Linnaeus, 1758) (*Decapoda*: *Portunidae*). *Sci Mar* 1993.

NOAA. Green Crab Detected in Alaska for the First Time. *NOAA* 2022.

Noisette F, Calosi P, Madeira D *et al.* Tolerant Larvae and Sensitive Juveniles: Integrating
Metabolomics and Whole-Organism Responses to Define Life-Stage Specific Sensitivity to
Ocean Acidification in the American Lobster. *Metabolites* 2021;**11**, DOI:
10.3390/metabo11090584.

Oliver MK, Piertney SB. Selection maintains MHC diversity through a natural population
bottleneck. *Mol Biol Evol* 2012;**29**:1713–20.

Pazzaglia J, Badalamenti F, Bernardeau-Esteller J *et al.* Thermo-priming increases heat-stress
tolerance in seedlings of the Mediterranean seagrass *P. oceanica*. *Mar Pollut Bull*
2021;**174**:113164.

Pörtner HO. Climate variations and the physiological basis of temperature dependent
biogeography: systemic to molecular hierarchy of thermal tolerance in animals. *Comp*
Biochem Physiol A Mol Integr Physiol 2002;**132**:739–61.

Pörtner H-O, Bock C, Mark FC. Oxygen- and capacity-limited thermal tolerance: bridging
ecology and physiology. *J Exp Biol* 2017;**220**:2685–96.

Pörtner HO, Langenbuch M, Michaelidis B. Synergistic effects of temperature extremes,
hypoxia, and increases in CO_2 on marine animals: From Earth history to global change. *J*
Geophys Res C: Oceans 2005;**110**.

Pörtner HO, Playle RC. Temperature and energetics: an introduction to cold ocean physiology.
In: Pörtner HO, Playle RC (eds.). *Cold Ocean Physiology*. Cambridge: Cambridge
University Press, 1998, 3–30.

R Core Team. *R: A Language and Environment for Statistical Computing*. R Foundation for
Statistical Computing, Vienna, Austria, 2024.

Riginos C, Crandall ED, Liggins L *et al.* Navigating the currents of seascape genomics: how
spatial analyses can augment population genomic studies. *Curr Zool* 2016;**62**:581–601.

Rivers ML, McKenzie CH, McGaw IJ. Effects of low temperature exposure and acclimation on
the behavioural responses of the green crab (*Carcinus maenas*) from Newfoundland,
Canada. *J Therm Biol* 2025:104071.

Rocca KAC, Gray EM, Costantini C *et al.* 2La chromosomal inversion enhances thermal
tolerance of Anopheles gambiae larvae. *Malar J* 2009;**8**:147.

Roman J. Diluting the founder effect: cryptic invasions expand a marine invader's range. *Proc*
Biol Sci 2006;**273**:2453–9.

Roman J, Darling JA. Paradox lost: genetic diversity and the success of aquatic invasions.
Trends Ecol Evol 2007;**22**:454–64.

Roman J, Palumbi SR. A global invader at home: population structure of the green crab,
Carcinus maenas, in Europe. *Mol Ecol* 2004;**13**:2891–8.

Sanford E, Kelly MW. Local adaptation in marine invertebrates. *Ann Rev Mar Sci* 2011;**3**:509–
35.

Sasaki MC, Dam HG. Negative relationship between thermal tolerance and plasticity in
tolerance emerges during experimental evolution in a widespread marine invertebrate. *Evol*
Appl 2021;**14**:2114–23.

Schwander T, Libbrecht R, Keller L. Supergenes and complex phenotypes. *Curr Biol*
2014;**24**:R288–94.

Sejerkilde M, Sørensen JG, Loeschcke V. Effects of cold- and heat hardening on thermal
resistance in *Drosophila melanogaster*. *J Insect Physiol* 2003;**49**:719–26.

Somero GN. The physiology of climate change: how potentials for acclimatization and genetic
adaptation will determine "winners" and "losers." *J Exp Biol* 2010;**213**:912–20.

Somero GN. The physiology of global change: linking patterns to mechanisms. *Ann Rev Mar Sci*
2012;**4**:39–61.

Somero GN, Lockwood B, Tomanek L. *Biochemical Adaptation*. 1st ed. New York, NY: Oxford
University Press, 2017.

Sørensen JG, Sarup P, Kristensen TN *et al.* Temperature-induced hormesis in *Drosophila*. *Mild
Stress and Healthy Aging*. Dordrecht: Springer Netherlands, 2008, 65–79.

Sorte CJB, Ibáñez I, Blumenthal DM *et al.* Poised to prosper? A cross-system comparison of
climate change effects on native and non-native species performance. *Ecol Lett*
2013;**16**:261–70.

Spence-Jones HC, Scheibl C, Pein CM *et al.* Do you remember? Within-generation and
transgenerational heat stress memory of recurring marine heatwaves in threespine
stickleback. *Proc Biol Sci* 2025;**292**:20242913.

Styrishave B, Rewitz K, Andersen O. Frequency of moulting by shore crabs *Carcinus maenas*
(L.) changes their colour and their success in mating and physiological performance. *J Exp
Mar Bio Ecol* 2004;**313**:317–36.

Swindell WR, Bouzat JL. Modeling the adaptive potential of isolated populations: experimental
simulations using *Drosophila*. *Evolution* 2005;**59**:2159–69.

Tam V, Patel N, Turcotte M *et al.* Benefits and limitations of genome-wide association studies.
Nat Rev Genet 2019;**20**:467–84.

Teixeira JC, Huber CD. The inflated significance of neutral genetic diversity in conservation
genetics. *Proc Natl Acad Sci U S A* 2021;**118**, DOI: 10.1073/pnas.2015096118.

Tepolt C. Thermal biology. In: Weihrauch D, McGaw IJ (eds.). *Ecophysiology of the European
Green Crab (Carcinus Maenas) and Related Species*. Elsevier, 2024, 231–47.

Tepolt CK, Darling JA, Bagley MJ *et al.* European green crabs (*Carcinus maenas*) in the
northeastern Pacific: genetic evidence for high population connectivity and current-
mediated expansion from a single introduced source population. *Diversity and Distributions*
2009;**15**:997–1009.

Tepolt CK, Grosholz ED, de Rivera CE *et al.* Balanced polymorphism fuels rapid selection in an
invasive crab despite high gene flow and low genetic diversity. *Mol Ecol* 2022, DOI:
10.1111/mec.16143.

Tepolt CK, Palumbi SR. Transcriptome sequencing reveals both neutral and adaptive genome
dynamics in a marine invader. *Mol Ecol* 2015;**24**:4145–58.

Tepolt CK, Palumbi SR. Rapid Adaptation to Temperature via a Potential Genomic Island of
Divergence in the Invasive Green Crab, *Carcinus maenas*. *Frontiers in Ecology and*

Evolution 2020;**8**:411.

Tepolt CK, Somero GN. Master of all trades: thermal acclimation and adaptation of cardiac
function in a broadly distributed marine invasive species, the European green crab,
Carcinus maenas. *J Exp Biol* 2014;**217**:1129–38.

Walther G-R, Roques A, Hulme PE *et al.* Alien species in a warmer world: risks and
opportunities. *Trends Ecol Evol* 2009;**24**:686–93.

Wang H, Yang Z, Wang S *et al.* Genome-wide association analysis reveals the genetic basis of
thermal tolerance in dwarf surf clam *Mulinia lateralis*. *Genomics* 2024;**116**:110904.

Wasson K, Cressman K, Beheshti K *et al.* Setting the limit: Cold rather than hot temperatures
limit intertidal distribution of a coastal foundation species. *Mar Environ Res* 2025:107149.

Weihrauch D, McGaw I eds. *Ecophysiology of the European Green Crab (*Carcinus maenas*)*
and Related Species: Mechanisms behind the Success of a Global Invader. San Diego,
CA: Academic Press, 2024.

Wickham H, Kuhn M, Vaughan D. *Generics: Common S3 Generics Not Provided by Base R*
Methods Related to Model Fitting., 2022.

Wittmann AC, Held C, Pörtner HO *et al.* Ion regulatory capacity and the biogeography of
Crustacea at high southern latitudes. *Polar Biol* 2010;**33**:919–28.

Yao C-L, Somero GN. The impact of ocean warming on marine organisms. *Chin Sci Bull*
2014;**59**:468–79.

Young JS, Peck LS, Matheson T. The effects of temperature on walking and righting in
temperate and Antarctic crustaceans. *Polar Biol* 2006;**29**:978–87.

Zerebecki RA, Sorte CJB. Temperature tolerance and stress proteins as mechanisms of
invasive species success. *PLoS One* 2011;**6**:e14806.

Zhang Y, Liu S, De Meyer M *et al.* Genomes of the cosmopolitan fruit pest *Bactrocera dorsalis*
(*Diptera*: *Tephritidae*) reveal its global invasion history and thermal adaptation. *J Adv Res*
2023;**53**:61–74.

Zou D, Cao W, Liu G *et al.* Effects of acute and chronic thermal stress on survival, apoptosis,
and transcriptional responses of *Scapharca broughtonii*. *J Oceanol Limnol* 2023, DOI:
10.1007/s00343-022-2092-3.

Figure Captions

Figure 1. **A**) Green crab collection sites for experiments (purple diamond = Willapa Bay, WA; blue triangle = Buzzards Bay, MA; and green square = Harpswell Sound, ME). **B**) Average sea surface temperature (SST) for 2021-2024 near collection sites (solid purple line = Willapa Bay, WA; long blue dashes = Buzzards Bay, MA; small green dashes = Harpswell Sound, ME). SST data were obtained from NOAA's OI SST V2 High Resolution Dataset provided by the NOAA PSL, Boulder, Colorado, USA, from their website at https://psl.noaa.gov (Huang *et al.* 2021). A four-year mean, SST minimum, and SST maximum were calculated for each month. Solid lines represent four-year averages, while ribbons represent ranges.

Figure 2. Righting response before (0 hours) and after (22 hours) heat shock by **A**) population and **B**) population and genotype. In all panels, TTR is represented by green squares for ME *C. maenas* and purple triangles for WA *C. maenas*. Significantly different pairwise comparisons are indicated on the figure (**** = *P*-value < 0.0001).

Figure 3. **A**) Average TTR across all timepoints for control (grey) and treatment (red) crabs. The inset shows data for crabs whose TTR was < 4 seconds. Timing of the heat shocks are indicated by transparent red boxes. Day 1 measurements were taken after the first heat shock for the treatment crabs, and day 4 measurements were taken after the second heat shock experienced by all crabs. Pairwise differences between days are indicated within control (grey uppercase) and treatment (red lowercase) treatments. **B**) Average TTR by genotype.

Figure 4. **A**) Average time-to-right for MA *C. maenas* in 1.5ºC (dark blue squares) and 5ºC (light blue circles) temperature treatments; crabs that failed to right in 90 s were not included in these data Righting response was significantly impacted by temperature, time, and their interaction. Pairwise differences between timepoints are indicated within 1.5ºC (dark blue lowercase) and 5ºC (light blue uppercase) treatments. **B**) Average TTR by genotype. **C**) Proportion of MA crabs that righted within 90 seconds. **D**) Proportion of MA crabs at 1.5ºC that failed to right with and without a T allele. In panels **C**) and **D**), the dark blue color represents crabs that failed to right in 90 s. Failure to right was significantly impacted by temperature and time.

Figure 5. **A**) Average time-to-right for WA *C. maenas* in 1.5ºC (dark blue squares) and 5ºC (light blue circles) temperature treatments; crabs that failed to right in 90 s were not included in these data. Righting response was significantly impacted by temperature, timepoint, and their interaction. Pairwise differences between timepoints are indicated within 1.5ºC (dark blue lowercase) and 5ºC (light blue uppercase) treatments. **B**) Average TTR by genotype. **C**) Proportion of MA crabs that righted within 90 seconds. **D**) Proportion of MA crabs at 1.5ºC that failed to right with and without a T allele. In panels **C**) and **D**), the dark blue color represents crabs that failed to right in 90 s. Failure to right was significantly impacted by temperature and time.

A. Collection Sites

B. Average SST (2021–2024)

Willapa Bay, WA
Buzzards Bay, MA
Harpswell Sound, ME

A. Average Righting Response

B. Righting Response by Genotype

A. Righting Response for MA Crabs

B. Righting Response by Genotype

C. Proportion Righted

D. Righting by Genotype

A. Righting Response for WA Crabs

B. Righting Response by Genotype

C. Proportion Righted

D. Righting by Genotype