Structural modeling reveals the allosteric switch controlling the chitin utilization program of *Vibrio cholerae*

Holly S. Anderson[1] and Ankur B. Dalia[1,*]

[1]Department of Biology, Indiana University, Bloomington, Indiana, United States of America
*Author for correspondence: ankdalia@iu.edu

ABSTRACT

Signal transduction by histidine kinases (HKs) is nearly ubiquitous in bacterial species. HKs can either sense ligands directly or indirectly via a cognate substrate binding protein (SBP). The molecular basis for SBP-dependent signal reception, however, remains poorly understood in most cases. CBP and ChiS are the SBP-HK pair that activate the chitin utilization program of *Vibrio cholerae*. Here, we elucidate the molecular basis for allosteric regulation of CBP-ChiS by generating structural models of this complex in the unliganded and liganded states, which we support with extensive genetic, biochemical, and cell biological analysis. Our results reveal that ligand-binding induces a large conformational interface switch that is distinct from previously described SBP-HKs. Structural modeling suggests that similar interface switches may also regulate other uncharacterized SBP-HKs. Together, these results extend our understanding of signal transduction in bacterial species and highlight a new approach for uncovering the molecular basis of allostery in protein complexes.

INTRODUCTION

The ability to sense and respond appropriately to environmental stimuli is critical in all domains of life. One sensory system that is ubiquitous in bacterial species is signal transduction by histidine kinases (HKs)[1,2]. HKs can sense periplasmic signals either directly[3] or indirectly via a solute binding protein (SBP)[4]. While genetic studies have linked SBPs to activation of cognate HKs[4-8], the molecular mechanism for activation remains poorly understood in most cases. This is due, in part, to the challenges associated with structural characterization of transmembrane proteins like HKs. As a result, the current understanding of SBP-HK activation is informed by a relatively limited number of structural studies[9-12]. This work has revealed that the molecular basis of SBP-HK interactions differs greatly between systems, indicating the lack of a broadly conserved allosteric activation mechanism.

While activation of HKs by liganded SBPs is well-studied, the influence of unliganded SBPs on HK regulation has been overlooked in most cases. Paradigmatically, unliganded SBPs interact passively with their cognate HKs and then allosterically induce activation through a conformational change upon ligand-binding[9-12]. The chitin-sensing HK ChiS, which regulates the expression of genes required for chitin catabolism and natural transformation in *V. cholerae*[13-15], offers a unique and unexplored example of SBP-HK regulation. Specifically, prior genetic and cell biological analysis indicates that ChiS is subject to dual allosteric regulation by the SBP chitin-binding protein (CBP) wherein ChiS is repressed by unliganded CBP[15] and is conversely activated by liganded (chitin-bound) CBP[6]. Here, we sought to define the molecular basis underlying this dual allosteric regulation to expand our understanding of bacterial signal transduction.

RESULTS
AlphaFold predicts the CBP-ChiS complex in two distinct conformations

To help define the molecular basis for CBP-dependent allosteric regulation of ChiS activity, we sought to generate structural models of CBP-ChiS using AlphaFold-Multimer v3 (AF-Mv3)[16-18] and AlphaFold3 (AF3)[19]. Because HKs almost ubiquitously form homodimers[1], we modeled two copies of ChiS, and we empirically found that structural modeling with one CBP yielded the highest confidence models for both algorithms. AF-Mv3 and AF3 modeled ChiS as a homodimer, as expected, with CBP cradled at its apex (**Fig. S1**). Interestingly, the CBP-ChiS[AF-M] and CBP-ChiS[AF3] models predicted distinct orientations for CBP (**Fig. S1A-B, Movie S1**) despite having similar confidence scores (**Fig. S1D**). Since CBP differentially regulates ChiS activity depending on its liganded state[6], we hypothesized that these two CBP-ChiS models represented the unliganded (ChiS repressed) and liganded (ChiS active) conformations. To test this hypothesis, we aligned CBP from the top-ranked CBP-ChiS[AF-M] and CBP-ChiS[AF3] models to crystal structures of unliganded (PDB: 4gf8) and liganded (PBD: 4gfr) CBP. This analysis revealed that CBP[AF-M] closely aligned to liganded CBP, while CBP[AF3] aligned to unliganded CBP (**Fig. S2**). Thus, we hypothesized that the CBP-ChiS[AF-M] model represents the liganded (chitin-bound) conformation where ChiS is active, while the CBP-ChiS[AF3] model represents the unliganded conformation where ChiS is repressed (**Fig. 1**).

Comparison of the CBP-ChiS[AF3] and CBP-ChiS[AF-M] models reveals electrostatic interactions unique to each model

To formally test this hypothesis, we sought to identify interactions unique to each model for targeted mutagenesis. We focused on identifying residue pairs that putatively form electrostatic interactions, because they allow for a high degree of specificity in structure-function analysis. Specifically, mutating these residues to the opposite charge should break interactions through electrostatic repulsion. However, if these residues truly interact, flipping the charge on both sides of the interface should restore the electrostatic interaction. Thus, we analyzed the top-ranked models from five independent CBP-ChiS[AF-M] and CBP-ChiS[AF3] sessions to identify electrostatic interactions at the CBP-ChiS interface. Interactions predicted in ≥3 models that were <4 Å in one conformation but separated by >10 Å in the other conformation (**Fig. S3A, Dataset S1**) were selected for further characterization. This analysis revealed one electrostatic interaction specific to the CBP-ChiS[AF3] model (unliganded / repressed conformation) and five interactions specific to the CBP-ChiS[AF-M] model (liganded / active conformation) (**Fig. S3B-C**).

Interactions specific to the CBP-ChiS[AF3] (unliganded) model are critical for CBP to allosterically repress ChiS in the absence of chitin

In the absence of chitin, unliganded CBP interacts with ChiS to allosterically repress its activity[6,15]. We hypothesized that interactions specific to the CBP-ChiS[AF3] model (**Fig. 1, S3**) are necessary for CBP-dependent repression of ChiS in the absence of chitin. Thus, flipping the charge of residues unique to the CBP-ChiS[AF3] interface should disrupt CBP's ability to repress ChiS, resulting in increased ChiS activity in rich media (*i.e.*, absence of chitin). Since CBP is directly regulated by ChiS[6], a functional CBP-mCherry fusion under the control of its native promoter (P$_{chb}$-*cbp-mCherry*) is used as a reporter for ChiS activity throughout this manuscript. The mCherry fluorescence is normalized to a constitutively expressed mTFP1 construct (P$_{const2}$-*mTFP1*) to control for intrinsic noise in gene expression. Consistent with unliganded CBP repressing ChiS in the absence of chitin, ChiS activity is low when CBP is intact and inactivation of CBP (CBP$^\Delta$) results in elevated ChiS activity (**Fig. 2A**).

Mutating either side of the electrostatic interaction specific to CBP-ChiS[AF3] (unliganded) (*cbp*[E401R] or *chiS*[R141E,K145E]) results in a significant increase in ChiS activity (**Fig. 2A-B**), which is consistent with these residues playing an important role in CBP-dependent repression of ChiS. To test whether these residues interact, we flipped the charge on both sides of the interface to restore the electrostatic interaction. In *cbp*[E401R] *chiS*[R141E,K145E], CBP-dependent repression of

ChiS is almost completely restored (**Fig. 2A**). These data strongly support the hypothesis that CBP[E401]-ChiS[R141,K145] form an electrostatic interaction and that CBP-ChiS[AF3] represents the unliganded (repressed) complex conformation.

An electrostatic interaction specific to CBP-ChiS[AF3] (unliganded) contributes to CBP-ChiS interactions

Unliganded CBP directly interacts with the periplasmic domain of ChiS to repress its activity[6]. Thus, we hypothesized that mutating the CBP[E401]-ChiS[R141,K145] interface might prevent CBP-dependent repression by disrupting CBP-ChiS interactions. To test this, we used PopZ-linked apical recruitment (POLAR) assays[20] to assess CBP-ChiS interactions by fluorescence microscopy[6]. In these assays, the "bait" and "prey" proteins are tagged with a fluorescent protein and the bait also contains an H3H4 PopZ-interaction domain. Upon induction of PopZ, the bait (ChiS-msfGFP-H3H4) relocalizes to the cell pole and the coincident relocalization of the prey (CBP-mCherry) demonstrates a direct interaction[20]. Consistent with CBP-ChiS interactions, we see that CBP-mCherry and ChiS-msfGFP-H3H4 colocalize and are both relocalized to the cell pole when PopZ is induced (**Fig. 2C**). A truncated allele of CBP (CBP$^\Delta$ = CBP signal sequence-mCherry fusion) does not interact with ChiS and remains peripherally localized, consistent with its periplasmic localization (**Fig. 2C**). When we flip the charge on either side of the putative unliganded interface (*cbp[E401R]* or *chiS[R141E,K145E]*), CBP-ChiS interactions are partially disrupted (**Fig. 2C**). Importantly, CBP-ChiS interactions are restored when the charge is flipped on both sides of the interface (*cbp[E401R] chiS[R141E,K145E]*), which is consistent with restoration of the electrostatic interaction (**Fig. 2C**). Together, these results suggest that the electrostatic interaction between CBP[E401]-ChiS[R141,K145] is necessary for CBP-dependent repression in the absence of chitin because it contributes to CBP-ChiS interactions.

Interactions specific to the ChiS[AF-M] (liganded) model are necessary for CBP to allosterically activate ChiS in the presence of chitin

If CBP-ChiS[AF-M] represents the liganded (active) conformation, we hypothesize that interactions unique to this model (**Fig. S3**) should be dispensable for CBP-dependent repression of ChiS in the absence of chitin. Consistent with this, we found that flipping the charge of residues uniquely found at the CBP-ChiS[AF-M] interface (*cbp[D231R]*, *cbp[D251R]*, *cbp[R254E]*, *cbp[E529R,E530R]*, *chiS[R81E]*, *chiS[R89E]*, *chiS[R99D]*, *chiS[R146D]*) did not impact CBP-dependent repression of ChiS in the absence of chitin with one exception (*chiS[R139E]*), which partially disrupted repression (**Fig. S4**). Thus, mutations to the CBP-ChiS[AF-M] interface do not disrupt CBP-ChiS interactions in the unliganded state.

In the presence of chitin, liganded CBP allosterically activates ChiS activity[6]. If CBP-ChiS[AF-M] represents the liganded (active) conformation, then mutating interactions specific to this interface should prevent chitin-dependent activation of ChiS. One potentially confounding issue in testing this hypothesis is that any disruption to CBP-ChiS interactions inherently increases ChiS activity, even in the absence of chitin (**Fig. 2A**). Contrary to most HKs, ChiS is active when it is unphosphorylated, and prior work demonstrates that liganded CBP allosterically activates ChiS by inhibiting its kinase activity[6]. Specifically, a ChiS phosphatase mutant (*chiS[T473A]*) can only be activated in the presence of both CBP and chitin[6], a finding that we recapitulate here (**Fig. S5**). Additionally, mutations to the CBP ligand-binding pocket (*cbp[W389A,W539A]*), that inhibit chitin-binding[21], prevent chitin-dependent activation of *chiS[T473A]* (**Fig. S5**). This strongly suggests that ChiS[T473A] requires allosteric activation by liganded CBP (whereas ChiS[WT] does not). Thus, the *chiS[T473A]* allele was used to assess allosteric, chitin-dependent activation of ChiS moving forward.

Consistent with CBP-ChiS^{AF-M} representing the liganded (active) conformation, we found that disrupting most of the electrostatic interactions unique to this model inhibited chitin-dependent activation (*cbp^{D231R}*, *cbp^{D251R}*, *cbp^{E401K}*, *chiS^{T473A,R89E}*, *chiS^{T473A,R99D}*, *chiS^{T473A,R139E}*, *chiS^{T473A,R146D}*) (**Fig. 3, S7**). Furthermore, flipping the charge on both sides of the interface restored chitin-dependent activation in all cases. For example, flipping the charge on either side of the CBP^{D251}-ChiS^{R146} interface ablates chitin-dependent activation, while flipping the charge on both sides of the interface restores chitin-dependent activation to parental levels (**Fig. 3**). Similarly, flipping the charge on both sides of the interface restored chitin-dependent activation for CBP^{D231}-ChiS^{R139}, CBP^{E529,E530}-ChiS^{R99}, and CBP^{E401}-ChiS^{R89} (**Fig. S7**). Importantly, the ChiS mutations assessed are not simply gain-of-function alleles (**Fig. S6**), indicating that this restoration is indicative of a direct electrostatic interaction between these residues.

Together, these results suggest that interactions specific to CBP-ChiS^{AF-M} are dispensable for CBP-dependent allosteric repression of ChiS when chitin is absent, but necessary for allosteric activation of ChiS in the presence of chitin. This genetic analysis helps validate the hypothesis that CBP-ChiS^{AF-M} represents the conformation where ChiS is activated by chitin-bound CBP.

Residue partner-switching facilitates CBP-dependent allosteric dual-regulation of ChiS
Thus far, we provide strong evidence for the residue-level accuracy of both the CBP-ChiS^{AF3} (unliganded) and CBP-ChiS^{AF-M} (liganded) models (**Fig. 2, Fig. 3**). Since ChiS activity is induced when cells are exposed to chitin, the CBP-ChiS complex must transition from the unliganded conformation to the liganded conformation when CBP binds chitin (**Movie S1**). Like many SBPs, CBP undergoes a large conformational change when binding to its chitin ligand[22,23], which may facilitate this transition. CBP^{E401} participates at the interface of both the unliganded and liganded conformations (**Fig. 2, Fig. S7**), but maintains unique interacting partners in each model: ChiS$^{R141\ K145}$ in CBP-ChiS^{AF3} (unliganded) (**Fig. 2, 4A**) and ChiS^{R89} in CBP-ChiS^{AF-M} (**Fig. S7, 4A**). Thus, CBP^{E401} must "partner-switch", which we hypothesize is critical for the transition between unliganded (repressed) and liganded (active) conformations.

If CBP^{E401} undergoes partner-switching, then favoring an electrostatic interaction in the unliganded conformation should prevent the complex from transitioning to the liganded (active) state. To test this, we revisited the charge-flip mutants for CBP^{E401}-ChiS^{R141,K145}. Flipping the charge on either side of the interface (*cbp^{E401R}* or *chiS^{R141E,K145E}*) discourages only the unliganded conformation (**Fig. 2, S7**). Accordingly, these mutations do not prevent chitin-dependent activation (**Fig. 4B**). However, flipping the charge on both sides of the interface (*cbp^{E401R} chiS^{R141E,K145E,T473A}*) should restore the interaction in the unliganded conformation (*i.e.*, restoring CBP^{E401}-ChiS^{R141,K145}) while simultaneously discouraging the interaction in the liganded conformation (*i.e.*, disrupting CBP^{E401}-ChiS^{R89}). This should skew the complex towards the unliganded (repressed) conformation by preventing the partner-switch mediated transition to the liganded (active) conformation. Indeed, chitin-dependent activation is abolished in *cbp^{E401R} chiS^{R141E,K145E,T473A}* (**Fig. 4B**). These results further validate the CBP-ChiS^{AF3} (unliganded) and CBP-ChiS^{AF-M} (liganded) models and suggest that residue partner-switching is a critical feature of the conformational transition associated with chitin-dependent allosteric activation.

Disulfide locking CBP-ChiS into the liganded (active) conformation induces ChiS activity in the absence of chitin
Thus far, our data supports the hypothesis that CBP and ChiS form a dynamic complex, that transitions between the unliganded (repressed) and liganded (active) conformations (**Movie S1**). Above, we show that skewing CBP-ChiS interactions in favor of the unliganded conformation likely prevents the structural transition associated with chitin-dependent activation. Conversely, we hypothesized that if we could skew CBP-ChiS interactions in favor of the liganded

conformation, we could artificially induce ChiS activity. To that end, we introduced cysteines in CBP and ChiS at positions that are closely interacting in CBP-ChiS^{AF-M} (liganded) to "lock" the complex into the liganded (active) conformation via a disulfide bond. Specifically, we introduced cysteines at CBP^{D310} and ChiS^{Q243}, whose α-carbons are <10 Å apart in CBP-ChiS^{AF-M}, but >20 Å apart in CBP-ChiS^{AF3} (**Dataset 1, Fig. 5A**). As a negative control, CBP^{D310C} was paired with ChiS^{Y245C}, a position that is not predicted to interact with CBP^{D310} in either the unliganded or liganded conformations (**Fig. 5A**).

First, disulfide crosslinking was assessed by non-reducing Western blots. When cells harbor only a single cysteine knock-in (CBP^{D310C} or ChiS^{Q243C}), disulfide crosslinking is not observed (**Fig. 5B**), despite robust CBP-ChiS interactions in these strains as evidenced by NHS-ester crosslinking with disuccinimidyl suberate (DSS) (**Fig. 8A**). However, in cells harboring both CBP^{D310C} and ChiS^{Q243C}, robust disulfide crosslinking is observed (**Fig. 5B**). Importantly, disulfide crosslinks are not observed in the CBP^{D310C} ChiS^{Y245C} negative control (**Fig. 5B**), despite robust CBP-ChiS interactions (**Fig. 8A**). Importantly, CBP^{D310C}-ChiS^{Q243C} disulfide crosslinking occurs in cells grown in rich medium (*i.e.,* in the absence of chitin). This suggests that CBP-ChiS dynamically samples the active conformation even in the absence of its chitin ligand, and that disulfide crosslinking of CBP^{D310C}-ChiS^{Q243C} locks the complex into the liganded conformation.

Next, we tested whether the liganded conformation is sufficient for ChiS activity. We found that *cbp^{D310C} chiS^{T473A,Q243C}* exhibited strong ChiS activity, even in the absence of chitin (**Fig. 5C**). Conversely, this was not observed for strains harboring only a single cysteine (*cbp^{D310C}* or *chiS^{T473A,Q243C}*) or the non-disulfide crosslinked cysteine pair (*cbp^{D310C}-chiS^{T473A,Y245C}*) (**Fig. 5C**). Importantly, the *chiS^{Q243C}* and *chiS^{Y245C}* cysteine mutations do not diminish ChiS activity (**Fig. S4**). Together, these results strongly suggest that stabilizing the CBP-ChiS^{AF-M} (liganded) conformation, even in the absence of chitin, is sufficient to induce ChiS activity. Moreover, they suggest that the CBP-ChiS complex dynamically samples both the unliganded and liganded conformations, with the latter likely being stabilized when CBP binds to its chitin ligand.

Transitioning from the unliganded to liganded conformation is critical for ChiS-induced behaviors
ChiS activates both the chitin utilization program[6,13,15] and a small RNA required for horizontal gene transfer by natural transformation[24]. Our results suggest that in the presence of chitin, CBP-ChiS transitions from the unliganded conformation to the liganded conformation to induce ChiS activity. To test whether this transition is required for the ChiS-regulated behaviors, we assessed our mutants for growth on insoluble chitin and for chitin-induced natural transformation. In particular, we tested the phenotype of charge flip mutations at both the liganded interface, CBP^{D251}-ChiS^{R146} (**Fig. 3**), and the unliganded interface, CBP^{E401}-ChiS^{R141,K145} (**Fig. 4**). We found that the ChiS-dependent activity of these mutants directly correlated with chitin growth (**Fig. 6A**) and natural transformation (**Fig. 6B**).

AF-Mv3 predicts distinct conformations for other SBP-HK signaling complexes
Our results suggest that AlphaFold can reveal the distinct conformations associated with allosteric activation of CBP-ChiS. Further analysis revealed that AF-Mv3 alone could reveal the two distinct conformations of CBP-ChiS (**Fig. S9**). We therefore sought to determine whether AF-Mv3 could reveal the allosteric activation mechanism for other SBP-HK signaling complexes that remain structurally uncharacterized[4]. We found that AF-Mv3 predicted two distinct conformations for *Agrobacterium tumefaciens* ChvE-VirA[25] and *Bordetella pertussis* BctC-BctE[8] (**Fig. S9**). Like CBP-ChiS, the orientation of the SBP changes with respect to the HK to produce two distinct conformations for ChvE-VirA and BctC-BctE (**Fig. S9, S10A, Movie S2-S3**). Thus, it

is tempting to speculate that these distinct conformations represent the unliganded and liganded states of these signaling complexes.

While the molecular mechanism of BctC-BctE signal transduction remains poorly understood, ChvE-VirA is relatively well-characterized. In particular, one study has defined a number of ChvE mutant alleles that alter allosteric activation of VirA[5]. Specifically, it uncovered ChvE mutations that either (1) prevented VirA activation in the presence of ligand or (2) induced VirA activity in the absence of ligand. We hypothesize that the former represent residues required for the liganded interface, while the latter represent residues that stabilize the unliganded interface (and thus, their mutation skews the complex into the liganded/active conformation). Consistent with this, these residues lie at the ChvE-VirA interface in the two conformations predicted by AF-Mv3 (**Fig. S10**). The residues required for ChvE-VirA activation in the presence of ligand are closer to the interface of one model, while the residues that, when mutated, induce ChvE-VirA activation in the absence of ligand are closer to the interface of the other model (**Fig. S10C**, **Movie S2**), which supports their opposing effects on ChvE-VirA activity. Furthermore, analyzing both ChvE-VirA conformations for predicted contacts reveals evidence for residue partner-switching (**Fig. S10D**) akin to what is observed for CBP-ChiS (**Fig. 4**). Importantly, mutation of the ChvE residues involved in this partner-switching have opposing phenotypic effects, where mutation of one residue, ChvE[L181], prevents activation in the presence of ligand, while mutation of the other, ChvE[A186], induces VirA activity in the absence of ligand[5]. All together, this analysis provides compelling evidence for the biological relevance of the two ChvE-VirA conformations predicted by AF-Mv3. More broadly, these results highlight how AF-M modeling can help reveal the molecular basis for allosteric signal transduction.

DISCUSSION

Through a combination of structural modeling and complementary genetic and biochemical approaches, this study has uncovered the molecular mechanism of the dual allosteric switch regulating the chitin utilization program of *V. cholerae*. Our results indicate that the CBP-ChiS complex undergoes a large conformational change upon ligand-binding to induce the activity of the ChiS sensor kinase. More broadly, this work highlights how structural modeling combined with structure-function analysis can rapidly reveal the molecular basis for signal transduction.

SBP-HK complexes facilitate signal transduction to a variety of environmental signals[4]. In these systems, liganded SBPs must induce a conformational change in their cognate HK to induce their activity. The molecular basis for this activity, however, varies widely[9-12], highlighting the lack of a broadly conserved allosteric activation mechanism. All currently characterized SBP-HK systems adopt a 2:2 heterotetrameric complex in the liganded form. By contrast, our data suggests that liganded CBP-ChiS uniquely adopts a 1:2 stoichiometry. This is supported by the higher model confidence of 1:2 models, and the fact that 2:2 models lacked the contacts we demonstrate are critical for chitin-dependent activation of ChiS. While this conformation has not been observed among other SBP-HKs, a similar arrangement is predicted for maltose binding protein (MBP)-dependent activation of the Tar chemoreceptor[26].

For all structurally characterized SBP-HK systems, the orientation of the SBP with respect to the HK remains relatively unchanged upon ligand binding. This includes LuxP-LuxQ[9], TorT-TorS[10], XylFII-LytS[12], and HptA-HptS[11]. By contrast, we show that CBP-ChiS undergo an interface switch upon ligand-binding, where the orientation of CBP exhibits a large rotation relative to the ChiS dimer (**Movie S1**). Structural modeling suggests that similar interface switches also facilitate the activation of ChvE-VirA and BctC-BctE (**Movie S2-S3**). Thus, interface switching may represent a novel mechanism for allosteric activation of signal transduction in SBP-HKs.

Our results suggest that structural modeling can rapidly reveal the allosteric activation mechanism of protein complexes. In particular, we show that AF-Mv3 can model distinct conformations for a number of SBP-HKs. This was not observed when using earlier versions of AF-M (v1 or v2) or AF3, which all generated CBP-ChiS models in only one conformation. Furthermore, AF3 yielded low confidence models for ChvE-VirA, suggesting that AF-Mv3 may be better at modeling some protein complexes. AF3, however, was able to model CBP in its unliganded state (**Fig. S2**) and reveal which CBP-ChiS complex represents the ChiS-repressed conformation. This would not have been possible with AF-Mv3 alone. While AF-Mv3 revealed multiple conformations for CBP-ChiS, the CBPs in these models were identical and matched the crystal structure of liganded CBP. So, employing multiple structural modeling algorithms (*e.g.*, AF-M[18], AF3[19], Chai-1[27], RosettaFold2[28], and RosettaFold All Atom[29]) and comparing between models may help reveal allosteric conformations in other protein complexes. Furthermore, recent work highlights that clustering and/or subsampling of multiple sequence alignments (MSAs) can help elucidate protein conformational landscapes[30-33], which may help accelerate the discovery of additional allosteric activation mechanisms moving forward.

METHODS

Structural modeling with AlphaFold

The initial CBP-ChiS models (**Fig. 1**) were generated using the AF-Mv3 algorithm[18] in ColabFold[16] on Indiana University's BigRed200 and Quartz Supercomputers or using the AlphaFold3 algorithm[19] on the AlphaFold webserver. For AF-Mv3, multiple sequence alignments (MSAs) were generated by MMseqs2[34] and five relaxed models were generated from a single seed using 50 recycles with tolerance set to 0.2. The 5 top ranked models from 5 independent runs were used for subsequent analysis. To identify distinct protein conformations using AF-Mv3 (**Fig. S9** and **S10**) each query generated 1 unrelaxed model for 25 independent random seeds using 30 recycles with tolerance set to 1.0.

Models were analyzed using ChimeraX[35] (v1.9), and contacts were determined using the "contacts" and "h-bonds" tools. Parameters for "contacts" included a VDW overlap of ≥ -0.4 Å and intramodel interactions only (intermodel, intramolecule, intraresidue contacts were excluded). Parameters for "h-bonds" included a distance tolerance of 5 Å, angle tolerance of 20°, and salt-bridges only. For a complete list of predicted electrostatic contacts, see **Dataset S1**. Structural models are available at ModelArchive (Accession: ma-dal-cbpchis).

Bacterial strains and culture conditions

Bacterial strains were grown in LB Miller broth at 30°C rolling and on LB Miller agar. When necessary, the medium was supplemented with spectinomycin (200 µg/ml), kanamycin (50 µg/mL), trimethoprim at 10 (µg/mL), chloramphenicol (1 µg/mL), carbenicillin (100 µg/mL), erythromycin (10 µg/mL), sulfamethoxazole (100 µg/mL), tetracycline (0.5 µg/mL), and/or zeocin (100 µg/mL) as appropriate.

All strains were derived from the *Vibrio cholerae* El Tor strain E7946. Mutant constructs were created using splicing-by-overlap-extension (SOE) PCR exactly as previously described[36]. All chromosomally integrated ectopic expression constructs were generated exactly as previously described[37]. Mutant constructs were transformed into *V. cholerae* using natural transformation. Competence was artificially induced in cells via ectopic expression of TfoX, the master regulator of competence, using either an IPTG-inducible plasmid (pMMB67EH-*tfoX*) or via a chromosomally integrated arabinose-inducible construct (P$_{BAD}$-*tfoX*). The ectopic *tfoX*

expression construct was either cured (for pMMB67EH-*tfoX*) or deleted (for P$_{BAD}$-*tfoX*) from all strains prior to their use in experiments. All mutations were verified by PCR and/or sequencing.

For a detailed list of all strains, see **Table S1**. For a complete list of oligos used to generate mutant constructs specific to this study, see **Table S2**. The CBP$^{\Delta}$ construct used throughout the study consists of the CBP signal sequence translationally fused to mCherry (*i.e.,* P$_{chb}$-*cbp$^{1-40}$*-*mCherry)*.

Microscopy
Cells were imaged on an inverted Nikon Ti-2 microscope with a Plan Apo 60x objective lens, the appropriate filter cubes (FITC, YFP, mCherry, and/or CFP), a Hamamatsu ORCA Flash 4.0 camera, and Nikon NIS Elements imaging software. All image analysis was performed using the MicrobeJ[38] plugin (v5.10m) in Fiji[39] (v1.53k).

ChiS activity assays
Single colonies were inoculated into LB medium and incubated at 30°C for 18 hours rolling to generate overnight cultures. To assess chitin-independent ChiS activity by plate reader, cells were washed and resuspended in Instant Ocean (IO) medium (7 g/L, Aquarian Systems) to an OD$_{600}$ of ~5.0. Then, 200 μL of cell suspension was transferred to two wells (technical replicates) per strain in a flat-bottom 96-well plate and fluorescence was determined on a Biotek H1M plate reader. The fluorescence monochromator settings for excitation/emission were: 446/491 for mTFP1 and 580/610 for mCherry. Background fluorescence was determined by analyzing a non-fluorescent strain and was subtracted from all experimental samples. Technical replicate measurements were averaged. Then, CBP-mCherry fluorescence was normalized by mTFP1 fluorescence to determine the ChiS activity for each biological replicate.

To assess chitin-independent ChiS activity by microscopy, overnight cultures were washed and resuspended in IO to an OD$_{600}$ of ~1.0. Then, 2 μL of this cell suspension was placed under a 0.3% Gelzan IO pad. Cells were imaged and the average fluorescence signal per cell was determined using MicrobeJ. Background fluorescence was determined by analyzing a non-fluorescent strain and was subtracted from all experimental samples. For each cell, the CBP-mCherry fluorescence was normalized by mTFP1 fluorescence to control for intrinsic noise in gene expression. The geometric mean of 300 cells was reported for each biological replicate as the ChiS activity.

To assess chitin-dependent ChiS activity by microscopy, overnight cultures were subcultured into fresh LB and grown to an OD$_{600}$ of ~1.0. Cells were then washed and resuspended to an OD$_{600}$ of 1.0 in IO. Then, 100 μL of cell suspension was mixed with 750 μL of IO and 150 μL of insoluble chitin slurry (5% w/v insoluble chitin flakes in IO). Reaction tubes were shaken at 30°C for 48 hours to allow for chitin-induction. Then, reactions were vortexed to release the cells from chitin and the supernatant was transferred to a fresh Eppendorf tube to concentrate the cells in IO. Next, cells were placed under a 0.3% Gelzan IO pad and imaged and analyzed exactly as described above for chitin-independent ChiS activity assays.

PopZ-linked apical recruitment (POLAR) assays
Single colonies were inoculated into LB medium supplemented with 0.1 μM IPTG to induce P$_{tac}$-*chiS-msfGFP-H3H4* expression. Where indicated, samples were also supplemented with 0.05% arabinose to induce P$_{BAD}$-PopZ expression to induce polar relocalization of the H3H4-tagged bait. In all strains, the cell curvature determinant *crvA* was deleted because ectopic expression of ChiS induces strong cell curvature[40], which confounds the analysis. Importantly, the *chiS*-msfGFP-H3H4 construct is nonfunctional for *P$_{chb}$*-induction, thus, the *P$_{chb}$-cbp-mCherry*

construct is expressed at a basal and equivalent level in all strains. Cultures were incubated at 30°C for 18 hours rolling. Then, cells were subcultured the following day in fresh medium and grown for 1 hour at 30°C rolling. Cells were then washed and resuspended in IO to an OD_{600} of ~1.0. Then, 2µL of cell suspension was placed under a 0.3% Gelzan IO pad and imaged. Fluorescence localization within cells was determined using MicrobeJ. Heatmaps of fluorescence localization were generated by analyzing 300 cells per replicate. To quantify polar localization, the distance of fluorescence to the nearest cell pole was measured. Fluorescence ≤0.2 µm from the cell pole was conservatively defined as polarly localized.

Western Blotting

To ensure that CBP expression levels were equivalent between strains, we used a previously described ChiS construct[24] where the native DNA-binding domain was deleted (∆DBD) and replace by TetR. Strains were grown in LB medium supplemented with 0.04% arabinose, to induce P_{tac}-$chiS^{\Delta DBD}$-$tetR$ expression, and incubated at 30°C for 18 hours rolling. Strains were subcultured into fresh medium and grown at 30°C to an OD_{600} of ~4.0. Cells were then washed in ice cold IO.

For disulfide crosslinked samples, cells were resuspended to an OD_{600} of 100 in ice cold IO. Next, cells were chemically lysed by adding 10% FastBreak, lysozyme (0.5 mg/mL), and Benzonase (0.5 unit/µL) and incubating for 15 minutes at room temperature. To assess disulfide crosslinking, samples were mixed 1:1 with 2X non-reducing SDS-PAGE sample buffer (222 mM Tris HCl pH 6.8, 26 % glycerol, 3.9 % SDS, 0.022 % bromophenol blue). To reduce disulfde crosslinks prior to gel electrophoresis, samples were mixed 1:1 with 2X reducing SDS-PAGE sample buffer (222 mM Tris HCl pH 6.8, 26 % glycerol, 3.9 % SDS, 0.022 % bromophenol blue, 0.715 M beta-mercaptoethanol, 160 mM DTT).

For DSS crosslinking, cells were resuspended in ice cold IO, mixed with 5 mM DSS, and nutated at room temperature for 30 minutes. Samples were then quenched with 45mM Tris HCl pH 7.5 and nutated at room temperature for 15 minutes. Samples were then washed and resuspended to an OD_{600} of ~100 in ice cold IO. Next, cells were chemically lysed exactly as described above. Then, lysates were mixed 1:1 with 2X reducing SDS-PAGE sample buffer.

Next, 5µL of each sample was electrophoretically separated on a 7.5% SDS-PAGE gel. Proteins were then transferred to a polyvinylidene difluoride (PVDF) membrane. Membranes were blocked in TBST (20 mM Tris HCl pH 7.5, 50 mM NaCl, 0.05% Tween 20) containing 5% milk powder for 1 hour rocking at room temperature. Membranes were then incubated with 1:10,000 rabbit polyclonal α-mCherry primary antibody in TBST supplemented with 0.02% sodium azide and 2% milk powder rocking at room temperature overnight. Membranes were then washed with TBST before a 2-hour incubation with 1:10,000 α-rabbit horseradish peroxidase-conjugated secondary antibody. Membranes were then washed and incubated with Pierce ECL Western Blotting Substrate before chemiluminescence imaging on a ProteinSimple FluorChem R instrument.

Insoluble chitin growth assays

Single colonies were inoculated into LB medium and incubated at 30°C for 18 hours rolling. Then, cells were subcultured into fresh LB and grown to an OD_{600} of ~1.0. Cells were washed and resuspended to an OD_{600} of 1.0 in M9 minimal medium (1X M9 salts, 2 mM $MgSO_4$, 0.1 mM $CaCl_2$, 0.03 mM $FeSO_4$). Then, 10 µL of cell suspension was added into 1 mL reactions containing 850 µL of M9 minimal medium and 150 µL of chitin slurry. Reactions were shaken at 30°C for a total of 72 hours. Growth was assessed at 0, 24, 48, and 72 hours by quantitative dilution plating.

Chitin-induced natural transformation assays

Single colonies were inoculated into LB medium and incubated at 30°C for 18 hours rolling. Cells were then subcultured into fresh LB and grown to an OD_{600} of ~1.0. Cells were washed and resuspended to an OD_{600} of 1.0 in IO. Then, 100 µL of cell suspension was added into 1mL reactions containing 750 µL of IO and 150 µL of chitin slurry. Reaction tubes were incubated statically at 30°C for 18 hours. Then, 550 µL of the supernatant was removed without disturbing the settled chitin. Next, 100 ng of transforming DNA (ΔVC1807::Spec^R) was added. For each strain, control reactions were performed where no DNA was added. Reaction tubes were incubated statically at 30°C for 5 hours. Next, 500 µL of LB medium was added to each reaction, and outgrown by incubating at 37°C for 3 hours shaking. Reactions were then dilution plated on LB supplemented with spectinomycin to quantify transformants, and plain LB to quantify total viable counts. Transformation frequency is defined as the CFU/mL of transformants divided by the CFU/mL of total viable counts.

Statistical comparisons

All statistical comparisons were made in Graphpad Prism (v10) and the statistical tests used are indicated in each figure legend. See **Dataset S2** for a comprehensive list of descriptive statistics and statistical comparisons.

ACKNOWLEDGEMENTS

We thank Virginia Green and Catherine Klancher for constructive comments on the manuscript and helpful discussions. This work was supported by grant R35GM128674 from the National Institutes of Health to ABD.

REFERENCES

Bhate, M. P., Molnar, K. S., Goulian, M. & DeGrado, W. F. Signal transduction in histidine kinases: insights from new structures. *Structure* **23**, 981-994 (2015). https://doi.org/10.1016/j.str.2015.04.002

Wolanin, P. M., Thomason, P. A. & Stock, J. B. Histidine protein kinases: key signal transducers outside the animal kingdom. *Genome Biol* **3**, REVIEWS3013 (2002). https://doi.org/10.1186/gb-2002-3-10-reviews3013

Mascher, T., Helmann, J. D. & Unden, G. Stimulus perception in bacterial signal-transducing histidine kinases. *Microbiol Mol Biol Rev* **70**, 910-938 (2006). https://doi.org/10.1128/MMBR.00020-06

Matilla, M. A., Ortega, A. & Krell, T. The role of solute binding proteins in signal transduction. *Comput Struct Biotechnol J* **19**, 1786-1805 (2021). https://doi.org/10.1016/j.csbj.2021.03.029

He, F. *et al.* Molecular basis of ChvE function in sugar binding, sugar utilization, and virulence in Agrobacterium tumefaciens. *J Bacteriol* **191**, 5802-5813 (2009). https://doi.org/10.1128/JB.00451-09

Klancher, C. A., Yamamoto, S., Dalia, T. N. & Dalia, A. B. ChiS is a noncanonical DNA-binding hybrid sensor kinase that directly regulates the chitin utilization program in Vibrio cholerae. *Proc Natl Acad Sci U S A* **117**, 20180-20189 (2020). https://doi.org/10.1073/pnas.2001768117

Xu, C., Cao, Q. & Lan, L. Glucose-Binding of Periplasmic Protein GltB Activates GtrS-GltR Two-Component System in Pseudomonas aeruginosa. *Microorganisms* **9** (2021). https://doi.org/10.3390/microorganisms9020447

Antoine, R. *et al.* The periplasmic binding protein of a tripartite tricarboxylate transporter is involved in signal transduction. *J Mol Biol* **351**, 799-809 (2005). https://doi.org/10.1016/j.jmb.2005.05.071

Neiditch, M. B. *et al.* Ligand-induced asymmetry in histidine sensor kinase complex regulates quorum sensing. *Cell* **126**, 1095-1108 (2006). https://doi.org/10.1016/j.cell.2006.07.032

Moore, J. O. & Hendrickson, W. A. An asymmetry-to-symmetry switch in signal transmission by the histidine kinase receptor for TMAO. *Structure* **20**, 729-741 (2012). https://doi.org/10.1016/j.str.2012.02.021

Wang, M. *et al.* Interface switch mediates signal transmission in a two-component system. *Proc Natl Acad Sci U S A* **117**, 30433-30440 (2020). https://doi.org/10.1073/pnas.1912080117

Li, J. *et al.* Molecular mechanism of environmental d-xylose perception by a XylFII-LytS complex in bacteria. *Proc Natl Acad Sci U S A* **114**, 8235-8240 (2017). https://doi.org/10.1073/pnas.1620183114

Meibom, K. L. *et al.* The Vibrio cholerae chitin utilization program. *Proc Natl Acad Sci U S A* **101**, 2524-2529 (2004). https://doi.org/101/8/2524

Meibom, K. L., Blokesch, M., Dolganov, N. A., Wu, C. Y. & Schoolnik, G. K. Chitin induces natural competence in Vibrio cholerae. *Science* **310**, 1824-1827 (2005). https://doi.org/310/5755/1824

Li, X. & Roseman, S. The chitinolytic cascade in Vibrios is regulated by chitin oligosaccharides and a two-component chitin catabolic sensor/kinase. *Proc Natl Acad Sci U S A* **101**, 627-631 (2004). https://doi.org/10.1073/pnas.0307645100

Mirdita, M. *et al.* ColabFold: making protein folding accessible to all. *Nat Methods* **19**, 679-682 (2022). https://doi.org/10.1038/s41592-022-01488-1

Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583-589 (2021). https://doi.org/10.1038/s41586-021-03819-2

Evans, R. *et al.* Protein complex prediction with AlphaFold-Multimer. *bioRxiv*, 2021.2010.2004.463034 (2022). https://doi.org/10.1101/2021.10.04.463034

Abramson, J. *et al.* Accurate structure prediction of biomolecular interactions with AlphaFold 3. *Nature* **630**, 493-500 (2024). https://doi.org/10.1038/s41586-024-07487-w

Lim, H. C. & Bernhardt, T. G. A PopZ-linked apical recruitment assay for studying protein-protein interactions in the bacterial cell envelope. *Mol Microbiol* **112**, 1757-1768 (2019). https://doi.org/10.1111/mmi.14391

Kitaoku, Y. *et al.* A structural model for (GlcNAc)2 translocation via a periplasmic chitooligosaccharide-binding protein from marine Vibrio bacteria. *J Biol Chem* **297**, 101071 (2021). https://doi.org/10.1016/j.jbc.2021.101071

Fukamizo, T., Kitaoku, Y. & Suginta, W. Periplasmic solute-binding proteins: Structure classification and chitooligosaccharide recognition. *Int J Biol Macromol* **128**, 985-993 (2019). https://doi.org/10.1016/j.ijbiomac.2019.02.064

Suginta, W. *et al.* Structure and function of a novel periplasmic chitooligosaccharide-binding protein from marine Vibrio bacteria. *J Biol Chem* **293**, 5150-5159 (2018). https://doi.org/10.1074/jbc.RA117.001012

Hullinger, A. C., Green, V. E., Klancher, C. A., Dalia, T. N. & Dalia, A. B. Two transmembrane transcriptional regulators coordinate to activate chitin-induced natural transformation in Vibrio cholerae. *PLoS Genet* **21**, e1011606 (2025). https://doi.org/10.1371/journal.pgen.1011606

Cangelosi, G. A., Ankenbauer, R. G. & Nester, E. W. Sugars induce the Agrobacterium virulence genes through a periplasmic binding protein and a transmembrane signal protein. *Proc Natl Acad Sci U S A* **87**, 6708-6712 (1990). https://doi.org/10.1073/pnas.87.17.6708

Zhang, Y. *et al.* Model of maltose-binding protein/chemoreceptor complex supports intrasubunit signaling mechanism. *Proc Natl Acad Sci U S A* **96**, 939-944 (1999). https://doi.org/10.1073/pnas.96.3.939

Discovery, C. *et al.* Chai-1: Decoding the molecular interactions of life. *bioRxiv*, 2024.2010.2010.615955 (2024). https://doi.org/10.1101/2024.10.10.615955

Baek, M. *et al.* Efficient and accurate prediction of protein structure using RoseTTAFold2. *bioRxiv* (2023). https://doi.org/10.1101/2023.05.24.542179

Krishna, R. *et al.* Generalized biomolecular modeling and design with RoseTTAFold All-Atom. *Science* **384**, eadl2528 (2024). https://doi.org/10.1126/science.adl2528

Chakravarty, D. *et al.* AlphaFold predictions of fold-switched conformations are driven by structure memorization. *Nat Commun* **15**, 7296 (2024). https://doi.org/10.1038/s41467-024-51801-z

Schafer, J. W. *et al.* Sequence clustering confounds AlphaFold2. *Nature* **638**, E8-E12 (2025). https://doi.org/10.1038/s41586-024-08267-2

Wayment-Steele, H. K. *et al.* Predicting multiple conformations via sequence clustering and AlphaFold2. *Nature* **625**, 832-839 (2024). https://doi.org/10.1038/s41586-023-06832-9

Rustamov, K. R. & Baev, A. Y. MSA clustering enhances AF-Multimer's ability to predict conformational landscapes of protein-protein interactions. *Bioinform Adv* **5**, vbae197 (2025). https://doi.org/10.1093/bioadv/vbae197

Steinegger, M. & Soding, J. MMseqs2 enables sensitive protein sequence searching for the analysis of massive data sets. *Nat Biotechnol* **35**, 1026-1028 (2017). https://doi.org/10.1038/nbt.3988

Meng, E. C. *et al.* UCSF ChimeraX: Tools for structure building and analysis. *Protein Sci* **32**, e4792 (2023). https://doi.org/10.1002/pro.4792

Dalia, A. B. Natural Cotransformation and Multiplex Genome Editing by Natural Transformation (MuGENT) of Vibrio cholerae. *Methods Mol Biol* **1839**, 53-64 (2018). https://doi.org/10.1007/978-1-4939-8685-9_6

Dalia, T. N., Chlebek, J. L. & Dalia, A. B. A modular chromosomally integrated toolkit for ectopic gene expression in Vibrio cholerae. *Sci Rep* **10**, 15398 (2020). https://doi.org/10.1038/s41598-020-72387-8

Ducret, A., Quardokus, E. M. & Brun, Y. V. MicrobeJ, a tool for high throughput bacterial cell detection and quantitative analysis. *Nat Microbiol* **1**, 16077 (2016). https://doi.org/10.1038/nmicrobiol.2016.77

Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**, 676-682 (2012). https://doi.org/10.1038/nmeth.2019

Green, V. E., Klancher, C. A., Yamamoto, S. & Dalia, A. B. The molecular mechanism for carbon catabolite repression of the chitin response in Vibrio cholerae. *PLoS Genet* **19**, e1010767 (2023). https://doi.org/10.1371/journal.pgen.1010767

Fig. 1. *Structural modeling of CBP-ChiS reveals distinct conformations.* (**A**) Schematic representation of CBP dual allosteric regulation of ChiS activity. Unliganded CBP represses ChiS activity, while liganded CBP stimulates ChiS activity. (**B**) Top ranked models of CBP-ChiS^{AF3} and CBP-ChiS^{AF-M}. CBP is colored blue, while the two copies of the ChiS periplasmic domain (residues 1-427) are colored yellow & orange.

Fig. 2. *An electrostatic interaction specific to CBP-ChiS^{AF3} (unliganded) is necessary for allosteric repression of ChiS in the absence of chitin.* (**A**) Transcriptional reporter assays to measure ChiS activity in strains harboring the indicated mutations. All strains harbored P$_{chb}$-*cbp-mCherry* and P$_{const2}$-*mTFP1* reporter constructs. The normalized mCherry fluorescence (see **Methods** for details) is reported as ChiS activity. Cells were grown in rich medium (*i.e.*, in the absence of chitin) and fluorescence was determined by microscopy. Data are from three independent biological replicates (*n* = 300 cells analyzed per replicate) and shown as the mean ± SD. Statistical comparisons were made by one-way ANOVA with Tukey's multiple comparison test on the log-transformed data. Statistical identifiers directly above bars represent comparisons to the parent. (**B**) Positions of CBP^{E401} (cyan) and ChiS^{R141,K145} (magenta) are highlighted in the CBP-ChiS^{AF3} (unliganded) model to show the electrostatic interaction (yellow dashed line) formed. CBP is shown in light grey, while ChiS is shown in dark grey. (**C**) CBP-ChiS interactions were assessed by POLAR assays. Cells contained *cbp-mCherry* and *chiS-msfGFP-H3H4* constructs with the indicated mutations as well as an inducible PopZ construct (P$_{BAD}$-*popZ*). CBP^{Δ} = the CBP signal sequence translationally fused to mCherry (P$_{chb}$-*cbp$^{1-40}$-mCherry)*. Fluorescence microscopy was performed on cells without (-) and with (+) PopZ

induction. Representative images are shown. Scale bars, 2 μm. Heat maps represent the localization of mCherry and GFP fluorescence in 300 cells. The proportion of CBP-mCherry (cyan bars) and ChiS-msfGFP-H3H4 (purple bars) fluorescence within the polar region (≤0.2 μm from the cell pole) was quantified. For each biological replicate ($n = 3$), fluorescence localization was determined by analyzing 300 individual cells. Statistical comparisons were made by Student's t-test on the log-transformed data. ns = not significant, * = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.001$.

Fig. 3. *An electrostatic interaction specific to CBP-ChiS^{AF-M} (liganded) is necessary for allosteric activation of ChiS in the presence of chitin.* (**A**) The positions of CBP^{D251} (cyan) and ChiS^{R146} (magenta) are highlighted in the CBP-ChiS^{AF3} (unliganded) and CBP-ChiS^{AF-M} (liganded) models to show that these residues only form an electrostatic interaction (yellow dashed line) in the latter. CBP is shown in light grey, while ChiS is shown in dark grey. (**B**) Transcriptional reporter assays to assess ChiS activity in strains harboring the indicated mutations. All strains harbored P$_{chb}$-*cbp-mCherry* and P$_{const2}$-*mTFP1* reporter constructs. CBP* = CBP$^{W389A\ W539A}$, a mutant allele that cannot bind to chitin. Cells were grown in either rich medium (No chitin; dark grey bars) or on chitin flakes (chitin; light grey bars) and fluorescence was determined by microscopy. Data are from three independent biological replicates (*n* = 300 cells analyzed per replicate) and shown as the mean ± SD. Statistical comparisons were made by one-way ANOVA with Tukey's multiple comparison test on the log-transformed data. ns = not significant, *** = *p* < 0.001. Statistical identifiers directly above bars represent comparisons to the parent from the same experimental condition.

Fig. 4. *Residue partner-switching facilitates CBP-dependent allosteric dual-regulation of ChiS.* CBP[E401] undergoes partner switching to form distinct electrostatic interactions when transitioning from the unliganded (repressed) to liganded (active) conformations. (**A**) Models highlight that CBP[E401] (cyan) interacts with ChiS[R141,K145] (magenta) in CBP-ChiS[AF3] (unliganded) and ChiS[R89] (orange) in CBP-ChiS[AF-M] (liganded). CBP is shown in light grey, while ChiS is shown in dark grey. (**B**) Transcriptional reporter assays to assess ChiS activity in strains harboring the indicated mutations. All strains harbored P*chb*-*cbp-mCherry* and P*const2*-*mTFP1* reporter constructs. CBP* = CBP[W389A W539A], a mutant allele that cannot bind to chitin. Cells were grown in either rich medium (No chitin; dark grey bars) or on chitin flakes (chitin; light grey bars) and fluorescence was determined by microscopy. Data are from three independent biological replicates (*n* = 300 cells analyzed per replicate) and shown as the mean ± SD. Statistical comparisons were made by one-way ANOVA with Tukey's multiple comparison test on the log-transformed data. ns = not significant, *** = $p < 0.001$. Statistical identifiers directly above bars represent comparisons to the parent from the same experimental condition.

Fig. 5. *Disulfide locking CBP-ChiS into the liganded (active) conformation induces ChiS activity in the absence of chitin.* (**A**) Models highlight the positions of CBP^{D310} (cyan), ChiS^{Q243} (magenta), and ChiS^{Y245} (orange) in the CBP-ChiS^{AF3} (unliganded) and CBP-ChiS^{AF-M} (liganded) models. The Inset shows that CBP^{D310} is predicted to interact with ChiS^{Q243} (yellow dashed line), but not ChiS^{Y245C}. CBP is shown in light grey, while ChiS is shown in dark grey. (**B**) Non-reducing Western blots to assess CBP-ChiS disulfide crosslinking. All strains harbor P_{tac}-*chiS$^{\Delta DBD}$-tetR* and P_{chb}-*cbp-mCherry* constructs with the indicated mutations. (**C**) Transcriptional reporter assays to assess ChiS activity in strains harboring the indicated mutations. All strains harbored P_{chb}-*cbp-mCherry* and P_{const2}-*mTFP1* reporter constructs. Cells were grown in rich medium (*i.e.*, in the absence of chitin) and fluorescence was determined by microscopy. The data for the CBP$^{\Delta}$ ChiS^{WT}, CBP^{WT} ChiS^{T473A}, and CBP$^{\Delta}$ ChiS^{T473A} controls were duplicated from **Fig. 2A** and **4B**, and included here for ease of comparison. Data are from three independent biological replicates (*n* = 300 cells analyzed per replicate) and shown as the mean ± SD. Statistical comparisons were made by one-way ANOVA with Tukey's multiple comparison test on the log-transformed data. ns = not significant, *** = $p < 0.001$. Statistical identifiers directly above bars represent comparisons to the first sample (CBP$^{\Delta}$ ChiS^{WT}).

Fig. 6. *Transitioning from the unliganded (repressed) to liganded (active) conformation is critical for ChiS-induced behaviors.* (**A**) The indicated strains were grown in M9 minimal media supplemented with insoluble chitin as the sole carbon source. Growth was assessed by quantitative culture at the indicated timepoints. (**B**) Chitin-induced natural transformation assays. CBP* = $CBP^{W389A\ W539A}$, a mutant allele that cannot bind to chitin. All data are from at least three independent biological replicates and shown as the mean ± SD. Statistical comparisons were made by one-way ANOVA with Tukey's multiple comparison test on the log-transformed data. ns = not significant, *** = $p < 0.001$. Statistical identifiers directly above bars represent comparisons to the parent from the same experimental condition.

Fig. S1. *Different AlphaFold algorithms predict two distinct conformations for CBP-ChiS.* (**A**) Top ranked models of CBP-ChiS[AF-M] and CBP-ChiS[AF3]. CBP's orientation is rotated relative to ChiS dimer. (**B**) Overlay of CBP-ChiS[AF-M] and CBP-ChiS[AF3] models to highlight the distinct orientation of CBP within each model. The CBP-ChiS[AF-M] model is colored in cyan, while the CBP-ChiS[AF3] model is colored in magenta. (**C**) PAE scores for the CBP-ChiS[AF-M] (left) and CBP-ChiS[AF3] (right) models. (**D**) Quantitative metrics of the AlphaFold models. Model confidence score is defined as: (0.8*pTM) + (0.2*ipTM).

Model	pTM	ipTM	Model Confidence Score
CBP-ChiS[AF-M]	0.73	0.68	0.720
CBP-ChiS[AF3]	0.69	0.62	0.676

Fig. S2. *Alignment of CBP from CBP-ChiS^{AF3} and CBP-ChiS^{AF-M} to solved crystal structures reveals the liganded state of each AlphaFold model.* CBP extracted from either the CBP-ChiS^{AF-M} or CBP-ChiS^{AF3} model (blue) were aligned to either unliganded CBP (4gf8, salmon) or liganded CBP (4gfr, magenta) as indicated. Untrimmed RMSD scores are given for each alignment.

A

CBP Residue	ChiS Residue	Unliganded Distance (Å) ± SD	Liganded Distance (Å) ± SD
E401	R141	2.745±0.092	26.503±0.210
E401	K145	3.597±0.414	20.869±0.209
D231	R99	12.941±1.408	3.482±0.261
D231	R139	14.771±1.555	2.692±0.013
D251	R146	19.226±0.444	2.725±0.011
R254	E81	25.215±0.956	2.976±0.314
E401	E89	27.296±0.241	2.746±0.045
E529	R99	19.062±0.274	2.765±0.081
E530	R99	22.108±0.932	2.662±0.021

Fig. S3. *Defining putative electrostatic interactions unique to each Alpahfold model.* The top models from five independent CBP-ChiS[AF-M] and CBP-ChiS[AF3] runs were analyzed to define putative electrostatic interactions unique to each model. (**A**) The average of the smallest distance between side chain atoms of the indicated residue pairs are shown ± SD. Residue pairs unique to (**B**) CBP-ChiS[AF3] and (**C**) CBP-ChiS[AF-M] residue pairs are mapped onto each AlphaFold model. CBP is colored light grey (indicated residues in cyan), while ChiS is colored dark grey (indicated residues in magenta).

Fig. S4. *Mutating residues specific to the CBP-ChiS^{AF-M} (liganded) model do not affect CBP-dependent repression of ChiS in the absence of chitin.* Transcriptional reporter assays to assess ChiS activity in strains harboring the indicated mutations that specifically disrupt the CBP-ChiS interface in the CBP-ChiS^{AF-M} (liganded) model (see **Fig. S3C** for details). All strains harbored P$_{chb}$-*cbp-mCherry* and P$_{const2}$-*mTFP1* reporter constructs. CBP$^\Delta$ = the CBP signal sequence translationally fused to mCherry (P$_{chb}$-*cbp$^{1-40}$-mCherry).* Cells were grown in rich medium (*i.e.,* in the absence of chitin) and fluorescence was determined by plate reader. Data are from three independent biological replicates and shown as the mean ± SD.

Fig. S5. *The phosphatase-deficient ChiS^{T473A} allele requires liganded (chitin-bound) CBP for allosteric activation.* Transcriptional reporter assays to assess ChiS activity in strains harboring the indicated mutations. All strains harbored P_{chb}-*cbp-mCherry* and P_{const2}-*mTFP1* reporter constructs. CBP* = CBP^{W389A W539A}, a mutant allele that cannot bind to chitin. CBP^Δ = the CBP signal sequence translationally fused to mCherry (P_{chb}-*cbp^{1-40}-mCherry*). Cells were grown in either rich medium (No chitin; dark grey bars) or on chitin flakes (chitin; light grey bars) and fluorescence was determined by microscopy. Data are from three independent biological replicates (*n* = 300 cells analyzed per replicate) and shown as mean ± SD. Statistical comparisons were made by one-way ANOVA with Tukey's multiple comparison test on the log-transformed data. ns = not significant, *** = $p < 0.001$.

Fig. S6. *Testing CBP-ChiS^{AF-M} interface mutants for a gain-of-function phenotype.* Transcriptional reporter assays to assess ChiS activity in strains harboring the indicated mutations. All strains harbored CBP$^{\Delta}$ (P$_{chb}$-*cbp$^{1-40}$*-*mCherry*) and P$_{const2}$-*mTFP1* reporter constructs. Cells were grown in rich medium (*i.e.*, in the absence of chitin) and fluorescence was determined by plate reader. Data are from three independent biological replicates and shown as mean ± SD. The ChiS^{T473A,E81R} allele exhibited a gain-of-function phenotype, thus, this interaction (**Fig. S3C**) was excluded from further analysis.

Fig. S7. *Multiple electrostatic interactions specific to CBP-ChiS^{AF-M} (liganded) are necessary for chitin-induced activation.* (**A**) The positions of CBP^{D231}, CBP$^{E529\,E530}$, CBP^{E401}, ChiS^{R99,R139}, and ChiS^{R89} are highlighted in the CBP-ChiS^{AF-M} (liganded) model to show the electrostatic interactions formed. CBP is shown in light grey, while ChiS is shown in dark grey. Insets highlight the position of side chains with CBP residues colored cyan, ChiS residues colored magenta, and their associated electrostatic interactions denoted by a yellow dashed line. (**B**) Transcriptional reporter assays to assess ChiS activity in strains harboring the indicated mutations. All strains harbored P$_{chb}$-*cbp-mCherry* and P$_{const2}$-*mTFP1* reporter constructs. CBP* = CBP$^{W389A\,W539A}$, a mutant allele that cannot bind to chitin. CBP$^{\Delta}$ = the CBP signal sequence translationally fused to mCherry (P$_{chb}$-*cbp$^{1-40}$-mCherry)*. Cells were grown in either rich medium (No chitin; dark grey bars) or on chitin flakes (chitin; light grey bars) and fluorescence was determined by microscopy. The "No Chitin" data for CBP^{WT} ChiS^{T473A}, CBP$^{\Delta}$ ChiS^{T473A} and CBP$^{W389A\,W539A}$ ChiS^{WT} were duplicated from **Fig. 4B** and included here for ease of comparison. Data are from three independent biological replicates (*n* = 300 cells analyzed per replicate) and shown as the mean ± SD. Statistical comparisons were made by one-way ANOVA with Tukey's multiple comparison test on the log-transformed data. ns = not significant, *** = *p* < 0.001. Statistical identifiers directly above bars represent comparisons to the parent from the same experimental condition.

Fig. S8. *Cysteine knock-ins to CBP-ChiS do not inhibit CBP-ChiS interactions and form a sequence-specific disulfide lock that can be reversed under reducing conditions.* All strains harbored P_{tac}-*chiS*$^{\Delta DBD}$-*tetR* and P_{chb}-*cbp-mCherry* constructs with the indicated mutations and were grown aerobically. (**A**) NHS-ester crosslinking with DSS was used to assess CBP-ChiS interactions by Western Blot. Data are representative of two independent biological replicates. The shifted band absent in the *cbp*^{WT} $\Delta chiS$ sample defines the band representing the CBP-ChiS complex. (**B**) Non-reducing western blots to assess CBP-ChiS disulfide crosslinking. All strains harbor P_{tac}-*chiS*$^{\Delta DBD}$-*tetR* and P_{chb}-*cbp-mCherry* constructs with the indicated mutations. Lysates were either run directly (-DTT; oxidizing conditions) or treated with DTT to reduced disulfide bonds prior to gel electrophoresis. Data are representative of two independent biological replicates.

Fig. S9. AF-M *predicts distinct conformations for other SBP-HK signaling complexes.* AF-M reveals two distinct conformations for each SBP-HK pair. Models are aligned using one HK protein chain as the reference point. PAE confidence plots for the top ranked model in each conformation are shown below. (**A**) The top-ranked conformation for CBP-ChiS, ChvE-VirA, and BctC-BctE. Out of 25 independent seeds, this conformation was predicted in 11, 9, and 8 models, respectively. The models that shared this conformation have been aligned to show model-to-model agreement. SBPs are shown in teal and HKs shown in aquamarine. (**B**) The 2nd-ranked conformation for CBP-ChiS, ChvE-VirA, and BctC-BctE. Out of 25 independent seeds, this conformation was predicted in 9, 3, and 4 models, respectively. The models that shared this conformation have been aligned to show model-to-model agreement. SBPs shown in purple and HKs shown in violet. (**C**) Alignment of the top-ranked model in each conformation highlights the major differences between the orientation of the SBP in each conformation. Conformation 1 shown in teal (SBP) and aquamarine (HK). Conformation 2 shown in dark magenta (SBP) and violet (HK).

Fig. S10. *ChvE residues critical for VirA-dependent activity are predicted at the ChvE-VirA interface.* (**A**) Cartoon models (alpha-helices shown as cylinders) highlight the large differences in the ChvE-VirA interaction interface between the two predicted conformations of this complex (see **Fig. S9**). For conformation 1, ChvE is colored teal and VirA is colored aquamarine. For conformation 2, ChvE is colored purple and VirA is colored violet. (**B**) Cartoon models highlight the residues previously found to influence ChvE-VirA activity[5] in both conformations. Residues that, when mutated, decrease VirA activity in the presence of its glucose ligand are colored red. Residues that, when mutated, increase VirA activity in the absence of its glucose ligand are colored yellow. (**C**) The distance between the α-carbons of the indicated ChvE residues (highlighted in **B**) and the closest α-carbon in VirA are plotted for each conformation. (**D**) Predicted contacts for a subset of the ChvE residues shown in **B** highlights residues that undergo partner switching (VirA partner-switch residues are highlighted in grey).

Movie S1. A morph between the CBP-ChiS^{AF3} (unliganded) and CBP-ChiS^{AF-M} (liganded) conformations that highlights the large rotational shift of CBP relative to the ChiS dimer. The ChiS dimer is colored yellow and orange, while CBP is colored blue.

Movie S2. A morph between the two predicted conformations of ChvE-VirA that highlights the large rotational shift of ChvE relative to VirA. VirA is colored light gray, while ChvE is colored blue. Residues that, when mutated, decrease VirA activity in the presence of its glucose ligand are colored red. Residues that, when mutated, increase VirA activity in the absence of its glucose ligand are colored yellow.

Movie S3. A morph between the two predicted conformations of BctC-BctE that highlights the large rotational shift of BctC relative to BctE. BctE is colored light gray, while BctC is colored blue.

Table S1. *Strains used in this study*.

Strain*	Genotype	Figure(s)	Use in manuscript
SAD030	*V. cholerae* WT E7946 Sm[R]		Parent for all strains generated in this study
HSA0342 / SAD3734	Δ*lacZ*::Tet[R]- P$_{const2}$-*mTFP1*, igVCA0265-VCA0266::Cm[R]-P$_{chb}$-*cbp-mCherry*, Δ*chiS/cbp*::Erm[R], ΔVC1807::Zeo[R], igVCA0587-VCA0588::Kan[R]-P$_{chb}$-*gfp*	Fig. 2A, S8A	Δ*chiS cbp^{WT}* for ChiS activity and DSS crosslinking assays
HSA0343 / SAD3735	Δ*lacZ*::Tet[R]- P$_{const2}$-*mTFP1*, igVCA0265-VCA0266::Cm[R]-P$_{chb}$-*cbp-mCherry*, ΔVCA0692::Tm[R]-P$_{chiS}$-*chiS*, Δ*chiS/cbp*::Erm[R], ΔVC1807::Zeo[R], igVCA0587-VCA0588::Kan[R]-P$_{chb}$-*gfp*	Fig. 2A, S4, S5	*chiS^{WT} cbp^{WT}* for ChiS activity assays
HSA0344 / SAD3736	Δ*lacZ*::Tet[R]- P$_{const2}$-*mTFP1*, igVCA0265-VCA0266::Cm[R]-P$_{chb}$-*cbp^{E401R}-mCherry*, ΔVCA0692::Tm[R]-P$_{chiS}$-*chiS*, Δ*chiS/cbp*::Erm[R], ΔVC1807::Zeo[R], igVCA0587-VCA0588::Kan[R]-P$_{chb}$-*gfp*	Fig. 2A	*chiS^{WT} cbp^{E401R}* for ChiS activity assays
HSA0345 / SAD3737	Δ*lacZ*::Tet[R]- P$_{const2}$-*mTFP1*, igVCA0265-VCA0266::Cm[R]-P$_{chb}$-*cbp-mCherry*, ΔVCA0692::Tm[R]-P$_{chiS}$-*chiS$^{R141E\ K145E}$*, Δ*chiS/cbp*::Erm[R], ΔVC1807::Zeo[R], igVCA0587-VCA0588::Kan[R]-P$_{chb}$-*gfp*	Fig. 2A	*chiS^{R141E,K145E} cbp^{WT}* for ChiS activity assays
HSA0346 / SAD3738	Δ*lacZ*::Tet[R]- P$_{const2}$-*mTFP1*, igVCA0265-VCA0266::Cm[R]-P$_{chb}$-*cbp^{E401R}-mCherry*, ΔVCA0692::Tm[R]-P$_{chiS}$-*chiS$^{R141\ K145E}$*, Δ*chiS/cbp*::Erm[R], ΔVC1807::Zeo[R], igVCA0587-VCA0588::Kan[R]-P$_{chb}$-*gfp*	Fig. 2A	*chiS^{R141E,K145E} cbp^{E401R}* for ChiS activity assays
HSA0359 / SAD3739	Δ*lacZ*::Tet[R]- P$_{const2}$-*mTFP1*, igVCA0265-VCA0266::Cm[R]-P$_{chb}$-*cbp$^{[1-40]}$-mCherry*, ΔVCA0692::Tm[R]-P$_{chiS}$-*chiS*, Δ*chiS/cbp*::Erm[R], ΔVC1807::Zeo[R], igVCA0587-VCA0588::Kan[R]-P$_{chb}$-*gfp*	Fig. 2A, 5C S4, S5, S6	*chiS^{WT} cbp^{Δ}* for ChiS activity assays
HSA0928 / SAD3740	igVCA0265-VCA0266::Cm[R]-P$_{chb}$-*cbp$^{[1-40]}$-mCherry*, Δ*lacZ*::Tet[R]- P$_{const2}$-*mTFP1*, Δchis/cbp::Erm[R], ΔVC1807::Zeo[R], igVCA0587-VCA0588::Kan[R]-P$_{chb}$-*gfp*	Fig. 2A	Δ*chiS cbp^{Δ}* for ChiS activity assays
HSA0431 / SAD3741	Δ*lacZ*::Spec[R]-P$_{BAD}$-*popZ*, Δ*chiS/cbp*::Erm[R], Δ*crvA*::Zeo[R], igVCA0265-VCA0266::Cm[R]-P$_{chb}$-*cbp-mCherry*, ΔVCA0692::Tm[R]-P$_{tac}$-*chiS-msfGFP-H3H4*	Fig. 2C	*chiS^{WT} cbp^{WT}* for POLAR assays
HSA0432 / SAD3742	Δ*lacZ*::Spec[R]-P$_{BAD}$-*popZ*, Δ*chiS/cbp*::Erm[R], Δ*crvA*::Zeo[R], igVCA0265-VCA0266::Cm[R]-P$_{chb}$-*cbp$^{[1-40]}$-mCherry*, ΔVCA0692::Tm[R]-P$_{tac}$-*chiS-msfGFP-H3H4*	Fig. 2C	*chiS^{WT} cbp^{Δ}* for POLAR assays

Strain	Genotype	Figure	Description
HSA0491 / SAD3743	$\Delta lacZ$::Spec^R-P$_{BAD}$-$popZ$, $\Delta chiS/cbp$::Erm^R, $\Delta crvA$::Zeo^R, igVCA0265-VCA0266::Cm^R-P$_{chb}$-cbp^{E401R}-$mCherry$, ΔVCA0692::Tm^R-P$_{tac}$-$chiS$-$msfGFP$-$H3H4$	Fig. 2C	$chiS^{WT}$ cbp^{E401R} for POLAR assays
HSA0492 / SAD3744	$\Delta lacZ$::Spec^R-P$_{BAD}$-$popZ$, $\Delta chiS/cbp$::Erm^R, $\Delta crvA$:: Zeo^R, ΔVCA0692::Tm^R-P$_{tac}$-$chiS^{R141E\ K145E}$-$msfGFP$-$H3H4$, igVCA0265-VCA0266::Cm^R-P$_{chb}$-cbp-$mCherry$	Fig. 2C	$chiS^{R141E,K145E}$ cbp^{WT} for POLAR assays
HSA0493 / SAD3745	$\Delta lacZ$::Spec^R-P$_{BAD}$-$popZ$, $\Delta chiS/cbp$::Erm^R, $\Delta crvA$:: Zeo^R, ΔVCA0692::Tm^R-P$_{tac}$-$chiS^{R141E\ K145E}$-$msfGFP$-$H3H4$, igVCA0265-VCA0266::Cm^R-P$_{chb}$-cbp^{E401R}-$mCherry$	Fig. 2C	$chiS^{R141E,K145E}$ cbp^{E401R} for POLAR assays
HSA0356 / SAD3746	$\Delta lacZ$::Tet^R- P$_{const2}$-$mTFP1$, igVCA0265-VCA0266::Cm^R-P$_{chb}$-$cbp^{W389A\ W539A}$-$mCherry$, ΔVCA0692::Tm^R-P$_{chiS}$-$chiS^{T473A}$, $\Delta chiS/cbp$::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-gfp	Fig. 3B, 4B, 6, S5, S7B	$chiS^{T473A}$ $cbp*$ for ChiS activity, insoluble chitin growth, and chitin-induced natural transformation assays
HSA0360 / SAD3747	$\Delta lacZ$::Tet^R- P$_{const2}$-$mTFP1$, igVCA0265-VCA0266::Cm^R-P$_{chb}$-$cbp^{[1-40]}$-$mCherry$, ΔVCA0692::Tm^R-P$_{chiS}$-$chiS^{T473A}$, $\Delta chiS/cbp$::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-gfp	Fig. 3B, 4B, 5C, 6, S5, S6, S7B	$chiS^{T473A}$ cbp^{Δ} for ChiS activity, insoluble chitin growth, and chitin-induced natural transformation assays
HSA0362 / SAD3748	$\Delta lacZ$::Tet^R- P$_{const2}$-$mTFP1$, igVCA0265-VCA0266::Cm^R-P$_{chb}$-cbp-$mCherry$, ΔVCA0692::Tm^R-P$_{chiS}$-$chiS^{T473A}$, $\Delta chiS/cbp$::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-gfp	Fig. 3B, 4B, 5C, 6, S5, S7B	$chiS^{T473A}$ cbp^{WT} for ChiS activity, insoluble chitin growth, and chitin-induced natural transformation assays
HSA0369 / SAD3749	$\Delta lacZ$::Tet^R- P$_{const2}$-$mTFP1$, igVCA0265-VCA0266::Cm^R-P$_{chb}$-cbp^{D251R}-$mCherry$, ΔVCA0692::Tm^R-P$_{chiS}$-$chiS^{T473A}$, $\Delta chiS/cbp$::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-gfp	Fig. 3B, 6	$chiS^{T473A}$ cbp^{D251R} for ChiS activity, insoluble chitin growth, and chitin-induced natural transformation assays
HAS0763 / SAD3750	igVCA0265-VCA0266::Cm^R-P$_{chb}$-cbp-$mCherry$, ΔVCA0692::Tm^R-P$_{chiS}$-$chiS^{R146D\ T473A}$, $\Delta lacZ$::Tet^R- P$_{const2}$-$mTFP1$, $\Delta chiS/cbp$::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-gfp	Fig. 3B, 6	$chiS^{T473A,R146D}$ cbp^{WT} for ChiS activity, insoluble chitin growth, and chitin-induced natural transformation assays
HAS0764 / SAD3751	igVCA0265-VCA0266::Cm^R-P$_{chb}$-cbp^{D251R}-$mCherry$, ΔVCA0692::Tm^R-P$_{chiS}$-$chiS^{R146D\ T473A}$, $\Delta lacZ$::Tet^R- P$_{const2}$-$mTFP1$, $\Delta chiS/cbp$::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-gfp	Fig. 3B, 6	$chiS^{T473A,R146D}$ cbp^{D251R} for ChiS activity, insoluble chitin growth, and chitin-induced natural transformation assays
HSA0589 / SAD3752	igVCA0265-VCA0266::Cm^R-P$_{chb}$-cbp^{E401R}-$mCherry$, $\Delta lacZ$::Tet^R- P$_{const2}$-$mTFP1$, ΔVCA0692::Tm^R-P$_{chiS}$-$chiS^{T473A}$, $\Delta chiS/cbp$::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-gfp	Fig. 4B, 6, S7B	$chiS^{T473A}$ cbp^{E401R} for ChiS activity, insoluble chitin growth, and chitin-induced natural transformation assays

HSA0662 / SAD3753	igVCA0265-VCA0266::Cm^R-P$_{chb}$-*cbp-mCherry*, ΔVCA0692::Tm^R-P$_{chiS}$-*chiS^{R141E}$^{K145E\ T473A}$*, Δ*lacZ*::Tet^R- P$_{const2}$-*mTFP1*, Δ*chiS/cbp*::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-*gfp*	Fig. 4B, 6	*chiS^{T473A,R141E,K145E} cbp^{WT}* for ChiS activity, insoluble chitin growth, and chitin-induced natural transformation assays
HSA0696 / SAD3754	igVCA0265-VCA0266::Cm^R-P$_{chb}$-*cbp^{E401R}-mCherry*, ΔVCA0692::Tm^R-P$_{chiS}$-*chiS^{R141E,K145E,T473A}*, Δ*lacZ*::Tet^R- P$_{const2}$-*mTFP1*, Δ*chiS/cbp*::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-*gfp*	Fig. 4B, 6	*chiS^{T473A,R141E,K145E} cbp^{E401R}* for ChiS activity, insoluble chitin growth, and chitin-induced natural transformation assays
HSA0721 / SAD3755	VCA0692:: Tm^R-P$_{BAD}$-*chiS^{ΔDBD}-L5-tetR*, ΔVC1807::Spec^R, Δ*lacZ*::Tet^R- P$_{const2}$-*mTFP1*, igVCA0265-VCA0266::Cm^R-P$_{chb}$-*cbp-mCherry*, Δ*chis/cbp*::Erm^R	Fig. 5B, S8A-B	*chiS^{WT} cbp^{WT}* for disulfide and DSS crosslinking
HSA0850 / SAD3756	igVCA0265-VCA0266::Cm^R-P$_{chb}$-*cbp^{D310C}-mCherry*, ΔVCA0692:: Tm^R-P$_{BAD}$-*chiS^{ΔDBD}-L5-tetR*, ΔVC0622-VC0618::Smx^R, Δ*lacZ*::Tet^R- P$_{const2}$-*mTFP1*, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-*gfp*	Fig. 5B, S8A-B	*chiS^{WT} cbp^{D310C}* for disulfide and DSS crosslinking
HSA0854 / SAD3757	igVCA0265-VCA0266::Cm^R-P$_{chb}$-*cbp-mCherry*, ΔVCA0692:: Tm^R-P$_{BAD}$-*chiS^{Q243C,ΔDBD}-L5-tetR*, ΔVC0622-VC0618::Smx^R, Δ*lacZ*::Tet^R- P$_{const2}$-*mTFP1*, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-*gfp*	Fig. 5B, S8A-B	*chiS^{Q243C} cbp^{WT}* for disulfide and DSS crosslinking
HSA0855 / SAD3758	igVCA0265-VCA0266::Cm^R-P$_{chb}$-*cbp^{D310C}-mCherry*, ΔVCA0692:: Tm^R-P$_{BAD}$-*chiS^{Q243C,ΔDBD}-L5-tetR*, ΔVC0622-VC0618::Smx^R, Δ*lacZ*::Tet^R- P$_{const2}$-*mTFP1*, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-*gfp*	Fig. 5B, S8A-B	*chiS^{Q243C} cbp^{D310C}* for disulfide and DSS crosslinking
HSA0925 / SAD3759	igVCA0265-VCA0266::Cm^R-P$_{chb}$-*cbp^{D310C}-mCherry*, ΔVCA0692:: Tm^R-P$_{BAD}$-*chiS^{Y245C,ΔDBD}-L5-tetR*, ΔVC0622-VC0618::Smx^R, Δ*lacZ*::Tet^R- P$_{const2}$-*mTFP1*, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-*gfp*	Fig. 5B, S8A-B	*chiS^{Y245C} cbp^{D310C}* for disulfide and DSS crosslinking
HSA0822 / SAD3760	igVCA0265-VCA0266::Cm^R-P$_{chb}$-*cbp-mCherry*, ΔVCA0692::Tm^R-P$_{chiS}$-*chiS^{Q234C,T473A}*, Δ*lacZ*::Tet^R- P$_{const2}$-*mTFP1*, Δ*chis/cbp*::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-*gfp*	Fig. 5C	*chiS^{T473A,Q243C} cbp^{WT}* for ChiS activity assays
HSA0826 / SAD3761	igVCA0265-VCA0266::Cm^R-P$_{chb}$-*cbp-mCherry*, ΔVCA0692::Tm^R-P$_{chiS}$-*chiS^{Y245C,T473A}*, Δ*lacZ*::Tet^R- P$_{const2}$-*mTFP1*, Δchis/cbp::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-*gfp*	Fig. 5C	*chiS^{T473A,Y245C} cbp^{WT}* for ChiS activity assays

HSA0828 / SAD3762	igVCA0265-VCA0266::Cm^R-P$_{chb}$-cbp^{D310C}-mCherry, ΔlacZ::Tet^R- P$_{const2}$-mTFP1, ΔVCA0692::Tm^R-P$_{chiS}$-chiS^{T473A}, Δchis/cbp::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-gfp	Fig. 5C	chiS^{T473A} cbp^{D310C} for ChiS activity assays
HSA0863 / SAD3763	igVCA0265-VCA0266::Cm^R-P$_{chb}$-cbp^{D310C}-mCherry, ΔVCA0692::Tm^R-P$_{chiS}$-chiS^{Q234C,T473A}, ΔlacZ::Tet^R- P$_{const2}$-mTFP1, Δchis/cbp::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-gfp	Fig. 5C	chiS^{T473A,Q243C} cbp^{D310C} for ChiS activity assays
HSA0915 / SAD3764	igVCA0265-VCA0266::Cm^R-P$_{chb}$-cbp^{D310C}-mCherry, ΔVCA0692::Tm^R-P$_{chiS}$-chiS^{Y245C,T473A}, ΔlacZ::Tet^R- P$_{const2}$-mTFP1, Δchis/cbp::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-gfp	Fig. 5C	chiS^{T473A,Y245C} cbp^{D310C} for ChiS activity assays
HSA0371 / SAD3765	igVCA0265-VCA0266::Cm^R-P$_{chb}$-cbp^{D251R}-mCherry, ΔlacZ::Tet^R- P$_{const2}$-mTFP1, ΔVCA0692::Tm^R-P$_{chiS}$-chiS, ΔchiS/cbp::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-gfp	Fig. S4	chiS^{WT} cbp^{D251R} for ChiS activity assays
HSA0700 / SAD3766	igVCA0265-VCA0266::Cm^R-P$_{chb}$-cbp^{E529R}^{E530R}-mCherry, ΔlacZ::Tet^R- P$_{const2}$-mTFP1, ΔVCA0692::Tm^R-P$_{chiS}$-chiS, ΔchiS/cbp::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-gfp	Fig. S4	chiS^{WT} cbp^{E529R,E530R} for ChiS activity assays
HSA0584 / SAD3767	igVCA0265-VCA0266::Cm^R-P$_{chb}$-cbp^{D231R}-mCherry, ΔlacZ::Tet^R- P$_{const2}$-mTFP1, ΔVCA0692::Tm^R-P$_{chiS}$-chiS, ΔchiS/cbp::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-gfp	Fig. S4	chiS^{WT} cbp^{D231R} for ChiS activity assays
HSA0617 / SAD3768	igVCA0265-VCA0266::Cm^R-P$_{chb}$-cbp^{R254E}-mCherry, ΔlacZ::Tet^R- P$_{const2}$-mTFP1, ΔVCA0692::Tm^R-P$_{chiS}$-chiS, ΔchiS/cbp::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-gfp	Fig. S4	chiS^{WT} cbp^{R254E} for ChiS activity assays
HSA0373 / SAD3769	ΔlacZ::Tet^R- P$_{const2}$-mTFP1, igVCA0265-VCA0266::Cm^R-P$_{chb}$-cbp-mCherry, ΔVCA0692::Tm^R-P$_{chiS}$-chiS^{R99D}, ΔchiS/cbp::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-gfp	Fig. S4	chiS^{R99D} cbp^{WT} for ChiS activity assays
HSA0621 / SAD3770	igVCA0265-VCA0266::Cm^R-P$_{chb}$-cbp-mCherry, ΔVCA0692::Tm^R-P$_{chiS}$-chiS^{E81R}, ΔlacZ::Tet^R- P$_{const2}$-mTFP1, ΔchiS/cbp::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-gfp	Fig. S4	chiS^{E81R} cbp^{WT} for ChiS activity assays
HSA0717 / SAD3771	igVCA0265-VCA0266::Cm^R-P$_{chb}$-cbp-mCherry, ΔVCA0692::Tm^R-P$_{chiS}$-chiS^{R139E}, ΔlacZ::Tet^R- P$_{const2}$-mTFP1, ΔchiS/cbp::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-gfp	Fig. S4	chiS^{R139E} cbp^{WT} for ChiS activity assays

HSA0940 / SAD3772	igVCA0265-VCA0266::Cm^R-P$_{chb}$-*cbp*-*mCherry*, ΔVCA0692::Tm^R-P$_{chiS}$-*chiS^{R89E}*, Δ*lacZ*::Tet^R- P$_{const2}$-*mTFP1*, Δchis/cbp::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-*gfp*	Fig. S4	*chiS^{R89E} cbp^{WT}* for ChiS activity assays
HAS0761 / SAD3773	igVCA0265-VCA0266::Cm^R-P$_{chb}$-*cbp*-*mCherry*, ΔVCA0692::Tm^R-P$_{chiS}$-*chiS^{R146D}*, Δ*lacZ*::Tet^R- P$_{const2}$-*mTFP1*, Δchis/cbp::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-*gfp*	Fig. S4	*chiS^{R146D} cbp^{WT}* for ChiS activity assays
HSA0643 / SAD3774	igVCA0265-VCA0266::Cm^R-P$_{chb}$-*cbp$^{[1-40]}$*-*mCherry*, Δ*lacZ*::Tet^R- P$_{const2}$-*mTFP1*, ΔVCA0692::Tm^R-P$_{chiS}$-*chiS^{R99D}*, Δ*chiS/cbp*::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-*gfp*	Fig. S4	*chiS^{R99D} cbp$^\Delta$* for ChiS activity assays
HSA0697 / SAD3775	igVCA0265-VCA0266::Cm^R-P$_{chb}$-*cbp$^{[1-40]}$*-*mCherry*, ΔVCA0692::Tm^R-P$_{chiS}$-*chiS^{E81R}*, Δ*lacZ*::Tet^R- P$_{const2}$-*mTFP1*, Δ*chiS/cbp*::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::KanR-P$_{chb}$-*gfp*	Fig. S4	*chiS^{E81R} cbp$^\Delta$* for ChiS activity assays
HSA0716 / SAD3776	igVCA0265-VCA0266::Cm^R-P$_{chb}$-*cbp$^{[1-40]}$*-*mCherry*, ΔVCA0692::Tm^R-P$_{chiS}$-*chiS^{R139E}*, Δ*lacZ*::Tet^R- P$_{const2}$-*mTFP1*, Δ*chiS/cbp*::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-*gfp*	Fig. S4	*chiS^{R139E} cbp$^\Delta$* for ChiS activity assays
HSA0941 / SAD3777	igVCA0265-VCA0266::Cm^R-P$_{chb}$-*cbp$^{[1-40]}$*-*mCherry*, ΔVCA0692::Tm^R-P$_{chiS}$-*chiS^{R89E}*, Δ*lacZ*::Tet^R- P$_{const2}$-*mTFP1*, Δ*chiS/cbp*::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-*gfp*	Fig. S4	*chiS^{R89E} cbp$^\Delta$* for ChiS activity assays
HAS0767 / SAD3778	igVCA0265-VCA0266::Cm^R-P$_{chb}$-*cbp$^{[1-40]}$*-*mCherry*, ΔVCA0692::Tm^R-P$_{chiS}$-*chiS^{R146D}*, LacZ::Tet^R- P$_{const2}$-*mTFP1*, Δ*chiS/cbp*::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-*gfp*	Fig. S4	*chiS^{R146D} cbp$^\Delta$* for ChiS activity assays
HSA0936 / SAD3779	igVCA0265-VCA0266::Cm^R-P$_{chb}$-*cbp$^{[1-40]}$*-*mCherry*, ΔVCA0692::Tm^R-P$_{chiS}$-*chiS^{Q243C}*, Δ*lacZ*::Tet^R- P$_{const2}$-*mTFP1*, Δ*chiS/cbp*::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-*gfp*	Fig. S4	*chiS^{Q243C} cbp$^\Delta$* for ChiS activity assays
HSA0937 / SAD3780	igVCA0265-VCA0266::Cm^R-P$_{chb}$-*cbp$^{[1-40]}$*-*mCherry*, ΔVCA0692::Tm^R-P$_{chiS}$-*chiS^{Y245C}*, Δ*lacZ*::Tet^R- P$_{const2}$-*mTFP1*, Δ*chiS/cbp*::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-*gfp*	Fig. S4	*chiS^{Y245C} cbp$^\Delta$* for ChiS activity assays
HSA0355 / SAD3781	Δ*lacZ*::Tet^R- P$_{const2}$-*mTFP1*, igVCA0265-VCA0266::Cm^R-P$_{chb}$-*cbp$^{W389A\ W539A}$*-*mCherry*, ΔVCA0692::Tm^R-P$_{chiS}$-*chiS*, Δ*chiS/cbp*::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-*gfp*	Fig. S5	*chiS^{WT} cbp** for ChiS activity assays

HSA0714 / SAD3782	igVCA0265-VCA0266::Cm^R-P$_{chb}$-$cbp^{[1-40]}$-$mCherry$, ΔVCA0692::Tm^R-P$_{chiS}$-$chiS^{E81R}$ ^{T473A}, Δ$lacZ$::Tet^R- P$_{const2}$-$mTFP1$, Δ$chiS/cbp$::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-gfp	Fig. S6	$chiS^{T473A,E81R}$ cbp^{Δ} for ChiS activity assays
HSA0629 / SAD3783	igVCA0265-VCA0266::Cm^R-P$_{chb}$-$cbp^{[1-40]}$-$mCherry$, ΔVCA0692::Tm^R-P$_{chiS}$-$chiS^{R99D}$ ^{T473A}, Δ$lacZ$::Erm^R- P$_{const2}$-$mTFP1$, Δ$chiS/cbp$::Tet^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-gfp	Fig. S6	$chiS^{T473A,R99D}$ cbp^{Δ} for ChiS activity assays
HSA0665 / SAD3784	igVCA0265-VCA0266::Cm^R-P$_{chb}$-$cbp^{[1-40]}$-$mCherry$, ΔVCA0692::Tm^R-P$_{chiS}$-$chiS^{R139E,T473A}$, Δ$lacZ$::Tet^R- P$_{const2}$-$mTFP1$, Δ$chiS/cbp$::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-gfp	Fig. S6	$chiS^{T473A,R139E}$ cbp^{Δ} for ChiS activity assays
HAS0766 / SAD3785	igVCA0265-VCA0266::Cm^R-P$_{chb}$-$cbp^{[1-40]}$-$mCherry$, ΔVCA0692::Tm^R-P$_{chiS}$-$chiS^{R89E}$ ^{T473A}, Δ$lacZ$::Tet^R- P$_{const2}$-$mTFP1$, Δ$chiS/cbp$::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-gfp	Fig. S6	$chiS^{T473A,R89E}$ cbp^{Δ} for ChiS activity assays
HAS0768 / SAD3786	igVCA0265-VCA0266::Cm^R-P$_{chb}$-$cbp^{[1-40]}$-$mCherry$, ΔVCA0692::Tm^R-P$_{chiS}$-$chiS^{R146D}$ ^{T473A}, Δ$lacZ$::Tet^R- P$_{const2}$-$mTFP1$, Δ$chiS/cbp$::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-gfp	Fig. S6	$chiS^{T473A,R146D}$ cbp^{Δ} for ChiS activity assays
HSA0367 / SAD3787	Δ$lacZ$::Tet^R- P$_{const2}$-$mTFP1$, igVCA0265-VCA0266::Cm^R-P$_{chb}$-cbp-$mCherry$, ΔVCA0692::Tm^R-P$_{chiS}$-$chiS^{R99D\ T473A}$, Δ$chiS/cbp$::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-gfp	Fig. S7B	$chiS^{T473A,R99D}$ cbp^{WT} for ChiS activity assays
HSA0546 / SAD3788	igVCA0265-VCA0266::Cm^R-P$_{chb}$-cbp^{D231R}-$mCherry$, Δ$lacZ$::Tet^R- P$_{const2}$-$mTFP1$, ΔVCA0692::Tm^R-P$_{chiS}$-$chiS^{T473A}$, Δ$chiS/cbp$::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-gfp	Fig. S7B	$chiS^{T473A}$ cbp^{D231R} for ChiS activity assays
HSA0556 / SAD3789	igVCA0265-VCA0266::Cm^R-P$_{chb}$-cbp-$mCherry$, ΔVCA0692::Tm^R-P$_{chiS}$-$chiS^{R139E}$ ^{T473A}, Δ$lacZ$::Tet^R- P$_{const2}$-$mTFP1$, Δ$chiS/cbp$::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-gfp	Fig. S7B	$chiS^{T473A,R139E}$ cbp^{WT} for ChiS activity assays
HSA0638 / SAD3790	igVCA0265-VCA0266::Cm^R-P$_{chb}$-cbp^{D231R}-$mCherry$, ΔVCA0692::Tm^R-P$_{chiS}$-$chiS^{R99D}$ ^{T473A}, Δ$lacZ$::Erm^R- P$_{const2}$-$mTFP1$, Δ$chiS/cbp$::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-gfp	Fig. S7B	$chiS^{T473A,R99D}$ cbp^{D231R} for ChiS activity assays
HSA0651 / SAD3791	igVCA0265-VCA0266::Cm^R-P$_{chb}$-cbp^{E529R} ^{E530R}-$mCherry$, Δ$lacZ$::Tet^R- P$_{const2}$-$mTFP1$, ΔVCA0692::Tm^R-P$_{chiS}$-$chiS^{T473A}$, Δ$chiS/cbp$::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-gfp	Fig. S7B	$chiS^{T473A}$ $cbp^{E529R,E530R}$ for ChiS activity assays

HSA0896 / SAD3792	igVCA0265-VCA0266::Cm^R-P$_{chb}$-cbp^{E529R,E530R}-mCherry, ΔVCA0692::Tm^R-P$_{chiS}$-chiS^{R99D,T473A}, ΔlacZ::Erm^R- P$_{const2}$-mTFP1, Δchis/cbp::Tet^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-gfp	Fig. S7B	chiS^{T473A,R99D} cbp^{E529R,E530R} for ChiS activity assays
HSA0666 / SAD3793	igVCA0265-VCA0266::Cm^R-P$_{chb}$-cbp^{D231R}-mCherry, ΔVCA0692::Tm^R-P$_{chiS}$-chiS^{R139E,T473A}, ΔlacZ::Tet^R- P$_{const2}$-mTFP1, Δchis/cbp::Erm^R, ΔVC1807::ZeoR, igVCA0587-VCA0588::Kan^R-P$_{chb}$-gfp	Fig. S7B	chiS^{T473A,R139E} cbp^{D231R} for ChiS activity assays
HSA0871 / SAD3794	igVCA0265-VCA0266::Cm^R-P$_{chb}$-cbp^{E401K}-mCherry, ΔVCA0692::Tm^R-P$_{chiS}$-chiS^{R89E,T473A}, ΔlacZ::Tet^R- P$_{const2}$-mTFP1, ΔchiS/cbp::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-gfp	Fig. S7B	chiS^{T473A,R89E} cbp^{E401K} for ChiS activity assays
HSA0891 / SAD3795	ΔlacZ::Tet^R- P$_{const2}$-mTFP1, igVCA0265-VCA0266::Cm^R-P$_{chb}$-cbp^{E401K}-mCherry, ΔVCA0692::Tm^R-P$_{chiS}$-chiS^{T473A}, ΔchiS/cbp::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-gfp	Fig. S7B	chiS^{T473A} cbp^{E401K} for ChiS activity assays
HAS0759 / SAD3796	igVCA0265-VCA0266::Cm^R-P$_{chb}$-cbp-mCherry, ΔVCA0692::Tm^R-P$_{chiS}$-chiS$^{R89E\ T473A}$, ΔlacZ::Tet^R- P$_{const2}$-mTFP1, ΔchiS/cbp::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-gfp	Fig. S7B	chiS^{T473A,R89E} cbp^{WT} for ChiS activity assays
HAS0760 / SAD3797	igVCA0265-VCA0266::Cm^R-P$_{chb}$-cbp^{E401R}-mCherry, ΔVCA0692::Tm^R-P$_{chiS}$-chiS$^{R89E\ T473A}$, ΔlacZ::Tet^R- P$_{const2}$-mTFP1, ΔchiS/cbp::Erm^R, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-gfp	Fig. S7B	chiS^{T473A,R89E} cbp^{E401R} for ChiS activity assays
HAS0812 / SAD3798	VCA0692:: Tm^R-P$_{BAD}$-chiS^{ΔDBD}-L5-tetR, ΔVC0622-VC0618::Smx^R, ΔlacZ::Tet^R-P$_{const2}$-mTFP1, ΔVC1807::Zeo^R, igVCA0587-VCA0588::Kan^R-P$_{chb}$-gfp, pMMB-tfoX Carb^R	Fig. S8A	chiS^{WT} cbp^{Δ} for DSS crosslinking

*Note that duplicate strain identifiers signify the same strain stocked in two independent strain collections.

Table S2. *Primers used in this study.*

Primer	Sequence (5' → 3')*	Description
Primers for SOE deletions		
ABD123	ATTCCGGGGATCCGTCGAC	Amplify all Ab^R cassettes F
ABD124	TGTAGGCTGGAGCTGCTTC	Amplify all Ab^R cassettes R
BBC3226	GTATCGATCAGCAAGGACAAC	ΔVC1807 F1
ABD340	gtcgacggatccccggaatACGTTTCATTAGTCACCTCTATTGTTAACTTGTTC	ΔVC1807 R1
ABD341	gaagcagctccagcctacaTAGTCGAAAATAAAAAAAAGAGGCTCGCCTC	ΔVC1807 F2
BBC2412	CAATTTTGCTTTTGGACCATCC	ΔVC1807 R2
BBC4889	ATCAATAAGCAGGCAGATGCG	Δ*crvA* (VCA1075) F1
BBC1783	gtcgacggatccccggaatCCACATAAAGTGGGAAAGACAAAC	Δ*crvA* (VCA1075) R1
BBC1784	gaagcagctccagcctacaGTCAATGGCAATGACACGG	Δ*crvA* (VCA1075) F2
BBC4890	GTTGACTTCTTGCTCAAAAGTG	Δ*crvA* (VCA1075) R2
CKP243	GACAGTGACTTGCTATGAAG	Δ*chiS/cbp* (ΔVC0622/VC0620) or ΔVC0622-VC0618 F1
ABD768	gtcgacggatccccggaatCAAAAAACGTGAGGAGAATGCC	Δ*chiS/cbp* (ΔVC0622/VC0620) or ΔVC0622-VC0618 R1
ABD798	gaagcagctccagcctacaGTACTGGATCTGAAACCAGTTAAG	Δ*chiS/cbp* (ΔVC0622/VC0620) F2
BBC3154	GTCATGATTATTTATCTCCGCTCC	Δ*chiS/cbp* (ΔVC0622/VC0620) R2
BBC0083	gaagcagctccagcctacaGGAGATAAATAATCATGACTACGCC	ΔVC0622-VC0618 F2
BBC0084	TAAAGTTCGCAACACGCC	ΔVC0622-VC0618 R2
Primers for CBP, ChiS & reporter constructs		
HAP0007	GGCAGCACTTCGTTCATAG	Amplify igVCA0265-266::P$_{chb}$-*cbp-mCherry* constructs F1
HAP0012	GATACGCCTATAGTTCCCATG	Amplify igVCA0265-266::P$_{chb}$-*cbp-mCherry* constructs R2
HAP0043	tccaccacttccacctgcTGTAGGGTAGAAATCTGGCACG	*cbp*^{Δ} junction (*i.e.,* the *cbp* signal sequence fused to mCherry) R1
HAP0218	ACACAACGTCTTCTCGTAGG	Amplify ΔVCA0692::P$_{chiS}$-*chiS* constructs F1
VGP0918	GTCGTCGGTTTAGGGTATTCATC	Amplify ΔVCA0692::P$_{chiS}$-*chiS* constructs R2
ABD332	GGCTGAACGTGGTTGTCGAAAATGAC	Amplify Δ*lacZ*::P$_{const2}$-*mTFP1* constructs F1
ABD256	CCCAAATACGGCAACTTGGCG	Amplify Δ*lacZ*::P$_{const2}$-*mTFP1* constructs R2
BBC2511	CTCATTAGGCACCCCAGGC	Insert *mTFP1* in P$_{const2}$ construct R1
ABD255	gaagcagctccagcctacaCCACAATAAGCCAGAGAGCCTTAAG	Insert *mTFP1* in P$_{const2}$ construct F2
BBC3287	GCCTGGGGTGCCTAATGAGcacagctaacaccacgtc	Amplify mTFP1 for *P$_{const2}$* construct F
BBC4272	tgtaggctggagctgcttcCTATTTATACAGTTCATCCATACCATC	Amplify mTFP1 for *P$_{const2}$* construct R

Primers for point mutations		
HAP0234	TCTCCAAACGGCgccACTGACTTTAACAACACAGTTCAACTGGCGG	*cbp*^{W389A} F2
HAP0235	TTAAAGTCAGTggcGCCGTTTGGAGATTGGATCAGCAGTTCAAAAGATTT	*cbp*^{W389A} R1
HAP0236	TGTTGTTAAAGTCAGcggc	*cbp*^{W389A} detect
HAP0022	CCAAGCGTTgcgGCTGGTATCCCAGAGCGTCTACTACACGTAC	*cbp*^{W539A} F2
HAP0023	CTGGGATACCAGCcgcAACGCTTGGACGGCCCTTCGGATTTTC	*cbp*^{W539A} R1
HAP0024	CGTCCAAGCGTggcg	*cbp*^{W539A} detect
CKP228	CACATGAATTACGCgCCCCATTGCATGGCATGATTG	*chiS*^{T473A} F
CKP229	GCCATGCAATGGGGcGCGTAATTCATGTGATGTGG	*chiS*^{T473A} R
CKP230	CACATCACATGAATTACGtg	*chiS*^{T473A} detect
HAP0083	ACTGGCGGTTcgtCAGCTACAAGAAGTGGGTATCAAAGCGAAAGCAC	*cbp*^{E401R} F2
HAP0084	CTTCTTGTAGCTGacgAACCGCCAGTTGAACTGTGTTGTTAAAGTCAGTC	*cbp*^{E401R} R1
HAP0091	CCACTTCTTGTAGCTaacg	*cbp*^{E401R} detect
HAP0153	TGAACGCTATgaACTGCTGCATgAgCGCTACCACTGGGCTTACTTAGA	*chiS*^{R141E,K145E} F2
HAP0154	GcTcATGCAGCAGTtcATAGCGTTCACTGCCCACGTAGCTGCTT	*chiS*^{R141E,K145E} R1
HAP0155	GGGCAGTGAACGCTAcga	*chiS*^{R141E,K145E} detect
HAP0077	ATTCGTTCCAcgTATCGACCGTACTTACGCCGCAGCGAACCCTAACCA	*cbp*^{D251R} F2
HAP0078	AAGTACGGTCGATAcgTGGAACGAATGAAGAAGTCCAGTCGAGCTCAG	*cbp*^{D251R} R1
HAP0088	TTCTTCATTCGTTCCgcg	*cbp*^{D251R} detect
HAP0297	GCTGCATAAAgatTACCACTGGGCTTACTTAGAACTGCTGAAACG	*chiS*^{R146D} F2
HAP0298	CCCAGTGGTAatcTTTATGCAGCAGTCGATAGCGTTCACTGCC	*chiS*^{R146D} R1
HAP0299	TAAGTAAGCCCAGTGGTgat	*chiS*^{R146D} detect
HAP0301	AACCATCATTtgcATCGCCTTCTACGGCAGCGGTACTGTGAAC	*cbp*^{D310C} F2
HAP0302	AGAAGGCGATgcaAATGATGGTTTGACGATCCAGAGCCATAGA	*cbp*^{D310C} R1
HAP0303	CGTCAAACCATCATctgc	*cbp*^{D310C} detect
HAP0066	GATTATTCAGtgtGGCTACTTGCATAGTTACGCGATGTTCCGCTTAC	*chiS*^{Q243C} F2
HAP0067	TATGCAAGTAGCCacaCTGAATAATCGGTGCGCCAAGCCAAGCGTATTGTTT	*chiS*^{Q243C} R1
HAP0073	CGTAACTATGCAAGTAGCtaca	*chiS*^{Q243C} detect
HAP0313	TCAGCAAGGCTgtTTGCATAGTTACGCGATGTTCCGCTTACCAA	*chiS*^{Y245C} F2
HAP0314	AACTATGCAAacAGCCTTGCTGAATAATCGGTGCGCCAAGCCAA	*chiS*^{Y245C} R1

HAP0315	ATCGCGTAACTATGCAgac	*chiS*^{Y245C} detect
HAP0240	CGCAAACAACcgtCAACTACTGGGCAAAATCGTTAACTCTGAG	*cbp*^{D231R} F2
HAP0241	CCAGTAGTTGacgGTTGTTTGCGATTTGTGGAACACGCAGACA	*cbp*^{D231R} R1
HAP0242	CAAATCGCAAACAAtcgt	*cbp*^{D231R} detect
HAP0261	AGATATCGACgaaACTTACGCCGCAGCGAACCCTAACCACCAC	*cbp*^{R254E} F2
HAP0262	CGGCGTAAGTttcGTCGATATCTGGAACGAATGAAGAAGTCCA	*cbp*^{R254E} R1
HAP0263	GTTCCAGATATCGAtgaa	*cbp*^{R254E} detect
HAP0252	GGTGGAGCcgtcgtAATCCGAAGGGCCGTCCAAGCGTTTGGGC	*cbp*^{E529R,E530R} F2
HAP0253	CTTCGGATTacgacgGCTCCACCAGCCAGTGAAACGAGTGGTG	*cbp*^{E529R,E530R} R1
HAP0254	GCCCTTCGGATTacgacg	*cbp*^{E529R,E530R} detect
HAP0264	TGTTCGCTCCcgtCTGGCTTATGCCAGTGGCGGGCGTTTTTAT	*chiS*^{E81R} F2
HAP0265	CATAAGCCAGacgGGAGCGAACAAAACCCATCACTTCCGACTC	*chiS*^{E81R} R1
HAP0266	ACTGGCATAAGCCAaacg	*chiS*^{E81R} detect
HAP0279	CAGTGGCGGGgaaTTTTATGGTCTGGTGAATGCTTTCCAGCGT	*chiS*^{R89E} F2
HAP0280	GACCATAAAAttcCCCGCCACTGGCATAAGCCAGCTCGGAGCG	*chiS*^{E81R} R1
HAP0281	CACCAGACCATAAAgttc	*chiS*^{E81R} detect
HAP0054	TGCTTTCCAGgaTTTAGACGTCAGTATTGAAGCGGCTCGTGAACAT	*chiS*^{R99D} F2
HAP0055	TACTGACGTCTAAAtcCTGGAAAGCATTCACCAGACCATAAAAACGCC	*chiS*^{R99D} R1
HAP0068	TGGTGAATGCTTTCCAaga	*chiS*^{R99D} detect
HAP0258	GGGCAGTGAAgaaTATCGACTGCTGCATAAACGCTACCACTGG	*chiS*^{R139E} F2
HAP0259	GCAGTCGATAttcTTCACTGCCCACGTAGCTGCTTGATTGGGG	*chiS*^{R139E} R1
HAP0260	ATGCAGCAGTCGATgttc	*chiS*^{R139E} detect
HAP0149	ACTGGCGGTTaAaCAGCTACAAGAAGTGGGTATCAAAGCGAAAGCAC	*cbp*^{E401K} F2
HAP0150	CTTCTTGTAGCTGtTtAACCGCCAGTTGAACTGTGTTGTTAAAGTCAGTC	*cbp*^{E401K} R1
HAP0151	CCACTTCTTGTAGCTGtct	*cbp*^{E401K} detect

*Lower nucleotides at the 5' end denote overlap sequences, while internal lowercase nucleotides denote point mutations

Dataset S1. *ChimeraX analysis of contact residues in CBP-ChiS^{AF3} and CBP-ChiS^{AF-M}.*

Dataset S2. *Summary statistics and statistical comparisons for all quantitative data.*