Mechanistic insights and clinical implications of cross-reactive anti-prophage antibodies and bacterial heteroresistance on phage therapeutic failure

Gordillo Altamirano F[1,2], Subedi D[2,3], Beiers M[1], Bucher M[3], Dahlman S[3], Patel DM[4], Parker M[1], Korneev D[5], Robinson MJ[6], Pragastis K[1], Wisniewski J[1], Rees C[7], Ramshaw H[4], Khan SF[1,8], Gardiner BJ[1], Hammerschlag Y[1], Keating D[9], Kotsimbos T[9], Hawkey J[1], Barr JJ[2,3], Peleg AY[1,2,10]*

[1]Department of Infectious Diseases, Alfred Health and School of Translational Medicine, Monash University, VIC, Australia

[2]Centre to Impact AMR, Monash University, VIC, Australia

[3]School of Biological Sciences, Monash University, VIC, Australia

[4]Antibody Discovery Platform, Monash University, VIC, Australia

[5]Ramaciotti Centre for Cryo-Electron Microscopy, Monash University, VIC, Australia

[6]Department of Immunology, School of Translational Medicine, Monash University, VIC, Australia

[7]Melbourne Water Corporation, VIC, Australia

[8]Microbiology Unit, Department of Pathology, The Alfred Hospital, Melbourne, VIC, Australia

[9]Department of Respiratory Medicine, The Alfred Hospital and School of Translational Medicine, Melbourne, VIC, Australia

[10]Infection Program, Monash Biomedicine Discovery Institute, Department of Microbiology, Monash University, VIC, Australia

*Corresponding author: Professor Anton Y. Peleg MBBS PhD MPH FRACP FAHMS
Department of Infectious Diseases
Alfred Health and School of Translational Medicine, Monash University
Alfred Lane House, Level 1, 55 Commercial Road, Melbourne, 3004, VIC, Australia
+61 3 9076 6076
anton.peleg@monash.au

Word count (main text): 4,094 words (excluding abstract, methods, references and figure legends).

Abstract

Phage therapy is an exciting strategy against antimicrobial-resistant bacterial infections, but critical knowledge gaps regarding its clinical application persist. Studying a patient with a life-threatening, chronic bacterial infection who failed phage therapy, we uncovered important biological concepts with direct translational impact. Using longitudinal clinical samples, we found that patients can harbour pre-existing antibodies against active prophages induced from the genome of the causative pathogen. Notably, these antibodies can contribute to clinical failure by cross-reacting with and effectively neutralising therapeutic phage. We also uncovered bacterial heteroresistance, characterised by bacterial subpopulations from the initial infection with reduced phage susceptibility, as a further contributor to treatment failure. These findings highlight the intricate interplay between host

immunology, bacterial genetic diversity and phage biology, bearing broad significance for clinical phage therapy. Future phage therapy patients, especially those with chronic infections, should be screened for antiphage immunity and bacterial heteroresistance prior to phage treatment.

Introduction

Antimicrobial resistance (AMR) is one of the greatest health threats we currently face. The latest global estimations suggest that in 2021, bacterial AMR directly caused over a million deaths, while playing a role in just under 5 million [1]. AMR is not only a leading cause of death but poses a significant burden on the global economy and healthcare systems [2]. These consequences are expected to increase unless a concerted, multi-disciplinary effort is deployed to fight AMR, including the development of therapeutic strategies for when antibiotics prove ineffective, such as phage therapy.

Phage therapy is the clinical use of bacteriophages, viruses that specifically kill bacteria, to treat bacterial infections [3]. Although phages were discovered and used before antibiotics, phage therapy is now experiencing a resurgence amidst the AMR crisis. In the last two decades, we have accumulated a significant body of evidence from preclinical studies suggesting that phage therapy is safe and potentially effective [4], encouraging translational efforts. As for clinical use, a recent systematic review examined 2,241 patients from 59 studies and deemed phage therapy a promising strategy in the fight against difficult-to-treat infections and AMR [5]. However, most of this clinical experience comes from case reports and case-series, with only a handful of clinical trials. Substantial knowledge gaps remain to be addressed before phage therapy becomes standardised and widespread, including questions around the pharmacology of phages, their interactions with antibiotics, and the tripartite relationship between phages, bacteria and the human host [6].

We present a patient with a chronic, extensively drug-resistant (XDR) bacterial (*Bordetella bronchialis*) infection associated with life-threatening infective pulmonary exacerbations and bacteraemia that was poorly controlled with traditional antibiotics. The patient qualified for compassionate use phage therapy. We report the isolation and detailed characterisation of the therapeutic phage, and the clinical course of the patient that resulted in phage therapeutic failure with bacteriological relapse. We explored the complex *in vivo* dynamics of host immunology, bacteriology, and phage biology, and have provided essential clinical insights into the future treatment of life-threatening bacterial infections with bacteriophages.

Results and Discussion

Recurrent life-threatening infection with extensively drug-resistant bacteria

A 22-year-old male patient with cystic fibrosis (CF) was referred to the Victorian Bacteriophage Therapy Program (VIC*Phage*) in 2022 for compassionate use phage treatment of recurrent, invasive *Bordetella bronchialis* infection, including pneumonia and bacteraemia. He had a history of a liver-pancreas transplant six years prior for CF-related liver disease and diabetes mellitus, and end-stage kidney

81 disease secondary to calcineurin inhibitor use, aminoglycoside exposure, and diabetes mellitus. He
82 was on maintenance immunosuppression with tacrolimus and prednisone.
83
B. bronchialis was first isolated from the patient's sputum at the age of 10 years. *B. bronchialis*, known
before 2015 as *Bordetella* genogroup 3, are clinically uncommon motile Gram-negative bacilli that are
facultatively aerobic and slow-growing [7]. Despite reported cases of this pathogen in patients with CF
[8,9], there is limited information regarding its pathogenesis, biology, or epidemiology. Through his
teenage years, the patient developed recurrent infective respiratory exacerbations with repeated
hospitalisation and isolation of *B. bronchialis*. Initially, the isolate tested susceptible to ceftazidime and
piperacillin-tazobactam (Table S1). However, repeated infections ensued and in the three years prior to
referral, he was admitted nine times for infective exacerbations, including three episodes of invasive
bacteraemia, an uncommon complication in patients with CF. To prevent or reduce the frequency of
these life-threatening exacerbations, ongoing minocycline and trimethoprim-sulfamethoxazole were
commenced in 2021.
95
In June 2022, the patient presented with fever, productive cough, and dyspnoea. Breakthrough *B.*
bronchialis was isolated from sputum and multiple sets of blood cultures. Chest imaging demonstrated
diffuse, severe bilateral bronchiectasis with extensive mucous plugging and patchy foci of consolidation.
He was commenced on empiric IV meropenem and inhaled colistin, and antimicrobial susceptibility
testing demonstrated that the isolate had become XDR (Table S1, Note S1). At this point, referral for
compassionate use phage treatment was received.
102

Isolation and characterisation of an active phage

Using wastewater samples from Melbourne, Australia, and cultures of a single *B. bronchialis* colony
obtained from the patient's sputum in June 2022 (wild type [wt] *B. bronchialis*), we successfully
discovered, isolated and amplified phage øSimón (Fig. 1a). Multiple efforts to isolate additional phages
were unsuccessful. In liquid coculture, øSimón inhibited bacterial growth for up to 40 hours at
multiplicities of infection (MOI) as low as 0.01 (Fig. 1b), and consistently achieved titres above 10^8
plaque forming units (pfu)/ml. In a one-step kill curve, we determined that øSimón had a latency time
of 2.5 h, more than three times faster than the calculated doubling time for the host strain (8.3 ± 2.6 h),
and a burst size of nearly 30 virions per cell (Fig. 1c). In artificial sputum media, the bactericidal activity
of øSimón was slower, less profound, but more sustained than in liquid culture (Fig. 1d), while also
resulting in slightly higher phage titres (Fig. 1e). øSimón displayed similar activity against five other
clinical *B. bronchialis* isolates (three from blood, two from sputum) obtained from the patient prior to
starting phage therapy (Fig. 1f). Transmission electron microscopy (TEM) showed that øSimón was
morphologically a siphovirus, with a total length of ~226 nm (Fig. 1g). øSimón's genome did not encode
for any markers of lysogeny, AMR, toxins or virulence (Fig. 1h, Note S2), making it an ideal candidate
for therapeutic use.

Figure 1. Characterisation of phage øSimón. a: Plaque morphology on HI agar, 48 h of incubation. Scale bar: 1 cm. **b:** Optical density at 600 nm (OD_{600}) growth curve (mean ± SD; n = 3) of *B. bronchialis* without phage (black) and with three different multiplicities of infection (MOI) of øSimón. **c:** One-step kill curve (median ± 95% CI; n = 3). **d and e**: Comparison of øSimón's bactericidal activity and amplification in liquid (heart infusion broth, black) and artificial sputum media (magenta) (median ± 95% CI; n = 3; dotted lines represent the limit of detection). **f:** Sensitivity of further *B. bronchialis* isolates from the patient against øSimón in liquid culture (growth curve, mean ±

SD; n = 3). Isolates are labelled by type of clinical sample and date of isolation. **g:** øSimón virion under transmission electron microscopy. Scale bar: 50 nm. **h:** Assembled øSimón genome (60,921 bp, 80 open reading frames).

Clinical administration of øSimón

A clinical-grade preparation of øSimón was produced locally at the Monash Phage Foundry. All product batches (Table S2) underwent internal and third-party validation for sterility and endotoxin levels suitable for IV administration [10]. In the setting of a repeated episode of invasive bacteraemia with XDR *B. bronchialis*, the patient received the first dose of IV øSimón (5.4×10^7 pfu/dose) (day 0) (Fig. 2). Phage therapy was administered twice daily and, given the severity of infection, was co-administered alongside IV meropenem and inhaled colistin. The patient stabilised clinically, with blood cultures clearing by day 5. After close clinical and laboratory monitoring for six days, the patient was transferred to our hospital-in-the-home program for ongoing IV phage management, completing 30 days [11]. Phage therapy was well tolerated, with no clinical or laboratory adverse effects reported (Fig. S1).

Alongside these clinical outcomes, we observed an almost 80% reduction in the amount of *B. bronchialis* in the patient's sputum at day 8 of phage treatment, and notably, this coincided with the finding of actively replicating øSimón (Fig. 2). However, by day 17, the *B. bronchialis* load increased back to pre-treatment levels, and øSimón was no longer isolated from the sputum. The patient went on to develop progressive complications of his end-stage renal disease. In the context of a repeat admission for worsening illness (day 68), he developed further episodes of *B. bronchialis* bacteraemia and passed away on day 93, two months after the last dose of øSimón. Given the undulating clinical and microbiological course, we investigated the possible mechanisms of phage therapy failure and the dynamic interactions between bacteria, phage and human host over time.

Pre-existing antiphage humoral immunity

Despite phages being non-pathogenic to humans, it has been confirmed that the human immune system produces antibodies against them [12]. Antiphage antibodies can neutralise phage activity [13], either directly, by targeting surface-exposed structural proteins involved in bacterial adsorption [14,15], or, to a lesser degree, indirectly by opsonising them for phagocytosis [12,15]. However, how much this immune response, or lack thereof, contributes to the outcomes of phage therapy is still debated [16-19]. This humoral response is T-cell dependent, and activated B-cells produce a specific, polyclonal, antiphage antibody response [12,20]. Circulating antibodies usually peak one to two weeks after initial contact with a phage [12,21], aligning with the time we observed a reduction of phage activity in our patient.

Using patient serum samples from days 0, 7, and 21, we first performed a phage neutralisation assay [19]. Briefly, we tested the ability of three successive 10-fold serum dilutions to neutralise a known dose of øSimón in 30 min. We used an unrelated, anti-*Pseudomonas aeruginosa* phage (øFaith) as a control under the same conditions. Notably, at day 0, before the patient had received the first dose of øSimón, the serum could neutralise ~25% of the phage dose (Fig. 3a). The neutralisation increased to ~60% on days 7 and 21. No neutralisation was seen against the control phage. We also used a more standardised metric known as the phage inactivation rate (k), which generates "high", "moderate", and "low" thresholds, inversely correlating with odds of treatment success [17,19]. We showed that the patient's pre-treatment serum had a "moderate" k value of 10.69, which increased to "high" at days 7 and 21 (27.67 and 30.60, respectively) (Fig. 3b). We also ruled out that the neutralisation seen by our assay could have been mediated by an effect of the serum aimed at the bacterial surface instead of the phage virions (Fig. S2a and S2b). Interestingly, we also observed low phage neutralisation capacity in the patient's sputum (Fig. 3c and 3d).

To confirm that the phage inactivation was mediated by anti-øSimón antibodies, we designed an indirect ELISA assay. Intact, highly purified [22] øSimón was used as the coating antigen. To further discriminate the presence of confounding antibodies against residual endotoxin or other bacterial products, we also used a vehicle control obtained by filtering out (0.02 μm) phage particles from the coating product. Sera from the patient were run in parallel with a serum control from a healthy patient, and all tests used an isotype pan-specific detection antibody. The ELISA confirmed the presence of anti-øSimón antibodies at days 0, 7, and 21, with no quantitative differences between days (Fig. 3e). Anti-øSimón antibodies were also detected in the patient's sputum, although the signal was markedly less intense than in serum (Fig. 3f). Interestingly, a significant peak of antibody was detected in the sputum on days 8 and 17, while

the patient was receiving IV phage therapy, compared to follow-up sputum from day 84, almost two
months after phage therapy had stopped. We next sought to visualise the phage structures being
targeted by the antibodies. To do this, we purified the IgG fraction from the patient's serum and used it
in an immuno-TEM experiment with øSimón and gold-labelled anti-human IgG detection antibodies. We
observed that the patient's antibodies were binding to multiple sites on the virion, including the capsid,
neck, tail and tail fibres (Fig. 3g, 3h and S3). Lastly, we used serial passages of a day 0 serum sample
on ELISA wells coated with øSimón to deplete the serum of anti-øSimón antibodies (Fig. S4a, Note S3).
We repeated the neutralisation assay with this depleted serum and observed that the neutralisation
effect was reversed (Fig. S4b), confirming the role of the antibodies in phage inactivation. To summarise,
we demonstrated the presence of antibodies capable of recognising and inactivating øSimón in both
serum and sputum of the patient even prior to phage therapy. These antibody levels were maintained
throughout the duration of therapy and beyond, but their neutralising effect increased over time,
suggesting affinity maturation [23].

208
209

Figure 3. The patient harboured pre-existing antiphage immunity. a: Serum neutralisation assay (10^6 pfu). The x-axis follows the treatment timeline, and the y-axis plots the percentage of active phage (soft overlay agar assay)

(median ± 95% CI; n = 3). **b:** Calculation of phage inactivation rate (*k*) with data from the previous assay (median ± 95% CI; n = 3). **c and d:** The experiments depicted in a and b were repeated using patient's filter-sterilised sputum instead of serum. **e and f:** Indirect ELISA looking for anti-øSimón antibodies in serum (e) and sputum (f). Controls were run with serum from a healthy donor or sputum from a patient with cystic fibrosis (median ± 95% CI; all replicates from two experiments plotted; n ≥ 6; Kruskal-Wallis test; if not plotted the comparison was not statistically significant). **g and h:** Immuno-TEM micrographs of øSimón with pre- and post-treatment patient serum, respectively, and gold-labelled anti-human IgG detection antibodies (black dots with magenta arrowheads). Scale bars: 50 nm. Additional micrographs and controls in Fig. S3.

Resident prophages as possible drivers of pre-existing humoral immunity to therapeutic phage

We investigated a possible explanation for the presence of pre-existing anti-øSimón antibodies. Given the seemingly infinite diversity of phages in the biosphere [24], we believe it was unlikely for the patient to have previously encountered øSimón. However, as the patient had harboured the *B. bronchialis* infection for over a decade, we hypothesised he may have been exposed to other phages with activity against the pathogen, sharing similar structural features to øSimón, and triggering a cross-reactive immune response. Furthermore, recent studies using cell-free DNA sequencing have demonstrated that patients with sepsis have an overrepresentation of pathogen-specific phages [25,26]. This could be explained by prophages, which are phages that have integrated into a bacterium's chromosome and occasionally re-enter the lytic cycle in periods of rapid bacterial expansion or stress [26]. Consequently, we sought to investigate whether prophages integrated into the patient's *B. bronchialis,* and reactivated at different times during the chronic infection, could have been responsible for cross-reactive immunity against øSimón.

We parsed the genome of the wt *B. bronchialis* for prophage regions. We identified three prophages with medium-to-high completeness scores [27]. On a nucleotide level, they did not show homology to øSimón (Fig. 4a, S5a). Next, we explored whether these prophages could re-enter the lytic lifecycle by the addition of an induction agent (mitomycin C) to a bacterial liquid culture, or spontaneously in the absence of an induction agent. After purification [22], we confirmed the presence of phage virions through TEM and DNA sequencing, but only in the culture with mitomycin C. TEM micrographs of the suspension detected phage particles at a concentration of ~10^6 virions/ml, with a single (siphovirus) morphology (Fig. 4b). After sequencing, two thirds of the obtained reads mapped to prophage region 2, a 51.5 kb-long sequence with the closest match being *Burkholderia* phage Bups phi1 (GenBank accession numbers EU307292.1-EU307295.1), also a siphovirus. This purified lysate, containing at least one of the induced prophages from the *B. bronchialis* host, was then used for subsequent experiments.

Using the induced prophage suspension, an ELISA was developed to detect anti-prophage antibodies, which were confirmed in the patient's serum but not sputum (Fig. 4c and S6). Although the antibody signal was weaker than in the assays for anti-øSimón antibodies, the two assays are not directly comparable. The concentration of virions in the solution used to coat the ELISA wells was 100-times lower for the induced prophage.

We then tested whether the anti-prophage and anti-øSimón antibodies were cross-reactive. To do this,
we modified our indirect ELISA by adding a pre-capture step; patient serum was first incubated on wells
coated with one of three conditions in parallel: 1) phage vehicle (control), 2) øSimón virions, or 3)
induced prophage virions. Following this pre-adsorption, we performed two standard indirect ELISAs:
one to detect anti-prophage antibodies (Fig. 4d) and another for anti-øSimón antibodies. Theoretically,
pre-adsorption with vehicle should yield the highest signal (baseline), while pre-adsorption with the
same phage as the ELISA target should result in the lowest signal due to specific antibody depletion. If
pre-adsorption with the opposite phage also reduced signal, this would indicate the presence of cross-
reactive antibodies. As shown in Fig. 4e, pre-adsorption with either øSimón or prophage virions led to
comparable reductions in antibody signal against both targets (mean reduction for anti-prophage
antibodies: 2.8% $\pm$ 1.2% with prophage vs. 12% $\pm$ 8.5% with øSimón; mean reduction for anti-øSimón
antibodies: 10.9% $\pm$ 10.8% with øSimón vs. 12.4% $\pm$ 6.8% with prophage). These results support the
presence of serum antibodies capable of simultaneously recognising øSimón and prophage virions.
264
Together, our data suggests that the human body can produce antibodies against induced prophages
from a resident bacterial pathogen. An alternative hypothesis we cannot fully discard is that a different,
unknown, immunogen that the patient had been exposed to drove the production of these antibodies.
Regardless of their immunological origin, these antibodies demonstrated prophage-to-Simón cross-
reactivity, and provide an explanation for the neutralisation activity of the patient's pre-treatment serum
and the potential impact on øSimón efficacy. Our results complement two important earlier findings.
First, Halperin *et al*. showed that patients with acute glomerulonephritis produced antibodies against a
secreted, prophage-encoded, streptococcal hyaluronidase [28]. Second, Fluckiger *et al*. demonstrated T-
cell-mediated immunity aimed at a specific tail length tape measure protein from an enterococcal
prophage in mice and humans, which cross-reacted with tumour-associated antigens [29]. Taken together,
there is a growing body of evidence showing the relevance of prophages and their interactions with the
immune system in various states of human health and disease.
277
Future endeavours in this area include establishing the precise proteins or epitopes responsible for the
observed cross-reactivity. We note, for instance, that at the amino acid level, prophage region 2 has a
gene coding for a minor tail protein, with a modest homology to øSimón (29% amino acid identity, 87%
query coverage) (Fig. S5). Berkson *et al*. have demonstrated that cross-reactive immunity between two
therapeutic myoviruses in a phage cocktail was possibly mediated by tail fibre, tail sheath, and major
capsid proteins as antigens [30]. Upon sequencing further pre-treatment isolates from the patient (see
section "Genetic analysis of bacterial heteroresistance" below), we also found three additional prophage
regions not present in the wt strain that could have played a role in the pre-existing immunity (Table
S3). Other areas worth exploring are the precise impacts of antibody titre and affinity on phage
pharmacokinetics at different body sites, quantifying their consequence on therapeutic effects, and the
influence of other components of the immune system, including phagocytes, on phage activity [21,31,32].

a Prophages in the wt *B. bronchialis* genome

Prophage Region	Length	Completeness Score	Most Similar Genome (MSG)	MSG Morphology	% Mapped reads after Induction
1	39.9 kb	77 (Medium)	Bordetella phage vB_BbrS_PHB09	Siphovirus	0%
2	51.5 kb	86 (Medium)	Burkholderia phage Bups phi1	Siphovirus	67%
3	48.5 kb	91 (High)	Ralstonia phage RSY1	Myovirus	33%

b Induced prophage virion morphology

50 nm

c Anti-Prophage Antibody ELISA in serum

d 1. Pre-capture 2. Transfer 3. Wash and Read

e Pre-capture ELISA (1:2,000 Day 0 serum in PBS)

Figure 4. *Bordetella bronchialis* prophages could have contributed to pre-existing immunity. a: Screening of prophage regions in the genome of wt *B. bronchialis.* Rings from innermost to outermost: genome length, GC%, GC skew and open reading frames. The adjacent chart provides information about the three prophages found. **b:** Transmission electron micrograph of the virions found after a culture of *B. bronchialis* was treated with mitomycin C for prophage induction. Scale bar = 50 nm. **c:** Indirect ELISA looking for anti-prophage antibodies in serum; (median ± 95% CI; all replicates from two experiments plotted; n ≥ 4; Kruskal-Wallis test; ns: not significant). **d:** Experimental setup for the pre-capture ELISA. Hypothetical antibodies colour-coded by their ability to recognise øSimón (blue), prophage (red), both (purple), or neither (green). Pre-capture step (1), serum is passed through wells that have been coated with either vehicle (blank, top), øSimón (blue virions, middle), or prophage (red virions, bottom). The circles beneath wells represent the colour-coded antibodies either binding to phages (dashed border) or remaining in suspension (solid border). Next, the serum is transferred to fresh, prophage-coated wells (2), left for incubation and binding, and discarded. All unbound antibodies are washed away, and the signal is read (3). The assay is interpreted by measuring the signal reduction between the top condition and each of the remaining two. Not shown in the figure, the assay was later repeated using øSimón to coat wells in step (2). **e:** Pre-capture ELISA. Percentage reduction was calculated using the reading for the vehicle pre-capture condition as baseline (bars are medians, all replicates from two experiments plotted; Kruskal-Wallis test for comparisons; ns: not significant, * p < 0.05, ** p < 0.005).

Bacterial heteroresistance as a risk for treatment failure

Bacteria can use a vast repertoire of mechanisms to resist phage infection [33,34], and previous reports have demonstrated emergence of phage resistance during phage therapy [35-38]. While this phenomenon can sometimes be beneficial, for example leading to collateral loss of antibiotic-resistance resulting in phage-antibiotic synergy [36,38,39], it can also be the cause for treatment adjustment [40], or failure [37]. Therefore, we investigated if resistance to øSimón emerged during treatment, possibly contributing to the observed therapeutic failure.

We first assessed the ability and mechanism of the patient's *B. bronchialis* to develop resistance to øSimón *in vitro*. After spotting øSimón on a lawn of wt *B. bronchialis*, we observed colonies growing in the middle of a lysis zone. We purified one of these colonies, confirmed it was phenotypically resistant to øSimón and sequenced its genome. Pairwise genomic comparison found that this isolate harboured a single mutation, a one-nucleotide insertion leading to a frameshift and premature stop codon in the gene *btuB,* coding for the outer membrane vitamin B12 receptor of the same name (Fig. 5a). The BtuB-deficient strain grew in the presence and absence of øSimón (Fig. 5b) and had no fitness defect in minimal media (Fig. S7). An adsorption assay revealed that øSimón virions did not adsorb to this bacterial host (Fig. 5c). BtuB has been previously confirmed as either a sole, dual, or co-receptor for phages infecting the genera *Salmonella, Yersinia* and *Escherichia* [41-44]. All of this suggests that øSimón needs BtuB for adsorption and that, as seen for many other phage-host pairs [45], loss of receptor function is a potential mechanism of *B. bronchialis* resistance to øSimón *in vitro*.

a **Phage øSimón putative receptor**

Outer membrane vitamin B12 receptor BtuB

b ***B. bronchialis* (10⁶ cfu) + øSimón**

c **øSimón Adsorption Assay**

Figure 5. *In vitro* mechanism of resistance against phage øSimón. a: Representation of the *btuB* gene sequence in wt *B. bronchialis* (top) and a mutant resistant to øSimón (bottom). The red arrow and box indicate the insertion of an alanine at position 599, leading to a premature stop codon in the phage-resistant bacterium. **b:** Bacterial growth curves of wt *B. bronchialis* and the BtuB-deficient variant presented in panel (a), with and without phage (mean ± SD; n = 3). **c:** Adsorption assay of øSimón to wt or BtuB-deficient *B. bronchialis* over 90 minutes (median ± 95% CI; n = 3; Kruskal-Wallis test).

We then analysed the patient's sputum samples over time for the presence of phage-resistant bacteria. Instead of starting our screening with individual colonies grown from each sputum sample, we took a population-wide approach. We adapted the inverted spot assay technique [46] whereby a high concentration of øSimón was used as a selection marker on semisolid media to detect putative phage-resistant *B. bronchialis*, with species confirmation performed using MALDI-ToF MS (Fig. 6a). This method allowed us to compare the number of *B. bronchialis* colonies that grew in the absence of øSimón (general population) against the number that grew in its presence (putative resistant population), thus calculating a ratio (Fig. 6b). Surprisingly, we discovered that before phage administration (day -1), a median of 2.3% (range: 1.7% to 11.6%) of the *B. bronchialis* population appeared to resist øSimón. This finding suggested baseline heteroresistance, defined as the presence of bacterial subpopulations with varied susceptibility to an antimicrobial (øSimón in this case), a phenomenon capable of contributing to treatment failure [47]. To further investigate for heteroresistance, we randomly chose and purified 14 bacterial colonies from the pre-treatment sputum sample (day -1) and tested their efficiency of plating (EOP) for øSimón. A range of EOPs were identified across these pre-treatment colonies with most of

352   them having values below 1, indicating reduced susceptibility to øSimón (Fig. 6c). These data supported
353   the findings that the patient was infected with a heteroresistant population of *B. bronchialis.*
354
During and after treatment, the proportion of sputum colonies able to grow in the presence of phage
remained relatively stable, but it reached a peak of 5.3% at day 17, coinciding with the bacteriological
relapse seen in the patient (Fig. 2 and 6b). We randomly picked 6 to 8 colonies from the phage-selection
plates (Fig. 6a) at each timepoint after initiating phage therapy (days 2, 6, 8, 17, 72 and 84, n = 44
colonies) and performed spotting and EOP assays with øSimón. In this subset, we also observed a
range of EOPs, including five strains that were completely resistant to øSimón (EOP below the limit of
detection of 10^{-7}) (Fig. 6d). Mirroring the range in EOPs, we also observed variations in the size and/or
clarity of the plaques øSimón produced on the isolates (Fig. 6d). Together, these findings revealed that
before, during and after phage therapy, the patient harboured a bacterial population that was not
uniformly sensitive to øSimón, and that instead of a sensitive/resistant dichotomy, the effectiveness of
øSimón against the patient's *B. bronchialis* fell within a spectrum. This solidified the hypothesis of
heteroresistance and prompted an exploration of its genetic basis.
367

Genetic analysis of bacterial heteroresistance

Using short-read whole genome sequencing of all 58 isolates (14 pre-treatment and 44 post-treatment)
compared to the assembled genome of the wt *B. bronchialis* (host of isolation of øSimón, June 2022),
we identified broad diversity, with two major clusters and several subclusters (Fig. 6e). Cluster 1 housed
the wt strain, 3 of the 14 (18%) pre-treatment isolates, and 9 of the 44 (20%) post-treatment isolates.
Cluster 2 was larger and included the remaining 11 pre-treatment and 35 post-treatment isolates. Using
a SNP distance matrix (Fig. 6f) we noticed that the cluster differentiation corresponded to the EOP of
the pre-treatment strains. Namely, all 3 strains belonging to cluster 1 had an EOP > 1 indicating high
susceptibility to øSimón, whereas 9 of the 11 strains from cluster 2 had an EOP < 1. We then screened
for specific genes and mutations that may explain the observed heteroresistance by comparing between
the clusters in the pre-treatment bacterial strains. Strains from cluster 2 had a nonsense mutation in
rcsC, which codes for a histidine kinase of the regulator of capsule synthesis (rcs) system. This system
is activated upon cell surface stress, including outer membrane damage, lipopolysaccharide (LPS)
synthesis defects, peptidoglycan perturbation and lipoprotein mislocalisation [48]. It subsequently acts as
a transcription regulator of associated genes, but also of those related to biofilm formation, motility and
virulence [49]. The system regulates the balance between innate and adaptive defence against phages
[50-52], and mutations affecting it can confer partial phage-resistance [53]. Cluster 2 also presented
missense mutations in genes involved in DNA replication (thymidylate kinase [*tmk*]), amino acid
metabolism (shikimate kinase [*aroK*]; 2-aminoadipate transaminase [*lysN*]), and the outer membrane
porin *ompP*, although their contribution to decreased susceptibility to øSimón is uncertain.
388

Finally, we looked for mutations in the post-treatment, phage-selected isolates and found that compared
to the wt *B. bronchialis* strain, there was a median of 27 SNPs per strain (range 4 – 32), with no
correlation between the number of SNPs and susceptibility to øSimón (Fig. S8 and Supplementary data

file). Notably, a bacterial isolate recovered from sputum after 2 days of phage therapy had a total
deletion of *btuB*, the gene for the vitamin B12 receptor that we showed was important for øSimón-
resistance *in vitro*. This isolate was completely resistant to øSimón. Another isolate from day 2 had a
missense mutation in the toxin gene *hipA*, which is part of a known toxin-antitoxin phage defence
system [54]. Additionally, we found that many of the phage-selected strains, regardless of their genetic
background (cluster 1 or 2), also harboured mutations in cell envelope-associated genes, including cell
wall synthesis (*anmK, mrcA, amgK, murB, murU*), LPS biosynthesis (*pgaB, ppgK, waaQ, wecE, gmhB*),
LPS export (*lptD, lptG*), and capsular and extracellular polysaccharide synthesis (*wzb*) (Fig. 6f, Table
S4), all potentially impacting their susceptibility to øSimón to varying degrees and through different
pathways [55-59]. Taken together, these data show that our patient was infected with genetic variants of
B. bronchialis that had a diverse spectrum of susceptibility to øSimón, even before phage treatment
started. After treatment initiation, strains with reduced susceptibility to øSimón persisted, with several
defined mutations, including loss of the putative receptor *btuB*, likely explaining the phenotype.
405
The pre-treatment phage heteroresistance we identified likely contributed to the failure of phage
therapy. However, our findings should be interpreted with nuance. First, the proportion of bacteria with
reduced sensitivity to øSimón did not dramatically increase during phage administration (Fig. 6b),
suggesting these subpopulations did not possess a significant advantage under phage selection *in vivo*.
Second, the isolates obtained during and after phage treatment were distributed across clusters 1 and
2 at a similar ratio as the pre-treatment isolates. This could indicate that the observed genomic changes
reflect natural pathogen variability or in-host evolution, rather than direct selection by phage treatment.
In any case, we argue that testing for heteroresistance prior to phage treatment would be advisable in
future patients.

Figure 6. The patient was infected with a heteroresistant bacterial population. a: Phage-selection experiment. Serial dilutions of sputum were mixed with soft agar and either PBS (top) or øSimón (bottom), poured over hard agar plates and incubated. *B. bronchialis* colonies were identified through morphology and MALDI-ToF MS and counted. **b:** The ratio between non-phage-selected (PBS, top in panel a) and phage-selected (øSimón, bottom in panel a) was calculated for each sputum sample timepoint. Dotted line represents the median false positive threshold for the assay calculated on a clonal wt *B. bronchialis* culture (n = 3). Values are median ± 95% CI; n = 3; Kruskal-Wallis test, only p values < 0.05 plotted. **c:** Efficiency of plating (EOP) of øSimón on 14 random colonies from the pre-treatment sputum sample. All replicates plotted; bars are medians. EOPs > 1 in yellow, EOPs < 1 in blue. **d:** Heat map of the sensitivity/resistance spectrum to øSimón of sputum *B. bronchialis*. EOPs (n = 3; median plotted) of 44 phage-selected colonies from 6 different days during and after phage therapy. On the right, representative extremes of plaque size and clarity of øSimón on these strains. The white and black dots at the top right corner of each picture correspond to the strains used in the assay (marked with the same dots in the EOP panel). Scale bar = 1 cm. **e:** Maximum-likelihood tree of the 59 *B. bronchialis* colonies that were sequenced for this study. The tree depicts two clear clusters (100% node support values). **f:** SNP distance matrix between pre-treatment (columns), post-treatment (rows) and wt *B. bronchialis* (upper left corner). Columns ordered left-to-right in descending order of mean SNP distance to post-treatment isolates. Rows ordered by day of treatment and, within days, by descending EOP. The additional vertical dashed line in white separates pre-treatment isolates with EOPs higher and lower than 1, labelled respectively. Gene names in each cell represent mutations in a specific isolate from the rows, discovered after comparison against the corresponding isolate from the columns. The five genes under the heat map are mutated in all isolates of cluster 2, intact in cluster 1. Underlined genes: nonsense mutations or complete deletions; genes in bold: gain-of-function mutations; all the rest: missense mutations. All mutations are further detailed in Table S4 and the Supplementary Data File.

Clinical implications

The resurgence of phage therapy in the context of challenging AMR bacterial infections has highlighted its many remaining unknowns. Here, we provide a detailed analysis of a clinical case of phage therapeutic failure and uncovered important insights with clinical implications for the future care of phage therapy patients. Using longitudinal clinical samples, we first identified the complex dynamics that are taking place between phages, bacteria and human immune responses. When administered intravenously, phage was able to reach the patient's sputum, however phage activity was short-lived. Two main factors appeared to be at play: pre-existing antiphage immunity and bacterial heteroresistance to phage. Importantly, we have shown that in the setting of a chronic bacterial infection, bacterial prophages could act as host immune stimulators, leading to pre-treatment cross-reactive antibodies that can target the therapeutic phage. This pre-treatment phage neutralisation capacity, combined with subpopulations of infecting bacteria with reduced susceptibility to the phage, create a recipe for clinical failure that has been under-appreciated. Arguably, these barriers could have potentially been overcome using a phage cocktail instead of a single-phage approach, a strategy we could not use despite a sustained effort to isolate additional phages against the patient's pathogen.

We recommend screening a patient's serum for antiphage immunity (serum neutralisation) well before undertaking clinical production of a specific phage. The current Phage Australia clinical protocol suggests testing for antiphage antibodies during and after phage therapy [11]. We suggest that, in the

presence of alternative phages suitable for therapy, those that are susceptible to pre-existing immunity
should be considered lower priority. In the absence of an alternative phage for a patient requiring urgent
treatment, however, pre-existing immunity should not be considered a contraindication. Second, we
recommend choosing a phage or combination of phages for clinical use by testing them against a
patient's bacterial population instead of a single colony. Homogenous colony morphology or growth
kinetics should not be used as evidence to rule out possible bacterial heteroresistance. Instead, the
inverted top agar assay described in this study could be used for that purpose in a cost-effective manner.
An increasing amount of evidence recognises that, especially in chronic infections, patients can harbour
heterogenous populations of a pathogen [60-62], and a single colony might not be representative of the
population's susceptibility to phage therapy. These clinical recommendations are easily applicable to all
prospective phage therapy patients but are particularly relevant in patients with recurrent, chronic or
multisite infections. Likewise, it will be important for those planning clinical trials to consider these
variables in trial design and interpretation, including the collection of appropriate samples throughout
the study. We hope that as global efforts to bring phage therapy into widespread clinical practice
continue, careful analysis of clinical cases and results from well-designed clinical trials will keep
providing useful and actionable insights to maximise the effectiveness of this promising antimicrobial
strategy.
475

Materials and Methods
Bacterial growth, quantification, and storage. *B. bronchialis* was grown on heart infusion (HI) broth
(Oxoid, CM1032), supplemented with 1.5% agar, when necessary, aerobically, with constant shaking
for liquid cultures, at 37°C for 48 h. Bacterial quantification was performed using the spread-plating
technique, with at least two successive 10-fold serial dilutions of the relevant sample yielding countable
colonies. When needed, bacterial cultures were diluted 2:3 in 80% glycerol for storage at -80°C.
482

Phage isolation, purification, quantification and efficiency of plating (EOP). Phage øSimón was
isolated from raw sewage samples (Melbourne, VIC, Australia, May 2022). Aliquots of 100 ml of
centrifuged and filtered sewage were combined with 1 ml of a turbid *B. bronchialis* liquid culture and
supplemented with 10× HI, and 10 mM each of $CaCl_2$ and $MgSO_4$. Mixtures were incubated as above,
the resulting lysates were purified with the Phage-on-Tap protocol [63], with or without buffer replacement,
for general lab stocks. Phage suspensions needed for Immunology work underwent further purification
as performed in [22]. All phage products were stored at 4 °C. Phage quantification and EOP testing were
performed with a slightly modified double overlay agar assay [64] using 4 ml of molten soft (0.4%) HI agar
and 1 ml of a *B. bronchialis* culture at an optical density (OD_{600}) of 0.3 (previously tested to correspond
to a bacterial density of 8×10^8 cfu/ml), per plate. Plates were always read after 48 h of incubation, and
the average number of plaques in at least two plates was recorded and compared as a percentage of
the plaques formed on the original host of isolation under the same conditions.
495

Bacterial and phage kinetics. Growth kinetics were determined using a 96-well plate format and
spectrophotometry (BioTek Epoch 2), measuring OD_{600} every 15 min with a starting inoculum of 10^6

cfu/well, and phage pfu at multiplicities of infection (MOI) of either 0, 0.01, 0.1, or 1. For experiments

comparing bactericidal activity in liquid and sputum, cocultures were set up in 10 ml volumes, at a

starting bacterial density of 10^7 cfu/ml and MOIs of either 0, 0.01, or 0.1. Artificial sputum medium was

commercially acquired (Biochemazone, Canada, BZ316), handled, stored and reconstituted according

to the manufacturer's instructions.

Phage one-step kill curve and adsorption assay. Bacteria from liquid cultures and phage from a

purified stock were mixed at an MOI of 0.01 (10^8 cfu/ml) in HI broth, with cation supplementation as

above. The assay was run in 10 ml total volumes. The suspensions were incubated, and at each

timepoint, 1 ml samples were taken and filtered with a 0.2 μm syringe filter (Pall, 4612). This filtrate was

serially diluted in PBS and plated in duplicate for quantification of free phage particles. A control with no

bacteria was routinely run in adsorption assays.

Transmission electron microscopy (TEM) and immuno-TEM. Droplets of phage suspension (10 μl)

were placed on copper TEM grids (200 mesh; SPI) with carbon-coated ultrathin formvar film. They were

left for 30 s and then dried using filter paper. For negative staining, the grids (face side down) were put

to droplets (10 μl) of uranyl acetate water solution (1% w/v) and incubated for 30 s. Finally, the grids

were dried with filter paper and examined under a transmission electron microscope (JEM-1400 Plus)

at an accelerating voltage of 80 kV. In the induced prophage suspension, imaging was hindered by

contamination and low virion concentration. For immuno-TEM experiments, the IgG fractions of the

patient's serum samples were first purified using protein G magnetic beads (New England Biolabs,

S1430S) largely according to the manufacturer's instructions. Briefly, after washing beads in binding

buffer (0.1 M sodium carbonate, pH = 8.0) three times, serum diluted 1:3 with binding buffer was added

to beads and then incubated with rolling and tilting for 15 min at room temperature. Beads were placed

on a magnetic stand for 30 s, the depleted serum removed and then the beads washed four times with

1 ml binding buffer. Following the washes, IgG was eluted in 0.2 M glycine pH = 2.5 and transferred to

a tube containing a neutralising volume of 1 M Tris-HCl pH = 9.0. The sample was diluted 1:4 in PBS,

filtered (0.2 μm) and concentrated sequentially through 10 kDa Amicon Ultra Centrifugal filters (Merck

Millipore, UFC8010 and UFC5010). The A280 of the concentrated, purified IgG was measured by

NanoDrop (NanoDrop Technologies LLC) and a 0.1% Ce of 1.36 applied to determine the IgG

concentration (at least 0.2 mg/ml for immuno-TEM). Phages were left to adsorb to the TEM grids for 10

min, followed by three 1-min washes on droplets of 1% bovine serum albumin (BSA) in PBS. Grids were

then incubated for 2 h on a mixture containing 10 μl of 1% BSA in PBS and 5 μl of patient-derived IgG.

After three additional 1-min washes, grids were incubated for 1 h on a mixture of 10 μl of 1% BSA in

PBS and 5 μl of gold-conjugated goat-anti human IgG antibodies (RRID: AB_2337747) (diluted 1:10 in

PBS). This was followed by three washes with PBS, and standard negative staining and examination.

Phage DNA extraction, sequencing and bioinformatics. One ml volumes of high-titre phage lysate

(> 10^8 pfu/ml whenever possible) were treated with DNase (1 mg/ml) and RNase A (12.5 mg/ml) for 2 h

at 37 °C, followed by an inactivation step of 5 min at 75 °C. We then used the Norgen phage DNA kit

(Norgen Biotek Corp., Thorold, ON, Canada) to extract DNA following the manufacturer's instructions.

DNA concentration was measured using a Qubit® Fluorometer, quality confirmed with the 260/280 nm

absorbance ratio, and absence of bacterial contamination was confirmed by 16S rRNA PCR and gel

visualisation of the DNA samples. Phage øSimón was sequenced on an Illumina HiSeq platform, and

with Oxford Nanopore Technologies (ONT) long-read sequencing with a GridION device on FLO-

MIN114 R10.4.1 flow cells. The reads were trimmed using trimmomatic (v0.39) [65], assembled with

Unicycler (v0.5.0) [66], and annotated using Prokka (v.1.14.6), the RAST server and the PHROG

database [67-69]. The genome was set to start at the large terminase subunit (*terL*) gene [70], polished with

Pilon (v.1.22) [71]. manually screened for lysogeny indicators and assessed for AMR determinants using

AMRFinderPlus (v.3.11.26) [72] and Abricate (v.1.0.1) (https://github.com/tseemann/abricate). Phylogeny

was investigated using vContact2 [73] and the gene voting approach [74], both using the Inphared database

(version 1Dec2021) [75]. Gene similarity between øSimón and *B. bronchialis* prophages was studied

using blastp (v.2.15.0+) and genome maps were visualised using the R package *gggenomes* (v.1.0.0).

Production of clinical-grade phage øSimón. We followed a slightly modified version of a previously

used protocol [76]. The phage was amplified in 1,000 ml HI broth. The lysate was centrifugated for 30 min

at 4,000 × g to remove larger bacterial debris, and then filtered through a depth filter from 5 μm to 0.2

μm, monitoring the pressure not to exceed and compromise the filter specifications (<2 bar). The lysate

was passed once again through a sterilisation-grade (0.2 μm) filter and transferred to a bacteria-free

production facility. The lysate was diluted 1:10 in PBS and flushed through an ÄKTA Flux tangential flow

filtration (TFF) system (Cytiva) equipped with a 300 kDa microfiltration hollow fibre cartridge. This

process was repeated until the HI broth was completely replaced by PBS. The lysate was filter-sterilised

again, and the phage titre measured. Endotoxin removal was performed using EndoTrap®HD affinity

columns, and the endotoxin concentration was assessed with Endosafe® PTS™ cartridge (Charles

River laboratories, MA, USA) kits. After achieving a satisfactory endotoxin level, we processed the

samples under strict sterile conditions, packed them in sealed glass vials with syringe access (Medisca,

9497-01), and sent them to Eurofins BioPharma Product Testing in Sydney, Australia for third-party

validation for sterility (USP71) and endotoxin (USP85).

Handling and storage of clinical specimens. Serum samples were aliquoted into 200 μl volumes and

kept at -20 °C, minimising freeze/thaw cycles. For sputum samples, we first measured their volume and

mixed them 1:1 with a 1:10 solution of dithiothreitol (Sputolysin®, CAS 578517) in sterile water with

vortexing for 1 min. Bacterial and phage quantification were performed at this stage. For long-term

storage for other experiments, each sample was mixed 1:1 with sterile glycerol broth (60 ml glycerol, 2

g tryptone, 1 g yeast extract, 2 g NaCl, in 200 ml deionised water) in cryovials and stored at -80 °C.

Serum and sputum neutralisation assays and calculation of the phage inactivation rate (*k*). The

clinical samples were diluted 1:10 in PBS, filter-sterilised (0.2 μm syringe filter), and then further diluted

in PBS in 10-fold serial dilutions. Working volumes of 1 ml of these dilutions, and a PBS-only control,

were spiked with 10^6 pfu each of phages øSimón and øFaith (unpublished phage, isolated in our lab

against *Pseudomonas aeruginosa* strains from patients with cystic fibrosis). The samples were
incubated at 37 °C for 30 min with constant shaking. At the endpoint, phage quantification using the
double overlay agar assay was performed for each phage on their respective hosts of isolation.
Neutralisation was calculated as the percentage of active phage at the endpoint compared to the initial
phage inoculum for each condition. The phage inactivation rate (*k*) was calculated with the formula:

583
$$k = 2.3 \times \frac{D}{T} \times log \frac{P0}{Pt}$$

Where D is the reciprocal of the serum or sputum dilution in PBS, T is the time of incubation, and P0
and Pt are the phage titres at the beginning and endpoint of the assay, respectively [17,19].

Indirect ELISA and cross-reactive ELISA. We based our protocol in [18] with some modifications.
Individual wells of MICROLON™ high binding 96-well plates (Greiner, 655061) were coated with 100 μl
of either pure coating buffer (carbonate-bicarbonate with pH = 9.6, Sigma C3041), phage øSimón
calculated at 10^5 pfu/well, induced prophage at 10^3 virions/well, or phage vehicle (obtained after filtering
phage solutions with a 0.02 μm syringe filter, Whatman 68091002); the latter three diluted in coating
buffer. Plates were covered with foil and incubated at 4 °C for 20 h. The wells were emptied, washed 4
times with 300 μl of PBS + 0.05% Tween 80 (PBST), blocked with 200 μl of 3% skim milk powder in
PBS, and the plates incubated as above. The blocking solution was shaken off, and the plates patted
dry with paper towels. A further blocking step was performed with 3% bovine serum albumin (BSA) in
PBS at 37 °C for 30 min, before washing with PBST. The patient samples (serum or filter-sterilised
sputum) were diluted in 3% BSA in PBS, added to the respective wells in volumes of 50 μl, incubated
at 37 °C for 1 h, and washed with PBST. Next, volumes of 50 μl of 1:500 goat anti-human IgA + IgG +
IgM-AP (alkaline phosphatase)-conjugated antibody (RRID: AB_2337605) were added and the plates
again incubated and later washed with PBST. Finally, the p-Nitrophenyl phosphate substrate
(Sigmafast™, N2770) was added in a volume of 50 μl and the reaction left to develop for 20 min. The
assay was stopped by adding 50 μl of 2 M NaOH and the plates were read with a spectrophotometer
at 450 nm. For the antibody cross-reactivity testing the same protocol was followed except for a "pre-
capture" step involving the passage of individual serum samples by wells covered with øSimón,
prophage, or phage vehicle in coating buffer, for 1 h before transferring them into new plates and wells
for a regular ELISA as previously described.

Prophage induction. A stationary phase *B. bronchialis* culture was diluted 1:10 in HI and left for
incubation for 6 h. The culture was split into two 10 ml aliquots, the first one received a PBS spike
(spontaneous induction control), and the second one a mitomycin C at 0.3 mg/ml spike, in equal
volumes. Both cultures were left for overnight incubation and purified as above [22]. Prophage induction
was confirmed through TEM and phage DNA sequencing.

Inverted soft agar assay for phage-selection. Sputum samples were 10-fold serially diluted in PBS.
Then, 100 μl of the dilutions were mixed with 4 ml of molten soft (0.4%) HI agar and 100 μl of either
PBS (no phage selection) or a stock of øSimón for a concentration of 10^7 pfu/plate (phage selection),

before being poured over plates containing a first layer of standard HI agar. Plates were left for 48 h of
incubation and, for counting, *B. bronchialis* colonies on each plate were identified through visual
assessment of their morphology, followed by species confirmation using MALDI-ToF MS (Bruker,
Biotyper microflex®). Normal microbiota from the upper respiratory tract was consistently seen at
concentrations 2 log lower than *B. bronchialis* and thus did not represent a significant confounder in this
process. Phage-selected and non-phage-selected colonies underwent two rounds of single-colony
purification before being grown in liquid culture and stored as above.

Bacterial DNA extraction and sequencing. Genomic DNA was extracted from *B. bronchialis* cultures
using the GenFind V3 Reagent Kit (Beckman Coulter) as per manufacturer's instructions. Illumina
libraries were prepared using the Illumina DNA Library Prep Kit, (M) Tagmentaton (Illumina) and
sequenced on the Illumina NovaSeq 6000 platform to produce paired-end reads. The wt *B. bronchialis*
isolate also underwent long-read Nanopore sequencing with a GridION device on FLO-MIN114 R10.4.1
flow cells (Oxford Nanopore Technologies). Sequencing libraries were barcoded with the native
barcoding kit SQK-RBK114.96 in accordance with the protocol RBK_9176_v114_revM_27Nov2022.

Bacterial genomics. A *de novo* assembly of the genome of wt *B. bronchialis* was first constructed
using Trycycler (v0.5.4) with long read data [77], polished with matching short reads using Polypolish
(v0.5.0) [78] and annotated with Bakta (v1.7) [79]. The short-read data from the other isolates also
underwent quality control (https://github.com/s-andrews/FastQC) before the genomes were assembled
with Unicycler (v0.5.0) [66] and annotated with Prokka (v.1.14.6) [67]. We identified SNPs between the wt
B. bronchialis reference genome and other sputum isolates using the Nextflow implementation of the
Reddog pipeline v1beta.11 using default parameters (https://github.com/scwatts/reddog-nf). The
resulting SNP alignment consisted of 250 sites. Complementarily, the wt genome was manually parsed
for insertion sequences, whose nucleotide sequences were then mapped against the short reads from
other isolates using ISMapper v2.0.0 [80], to detect alternative gene disruptions in these isolates. To
identify the absence of whole genes in pairwise comparisons between isolates we used panaroo
(v1.2.4) [81] using thresholds of ≥95% for gene coverage and ≥5x for depth. Phylogeny was studied by
constructing a maximum-likelihood tree from the SNP alignment obtained above, with the GTR+Γ
model, using RAxML (v8.2.9) [82] with 1,000 bootstrap replicates, then visualised with "ape" (v5.7.1) [83]
and "ggtree" (v3.6.2) [84] in R. The pairwise SNP distance matrix was created using "harrietr"
(https://github.com/andersgs/harrietr).

Graphing, statistics, and reproducibility. Graphing and statistical analyses were performed with
GraphPad Prism 9 (GraphPad Software, Inc.). Figures 4d and 6a were crafted using BioRender. All *in*
vitro experiments were performed in triplicate, with at least two technical replicates each. Where
appropriate, due to the small number of replicates, medians were presented. To decide between the
use of parametric or nonparametric statistical analyses, the data were tested for normality using the
Shapiro-Wilk method. The threshold value of two-tailed $p < 0.05$ was used for statistical significance.

Observational results, including those from TEM, and phage plaque morphology are representative of
at least 100 observations each.

Data availability. All bacterial and phage genomes, sequencing data, and metadata derived from this
study have been deposited on NCBI under the BioProject number PRJNA1180749 (Table S5).

Ethical statement. All procedures performed in this study were in accordance with the ethical
standards of the institutional research committee (Alfred Hospital Ethics Committee approval number
230/24) and with the Helsinki Declaration (as revised in 2013). Written informed consent to receive
phage therapy was obtained from the patient. Data from this clinical case forms part of the STAMP
Study[11], trial registration number (registered on ANZCTR 10 Nov 2021): ACTRN12621001526864;
WHO Universal Trial Number: U1111-1269-6000.

Acknowledgements
We thank Dr Ryan Burrows (University of Melbourne & Melbourne Water) for his help establishing the
collaboration that granted us access to the raw sewage samples used to isolate øSimón. Thanks to
Christina Rootes, Dr Christopher Pace, and Dr Duen Siauw for useful feedback on the structure and
readability of the manuscript, and to Ruzeen Patwa for her help in logistics and laboratory management.
Thanks to Simón Usúcar Gordillo, after whom phage øSimón was named (*que tu vida esté llena de
éxito y alegría, tu tío te ama*). The Monash Antibody Discovery Platform is supported by funding from
Bioplatforms Australia (enabled by NCRIS). This work was supported by Monash eResearch
capabilities, including M3, and the Alfred Hospital and Monash University funding of the Victorian
Bacteriophage Therapy Program (VIC*Phage*) and the Monash Phage Foundry. Jeremy J. Barr and
Anton Y. Peleg also acknowledge the Australian National Health and Medical Research Council
(NHMRC) Investigator L1 Grant (2023/GNT2026130) and Practitioner Fellowship (APP1117940),
respectively, and the Frontier Health Medical Research support (RFRHPI000017).

Author contributions
Conceptualisation: FGA, JJB, AYP.
Laboratory experimental work: FGA, DS, MBu, DMP, MP, DKo, MJR, KP, JW, HR.
Genomics and bioinformatics: DS, MBe, MBu, SD, JH.
Clinical monitoring and follow up: FGA, SK, BJG, YH, DKe, TK, AYP.
Data curation, analysis and visualisation: FGA, AYP.
Provision of resources: CR, JJB, AYP.
Original preparation of the manuscript: FGA, AYP.
All authors reviewed, edited and approved the manuscript.

Competing interests
None to declare.

References

1.   Collaborators, G.B.D.A.R. Global burden of bacterial antimicrobial resistance 1990-
2021: a systematic analysis with forecasts to 2050. *Lancet* **404**, 1199-1226 (2024).
2.   Dadgostar, P. Antimicrobial Resistance: Implications and Costs. *Infect Drug Resist* **12**,
3903-3910 (2019).
3.   Gordillo Altamirano, F.L. & Barr, J.J. Phage Therapy in the Postantibiotic Era. *Clin*
Microbiol Rev **32**(2019).
4.   Gomez-Ochoa, S.A.*, et al.* Efficacy of phage therapy in preclinical models of bacterial
infection: a systematic review and meta-analysis. *Lancet Microbe* **3**, e956-e968
(2022).
5.   Uyttebroek, S.*, et al.* Safety and efficacy of phage therapy in difficult-to-treat
infections: a systematic review. *Lancet Infect Dis* **22**, e208-e220 (2022).
6.   Marchi, J., Zborowsky, S., Debarbieux, L. & Weitz, J.S. The dynamic interplay of
bacteriophage, bacteria and the mammalian host during phage therapy. *iScience* **26**,
106004 (2023).
7.   Vandamme, P.A.*, et al.* Bordetella bronchialis sp. nov., Bordetella flabilis sp. nov. and
Bordetella sputigena sp. nov., isolated from human respiratory specimens, and
reclassification of Achromobacter sediminum Zhang et al. 2014 as Verticia
sediminum gen. nov., comb. nov. *Int J Syst Evol Microbiol* **65**, 3674-3682 (2015).
8.   Spilker, T., Darrah, R. & LiPuma, J.J. Complete Genome Sequences of Bordetella
flabilis, Bordetella bronchialis, and "Bordetella pseudohinzii". *Genome Announc*
4(2016).
9.   Bridel, S.*, et al.* A comprehensive resource for Bordetella genomic epidemiology and
biodiversity studies. *Nat Commun* **13**, 3807 (2022).
10.   Suh, G.A.*, et al.* Considerations for the Use of Phage Therapy in Clinical Practice.
Antimicrob Agents Chemother **66**, e0207121 (2022).
11.   Khatami, A.*, et al.* Standardised treatment and monitoring protocol to assess safety
and tolerability of bacteriophage therapy for adult and paediatric patients (STAMP
study): protocol for an open-label, single-arm trial. *BMJ Open* **12**, e065401 (2022).
12.   Gembara, K. & Dabrowska, K. Phage-specific antibodies. *Curr Opin Biotechnol* **68**,
186-192 (2021).
13.   Jerne, N.K. & Avegno, P. The development of the phage-inactivating properties of
serum during the course of specific immunization of an animal: reversible and
irreversible inactivation. *J Immunol* **76**, 200-208 (1956).
14.   Hodyra-Stefaniak, K.*, et al.* Natural and Induced Antibodies Against Phages in
Humans: Induction Kinetics and Immunogenicity for Structural Proteins of PB1-
Related Phages. *Phage (New Rochelle)* **1**, 91-99 (2020).
15.   Kazmierczak, Z.*, et al.* Immune Response to Therapeutic Staphylococcal
Bacteriophages in Mammals: Kinetics of Induction, Immunogenic Structural Proteins,
Natural and Induced Antibodies. *Front Immunol* **12**, 639570 (2021).
16.   Lusiak-Szelachowska, M., Miedzybrodzki, R., Fortuna, W., Borysowski, J. & Gorski, A.
Anti-phage serum antibody responses and the outcome of phage therapy. *Folia*
Microbiol (Praha) **66**, 127-131 (2021).
17.   Lusiak-Szelachowska, M.*, et al.* Phage neutralization by sera of patients receiving
phage therapy. *Viral Immunol* **27**, 295-304 (2014).

18.   Dedrick, R.M.*, et al.* Phage Therapy of Mycobacterium Infections: Compassionate Use
742   of Phages in 20 Patients With Drug-Resistant Mycobacterial Disease. *Clin Infect Dis*
743   **76**, 103-112 (2023).

744   19.   Lusiak-Szelachowska, M.*, et al.* Antiphage activity of sera during phage therapy in
745   relation to its outcome. *Future Microbiol* **12**, 109-117 (2017).

746   20.   Van Belleghem, J.D., Dabrowska, K., Vaneechoutte, M., Barr, J.J. & Bollyky, P.L.
747   Interactions between Bacteriophage, Bacteria, and the Mammalian Immune System.
748   *Viruses* **11**(2018).

749   21.   Popescu, M., Van Belleghem, J.D., Khosravi, A. & Bollyky, P.L. Bacteriophages and the
750   Immune System. *Annu Rev Virol* **8**, 415-435 (2021).

751   22.   Bichet, M.C., Patwa, R. & Barr, J.J. Protocols for studying bacteriophage interactions
752   with in vitro epithelial cell layers. *STAR Protoc* **2**, 100697 (2021).

753   23.   Doria-Rose, N.A. & Joyce, M.G. Strategies to guide the antibody affinity maturation
754   process. *Curr Opin Virol* **11**, 137-147 (2015).

755   24.   Dion, M.B., Oechslin, F. & Moineau, S. Phage diversity, genomics and phylogeny. *Nat*
756   *Rev Microbiol* **18**, 125-138 (2020).

757   25.   Haddock, N.L., Barkal, L.J. & Bollyky, P.L. Bacteriophage populations mirror those of
758   bacterial pathogens at sites of infection. *mSystems* **8**, e0049723 (2023).

759   26.   Haddock, N.L.*, et al.* Phage diversity in cell-free DNA identifies bacterial pathogens in
760   human sepsis cases. *Nat Microbiol* **8**, 1495-1507 (2023).

761   27.   Nayfach, S.*, et al.* CheckV assesses the quality and completeness of metagenome-
762   assembled viral genomes. *Nat Biotechnol* **39**, 578-585 (2021).

763   28.   Halperin, S.A., Ferrieri, P., Gray, E.D., Kaplan, E.L. & Wannamaker, L.W. Antibody
764   response to bacteriophage hyaluronidase in acute glomerulonephritis after group A
765   streptococcal infection. *J Infect Dis* **155**, 253-261 (1987).

766   29.   Fluckiger, A.*, et al.* Cross-reactivity between tumor MHC class I-restricted antigens
767   and an enterococcal bacteriophage. *Science* **369**, 936-942 (2020).

768   30.   Berkson, J.D.*, et al.* Phage-specific immunity impairs efficacy of bacteriophage
769   targeting Vancomycin Resistant Enterococcus in a murine model. *Nat Commun* **15**,
770   2993 (2024).

771   31.   Zborowsky, S.*, et al.* Macrophage-induced reduction of bacteriophage density limits
772   the efficacy of in vivo pulmonary phage therapy. *Nat Commun* **16**, 5725 (2025).

773   32.   Roach, D.R.*, et al.* Synergy between the Host Immune System and Bacteriophage Is
774   Essential for Successful Phage Therapy against an Acute Respiratory Pathogen. *Cell*
775   *Host Microbe* **22**, 38-47 e34 (2017).

776   33.   Egido, J.E., Costa, A.R., Aparicio-Maldonado, C., Haas, P.J. & Brouns, S.J.J.
777   Mechanisms and clinical importance of bacteriophage resistance. *FEMS Microbiol*
778   *Rev* **46**(2022).

779   34.   Labrie, S.J., Samson, J.E. & Moineau, S. Bacteriophage resistance mechanisms. *Nat*
780   *Rev Microbiol* **8**, 317-327 (2010).

781   35.   Lebeaux, D.*, et al.* A Case of Phage Therapy against Pandrug-Resistant
782   Achromobacter xylosoxidans in a 12-Year-Old Lung-Transplanted Cystic Fibrosis
783   Patient. *Viruses* **13**(2021).

784   36.   Blasco, L.*, et al.* Case report: Analysis of phage therapy failure in a patient with a
785   Pseudomonas aeruginosa prosthetic vascular graft infection. *Front Med (Lausanne)*
786   **10**, 1199657 (2023).

37. Jault, P.*, et al.* Efficacy and tolerability of a cocktail of bacteriophages to treat burn
wounds infected by Pseudomonas aeruginosa (PhagoBurn): a randomised,
controlled, double-blind phase 1/2 trial. *Lancet Infect Dis* **19**, 35-45 (2019).
38. Schooley, R.T.*, et al.* Development and Use of Personalized Bacteriophage-Based
Therapeutic Cocktails To Treat a Patient with a Disseminated Resistant Acinetobacter
baumannii Infection. *Antimicrob Agents Chemother* **61**(2017).
39. Gordillo Altamirano, F.L.*, et al.* Phage-antibiotic combination is a superior treatment
against Acinetobacter baumannii in a preclinical study. *EBioMedicine* **80**, 104045
(2022).
40. Zhvania, P., Hoyle, N.S., Nadareishvili, L., Nizharadze, D. & Kutateladze, M. Phage
Therapy in a 16-Year-Old Boy with Netherton Syndrome. *Front Med (Lausanne)* **4**, 94
(2017).
41. Bertozzi Silva, J., Storms, Z. & Sauvageau, D. Host receptors for bacteriophage
adsorption. *FEMS Microbiol Lett* **363**(2016).
42. Happonen, L.J., Pajunen, M.I., Jun, J.W. & Skurnik, M. BtuB-Dependent Infection of
the T5-like Yersinia Phage varphiR2-01. *Viruses* **13**(2021).
43. Hong, J.*, et al.* Identification of host receptor and receptor-binding module of a newly
sequenced T5-like phage EPS7. *FEMS Microbiol Lett* **289**, 202-209 (2008).
44. Golomidova, A.K.*, et al.* Branched Lateral Tail Fiber Organization in T5-Like
Bacteriophages DT57C and DT571/2 is Revealed by Genetic and Functional Analysis.
Viruses **8**(2016).
45. Gordillo Altamirano, F.L. & Barr, J.J. Unlocking the next generation of phage therapy:
the key is in the receptors. *Curr Opin Biotechnol* **68**, 115-123 (2021).
46. Gillis, A. & Mahillon, J. An improved method for rapid generation and screening of
Bacillus thuringiensis phage-resistant mutants. *J Microbiol Methods* **106**, 101-103
(2014).
47. Andersson, D.I., Nicoloff, H. & Hjort, K. Mechanisms and clinical relevance of bacterial
heteroresistance. *Nat Rev Microbiol* **17**, 479-496 (2019).
48. Meng, J., Young, G. & Chen, J. The Rcs System in Enterobacteriaceae: Envelope Stress
Responses and Virulence Regulation. *Front Microbiol* **12**, 627104 (2021).
49. Wall, E., Majdalani, N. & Gottesman, S. The Complex Rcs Regulatory Cascade. *Annu
Rev Microbiol* **72**, 111-139 (2018).
50. Chaudhry, W.*, et al.* Mucoidy, a general mechanism for maintaining lytic phage in
populations of bacteria. *FEMS Microbiol Ecol* **96**(2020).
51. Mutalik, V.K.*, et al.* High-throughput mapping of the phage resistance landscape in E.
coli. *PLoS Biol* **18**, e3000877 (2020).
52. Smith, L.M.*, et al.* The Rcs stress response inversely controls surface and CRISPR-Cas
adaptive immunity to discriminate plasmids and phages. *Nat Microbiol* **6**, 162-172
(2021).
53. Burmeister, A.R., Tewatia, H. & Skinner, C. A tradeoff between bacteriophage
resistance and bacterial motility is mediated by the Rcs phosphorelay in Escherichia
coli. *Microbiology (Reading)* **170**(2024).
54. LeRoux, M. & Laub, M.T. Toxin-Antitoxin Systems as Phage Defense Elements. *Annu
Rev Microbiol* **76**, 21-43 (2022).
55. Kortright, K.E., Chan, B.K. & Turner, P.E. High-throughput discovery of phage
receptors using transposon insertion sequencing of bacteria. *Proc Natl Acad Sci U S A*
117, 18670-18679 (2020).

56. Acton, L.*, et al.* Collateral sensitivity increases the efficacy of a rationally designed
bacteriophage combination to control Salmonella enterica. *J Virol*, e0147623 (2024).

57. Bai, J., Raustad, N., Denoncourt, J., van Opijnen, T. & Geisinger, E. Genome-wide
phage susceptibility analysis in Acinetobacter baumannii reveals capsule modulation
strategies that determine phage infectivity. *PLoS Pathog* **19**, e1010928 (2023).

58. Nyenhuis, D.A., Nilaweera, T.D. & Cafiso, D.S. Native Cell Environment Constrains
Loop Structure in the Escherichia coli Cobalamin Transporter BtuB. *Biophys J* **119**,
1550-1557 (2020).

59. Shearer, J., Jefferies, D. & Khalid, S. Outer Membrane Proteins OmpA, FhuA, OmpF,
EstA, BtuB, and OmpX Have Unique Lipopolysaccharide Fingerprints. *J Chem Theory*
Comput **15**, 2608-2619 (2019).

60. Dewachter, L., Fauvart, M. & Michiels, J. Bacterial Heterogeneity and Antibiotic
Survival: Understanding and Combatting Persistence and Heteroresistance. *Mol Cell*
76, 255-267 (2019).

61. Klein, S.*, et al.* Comparative Genomic Reveals Clonal Heterogeneity in Persistent
Staphylococcus aureus Infection. *Front Cell Infect Microbiol* **12**, 817841 (2022).

62. Bartell, J.A.*, et al.* Evolutionary highways to persistent bacterial infection. *Nat*
Commun **10**, 629 (2019).

63. Bonilla, N.*, et al.* Phage on tap-a quick and efficient protocol for the preparation of
bacteriophage laboratory stocks. *PeerJ* **4**, e2261 (2016).

64. Stachurska, X., Roszak, M., Jablonska, J., Mizielinska, M. & Nawrotek, P. Double-Layer
Agar (DLA) Modifications for the First Step of the Phage-Antibiotic Synergy (PAS)
Identification. *Antibiotics (Basel)* **10**(2021).

65. Bolger, A.M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**, 2114-2120 (2014).

66. Wick, R.R., Judd, L.M., Gorrie, C.L. & Holt, K.E. Unicycler: Resolving bacterial genome
assemblies from short and long sequencing reads. *PLoS Comput Biol* **13**, e1005595
(2017).

67. Seemann, T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* **30**, 2068-
2069 (2014).

68. Aziz, R.K.*, et al.* The RAST Server: rapid annotations using subsystems technology.
BMC Genomics **9**, 75 (2008).

69. Terzian, P.*, et al.* PHROG: families of prokaryotic virus proteins clustered using remote
homology. *NAR Genom Bioinform* **3**, lqab067 (2021).

70. Shen, W., Le, S., Li, Y. & Hu, F. SeqKit: A Cross-Platform and Ultrafast Toolkit for
FASTA/Q File Manipulation. *PLoS One* **11**, e0163962 (2016).

71. Walker, B.J.*, et al.* Pilon: an integrated tool for comprehensive microbial variant
detection and genome assembly improvement. *PLoS One* **9**, e112963 (2014).

72. Feldgarden, M.*, et al.* AMRFinderPlus and the Reference Gene Catalog facilitate
examination of the genomic links among antimicrobial resistance, stress response,
and virulence. *Sci Rep* **11**, 12728 (2021).

73. Bin Jang, H.*, et al.* Taxonomic assignment of uncultivated prokaryotic virus genomes
is enabled by gene-sharing networks. *Nat Biotechnol* **37**, 632-639 (2019).

74. Nayfach, S.*, et al.* Metagenomic compendium of 189,680 DNA viruses from the
human gut microbiome. *Nat Microbiol* **6**, 960-970 (2021).

75. Cook, R.*, et al.* INfrastructure for a PHAge REference Database: Identification of
Large-Scale Biases in the Current Collection of Cultured Phage Genomes. *Phage (New*
Rochelle) **2**, 214-223 (2021).
76. Subedi, D.*, et al.* Rational Design of Frontline Institutional Phage Cocktail for the
Treatment of Nosocomial Enterobacter cloacae Complex Infections.
bioRxiv, 2024.2006.2030.601436 (2024).
77. Wick, R.R.*, et al.* Trycycler: consensus long-read assemblies for bacterial genomes.
Genome Biol **22**, 266 (2021).
78. Wick, R.R. & Holt, K.E. Polypolish: Short-read polishing of long-read bacterial genome
assemblies. *PLoS Comput Biol* **18**, e1009802 (2022).
79. Schwengers, O.*, et al.* Bakta: rapid and standardized annotation of bacterial genomes
via alignment-free sequence identification. *Microb Genom* **7**(2021).
80. Hawkey, J.*, et al.* ISMapper: identifying transposase insertion sites in bacterial
genomes from short read sequence data. *BMC Genomics* **16**, 667 (2015).
81. Tonkin-Hill, G.*, et al.* Producing polished prokaryotic pangenomes with the Panaroo
pipeline. *Genome Biol* **21**, 180 (2020).
82. Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. *Bioinformatics* **30**, 1312-1313 (2014).
83. Paradis, E. & Schliep, K. ape 5.0: an environment for modern phylogenetics and
evolutionary analyses in R. *Bioinformatics* **35**, 526-528 (2019).
84. Yu, G. Using ggtree to Visualize Data on Tree-Like Structures. *Curr Protoc*
Bioinformatics **69**, e96 (2020).
901
902