1

Title: Hypothalamic deiodinase type-3 establishes the period of circannual interval timing in

mammals

Calum Stewart[1], T. Adam Liddle[1], Elisabetta Tolla[1], Jo Edward Lewis[2], Christopher Marshall[3],

Neil P Evans[1], Peter J Morgan[4], Fran JP Ebling[5], Tyler J Stevenson[1]*.

Corresponding Author: Professor Tyler Stevenson

Email: tyler.stevenson@glasgow.ac.uk

Affiliations:

[1] School of Biodiversity, One Health, and Veterinary Medicine, University of Glasgow;

Glasgow, G61 1QH, UK.

[2] Institute of Metabolic Science-Metabolic Research Laboratories & MRC-Metabolic

Diseases Unit, University of Cambridge, Cambridge, UK.

[3] School of Physiology, Pharmacology and Neuroscience, University of Bristol; Bristol, BS8

1TD, UK.

[4] Rowett Institute, University of Aberdeen; Aberdeen, AB25 2ZD, UK.

[5] School of Life Sciences, University of Nottingham; Nottingham, NG7 2UH, UK.

Competing Interest Statement: Authors declare that they have no competing interests.

Abstract

Animals respond to environmental cues to time phenological events, but the intrinsic

mechanism of circannual timing remains elusive. We used transcriptomic sequencing and

frequent sampling of multiple hypothalamic nuclei in Djungarian hamster to examine the

neural and molecular architecture of circannual interval timing. Our study identified three

distinct phases of transcript changes, with deiodinase type-3 (*Dio3*) expression activated

during the early induction phase. Subsequent work demonstrated that targeted mutation of

Dio3 using CRISPR-Cas resulted in a shorter period for circannual interval timing. Hamsters

that are non-responsive to short photoperiod and fail to show any winter adaptations do not

display changes in *Dio3* expression do not show any change in body mass or pelage. Our

work demonstrates that changes in *Dio3* induction is essential for setting the period of

circannual interval timing.

Introduction

Phenology of key life history traits is common across plant (Piao et al., 2019) and animal

kingdoms (Cohen et al., 2018). The annual changes in daylength are the predominant

environmental cue which animals use to time seasonal life history transitions (Liddle et al.,

2022; Pérez et al., 2019; Wood and Loudon, 2014). Plants and animals also exhibit

endogenous circannual timing in the absence of any change in environmental cues. For

example, bird migration (Gwinner and Dittami, 1990), mammalian hibernation (Pengelley

and Fisher, 1957), and reproduction (Woodfill et al., 1994; Lincoln et al., 2006) are all driven

by robust intrinsically generated *circannual clocks*, the cycle of which nearly match a 12-

month period. Some *circannual timers* estimate an interval period (e.g., 6-month) in which

programmed changes in physiology and morphology occur in anticipation of the next

season. Such interval timers are commonly observed as flowering in plants (Duncan et al.,

2015), diapause in insects (Denlinger, 1974) and spring emergence in rodents (Prendergast

et al., 2004). Interval timers are typically characterized by having light-dependent induction,

maintenance and recovery phases.

Annual mammalian life history transitions are typically associated with major changes

in energy demand (Ricklefs, 1991). While the melanocortin system, including neuropeptide y

(*Npy*), agouti-related peptide (*Agrp*), and melanocortin receptors are known to regulate

short-term changes in energy homeostasis (Yeo et al., 2021), the mechanisms and

anatomical structures implicated in long-term, circannual variation in energy rheostasis are

not well characterized. Somatostatin (*Sst*) (Marshall et al., 2024; Petri et al., 2016, 2014),

proopiomelanocortin (POMC) (Bao et al., 2019; Mercer et al., 2000), and VGF nerve growth

factor (Barrett et al., 2005) are known to be strong correlates of seasonal variation in

energetic state. Tanycyte somas are essential for the integration of environmental and

physiological signals required for circannual interval timing. Tanycytes are localized in the

ependymal layer of the third ventricle and are highly sensitive to nutrient state (Bolborea et

al., 2020) and receive photoperiodic signaling derived from thyrotropes in the pars tuberalis

(Hanon et al., 2008; Wood et al., 2020). Previous work has established deiodinase type-2

(*Dio2*) and type-3 (*Dio3*) expression is anatomically localized to tanycytes and coordinate

triiodothyronine-dependent annual transitions in physiological state (Bao et al., 2019; Ebling

and Lewis, 2018; Hanon et al., 2008; Murphy et al., 2012; Petri et al., 2016; Wood et al.,

2020). Here we delineate the molecular architecture of circannual interval timing by the

hypothalamus and pituitary gland and tested the conjecture that *Dio3*, in the Djungarian

hamster (*Phodopus sungorus*), is upregulated during the induction of circannual interval

timing for energy rheostasis, and functions to establish the duration (or period) of the

circannual interval timer.

Results

Male Djungarian hamsters were either kept in a long photoperiod (LP) control condition or

moved from LP to short photoperiod (SP) conditions (Fig.1). Djungarian hamsters remain in

LP phenotype unless exposed to SP. Pelage color, torpor and body mass were used to

monitor the induction, maintenance, and recovery phases of the circannual timer (Fig.1A and

C *SI Appendix* Movies S1 and 2). A full white pelage color was observed between 12-16

weeks after exposure to SP and gradually reversed to the LP agouti color after 28 weeks in

SP. Hamsters engaged in torpor between 12-20 weeks in SP (movie S2). Massive,

programmed changes in energy rheostasis were observed with a 30% reduction in daily food

intake and a 20% decrease in average body mass by 12 weeks SP (Fig.1C and D). Both

food intake and body mass started to reverse to LP conditions after 20 weeks in SP.

Epididymal adipose tissue mass and plasma insulin concentrations paralleled the change in

body mass (Fig. 1D and *SI Appendix* Fig. S1). These reversals in physiological state are

indicative of the recovery phase of the circannual interval timer. GLP-1 did not display any

change in circulating concentrations (Fig. 1E). Plasma glucose concentration jumped sharply

after 16 weeks in SP (Fig. 1E), concurrent with the development of torpor in these animals

(movie S2). The observed change in body mass, food intake, adipose tissue and plasma

insulin versus the lack of change in GLP-1 highlights the physiological distinction between a

programmed rheostatic mechanism characteristic of the circannual interval timing of energy

stability in the former, while the latter are driven by short-term homeostatic mechanisms

(Stevenson, 2024).

We then used Oxford Nanopore transcriptomic sequencing to characterize the molecular

changes associated with phases of the circannual interval timer within multiple individual

hypothalamic nuclei and the pituitary gland (Fig.1F to G; fig. *SI Appendix* S2 to 6 and

Dataset S1 to 4). The development of refractoriness to melatonin, indicative of the recovery

phase of the circannual timer, has been shown to develop independently in different

hypothalamic regions (Freeman and Zucker, 2001). Molecular data was collected from the

mediobasal hypothalamus (Bolborea et al., 2020), dorsomedial hypothalamus (Ebling et al.,

2008), and the paraventricular nucleus (Bittman et al., 1991), along with the pituitary gland

(Majumdar et al., 2023). This experimental approach provides a high-throughput and high-

frequency sampling resolution to comprehensively chart the induction, maintenance and

recovery of the mammalian circannual interval timer (Fig. 1H). Biodare2.0 identified 290

(Dataset S1) transcripts in the mediobasal hypothalamus as endogenously rhythmic. Density

heat-maps show an early, and robust upregulation of transcripts by 8 weeks SP associated

with the circannual interval induction (Fig. 1F). Consistent with previous reports (Petri et al.,

2016; Bao et al., 2019; Yoshimura et al., 2003), *Dio3* expression was significantly

upregulated and clustered with the initial wave of transcript expression (Fig.1F and *SI*

Appendix Fig. S1). A second cluster of transcripts were upregulated during the maintenance

phase occurring between 12- and 20-weeks in SP which primarily consisted of somatostatin

(*Sst*) and the cognate receptor subtype 5 (*Sstr5*). Increased *Sst* expression coincided with

the reduction in body and adipose tissue mass reflecting the conserved role in inhibiting

growth and metabolism (*SI Appendix* Figs. S1 and S2). Finally, a third wave of transcripts

became upregulated during the recovery phase, and largely resembled the molecular

landscape in LP hamsters (Fig.1F). qPCR assays for *Dio3* and *Sst* expression reflected the

sequencing transcript count pattern providing independent replication of the sequencing

analyses (Fig. S1). In the pituitary gland, 250 transcripts were identified as rhythmic and

included genes associated with secretagogue cells such as somatotrophs (*Gh*) and

lactotrophs (*Prl*) (Fig.1g and *SI Appendix* Fig S3 and Dataset S2). Similar rhythmic

expression was identified in other hypothalamic nuclei including the paraventricular nucleus

(518 transcripts) (*SI Appendix* Fig. S4 and S5 and Dataset S3), and the dorsomedial

hypothalamus (374 transcripts) (Fig. S4 and *SI Appendix* S6 and Dataset S4).

Dio3*-dependent changes in triiodothyronine induced body mass loss via *Sst

expression

Treatment of Djungarian hamster with the somatostatin agonist pasireotide decreases body

mass and is sufficient to drive changes similar to SP exposure (Dumbell et al., 2017, 2015).

Further, certain SST receptor subtypes appear to be involved in driving seasonal torpor in

Djungarian hamsters (Scherbarth et al., 2015). To determine if hypothalamic *Sst* expression

reflects the programmed rheostatic change in energy state governed by circannual interval

timing or homeostatic cues, tissues were collected from *ad libitum* (AL) fed hamsters under

either LP control condition or the maintenance phase (12-week SP). To induce a negative

energetic state, hamsters experienced an acute overnight food restriction (FR; 16hrs) (Fig.

2A). As anticipated, body mass decreased approximately 30% after exposure to SP (Fig.2B,

closed symbols); homeostatic negative energy state challenges by food restriction reduced

body mass by 1.5g on average for both LP (open symbols) and SP conditions (Fig.2C).

Epididymal adipose tissue mass was higher in LP compared to SP but did not decrease in

response to overnight FR (*SI Appendix* Fig. S7). Plasma insulin was significantly reduced in

response to SP treatment and FR suggesting circulating levels are an output of both

rheostatic and homeostatic signals (Fig. S7). The two manipulations permit the ability to

dissect the impact of rheostatic changes associated with circannual interval timing from

those involved in short-term homeostatic changes in energy balance. Mediobasal

hypothalamic *Sst* expression was significantly increased in SP conditions, but insensitive to

homeostatic challenges in energy state (Fig.2D). In contrast, mediobasal hypothalamic *Npy*

expression was significantly upregulated in food restricted hamsters but was similar across

LP and SP conditions (Fig.2E). *Prl* expression was higher in LP compared to SP conditions

(Fig. 2F). *Gh* expression did not change in response to either SP induced circannual

changes in energy state, or homeostatic manipulations (Fig. S7). To establish whether

mediobasal hypothalamic *Sst* expression is regulated by upstream *Dio3*-dependent changes

in local thyroid hormone catabolism, SP housed hamsters received sub-cutaneous injections

with either vehicle or triiodothyronine. A single triiodothyronine injection in SP hamsters was

sufficient to reduce *Sst* expression compared to vehicle controls (Fig. 2G). These data

indicate SP induced *Dio3* expression removes T3-dependent inhibition of *Sst* expression

leading to long-term inhibition of body mass during the maintenance phase of circannual

interval timing of rheostatic energy balance.

***Dio3* disfunction reduces the period of circannual interval timing**

Next, we sought to establish the functional role of *Dio3* signaling for circannual interval

timing of energy rheostasis in hamsters. Targeted genomic mutations of the *Dio3* gene

localized to the MBH were achieved via intracerebroventricular (ICV) injection of CRISPR-

Cas9 (*Dio3^{cc}*; Fig. S8) to assess the functional role for circannual interval timing. Control

hamsters were administered a blank crispr-cas9 construct (*Dio3^{wt}*). Two weeks following

surgery, hamsters were transferred to SP and programmed circannual changes in body

mass (Fig.3B) and pelage were monitored (*SI Appendix* Fig. S8). *Dio3^{cc}* hamsters had

slower body mass loss and regained body mass significantly quicker during the recovery

phase, compared to *Dio3^{wt}* controls. Similarly, the change in pelage color occurred later in

Dio3^{cc} compared *Dio3^{wt}*, and *Dio3^{cc}* was never observed to develop full white pelage (*SI*

Appendix Fig. S8). Lomb-Scargle period analysis of body mass identified that *Dio3^{cc}*

hamsters have notably shorter duration (period = 26.959) compared to *Dio3^{wt}* (period = 32)

(Figs. 3C and D). There was no significant difference in terminal white adipose tissue mass,

or daily food intake. A subpopulation of hamsters was physiologically nonresponsive (NR) to

the SP manipulation and maintained the higher LP body mass (Fig.3E). *Dio3* expression in

the mediobasal hypothalamus of these NR hamsters remained exceptionally low compared

to responsive SP hamsters at 8- and 12-weeks following exposure to SP (Fig.3F). These

data suggest that the inability to increase *Dio3* expression in NR hamsters acts as a form of

naturally occurring disfunction that prevents SP induction of circannual interval timing.

Discussion

The molecular and cellular basis of circannual clocks and timers in vertebrates is not well

characterized (Helm and Stevenson, 2014). Here, we sequenced the transcriptomes of three

hypothalamic regions and the pituitary gland, key neuroendocrine structures which govern

programmed circannual rheostatic and homeostatic processes. Our approach provided a

high sampling frequency of the circannual interval timer in Djungarian hamsters. The findings

produced a robust and comprehensive neural and molecular database which facilitates the

delineation of a circannual interval timer in mammals. Several novel transcripts displayed a

close correlation with the induction phase of the photoperiodic response. Among these was

Dio3, conducive with the removal of thyroid hormone (e.g., thyroxine, triiodothyronine) being

a key step in the transition to the winter phenotype. The photoinduced change in *Dio3*

expression induces a long-term change in physiology and morphology and our results

demonstrate that it is insensitive to short-term alterations in response to other environmental

cues (e.g., nutritional state). Functional manipulation of *Dio3* expression also delayed the

induction and enhanced the reversal of the winter phenotype suggesting that upregulation of

Dio3 acts to establish the period of circannual interval timing.

The transcriptomic data are consistent with evidence that shows *Sst* expression in

the arcuate nucleus is one critical event for the maintenance of a reduced rheostatic energy

state during the winter season (Marshall et al., 2024; Petri et al., 2016, 2014). In hamsters,

Sst expression is upregulated after *Dio3* induction and is associated with the low body mass

from week 12-20 before a downregulation in transcript levels coinciding with increased body

mass. By increasing circulating triiodothyronine in hamsters maintained in SP conditions, our

manipulation indicates that elevated hormone concentrations lead to the reduction in *Sst*

expression in the mediobasal hypothalamus (Fig.2). The overall expression patterns provide

the ability to develop a linear series of steps in which LP conditions have high local

concentrations of triiodothyronine in the mediobasal hypothalamus (Helfer and Stevenson,

2020; Murphy et al., 2012) which reduces the levels of *Sst* expression resulting in

consistently higher body mass. Exposure to SP induces *Dio3*, which catabolizes

triiodothyronine leading to the removal of the inhibition of *Sst* expression and reduction in

body mass. Local synthesis of triiodothyronine in the mediobasal hypothalamus by tanycytes

is an evolutionary conserved process for timing photoperiod-induced transitions across the

animal kingdom (Ebling and Lewis, 2018; Lewis and Ebling, 2017). Photoperiodic signals

derived from the pars tuberalis via thyrotropin-stimulating hormone are essential to regulate

Dio3 (and *Dio2*) expression in tanycytes leading to increased triiodothyronine content in LP

conditions for mammals (22,23), birds (Nakao et al., 2008) and fish (Lorgen et al., 2015).

The current limitation is understanding how SP signals regulate *Dio3* expression.

Undoubtedly, the duration of melatonin secretion during the daily nocturnal phase is

essential (Helfer et al., 2019). Future research is necessary to identify the conserved

upstream signals that govern the timing of *Dio3* expression.

These data support the existence of a neuroendocrine pathway for the long-term

rheostatic regulation, and another for short-term homeostatic control of energy stability

(Stevenson, 2024). The rheostatic pathway consists of triiodothyronine (T3) signaling

dependent on a *Dio3* switch to regulate *Sst* expression in the mediobasal hypothalamus

whereas short-term regulation of energy stability via nutrient availability is homeostatically

regulated by well-established orexigenic (e.g., *Npy*) (Yeo et al., 2021). In response to the

output of both pathways, peripheral effects may be mediated by acute and chronic changes

in pancreatic insulin secretion (Fig.4). Overall, the findings provide a clear neural and cellular

circuit for circannual interval timing of energy rheostasis and demonstrate *Dio3* as a gene

critical for the control of seasonal life history transitions.

Materials and Methods

Ethics

All procedures were in accordance with the National Centre for the Replacement,

Refinement and Reduction of Animals in Research ARRIVE guidelines

(https://www.nc3rs.org.uk/revision-arrive-guidelines). All procedures were approved by the

Animal Welfare and Ethics Review Board at the University of Glasgow and conducted under

the Home Office Project License PP5701950.

Animals

All animals were obtained from a colony of Djungarian hamsters (3-8 months of age) kept at

the University of Glasgow, Veterinary Research Facility. Animals were held at a room

temperature of 21°C, and a humidity of 50%. The colony room was kept on a long

photoperiod (16L:8D). All hamsters had food (SDS [BK001]) and water *ad libitum*.

Experiment 1 – Defining the circannual interval timer in hamsters

Adult male hamsters (N = 100) were required to assess the physiological and molecular

basis of circannual interval timing in hamsters. A photoperiod reference group was kept on

LP (n=14) (8D:16L) conditions for the duration of the study. The SP treatment groups

consisted of hamsters moved to SP (16D:8L) at 4 weeks intervals. The groups consisted of

SP for 4 (n=12), 8 (n=12), 12 (n=12), 16 (n=11), 20 (n=12), 24 (n=10), 28 (n=7) and 32

(n=10) weeks. This photoperiod schedule reliably captures the induction, maintenance, and

recovery of circannual interval timing in hamsters (Prendergast et al., 2002). Pelage score

was determined using visual assessment and defined scale in which a score of 1 is the

summer dark agouti coat; a score of 2 is a transition between agouti and white; and a score

of 3 is a full winter white pelage (Marshall et al., 2024). Body mass was monitored to the

nearest 0.1g using an Ohaus portable scale (TA301). Food intake was measured by

weighing the food pellets at the mid-point on two consecutive days to obtain a daily value.

Food weight was measured using an Ohaus scale (TA301). Hamsters were killed between 2-

4hr after lights on by cervical dislocation followed by exsanguination. Adipose tissue was

immediately dissected and measured using a Sartorius cp64 anatomical balance. A terminal

blood sample was collected to measure glucose (Accu-check Performa nano blood glucose

meter). Circulating levels of insulin and total glucagon-like peptide 1 (GLP1) using ELISA

(MesoScale Discovery, UK) at Core Biochemical Assays laboratories, Cambridge, UK. Brain

and pituitary gland tissue was quickly extracted and frozen on powdered dry ice and stored

at -70°C. Epididymal white adipose tissue was dissected and weighed using Sartorius cp64

anatomical balance and stored at -70°C.

Brains were cut into 200μm coronal sections using a Leica CM1520 cryostat. Anatomical

structures (optic tract to the infundibular stem; Bregma -2.12mm to -3.80mm) were used to

isolate the mediobasal hypothalamus (MBH), dorsomedial hypothalamus (DMH) and the

paraventricular nucleus (PVN). Bilateral tissue punches were performed using an integra

Miltex 1mm disposable biopsy punch. Tissue punches were stored at -70°C until

transcriptomic sequencing and confirmatory qPCR analyses (see below).

Experiment 2 – Dissociating mechanisms that govern circannual rheostatic versus

homeostatic energy stability.

A total of N=35 adult male hamsters were used to determine molecular signatures of

rheostatic versus homeostatic energy stability. A LP control group was kept on long

photoperiod for the duration of the study (n = 18). A subset of hamsters was moved to SP (n

= 17) for 12 weeks to obtain animals in the maintenance phase of circannual interval timing.

Food (SDS [BK001]) and tap water were provided *ad libitum* to both LP and SP animals for

the 12 weeks duration. Body mass was measured every two weeks using a Traveler Ohaus

portable balance. On the final night of the experiment, 50% of hamsters (i.e., LP = 9, SP = 9)

were kept on food and water *ad libitum*. The other 50% of hamsters (i.e., LP = 9, SS = 8)

served as the homeostatic treatment group and received an acute food restriction by

removing all food. Overnight food restriction induces a robust negative energy state that

reliably results in a 1–2-gram loss in hamster body mass (Bao *et al*., 2019). Body mass was

measured prior to food restriction, and again before tissue collection on the subsequent day.

Hamsters were killed between 2-4hr after lights on by cervical dislocation and then

exsanguination. A terminal blood sample was collected to measure insulin (MesoScale

Discovery, Core Biochemical Assays laboratories, Cambridge, UK). Body mass and

epididymal white adipose tissue mass were measured using a Sartorius cp64 anatomical

balance. Brains and pituitary glands were dissected and were immediately placed on dry ice

and stored at -70°C. Brains were cut into 200μM sections using a Leica CM1520 cryostat.

Anatomical structures (optic tract to the infundibular stem; Bregma -2.12mm to -3.80mm)

were used to isolate the MBH. Bilateral tissue punches were performed using an integra

Miltex 1mm disposable biopsy punch. Brain tissue punches and pituitary gland samples

were stored at -70°C until qPCR assay (see below).

Experiment 3 – Sufficiency of triiodothyronine to induce LP neuroendocrine state.

Adult male hamsters (n = 35) were used to determine the impact of triiodothyronine on *Sst*

expression. Hamsters were transferred from LP to SP for 12 weeks (n = 24) to initiate

circannual interval timing and establish the maintenance phase. Hamsters were divided into

two groups for the last week of SP treatment. A SP reference group received subcutaneous

injections of 0.9% w/v saline for 7 days (n = 10). To determine the sufficiency of

triiodothyronine to regulate *Sst* expression, hamsters (n=11) were administered a single

subcutaneous injection of 5µg/100ul triiodothyronine (Merck 102467157) prior to lights off on

the final night. Subcutaneous injections of triiodothyronine are well established to regulate

transcript expression in the hamster hypothalamus (Banks et al., 2016; Bao et al., 2019;

Stevenson et al., 2014). The subsequent day hamsters were sacrificed by cervical

dislocation followed by exsanguination between 4-5hr after lights on. Terminal body mass

and epididymal adipose tissue mass were recorded using a Sartorius cp64 anatomical

balance. Brains were dissected and stored immediately at -70°C until sectioning. Brains

were cut into 200µM sections using a Leica CM1520 cryostat. Anatomical structures (optic

tract to the infundibular stem; Bregma -2.12mm to -3.80mm) were used to isolate the MBH.

Bilateral tissue punches were performed using an integra Miltex 1mm disposable biopsy

punch. Tissue punches were stored at -70°C until qPCR assays (see below).

Experiment 4 – Functional manipulation of *Dio3* expression in hamsters

Targeted neuroanatomical injections, using highly precise Guide RNA (gRNA) against *Dio3*

and Crispr-Cas9 vectors was essential to ensure the long-term inhibition of *Dio3* function in

the MBH in hamsters; an approach necessary to reliably examine the functional significance

of *Dio3* for the duration of circannual interval timing. Custom-made Crispr-cas9 constructs

were packaged in a lentiviral vector by Merck Life science UK. Lentiviral vectors are

established to effectively transfect in adult Syrian hamsters (Gao M *et al.*, 2014) and in

Siberian hamsters (Munley K *et al.*, 2022). The lentiviral vector used was U6-gRNA:ef1a-

puro-2A-Cas9-2A-tGFP (Sigma All-in-One vector). gRNA for *Dio3* were designed using

CHOPCHOP (Labun *et al.,* 2019). Three gRNA were designed to target the *Mus musculus*

Dio3 gene (gRNA1: CGACAACCGTCTGTGCACCCTGG; gRNA2:

GTTCCCGCGCTTCCTAGGCACGG, and gRNA3: GACCCAGCCGTCGGATGGGTGGG).

Only gRNA 1 and 2 were identified to align with 5' end of the Siberian hamster *Dio3* gene

and were predicted to align with chromosome 12 at locations 110279543 and 110279375

respectively (fig. S8). To check specificity of gRNA sequences, we conducted BLAST using

the NCBI dataset. Both gRNA1 and gRNA2 were found to align 100% with *Dio3* and did not

align 23/23 with any other region.

Adult male hamsters (n = 10) were selected to assess the impact of *Dio3* genome

modification on circannual interval timing. Hamsters were kept on LP conditions to maintain

the summer phenotype. Hamsters were then divided into two treatment groups. The control

group included hamsters (n=6) that received intracerebroventricular (ICV) injections of blank

Crispr-Cas9 constructs that were packaged into a lentivirus (*Dio3^{wt}*; U6-gRNA:ef1a-puro-2A-

Cas9-2A-tGFP (Sigma All-in-One vector). The treatment group consisted of hamsters (n=4)

that received an ICV injection of Crispr-cas9 constructs that harbored both gRNA1 and

gRNA2 (*Dio3^{cc}*).

Intracerebroventricular injections provide the ability to induce neuroanatomically localized

manipulations in *Dio3* gene function in the adult brain. Crispr-Cas9 constructs were delivered

via stereotaxic ICV injection into the third ventricle to target the MBH tanycytes. Under

general anesthesia (5% induction, 2% maintenance of isoflurane mixed with oxygen) a SGE

series II syringe was positioned along the midline at Bregma -1.5mm, to a depth of -5.7mm.

The anatomical coordinates were refined based on estimates taken from adult hamster

brains (Steward et al., 2003) and tested using fresh hamster cadavers. Analgesia was

administered subcutaneously (5mg/kg Rimadyal and 0.1mg/kg buprenorphine). A small

~1cm incision was made by scalpel to expose the skull and a small hole was drilled using a

Bone Micro Drill for Brain surgery (Harvard Apparatus). Lentivirus was administered using a

Pump 11 Elite Programmable Syringe injection system (Harvard Apparatus) and 1ul of viral

vector (2-2.5 E+13 vg/ml) was delivered over 10 minutes at a rate of 0.1ul per minute. To

allow viral diffusion and prevent backflow the syringe was kept in place after injection for 1-

minute and then the syringe was raised by 1mm and kept in place for another 1-minute.

Animals were then moved to heated home cages and given mashed food to aid recovery.

Hamsters were provided with 2-weeks recovery and were monitored to ensure stable body

mass, locomotor activity and continued food and water intake.

To assess the impact of *Dio3[cc]* on circannual interval timing hamsters were moved to SP for

32 weeks and body mass and pelage were measured biweekly. Pelage score was

determined using visual assessment and defined scale in which a score of 1 is the summer

dark agouti coat; a score of 2 is a transition between agouti and white; and a score of 3 is a

full winter white pelage (Marshall et al., 2024). Body mass was monitored to the nearest 0.1g

using an Ohaus portable scale (TA301). After 32 weeks, hamsters were killed 4-5hr after

lights on by cervical dislocation followed by exsanguination. Brains were dissected and

stored immediately at -70°C. Brains were cut into 200μM sections using a Leica CM1520

cryostat. Anatomical structures (optic tract to the infundibular stem; Bregma -2.12mm to -

3.80mm) were used to isolate the MBH. Bilateral tissue punches were performed using an

integra Miltex 1mm disposable biopsy punch. Tissue punches were stored at -70°C. DNA

was extracted from dissected tissues using the DNeasy Blood & Tissue Kit (QIAGEN) as per

manufacturer's instructions. The *Dio3* gene was amplified using OneTaq Quick-Load 2X

mastermix, as per manufacturer's instructions (New England Biolabs). Amplification was

achieved using a SimpliAmp thermal cycler (Applied Biosystems) the following thermal

cycling conditions (94°C for 30 seconds, (94°C for 15 seconds, 62°C for 30 seconds, 68°C

for 3 minutes) for 45 cycles, 68°C for 5 minutes, 4°C until further analysis.). The resultant

amplicon was purified using the Qiaquick PCR purification kit (Qiagen) as per

manufacturer's instructions. Isolated *Dio3* PCR amplicons were sequenced using the

Eurofins LightRun sanger sequencing service. The forward primer provided to Eurofins was

CATGCTCCGCTCCCTGCTGCTTCA (5'-3'). CRISPR modified transcripts were aligned to a

reference wild type transcript (animal #821) using the Tracy sanger basecaller and aligner

(Rausch et al., 2020). Alignments were visualized using the Ugene integrated bioinformatics

tool (Okonechnikov et al., 2012). In case mutation was not obvious, an additional analysis

was carried out using the associated online resource SAGE (https://www.gear-

genomics.com/), to investigate the decomposition error of the modified transcripts with the

wild-type control (821). Four $Dio3^{wt}$ hamsters had $Dio3$ sequences that fully matched the

reference genome (fig. S8c). Two hamster $Dio3$ sequences did not automatically align, and

the raw reads were visually inspected. No mutations or sequence mutations were detected

in the raw sequence trace. Three $Dio3^{cc}$ treated hamsters had $Dio3$ sequences with

evidence of genomic mutation evidenced by low decomposition error (*SI Appendix* Figs. S8C

and D). One $Dio3^{cc}$ treated hamster did not show any genomic mutation (905) and was

removed from the final analysis. Hamster 905 had a body mass circannual interval period of

32 weeks and we propose the animal represents a false positive control.

Experiment 5 – Naturally occurring variation in *Dio3* expression and photoperiodic

response.

A subset of hamsters does not physiologically respond to SP treatment and are classified as

photoperiodically non-responsive (NR) (Przybylska-Piech and Jefimow, 2022). Adult

hamsters (n = 14) were held in LP or SP for 8 weeks. A group of male non responders (n =

4) were identified due to the absence of body mass and pelage change in response to SP

(Fig. 3e). A reference group of male SP responders (n = 10) were collected. MBH was

extracted, RNA isolated and cDNA synthesized. qPCR was carried out for *Dio3* and *Sst*

expression. Hamsters were killed between 2-4hr after lights on.

RNA extraction

RNA was extracted using the Qiagen RNEasy plus mini kit. Tissues were homogenized

using a Polytron PT 1200 E. RNA was then extracted from homogenized tissue following the

manufacturer's instructions. RNA was tested for quality and quantity using an ND-1000

Nanodrop.

Next Generation Sequencing and data processing

Oxford Nanopore Sequencing was used to carry out transcriptomic sequencing. RNA from

MBH, DMH and PVN samples was sequenced using the PCR-cDNA barcoding kit following

manufacturer's instructions (SQK-PCB109; Oxford Nanopore). RNA from the pituitary gland

samples was synthesized using the direct cDNA sequencing kit following manufacturer's

instructions (SQK-DCS109 with EXP-NBD104; Oxford Nanopore). Bioinformatic analysis

was carried out on Linux within a conda environment. Raw reads (Fast5) were demultiplexed

and basecalled using guppy basecaller (4.2.1). Porechop (0.2.4) was used to remove

adapters from reads and Filtlong (v0.2.0) was used to filter for quality and read length

(length: >25 base pairs, quality score > 9). Transcripts were aligned to Mus musculus

genome (GRCm39) using Minimap2 (Li, 2018). Previously, rodent genomes have been

demonstrated to show significant similarity, such that cross-species transcriptomic analysis

appears feasible (Przybylska-Piech and Jefimow, 2022). Transcript expression levels were

generated using Salmon (v0.14.2) (Patro et al., 2017) and EdgeR (v3.24.3) (Robinson et al.,

2010) was used to filter lowly expressed transcripts.

cDNA synthesis and qPCR

RNA was transformed into cDNA for qPCR analysis using superscript III (Invitrogen) as per

manufacturer's instructions. Quantification of transformed cDNA was achieved using Brilliant

II SYBR Mastermix (Agilent). Stock forward and reverse primers (100pmol/µl) were mixed

and diluted in nuclease free water to 20pmol/µl. A working SYBR mixture was prepared by

mixing 1-part primer mixture per 24-part SYBR mastermix. Reaction mixtures were prepared

in wells on a 96-well plate by mixing 4.8µl of normalised sample and 4.8µl of SYBR working

mix. All reactions were performed in duplicate. Primer and qPCR parameters are outlined in

Table S1. qPCR reactions were carried out in a Stratagene Mx3000P thermal cycler. Cycling

conditions utilized were in sequence; at 95°C for 5 minutes (denaturing), 40 cycles at (95°C

for 30 seconds, X°C for 1 minute – see table, 72°C for 30 seconds with fluorescent

measurement at end), 95°C for 1 minute, 55°C for 30 seconds increasing to 95°C (Melt

curve analysis). Melt cure analysis was used to determine specificity of amplification. Data

was analyzed for meanCT, efficiency and variability using PCR Miner (Zhao and Fernald,

2005). Logfold CT was calculated using the ΔΔCT method using 18s and/or *Hrpt* reference

transcripts as these transcripts are stable across photoperiodic conditions (Stewart et al.,

2022).

Statistical Analysis

Raw data is provided in Data file 1 and sequencing data is provided in Table S1-S4.

Physiological and qPCR data sets were tested for normality using the Shapiro test and

transformed by log transformation if normality was violated. Statistical significance was

determined using a two-way ANOVA. Post hoc analysis of significant data was achieved by

t-test. Type 1 error was minimized by utilizing the Bonferroni adjustment on post-hoc

analyses. Data were analyzed using non-linear regression for rhythmicity using the online

resource BioDare 2.0 (Zielinski et al., 2014) (biodare2.ed.ac.uk). The empirical JTK_CYCLE

method was used for detection of rhythmicity and the classic BD2 waveform set was used

for comparison testing. Rhythmicity was determined by a Benjamini-Hochberg controlled

false discovery rate (BH corrected FDR < 0.1). The false discovery rate of this data set was

controlled by applying the Benjamini-Hochberg method (Benjamini and Hochberg, 1995).

Clusters within the data set were identified using the gap-statistic and clustered using k-

means clustering. Gene ontology analysis was carried out using ShinyGO v0.77 (Ge et al.,

2020).

Data and materials availability

All data are available in Dataset S1-S5. The code is available via github repository

(https://github.com/CS-Gla/Hamster_seasonal_sci_cs_2024). Sequencing information was

deposited in GEO (GSE274003).

Acknowledgments

The authors thank Gerald Lincoln for helpful discussions which contributed to the

conceptualization of the research. The authors thank the technical assistance provided by

Nicola Munro, Ana Monterio, and Fallon Cuthill. Biorender.com was used for icons used in

Fig. 1-3 and fig. S9. Funding was provided via Leverhulme Trust award LT-RL-2019-06 (TJS)

and ISSF Wellcome Trust award (TJS, PJM, FJPE).

References

Banks R, Delibegovic M, Stevenson TJ. 2016. Photoperiod- and Triiodothyronine-dependent

Regulation of Reproductive Neuropeptides, Proinflammatory Cytokines, and Peripheral

Physiology in Siberian Hamsters (*Phodopus sungorus*). *J Biol Rhythms* **31**:299–307.

doi:10.1177/0748730416637707

Bao R, Onishi KG, Tolla E, Ebling FJP, Lewis JE, Anderson RL, Barrett P, Prendergast BJ,

Stevenson TJ. 2019. Genome sequencing and transcriptome analyses of the Siberian

hamster hypothalamus identify mechanisms for seasonal energy balance. *Proc Natl Acad*

Sci USA **116**:13116–13121. doi:10.1073/pnas.1902896116

Barrett P, Ross AW, Balik A, Littlewood PA, Mercer JG, Moar KM, Sallmen T, Kaslin J,

Panula P, Schuhler S, Ebling FJ, Ubeaud C, Morgan PJ. 2005. Photoperiodic Regulation of

Histamine H3 Receptor and VGF Messenger Ribonucleic Acid in the Arcuate Nucleus of the

Siberian Hamster. *Endocrinology* **146**:1930–1939. doi:10.1210/en.2004-1452

Benjamini Y, Hochberg Y. 1995. Controlling the False Discovery Rate: A Practical and

Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society Series B:*

Statistical Methodology **57**:289–300. doi:10.1111/j.2517-6161.1995.tb02031.x

Bittman EL, Bartness TJ, Goldman BD, DeVries GJ. 1991. Suprachiasmatic and

paraventricular control of photoperiodism in Siberian hamsters. *American Journal of*

Physiology-Regulatory, Integrative and Comparative Physiology **260**:R90–R101.

doi:10.1152/ajpregu.1991.260.1.R90

Bolborea M, Pollatzek E, Benford H, Sotelo-Hitschfeld T, Dale N. 2020. Hypothalamic

tanycytes generate acute hyperphagia through activation of the arcuate neuronal network.

Proc Natl Acad Sci USA **117**:14473–14481. doi:10.1073/pnas.1919887117

Cohen JM, Lajeunesse MJ, Rohr JR. 2018. A global synthesis of animal phenological

responses to climate change. *Nature Clim Change* **8**:224–228. doi:10.1038/s41558-018-

0067-3

Denlinger DL. 1974. Diapause potential in tropical flesh flies. *Nature* **252**:223–224.

doi:10.1038/252223a0

Dumbell R, Petri I, Scherbarth F, Diedrich V, Schmid HA, Steinlechner S, Barrett P. 2017.

Somatostatin Agonist Pasireotide Inhibits Exercise-Stimulated Growth in the Male Siberian

Hamster (Phodopus sungorus). *J Neuroendocrinology* **29**:jne.12448. doi:10.1111/jne.12448

Dumbell RA, Scherbarth F, Diedrich V, Schmid HA, Steinlechner S, Barrett P. 2015.

Somatostatin Agonist Pasireotide Promotes a Physiological State Resembling Short-Day

Acclimation in the Photoperiodic Male Siberian Hamster (Phodopus sungorus). *J*

Neuroendocrinology **27**:588–599. doi:10.1111/jne.12289

Duncan S, Holm S, Questa J, Irwin J, Grant A, Dean C. 2015. Seasonal shift in timing of

vernalization as an adaptation to extreme winter. *eLife* **4**:e06620. doi:10.7554/eLife.06620

Ebling FJP, Lewis JE. 2018. Tanycytes and hypothalamic control of energy metabolism. *Glia*

66:1176–1184. doi:10.1002/glia.23303

Ebling FJP, Wilson D, Wood J, Hughes D, Mercer JG, Morgan PJ, Barrett P. 2008. The

Thyrotropin-Releasing Hormone Secretory System in the Hypothalamus of the Siberian

Hamster in Long and Short Photoperiods. *J Neuroendocrinology* **20**:576–586.

doi:10.1111/j.1365-2826.2008.01702.x

Freeman DA, Zucker I. 2001. Refractoriness to melatonin occurs independently at multiple

brain sites in Siberian hamsters. *Proc Natl Acad Sci USA* **98**:6447–6452.

doi:10.1073/pnas.111140398

Gao M, Zhang B, Liu J, Guo X, Li H, Wang T, Zhang Z, Liao J, Cong N, Wang Y, Yu L, Zhao

D, Liu G. 2014. Generation of transgenic golden Syrian hamsters. *Cell Res* **24**:380–382.

doi:10.1038/cr.2014.2

Ge SX, Jung D, Yao R. 2020. ShinyGO: a graphical gene-set enrichment tool for animals
and plants. *Bioinformatics* **36**:2628–2629. doi:10.1093/bioinformatics/btz931

Gwinner E, Dittami J. 1990. Endogenous Reproductive Rhythms in a Tropical Bird. *Science*
249:906–908. doi:10.1126/science.249.4971.906

Hanon EA, Lincoln GA, Fustin J-M, Dardente H, Masson-Pévet M, Morgan PJ, Hazlerigg
DG. 2008. Ancestral TSH Mechanism Signals Summer in a Photoperiodic Mammal. *Current*
Biology **18**:1147–1152. doi:10.1016/j.cub.2008.06.076

Helfer G, Barrett P, Morgan PJ. 2019. A unifying hypothesis for control of body weight and
reproduction in seasonally breeding mammals. *J Neuroendocrinology* **31**:e12680.
doi:10.1111/jne.12680

Helfer G, Stevenson TJ. 2020. Pleiotropic effects of proopiomelanocortin and VGF nerve
growth factor inducible neuropeptides for the long-term regulation of energy balance.
Molecular and Cellular Endocrinology **514**:110876. doi:10.1016/j.mce.2020.110876

Helm B, Stevenson TJ. 2014. Circannual rhythms: history, present challenges, future
directions. In *Annual, Lunar, and Tidal Clocks: patterns and mechanisms of nature's*
enigmatic rhythms. Spring Japan, pg 203-225.

Labun K, Montague TG, Krause M, Torres Cleuren YN, Tjeldnes H, Valen E. 2019.
CHOPCHOP v3: expanding the CRISPR web toolbox beyond genome editing. *Nucleic Acids*
Research **47**:W171-174

Lewis JE, Ebling FJP. 2017. Tanycytes As Regulators of Seasonal Cycles in Neuroendocrine
Function. *Front Neurol* **8**. doi:10.3389/fneur.2017.00079

Li H. 2018. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**:3094–
3100. doi:10.1093/bioinformatics/bty191

Lincoln GA, Clarke IJ, Hut RA, Hazlerigg DG. 2006. Characterizing a Mammalian Circannual
Pacemaker. *Science* **314**:1941–1944. doi:10.1126/science.1132009

Lorgen M, Casadei E, Król E, Douglas A, Birnie MJ, Ebbesson LOE, Nilsen TO, Jordan WC,
Jørgensen EH, Dardente H, Hazlerigg DG, Martin SAM. 2015. Functional Divergence of
Type 2 Deiodinase Paralogs in the Atlantic Salmon. *Current Biology* **25**:936–941.
doi:10.1016/j.cub.2015.01.074

Majumdar G, Liddle TA, Stewart C, Marshall CJ, Bain M, Stevenson T. 2023. FSHβ links
photoperiodic signaling to seasonal reproduction in Japanese quail. *eLife* **12**:RP87751.
doi:10.7554/eLife.87751.3

Marshall CJ, Blake A, Stewart C, Liddle TA, Denizli I, Cuthill F, Evans NP, Stevenson TJ.
2024. Prolactin Mediates Long-Term, Seasonal Rheostatic Regulation of Body Mass in
Female Mammals. *Endocrinology* **165**:bqae020. doi:10.1210/endocr/bqae020

Mercer JG, Moar KM, Ross AW, Hoggard N, Morgan PJ. 2000. Photoperiod regulates
arcuate nucleus POMC, AGRP, and leptin receptor mRNA in Siberian hamster
hypothalamus. *American Journal of Physiology-Regulatory, Integrative and Comparative*
Physiology **278**:R271–R281. doi:10.1152/ajpregu.2000.278.1.R271

Munley KM, Dutta S, Jasnow AM, Demas GE. 2022. Adrenal MT1 melatonin receptor
expression is linked with seasonal variation in social behavior in male Siberian hamsters.
Hormones and Behavior **138**:105099. doi:10.1016/j.yhbeh.2021.105099

Murphy M, Jethwa PH, Warner A, Barrett P, Nilaweera KN, Brameld JM, Ebling FJP. 2012.
Effects of Manipulating Hypothalamic Triiodothyronine Concentrations on Seasonal Body
Weight and Torpor Cycles in Siberian Hamsters. *Endocrinology* **153**:101–112.
doi:10.1210/en.2011-1249

Nakao N, Ono H, Yamamura T, Anraku T, Takagi T, Higashi K, Yasuo S, Katou Y, Kageyama
S, Uno Y, Kasukawa T, Iigo M, Sharp PJ, Iwasawa A, Suzuki Y, Sugano S, Niimi T, Mizutani
M, Namikawa T, Ebihara S, Ueda HR, Yoshimura T. 2008. Thyrotrophin in the pars tuberalis
triggers photoperiodic response. *Nature* **452**:317–322. doi:10.1038/nature06738

Okonechnikov K, Golosova O, Fursov M, the UGENE team. 2012. Unipro UGENE: a unified
bioinformatics toolkit. *Bioinformatics* **28**:1166–1167. doi:10.1093/bioinformatics/bts091

Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. 2017. Salmon provides fast and bias-
aware quantification of transcript expression. *Nat Methods* **14**:417–419.
doi:10.1038/nmeth.4197

Pengelley ET, Fisher KC. 1957. Onset and Cessation of Hibernation under Constant
Temperature and Light in the Golden-mantled Ground Squirrel (Citellus lateralis). *Nature*
180:1371–1372. doi:10.1038/1801371b0

Pérez JH, Tolla E, Dunn IC, Meddle SL, Stevenson TJ. 2019. A Comparative Perspective on
Extra-retinal Photoreception. *Trends in Endocrinology & Metabolism* **30**:39–53.
doi:10.1016/j.tem.2018.10.005

Petri I, Diedrich V, Wilson D, Fernández-Calleja J, Herwig A, Steinlechner S, Barrett P. 2016.
Orchestration of gene expression across the seasons: Hypothalamic gene expression in
natural photoperiod throughout the year in the Siberian hamster. *Sci Rep* **6**:29689.
doi:10.1038/srep29689

Petri I, Dumbell R, Scherbarth F, Steinlechner S, Barrett P. 2014. Effect of Exercise on
Photoperiod-Regulated Hypothalamic Gene Expression and Peripheral Hormones in the
Seasonal Dwarf Hamster Phodopus sungorus. *PLoS ONE* **9**:e90253.
doi:10.1371/journal.pone.0090253

Piao S, Liu Q, Chen A, Janssens IA, Fu Y, Dai J, Liu L, Lian X, Shen M, Zhu X. 2019. Plant
phenology and global climate change: Current progresses and challenges. *Global Change*
Biology **25**:1922–1940. doi:10.1111/gcb.14619

Prendergast BJ, Mosinger B, Kolattukudy PE, Nelson RJ. 2002. Hypothalamic gene
expression in reproductively photoresponsive and photorefractory Siberian hamsters. *Proc*
Natl Acad Sci USA **99**:16291–16296. doi:10.1073/pnas.232490799

Prendergast BJ, Renstrom RA, Nelson RJ. 2004. Genetic Analyses of a Seasonal Interval
Timer. *J Biol Rhythms* **19**:298–311. doi:10.1177/0748730404266626

Przybylska-Piech AS, Jefimow M. 2022. Siberian hamsters nonresponding to short
photoperiod use fasting-induced torpor. *Journal of Experimental Biology* **225**:jeb244222.
doi:10.1242/jeb.244222

Rausch T, Fritz MH-Y, Untergasser A, Benes V. 2020. Tracy: basecalling, alignment,
assembly and deconvolution of sanger chromatogram trace files. *BMC Genomics* **21**:230.
doi:10.1186/s12864-020-6635-8

Ricklefs RE. 1991. Structures and Transformations of Life Histories. *Functional Ecology*
5:174. doi:10.2307/2389255

Robinson MD, McCarthy DJ, Smyth GK. 2010. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics* **26**:139–140.
doi:10.1093/bioinformatics/btp616

Scherbarth F, Diedrich V, Dumbell RA, Schmid HA, Steinlechner S, Barrett P. 2015.
Somatostatin receptor activation is involved in the control of daily torpor in a seasonal
mammal. *American Journal of Physiology-Regulatory, Integrative and Comparative*
Physiology **309**:R668–R674. doi:10.1152/ajpregu.00191.2015

Stevenson TJ. 2024. Defining the brain control of physiological stability. *Hormones and*
Behavior **164**:105607. doi:10.1016/j.yhbeh.2024.105607

Stevenson TJ, Onishi KG, Bradley SP, Prendergast BJ. 2014. Cell-autonomous
iodothyronine deiodinase expression mediates seasonal plasticity in immune function. *Brain,*
Behavior, and Immunity **36**:61–70. doi:10.1016/j.bbi.2013.10.008

Steward CA, Horan TL, Schuhler S, Bennett GW, Ebling FJP. 2003. Central administration of
thyrotropin releasing hormone (TRH) and related peptides inhibits feeding behavior in the
Siberian hamster: *NeuroReport* **14**:687–691. doi:10.1097/00001756-200304150-00006

Stewart C, Liddle TA, Stevenson TJ. 2022. Abundance, efficiency, and stability of reference
transcript expression in a seasonal rodent: The Siberian hamster. *PLoS ONE* **17**:e0275263.
doi:10.1371/journal.pone.0275263

Wood S, Loudon A. 2014. Clocks for all seasons: unwinding the roles and mechanisms of
circadian and interval timers in the hypothalamus and pituitary. *Journal of Endocrinology*
222:R39–R59. doi:10.1530/JOE-14-0141

Wood SH, Hindle MM, Mizoro Y, Cheng Y, Saer BRC, Miedzinska K, Christian HC, Begley N,
McNeilly J, McNeilly AS, Meddle SL, Burt DW, Loudon ASI. 2020. Circadian clock
mechanism driving mammalian photoperiodism. *Nat Commun* **11**:4291. doi:10.1038/s41467-
020-18061-z

Woodfill CJI, Wayne NL, Moenter SM, Karsch FJ. 1994. Photoperiodic Synchronization of a
Circannual Reproductive Rhythm in Sheep: Identification of Season-Specific Time Cues1.
Biology of Reproduction **50**:965–976. doi:10.1095/biolreprod50.4.965

Yeo GSH, Chao DHM, Siegert A-M, Koerperich ZM, Ericson MD, Simonds SE, Larson CM,
Luquet S, Clarke I, Sharma S, Clément K, Cowley MA, Haskell-Luevano C, Van Der Ploeg L,
Adan RAH. 2021. The melanocortin pathway and energy homeostasis: From discovery to
obesity therapy. *Molecular Metabolism* **48**:101206. doi:10.1016/j.molmet.2021.101206

Yoshimura T, Yasuo S, Watanabe M, Iigo M, Yamamura T, Hirunagi K, Ebihara S. 2003.
Light-induced hormone conversion of T4 to T3 regulates photoperiodic response of gonads
in birds. *Nature* **426**:178–181. doi:10.1038/nature02117

Zhao S, Fernald RD. 2005. Comprehensive Algorithm for Quantitative Real-Time
Polymerase Chain Reaction. *Journal of Computational Biology* **12**:1047–1064.
doi:10.1089/cmb.2005.12.1047

Zielinski T, Moore AM, Troup E, Halliday KJ, Millar AJ. 2014. Strengths and Limitations of
Period Estimation Methods for Circadian Data. *PLoS ONE* **9**:e96462.
doi:10.1371/journal.pone.0096462

647   **Figures**

648
649

Figure 1. Molecular basis of circannual interval timing for morphology and

physiology. Experimental design in which Djungarian hamsters were kept in long

photoperiod (LP) or transferred to short photoperiod (SP) (a). SP induced pelage color

change from LP agouti to winter white after 12-weeks which reversed to agouti by 28-weeks

($F_{8,91}$ = 50.77; P < 0.001) (b). Torpor was identified in hamster between 12- and 20-weeks

SP exposure indicate by red line (b). SP exposure induced significant reduction in body

mass ($F_{8,91}$ = 13.428; P < 0.001) (c), food intake ($F_{8,91}$ = 10.860; P < 0.001; denoted as

downward triangles) (d) and adipose mass ($F_{8,91}$ = 27.929; P < 0.001; denoted as upward

triangles) (d). Plasma GLP-1 did not vary in response to SP manipulation; denoted as

upward triangles (e). SP exposure resulted in significant changes in plasma glucose around

the onset of torpor; denoted as downward triangles ($F_{8,91}$ = 3.117; P < 0.05) (e). BioDare2.0

heatmaps of mediobasal hypothalamus (f) and pituitary gland (g) transcripts from Djungarian

hamster collected at 4 -week SP intervals. Transcripts identified as highly rhythmic (FDR <

0.1) showed three distinct phases of expression that coincide with the induction,

maintenance and recovery of circannual interval timing. Deiodinase type-3 (*Dio3*) was

upregulated during the induction phase, whereas transcripts associated with energy stability

(e.g., somatostatin [Sst and Sstr5]) were upregulated during the maintenance phase. All

rhythmic transcripts reverted to the LP condition by 28-weeks SP exposure. Polar scatter

chart of significant transcripts from mediobasal hypothalamus and pituitary gland provide a

comprehensive seasonal clock for mammalian circannual interval timing across

neuroendocrine tissues (h). Green line indicates the induction phase, red line indicates the

maintenance phase, and the blue line represents the recovery phase. Letters denote

significant differences between treatment groups ($P < 0.05$) (b-e). In panel f and g, the scale

bar represents transcript expression as z-scores from 1 (upregulation in red) to -1 (down

regulation in blue). Density heatmaps are adjacent to transcript expression heat maps and

display the average z-score expression of each individual cluster on the y-axis, and the

graph shows percentile lines and density of z-score expression.

Figure 2. Somatostatin expression reflects programmed circannual interval timing
that is dependent on *Dio3* regulation of local triiodothyronine signaling. The
experimental design in which SP exposure induced rheostatic reduction in body mass after
12-weeks, and then hamsters experienced either an overnight food restriction (FR) for 16
hours or maintained food ad libitum (AL) (a). SP induced a significant reduction in body
mass ($F_{1,22}$ = 3.85; P < 0.01) (b). Food restriction further reduced body mass ($F_{1,22}$ = 25.46; P
< 0.001) (c). *Sst* expression in the mediobasal hypothalamus was significantly increased in
SP ($F_{1,15}$ = 5.59; P < 0.05) but did not change after manipulations in nutritional availability
(d). *Npy* expression was increased after food restriction ($F_{1,15}$ = 6.12; P < 0.05) but did not
change with SP exposure (e). Prolactin (*Prl*) expression in the pituitary gland was
downregulated in response to SP ($F_{1,20}$ = 28.53; P < 0.001), and insensitive to food
restriction. 12-weeks of SP were found to increase *Sst* expression in the hamster
mediobasal hypothalamus and levels were significantly reduced in response to a single
triiodothyronine (T3) injection (g). *P < 0.05 and **P < 0.01 denote significant difference
between SP and LP conditions (b). Letters denote significant difference between treatment
groups (P < 0.05) (c-g).

Figure 3. *Dio3* functions to time circannual interval duration in hamsters. Hamsters received intracerebroventricular injections to target *Dio3* expressing cells in the tanycytes localized to the mediobasal hypothalamus (a). Crispr-Cas9 constructs were packaged into lentiviral vectors to generate blank control hamsters (*Dio3^wt^*) or contain gRNAs that mutated the *Dio3* gene (*Dio3^cc^*). Hamsters were then exposed to SP conditions and the circannual interval timer was assessed by monitoring body mass (b). *Dio3^cc^* hamsters were slower to initiate interval timing as evidenced by higher body mass at 10-weeks SP exposure ($P < 0.001$) and recovered body mass quicker at 26 to 32 ($P < 0.05$) weeks. Lomb-Scargle analyses identified a significant reduction in the period of the circannual interval timer (Welch's t-test; $t_3 = 2.62$; $P < 0.05$) (c,d). Example *Dio3^cc^* (hamster #903) and *Dio3^wt^* (hamster #825) are presented in (c). A subpopulation of hamsters did not decrease body mass in response to SP exposure and termed non-responsive (NR) ($t_{12} = 7.12$; $P < 0.001$) (e). *Dio3* expression in the mediobasal hypothalamus of NR hamster was nearly nondetectable after exposure to 8-12 weeks of SP exposure compared to SP control hamsters ($t_{12} = -3.78$; $P < 0.005$) (f). *$P < 0.05$, **$P < 0.01$, and ***$P < 0.001$.

Fig.4. Schematic model for the mechanisms of circannual interval timing. (a) Hamsters held in long photoperiod (LP) have high body mass, thyrotropin-stimulating hormone beta (*Tshβ*) and low somatostatin (*Sst*) and deiodinase type-3 (*Dio3*). Transfer to short photoperiod (SP) results in the induction of circannual interval timing characterized by a reduction in body mass, low *Tshβ* in the pars tuberalis and increased *Dio3* expression in tanycytes. The maintenance phase is associated with increased *Sst* expression which serves to inhibit growth. The onset of the recovery phase is associated with a complete reversal in whole organismal physiology to the LP phenotype. (b) tanycytes along the third ventricle in the mediobasal hypothalamus integrate photic cues derived from *Tshβ* and are sensitive to nutritional cues. The triiodothyronine (T3) output signal from tanycytes regulates *Sst* expression leading to long-term programmed rheostatic changes in body mass that serve to control circannual interval timing in multiple physiological systems. Conversely, homeostatic stability is maintained despite large scale seasonal rhythms in body mass. Homeostatic energy balance is established by well characterized circuits that include neuropeptide Y (*Npy*), agouti-related peptide (*Agrp*) and the melanocortin receptor 4 (*Mc4r*).

Supplementary Figures:

732
Fig. S1. Plasma insulin reflects changes in transcript expression. (A) Plasma insulin of

Siberian hamsters across 32-weeks of short photoperiod exposure (H_8 = 20.401, P = 0.009).

(B) Expression of *Dio3* from the MBH of Siberian hamsters, assessed by qPCR, across 32-

weeks of short photoperiod exposure (H_8 = 39.771, P < 0.001). (C) Expression of *Sst* from

the MBH of Siberian hamsters, assessed by qPCR, across 32-weeks of short photoperiod

exposure (H_8 =16.002, P = 0.042). Letters denote significance between groups.

740
Fig. S2. Gene ontology analysis of MBH sequencing reveals well known seasonal

pathways. (A) Gene ontology enrichment analysis of significant transcripts from sequencing

of Djungarian hamster MBH across 32-weeks of short photoperiod exposure. (B) Interaction

network of enriched gene ontology terms. Size of individual points represent the number of

transcripts in each term. Thickness of interactions represents shared transcripts between

terms.

747
748

Fig. S3. Gene ontology analysis of pituitary gland unveils potential mechanisms for

seasonal changes in protein processing and release. (A) Gene ontology enrichment

analysis of significant transcripts from sequencing of Djungarian hamster pituitary gland

across 32-weeks of short photoperiod exposure. (B) Interaction network of enriched gene

ontology terms. Size of individual points represents the number of transcripts in each term.

Thickness of interactions represents shared transcripts between terms.

Fig. S4. Paraventricular and dorsomedial hypothalamus sequencing unveils novel transcripts and widespread seasonal interval timing within the hypothalamus. (A-B) Heatmaps of significant (FDR < 0.1) Djungarian hamster transcripts displaying sine or cosine rhythmicity were selected from (A) paraventricular hypothalamus and (B) dorsomedial hypothalamus. (C) Polar scatter chart of significant transcripts from Djungarian hamster paraventricular and dorsomedial hypothalamus, displaying peak of expression and -log(FDR).

765
766

768 **Fig. S5. Gene ontology analysis of paraventricular hypothalamic sequencing.** (A) Gene
769 ontology enrichment analysis of significant transcripts from sequencing of Djungarian
770 hamster paraventricular hypothalamus across 32-weeks of short photoperiod exposure.

771
772

Fig. S6. Gene ontology analysis of dorsomedial hypothalamus suggests widespread

immune involvement and cellular differentiation. (A) Gene ontology enrichment analysis

of significant transcripts from sequencing of Djungarian hamster dorsomedial hypothalamus

across 32-weeks of short photoperiod exposure. (B) Interaction network of enriched gene

ontology terms. Size of individual points represents the number of transcripts in each term.

Thickness of interactions represents shared transcripts between terms.

Fig. S7. Rheostatic mechanism controlling body mass change in Djungarian

hamsters. (A) Epidydimal fat mass in long- and short- photoperiod exposed Djungarian

hamsters ($F_{1,20}$ = 59.136; P < 0.001) after an overnight food restriction, or ad libitum, feeding

protocol ($F_{1,22}$ = 2.738; P = 0.114). (B) Pituitary gland expression of *Gh* from in long- and

short- photoperiod exposed Djungarian hamsters ($F_{1,20}$ = 2.738; P = 0.114) after an

overnight food restriction, or ad libitum, feeding protocol, assessed by qPCR ($F_{1,20}$ = 0.518; P

= 0.480). (C) Plasma insulin in long- and short-photoperiod exposed Djungarian hamsters

($F_{1,20}$ = 29.865; P < 0.001) after an overnight food restriction, or ad libitum, feeding protocol

($F_{1,22}$ = 6.863; P < 0.05). Letters denote significant main effect of photoperiod (a). Asterisk

indicates significant effect of photoperiod (P < 0.005), and hashtag denotes significant main

effect of food restriction (P < 0.05).

793
794

Fig. S8. Evidence of CRISPR modification in *Dio3^cc* and *Dio3^wt* Djungarian hamsters.

(A) Binding positions of gRNA1 and gRNA2, injected into Djungarian hamster hypothalamus

(B) Sequence of gRNA1 and gRNA2 binding and PAM sites. (C) Representative *Dio3*

sequence alignment with gRNA1 and PAM sites indicated for a representative *Dio3^wt*

(control) and *Dio3^cc* (CRISPR) hamster. (D) Decomposition error difference for *Dio3*

sequence alignment to determine genomic modification ($t_5 = 2.66$, $p < 0.05$) (E) Pelage

change in *Dio3^wt* and *Dio3^cc* Djungarian hamsters across 32 weeks of SP exposure, *Dio3^wt*

showed significantly increased pelage score at 13 ($P < 0.001$) and 14 ($p < 0.001$).

Gene	Primer	Size	Temp	Melt
Reference	-	-	-	-
18s	GCTCCTCTCCTACTTGGATAACTGTG CGGGTTGGTTTTGATCTGATAAATGCA	111	62˚C	80˚C
Hrpt	AGTCCCAGCGTCGTGATTAGTGATG CGAGCAAGTCTTTCAGTCCTGTCCA	141	62˚C	76˚C
Mediobasal hypothalamus				
Npy	CCAGGCAGAGATACGGCAAGAGATC CCATCACCACATGGAAGGGTCC	119	60˚C	81˚C
Sst	GAAGTCTCTGGCGGCTGCTG CAGCCTCATTTCATCCTGCTCCG	145	60˚C	85˚C
Dio3	CATGCTCCGCTCCCTGCTGCTTCA CAGGGTGCACAGACGGTTGTC	251	62˚C	85˚C
Pituitary Gland				
Prl	TCCGGAAGTCCTTCTGAACC CGCAGGCAGCGAATCTTATTG	300	60˚C	82˚C
Gh	ACCTACAAAGAGTTTGAGCGTG ATGAGCAGCAGCGAGAATCG	167	58˚C	85˚C

807 **Table S1.** Djungarian hamster qPCR primers sequences and thermal cycling profiles.
808

a)

LP | Induction | Maintenance | Recovery
 | SP

Body mass
Tshβ
Dio3
Npy
Sst
Time

b)

Mediobasal hypothalamus

3rdV

Rheostasis

Sst

Tanycytes

Energetic cues

Dio3

T3

Photic cues

Npy
AgRP
Mc4r

Homeostasis

Tshβ

Pars tuberalis

Systemic:
- body mass
- pelage

Adipose tissue:
- mass
- morphology

Pancreas:
- insulin