1

Title: Hypothalamic deiodinase type-3 establishes the period of circannual interval timing in mammals

Calum Stewart[1], T. Adam Liddle[1], Elisabetta Tolla[1], Jo Edward Lewis[2], Christopher Marshall[3], Neil P Evans[1], Peter J Morgan[4], Fran JP Ebling[5], Tyler J Stevenson[1]*.

Corresponding Author: Professor Tyler Stevenson

Email: tyler.stevenson@glasgow.ac.uk

Affiliations:

[1] School of Biodiversity, One Health, and Veterinary Medicine, University of Glasgow; Glasgow, G61 1QH, UK.

[2] Institute of Metabolic Science-Metabolic Research Laboratories & MRC-Metabolic Diseases Unit, University of Cambridge, Cambridge, UK.

[3] School of Physiology, Pharmacology and Neuroscience, University of Bristol; Bristol, BS8 1TD, UK.

[4] Rowett Institute, University of Aberdeen; Aberdeen, AB25 2ZD, UK.

[5] School of Life Sciences, University of Nottingham; Nottingham, NG7 2UH, UK.

Competing Interest Statement: Authors declare that they have no competing interests.

Abstract

Animals respond to environmental cues to time phenological events, but the intrinsic

mechanism of circannual timing remains elusive. We used transcriptomic sequencing and

frequent sampling, during three distinct phases (induction, maintenance and recovery) of

circannual interval timing for Djungarian hamster energy balance, to investigate the

molecular architecture of a neuroendocrine seasonal clock. Our study identified three distinct

phases of *transcript changes, with deiodinase type-3 (Dio3) expression activated during the*

early induction phase. Subsequent work demonstrated that targeted mutation of *Dio3*

resulted in a shorter period for circannual interval timing. Hamsters that exhibit naturally

disrupted *Dio3* expression do not show any change in body mass or pelage. Our work

demonstrates that changes in Dio3 induction is essential for setting the period of circannual

interval timing.

Introduction

Phenology of key life history traits is common across plant (Piao S *et al.*, 2019) and animal kingdoms (Cohen J, Lajeunesse MJ and Rohr JR, 2018). The annual changes in daylength are the predominant environmental cue which animals use to time seasonal life history transitions (Perez JH *et al.*, 2019; Wood S and Loudon A. 2014). Plants and animals also exhibit endogenous circannual timing in the absence of any change in environmental cues. For example, bird migration (Gwinner E and Dittami J, 1990), mammalian hibernation (Pengelley ET and Fisher KC, 1957), and reproduction (Woodfill CJ *et al.*, 1994; G *et al.*, 2006) are all driven by robust intrinsically generated circannual timers, the cycle of which nearly match a 12-month period. Some circannual timers estimate an interval period (e.g., 6-month) in which programmed changes in physiology and morphology occur in anticipation of the next season. Such interval timers are commonly observed as flowering in plants (Lincoln Duncan S *et al.*, 2015), diapause in insects (Denlinger DL, 1974) and spring emergence in rodents (Prendergast BJ, Renstrom RA and Nelson RJ, 2004). These interval timers are typically characterized by having light-dependent induction, maintenance and recovery phases.

Annual mammalian life history transitions are typically associated with major changes in energy demand (Ricklets RE, 1991). While the melanocortin system, including neuropeptide y (NPY), agouti-related peptide (AGRP), and melanocortin receptors are known to regulate short-term changes in energy homeostasis (Yeo GSH *et al.*, 2021), the neuropeptides implicated in long-term, circannual variation in energy rheostasis are not well characterized. Somatostatin (SST) (Marshall CJ *et al.*, 2024; Petri I *et al.*, 2016; Petri I *et al.*, 2014), proopiomelanocortin (POMC) (Bao R *et al.*, 2019; Mercer JG *et al.*, 2000), and VGF nerve growth factor (Barrett P *et al.*, 2005) are known to be strong correlates of seasonal variations in energetic state. Tanycyte somas are essential for the integration of environmental and physiological signals required for circannual interval timing. Tanycytes are localized in the ependymal layer of the third ventricle and are highly sensitive to nutrient state (Bolborea M

et al., 2020) and receive photoperiodic signaling derived from thyrotropes in the pars

tuberalis (Wood SH *et al.*, 2020; Hanon EA *et al.*, 2008). Previous work has established

deiodinase type-2 (*Dio2*) and type-3 (*Dio3*) expression is anatomically localized to tanycytes

and coordinate triiodothyronine-dependent annual transitions in physiological state (Petri I *et*

al., 2016; Bao R *et al.*, 2019, Wood SH *et al.*, 2020; Hanon EA *et al.*, 2008; Ebling FJ and

Lewis JE, 2018; Murphy M *et al.*, 2012). Here we delineate the molecular architecture of

circannual interval timing by the hypothalamus and pituitary gland and tested the conjecture

that *Dio3*, in the Djungarian hamster (*Phodopus sungorus*), is upregulated during the

induction of circannual interval timing for energy rheostasis, and functions to establish the

duration (or period) of the circannual interval timer.

Results

Male Djungarian hamsters were either kept in a long photoperiod (LP) control condition or

moved from LP to short photoperiod (SP) conditions (Fig.1). Djungarian hamsters remain in

LP phenotype unless exposed to SP. Pelage color, torpor and body mass were used to

monitor the induction, maintenance, and recovery phases of the circannual timer (Fig.1A and

C *SI Appendix* Movies S1 and 2). A full white pelage color was observed between 12-16

weeks after exposure to SP and gradually reversed to the LP agouti color after 28 weeks in

SP. Hamsters engaged in torpor between 12-20 weeks in SP (movie S2). Massive,

programmed changes in energy rheostasis were observed with a 30% reduction in daily food

intake and a 20% decrease in average body mass by 12 weeks SP (Fig.1C and D). Both

food intake and body mass started to reverse to LP conditions after 20 weeks in SP.

Epidydimal adipose tissue mass and plasma insulin concentrations paralleled the change in

body mass (Fig. 1D and *SI Appendix* Fig. S1). These reversals in physiological state are

indicative of the recovery phase of the circannual interval timer. GLP-1 did not display any

change in circulating concentrations (Fig. 1E). Plasma glucose concentration jumped sharply

after 16 weeks in SP (Fig. 1E), concurrent with the development of torpor in these animals

(movie S2). The observed change in body mass, food intake, adipose tissue and plasma

insulin versus the lack of change in GLP-1 highlights the physiological distinction between a

programmed rheostatic mechanism characteristic of the circannual interval timing of energy

stability in the former, while the latter are driven by short-term homeostatic mechanisms

(Stevenson TJ, 2024).

Figure 1. Molecular basis of circannual interval timing for morphology and physiology. Experimental design in which Djungarian hamsters were kept in long photoperiod (LP) or transferred to short photoperiod (SP) (a). SP induced pelage color change from LP agouti to winter white after 12-weeks which reversed to agouti by 28-weeks (F8, 91 = 50.77; P < 0.001) (b). Torpor was identified in hamster between 12- and 20-weeks SP exposure indicate by red line (b). SP exposure induced significant reduction in body mass (F8, 91 = 13.428; P < 0.001) (c), food intake (F8, 91 = 10.860; P < 0.001; denoted as downward triangles) (d) and adipose mass (F8, 91 = 27.929; P < 0.001; denoted as upward triangles) (d). Plasma GLP-1 did not vary in response to SP manipulation; denoted as upward triangles (e). SP exposure resulted in significant changes in plasma glucose around the onset of torpor; denoted as downward triangles (F8, 91 = 3.117; P < 0.05) (e). BioDare2.0 heatmaps of mediobasal hypothalamus (f) and pituitary gland (g) transcripts from Djungarian hamster collected at 4 -week SP intervals. Transcripts identified as highly rhythmic (FDR < 0.1) showed three distinct phases of expression that coincide with the induction, maintenance and recovery of circannual interval timing. Deiodinase type-3 (Dio3) was upregulated during the induction phase, whereas transcripts associated with energy

stability (e.g., somatostatin [Sst and Sstr5]) were upregulated during the maintenance phase.

All rhythmic transcripts reverted to the LP condition by 28-weeks SP exposure. Polar scatter

chart of significant transcripts from mediobasal hypothalamus and pituitary gland provide a

comprehensive seasonal clock for mammalian circannual interval timing across

neuroendocrine tissues (h). Green line indicates the induction phase, red line indicates the

maintenance phase, and the blue line represents the recovery phase. Letters denote

significant differences between treatment groups ($P < 0.05$) (b-e).

We then used Oxford Nanopore transcriptomic sequencing to characterize the molecular
changes associated with phases of the circannual interval timer within multiple individual
hypothalamic nuclei and the pituitary gland (Fig.1F to G; fig. *SI Appendix* S2 to 6 and
Dataset S1 to 4). The development of refractoriness to melatonin, indicative of the recovery
phase of the circannual timer, has been shown to develop independently in different
hypothalamic regions (Freeman DA and Zucker I, 2001). Molecular data was collected from
the mediobasal hypothalamus (Bolborea M *et al.*, 2020), dorsomedial hypothalamus (Ebling
FJP *et al.*, 2008), and the paraventricular nucleus (Bittman EL *et al.*, 1991), along with the
pituitary gland (Majumdar G *et al.*, 2023). This experimental approach provides a high-
throughput and high-frequency sampling resolution to comprehensively chart the induction,
maintenance and recovery of the mammalian circannual interval timer (Fig. 1H). Biodare2.0
identified 290 (Dataset S1) transcripts in the mediobasal hypothalamus as endogenously
rhythmic. Density heat-maps show an early, and robust upregulation of transcripts by 8
weeks SP associated with the circannual interval induction (Fig. 1F). Consistent with
previous reports (Petri I *et al.*, 2016; Bao R *et al.*, 2019; Yoshimura T *et al.*, 2003), Dio3
expression was significantly upregulated and clustered with the initial wave of transcript
expression (Fig.1F and *SI Appendix* Fig. S1). A second cluster of transcripts were
upregulated during the maintenance phase occurring between 12- and 20-weeks in SP
which primarily consisted of somatostatin (*Sst*) and the cognate receptor subtype 5 (*Sstr5*).
Increased *Sst* expression coincided with the reduction in body and adipose tissue mass
reflecting the conserved role in inhibiting growth and metabolism (*SI Appendix* Figs. S1 and
S2). Finally, a third wave of transcripts became upregulated during the recovery phase, and
largely resembled the molecular landscape in LP hamsters (Fig.1f). qPCR assays for *Dio3*
and *Sst* expression reflected the sequencing transcript count pattern providing independent
replication of the sequencing analyses (Fig. S1). In the pituitary gland, 250 transcripts were
identified as rhythmic and included genes associated with secretagogue cells such as
somatotrophs (*Gh*) and lactotrophs (*Prl*) (Fig.1g and *SI Appendix* Fig S3 and Dataset S2).
Similar rhythmic expression was identified in other hypothalamic nuclei including the
paraventricular nucleus (518 transcripts) (*SI Appendix* Fig. S4 and S5 and Dataset S3), and
the dorsomedial hypothalamus (374 transcripts) (Fig. S4 and *SI Appendix* S6 and Dataset
S4).

Dio3-dependent changes in triiodothyronine induced body mass loss via _Sst_

expression

Treatment of Djungarian hamster with the somatostatin agonist pasireotide decreases body

mass and is sufficient to drive changes similar to SP exposure (Dumbell RA *et al.*, 2015 and

2017). Further, certain SST receptor subtypes appear to be involved in driving seasonal

torpor in Djungarian hamsters (Scherbarth F *et al.*, 2014). To determine if hypothalamic *Sst*

expression reflects the programmed rheostatic change in energy state governed by

circannual interval timing or homeostatic cues, tissues were collected from *ad libitum* (AL)

fed hamsters under either LP control condition or the maintenance phase (12-week SP). To

induce a negative energetic state, hamsters experienced an acute overnight food restriction

(FR; 16hrs) (Fig. 2A). As anticipated, body mass decreased approximately 30% after

exposure to SP (Fig.2B, closed symbols); homeostatic negative energy state challenges by

food restriction reduced body mass by 1.5g on average for both LP (open symbols) and SP

conditions (Fig.2C). Epididymal adipose tissue mass was higher in LP compared to SP but

did not decrease in response to overnight FR (*SI Appendix* Fig. S7). Plasma insulin was

significantly reduced in response to SP treatment and FR suggesting circulating levels are

an output of both rheostatic and homeostatic signals (Fig. S7). The two manipulations permit

the ability to dissect the impact of rheostatic changes associated with circannual interval

timing from those involved in short-term homeostatic changes in energy balance.

Mediobasal hypothalamic *Sst* expression was significantly increased in SP conditions, but

insensitive to homeostatic challenges in energy state (Fig.2D). In contrast, mediobasal

hypothalamic *Npy* expression was significantly upregulated in food restricted hamsters but

was similar across LP and SP conditions (Fig.2E). *Prl* expression was higher in LP

compared to SP conditions (Fig. 2F). *Gh* expression did not change in response to either SP

induced circannual changes in energy state, or homeostatic manipulations (Fig. S7). To

establish whether mediobasal hypothalamic *Sst* expression is regulated by upstream *Dio3*-

dependent changes in local thyroid hormone catabolism, SP housed hamsters received sub-

cutaneous injections with either vehicle or triiodothyronine. A single triiodothyronine injection

in SP hamsters was sufficient to reduce *Sst* expression compared to vehicle controls (Fig.

2G). These data indicate SP induced *Dio3* expression removes T3-dependent inhibition of

Sst expression leading to long-term inhibition of body mass during the maintenance phase of

circannual interval timing of rheostatic energy balance.

Figure 2. Somatostatin expression reflects programmed circannual interval timing that is dependent on Dio3 regulation of local triiodothyronine signaling. The experimental design in which SP exposure induced rheostatic reduction in body mass after 12-weeks, and then hamsters experienced either an overnight food restriction (FR) or maintained food ad libitum (AL) (a). SP induced a significant reduction in body mass ($F_{1, 22}$ = 3.85; $P < 0.01$) (b). Food restriction further reduced body mass ($F_{1, 22}$ = 25.46; $P < 0.001$) (c). Sst expression in the mediobasal hypothalamus was significantly increased in SP ($F_{1, 15}$ = 5.59; $P < 0.05$) but did not change after manipulations in nutritional availability (d). Npy expression was increased after food restriction ($F_{1, 15}$ = 6.12; $P < 0.05$) but did not change with SP exposure (e). Prolactin (Prl) expression in the pituitary gland was downregulated in response to SP ($F_{1, 20}$ = 28.53; $P < 0.001$), and insensitive to food restriction. 12-weeks of SP were found to increase Sst expression in the hamster mediobasal hypothalamus and levels were significantly reduced in response to a single triiodothyronine (T3) injection (g). *$P < 0.05$ and **$P < 0.01$ denote significant difference between SP and LP conditions (b). Letters denote significant difference between treatment groups ($P < 0.05$) (c-g).

Dio3 disfunction reduces the period of circannual interval timing

Next, we sought to establish the functional role of *Dio3* signaling for circannual interval

timing of energy state in hamsters. Targeted genomic mutations of the *Dio3* gene localized to

the MBH were achieved via intracerebroventricular (ICV) injection of CRISPR-Cas9 (*Dio3cc*;

Fig. S8) to assess the functional role for circannual interval timing. Control hamsters were

administered a blank crispr-cas9 construct (*Dio3wt*). Two weeks following surgery, hamsters

were transferred to SP and programmed circannual changes in body mass (Fig.3B) and

pelage monitored (*SI Appendix* Fig. S8). *Dio3cc* hamsters delayed body mass loss and

regained body mass significantly quicker during the recovery phase, compared to *Dio3wt*

controls. Similarly, the change in pelage color occurred later in *Dio3cc* compared *Dio3wt*,

and *Dio3cc* was never observed to develop full white pelage (*SI Appendix* Fig. S8). Lomb-

Scargle period analysis of body mass identified that *Dio3cc* hamsters have notably shorter

duration (period = 26.959) compared to *Dio3wt* (period = 32) (Figs. 3C and D). A

subpopulation of hamsters was physiologically nonresponsive (NR) to the SP manipulation

and maintained the higher LP body mass (Fig.3E). *Dio3* expression in the mediobasal

hypothalamus of these NR hamsters remained exceptionally low compared to responsive SP

hamsters at 8- and 12-weeks following exposure to SP (Fig.3F). These data suggest that the

inability to increase *Dio3* expression in NR hamsters acts as a form of naturally occurring

disfunction that prevents SP induction of circannual interval timing.

Figure 3. Dio3 functions to time circannual interval duration in hamsters. Hamsters

received intracerebroventricular injections to target Dio3 expressing cells in the tanycytes

localized to the mediobasal hypothalamus (a). Crispr-Cas9 constructs were packaged into

lentiviral vectors to generate blank control hamsters (*Dio3wt*) or contain gRNAs that mutated

the Dio3 gene (Dio3cc). Hamsters were then exposed to SP conditions and the circannual

interval timer was assessed by monitoring body mass (b). *Dio3cc* hamsters delayed interval

timing as evidenced by higher body mass at 10-weeks SP exposure (P < 0.001) and

recovered body mass quicker at 26 to 32 (P < 0.05) weeks. Lomb-Scargle analyses

identified a significant reduction in the period of the circannual interval timer (t6 = 6.975; P <

0.05) (c,d). Example Dio3cc (hamster #903) and Dio3wt (hamster #825) are presented in (c).

A subpopulation of hamsters did not decrease body mass in response to SP exposure and

termed non-responsive (NR) (t12 = 7.12; P < 0.001) (e). *Dio3* expression in the mediobasal

hypothalamus of NR hamster was nearly nondetectable after exposure to 8-12 weeks of SP

exposure compared to SP control hamsters (t12 = -3.78; P < 0.005) (f). *P < 0.05, **P <

0.01, and ***P < 0.001.

Discussion

Here, we sequenced the transcriptomes of three hypothalamic nuclei and the pituitary gland, key neuroendocrine structures which govern programmed circannual rheostatic and homeostatic processes. Our approach provided a high sampling frequency of the circannual interval timer in Djungarian hamsters. The findings produced a robust and comprehensive neural and molecular database which facilitates the delineation of a seasonal clock in mammals. Several novel transcripts displayed a close correlation with the induction phase of the seasonal response. Among these was *Dio3*, conducive with the removal of thyroid hormone (e.g., thyroxine, triiodothyronine) being a key step in the transition to the winter phenotype. The photoinduced change in *Dio3* expression induces a long-term change in physiology and morphology and our results demonstrate that it is insensitive to short-term alterations in response to other environmental cues (e.g., nutritional state). Functional manipulation of *Dio3* expression also delayed the induction and enhanced the reversal of the winter phenotype suggesting that upregulation of *Dio3* acts to establish the period of circannual interval timing.

The transcriptomic data are consistent with evidence that shows *Sst* expression is one critical event for the maintenance of a reduced rheostatic energy state during the winter season (Marshall CJ *et al.*, 2024; Petri I *et al.*, 2016 and 2014). In hamsters, *Sst* expression is upregulated after Dio3 induction and is associated with the low body mass from week 12-20 before a downregulation in transcript levels coinciding with increased body mass. By increasing circulating triiodothyronine in hamsters maintained in SP conditions, our manipulation indicates that elevated hormone concentrations lead to the reduction in *Sst* expression in the mediobasal hypothalamus (Fig.2). The overall expression patterns provide the ability to develop a linear series of steps in which LP conditions have high local concentrations of triiodothyronine in the mediobasal hypothalamus (Murphy M *et al.*, 2012; Helfer G and Stevenson TJ, 2020) which reduces the levels of *Sst* expression resulting in consistently higher body mass. Exposure to SP induces *Dio3*, which catabolizes

triiodothyronine leading to the removal of the inhibition of *Sst* expression and reduction in

body mass. Local synthesis of triiodothyronine in the mediobasal hypothalamus by tanycytes

is an evolutionary conserved process for timing photoperiod-induced transitions across the

animal kingdom (Lewis JE and Ebling FJP, 2017; Ebling FJP and Lewis JE, 2018).

Photoperiodic signals derived from the pars tuberalis via thyrotropin-stimulating hormone are

essential to regulate *Dio3* (and *Dio2*) expression in tanycytes leading to increased

triiodothyronine content in LP conditions for mammals (22,23), birds (Nakao N *et al.*, 2008)

and fish (Lorgen M *et al.*, 2015). The current limitation is understanding how SP signals

regulate Dio3 expression. Undoubtedly, the duration of melatonin secretion during the daily

nocturnal phase is essential (Helfer G, Barrett P and Morgan PJ, 2019). Future research is

necessary to identify the conserved upstream signals that govern the timing of *Dio3*

expression.

These data support the existence of a neuroendocrine pathway for the long-term rheostatic

regulation, and another for short-term homeostatic control of energy stability. The rheostatic

pathway consists of triiodothyronine (T3) signaling dependent on a DIO3 switch to regulate

Sst expression in the mediobasal hypothalamus whereas short-term regulation of energy

stability via nutrient availability is homeostatically regulated by well-established orexigenic

(e.g., *Npy*) and anorexigenic neuropeptides (*Pomc*) (Yeo GSH *et al.*, 2021). In response to

the output of both pathways, peripheral effects may be mediated by acute and chronic

changes in pancreatic insulin secretion (*SI Appendix* Fig. S9). Overall, the findings provide a

clear neural and cellular circuit for circannual interval timing of energy stability and

demonstrate *Dio3* as a gene critical for the control of seasonal life history transitions.

Materials and Methods

Ethics

All procedures were in accordance with the National Centre for the Replacement,

Refinement and Reduction of Animals in Research ARRIVE guidelines

(https://www.nc3rs.org.uk/revision-arrive-guidelines). All procedures were approved by the

Animal Welfare and Ethics Review Board at the University of Glasgow and conducted under

the Home Office Project License PP5701950.

Animals

All animals were obtained from a colony of Djungarian hamsters (3-8 months of age) kept at

the University of Glasgow, Veterinary Research Facility. Animals were held at a room

temperature of 21°C, and a humidity of 50%. The colony room was kept on a long

photoperiod (16L:8D). All hamsters had food (SDS [BK001]) and water *ad libitum*.

Experiment 1 – Defining the circannual interval timer in hamsters

Adult male hamsters (N = 100) were required to assess the physiological and molecular

basis of circannual interval timing in hamsters. A photoperiod reference group were kept on

LP (n=14) (8D:16L) conditions for the duration of the study. The SP treatment groups

consisted of hamsters moved to SP (16D:8L) at 4 weeks intervals. The groups consisted of

SP for 4 (n=12), 8 (n=12), 12 (n=12), 16 (n=11), 20 (n=12), 24 (n=10), 28 (n=7) and 32

(n=10) weeks. This photoperiod schedule reliably captures the induction, maintenance, and

recovery of circannual interval timing in hamsters (Prendergast BJ *et al.*, 2002). Pelage

score was determined using visual assessment and defined scale in which a score of 1 is

the summer dark agouti coat; a score of 2 is a transition between agouti and white; and a

score of 3 is a full winter white pelage (Marshall CJ *et al.*, 2024). Body mass was monitored

to the nearest 0.1g using an Ohaus portable scale (TA301). Food intake was measured by

weighting the food pellets at the mid-point on two consecutive days to obtain a daily value.

Food weight was measured using an Ohaus scale (TA301). Hamsters were killed between 2-

4hr after lights on by cervical dislocation followed by exsanguination. Adipose tissue was

immediately dissected and measured using a Sartorius cp64 anatomical balance. A terminal

blood sample was collected to measure glucose (Accu-check Performa nano blood glucose

meter). Circulating levels of insulin and total glucagon-like peptide 1 (GLP1) using ELISA

(MesoScale Discovery, UK) at Core Biochemical Assays laboratories, Cambridge, UK. Brain

and pituitary gland tissue was quickly extracted and frozen on powdered dry ice and stored

at -70°C. Epididymal white adipose tissue was dissected and weighed using Sartorius cp64

anatomical balance and stored at -70°C.

Brains were cut into 200μM coronal sections using a Leica CM1520 cryostat. Anatomical

structures (optic tract to the infundibular stem; Bregma -2.12mm to -3.80mm) were used to

isolate the mediobasal hypothalamus (MBH), dorsomedial hypothalamus (DMH) and the

paraventricular nucleus (PVN). Bilateral tissue punches were performed using an integra

Miltex 1mm disposable biopsy punch. Tissue punches were stored at -70°C until

transcriptomic sequencing and confirmatory qPCR analyses (see below).

Experiment 2 – Dissociating mechanisms that govern circannual rheostatic versus

homeostatic energy stability.

A total of N=35 adult male hamsters were used to determine molecular signatures of

rheostatic versus homeostatic energy stability. A LP control group was kept on long

photoperiod for the duration of the study (n = 18). A subset of hamsters was moved to SP (n

= 17) for 12 weeks to obtain animals in the maintenance phase of circannual interval timing.

Food (SDS [BK001]) and tap water were provided *ad libitum* to both LP and SP animals for

the 12 weeks duration. Body mass was measured every two weeks using a Traveler Ohaus

portable balance. On the final night of the experiment, 50% of hamsters (i.e., LP = 9, SP = 9)

were kept on food and water *ad libitum*. The other 50% of hamsters (i.e., LP = 9, SS = 8)

served as the homeostatic treatment group and received an acute food restriction by

removing all food. Overnight food restriction induces a robust negative energy state that

reliably results in a 1–2-gram loss in hamster body mass (Bao R *et al.*, 2019). Body mass

was measured prior to food restriction, and again before tissue collection on the subsequent

day. Hamsters were killed between 2-4hr after lights on by cervical dislocation and then

exsanguination. A terminal blood sample was collected to measure insulin (MesoScale

Discovery, Core Biochemical Assays laboratories, Cambridge, UK). Body mass and

epididymal white adipose tissue mass were measured using a Sartorius cp64 anatomical

balance. Brains and pituitary glands were dissected and were immediately placed on dry ice

and stored at -70°C. Brains were cut into 200µM sections using a Leica CM1520 cryostat.

Anatomical structures (optic tract to the infundibular stem; Bregma -2.12mm to -3.80mm)

were used to isolate the MBH. Bilateral tissue punches were performed using an integra

Miltex 1mm disposable biopsy punch. Brain tissue punches and pituitary gland samples

were stored at -70°C until qPCR assay (see below).

Experiment 3 – Sufficiency of triiodothyronine to induce LP neuroendocrine state.

Adult male hamsters (n = 35) were used to determine the impact of triiodothyronine on *Sst*

expression. Hamsters were transferred from LP to SP for 12 weeks (n = 24) to initiate

circannual interval timing and establish the maintenance phase. Hamsters were divided into

two groups for the last week of SP treatment. A SP reference group received subcutaneous

injections of 0.9% w/v saline for 7 days (n = 10). To determine the sufficiency of

triiodothyronine to regulate *Sst* expression, hamsters (n=11) were administered a single

subcutaneous injection of 5µg/100ul triiodothyronine (Merck 102467157) prior to lights off on

the final night. Subcutaneous injections of triiodothyronine are well established to regulate

transcript expression in the hamster hypothalamus (Bao R *et al.*, 2019; Banks R, Delibegovic

M and Stevenson TJ, 2016; Stevenson TJ *et al.*, 2014). The subsequent day hamsters were

sacrificed by cervical dislocation followed by exsanguination between 4-5hr after lights on.

Terminal body mass and epididymal adipose tissue mass were recorded using a Sartorius

cp64 anatomical balance. Brains were dissected and stored immediately at -70°C until

sectioning. Brains were cut into 200µM sections using a Leica CM1520 cryostat. Anatomical

structures (optic tract to the infundibular stem; Bregma -2.12mm to -3.80mm) were used to

isolate the MBH. Bilateral tissue punches were performed using an integra Miltex 1mm

disposable biopsy punch. Tissue punches were stored at -70°C until qPCR assays (see

below).

Experiment 4 – Functional manipulation of *Dio3* expression in hamsters

Targeted neuroanatomical injections, using highly precise Guide RNA (gRNA) against *Dio3*

and Crispr-Cas9 vectors was essential to ensure the long-term inhibition of *Dio3* function in

the MBH in hamsters; an approach necessary to reliably examine the functional significance

of *Dio3* for the duration of circannual interval timing. Custom-made Crispr-cas9 constructs

were packaged in a lentiviral vector by Merck Life science UK. The lentiviral vector used was

U6-gRNA:ef1a-puro-2A-Cas9-2A-tGFP (Sigma All-in-One vector). gRNA for *Dio3* were

designed using CHOPCHOP (43). Three gRNA were designed to target the *Dio3* gene

(gRNA1: CGACAACCGTCTGTGCACCCTGG; gRNA2:

GTTCCCGCGCTTCCTAGGCACGG, and gRNA3: GACCCAGCCGTCGGATGGGTGGG).

Only gRNA 1 and 2 were identified to align with 5' end of the *Dio3* gene and were predicted

to align with chromosome 12 at locations 110279543 and 110279375 respectively (fig. S8).

The efficiency for gRNA1 was 61.24 and gRNA2 was 57.4. To check specificity of gRNA

sequences, we conducted BLAST using the NCBI dataset. Both gRNA1 and gRNA2 were

found to align 100% with *Dio3* and did not align 23/23 with any other region.

Adult male hamsters (n = 10) were selected to assess the impact of *Dio3* genome

modification on circannual interval timing. Hamsters were kept on LP conditions to maintain

the summer phenotype. Hamsters were then divided into two treatment groups. The control

group included hamsters (n=6) that received intracerebroventricular (ICV) injections of blank

Crispr-Cas9 constructs that were packaged into a lentivirus (*Dio3wt*; U6-gRNA:ef1a-puro-

2A-Cas9-2A-tGFP (Sigma All-in-One vector). The treatment group consisted of hamsters

(n=4) that received an ICV injection of Crispr-cas9 constructs that harbored both gRNA1 and

gRNA2 (*Dio3cc*).

Intracerebroventricular injections provide the ability to induce neuroanatomically localized

manipulations in *Dio3* gene function in the adult brain. Crispr-Cas9 constructs were delivered

via stereotaxic ICV injection into the third ventricle to target the MBH tanycytes. Under

general anesthesia (5% induction, 2% maintenance of isoflurane mixed with oxygen) a SGE

series II syringe was positioned along the midline at Bregma -1.5mm, to a depth of -5.7mm.

The anatomical coordinates were refined based on estimates taken from adult hamster

brains (Steward CA *et al.*, 2003) and tested using fresh hamster cadavers. Analgesia was

administered subcutaneously (5mg/kg Rimadyal and 0.1mg/kg buprenorphine). A small

~1cm incision was made by scalpel to expose the skull and a small hole was drilled using a

Bone Micro Drill for Brain surgery (Harvard Apparatus). Lentivirus was administered using a

Pump 11 Elite Programmable Syringe injection system (Harvard Apparatus) and 1ul of viral

vector (2-2.5 E+13 vg/ml) was delivered over 10 minutes at a rate of 0.1ul per minute. To

allow viral diffusion and prevent backflow the syringe was kept in place after injection for 1-

minute and then the syringe was raised by 1mm and kept in place for another 1-minute.

Animals were then moved to heated home cages and given mashed food to aid recovery.

Hamsters were provided with 2-weeks recovery and were monitored to ensure stable body

mass, locomotor activity and continued food and water intake.

To assess the impact of *Dio3cc* on circannual interval timing hamsters were moved to SP for

32 weeks and body mass and pelage were measured biweekly. Pelage score was

determined using visual assessment and defined scale in which a score of 1 is the summer

dark agouti coat; a score of 2 is a transition between agouti and white; and a score of 3 is a

full winter white pelage (Marshall CJ *et al.*, 2024). Body mass was monitored to the nearest

0.1g using an Ohaus portable scale (TA301). After 32 weeks, hamsters were killed 4-5hr

after lights on by cervical dislocation followed by exsanguination. Brains were dissected and

stored immediately at -70°C. Brains were cut into 200µM sections using a Leica CM1520

cryostat. Anatomical structures (optic tract to the infundibular stem; Bregma -2.12mm to -

3.80mm) were used to isolate the MBH. Bilateral tissue punches were performed using an

integra Miltex 1mm disposable biopsy punch. Tissue punches were stored at -70°C. DNA

was extracted from dissected tissues using the DNeasy Blood & Tissue Kit (QIAGEN) as per

manufacturer's instructions. Isolated DNA was sequenced using the Eurofins LightRun

sanger sequencing service. The forward primer provided to Eurofins was

CATGCTCCGCTCCCTGCTGCTTCA (5'-3'). CRISPR modified transcripts were aligned to a

reference wild type transcript (animal #821) using the Tracy sanger basecaller and aligner

(Rausch *et al.*, 2020). Alignments were visualized using the Ugene integrated bioinformatics

tool (Okonechnikov K, Golosova O and Fursov M, 2012). In case mutation was not obvious,

an additional analysis was carried out using the associated online resource SAGE

(https://www.gear-genomics.com/), to investigate the decomposition error of the modified

transcripts with the wild-type control (821). Four *Dio3wt* hamsters had *Dio3* sequences that

fully matched the reference genome (fig. S8c). Two hamster *Dio3* sequences did not

automatically align, and the raw reads were visually inspected. No mutations or *sequence*

mutations were detected in the raw sequence trace. Three *Dio3cc* treated hamsters had

Dio3 sequences with evidence of genomic mutation evidenced by low decomposition error

(*SI Appendix* Figs. S8C and D). One *Dio3cc* treated hamster did not show any genomic

mutation (#905) and was removed from the final analysis. Hamster #905 had a body mass

circannual interval period of 32 and we propose the animal represents a false positive

control.

Experiment 5 – Naturally occurring variation in *Dio3* expression and photoperiodic

response.

A subset of hamsters does not physiologically respond to SP treatment and are classified as

photoperiodically non-responsive (NR) (Przybylska-Piech AS and Jefimow M, 2022). Adult

hamsters (n = 14) were held in LP or SP for 8 weeks. A group of male non responders (n =

4) were identified due to the absence of body mass and pelage change in response to SP

(Fig. 3e). A reference group of male SP responders (n = 10) were collected. MBH was

extracted, RNA isolated and cDNA synthesized. qPCR was carried out for *Dio3* and *Sst*

expression. Hamsters were killed between 2-4hr after lights on.

RNA extraction

RNA was extracted using the Qiagen RNEasy plus mini kit. Tissues were homogenized

using a Polytron PT 1200 E. RNA was then extracted from homogenized tissue following the

manufacturer's instructions. RNA was tested for quality and quantity using an ND-1000

Nanodrop.

Next Generation Sequencing and data processing

Oxford Nanopore Sequencing was used to carry out transcriptomic sequencing. RNA from

MBH, DMH and PVN samples was sequenced using the PCR-cDNA barcoding kit following

manufacturer's instructions (SQK-PCB109; Oxford Nanopore). RNA from the pituitary gland

samples was synthesized using the direct cDNA sequencing kit following manufacturer's

instructions (SQK-DCS109 with EXP-NBD104; Oxford Nanopore). Bioinformatic analysis

was carried out on Linux within a conda environment. Raw reads (Fast5) were demultiplexed

and basecalled using guppy basecaller (4.2.1). Porechop (0.2.4) was used to remove

adapters from reads and Filtlong (v0.2.0) was used to filter for quality and read length

(length: >25 base pairs, quality score > 9). Transcripts were aligned to Mus musculus

genome (GRCm39) using Minimap2 (Li H, 2018). Previously, rodent genomes have been

demonstrated to show significant similarity, such that cross-species transcriptomic analysis

appears feasible (Przybylska-Piech AS and Jefimow M, 2022). Transcript expression levels

were generated using Salmon (v0.14.2) (Patro R *et al.*, 2017) and EdgeR (v3.24.3)

(Robinson MD, McCarthy DJ and Smyth GK, 2010) was used to filter lowly expressed

transcripts.

cDNA synthesis and qPCR

RNA was transformed into cDNA for qPCR analysis using superscript III (Invitrogen) as per

manufacturer's instructions. Quantification of transformed cDNA was achieved using Brilliant

II SYBR Mastermix (Agilent). Stock forward and reverse primers (100pmol/µl) were mixed

and diluted in nuclease free water to 20pmol/µl. A working SYBR mixture was prepared by

mixing 1-part primer mixture per 24-part SYBR mastermix. Reaction mixtures were prepared

in wells on a 96-well plate by mixing 4.8µl of normalised sample and 4.8µl of SYBR working

mix. All reactions were performed in duplicate. Primer and qPCR parameters are outlined in

Table S1. qPCR reactions were carried out in a Stratagene Mx3000P thermal cycler. Cycling

conditions utilized were in sequence; at 95°C for 5 minutes (denaturing), 40 cycles at (95°C

for 30 seconds, X°C for 1 minute – see table, 72°C for 30 seconds with fluorescent

measurement at end), 95°C for 1 minute, 55°C for 30 seconds increasing to 95°C (Melt

curve analysis). Melt cure analysis was used to determine specificity of amplification. Data

was analyzed for meanCT, efficiency and variability using PCR Miner (Zhao S and Fernald

RD, 2005). Logfold CT was calculated using the ΔΔCT method using 18s and/or *Hrpt*

reference transcripts as these transcripts are stable across photoperiodic conditions (Stewart

C, Liddle TA and Stevenson TJ, 2022).

Statistical Analysis

Raw data is provided in Data file 1 and sequencing data is provided in Table S1-S4. Physiological and qPCR data sets were tested for normality using the Shapiro test and transformed by log transformation if normality was violated. Statistical significance was determined using a two-way ANOVA. Post hoc analysis of significant data was achieved by t-test. Type 1 error was minimized by utilizing the Bonferroni adjustment on post-hoc analyses. Data were analyzed using non-linear regression for rhythmicity using the online resource BioDare 2.0 (Zielinski T *et al.*, 2014) (biodare2.ed.ac.uk). The empirical JTK_CYCLE method was used for detection of rhythmicity and the classic BD2 waveform set was used for comparison testing. Rhythmicity was determined by a Benjamini-Hochberg controlled false discovery rate (BH corrected FDR < 0.1). The false discovery rate of this data set was controlled by applying the Benjamini-Hochberg method (Benjamini Y and Hochberg Y, 1995). Clusters within the data set were identified using the gap-statistic and clustered using k-means clustering. Gene ontology analysis was carried out using ShinyGO v0.77 (Ge SX, Jung D and Yao R, 2020).

Data and materials availability

All data are available in Dataset S1-S5. The code is available via github repository (https://github.com/CS-Gla/Hamster_seasonal_sci_cs_2024). Sequencing information was deposited in GEO (GSE274003).

Acknowledgments

The authors thank Gerald Lincoln for helpful discussions which contributed to the conceptualization of the research. The authors thank the technical assistance provided by Nicola Munro, Ana Monterio, and Fallon Cuthill. Biorender.com was used for icons used in

528   Fig. 1-3 and fig. S9. Funding was provided via Leverhulme Trust award LT-RL-2019-06 (TJS)

529   and ISSF Wellcome Trust award (TJS, PJM, FJPE).

References

1.   S. Piao, Q. Liu, A. Chen, I.A. Janssens, Y. Fu, J. Dai, L. Liu, X. Lian, M. Shen, X. Zhu, Plant phenology and global climate change: current progresses and challenges. *Global Change Biology* **25**, 1922-1940 (2019).

2.   J. Cohen, M.J. Lajeunesse, J.R. Rohr, A global synthesis of animal phenological responses to climate change. *Nature Climate Change* **8**, 224-228 (2018).

3.   J.H. Perez, E. Tolla, I.C. Dunn, S.L. Meddle, T.J. Stevenson, A comparative perspective on extra-retinal photoreception. *Trends Endocrinol Metab* **30**, 39-53 (2019).

4.   S. Wood, A. Loudon, Clocks for all seasons: unwinding the roles and mechanisms of circadian and interval timers in the hypothalamus and pituitary. J*. Endocrinol.* **222**, R39-R59 (2014).

5.   E. Gwinner, J. Dittami, Endogenous reproductive rhythms in a tropical bird. *Science* **249**, 906-908 (1990).

6.   E.T. Pengelley, K.C. Fisher, Onset and cessation of hibernation under constant temperature and light in the Golden-mantled ground squirrel (Citellus lateralis). *Nature* **180**, 1371-1372 (1957).

7.   C.J. Woodfill, N.L. Wayne, S.M. Moenter, F.J. Karsch, Photoperiodic synchronization of a circannual reproductive rhythm in sheep: identification of season-specific time cues. *Biol Reprod.* **50**, 965-976 (1994).

8.   G. Lincoln, I.J. Clarke, R.A. Hut, D.G. Hazlerigg, Characterizing a mammalian circannual pacemaker. *Science* **314**, 1941-1944 (2006).

9.   S. Duncan, S. Holm, J. Questa, J. Irwin, A. Grant, C. Dean, Seasonal shift in timing of vernalization as an adaptation to extreme winter. *eLife* **23**, e06620 (2015).

10.   D.L. Denlinger, Diapause potential in tropical flesh flies. *Nature* **252**, 223-224 (1974).

11.   B.J. Prendergast, R.A. Renstrom, R.J. Nelson, Genetic analyses of a seasonal

interval timer *J. Biol. Rhythms* **19**, 298-311 (2004).

12.   R.E. Ricklets, Structures and transformations of life histories. *Func. Ecol.* **5**, 174-183

(1991).

13.   G.S.H. Yeo, D.H.M. Chao, A.M. Siegert, Z.M. Koerperich, M.D. Ericson, S.E.

Simonds, C.M. Larson, S. Luquet, I. Clarke, S. Sharma, K. Clement, M.A. Cowley, C.

Haskell-Luevano, L. Van Der Ploeg, R.A.H. Adan, The melanocortin pathway and energy

homeostasis: from discovery to obesity therapy. *Mol. Metabol.* **48**, 101206 (2021).

14.   C.J. Marshall, A. Blake, C. Stewart, T.A. Liddle, I. Denizli, F. Cuthill, N.P. Evans, T.J.

Stevenson, Prolactin mediates long-term, seasonal rheostatic regulation of body mass in

female mammals. *Endocrinol.* **165**, bqae020 (2024).

15.   I. Petri, V. Diedrich, D. Wilson, J. Fernandez-Calleja, A. Herwig, S. Steinlechner, P.

Barrett, Orchestration of gene expression across the seasons: hypothalamic gene

expression in natural photoperiod throughout the year in the Siberian hamster. *Sci Rep.* **6**,

29689 (2016).

16.   I. Petri, R. Dumbell, F. Scherbarth, S. Steinlechner, P. Barrett, Effect of exercise on

photoperiod-regulated hypothalamic gene expression and peripheral hormones in the

seasonal Dwarf Hamster Phodopus sungorus. *Plos One* **9**, e90253 (2014).

17.   R. Bao, K.G. Onishi, E. Tolla, F.J.P. Ebling, J.E. Lewis, R.L. Anderson, P. Barrett, B.J.

Prendergast, T.J. Stevenson, Genome sequencing and transcriptome analyses of the

Siberian hamster hypothalamus identify mechanisms for seasonal energy balance. *Proc.*

Natl Acad. Sci. **116**, 13116-13121 (2019).

18.   J.G. Mercer, K.M. Moar, A.W. Ross, N. Hoggard, P.J. Morgan, Photoperiod regulates

arcuate nucleus POMC AGRP, and leptin receptor mRNA in Siberian hamster hypothalamus.

Am. *J. Physiol. Reg. Inte. Comp. Physiol.* **278**, R271-R281 (2000).

19. P. Barrett, A.W. Ross, A. Balik, P.A. Littlewood, J.G. Mercer, K.M. Moar, T. Sallmen, J.

Kaslin, P. Panula, S. Schuhler, F.J. Ebling, C. Ubeaud, P.J. Morgan, Photoperiodic regulation

of histamine H3 receptor and VGF messenger ribonucleic acid in the arcuate nucleus of the

Siberian hamster. *Endocrinol* **146**, 1930-1939 (2005).

20. J.E. Lewis, F.J.P. Ebling, Tanycytes as regulators of seasonal cycles in

neuroendocrine function. *Front. Neurol.* **8**, 79 (2017).

21. M. Bolborea, E. Pollatzek, H. Benford, T. Sotelo-Hitschfeld, N. Dale, Hypothalamic

tanycytes generate acute hyperphagia through activation of the arcuate neuronal network*.*

Proc. Natl Acad Sci **117**, 14473-14481 (2020).

22. S.H. Wood, M.M. Hindle, Y. Mizoro, Y. Cheng, B.R.C. Saer, K. Miedzinska, H.C.

Christian, N. Begley, J. McNeilly, A.S. McNeilly, S.L. Meddle, D.W. Burt, A.S.I. Loudon,

Circadian clock mechanism driving mammalian photoperiodism. *Nature Commun.* **11**, 4291

(2020).

23. E.A. Hanon, G.A. Lincoln, J.M. Fustin, H. Dardente, M.Masson-Pevet, P.J. Morgan,

D.G. Hazlerigg, Ancestral TSH mechanism signals summer in a photoperiodic mammal. *Curr*

Biol **18**, 1147-1152 (2008).

24. F.J. Ebling, J.E. Lewis, Tanycytes and hypothalamic control of energy metabolism.

Glia. **66**, 1176-1184 (2018).

25. M. Murphy, P.H. Jethwa, A. Warner, P. Barrett, K.N. Nilaweera, J.M. Brameld, F.J.

Ebling, Effects of manipulating hypothalamic triiodothyronine concentrations on seasonal

body weight and torpor cycles in Siberian hamsters. *Endocrinol.* **153**, 101-112 (2012).

26. T.J. Stevenson, Defining the brain control of physiological stability. *Horm Behav.*

(2024).

27. D. A. Freeman, I. Zucker, Refractoriness to melatonin occurs independently at

multiple brain sites in Siberian hamsters. *Proc. Natl. Acad. Sci.* **98**, 6447–6452 (2001).

28.   F. J. P. Ebling, D. Wilson, J. Wood, D. Hughes, J. G. Mercer, P. J. Morgan, P. Barrett,

The Thyrotropin-Releasing Hormone Secretory System in the Hypothalamus of the Siberian

Hamster in Long and Short Photoperiods. *J Neuroendocrinol* **20**, 576–586 (2008).

29.   E. L. Bittman, T. J. Bartness, B. D. Goldman, G. J. DeVries, Suprachiasmatic and

paraventricular control of photoperiodism in Siberian hamsters. *Amer. J. Physiol. Reg Intgr.*

Comp. Physiol. **260**, R90–R101 (1991).

30.   G. Majumdar, T.A. Liddle, C. Stewart, C.J. Marshall, M. Bain, T.J. Stevenson. FSH

links photoperiodic signaling to seasonal reproduction in Japanese quail. *eLife* **12**, RP87751

(2023).

31.   T. Yoshimura, S. Yasuo, M. Watanabe, M. Iigo, T. Yamamura, K. Hirunagi, S. Ebihara,

Light-induced hormone conversion of T4 to T3 regulates photoperiodic response of gonads

in birds. *Nature* **426**, 178-181 (2003).

32.   R. A. Dumbell, F. Scherbarth, V. Diedrich, H. A. Schmid, S. Steinlechner, P. Barrett,

Somatostatin Agonist Pasireotide Promotes a Physiological State Resembling Short-Day

Acclimation in the Photoperiodic Male Siberian Hamster (Phodopus sungorus). *J*

Neuroendocrinol. **27**, 588–599 (2015).

33.   R. A. Dumbell, I. Petri, F. Scherbarth, V. Diedrich, H. A. Schmid, S. Steinlechner, P.

Barrett, Somatostatin Agonist Pasireotide Inhibits Exercise-Stimulated Growth in the Male

Siberian Hamster (Phodopus sungorus). *J Neuroendocrinol.* **29**, JNE.12448 (2017).

34.   F. Scherbarth, V. Diedrich, R. A. Dumbell, H. A. Schmid, S. Steinlechner, P. Barrett,

Somatostatin receptor activation is involved in the control of daily torpor in a seasonal

mammal. *Amer. J. Physiol. Reg. Intgr. Comp. Physiol.* **309**, R668–R674 (2015).

35.   G. Helfer, T.J. Stevenson, Pleiotropic effects of proopiomelanocortin and VGF nerve

growth factor inducible neuropeptides for the long-term regulation of energy balance. *Mol.*

Cell. Endocrinol. **514**, 110876 (2020).

36.   N. Nakao, H. Ono, T. Yamamura, T. Anraku, T. Takagi, K. Higashi, S. Yasuo, Y. Katou,

S. Kageyama, Y. Uno, T. Kasukawa, M. Iigo, P.J. Sharp, A. Iwasawa, Y. Suzuki, S. Sugano,

T. Niimi, M. Mizutani, T. Namikawa, S. Ebihara, H.R. Ueda, T. Yoshimura, Thyrotropin in the

pars tuberalis triggers photoperiodic response. *Nature* **452**, 317-322 (2008).

37.   M. Lorgen, E. Casadei, E. Krol, A. Douglas, M.J. Birnie, L.O. Ebbesson, T.O. Nilsen,

W.C. Jordan, E.H. Jorgensen, H. Dardente, D.G. Hazlerigg, S.A. Martin, Functional

divergence of type 2 deiodinase paralogs in the Atlantic salmon. *Curr. Biol.* **25**, 936-941

(2015).

38.   G. Helfer, P. Barrett, P.J. Morgan, A unifying hypothesis for control of body weight and

reproduction in seasonally breeding species. *J. Neuroendocrinol.* **31**, e12680 (2019).

39.   B.J. Prendergast, B. Mosinger, P.E. Kolattukudy, R.J. Nelson, Hypothalamic gene

expression in reproductively photoresponsive and photorefractory Siberian hamsters. *Proc.*

Natl. Acad. Sci. **99**, 16291-16296 (2002).

40.   R. Banks, M. Delibegovic, T.J. Stevenson, Photoperiod- and triiodothyronine-

dependent regulation of reproductive neuropeptides, proinflammatory cytokines, and

peripheral physiology in Siberian hamsters (Phodopus sungorus). *J. Biol. Rhythms*, **31**, 299-

307 (2016).

41.   T.J. Stevenson, K.G. Onishi, S.P. Bradley, B.J. Prendergast. Cell-autonomous

iodothyronine deiodinase expression mediates seasonal plasticity in immune function. *Brain*

Behav. Immun. **36**, 61-70 (2014).

42.   C.A. Steward, T.L. Horan, S. Schuhler, G.W. Bennett, F.J.P. Ebling, Central

administration of thyrotropin releasing hormone (TRH) and related peptides inhibits feeding

behavior in the Siberian hamster. *NeuroReport,* **14**, 687-691 (2003).

43. K. Labun, T.G. Montague, M. Krause, Y.N.T. Cleuren, H. Tjeldnes, E. Valen,

CHOPCHOP v3: expanding the CRISPR web toolbox beyond genome editing. *Nucl. Acid.*

Res. **47**, W171-W174 (2019).

44. T. Rausch, M. H.-Y. Fritz, A. Untergasser, V. Benes, Tracy: basecalling, alignment,

assembly and deconvolution of sanger chromatogram trace files. BMC Genomics 21, 230

(2020).

45. K. Okonechnikov, O. Golosova, M. Fursov, the UGENE team, Unipro UGENE: a

unified bioinformatics toolkit. *Bioinformatics* **28**, 1166–1167 (2012).

46. A.S. Przybylska-Piech, M. Jefimow, Siberian hamster nonresponding to short

photoperiod use fasting-induced torpor. *J. Exp. Biol.* **225**, jeb244222 (2022).

47. H. Li, Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**,

3094–3100 (2018).

48. W. Ernst, E. Trummer, J. Mead, C. Bessant, H. Strelec, H. Katinger, F. Hesse,

Evaluation of a genomics platform for cross-species transcriptome analysis of recombinant

CHO cells. *Biotechnol. J.* **1**, 639–650 (2006).

49. R. Patro, G. Duggal, M. I. Love, R. A. Irizarry, C. Kingsford, Salmon provides fast and

bias-aware quantification of transcript expression. *Nat. Methods* **14**, 417–419 (2017).

50. M. D. Robinson, D. J. McCarthy, G. K. Smyth, edgeR: a Bioconductor package for

differential expression analysis of digital gene expression data. Bioinformatics 26, 139–140

(2010).

51. S. Zhao, R.D. Fernald, Comprehensive algorithm for quantitative real-time

polymerase chain reaction. *J. Comput. Biol.* **12**, 1047-1064 (2005).

52. C. Stewart, T.A. Liddle, T.J. Stevenson. Abundance, efficiency and stability of

reference transcript expression in a seasonal rodent: the Siberian hamster. *PLoS One* **17**,

e0275263 (2022).

53.   T. Zielinski, A.M. Moore, E. Troup, K.J. Halliday, A.J. Millar, Strengths and limitations

of period estimation methods for circadian data. *PLoS One* **9**, e96462 (2014).

54.   Y. Benhamini, Y. Hochberg, Controlling the false discovery rate: a practical and

powerful approach to multiple testing. *J. Roy. Stat. Soc. B.* **57**, 289-300 (1995).

55.   S.X. Ge, D. Jung, R. Yao, ShinyGO: a graphical gene-set enrichment tool for animals

and plants. *Bioinformatic* **36**, 2628-2629 (2020).

Supplementary Figures:

Fig.

S1. Plasma insulin reflects changes in transcript expression. (A) Plasma insulin of Siberian
hamsters across 32-weeks of short photoperiod exposure (H (8) = 20.401, P = 0.009). (B) Expression
of *Dio3* from the MBH of Siberian hamsters, assessed by qPCR, across 32-weeks of short
photoperiod exposure (H (8) = 39.771, P < 0.001). (C) Expression of *Sst* from the MBH of Siberian
hamsters, assessed by qPCR, across 32-weeks of short photoperiod exposure (H (8) =16.002, P =
0.042). Letters denote significance between groups.
706

a)

b)

Fig. S2. **Gene ontology analysis of MBH sequencing reveals well known seasonal pathways.**
(A) Gene ontology enrichment analysis of significant transcripts from sequencing of Djungarian
hamster MBH across 32-weeks of short photoperiod exposure. (B) Interaction network of enriched
gene ontology terms. Size of individual points represent the number of transcripts in each term.
Thickness of interactions represents shared transcripts between terms.

a)

b)

Fig. S3. Gene ontology analysis of pituitary gland unveils potential mechanisms for seasonal
changes in protein processing and release. (A) Gene ontology enrichment analysis of significant
transcripts from sequencing of Djungarian hamster pituitary gland across 32-weeks of short
photoperiod exposure. (B) Interaction network of enriched gene ontology terms. Size of individual
points represent the number of transcripts in each term. Thickness of interactions represents shared
transcripts between terms.
722

Fig. S4. Paraventricular and dorsomedial hypothalamus sequencing unveils novel transcripts and widespread seasonal interval timing within the hypothalamus. (A-B) Heatmaps of significant (FDR < 0.1) Djungarian hamster transcripts displaying sine or cosine rhythmicity were selected from (A) paraventricular hypothalamus and (B) dorsomedial hypothalamus. (C) Polar scatter chart of significant transcripts from Djungarian hamster paraventricular and dorsomedial hypothalamus, displaying peak of expression and -log(FDR).

730
731

733 **Fig. S5. Gene ontology analysis of paraventricular hypothalamic sequencing.** (A) Gene ontology
734 enrichment analysis of significant transcripts from sequencing of Djungarian hamster paraventricular
735 hypothalamus across 32-weeks of short photoperiod exposure.
736
737

Fig. S6. Gene ontology analysis of dorsomedial hypothalamus suggests widespread immune involvement and cellular differentiation. (A) Gene ontology enrichment analysis of significant transcripts from sequencing of Djungarian hamster dorsomedial hypothalamus across 32-weeks of short photoperiod exposure. (B) Interaction network of enriched gene ontology terms. Size of individual points represent the number of transcripts in each term. Thickness of interactions represents shared transcripts between terms.

Fig. S7. Rheostatic mechanism controlling body mass change in Djungarian hamsters. (A)
Epidydimal fat mass in long- and short- photoperiod exposed Djungarian hamsters (F1, 20 = 59.136;
P < 0.001) after an overnight food restriction, or ad libitum, feeding protocol (F1, 22 = 2.738; P =
0.114). (B) Pituitary gland expression of *Gh* from in long- and short- photoperiod exposed Djungarian
hamsters (F1, 20 = 2.738; P = 0.114) after an overnight food restriction, or ad libitum, feeding
protocol, assessed by qPCR (F1, 20 = 0.518; P = 0.480). (C) Plasma insulin in long- and short-
photoperiod exposed Djungarian hamsters (F1, 31 = 29.865; P < 0.001) after an overnight food
restriction, or ad libitum, feeding protocol (F1, 31 = 6.863; P = 0.014). Letters denote significant main
effect of photoperiod (a). Asterisk indicates significant effect of photoperiod (P < 0.005), and hashtag
denotes significant main effect of food restriction (P < 0.05).

Fig. S8. Evidence of CRISPR modification in Dio3cc and Dio3wt Djungarian hamsters. (A)
Binding positions of gRNA1 and gRNA2, injected into Djungarian hamster hypothalamus (B)
Sequence of gRNA1 and gRNA2 binding and PAM sites. (C) Representative Dio3 sequence
alignment with gRNA1 and PAM sites indicated for a representative Dio3wt (control) and Dio3cc
(CRISPR) hamster. (D) Decomposition error difference for Dio3 sequence alignment to determine
genomic modification (t5 = 2.66, p < 0.05) (E) Pelage change in Dio3wt and Dio3cc Djungarian
hamsters across 32 weeks of SP exposure, Dio3wt showed significantly increased pelage score at 13
(P < 0.001) and 14 (p < 0.001).

768
769

a)

b)

Fig. S9. Schematic model for the mechanisms of circannual interval timing. (a) Hamsters held in
long photoperiod (LP) have high body mass, thyrotropin-stimulating hormone beta (Tshβ) and low
somatostatin (Sst) and deiodinase type-3 (Dio3). Transfer to short photoperiod (SP) results in the
induction of circannual interval timing characterized by a reduction in body mass, low Tshβ in the pars
tuberalis and increased Dio3 expression in tanycytes. The maintenance phase is associated with
increased Sst expression which serves to inhibit growth. The onset of the recovery phase is
associated with a complete reversal in whole organismal physiology to the LP phenotype. (b)
tanycytes along the third ventricle in the mediobasal hypothalamus integrate photic cues derived from
Tshβ and are sensitive to nutritional cues. The triiodothyronine (T3) output signal from tanycytes
regulates Sst expression leading to long-term programmed rheostatic changes in body mass that
serve to control circannual interval timing in multiple physiological systems. Conversely, homeostatic
stability is maintained despite large scale seasonal rhythms in body mass. Homeostatic energy
balance is established by well characterized circuits that include neuropeptide Y (Npy), agouti-related
peptide (Agrp) and the melanocortin receptor 4 (Mc4r).

Gene	Primer	Size	Temp	Melt
Reference	-	-	-	-
18s	GCTCCTCTCCTACTTGGATAACTGTG	111	62˚C	80˚C
	CGGGTTGGTTTTGATCTGATAAATGCA			
Hrpt	AGTCCCAGCGTCGTGATTAGTGATG	141	62˚C	76˚C
	CGAGCAAGTCTTTCAGTCCTGTCCA			
Mediobasal hypothalamus				
Npy	CCAGGCAGAGATACGGCAAGAGATC	119	60˚C	81˚C
	CCATCACCACATGGAAGGGTCC			
Sst	GAAGTCTCTGGCGGCTGCTG	145	60˚C	85˚C
	CAGCCTCATTTCATCCTGCTCCG			
Dio3	CATGCTCCGCTCCCTGCTGCTTCA	251	62˚C	85˚C
	CAGGGTGCACAGACGGTTGTC			
Pituitary Gland				
Prl	TCCGGAAGTCCTTCTGAACC	300	60˚C	82˚C
	CGCAGGCAGCGAATCTTATTG			
Gh	ACCTACAAAGAGTTTGAGCGTG	167	58˚C	85˚C
	ATGAGCAGCAGCGAGAATCG			

789 **Table S1.** Djungarian hamster qPCR primers sequences and thermal cycling profiles.

790