A role of villin-dependent F-actin organization in peroxisome motility in Arabidopsis cells

Calvin H. Huang [a], Amanda Koenig [b], Yuh-Ru Julie Lee [a], Yibo Shi [a], Jianping Hu [b,c,1], Bo Liu [a,1]

[a] Department of Plant Biology, College of Biological Sciences, University of California, Davis, CA
95616, U.S.A.

[b] Michigan State University-Department of Energy Plant Research Laboratory, Michigan State
University, East Lansing, MI 48824, U.S.A.

[c] Department of Plant Biology, Michigan State University, East Lansing, MI 48824, U.S.A.

[1] Corresponding authors: Bo Liu, bliu@ucdavis.edu; Jianping Hu, huji@msu.edu

Short title: Villins and peroxisome motility

ABSTRACT

Actin microfilaments (F-actin) serve as the track for directional movement of organelles in plant

cells. In actively growing plant cells, F-actin often form robust bundles that trespass the cellular

dimension. To test how the F-actin network was employed for peroxisome movement, we

wished to disturb actin organization by genetically compromising the function of villin (VLN)

proteins that serve as the primary bundling factor in *Arabidopsis thaliana* cells. To do so, we

isolated T-DNA insertional mutants in three *VLN* genes that were most actively expressed in

vegetative tissues. We found that the *vln4* mutation greatly enhanced the growth defects caused

by the *vln2 vln3* double mutant as the *vln2 vln3 vln4* triple mutant had a great reduction of organ

growth and formed heavily deformed tissues. Both VLN2 and VLN4 proteins were detected on

bundled F-actin filaments. Compared to the wild-type cells, the double and triple mutants

exhibited progressively reduction of stable F-actin bundles and had fine F-actin filaments

undergo rapid remodeling. The defective F-actin network did not prevent peroxisomes from

taking on both rapid and slow movements along the F-actin tracks. However, we found that

compromised F-actin bundling caused significant reductions in the speed of peroxisome

movement and the displacement distance of peroxisome positions. Using a correlation analysis

method, we also demonstrated that the complex heterogeneous peroxisome movement may be

classified into clusters reflecting the directionality of peroxisome movement. The triple mutant

suffered from a significant reduction of peroxisomes exhibiting long-range and linear movement.

Our results provided insights into how VLN-dependent F-actin organization was coupled with

the complex patterns of peroxisome movement.

INTRODUCTION

The single membrane organelle peroxisomes display rapid directional movement in the cytoplasm while exercising their essential functions of detoxification, β oxidation, and phytohormone synthesis. All these functions probably require peroxisome translocation throughout the cell to scavenge cytotoxins and coordinate with other cellular compartments for versatile metabolic reactions (Sandalio et al., 2021). In mammals, peroxisomes interact with the endoplasmic reticulum (ER), lipid droplets, lysozymes, and mitochondria during metabolic processes like fatty acid catabolism, and defects in peroxisomal distribution and inter-organellar interactions are the underlying cause of several diseases in humans (Wanders et al., 2023). In plants, peroxisomes also exhibit critical interactions with chloroplasts for detoxification, specialized metabolite synthesis, and photorespiration besides the interactions with the ER and mitochondria (Koenig et al., 2023; Pan et al., 2020). Additionally, peroxisomes participate in functions spanning subcellular topography, and their trafficking and spatial arrangement often are linked to the fluctuating demands of the cell during development or under stresses (He et al., 2021). The dynamic distribution of peroxisomes, therefore, requires agile motility along the cytoskeletal filaments and their associated motor proteins (Neuhaus et al., 2016).

In fungi and animals, peroxisome trafficking is primarily governed by microtubules and the associated motors of dynein and kinesins (Covill-Cooke et al., 2021; Salogiannis and Reck-Peterson, 2017). In plants, however, active movement of peroxisomes and other organelles predominately takes place on the F-actin tracks and is dependent on the motors in the Myosin XI subfamily (Avisar et al., 2008). In *Arabidopsis thaliana*, for example, actin depolymerization or inactivation of multiple Myosin XI genes causes nearly complete cessation of organelle movement (Mathur et al., 2002; Peremyslov et al., 2010). However, research to date has barely explored the molecular mechanisms that regulate actomyosin-dependent organelle movement in plants (Koenig et al., 2023). Typical differentiated plant cells have F-actin organized into two distinct populations of long and stable bundles and highly dynamic filaments (Henty-Ridilla et al., 2013). It is unclear whether organelles like peroxisomes employ either or both types of F-actin for their directional movement.

F-actin bundling in plants is primarily mediated by the villin (VLN) family proteins, which were initially isolated from lily pollen as abundant actin-binding proteins (Huang et al., 2015; Yokota et al., 2003). In Arabidopsis, VLN proteins encoded by five paralogous genes show various degrees of actin bundling capabilities both *in vitro* and *in vivo* (Li et al., 2025). Of the five

paralogs, *VLN2*, *VLN3* and *VLN4* are most broadly expressed in vegetative tissues. The *vln2*

vln3 double mutants display a reduction in F-actin bundling and altered growth patterns with

twisted organs, but no significant compromise in vegetative growth (Bao et al., 2012; van der

Honing et al., 2012). Therefore, we hypothesized that VLN4 might function redundantly with

VLN2 and VLN3 in mediating organelle movement and the coupled growth processes in plants.

Here, we analyzed F-actin organization and peroxisome movement in the mutants of these

three Arabidopsis *VLN* genes that are highly expressed in vegetative tissues. Here we found

that the *vln2 vln3 vln4* triple mutant showed greatly reduced F-actin bundling and gross defects

in vegetative growth. The triple *vln* mutant allowed us to interrogate how compromised F-actin

bundling, without disassembling the entire F-actin network, impacted patterns of peroxisome

movement. Our results demonstrated that peroxisomes showed significantly reduced movement

speed and the displacement distance in the triple mutant. We also developed novel quantitative

methods to differentiate features associated with peroxisomal movement modes and correlate

such differences to the varied F-actin organization patterns.

RESULTS

Deficient F-actin organization in the *vln* mutants

Although peroxisomes are known to move along F-actin tracks by Myosin XI motors in plant

cells, it is unknown how the F-actin network is employed for the directional movement. We

sought to understand the role of actin organization, particularly F-actin bundling, in peroxisome

movement. Since the VLN proteins have been shown to contribute significantly to actin bundling

in plants (Huang et al., 2015), we aimed to employ the *vln* mutants in Arabidopsis to address the

potential role of F-actin bundling in peroxisome movement. Of the five Arabidopsis *VLN*

paralogs, *VLN2*, *VLN3* and *VLN4* are most broadly expressed in vegetative tissues according to

community-generated transcriptional data (https://bar.utoronto.ca/eplant/). It has recently been

reported that VLN2, VLN3, and VLN4 function redundantly in hypocotyl elongation, with VLN2

as the most abundant one in the hypocotyl, and that higher order *vln* mutants display aberrant

actin organization in their root and leaf cell (Li et al., 2025). Therefore, we predicted that these

three VLN proteins function redundantly to regulate organelle movement and the coupled

growth processes. To determine the role of VLN in peroxisome dynamics, we isolated the *vln4*

mutant and crossed it with the *vln2 vln3* (*vln2/3*) double mutant (Bao et al., 2012). As a result, a

vln2 vln3 vln4 (*vln2/3/4*) triple mutant was independently generated, which displayed growth

defects (Suppl. Figure 1) similar to what was described recently (Li et al., 2025).

Because of the impact of actomyosin-dependent cell growth and the defect of shoot cell

elongation in the *vln2/3/4* mutant, we used hypocotyl cells to characterize the actin defects in

more detail before investigating peroxisomal dynamics in the mutants. We first confirmed that F-

actin cable formation is compromised in hypocotyl cells in both the double and triple *vln* mutants

by live-cell imaging of F-actin labeled by a LifeAct-GFP (green fluorescent protein) fusion

protein (Figure 1A). Compared to the wild-type (WT) cells in which thick, continuous F-actin

bundles spun the cell length, the *vln* mutant cells showed that fine F-actin filaments were

organized in mesh-like networks and F-actin bundles were lessened in the *vln2/3* double mutant

or completely absent in the *vln2/3/4* triple mutant (Figure 1A). In addition, we quantified the

reduction of actin bundles in *vln* hypocotyl cells using the parameter skewness, a standard

metric used in assessing actin filament bundling in which higher fluorescence intensity of the

LifeAct-GFP signal in bundled filaments result in a higher skewness value (Higaki, 2017; Higaki

et al., 2010). The *vln2/3* and *vln2/3/4* mutants exhibited progressively reduced skewness

compared to the WT, indicative of decreased F-actin bundling in the mutant cells (Figure 1B).

In WT cells, prominent F-actin cables often are oriented perpendicularly, in parallel to the cell

elongation axis, likely underlying the cell elongation phenotype observed in the hypocotyls in the

recent report (Li et al., 2025). To further assess the deficient F-actin network in the hypocotyl of

the *vln* mutants, we quantified the average angle of F-actin filaments relative to the horizon, with

0° being in the transverse direction and 90° being parallel to the growth direction. In epidermal

cells of the WT hypocotyl, due to the presence of longitudinal bundles along the growth

direction, F-actin were more parallel to the direction of growth, indicated by a higher average

angle. In the *vln2/3* and *vln2/3/4* mutant cells, however, the overall angles become shallower,

concomitant with the decrease in these long cables and the increasingly abundant appearance

of mesh networks of F-actin (Figure 1C).

The dynamic and cell-spanning organization of F-actin is maintained in *vln* mutants.

To evaluate the specific contribution of actin bundling to peroxisome motility, ideally, we should

conduct the assessment in a system that retains functional albeit unbundled F-actin filaments.

To this end, we characterized the F-actin network in the *vln* mutants using stability, coherency,

and density as the metrics. Generally, thick F-actin fibers tend to be stable over time, whereas

actin filaments in the mesh network are more dynamic (Henty-Ridilla et al., 2013). Because the

F-actin mesh network is pronounced in the *vln2/3* and *vln2/3/4* mutants, we tested whether the

altered array organization was associated with rapid actin remodeling in these mutants. Images

of F-actin were captured at 5-sec intervals, after which pseudo-colored images from different

timepoints were merged. Noticeably, while F-actin in WT cells predominantly appeared in white

in the merged images, which is indicative of little if any remodeling, the mutant cells showed

more dynamic changes in F-actin organization with each timepoint observed in discrete colors

(Figure 2A). Further, the white color in the merged images often correlated with thick, long F-

actin bundles that are more static than those filaments in the mesh networks in the same WT

cells (Figure 2A). In the *vln2/3* and *vln2/3/4* mutant cells, the three distinct colors were more

evidently discerned (Figure 2A). Likewise, quantification of this F-actin stability showed that a

significantly higher proportion of F-actin in the *vln* mutants underwent remodeling when

compared to those of the WT and the *vln2/3/4* cells had significantly fewer F-actin cables than

vln2/3 (Figure 2B). So, our further characterization of the *vln* mutants supported previous

conclusions that loss of these three VLN proteins severely impaired F-actin bundling while

leaving a diffuse, dynamic mesh network of F-actin.

Coherency is a measure of the degree of isotropy of the F-actin distribution, where 0 indicates

entirely random orientation and 1 indicates uniform orientation, characteristic of cells enriched

with actin cables. F-actin filaments in WT cells are organized in a pattern with higher coherence

values compared to more disordered filaments with coherence closer to 0 in the *vln2/3* and

vln2/3/4 mutants (Figure 2C). The decreased coherency in the *vln* mutants suggests dispersed,

mesh F-actin, as a consequence of the reduced presence of cell-spanning F-actin bundles.

To further characterize the more diffuse, less linear F-actin network in *vln* mutants, we quantified

the density of F-actin by measuring the percentage of F-actin occupancy in the cytoplasm. The

vln2/3 cells exhibited significantly higher F-actin occupancy than the WT cells, and the F-actin

density in the higher order *vln2/3/4* mutant approximately doubled from that of the WT (Figure

2D). The increased F-actin occupancy suggests that the F-actin network, despite being devoid

of conspicuous bundles, still spans the cortex, and therefore the cytoskeletal apparatus capable

of supporting organelle motility across the whole cell should persist in the mutants.

To learn intracellular localization of the VLN proteins, we first had a VLN2-GFP fusion protein

expressed under the control of its own promoter in the *vln2/3* double mutant. Because growth

phenotype was suppressed by VLN2-GFP expression (Supple. Figure 2A), this fusion protein

was functional. When this functional VLN2-GFP was co-expressed with LifeAct-mScarlet-I, the

VLN2 was most conspicuously detected on F-actin bundles (Supple. Figure 2B). A mCherry-

VLN4 fusion protein also decorated F-actin bundles marked by LifeAct-GFP (Supple. Figure

2B).

Taken together, our data showed that in the *vln* mutants, F-actin dynamic polymerization and

cell-spanning organization is maintained while only cable formation is deficient. As such, these

knockout mutants can serve as an informative genetic material to study the role of actin bundles

in peroxisome motility.

VLN-dependent actin bundling facilitates peroxisome movement

To determine the impact of actin bundling in peroxisome movement and distribution in

Arabidopsis, we transformed WT, *vln2/3*, and *vln2/3/4* plants with constructs for the markers of

F-actin (LifeAct-GFP) and peroxisome (mScarlet-I-SRL) and visualized them simultaneously in

the hypocotyl cells of plants co-expressing the two fusion proteins. The transgenic *vln* double

and triple mutants exhibited the progressive deficiencies in F-actin bundling, resembling what

was observed earlier in this study. These defects were concomitant with the progressive

aggregation of peroxisomes, suggesting an inability of peroxisomes to be distributed uniformly

across the cell (Figure 3A). Time-lapse images, taken at 5-sec intervals in epidermal cells of the

hypocotyl, revealed that peroxisomes maintained motility along the F-actin tracks in both *vln2/3*

double and *vln2/3/4* triple mutants (Figure 3A, Movie S1). Moreover, each track of peroxisome

movement was heterogenous, with some rapid movements (>1.5 μm/s) as well as more

frequently observed slower movements (<0.5 μm/s), similar to what was observed in the WT

(Figure 3B), indicating that the mechanisms of actomyosin-dependent peroxisome movement

were at least partially maintained in the *vln* mutants.

We first compared the speed of peroxisome movement among the three genotypes of WT,

vln2/3 and *vln2/3/4*. Despite variations in instantaneous speed during imaging duration, the

mean speed of each peroxisome movement revealed that only the severe inhibition of F-actin

bundling in *vln2/3/*4 caused a significant decrease in the mean speed, while intermediate

bundling in *vln2/3* was sufficient to maintain peroxisome speed comparable to that of the WT

(Figure 3B). These data suggest that peroxisome movement is facilitated by both discrete F-

actin filaments and densely bundled actin cables. To exclude peroxisomes that potentially were

unable to associate with F-actin via myosin in the mutants, we calculated the maximum speed

of peroxisome movement, defined as the highest instantaneous speed recorded between two

consecutive timepoints, to assess the movement of peroxisomes already associated with

myosin motors. Again, the maximum speed distributions were not significantly different between

WT and *vln2/3*, whereas peroxisome movement was more frequently observed at slower

maximum speeds in *vln2/3/4* (Figure 3C). We therefore concluded that the actin dynamics and

organization influence peroxisome velocity.

Fast, directional peroxisome movement is regulated by F-actin cables

Since the heterogenous patterns of peroxisome motility biased towards slower speeds in the

absence of longitudinal actin bundles, as shown in the *vln2/3/4* triple mutants, we hypothesized

that F-actin cables specifically facilitate fast, linear movement, which consequently enables

uniform peroxisome distribution across the cell. To test this, we evaluated total moving distance,

displacement, straight-line speed, overall linearity of movement, and directional change rate

(Figure 4A-E). We first measured the total moving distance and found most peroxisomes

traversed similar distances with no significant difference among WT, *vln2/3,* and *vln2/3/4* mutant

cells (Figure 4A). However, whereas peroxisomes in *vln2/3/4* have a significantly shorter

displacement, measured as the shortest distance between the start and end points of the

tracked peroxisome (Figure 4B). Furthermore, a consistent phenotype was observed when the

displacement distance was normalized by track duration, as represented by "straight-line

speed", which accounted for any differences in the number of motions tracked (Figure 4C).

Together these data suggest that, while peroxisomes still move in short bursts that sum to a

similar distance observed between *vln* mutants and WT, peroxisomes did not travel as far

distance in the *vln2/3/4* mutant in the absence of prominent F-actin bundles or cables when

compared to WT or the *vln2/3* mutant. In fact, peroxisomes in *vln2/3/4* mutants shifted direction

more frequently than those in the WT cells and their motility was overall less linear and more

frequently changed directions (Figure 4D, E), likely because their movement predominantly

employed the dynamic F-actin mesh network rather than stable, longitudinal bundles as the

tracks.

To better assess the heterogenous motility patterns of peroxisomes, we employed a Pearson's

correlation matrix to conduct pairwise analysis of the influence of the aforementioned motility

parameters on each other within individual genotypes. While both *vln* mutant genotypes showed

correlations comparable to wildtype between average velocity, maximum speed, total distance

and track displacement, the correlations between parameters that account for directional

movement (straightforward speed, linearity, and directional change rate) varied across

genotypes (Supple. Figure S3A). We then plotted the data of straight-line speed, linearity, and

directional change rate from all genotypes in a 3-dimensional scatterplot and used k-means

clustering to group motility according to these multidimensional criteria (Figure 5A). The 3

clusters, determined using the Elbow Plot method (Supple. Figure S3B), comprised high

straightforward speed and linearity but low directional change rate (Cluster 1); intermediate

straightforward speed, linearity, and directional change (Cluster 2); and low straightforward

speed and linearity but high directional change rate (Cluster 3). Cluster 1, therefore, was

enriched for peroxisomes that rapidly and directionally traversed the cell with minimal turns,

while Cluster 3 represented peroxisomes that move in short bursts with frequent changes in

direction. Interestingly, WT peroxisomes exhibited uniformly heterogeneous movement, with

roughly one third of their tracked movements represented by each of the three clusters (Figure

5B). However, in the vln2/3 mutants, a smaller proportion of peroxisomes move directionally

(29.6%), as characterized by Cluster 1, compared to WT (32.8%), and the peroxisome motility

profiles shifted towards intermediate (Clusters 2) and short burst (Clusters 3) movements

(Figure 5B). In the *vln2/3/4* triple mutant, the reduction in fast, linear peroxisome movements

was further exacerbated (20.6%), with more peroxisomes moving in short bursts (39.8%),

compared to the WT (30.5%). This shift in cluster populations across genotypes suggests that a

lesser proportion of peroxisomes move quickly and linearly when F-actin bundling is deficient in

the *vln* mutants.

Taken together, our results showed that densely bundled F-actin cables were necessary for

rapid, directional movement of peroxisomes, which enabled the subcellular distribution of

peroxisomes required for their diverse metabolic functions.

DISCUSSION

We used the independently generated *vln2/3/4* triple mutant to investigate the synergistic roles

of three F-actin bundling VLN proteins in the dynamics and organization of the F-actin network.

Our quantitative analysis of peroxisome motility in the WT and *vln* mutants showed that such

VLN-dependent F-actin organization is required for rapid directional movement of peroxisomes

in the epidermal cells of the Arabidopsis hypocotyls.

The plant VLN proteins are grouped into three classes and VLN4 belongs to the class III while
VLN2 and VLN3 are in class II (Khurana et al., 2010). Consistent with what was reported
recently (Li et al., 2025), the loss of VLN4 greatly enhanced the defects observed in the *vln2/3*
double mutant. Therefore, these results collectively indicated that the function of these three
VLN proteins, despite being in two different classes, converged towards stabilizing F-actin and
promoting high order F-actin bundle formation in vegetative tissues. However, when the three
VLN proteins were examined individually without an F-actin marker, they exhibited different
extents of F-actin decoration in different tissues (Li et al., 2025). It would be interesting to learn
how these proteins localize and function in concert along F-actin filaments *in vivo*. Surprisingly,
additional mutations in two other *VLN* genes encoding class I and III VLN proteins did not
further enhance the growth phenotypes observed in the *vln2/3/4* triple mutant (Li et al., 2025).
Nevertheless, this finding suggested that the three VLN proteins included in our study played
the primary role in F-actin bundling.

Longitudinally oriented, prominent F-actin cables are found in interphase plant cells and often
are considered the primary tracks of long-distance transport of organelles for their robust
anisotropic expansion (Garcia-González and van Gelderen, 2021). Here, the *vln2/3/4* triple
mutant informed us that fine filaments of F-actin are equally qualified for directional transport of
peroxisomes. But the long, stable F-actin bundles exhibited great advantages of promoting
rapid, directional movement. Compromises of such rapid, directional movement have a
profound consequence in cell expansion.

Phenotypic analysis in myosin mutants unequivocally demonstrated that directional movement
of organelles like peroxisomes is dependent on Myosin XI motors (Peremyslov et al., 2010;
Perico and Sparkes, 2018). It is intriguing how specific organelle would be actively transported
with such a wide range of speeds. We hypothesize that the heterogeneity of peroxisome motility
is probably brought about by different Myosin XI motors that exhibit different motile activities
(Haraguchi et al., 2016). At least 7 of 13 Myosin XI motors in *A*. *thaliana* had their DIL (dilute)
domains that exhibit peroxisome-binding activities when truncations including the domains are
ectopically expressed in a heterologous system (Sattarzadeh et al., 2011). However, it is yet to
be determined whether the full-length myosin proteins are associated with the organelle when
expressed under their native promoters. Alternatively, multiple motors could bind to an organelle
to increase the movement processivity and velocity so that it can be transported for long
distance, especially in giant cells of organisms like *Chara corallina* (Shimmen and Yokota,
2004). Varied numbers of active motors per peroxisome, which is hard to be quantified *in vivo*,

likely contribute to the heterogeneity in transport velocities as well. The significance of the
heterogeneous peroxisome movement is unclear. Perhaps peroxisome motility may be coupled
with the position of other organelles to facilitate inter-organelle communication. So, it would be
interesting to determine whether certain patterned movement events were shared by other
organelles. Alternatively, peroxisome velocities could vary across the cytoplasm and were
coupled with geometrically determined space. In addition, various intrinsic or extrinsic factors
cause each peroxisome to display distinct velocities and directionalities.

METHODS AND MATERIALS

Plant genotype and growth

Arabidopsis thaliana plants were grown in soil in a growth chamber at 21°C with cycles of 16hr
light and 8hr dark. *Arabidopsis* seedlings that were used for imaging were sterilized and sowed
onto ½ Murashige & Skoog Basal media with 0.8% phytagel. Seeds in solid media were cold
stratified for 2 days then moved into the growth chamber for 6 days before imaging. All
Arabidopsis thaliana mutants and wild-type plants are in the Col-0 genetic background. Floral
dipping method was used to deliver transgenes into the plant, following established protocols.

Mutants of *VLN4* were isolated from seed stock of SALK_109099C (Arabidopsis Biological
Resource Center, Ohio State University), which carried a T-DNA insertion in the 12[th] intron of the
VLN4 gene. The *vln2/3* double mutant was described in an earlier publication (Bao et al., 2012).

Plasmid construction

To produce an expression vector carrying cassettes for expression of both mScarlet-I-PTS1 and
LifeAct-GFP, we first had the codon-optimized mScarlet-I-encoding DNA fragment synthesized
commercially (Integrated DNA Technologies, Inc.). The SRL-coding sequence was added to its
3' end by PCR using primers of 5'- GTA CCG AAT GGT GAG CAA GGG AGA AG-3' and 5'-CGG
CGC GCC TTA TAG CCG GCT TTT ATA CAG TTC GTC CAT ACC GCC G-3'. The fragment

was inserted into a plasmid containing the promoter sequence of EF1α (AT1G07920).

Separately, the LifeAct-coding sequence was synthesized and inserted into pENTR4 (Thermo

Fisher). The resulting plasmid was used to generate a plasmid expressing LifeAct-GFP under

the viral 35S promoter by Gateway LR cloning with the pGWB5 plasmid (Nakagawa et al.,

2007). The resulting plasmid was linearized by ApaI where the DNA fragment for

EF1α(p)::mScarlet-I-PTS1, amplified by the primers of 5'- GCG ACA ATA TGA TCG GGC CTC

CGG TGT CAA TCT CAG G-3' and 5'- TGT GGA CGC CGG GCC TTA TAG CCG GCT TTT ATA

CAG TTC G-3' from the plasmid described above, was inserted by Gibson reaction to give rise

to the plasmid pLB187 used for transformation.

To generate the *VLN2* expression plasmid, a 7-kb genomic *VLN2* genomic fragment containing

1.1-kb of the predicted promoter region was amplified by using the primers 5'-GAA CCA ATT

CAG TCG ACT GGA TCC GGT ACC TCT TTT GAG CTC TAA TTA ATT TTG GAT GAC-3' and

5'-TCC AGA GGA TCC AGA TGA ACC AGA AGA ACC GAA CAA GTC GAA CTT CTT CTT AAG

CAG-3'. The amplified fragment was introduced into a modified entry vector containing a C-

terminus GFP and PA tag (Huang et al., 2024) by Gibson Assembly. The resulting *VLN2* entry

plasmid was recombined with the destination vector pGWB1 (Nakagawa et al., 2007) by an LR

reaction to produce the *VLN2* expression plasmid.

To generate the *VLN4* expression plasmid, a 5-kb *VLN4* genomic CDS fragment was amplified

by using the primers 5'-TGT ACA AGG GCG CAG GCG CGA TGT CTG TTT CTA TGA GAG

ATT TAG ATC-3' and 5'-TTG TAC AAG AAA GCT GGG TCT AGA TTC AGA AAA GCT GAA CAG

CCA TTT TG-3'. The amplified fragment was inserted into a modified entry vector containing an

N-terminus mCherry-coding sequence by Gibson Assembly. The resulting plasmid was then

linearized by PCR with the primers 5'-TAT GGT GAG CAA GGG CGA GGA G-3' and 5'-CAG

TCG ACT GAA TTG GTT CCC ATG G-3'. A 2.5-kb predicted promoter fragment of the *VLN4*

gene is amplified by using the primers 5'-GGG AAC CAA TTC AGT CGA CTG ATT TGA GGT

ACG AAG CAG ATG-3' and 5'-TCC TCG CCC TTG CTC ACC ATA CTT AAG CAG TCT TTA

CCA CTT TCT C-3'. This *VLN4* promoter fragment and the linearized plasmid described above

were assembled together by Gibson Assembly. The resulting *VLN4* entry vector was

recombined with pGWB1 by LR reaction to produce the *VLN4* expression plasmid.

Plasmid were transformed into the agrobacteria strain Gv3101 prior to floral dipping in the WT,

vln2/3 and *vln2/3/4* plants.

Confocal microscopy

After germination and growth in the chambers, the seedlings were mounted in water for imaging. Live-cell imaging was performed on an LSM980 laser scanning confocal microscope with the Airyscan2 function using a 40X water immersion objective (Carl Zeiss). Single frame images of F-actin were used to measure orientation, coherency, density and skewness. For measurements of F-actin stability, time-lapse images with 5sec/frame were obtained. Similarly, to image peroxisome movement, both markers for F-actin and peroxisomes were simultaneously imaged with a time-lapsed of 5sec/frame. At least 3 independent transformant lines were imaged per genotype.

Quantification of F-actin

Images were first rotated such that the direction of growth in along the vertical axis. The images were thresholded, binarized and skeletonized for quantifying F-actin (Higaki, 2017). A region of interest was drawn to select only the cell of interest. The imageJ plugin OrientationJ was used to measure orientation, and coherency (Püspöki et al., 2016). Skewness and density were measured by using the LPX plugin package (Higaki et al., 2010; Schneider et al., 2012).

To measure percentage of stable F-actin, we imaged F-actin for 10 seconds at 5 sec/frame. The first and last frames were merged. The assumption is that stable filaments should still be present or unchanged in position after 10 seconds. A stable filament from the last frame should colocalize with a filament from the first frame. The merged image was thresholded for colocalized filaments. The percentage of stable F-actin was calculated using a ratio of colocalized F-actin signal to total F-actin signal.

Quantification of peroxisome motility

Quantitative measurement of F-actin-dependent peroxisome motility was carried out by imaging both F-actin and peroxisomes simultaneously with 5-sec intervals. Peroxisomes that were

unattached from F-actin were not measured. Peroxisomes were tracked using the ImageJ

plugin, TrackMate (Ershov et al., 2022). Previous reports have mentioned that slow moving

peroxisomes are usually at least 0.2 µm/sec (Mathur et al., 2002). we excluded peroxisomes

that were moving at velocities below 0.1 µm/sec as those appear to be more likely due to

Brownian movements. We manually reevaluated the tracks with velocities below 0.1 µm/sec to

confirm that these movements appeared oscillatory. The output track data summarized the

overall movement of individual peroxisome during the imaging duration, and provided the track

features regarding: (1) mean speed for the average speed values over every 5-second intervals

along a track; (2) max speed for the highest instantaneous speed between two consecutive

timepoints along a track; (3) straight-line speed, for track displacement divided by track duration;

(4) total distance traveled; (5) track displacement; (6) linearity of forward progression for the

ratio between straight-line speed and track mean speed, when tracks that are completely linear

would have a value of 1; and (7) directional change rate for the average angle between

consecutive timepoints along a track divided by the track duration.

Acknowledgements

We thank Dr. Shanjin Huang for providing the vln2/3 mutant seeds and Dr. Tsuyoshi Nakagawa

for sharing the pGWB plasmids. We also thank Dr. Felicia L. Peng and Dr. Chenxin Li for their

advice and comments on this work, and Dr. Thomas Wilkop for his superb microscopic support.

Competing interests

The authors declare no other competing or financial interests.

Author contributions

Conceptualization: C.H.H., Y.-R.J.L., J.H., B.L.; Methodology: C.H.H., A.M.K., Y.-R.J.L., J.H.,

B.L.; Investigation: C.H.H., Y.-R.J.L.; Formal analysis: C.H.H., A.M.K., Y.-R.J.L.; Visualization:

C.H.H., Y.-R.J.L.; Data curation: C.H.H.; Supervision: B.L., J.H.; Project administration: B.L.;

Funding acquisition: J.H., B.L.; Writing (original draft): C.H.H., A.M.K.; Writing (review and

editing): C.H.H., A.M.K., Y.-R.J.L., J.H., B.L.

Funding

This work was supported by grants from the National Science Foundation (2148206 &

2148207). B.L. was sponsored by the U.S. Department of Agriculture (USDA)-the National

Institute of Food and Agriculture (NIFA) under an Agricultural Experiment Station (AES) hatch

project (CA-D-PLB-2536-H).

Data and resource availability

All data acquired from this study are included within the main text and Supplementary

information.

FIGURE LEGENDS

Figure 1. The *vln4* mutation enhances the phenotype of F-actin organization caused by the *vln2*
and *vln3* mutations in Arabidopsis. (A) F-actin, labeled by the LifeAct-GFP fusion in transgenic
wild-type (WT), *vln2/3*, *vln2/3/4* plants, is imaged at the cell cortex of hypocotyl epidermal cells.
The WT cell has cortical F-actin in two populations of long, thick bundles and finer filaments in
mesh network. The prominent bundles are reduced in the *vln2/3* double mutant and almost
completely lost in the *vln2/3/4* triple mutant. Maximum projections of the LifeAct-GFP images
are shown. (B) The skewness of the F-actin is measured to reflect how well the filaments are
bundled. Values were measured from n>35 cells of each genotype. (C) Quantitative assessment
of F-actin orientation. The orientation is determined with a 90° angle perpendicular to cell
elongation axis. Significance is determined by a two-tailed t-test with $p < 0.01$ (****). Scale bar: 5
µm.

Figure 2. Effects of *vln* mutations on the dynamics of F-actin in Arabidopsis. (A) Three time-
lapse images of F-actin are taken at 5-second intervals in the WT, *vln2/3*, and *vln2/3/4* cells.
Each frame is pseudo-colored and merged to reflect the changes of F-actin filaments. Stable
filaments that experience no changes after 10 seconds appear in white in the merged images.
WT cells appear to have more of these stable filaments. The double mutant *vln2/3* has fewer
stable filaments and the triple mutant *vln2/3/4* mostly has rapidly remodeling F-actin filaments.
(B) Quantification of the amount of stable F-actin shows that WT has the highest amount while
the *vln2/3* and *vln2/3/4* mutants progressively have fewer. Values are measured from n>35 cells
of each genotype. (C) Coherency of F-actin filaments is measured to determine how F-actin
filaments are aligned. Higher coherency values represent more ordered filaments in a preferred
a specific direction. (D) Comparison of F-actin density in WT, *vln2/3*, and *vln2/3/4* cells. The
density value is determined by the proportion (%) of the cell cortex occupied by F-actin.
Significance is determined by a two-tailed t-test with $p < 0.01$ (****). Scale bar: 5 µm.

Figure 3. Peroxisome motility is affected in the *vln2/3/4* mutant lacking F-actin bundling. (A)
Peroxisome and F-actin are simultaneously imaged with 5-seconds intervals. In the *vln2/3* and
vln2/3/4 mutant cells that have reduced long F-actin bundles, peroxisomes show reduced linear
movement. (B-C) Quantitative assessments of average peroxisome speed and maximum speed
in WT and *vln* mutants. Values are measured from n>300 peroxisome tracks from each
genotype and plotted along a raincloud plot with a histogram on top and a boxplot below. Yellow
diamonds represent the mean position. Significance is determined by a Wilcoxon rank sum test
with p-values shown in the figures.

Figure 4. Quantitative comparisons of peroxisome tracks in the WT, *vln2/3* and *vln2/3/4* mutant
cells. (A-E) Parameters include total distance, track displacement, average straight-line speed,

linearity, and directional change rate. Values are measured from n>300 peroxisome tracks from
each genotype and plotted along a raincloud plot with a histogram on top and a boxplot below.
Yellow diamonds represent the mean values. Significance is determined by a Wilcoxon rank
sum test with p-values shown in the figures.

Figure 5. Pairwise correlation analysis of peroxisome movement in terms of directional change
rate, average straightforward speed, and linearity. (A) Formation of three clusters following the
pairwise correlation analysis. (B) Relative proportions of peroxisome movement in the three
clusters in the WT, *vln2/3*, and *vln2/3/4* cells.

REFERENCES

Avisar, D., Prokhnevsky, A. I., Makarova, K. S., Koonin, E. V. and Dolja, V. V.
(2008). Myosin XI-K Is required for rapid trafficking of Golgi stacks, peroxisomes, and
mitochondria in leaf cells of Nicotiana benthamiana. *Plant Physiol* **146**, 1098-108.
Bao, C. C., Wang, J., Zhang, R. H., Zhang, B. C., Zhang, H., Zhou, Y. H. and Huang,
S. J. (2012). Arabidopsis VILLIN2 and VILLIN3 act redundantly in sclerenchyma
development via bundling of actin filaments. *Plant Journal* **71**, 962-975.
Covill-Cooke, C., Toncheva, V. S. and Kittler, J. T. (2021). Regulation of
peroxisomal trafficking and distribution. *Cellular And Molecular Life Sciences* **78**, 1929-
1941.
Ershov, D., Phan, M.-S., Pylvänäinen, J. W., Rigaud, S. U., Le Blanc, L., Charles-
Orszag, A., Conway, J. R. W., Laine, R. F., Roy, N. H., Bonazzi, D. et al. (2022). TrackMate
7: integrating state-of-the-art segmentation algorithms into tracking pipelines. *Nature*
Methods **19**, 829-832.
Garcia-González, J. and van Gelderen, K. (2021). Bundling up the Role of the Actin
Cytoskeleton in Primary Root Growth. *Front Plant Sci* **12**.
Haraguchi, T., Tominaga, M., Nakano, A., Yamamoto, K. and Ito, K. (2016). Myosin
XI-I is Mechanically and Enzymatically Unique Among Class-XI Myosins in Arabidopsis.
Plant Cell Physiol **57**, 1732-43.
He, A. Y., Dean, J. M. and Lodhi, I. J. (2021). Peroxisomes as cellular adaptors to
metabolic and environmental stress. *Trends in Cell Biology* **31**, 656-670.
Henty-Ridilla, J. L., Li, J. J., Blanchoin, L. and Staiger, C. J. (2013). Actin dynamics
in the cortical array of plant cells. *Current Opinion in Plant Biology* **16**, 678-687.
Higaki, T. (2017). Quantitative evaluation of cytoskeletal organizations by
microscopic image analysis. *PLANT MORPHOLOGY* **29**, 15-21.
Higaki, T., Kutsuna, N., Sano, T., Kondo, N. and Hasezawa, S. (2010).
Quantification and cluster analysis of actin cytoskeletal structures in plant cells: role of
actin bundling in stomatal movement during diurnal cycles in Arabidopsis guard cells.
Plant Journal **61**, 156-165.
Huang, C. H., Peng, F. L., Lee, Y. R. J. and Liu, B. (2024). The microtubular
preprophase band recruits Myosin XI to the cortical division site to guide phragmoplast
expansion during plant cytokinesis. *Developmental Cell* **59**, 2333-2346.e6.
Huang, S., Qu, X. and Zhang, R. (2015). Plant villins: versatile actin regulatory
proteins. *J Integr Plant Biol* **57**, 40-9.
Khurana, P., Henty, J. L., Huang, S. J., Staiger, A. M., Blanchoin, L. and Staiger, C.
J. (2010). VILLIN1 and VILLIN3 Have Overlapping and Distinct Activities in Actin Bundle
Formation and Turnover. *Plant Cell* **22**, 2727-2748.
Koenig, A. M., Liu, B. and Hu, J. P. (2023). Visualizing the dynamics of plant energy
organelles. *Biochemical Society Transactions* **51**, 2029-2040.
Li, X., Zhuang, Y. H., Zhao, W. Y., Qu, X. L., Wang, J., Chang, M., Shen, J. F., Chen,
N. Z. and Huang, S. J. (2025). Molecular and functional adaption of Arabidopsis villins.
New Phytologist **245**, 1158-1179.

Mathur, J., Mathur, N. and Hulskamp, M. (2002). Simultaneous visualization of
peroxisomes and cytoskeletal elements reveals actin and not microtubule-based
peroxisome motility in plants. *Plant Physiol* **128**, 1031-45.

**Nakagawa, T., Kurose, T., Hino, T., Tanaka, K., Kawamukai, M., Niwa, Y., Toyooka,
K., Matsuoka, K., Jinbo, T. and Kimura, T.** (2007). Development of series of gateway binary
vectors, pGWBs, for realizing efficient construction of fusion genes for plant
transformation. *J Biosci Bioeng* **104**, 34-41.

Neuhaus, A., Eggeling, C., Erdmann, R. and Schliebs, W. (2016). Why do
peroxisomes associate with the cytoskeleton? *Biochimica Et Biophysica Acta-Molecular
Cell Research* **1863**, 1019-1026.

Pan, R. H., Liu, J., Wang, S. S. and Hu, J. P. (2020). Peroxisomes: versatile
organelles with diverse roles in plants. *New Phytologist* **225**, 1410-1427.

Peremyslov, V. V., Prokhnevsky, A. I. and Dolja, V. V. (2010). Class XI myosins are
required for development, cell expansion, and F-Actin organization in Arabidopsis. *Plant
Cell* **22**, 1883-97.

Perico, C. and Sparkes, I. (2018). Plant organelle dynamics: cytoskeletal control
and membrane contact sites. *New Phytologist* **220**, 381-394.

Püspöki, Z., Storath, M., Sage, D. and Unser, M. (2016). Transforms and Operators
for Directional Bioimage Analysis: A Survey. *Focus on Bio-Image Informatics* **219**, 69-93.

Salogiannis, J. and Reck-Peterson, S. L. (2017). Hitchhiking: A Non-Canonical
Mode of Microtubule-Based Transport. *Trends in Cell Biology* **27**, 141-150.

Sandalio, L. M., Peláez-Vico, M. A., Molina-Moya, E. and Romero-Puertas, M. C.
(2021). Peroxisomes as redox-signaling nodes in intracellular communication and stress
responses. *Plant Physiology* **186**, 22-35.

Sattarzadeh, A., Schmelzer, E. and Hanson, M. R. (2011). Analysis of Organelle
Targeting by DIL Domains of the Arabidopsis Myosin XI Family. *Front Plant Sci* **2**, 72.

Schneider, C. A., Rasband, W. S. and Eliceiri, K. W. (2012). NIH Image to ImageJ:
25 years of image analysis. *Nat Methods* **9**, 671-675.

Shimmen, T. and Yokota, E. (2004). Cytoplasmic streaming in plants. *Curr Opin
Cell Biol*. **16**, 68-72.

van der Honing, H. S., Kieft, H., Emons, A. M. C. and Ketelaar, T. (2012).
Arabidopsis VILLIN2 and VILLIN3 Are Required for the Generation of Thick Actin Filament
Bundles and for Directional Organ Growth. *Plant Physiology* **158**, 1426-1438.

Wanders, R. J. A., Baes, M., Ribeiro, D., Ferdinandusse, S. and Waterham, H. R.
(2023). The physiological functions of human peroxisomes. *Physiol Rev* **103**, 957-1024.

**Yokota, E., Vidali, L., Tominaga, M., Tahara, H., Orii, H., Morizane, Y., Hepler, P. K.
and Shimmen, T.** (2003). Plant 115-kDa actin-filament bundling protein, P-115-ABP, is a
homologue of plant villin and is widely distributed in cells. *Plant and Cell Physiology* **44**,
1088-1099.