Dietary Protein Source Shapes Gut Microbial Structure and Predicted Function: A Meta-Analysis with Machine Learning

Samson Adejumo[1], Pritha Das[1], Casey Kin Yun Lim[2], Judy Malas[1], Jacob M. Allen[2], Jarrad Hampton-Marcell[1]

[1]Department of Biological Sciences, University of Illinois Chicago, Chicago, IL 60607

[2]Department of Health and Kinesiology, University of Illinois at Urbana-Champaign, Urbana, IL

Corresponding Author

Jarrad Hampton-Marcell

950 S. Halsted St.

Science & Engineering Laboratory East, Room 4283

Chicago, IL 60607

marcell2@uic.edu

Abstract

Background: Dietary proteins are major modulators of gut microbial ecology, yet the microbial signatures and functional consequences of plant- versus animal-based proteins remain poorly defined. Although digestibility and fermentation profiles differ by protein type, a systematic evaluation of how these differences shape microbial diversity, structure, and metabolic capacity is still lacking. This meta-analysis integrates machine learning and functional inference to identify microbial biomarkers and metabolic signatures associated with dietary protein source in murine models

Methods: Following PRISMA guidelines, we analyzed 16S rRNA gene sequencing data from 10 murine studies (n = 187) comparing animal and plant protein diets. Alpha diversity was assessed using multiple indices, including Shannon, Inverse Simpson, and Chao1, to capture richness and evenness across samples. Beta diversity was evaluated using Aitchison distances. LEfSe and class-weighted Random Forest models were applied to identify differentially abundant taxa and predictive microbial features. Functional potential was inferred using PICRUSt2, and taxon–pathway relationships were explored through correlation and network analysis.

Results: Plant-protein diets significantly increased gut microbial diversity across all assessed alpha diversity metrics: Chao1 (p = 5.7×10^{-11}), Shannon (p = 3.3×10^{-8}), and Inverse Simpson (p = 0.0013), reflecting enhanced species richness and evenness. These diets are associated with increased richness of 33 genera, including *Bacteroides*, *Muribaculaceae*, and *Allobaculum*, associated with SCFA production, nitrogen recycling, and bile acid metabolism. In contrast, animal-protein diets favored taxa such as *Clostridium sensu stricto* 1 and *Colidextribacter*, linked to proteolytic fermentation, ammonia, and sulfur metabolism. PCA of Aitchison distances revealed distinct microbial community structures between groups (ANOSIM R = 0.663, p < 0.001). Random Forest models achieved >88% accuracy (AUC = 0.995) in predicting dietary groups based on microbial composition. LEfSe confirmed key taxa distinguishing the dietary groups. Functional profiling revealed that plant-based diets enriched

pathways for short-chain fatty acid (SCFA) production and aromatic amino acid degradation,
while animal-based diets favored pathways involved in sulfur metabolism and branched-chain
amino acid degradation. Network analysis identified *Muribaculaceae* as a central hub in plant-
associated subnetworks linked to carbohydrate and SCFA metabolism, whereas *Lactobacillus*
dominated animal-associated subnetworks enriched for proteolysis and sulfur-related functions.
However, variation in dietary fat and fiber across studies may partially contribute to these
microbial patterns, as these covariates were not consistently reported and therefore not included
in the model.

Conclusion: Dietary protein source significantly influences gut microbiota composition and
function in murine models. This study identifies distinct taxonomic and functional signatures
associated with plant and animal protein diets, with implications for microbial nitrogen
utilization, sulfur metabolism, host–microbe interactions, and dietary modulation strategies.
However, because other diet components, such as fat and fiber content, varied across studies and
were not consistently reported, we could not include them as covariates. This represents a
limitation in isolating protein-specific effects. Future studies should use controlled designs with
matched non-protein components to address this gap

Keywords: Gut microbiota, dietary protein, plant-based diet, animal-based diet, 16S rRNA
sequencing, PICRUSt2, Random Forest, LEfSe, meta-analysis, microbial metabolism.

1.0 Introduction
The gut microbiome is a complex and dynamic ecosystem that engages in bidirectional
interactions with host nutrition and immunity and plays a critical role in metabolic health (Wu *et*
al., 2011). While the effects of dietary carbohydrates, particularly in promoting saccharolytic
fermentation and short-chain fatty acid (SCFA) production, are well established, the influence of
dietary protein on gut microbial ecology is less clearly defined (Rowland *et al.,* 2018). Dietary
proteins that escape digestion in the upper gastrointestinal tract enter the colon, where they
undergo fermentation by resident microbes. The outcomes of this process vary by protein source
and depend on the compositional and functional capacities of the gut microbiota (Scott *et al.,*
2013; Beaumont *et al.,* 2017).

Microbial protein metabolism can proceed through saccharolytic- or proteolytic-associated
pathways. Saccharolytic fermentation of amino acids can generate SCFAs that support epithelial
barrier integrity and modulate host immunity (Canfora *et al.,* 2019). In contrast, proteolytic
degradation produces nitrogenous compounds such as ammonia, hydrogen sulfide, and phenolic
metabolites. These byproducts have been associated with microbial imbalance, epithelial
damage, and inflammatory responses (Windey *et al.,* 2012; Rooks & Garrett, 2016). Together,
these distinct metabolic routes suggest that the net impact of protein fermentation is shaped by
both substrate availability and microbial community composition.

Importantly, not all protein sources influence the microbiota equally. Plant proteins are often
accompanied by fiber and polyphenols, which foster saccharolytic fermentation and SCFA
production, contributing to anti-inflammatory effects and gut barrier reinforcement (den Besten
et al., 2013; Vinelli *et al.,* 2022). However, plant proteins may also be less digestible and lower
in certain essential amino acids, potentially limiting their nutritional completeness (Scott *et al.,*
2013; Wang *et al.,* 2020; Rutherfurd *et al.,* 2015). In contrast, animal proteins provide a more

complete amino acid profile, beneficial for muscle synthesis and maintenance, particularly in
aging populations (Phillips *et al.,* 2016) but are typically higher in sulfur-containing amino acids
like methionine and cysteine. These amino acids can fuel proteolytic fermentation and lead to
elevated levels of hydrogen sulfide and ammonia, which have been associated with mucosal
inflammation and impaired gut barrier function (Kitada *et al.,* 2019; Ling *et al.,* 2023).

Despite growing interest, relatively few studies have systematically compared the microbial
responses to plant- versus animal-based proteins, particularly at both the taxonomic and
functional levels. Many investigations focus on a single protein type or infer function without
linking it to compositional shifts. This limits our ability to identify microbial biomarkers of
protein source or to understand how these shifts contribute to gut health and disease.

To address this gap, we conducted a meta-analysis of 16S rRNA gene amplicon sequencing data
from ten murine studies (n = 187), integrating statistical, machine learning, and predictive
functional profiling approaches. We assessed microbial alpha and beta diversity, taxonomic
composition, and classification accuracy using Random Forest and LEfSe. Functional inference
via PICRUSt2 was used to identify enriched metabolic pathways, while correlation and network
analyses explored taxa–function relationships. We report that plant protein consumption is
associated with greater microbial diversity and enrichment of taxa involved in SCFA production
and nitrogen recycling. Conversely, animal protein favors the enrichment of proteolytic taxa
linked to ammonia, sulfur, and phenolic metabolite production. Together, these findings offer a
taxonomic and functional framework for understanding how dietary protein source modulates the
gut microbiome, and they provide a foundation for future mechanistic studies linking diet,
microbial metabolism, and host health.

2.0 Materials and Methods

2.1 Search Strategy

This meta-analysis followed the Preferred Reporting Items for Systematic Review and Meta-
Analysis (PRISMA) guidelines (Page *et al.,* 2021) to ensure a transparent, systematic, and
reproducible study selection process. PRISMA provides a structured framework for identifying,
screening, and selecting studies in a manner that minimizes bias and enhances the reliability of
findings. A comprehensive literature search was conducted to identify murine studies published
in different databases from inception to March 31, 2024. Peer-reviewed, English-written articles
were retrieved from (1) Web of Science, (2) PubMed, (3) MEDLINE Complete, (4) Scopus, (5)
Google Scholar, and (6) manual searching. The Boolean search string used was: *("protein*
fermentation" OR "protein digestion" OR "protein intake" OR "dietary protein" OR "protein
metabolism" OR "protein consumption" OR "protein diet") AND ("microbiome" OR
"microflora" OR "microbial" OR "commensal" OR "microbiota" OR "bacteria") AND (
"gut" OR "intestinal" OR "gastrointestinal" OR "intestine" OR "digestive tract" OR "GIT"
OR "enteric" OR "ileum" OR "colon" OR "colonic" OR "duodenum" OR "jejunum" OR
"fecal" OR "faecal") AND ("mice" OR "rat" OR "rats" OR "mouse" OR "rodent" OR
"rodents" OR "murine").

2.2 Selection Criteria

The study selection process followed PRISMA guidelines and consisted of four phases: (1)
Identification of relevant records, (2) Screening of abstracts for eligibility, (3) Full-text eligibility
assessment, and (4) Final inclusion in the meta-analysis. The PRISMA flow diagram (**Figure 1**)

summarizes the number of records identified, screened, excluded, and retained for analysis.
Studies were included if they met the following criteria: (1) involved murine models (mice or
rats), (2) examined dietary protein interventions, (3) utilized 16S rRNA gene sequencing, (4) had
clearly distinguishable treatment and control groups, and (5) provided gut microbiota
composition data. Studies were excluded if they were editorials, letters, reviews, comments, or
other meta-analyses, lacked a protein intervention, involved non-murine or in vitro studies, or
had unpublished data or unclear dietary group differentiation. Search results were exported to
Mendeley for initial deduplication, followed by a second deduplication in Zotero. Two reviewers
independently screened titles, abstracts, and full-text articles for eligibility. Data were extracted
in duplicate using a standardized form. Discrepancies were resolved through discussion until
consensus was reached.

2.3 Bioinformatics and Data Processing

Raw sequencing data (FASTQ files) were retrieved from the NCBI and ENA repositories using
SRA accession numbers reported in the included studies. Amplicon sequence processing was
conducted using QIIME2-amplicon-2024.5 (Bolyen *et al.,* 2019; Ziemski *et al.,* 2022). Reads
were demultiplexed and quality-filtered using the q2-demux plugin, followed by denoising with
DADA2 (v1.28.0) to generate high-resolution amplicon sequence variants (ASVs) (Callahan *et*
al., 2016). Taxonomic assignment was performed against the SILVA-138 reference database
using a Naïve Bayes classifier implemented via the q2-feature-classifier plugin (Bokulich *et al.,*
2018). ASVs with confidence scores below 70% were discarded to improve taxonomic
reliability.

Processed data, including feature tables, phylogenetic trees, and taxonomic classifications, were
imported into RStudio (v4.4.1) for downstream analyses. Microbiome data were integrated using
the phyloseq package (v1.48.0), and statistical analyses were conducted using ggplot2 (v3.5.1),
patchwork (v1.3.0), ggpubr (v0.6.0), pheatmap (v1.0.12), and dplyr (v1.1.4) (McMurdie &
Holmes, 2012; Wen et al., 2023).

Functional metagenomic prediction was performed using PICRUSt2 (v2.5.2), which infers
Kyoto Encyclopedia of Genes and Genomes (KEGG) Orthologs (KOs) and MetaCyc pathways
from 16S rRNA gene data via evolutionary modeling. The predicted KO and pathway tables
were normalized to relative abundance and exported for downstream differential abundance
testing and network analysis

Among the included studies, nine targeted the V3–V4 region of the 16S rRNA gene using either
the 338F/806R or 515F/806R primer pairs. Both forward primers amplify the V3 region, but
338F (5′-ACTCCTACGGGAGGCAGCAG-3′) is known for its broad bacterial coverage in
murine gut microbiota, while 515F (5′-GTGCCAGCMGCCGCGG-3′) is commonly used in
high-throughput studies due to its compatibility with barcoding and multiplex sequencing
(Parada et al., 2016). The reverse primer 806R (5′-GGACTACHVGGGTWTCTAAT-3′) was
used consistently across all V3–V4 studies. One study amplified the V1–V2 region using 27F
(5′-AGAGTTTGATCMTGGCTCAG-3′) and 338R (5′-TGCTGCCTCCCGTAGGAGT-3′),
which target a more upstream and variable portion of the gene. Primer selection can influence
sequencing depth, taxonomic bias, and classification accuracy (Klindworth *et al.,* 2013; Parada *et*
al., 2016), so all primer sequences and target regions are summarized in **Supplementary Table**
S1

2.4 Statistical Analysis

2.4.1 Alpha and Beta Diversity

Alpha diversity indices (Shannon, Simpson, Chao1) were used to assess microbial richness and diversity (Li *et al.,* 2022; Zaheer *et al.,* 2018; Deng *et al.,* 2022). Comparisons between plant- and animal-protein diets were performed using the Wilcoxon Rank-Sum test. Beta diversity was assessed using Aitchison distance (Euclidean distance on centered log-ratio [CLR]–transformed data) and visualized via Principal Component Analysis (PCA). Analysis of Similarities (ANOSIM) tested whether microbial composition varied more between dietary groups than within groups (Samuthpongtorn *et al.,* 2022). Permutational analysis of variance (PERMANOVA) was conducted to quantify the variance in microbial composition explained by dietary protein sources (Tang *et al.,* 2016).

2.4.2 Generalized Linear Mixed Model (GLMM)

To account for inter-study heterogeneity and sampling variability, a Generalized Linear Mixed Model (GLMM) was implemented using the lme4 (v1.1-34) package in R. Dietary protein source and protein concentration were modeled as fixed effects, while random intercepts were specified for country of origin, sample isolation site (fecal, colonic, ileal, small intestine), and animal age (in weeks) to control for study-level and anatomical variation. Different model configurations were evaluated, and final model selection was guided by goodness-of-fit statistics, including Akaike Information Criterion (AIC) and Bayesian Information Criterion (BIC). This modeling approach enabled robust estimation of microbial diversity metrics while accounting for clustering and potential confounding across datasets (Madi *et al.,* 2023; Sweeny *et al.,* 2023).

2.4.3 Machine Learning Classification and Microbial Feature Selection

To classify dietary protein source and identify microbial biomarkers, a supervised Random Forest (RF) model was implemented using the randomForest package in R (Manandhar *et al.,* 2021). Three RF approaches were compared: (1) unadjusted RF, trained on the original imbalanced dataset; (2) class-weighted RF, which accounted for class imbalance by assigning inverse frequency weights to underrepresented groups; and (3) synthetic sampling RF, using the ROSE (Random OverSampling Examples) technique to balance the dataset.

Microbial relative abundance data at the genus level were used as input features. The dataset was randomly partitioned into training (70%) and testing (30%) sets. Model performance was evaluated using the out-of-bag (OOB) error rate, confusion matrices, and area under the receiver operating characteristic curve (AUC). Feature importance was determined using the Mean Decrease in Accuracy (MDA), which ranked genera based on their contribution to classification performance

To complement the RF analysis, LEfSe (Linear Discriminant Analysis Effect Size) was performed using the microbiomeMarker package (v1.10.0) in R to independently identify microbial taxa significantly enriched in plant- versus animal-protein diets. LEfSe was applied to genus-level relative abundance data, using an LDA score threshold >3.0 and adjusted p-value <0.05. A taxonomic cladogram was generated to visualize differentially enriched microbial features between dietary groups (Segata *et al.,* 2011)

2.4.4 Receiver Operating Characteristic (ROC) Analysis

Receiver Operating Characteristic (ROC) analysis was used to assess the predictive performance of the three Random Forest (RF) models. ROC curves were generated using the pROC package (v1.18.0) in R. Each model was trained to classify diets based on microbial genus-level profiles, distinguishing between plant and animal protein sources. Predictions from the test dataset were used to generate ROC curves, and the Area Under the Curve (AUC) was computed to quantify model performance. Higher AUC values indicated stronger sensitivity–specificity trade-offs and improved classification capability. Overall classification accuracy was also reported to compare performance across sampling strategies

ROC analysis is commonly used in microbiome studies to assess classification models, particularly in dietary and clinical interventions (Zhou *et al.,* 2019; Yassour *et al.,* 2016). Evaluating multiple sampling strategies—including the original data distribution, class weighting, and synthetic resampling—provides a more robust assessment of model reliability and reduces the risk of overfitting (Topçuoğlu *et al.,* 2020)

2.4.5 Functional Profiling and Pathway Analysis

Functional metagenomic predictions were generated using PICRUSt2 (Douglas *et al.,* 2020), which infers gene family abundances from 16S rRNA data by leveraging ancestral-state reconstruction of known reference genomes. KEGG Orthologs (KOs) were predicted and mapped to MetaCyc and KEGG pathway hierarchies to enable downstream pathway-level analysis.

Differential pathway abundance between plant and animal protein diets was analyzed using MaAsLin2 (Mallick *et al.,* 2021), a multivariable linear modeling approach that allows covariate adjustment and correction for multiple testing. Relative abundances were log transformed, and protein source was included as a fixed effect. Pathways with false discovery rate (FDR)–adjusted p-values below 0.05 were considered statistically significant.

To visualize functional differences, heatmaps were generated using the pheatmap R package, focusing on the top-ranked differentially abundant pathways. Additional analysis focused on metabolic pathways related to sulfur metabolism, short chain fatty acid (SCFA) production, and nitrogen cycling. Taxa and pathway relationships were examined using Spearman correlation, and network topology metrics such as node centrality and clustering coefficients were used to characterize ecological interactions.

2.4.6 Network Analysis

To investigate ecological interactions between microbial taxa and predicted functional pathways, correlation-based co-occurrence networks were constructed. Genus-level relative abundances and KEGG Orthologs predicted by PICRUSt2 were used as input. Pairwise Pearson correlation coefficients were computed, and significant associations were retained based on a p-value threshold of < 0.05 and an absolute correlation coefficient ($|r|$) ≥ 0.4, consistent with prior network-based microbiome studies (Kuntal *et al.,* 2019; Liu *et al.,* 2023).

Networks were visualized using the igraph and ggraph R packages. In each network, nodes represented microbial genera or functional pathways, and edges indicated significant positive or negative correlations. Edge thickness corresponded to the strength of the correlation.

To further characterize the structure of these networks, topological features were assessed using centrality measures, including *degree*, *betweenness*, and *eigenvector centrality*. Degree centrality reflects the number of direct connections a node has, indicating its immediate influence or connectivity within the network. Betweenness centrality captures how often a node lies on the shortest paths between other nodes, thus highlighting its role as a potential bridge or bottleneck in network flow. Eigenvector centrality assigns importance based not only on the number of connections but also on the influence of connected neighbors, identifying nodes with broader systemic influence. These measures were used to identify potential keystone taxa and central metabolic pathways that may play critical roles in shaping the microbiome's structure and function (Faust & Raes, 2016; Liu *et al.,* 2019).

This integrative approach allowed for the simultaneous analysis of microbial community structure and functional potential, revealing clusters of co-associated taxa and pathways shaped by dietary protein source.

3.0 Results

3.1 Characteristics of Selected Studies

The systematic search identified a total of 5,380 articles from PubMed (n = 339), EMBASE (n = 540), Scopus (n = 2,741), Web of Science (n = 1,710), and manual searches (n = 50). After de-duplication and preliminary screening, 2,696 abstracts were excluded due to duplicates (n = 982), review articles (n = 1,188), conference abstracts (n = 309), meta-analyses (n = 8), and letters or editorials (n = 209).

Subsequently, 2,684 full-text articles were assessed for eligibility. Of these, 2,668 were excluded for the following reasons: lack of an appropriate control group (n = 237), use of non-murine models (n = 1,761), inability to retrieve sequencing data (n = 534), or absence of a defined protein intervention (n = 136).

Ultimately, 10 studies met all inclusion criteria and were included in the meta-analysis (Masuoka *et al.,* 2020; Xie *et al.,* 2020; Ijaz *et al.,* 2020; Ruocco *et al.,* 2020; Choi *et al.,* 2021; Tan *et al.,* 2022; Yin *et al.,* 2022; Zhu *et al.,* 2022; Zhong *et al.,* 2022; Zhang *et al.,* 2022). These studies provided accessible 16S rRNA sequencing data, clearly defined animal or plant-based protein interventions, and complete sample metadata. Although nine distinct dietary protein sources were represented—branched chain amino acids (BCAAs), beef, casein, chicken, essential amino acids (EAAs), low protein diets, pork, sausage, and soy protein—they were categorized into two main groups for analysis: animal based proteins (beef, casein, chicken, pork, sausage, and BCAAs/EAAs when derived from animal sources) and plant based proteins (soy and low protein diets, as applicable). This binary classification enabled a focused comparison of microbial responses to protein origin across studies. The study selection process is summarized in Figure 1.

3.2 Composition of Core Gut Bacteria in Protein-fed mice

A total of 9,575,175 sequencing reads were generated from 187 protein fed mouse samples, yielding 16,630 amplicon sequence variants (ASVs), with an average of 51,204 sequence reads per sample. After denoising and chimera removal, 10,123 ASVs remained. Across all samples, Firmicutes (58.48%) and Bacteroidota (21.88%) were the dominant phyla, followed by Proteobacteria (7.08%), Actinobacteriota (4.13%), Acidobacteriota (1.34%), and Verrucomicrobiota (1.28%). The Firmicutes to Bacteroidota (F:B) ratio, a traditional indicator of

metabolic health, was significantly higher in animal protein fed mice (63.3:19.9) compared to plant protein fed mice (47.3:23.5) (**Supplementary Table 1**, **Supplementary Figure 2**).

Among the 685 identified genera, plant protein diets supported a significantly higher number of unique microbial taxa (408 genera, 59.6%) compared to animal protein diets (71 genera, 10.4%) (**Figure 2A**), indicating greater microbial diversity. Despite this difference, both dietary groups shared a core set of dominant taxa, with *Muribaculaceae*, *Faecalibaculum*, and *Akkermansia* highly abundant across all samples, though their relative abundance varied by diet.

Muribaculaceae, a key saccharolytic fermenter, was more abundant in plant protein fed mice (35.64%) than in animal protein fed mice (21.93%), suggesting a preference for plant derived substrates. Conversely, animal-protein diets were enriched in *Lactococcus* (7.34%), *Ligilactobacillus* (4.24%), and *Staphylococcus* (3.25%), taxa associated with proteolytic fermentation and potential pro-inflammatory metabolite production. Plant-protein diets also exhibited higher relative abundances of *Dubosiella* (4.79%), *Allobaculum* (4.79%), *Bacteroides* (3.02%), and *Alloprevotella* (2.15%), taxa known for fermenting complex carbohydrates and supporting gut barrier integrity (Figure 2D; Table 2). Furthermore, plant-protein diets supported a greater number of low-abundance taxa, indicating enhanced ecological richness and metabolic versatility (**Figure 2B**).

3.3 Dietary Protein Sources Modulate Microbial Diversity

To assess how dietary protein source, protein concentration, and other contextual factors affect microbial diversity, we employed a generalized linear mixed model (GLMM) framework as described in Section 2.4.2. The best-fitting model included dietary protein source and protein concentration as fixed effects, with random intercepts specified for country of origin, sample isolation site, and animal age (in weeks). This model achieved the lowest AIC (252.88) and BIC (278.20), and the highest log-likelihood (-118.44) (Supplementary Table 2). Model selection was further validated by likelihood ratio tests and R^2 metrics, revealing a marginal R^2 of 0.071 and a conditional R^2 of 0.893, indicating substantial contribution from both fixed and random effects **(Figure 3A).** Protein source had the strongest positive association with Shannon diversity ($\beta = 1.25$), followed by protein level ($\beta = 0.40$). Sequencing depth did not significantly impact diversity estimates ($\rho = 0.107$, p = 0.145), suggesting biological rather than technical drivers of variation **(Figure 3B).**

Consistent with these model outputs, microbial alpha diversity was significantly higher in plant protein diets across all three diversity indices: Chao1 ($p = 5.7 \times 10^{-14}$), Shannon ($p = 3.3 \times 10^{-11}$), and Inverse Simpson ($p = 0.0013$) (**Figure 3C**). A notable interaction between protein source and concentration was detected (Supplementary Figure 3), with Shannon diversity highest in high plant-protein diets ($p = 7.39 \times 10^{-12}$), while differences were not significant at low protein levels ($p = 0.722$). These results suggest that both the source and concentration of dietary protein contribute to microbial richness and evenness.

To further examine shifts in overall microbial community composition, we analyzed beta diversity using Aitchison distance-based Principal Component Analysis (PCA). The ordination revealed distinct clustering by protein source, with the first two components explaining 10.58% and 10.37% of the total variance, respectively (**Figure 3D**). These compositional differences were confirmed by PERMANOVA ($R^2 = 0.055$, $p = 0.001$) and ANOSIM ($R = 0.663$, $p = 0.001$),

underscoring significant divergence in microbial communities between plant- and animal-protein
diets (**Figure 3E**).

Together, these findings demonstrate that dietary protein source exerts a dominant influence on
both within-sample (alpha) and between-sample (beta) microbial diversity, independent of
sequencing depth or study-specific factors. High plant protein intake supports a more diverse and
compositionally distinct gut microbiota, implicating protein source as key in modulating the gut
microbial ecosystem.

3.4 Machine Learning Identifies Microbial Predictors of Protein Source

Supervised machine learning confirmed that gut microbial composition is highly predictive of
dietary protein source. Among the three Random Forest (RF) models evaluated, the unadjusted
RF yielded the highest classification performance (AUC = 1.00, Accuracy = 98%), followed by
the ROSE-sampled RF (AUC = 0.997, Accuracy = 92.2%) and the class-weighted RF (AUC =
0.995, Accuracy = 88.2%) (**Figure 4A**).

Despite its slightly lower accuracy, the class-weighted model was selected for downstream
feature selection to correct for class imbalance between plant- and animal-protein groups.
Feature importance analysis based on Mean Decrease Accuracy (MDA) identified key taxa
contributing to classification, including *Lactococcus* (5.34%), *Streptococcus* (5.09%, 4.34%),
Peptococcus (4.71%), *Bifidobacterium* (4.56%), *Parasutterella* (4.47%), *Faecalibaculum*
(4.43%), *Coriobacteriaceae UCG-002* (4.44%), and *Muribaculaceae* (4.10%, 4.06%) (**Figure
4B**).

Model robustness was supported by cross-validation and comparison with a randomized control
model, which achieved only 38.2% accuracy (**Supplementary Figure 4A, 4C**). Confusion
matrix analysis showed high specificity for plant-protein predictions, with modest
misclassification among animal-protein samples (Supplementary Figure 4B).

LEfSe analysis further validated these findings, identifying 33 genera enriched in plant-protein
diets and 7 in animal-protein diets. Notably, plant-associated genera included *Muribaculaceae,*
Bacteroides, Parabacteroides, and *Allobaculum*, while animal-protein diets were characterized
by genera such as *Clostridium sensu stricto* 1*, Colidextribacter,* and *Rikenellaceae.* A LEfSe-
derived cladogram illustrated the phylogenetic structure of these associations, with plant-protein
diets favoring Bacteroidota and Proteobacteria, and animal-protein diets skewed toward
Firmicutes dominance (**Figure 4C–D**). These results demonstrate the use of machine learning for
uncovering diet-responsive microbial biomarkers and highlight the biological relevance of
microbial shifts linked to dietary protein type.

3.5 Functional Pathways Enriched by Protein Type

Functional profiling using PICRUSt2 revealed distinct metabolic signatures between plant- and
animal-protein diets. Differential pathway enrichment analysis identified 10 plant-enriched and
10 animal-enriched MetaCyc pathways with significant log□ fold changes (**Figure 5A**). Plant-
based diets were associated with pathways related to carbohydrate metabolism and amino acid
degradation, including the Entner-Doudoroff pathway, nitrate reduction, aromatic biogenic
amine degradation, and L-tryptophan degradation. Conversely, animal-protein diets were

enriched in fermentation and biosynthesis pathways such as creatinine degradation, propanoate fermentation, and L-isoleucine biosynthesis. These findings highlight distinct metabolic capacities of the microbiota shaped by dietary protein source.

3.6 Correlation and Network Analysis Reveal Functional Interactions

To investigate how enriched pathways relate to specific microbial taxa, a correlation analysis was performed between predicted KEGG Ortholog functions and the top differential genera. The resulting heatmap **(Figure 5B)** highlighted strong associations between specific microbial genera and functional pathways involved in SCFA, BCFA, sulfur, and amino acid metabolism. *Lactobacillus* correlated with cysteine biosynthesis ($r = 0.693$, $p = 6.5 \times 10^{-2\square}$), sulfate reduction ($r = 0.647$, $p = 7.1 \times 10^{-23}$), and methionine biosynthesis ($r = 0.566$, $p = 2.4 \times 10^{-1\square}$). *Faecalibaculum* showed positive correlations with methionine biosynthesis ($r = 0.606$, $p = 1.4 \times 10^{-2\square}$) and cysteine biosynthesis ($r = 0.602$, $p = 2.7 \times 10^{-2\square}$). *Dubosiella* was linked to glutamate degradation ($r = 0.440$, $p = 2.5 \times 10^{-1\square}$) and cysteine biosynthesis ($r = 0.537$, $p = 1.8 \times 10^{-1\square}$). *Bifidobacterium* was moderately correlated with methionine pathways, including S-adenosylmethionine biosynthesis ($r = 0.503$, $p = 3.8 \times 10^{-13}$). In contrast, *Alistipes* and *Muribaculaceae* were associated with SCFA-related pathways, including lysine fermentation ($r = 0.619$, $p = 8.5 \times 10^{-22}$ and $r = 0.582$, $p = 5.6 \times 10^{-1\square}$, respectively), glutamate degradation ($r = 0.509$, $p = 2.3 \times 10^{-1\square}$ and $r = 0.855$, $p = 2.1 \times 10^{\square\square\square}$), and valine biosynthesis ($r = 0.422$, $p = 4.3 \times 10^{-1\square}$ and $r = 0.858$, $p = 7.6 \times 10^{\square\square\square}$). These patterns suggest that distinct microbial taxa specialize in different metabolic niches depending on dietary protein source.

Network analysis based on pairwise Pearson correlations ($|r| \geq 0.4$, $p < 0.05$) visualized the co-occurrence structure of microbial taxa and metabolic functions **(Figure 5C).** Nodes represented taxa or pathways, and edge thickness indicated correlation strength. Highly connected nodes such as *Faecalibaculum* (eigenvector centrality = 0.97) and *Lactobacillus* (0.97) exhibited strong topological influence within sulfur and methionine metabolic subnetworks, suggesting they may serve as functional keystone taxa in modulating amino acid–related metabolic functions.

Similarly, *Muribaculaceae* demonstrated the highest eigenvector centrality (1.00) **(Supplementary Figure 5),** underscoring its potential integrative role across diverse metabolic modules, particularly those involved in glutamate degradation and SCFA production. These network metrics reinforce the centrality of select taxa in shaping key nutrient pathways influenced by dietary protein source.

4.0 Discussion

This meta-analysis integrates 16S rRNA datasets across multiple murine studies to provide a comprehensive overview of how dietary protein sources are associated with gut microbial composition, diversity, and functional potential. In addition to compositional differences, we identify consistent ecological and metabolic patterns linked to saccharolytic versus proteolytic fermentation, highlighting microbial taxa and pathways that co-vary with protein source (De Filippis *et al.,* 2016; Canfora *et al.,* 2019). These findings deepen our understanding of how variation in dietary nitrogen and carbon availability corresponds with shifts in microbial community structure and function, with potential implications for host health.

4.1 Dietary Protein Source Impacts Microbial Diversity and Community Structure

Consistent with prior findings (Singh *et al.,* 2017; Gilbert *et al.,* 2018), plant-protein diets were
associated with greater microbial alpha diversity, reflecting richer and more evenly distributed
microbial communities. Generalized Linear Mixed Model (GLMM) analysis identified protein
source as the strongest predictor of alpha diversity, while protein level (% kcal), country, sample
site, and host age also contributed significantly. The high conditional R^2 underscores the
combined influence of biological and environmental variables on microbial richness (Madi *et al.,*
2023; Sweeny *et al.,* 2023). Importantly, sequencing depth did not significantly impact diversity
estimates, confirming that observed patterns were biologically meaningful rather than artifacts of
sampling depth (Zaheer *et al.,* 2018; Deng *et al.,* 2022).

Beta diversity analysis revealed further separation in community structure between diet groups.
Principal component analysis (PCA) showed distinct clustering, supported by significant
PERMANOVA and ANOSIM results. These findings align with previous studies reporting that
plant-based diets promote more heterogeneous microbial communities, whereas animal-protein
diets tend to support a narrower microbiome enriched in nitrogen-metabolizing taxa (Wu *et al.,*
2011; Johnson *et al.,* 2019). Together, these results suggest that protein source not only shapes
microbial richness but also drives functionally distinct ecological communities.

4.2 Microbial Biomarkers Reflect Functional Adaptation to Protein Type

Using both Random Forest and LEfSe analyses, we identified microbial taxa that reliably
distinguished between plant- and animal-protein diets. For instance, *Muribaculaceae*,
Faecalibaculum, and *Parasutterella* were consistently enriched under plant-protein conditions,
consistent with their known roles in saccharolytic fermentation and nitrogen salvage.
Muribaculaceae is a dominant and functionally diverse group in murine gut ecosystems, often
associated with polysaccharide utilization and propionate production (Ormerod *et al.,* 2016). Its
enrichment here supports prior reports linking it to fiber-rich and lower-protein diets.
Faecalibaculum, another SCFA-producing genus, has been implicated in carbohydrate
fermentation and maintenance of gut barrier function (Zhu *et al.,* 2020). *Parasutterella*, though
less frequently studied, has been linked to bile acid metabolism and appears responsive to plant-
derived nitrogen sources, suggesting potential roles in amino acid salvage under lower-protein
conditions.

In contrast, genera such as *Bifidobacterium*, *Streptococcus*, and *Colidextribacter* were more
abundant in animal-protein-fed mice, likely reflecting a microbial community adapted to
proteolytic substrates. While *Bifidobacterium* is traditionally associated with fiber metabolism,
certain species are capable of amino acid fermentation and expand in protein-rich, low-fiber
contexts (Scott *et al.,* 2017). *Streptococcus* may utilize simple nitrogenous compounds and
peptides from dietary or host sources, while *Colidextribacter* has recently been associated with
branched-chain and aromatic amino acid fermentation in murine models.

While Random Forest provided high predictive accuracy, LEfSe highlighted taxa with consistent
enrichment and biological effect sizes (Segata *et al.,* 2011). Together, these findings suggest that
dietary protein type selectively enriches microbial lineages with distinct metabolic strategies.
Our study builds on prior research by linking taxonomic signatures to predicted metabolic
pathways, highlighting these taxa as potential biomarkers of diet-associated functional shifts in
the gut ecosystem (Topçuoğlu *et al.,* 2020).

4.3 Dietary Protein Source Shapes Gut Microbial Metabolism and Functional Networks

Functional prediction using PICRUSt2 showed that plant-based and animal-based protein diets influence the metabolic potential of gut microbes in distinct ways. Diets rich in plant protein increased the abundance of pathways related to carbohydrate metabolism, nitrogen recycling, and the degradation of aromatic amino acids. These included the Entner–Doudoroff pathway, nitrate reduction, and L-tryptophan degradation, suggesting a microbial community better adapted to environments with more fiber and limited nitrogen (Fu *et al.,* 2022; Diether *et al.,* 2019). These functions are often associated with enhanced metabolic flexibility, allowing microbes to utilize a broader range of substrates, especially under conditions of nutrient limitation (Zhao *et al.,* 2019; Scott *et al.,* 2013). Additionally, pathways such as nitrate reduction provide alternative electron acceptors, while ArAA degradation contributes to detoxification and redox balancing, potentially reducing the accumulation of harmful fermentation byproducts such as ammonia, hydrogen sulfide, and p-cresol, which are more commonly associated with proteolytic metabolism (Windey *et al.,* 2012; Dai *et al.,* 2011). Thus, the enrichment of these pathways in plant-protein diets may reflect a functional shift toward a more ecologically balanced and less putrefactive microbial community. In contrast, diets enriched in animal protein promoted microbial pathways involved in proteolytic fermentation and adaptation to nitrogen-rich environments. These included creatinine degradation, propionate fermentation, and branched-chain amino acid (BCAA) biosynthesis—metabolic routes that have been linked to the generation of pro-inflammatory metabolites such as ammonia, branched-chain fatty acids, and p-cresol (Wu *et al.,* 2022; Ridlon *et al.,* 2016; Windey *et al.,* 2012). The enrichment of these functions suggests a microbial metabolic profile that favors nitrogen disposal and amino acid fermentation, processes often associated with gut barrier disruption and systemic inflammation in both animal and human studies

Network analysis further illustrated the ecological organization of these functional traits. Correlation-based mapping identified *Muribaculaceae* and *Desulfovibrio* as central nodes within plant-protein-associated subnetworks enriched in SCFA and sulfate metabolism, while *Lactobacillus* emerged as a key hub in amino acid and sulfur biosynthesis modules under animal-protein conditions (Faust & Raes, 2016; Liu *et al.,* 2021).These results suggest that dietary protein source not only alters microbial composition but reprograms functional networks, shaping key outputs like SCFA, BCFA, p-cresol, and ammonia (Neis et al., 2015; Canfora et al., 2019). Understanding these specific pathways is critical for linking diet to gut microbial ecology and host physiology

4.4 Functional Implications of Protein-Driven Microbial Shifts

Distinct microbial communities shaped by plant- and animal-based protein diets exhibited markedly different metabolic profiles, particularly in pathways involved in short-chain fatty acid (SCFA), branched-chain fatty acid (BCFA), ammonia, sulfur, and phenol metabolism. These differences reflect not only microbial adaptation to the available nutrient substrates but also carry important consequences for host metabolic health and immune regulation.

SCFA metabolism was most strongly enriched in plant-protein-fed microbiomes. Taxa such as *Muribaculaceae*, *Alistipes*, and *Bacteroides*, which were more abundant under plant-protein diets, correlated positively with glutamate degradation and lysine fermentation pathways—key routes for the production of propionate and butyrate. SCFAs are essential for maintaining intestinal barrier integrity, supporting immune regulation, and providing energy to colonocytes

(Canfora *et al., 2019*; Koh *et al.,* 2016). Their anti-inflammatory and antitumor properties have
been well documented, especially in the context of fiber-rich and polyphenol-containing diets
(Ríos-Covián *et al.,* 2016; Louis & Flint, 2017). These findings suggest that plant-based proteins
potentially promote a more resilient and protective microbial environment.

In contrast, BCFA metabolism was predominantly linked to microbiomes from animals fed
animal-based protein diets. Microbes such as *Clostridium sensu stricto* 1, *Colidextribacter*, and
Lactobacillus were associated with the fermentation of isoleucine and methionine—branched-
chain and sulfur-containing amino acids that serve as substrates for proteolytic fermentation.
Clostridium sensu stricto 1, a genus within the *Clostridia* class, harbors enzymes such as
branched-chain aminotransferases and dehydrogenases that enable the conversion of BCAAs into
short-chain fatty acids like isobutyrate and isovalerate, with ammonia as a byproduct via
Stickland fermentation (Neis et al., 2015; Bahl et al., 2022). *Colidextribacter*, though recently
classified, has been associated in murine studies with the enrichment of genes involved in
aromatic and branched-chain amino acid degradation, suggesting its role in protein catabolism
under high-protein diets (Zhao et al., 2021). Likewise, certain *Lactobacillus* species can
catabolize amino acids such as methionine and isoleucine under carbohydrate-limited conditions,
utilizing amino acid deamination pathways to generate energy while releasing BCFA and
ammonia as end-products (Pessione, 2012).

While BCFAs such as isobutyrate and isovalerate can participate in host signaling, excessive
accumulation often coincides with proteolytic fermentation and mucosal stress (Smith &
Macfarlane, 1997). These metabolites tend to increase in the absence of carbohydrate substrates,
highlighting the trade-off between microbial fermentation strategies and host health outcomes.

Ammonia metabolism followed a similar trend. Animal-protein diets supported the enrichment
of Stickland-fermenting taxa, which deaminate amino acids and increase luminal ammonia
concentrations. Excessive ammonia is cytotoxic and has been associated with impaired
colonocyte metabolism and barrier dysfunction (Ivnitsky *et al.,* 2022; Windey *et al.,* 2012). On
the other hand, plant-protein diets appeared to favor nitrogen recycling through pathways such as
glutamine synthesis. The increased abundance of *Muribaculaceae* and *Parasutterella*, both
linked to ammonia assimilation, suggests a microbial adaptation that reduces nitrogen waste and
recycles ammonia into biomass (Smith *et al.,* 2022; Ju *et al.,* 2019).

Sulfur metabolism further illustrates the contrast between dietary protein sources. Diets rich in
animal protein increased the abundance of microbial genes involved in cysteine and methionine
breakdown, which are important sources of hydrogen sulfide (H_2S). When produced in excess,
H_2S can interfere with butyrate metabolism, damage cellular DNA, and weaken the intestinal
barrier by disrupting tight junctions (Carbonero *et al.,* 2012; Teigen *et al.,* 2022). Microbes such
as *Faecalibaculum* and *Lactobacillus*, which played central roles in sulfur-associated networks,
likely contributed to elevated H_2S levels in animals consuming animal-based proteins. In
contrast, the reduced presence of these pathways in mice fed plant protein may reflect a
microbial community that favors alternative fermentation routes, which are less dependent on
sulfur metabolism and more supportive of gut barrier function.

Finally, phenol metabolism varied depending on the source of dietary protein. *Colidextribacter*,
which was more abundant in mice fed with animal-derived protein, showed strong positive

correlations with tryptophan and tyrosine degradation pathways (Figure 5B). These pathways are known to produce phenolic metabolites such as *p*-cresol and indoxyl sulfate, which are classified as uremic toxins and have been associated with systemic inflammation, gut barrier dysfunction, and kidney injury (Thuy-Boun *et al.,* 2022; Wikoff *et al.,* 2009; Mishima and Abe, 2022). These compounds can disrupt gut–liver axis signaling and are increasingly recognized as biomarkers of excessive protein fermentation and microbial imbalance (Vaziri *et al.,* 2013). In contrast, *Bacteroides*, enriched under plant-based protein diets, was correlated with pathways linked to aromatic amino acid metabolism, including those involved in the production of 4-hydroxyphenylacetate (Figure 5B, 5C). This compound has been reported to exert anti-inflammatory effects and may contribute to metabolic resilience and reduced disease risk (De Filippis *et al.,* 2016; Canfora *et al.,* 2019).

Collectively, these findings emphasize that dietary protein type not only shapes microbial taxonomic profiles but also reprograms gut metabolic outputs. Plant-protein diets foster saccharolytic and nitrogen-recycling functions, while animal-protein diets enrich for proteolytic and potentially proinflammatory metabolite production. These functional shifts provide mechanistic insights into how dietary protein may influence host physiology, inflammation, and chronic disease risk.

4.5 Implications for Diet-Microbiome Interactions

These findings highlight the ecological and functional adaptability of the gut microbiota in response to dietary protein source. Plant-based proteins appear to support metabolic flexibility, nitrogen recycling, and anti-inflammatory capacities through enrichment of saccharolytic and amino acid degradation pathways. In contrast, animal-based proteins promote proteolytic fermentation and biosynthetic routes often associated with pro-inflammatory and potentially dysbiotic profiles. Notably, animal-protein diets are typically richer in sulfur-containing amino acids such as methionine and cysteine, which are metabolized by taxa like *Clostridium*, *Desulfovibrio*, and *Colidextribacter* into hydrogen sulfide and ammonia—metabolites associated with mucosal inflammation and compromised gut barrier integrity (Zhao *et al.,* 2021; Ridlon *et al.,* 2016).

These patterns suggest that the quality of dietary protein is potentially a key modulator of microbiome-associated metabolic health. However, caution is warranted when interpreting causality, given the potential confounding effects of co-varying dietary factors such as fiber, fat, and phytochemicals that differ across studies (Johnson et al., 2019; Agus et al., 2016; Holmes et al., 2008). Although this meta-analysis identifies consistent associations between protein type and microbial ecology, differences in overall diet composition likely contribute to the observed microbial signatures. To isolate protein-specific effects, future studies should prioritize controlled interventions using matched isocaloric, isonitrogenous diets with standardized non-protein components.

Future work should also examine how these microbial shifts impact host physiology— particularly in relation to insulin sensitivity, lipid metabolism, immune function, and gut barrier integrity under conditions of metabolic or inflammatory stress. Integrative multi-omics approaches, including shotgun metagenomics, untargeted metabolomics, and host

transcriptomics, will be essential for establishing causal links between dietary protein, microbial
metabolism, and host health

4.6 Limitations and Recommendations for Future Research

While this meta-analysis offers valuable insights into how dietary protein sources shape the gut
microbiome, several limitations should be acknowledged. First, the use of murine models limits
direct translatability to humans, as differences in gut microbial composition, immune responses,
and host physiology can influence microbial interactions (Nguyen et al., 2015). However, murine
models remain a cornerstone in microbiome research due to their controlled genetics, defined
dietary environments, and tractable experimental design. By focusing on mice, we were able to
isolate the effects of dietary protein while minimizing the confounding variables, such as
baseline diet variability, medication use, and lifestyle factors, that often complicate human
studies. Thus, this analysis provides a foundational framework for identifying microbial and
functional patterns that warrant further investigation in human cohorts.

Second, the reliance on 16S rRNA gene sequencing and functional prediction tools such as
PICRUSt2 limits taxonomic resolution and cannot directly confirm gene expression or strain-
level variation. While these tools are useful for comparative functional inference, they do not
capture actual microbial activity. Additionally, the absence of fecal or systemic metabolomic
data precludes validation of microbial metabolic products such as short-chain fatty acids,
ammonia, hydrogen sulfide, and p-cresol. As such, pathway-level inferences should be
interpreted cautiously and ideally confirmed using direct metabolite profiling.

Future research should incorporate longitudinal human dietary interventions that leverage strain-
resolved metagenomics, metatranscriptomics, and untargeted metabolomics to more precisely
characterize microbiota function and host–microbiota interactions (Zhao et al., 2019). Exploring
sex-specific and age-dependent responses will also improve our understanding of how life stage
and biological sex modulate microbial responses to dietary protein.

Finally, minor methodological differences, such as variation in primer pairs (e.g., 341F vs.
515F), may introduce slight biases in taxonomic profiling. However, the consistent targeting of
the V3–V4 region across studies and standardized data processing pipelines likely mitigate such
effects on broader community and functional trends. Together, despite these limitations, our
meta-analysis offers a valuable comparative resource and highlights reproducible microbial
features and functional shifts associated with dietary protein source across diverse mouse
models.

5.0 Conclusion

This meta-analysis demonstrates that dietary protein source plays a pivotal role in shaping gut
microbial diversity, community composition, and functional capacity. Distinct taxonomic and
metabolic signatures consistently differentiate plant-based from animal-based protein diets: plant
proteins enrich saccharolytic and nitrogen-recycling taxa, whereas animal proteins promote
proteolytic and sulfur-metabolizing communities. These functional differences were most
pronounced in pathways related to short-chain fatty acid (SCFA) biosynthesis, ammonia
metabolism, and hydrogen sulfide production.

The enrichment of sulfur-metabolizing and proteolytic microbes under animal-protein diets highlights the need to consider not only overall protein type but also amino acid composition, particularly sulfur-containing amino acids, when assessing diet–microbiome interactions. Given the variability in co-varying dietary components across studies, future research should prioritize controlled feeding trials using isocaloric, isonitrogenous diets to isolate protein-specific effects.

Altogether, these findings highlight the potential of dietary protein modulation as a strategy to shape the gut microbiota in ways that support gut barrier integrity, lower inflammation risk, and enhance host metabolic health

Author Contributions

JHM and SA designed the study. SA, and PD were involved in data processing. SA analyzed the data, created figures, and wrote the manuscript. JM, CKYH and JMA reviewed the manuscript. All authors participated in interpreting data, revised the manuscript, and agreed to the published version of the manuscript.

Funding

This research received no external funding.

Data Availability Statement

Data and bioinformatic workflows are publicly available here:

https://github.com/deejayprof/META_ANALYSIS

Acknowledgements

We acknowledge the support provided by various R libraries that were instrumental in the analysis conducted for this study. Comprehensive details about the data, code, and resources utilized are provided in the data availability section.

Conflicts of Interest

The authors declare no conflicts of interest and affirm that the study's results are presented transparently, accurately, and free from any fabrication, falsification, or inappropriate data manipulation.

Abbreviations

PRISMA: Preferred Reporting Items for Systematic Review and Meta-Analysis

RF: Random Forest

LDA: Linear Discriminant Analysis

SCFAs: Short-Chain Fatty Acids

BCFAs: Branched-Chain Fatty Acids

ANOVA: Analysis of Variance

ROC: Receiver Operating Characteristic

AUC: Area Under the Curve

ACE: Abundance-based Coverage Estimator

PERMANOVA: Permutational Multivariate Analysis of Variance

PCoA: Principal Coordinates Analysis

GLMM: Generalized Linear Mixed Model

LEfSe: Linear Discriminant Analysis Effect Size

qPCR: Quantitative Polymerase Chain Reaction

BCAAs: Branched-Chain Amino Acids

ArAAs: Aromatic Amino Acids

ANOSIM - Analysis of Similarities

ASVs - Amplicon Sequence Variants

rRNA: Ribosomal Ribonucleic Acid

NCBI: National Center for Biotechnology Information

SRA: Sequence Read Archive

ENA: European Nucleotide Archive

QIIME2 - Quantitative Insights Into Microbial Ecology 2

DADA2 - Divisive Amplicon Denoising Algorithm 2

GEMs: Genome-scale metabolic models

MDA: Mean Decrease Accuracy

EAAs: Essential amino acids

PICRUST: Phylogenetic Investigation of Communities by Reconstruction of Unobserved States

Figure Legends:

Figure 1. PRISMA Flow Diagram for Study Selection summarizing the systematic search and
screening process for identifying eligible studies included in the meta-analysis.

Figure 2. Microbial composition and taxonomic uniqueness across dietary protein sources.

Figure 3. Microbial Diversity and Community Structure in Protein-Based Diets.

Figure 4. Machine Learning performance and key microbial predictors of dietary protein source

Figure 5. Functional profiling of microbial communities by protein source using PICRUSt2 predictions

Table Legends:

Table 1. Summary of Included Studies. Study characteristics, dietary interventions, sequencing platforms, and sample sizes.

Table 2. Core microbiota composition at the genus level across dietary protein sources.

References

1. Rowland, I., Gibson, G., Heinken, A., Scott, K., Swann, J., Thiele, I., & Tuohy, K. (2018). Gut microbiota functions: metabolism of nutrients and other food components. *European journal of nutrition*, *57*(1), 1–24. https://doi.org/10.1007/s00394-017-1445-8

2. Gilbert, M. S., Ijssennagger, N., Kies, A. K., & van Mil, S. W. C. (2018). Protein fermentation in the gut; implications for intestinal dysfunction in humans, pigs, and poultry. *American journal of physiology. Gastrointestinal and liver physiology*, *315*(2), G159–G170. https://doi.org/10.1152/ajpgi.00319.2017

3. Zhao, J., Zhang, X., Liu, H., Brown, M. A., & Qiao, S. (2019). Dietary Protein and Gut Microbiota Composition and Function. Current protein & peptide science, 20(2), 145–154. https://doi.org/10.2174/1389203719666180514145437

4. Ling, ZN., Jiang, YF., Ru, JN. *et al.,* Amino acid metabolism in health and disease. Sig Transduct Target Ther 8, 345 (2023). https://doi.org/10.1038/s41392-023-01569-3

5. Kitada, M., Ogura, Y., Monno, I., & Koya, D. (2019). The impact of dietary protein intake on longevity and metabolic health. *EBioMedicine*, *43*, 632–640. https://doi.org/10.1016/j.ebiom.2019.04.005

6. Vinelli, V., Biscotti, P., Martini, D., Del Bo', C., Marino, M., Meroño, T., Nikoloudaki, O., Calabrese, F. M., Turroni, S., Taverniti, V., Unión Caballero, A., Andrés-Lacueva, C., Porrini, M., Gobbetti, M., De Angelis, M., Brigidi, P., Pinart, M., Nimptsch, K., Guglielmetti, S., & Riso, P. (2022). Effects of Dietary Fibers on Short-Chain Fatty Acids and Gut Microbiota Composition in Healthy Adults: A Systematic Review. *Nutrients*, *14*(13), 2559. https://doi.org/10.3390/nu14132559

7. den Besten, G., van Eunen, K., Groen, A. K., Venema, K., Reijngoud, D. J., & Bakker, B. M. (2013). The role of short-chain fatty acids in the interplay between diet, gut microbiota, and host energy metabolism. Journal of lipid research, 54(9), 2325–2340. https://doi.org/10.1194/jlr.R036012

8. Ríos-Covián, D., Ruas-Madiedo, P., Margolles, A., Gueimonde, M., de Los Reyes-Gavilán, C. G., & Salazar, N. (2016). Intestinal Short Chain Fatty Acids and their Link with Diet and Human Health. *Frontiers in microbiology*, *7*, 185. https://doi.org/10.3389/fmicb.2016.00185

9. Page, M. J., McKenzie, J. E., Bossuyt, P. M., Boutron, I., Hoffmann, T. C., Mulrow, C. D., Shamseer, L., Tetzlaff, J. M., Akl, E. A., Brennan, S. E., Chou, R., Glanville, J., Grimshaw, J. M., Hróbjartsson, A., Lalu, M. M., Li, T., Loder, E. W., Mayo-Wilson, E., McDonald, S.,

McGuinness, L. A., … Moher, D. (2021). The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ (Clinical research ed.)*, *372*, n71. https://doi.org/10.1136/bmj.n71

10. Bolyen, E., Rideout, J. R., Dillon, M. R., Bokulich, N. A., Abnet, C. C., Al-Ghalith, G. A., Alexander, H., Alm, E. J., Arumugam, M., Asnicar, F., Bai, Y., Bisanz, J. E., Bittinger, K., Brejnrod, A., Brislawn, C. J., Brown, C. T., Callahan, B. J., Caraballo-Rodríguez, A. M., Chase, J., Cope, E. K., … Caporaso, J. G. (2019). Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nature biotechnology*, *37*(8), 852–857. https://doi.org/10.1038/s41587-019-0209-9

11. Ziemski, M., Adamov, A., Kim, L., Flörl, L., & Bokulich, N. A. (2022). Reproducible acquisition, management and meta-analysis of nucleotide sequence (meta)data using q2-fondue. *Bioinformatics (Oxford, England)*, *38*(22), 5081–5091. https://doi.org/10.1093/bioinformatics/btac639

12. Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J., & Holmes, S. P. (2016). DADA2: High-resolution sample inference from Illumina amplicon data. *Nature methods*, *13*(7), 581–583. https://doi.org/10.1038/nmeth.3869

13. Bokulich, N. A., Kaehler, B. D., Rideout, J. R., Dillon, M., Bolyen, E., Knight, R., Huttley, G. A., & Gregory Caporaso, J. (2018). Optimizing taxonomic classification of marker-gene amplicon sequences with QIIME 2's q2-feature-classifier plugin. *Microbiome*, *6*(1), 90. https://doi.org/10.1186/s40168-018-0470-z

14. Wen, T., Niu, G., Chen, T., Shen, Q., Yuan, J., & Liu, Y. X. (2023). The best practice for microbiome analysis using R. *Protein & cell*, *14*(10), 713–725. https://doi.org/10.1093/procel/pwad024

15. McMurdie, P. J., & Holmes, S. (2012). Phyloseq: a bioconductor package for handling and analysis of high-throughput phylogenetic sequence data. *Pacific Symposium on Biocomputing. Pacific Symposium on Biocomputing*, 235–246.

16. Samuthpongtorn, C., Nopsopon, T., & Pongpirul, K. (2022). Gut microbiome diversity measures for metabolic conditions: a systematic scoping review. *Journal of Physiological and Biomedical Sciences*, *33*(2). retrieved from https://li01.tci-thaijo.org/index.php/j-pbs/article/view/254538

17. Manandhar, I., Alimadadi, A., Aryal, S., Munroe, P. B., Joe, B., & Cheng, X. (2021). Gut microbiome-based supervised machine learning for clinical diagnosis of inflammatory bowel diseases. *American journal of physiology. Gastrointestinal and liver physiology*, *320*(3), G328–G337. https://doi.org/10.1152/ajpgi.00360.2020

18. Ley, R. E., Turnbaugh, P. J., Klein, S., & Gordon, J. I. (2006). Microbial ecology: human gut microbes associated with obesity. *Nature*, *444*(7122), 1022–1023. https://doi.org/10.1038/4441022a

19. Walters, W. A., Xu, Z., & Knight, R. (2014). Meta-analyses of human gut microbes associated with obesity and IBD. *FEBS letters*, *588*(22), 4223–4233. https://doi.org/10.1016/j.febslet.2014.09.039

20. De Filippis, F., Pellegrini, N., Vannini, L., Jeffery, I. B., La Storia, A., Laghi, L., Serrazanetti, D. I., Di Cagno, R., Ferrocino, I., Lazzi, C., Turroni, S., Cocolin, L., Brigidi, P.,

Neviani, E., Gobbetti, M., O'Toole, P. W., & Ercolini, D. (2016). High-level adherence to a Mediterranean diet beneficially impacts the gut microbiota and associated metabolome. *Gut*, *65*(11), 1812–1821. https://doi.org/10.1136/gutjnl-2015-309957

21. Canfora, E. E., Meex, R. C. R., Venema, K., & Blaak, E. E. (2019). Gut microbial metabolites in obesity, NAFLD and T2DM. *Nature reviews. Endocrinology*, *15*(5), 261–273. https://doi.org/10.1038/s41574-019-0156-z

22. Sweeny, A. R., Lemon, H., Ibrahim, A., Watt, K. A., Wilson, K., Childs, D. Z., Nussey, D. H., Free, A., & McNally, L. (2023). A mixed-model approach for estimating drivers of microbiota community composition and differential taxonomic abundance. *mSystems*, *8*(4), e0004023. https://doi.org/10.1128/msystems.00040-23

23. Madi, N., Chen, D., Wolff, R., Shapiro, B. J., & Garud, N. R. (2023). Community diversity is associated with intra-species genetic diversity and gene loss in the human gut microbiome. *eLife*, *12*, e78530. https://doi.org/10.7554/eLife.78530

24. Zaheer, R., Noyes, N., Ortega Polo, R., Cook, S. R., Marinier, E., Van Domselaar, G., Belk, K. E., Morley, P. S., & McAllister, T. A. (2018). Impact of sequencing depth on the characterization of the microbiome and resistome. *Scientific reports*, *8*(1), 5890. https://doi.org/10.1038/s41598-018-24280-8

25. Deng, X. L., Frandsen, P. B., Dikow, R. B., Favre, A., Shah, D. N., Shah, R. D. T., Schneider, J. V., Heckenhauer, J., & Pauls, S. U. (2022). The impact of sequencing depth and relatedness of the reference genome in population genomic studies: A case study with two caddisfly species (Trichoptera, Rhyacophilidae, *Himalopsyche*). *Ecology and evolution*, *12*(12), e9583. https://doi.org/10.1002/ece3.9583

26. Li, Z., Zhou, J., Liang, H., Ye, L., Lan, L., Lu, F., Wang, Q., Lei, T., Yang, X., Cui, P., & Huang, J. (2022). Differences in Alpha Diversity of Gut Microbiota in Neurological Diseases. *Frontiers in neuroscience*, *16*, 879318. https://doi.org/10.3389/fnins.2022.879318

27. Singh, R. K., Chang, H. W., Yan, D., Lee, K. M., Ucmak, D., Wong, K., Abrouk, M., Farahnik, B., Nakamura, M., Zhu, T. H., Bhutani, T., & Liao, W. (2017). Influence of diet on the gut microbiome and implications for human health. *Journal of translational medicine*, *15*(1), 73. https://doi.org/10.1186/s12967-017-1175-y

28. Gilbert, J. A., Blaser, M. J., Caporaso, J. G., Jansson, J. K., Lynch, S. V., & Knight, R. (2018). Current understanding of the human microbiome. *Nature medicine*, *24*(4), 392–400. https://doi.org/10.1038/nm.4517

29. Segata, N., Izard, J., Waldron, L., Gevers, D., Miropolsky, L., Garrett, W. S., & Huttenhower, C. (2011). Metagenomic biomarker discovery and explanation. *Genome biology*, *12*(6), R60. https://doi.org/10.1186/gb-2011-12-6-r60

30. Ju, T., Kong, J. Y., Stothard, P., & Willing, B. P. (2019). Defining the role of *Parasutterella*, a previously uncharacterized member of the core gut microbiota. The ISME journal, 13(6), 1520–1534. https://doi.org/10.1038/s41396-019-0364-5

31. Johnson, A. J., Vangay, P., Al-Ghalith, G. A., *et al.,* (2019). Daily sampling reveals personalized diet-microbiome associations in humans. Cell Host & Microbe, 25(6), 789-802.e5. https://doi.org/10.1016/j.chom.2019.05.005

32. Tang, Z. Z., Chen, G., & Alekseyenko, A. V. (2016). PERMANOVA-S: association test for
microbial community composition that accommodates confounders and multiple distances.
Bioinformatics   (Oxford,   England),   32(17),   2618–2625.
https://doi.org/10.1093/bioinformatics/btw311

33. Scott, K. P., Gratz, S. W., Sheridan, P. O., Flint, H. J., & Duncan, S. H. (2013). The influence
of   diet   on   the   gut   microbiota. *Pharmacological   research*, *69*(1),   52–60.
https://doi.org/10.1016/j.phrs.2012.10.020

34. Wu, G. D., Chen, J., Hoffmann, C., Bittinger, K., Chen, Y. Y., Keilbaugh, S. A., Bewtra, M.,
Knights, D., Walters, W. A., Knight, R., Sinha, R., Gilroy, E., Gupta, K., Baldassano, R.,
Nessel, L., Li, H., Bushman, F. D., & Lewis, J. D. (2011). Linking long-term dietary patterns
with   gut   microbial   enterotypes. *Science   (New   York,   N.Y.)*, *334*(6052),   105–108.
https://doi.org/10.1126/science.1208344

35. Beaumont, M., Portune, K. J., Steuer, N., Lan, A., Cerrudo, V., Audebert, M., Dumont, F.,
Mancano, G., Khodorova, N., Andriamihaja, M., Airinei, G., Tomé, D., Benamouzig, R.,
Davila, A. M., Claus, S. P., Sanz, Y., & Blachier, F. (2017). Quantity and source of dietary
protein influence metabolite production by gut microbiota and rectal mucosa gene
expression: a randomized, parallel, double-blind trial in overweight humans. *The American*
journal of clinical nutrition, *106*(4), 1005–1019. https://doi.org/10.3945/ajcn.117.158816

36. Rooks, M. G., & Garrett, W. S. (2016). Gut microbiota, metabolites and host
immunity. *Nature reviews. Immunology*, *16*(6), 341–352. https://doi.org/10.1038/nri.2016.42

37. Topçuoğlu BD, Lesniak NA, Ruffin MT, Wiens J, Schloss PD. 2020. A   Framework   for
Effective Application of Machine Learning to Microbiome-Based Classification Problems.
mBio11:10.1128/mbio.00434-20. https://doi.org/10.1128/mbio.00434-20

38. Ridlon, J. M., Harris, S. C., Bhowmik, S., Kang, D. J., & Hylemon, P. B. (2016).
Consequences of bile salt biotransformations by intestinal bacteria. *Gut microbes*, *7*(1), 22–
39. https://doi.org/10.1080/19490976.2015.1127483

39. Tan, J., Ni, D., Taitz, J., Pinget, G. V., Read, M., Senior, A., Wali, J. A., Elnour, R.,
Shanahan, E., Wu, H., Chadban, S. J., Nanan, R., King, N. J. C., Grau, G. E., Simpson, S. J.,
& Macia, L. (2022). Dietary protein increases T-cell-independent sIgA production through
changes in gut microbiota-derived extracellular vesicles. *Nature communications*, *13*(1),
4336. https://doi.org/10.1038/s41467-022-31761-y

40. Masuoka, H., Suda, W., Tomitsuka, E., Shindo, C., Takayasu, L., Horwood, P., Greenhill, A.
R., Hattori, M., Umezaki, M., & Hirayama, K. (2020). The influences of low protein diet on
the   intestinal   microbiota   of   mice. *Scientific   reports*, *10*(1),   17077.
https://doi.org/10.1038/s41598-020-74122-9

41. Yin, Y., Cai, J., Zhou, L., Xing, L., & Zhang, W. (2022). Dietary oxidized beef protein alters
gut microbiota and induces colonic inflammatory damage in C57BL/6 mice. *Frontiers in*
nutrition, *9*, 980204. https://doi.org/10.3389/fnut.2022.980204

42. Zhu, Y., He, H., Tang, Y., Peng, Y., Hu, P., Sun, W., Liu, P., Jin, M., & Xu, X. (2022).
Reno-Protective Effect of Low Protein Diet Supplemented With α-Ketoacid Through Gut
Microbiota and Fecal Metabolism in 5/6 Nephrectomized Mice. *Frontiers in nutrition*, *9*,
889131. https://doi.org/10.3389/fnut.2022.889131

43. Xie, Y., Wang, C., Zhao, D., Zhou, G., & Li, C. (2020). Processing Method Altered Mouse Intestinal Morphology and Microbial Composition by Affecting Digestion of Meat Proteins. *Frontiers in microbiology*, *11*, 511. https://doi.org/10.3389/fmicb.2020.00511

44. Choi, B. S., Daniel, N., Houde, V. P., Ouellette, A., Marcotte, B., Varin, T. V., Vors, C., Feutry, P., Ilkayeva, O., Ståhlman, M., St-Pierre, P., Bäckhed, F., Tremblay, A., White, P. J., & Marette, A. (2021). Feeding diversified protein sources exacerbates hepatic insulin resistance via increased gut microbial branched-chain fatty acids and mTORC1 signaling in obese mice. *Nature communications*, *12*(1), 3377. https://doi.org/10.1038/s41467-021-23782-w

45. Ijaz, M. U., Ahmad, M. I., Hussain, M., Khan, I. A., Zhao, D., & Li, C. (2020). Meat Protein in High-Fat Diet Induces Adipogensis and Dyslipidemia by Altering Gut Microbiota and Endocannabinoid Dysregulation in the Adipose Tissue of Mice. *Journal of agricultural and food chemistry*, *68*(13), 3933–3946. https://doi.org/10.1021/acs.jafc.0c00017

46. Zhong, W., Wang, H., Yang, Y., Zhang, Y., Lai, H., Cheng, Y., Yu, H., Feng, N., Huang, R., Liu, S., Yang, S., Hao, T., Zhang, B., Ying, H., Zhang, F., Guo, F., & Zhai, Q. (2022). High-protein diet prevents fat mass increase after dieting by counteracting Lactobacillus-enhanced lipid absorption. *Nature metabolism*, *4*(12), 1713–1731. https://doi.org/10.1038/s42255-022-00687-6

47. Zhang, R., Mu, H., Li, Z., Zeng, J., Zhou, Q., Li, H., Wang, S., Li, X., Zhao, X., Sun, L., Chen, W., Dong, J., & Yang, R. (2022). Oral administration of branched-chain amino acids ameliorates high-fat diet-induced metabolic-associated fatty liver disease *via* gut microbiota-associated mechanisms. *Frontiers in microbiology*, *13*, 920277. https://doi.org/10.3389/fmicb.2022.920277

48. Ruocco, C., Ragni, M., Rossi, F., Carullo, P., Ghini, V., Piscitelli, F., Cutignano, A., Manzo, E., Ioris, R. M., Bontems, F., Tedesco, L., Greco, C. M., Pino, A., Severi, I., Liu, D., Ceddia, R. P., Ponzoni, L., Tenori, L., Rizzetto, L., Scholz, M., … Nisoli, E. (2020). Manipulation of Dietary Amino Acids Prevents and Reverses Obesity in Mice Through Multiple Mechanisms That Modulate Energy Homeostasis. *Diabetes*, *69*(11), 2324–2339. https://doi.org/10.2337/db20-0489

49. Douglas, G. M., Maffei, V. J., Zaneveld, J., Yurgel, S. N., Brown, J. R., Taylor, C. M., Huttenhower, C., & Langille, M. G. I. (2020). PICRUSt2 for prediction of metagenome functions. *Nature Biotechnology*, 38(6), 685–688. https://doi.org/10.1038/s41587-020-0548-6

50. Mallick, H., Rahnavard, A., McIver, L. J., Ma, S., Zhang, Y., Nguyen, L. H., Tickle, T. L., Weingart, G., Ren, B., Schwager, E. H., Thompson, K. N., Wilkinson, J. E., Subramanian, A., Lu, Y., Waldron, L., Paulson, J. N., & Huttenhower, C. (2021). Multivariable association discovery in population-scale meta-omics studies. *PLOS Computational Biology*, 17(11), e1009442. https://doi.org/10.1371/journal.pcbi.1009442

51. Faust, K., & Raes, J. (2016). CoNet app: Inference of biological association networks using Cytoscape. *F1000Research*, 5, 1519. https://doi.org/10.12688/f1000research.9050.2

52. Kuntal, B. K., Mande, S. S., & Dutta, A. (2019). CompNet: A GUI based tool for comparison of multiple biological interaction networks. *BMC Bioinformatics*, 20(1), 185. https://doi.org/10.1186/s12859-019-2721-3

53. Liu, Y., Amit, G., Zhao, X., Wu, N., Li, D., & Bashan, A. (2023). Individualized network analysis reveals a link between the gut microbiome, diet intervention and Gestational Diabetes Mellitus. PLoS computational biology, 19(6), e1011193. https://doi.org/10.1371/journal.pcbi.1011193

54. Liu, F., Li, Z., Wang, X., Xue, C., Tang, Q., & Li, R. W. (2019). Microbial Co-Occurrence Patterns and Keystone Species in the Gut Microbial Community of Mice in Response to Stress and Chondroitin Sulfate Disaccharide. *International journal of molecular sciences*, *20*(9), 2130. https://doi.org/10.3390/ijms20092130

55. Windey, K., De Preter, V., & Verbeke, K. (2012). Relevance of protein fermentation to gut health. *Molecular nutrition & food research*, *56*(1), 184–196. https://doi.org/10.1002/mnfr.201100542

56. Rutherfurd, S. M., Fanning, A. C., Miller, B. J., & Moughan, P. J. (2015). Protein digestibility-corrected amino acid scores and digestible indispensable amino acid scores differentially describe protein quality in growing male rats. *The Journal of nutrition*, *145*(2), 372–379. https://doi.org/10.3945/jn.114.195438

57. Phillips, S. M., Chevalier, S., & Leidy, H. J. (2016). Protein "requirements" beyond the RDA: implications for optimizing health. *Applied physiology, nutrition, and metabolism = Physiologie appliquee, nutrition et metabolisme*, *41*(5), 565–572. https://doi.org/10.1139/apnm-2015-0550

58. Zhou, W., Sailani, M. R., Contrepois, K., Zhou, Y., Ahadi, S., Leopold, S. R., Zhang, M. J., Rao, V., Avina, M., Mishra, T., Johnson, J., Lee-McMullen, B., Chen, S., Metwally, A. A., Tran, T. D. B., Nguyen, H., Zhou, X., Albright, B., Hong, B. Y., Petersen, L., … Snyder, M. (2019). Longitudinal multi-omics of host-microbe dynamics in prediabetes. *Nature*, *569*(7758), 663–671. https://doi.org/10.1038/s41586-019-1236-x

59. Yassour, M., Vatanen, T., Siljander, H., Hämäläinen, A. M., Härkönen, T., Ryhänen, S. J., Franzosa, E. A., Vlamakis, H., Huttenhower, C., Gevers, D., Lander, E. S., Knip, M., DIABIMMUNE Study Group, & Xavier, R. J. (2016). Natural history of the infant gut microbiome and impact of antibiotic treatment on bacterial strain diversity and stability. *Science translational medicine*, *8*(343), 343ra81. https://doi.org/10.1126/scitranslmed.aad0917

60. Ridlon, J. M., Wolf, P. G., & Gaskins, H. R. (2016). Taurocholic acid metabolism by gut microbes and colon cancer. *Gut microbes*, *7*(3), 201–215. https://doi.org/10.1080/19490976.2016.1150414

61. Wu, S., Bhat, Z. F., Gounder, R. S., Mohamed Ahmed, I. A., Al-Juhaimi, F. Y., Ding, Y., & Bekhit, A. E. A. (2022). Effect of Dietary Protein and Processing on Gut Microbiota-A Systematic Review. *Nutrients*, *14*(3), 453. https://doi.org/10.3390/nu14030453

62. Blakeley-Ruiz, J. A., Bartlett, A., McMillan, A. S., Awan, A., Walsh, M. V., Meyerhoffer, A. K., Vintila, S., Maier, J. L., Richie, T. G., Theriot, C. M., & Kleiner, M. (2025). Dietary protein source alters gut microbiota composition and function. *The ISME journal*, *19*(1), wraf048. https://doi.org/10.1093/ismejo/wraf048

63. Fu, J., Zheng, Y., Gao, Y., & Xu, W. (2022). Dietary Fiber Intake and Gut Microbiota in Human Health. *Microorganisms*, *10*(12), 2507. https://doi.org/10.3390/microorganisms10122507

64. Diether, N. E., & Willing, B. P. (2019). Microbial Fermentation of Dietary Protein: An
Important Factor in Diet☐Microbe☐Host Interaction. *Microorganisms*, *7*(1), 19.
https://doi.org/10.3390/microorganisms7010019

65. Ivnitsky, J. J., Schäfer, T. V., Rejniuk, V. L., & Vakunenkova, O. A. (2022). Secondary
Dysfunction of the Intestinal Barrier in the Pathogenesis of Complications of Acute
Poisoning. *Journal of evolutionary biochemistry and physiology*, *58*(4), 1075–1098.
https://doi.org/10.1134/S0022093022040123

66. Carbonero, F., Benefiel, A. C., Alizadeh-Ghamsari, A. H., & Gaskins, H. R. (2012).
Microbial pathways in colonic sulfur metabolism and links with health and disease. *Frontiers*
in physiology, *3*, 448. https://doi.org/10.3389/fphys.2012.00448

67. Koh, A., De Vadder, F., Kovatcheva-Datchary, P., & Bäckhed, F. (2016). From dietary fiber
to host physiology: Short-chain fatty acids as key bacterial metabolites. *Cell*, 165(6), 1332–
1345. https://doi.org/10.1016/j.cell.2016.05.041

68. Ríos-Covián, D., Ruas-Madiedo, P., Margolles, A., Gueimonde, M., de los Reyes-Gavilán,
C. G., & Salazar, N. (2016). Intestinal short chain fatty acids and their link with diet and
human health. *Frontiers in Microbiology*, 7, 185. https://doi.org/10.3389/fmicb.2016.00185

69. Louis, P., & Flint, H. J. (2017). Formation of propionate and butyrate by the human colonic
microbiota. *Environmental Microbiology*, 19(1), 29–41. https://doi.org/10.1111/1462-
2920.13589

70. Zhao, L., Zhang, F., Ding, X., Wu, G., & Wu, C. (2021). Gut bacteria selectively promoted
by dietary fibers alleviate type 2 diabetes. Science, 359(6380), 1151–1156.
https://doi.org/10.1126/science.aao5771

71. Neis, E. P. J. G., Dejong, C. H. C., & Rensen, S. S. (2015). The role of microbial amino acid
metabolism in host metabolism. Nutrients, 7(4), 2930–2946.
https://doi.org/10.3390/nu7042930

72. Smith, E. A., & Macfarlane, G. T. (1997). Formation of phenolic and indolic compounds by
anaerobic bacteria in the human large intestine. Microbial Ecology, 33(3), 180–188.
https://doi.org/10.1007/s002489900048

73. Windey, K., De Preter, V., & Verbeke, K. (2012). Relevance of protein fermentation to gut
health. Molecular Nutrition & Food Research, 56(1), 184–196.
https://doi.org/10.1002/mnfr.201100507

74. Teigen, L., Biruete, A., & Khoruts, A. (2023). Impact of diet on hydrogen sulfide production:
implications for gut health. *Current opinion in clinical nutrition and metabolic care*, *26*(1),
55–58. https://doi.org/10.1097/MCO.0000000000000881

75. Thuy-Boun, P. S., Wang, A. Y., Crissien-Martinez, A., Xu, J. H., Chatterjee, S., Stupp, G. S.,
Su, A. I., Coyle, W. J., & Wolan, D. W. (2022). Quantitative Metaproteomics and Activity-
based Protein Profiling of Patient Fecal Microbiome Identifies Host and Microbial Serine-
type Endopeptidase Activity Associated With Ulcerative Colitis. *Molecular & cellular*
proteomics : MCP, *21*(3), 100197. https://doi.org/10.1016/j.mcpro.2022.100197

76. Mishima, E., & Abe, T. (2022). Gut Microbiota Dynamics and Uremic Toxins. *Toxins*, *14*(2),
146. https://doi.org/10.3390/toxins14020146

77. Vaziri, N. D., Wong, J., Pahl, M., Piceno, Y. M., Yuan, J., DeSantis, T. Z., Ni, Z., Nguyen,
T. H., & Andersen, G. L. (2013). Chronic kidney disease alters intestinal microbial
flora. *Kidney international*, *83*(2), 308–315. https://doi.org/10.1038/ki.2012.345

78. Parada, A. E., Needham, D. M., & Fuhrman, J. A. (2016). Every base matters: assessing
small subunit rRNA primers for marine microbiomes with mock communities, time series
and global field samples. *Environmental microbiology*, *18*(5), 1403–1414.
https://doi.org/10.1111/1462-2920.13023

79. Klindworth, A., Pruesse, E., Schweer, T., Peplies, J., Quast, C., Horn, M., & Glöckner, F. O.
(2013). Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-
generation sequencing-based diversity studies. *Nucleic acids research*, *41*(1), e1.
https://doi.org/10.1093/nar/gks808

80. Guoyao Wu. Amino acids in nutrition, health, and disease. *Front. Biosci. (Landmark Ed)*
2021, 26(12), 1386–1392. https://doi.org/10.52586/5032

81. Holmes, E., Loo, R. L., Stamler, J., Bictash, M., Yap, I. K., Chan, Q., Ebbels, T., De Iorio,
M., Brown, I. J., Veselkov, K. A., Daviglus, M. L., Kesteloot, H., Ueshima, H., Zhao, L.,
Nicholson, J. K., & Elliott, P. (2008). Human metabolic phenotype diversity and its
association with diet and blood pressure. *Nature*, *453*(7193), 396–400.
https://doi.org/10.1038/nature06882

82. Agus, A., Denizot, J., Thévenot, J., Martinez-Medina, M., Massier, S., Sauvanet, P.,
Bernalier-Donadille, A., Denis, S., Hofman, P., Bonnet, R., Billard, E., & Barnich, N. (2016).
Western diet induces a shift in microbiota composition enhancing susceptibility to Adherent-
Invasive E. coli infection and intestinal inflammation. *Scientific reports*, *6*, 19032.
https://doi.org/10.1038/srep19032

83. Johnson, A. J., Vangay, P., Al-Ghalith, G. A., Hillmann, B. M., Ward, T. L., Shields-Cutler,
R. R., Kim, A. D., Shmagel, A. K., Syed, A. N., Personalized Microbiome Class Students,
Walter, J., Menon, R., Koecher, K., & Knights, D. (2019). Daily Sampling Reveals
Personalized Diet-Microbiome Associations in Humans. *Cell host & microbe*, *25*(6), 789–
802.e5. https://doi.org/10.1016/j.chom.2019.05.005

1013

Figure 1: PRISMA Flow Diagram for Study Selection summarizing the systematic search and screening process for identifying eligible studies included in the meta-analysis.

2A

Plant   Animal

408   206   71
(59.6%)   (30.1%)   (10.4%)

Unique Plant Taxa

2B

Genus

■ Anaerococcus (1.1%)	■ Chujaibacter (1.3%)	■ KF-JG30-C25 (1.2%)	■ Phascolarctobacterium (1.4%)	■ Rokubacteriales (1.6%)
■ Candidatus_Solibacter (1.2%)	■ Gemmatimonas (3%)	■ LWQ8 (1.1%)	■ Prevotellaceae_UCG-003 (1.2%)	■ Sphingobacterium (1.4%)
■ Candidatus_Udaeobacter (1.9%)	■ JG30-KF-AS9 (1.3%)	■ MND1 (1.2%)	■ RB41 (3.2%)	■ Subgroup_2 (1.3%)
■ Cetobacterium (2.5%)	■ KD4-96 (1.8%)	■ Others (69.9%)	■ Rhodanobacter (1.1%)	■ Subgroup_7 (1.3%)

Unique Animal Taxa

2C

Genus

[Acetivibrio]_ethanolgignens_group (1.7%)	Candidatus_Competibacter (6.3%)	Defluviicoccus (1.4%)
[Clostridium]_methylpentosum_group (1.4%)	Candidatus_Stoquefichus (1.9%)	Gordonibacter (4.8%)
Aeromonas (1.2%)	Caulobacter (6.7%)	Lachnospiraceae_UCG-004 (6.5%)
Bhargavaea (3.6%)	Christensenella (1.9%)	Micropruina (2.7%)

Others (25.5%)	Solibacillus (3.9%)
PeM15 (2.7%)	Sphingomonas (10.1%)
Propioniciclava (5.1%)	Sphingopyxis (8.4%)
Rhodococcus (2.2%)	Tessaracoccus (1.9%)

2D

Genus

- Muribaculaceae
- Faecalibaculum
- Akkermansia
- Lactococcus
- Ligilactobacillus
- Dubosiella
- Bifidobacterium
- Alistipes
- Bacteroides
- Staphylococcus
- HT002
- Allobaculum
- Rikenellaceae_RC9_gut_group
- Desulfovibrio
- Coriobacteriaceae_UCG-002
- Ileibacterium
- Desulfovibrionaceae Family
- Corynebacterium
- Alloprevotella
- Other

Relative Abundance

Protein Source

Animal   Plant

Figure 2. Microbial composition and taxonomic uniqueness across dietary protein sources.
(A) Venn diagram showing the number and percentage of microbial genera unique to plant-fed and animal-fed mice, as well as genera shared between both groups. (B) Pie chart illustrating the proportional abundance of genera uniquely found in the plant-protein group. Low abundance taxa are grouped under "Others". (C) Pie chart showing the relative abundance of genera unique to the animal-protein group. (D) Stacked bar plot comparing the relative abundance of dominant genera across animal- and plant-protein diets. Taxa such as *Muribaculaceae*, *Faecalibaculum*, *Lactococcus*, and *Dubosiella* differ in prevalence between the two dietary groups.

3B

Relationship Between Sequencing Depth and Microbial Diversity

Spearman's rho = -0.115
P-value = 0.116

Shannon Diversity Index

Sequencing Depth (Total Reads)

3C

Figure 3: Microbial Diversity and Community Structure in Protein-Based Diets
(A) Linear mixed model comparing predicted versus observed Shannon diversity, with marginal and conditional R² values indicating variance explained by fixed effects alone (0.071) versus

fixed plus random effects (0.893), respectively. (B) Spearman correlation between sequencing depth and Shannon diversity. (C) Boxplots showing significantly higher microbial richness and evenness in plant-based protein diets. (D) Principal Coordinates Analysis (PCoA) based on Aitchison distance reveals clear separation of microbial communities by protein source. Ellipses represent 95% confidence intervals. (E) ANOSIM analysis confirms significant differences in microbial community structure between diet groups (R = 0.663, P < 0.001).

4D

a: g__Corynebacterium
b: g__Bacteroides
c: g__Muribaculaceae
d: g__Muribaculum
e: g__Alloprevotella
f: g__Parabacteroides
g: g__Tannerellaceae
h: g__Erysipelatoclostridium
i: g__Allobaculum
j: g__Atopostipes
k: g__HT002
l: g__Lactobacillus
m: g__Mycoplasma
n: g__Clostridium_sensu_stricto_1
o: g__Lachnospiraceae_UCG-001
p: g__Colidextribacter
q: g__Intestinimonas
r: g__Parasutterella
s: g__Escherichia-Shigella
t: o__Corynebacteriales
u: o__Bacteroidales
v: o__Campylobacterales
w: o__Erysipelotrichales
x: o__Mycoplasmatales
y: o__Peptostreptococcales-Tissierellales
z: o__Burkholderiales
A: f__Corynebacteriaceae
B: f__Bacteroidaceae
C: f__Muribaculaceae
D: f__Prevotellaceae
E: f__Tannerellaceae
F: f__Erysipelotrichaceae
G: f__Mycoplasmataceae
H: f__Peptostreptococcaceae
I: f__Sutterellaceae

Protein_source
● Animal
● Plant

-log10(pvalue)
○ 5
○ 10

Figure 4. Machine Learning performance and key microbial predictors of dietary protein source

(A) Receiver Operating Characteristic (ROC) curves showing percentage accuracy and area under curve (AUC) comparing three Random Forest (RF) models: unadjusted (Original), class-weighted (Weighted), and balanced via ROSE sampling. The weighted RF model, while slightly lower in accuracy, avoids overfitting and offers more realistic performance, with an AUC of 0.995 and accuracy of 88.2%. (B) Top 10 microbial genera contributing most to the weighted RF model's classification of animal vs plant protein diets, based on Mean Decrease in Accuracy (MDA). (C). LDA using LEfSe highlights taxa enriched in each diet (LDA score ≥ 3.0, p < 0.05) (D). Cladogram shows microbial taxa enriched in plant-based (blue) and animal-based (red) diets, arranged hierarchically from phylum to genus with node sizes reflecting statistical significance (-log10[p-value]).

.

5A

5B Correlation of Metabolic Pathways and Key Genera

Functional Interaction Network Between Gut Microbiota and Metabolic Pathways

Figure 5. Functional profiling of microbial communities by protein source using PICRUSt2 predictions. (A) Top 10 KEGG pathways significantly enriched in mice fed animal- or plant-based protein diets (|log₂ fold change| > 2, FDR < 0.05). Red bars indicate enrichment in the animal group; blue bars indicate enrichment in the plant group. (B) Heatmap showing Pearson correlation between microbial genera and selected predicted metabolic pathways involved in short-chain fatty acid (SCFA), branched-chain fatty acid (BCFA), phenol/p-cresol, ammonia, and sulfur metabolism. Functions are color-coded by metabolic category, and genera are annotated by phylum. (C) Network diagram of strong correlations (|r| ≥ 0.6) between microbial genera and selected predicted metabolic pathways. Node size reflects the number of connections (degree centrality), and edge thickness corresponds to the strength of correlation.

Table 1: Characteristics of studies included in the meta-analysis

Country, Year, First Author	Title	Bioproject accession	Protein source	Sample number	Age of mice (weeks)
Australia, 2022 Jian Tan	Dietary protein increases T-cell-independent sIgA production through changes in gut microbiota-derived extracellular vesicles	PRJEB39583	Casein	8	6
Japan, 2020 Hiroaki Masuoka	The influences of low protein diet on the intestinal microbiota of mice	PRJDB8898	Casein	30	9
China, 2022 Yantao Yin	Dietary oxidized beef protein alters gut microbiota and induces colonic inflammatory damage in C57BL/6 mice	PRJNA872483	Beef	24	5
China, 2022 Yifan Zhu	Reno-Protective Effect of Low Protein Diet Supplemented With a-Ketoacid Through Gut Microbiota and Fecal Metabolism in 5/6 Nephrectomized Mice	PRJNA810785	Low Protein diet	10	8
China, 2020 Yunting Xie	Processing Method Altered Mouse Intestinal Morphology and Microbial Composition by Affecting Digestion of Meat Proteins	PRJNA607338	Casein, Pork, Sausage, Soy protein	30	4
Canada, 2021 Béatrice S.-Y. Choi	Feeding diversified protein sources exacerbates hepatic insulin resistance via increased gut microbial branched-chain fatty acids and mTORC1 signaling in obese mice	PRJEB37442	Casein	12	10
China, 2020 Muhammad Umair Ijaz	Meat Protein in High-Fat Diet Induces Adipogenesis and Dyslipidemia by Altering Gut Microbiota and Endocannabinoid Dysregulation in the Adipose Tissue of Mice	PRJNA548036	Beef, Casein, Chicken, Pork	27	7
China, 2022 Wuling Zhong	High-protein diet prevents fat mass increase after dieting by counteracting Lactobacillus-enhanced lipid absorption	PRJNA868153	Casein	14	8
China, 2022 Ranran Zhang	Oral administration of branched-chain amino acids ameliorates high-fat diet-induced metabolic-associated fatty liver disease via gut microbiota-associated mechanisms	PRJNA852193	Branched-chain amino acids	12	6
Italy, 2020 Chiara Ruocco	Manipulation of Dietary Amino Acids Prevents and Reverses Obesity in Mice Through Multiple Mechanisms That Modulate Energy Homeostasis	PRJEB25686	Essential amino acids, Casein	20	8

Table 2: Core microbiota based on protein source at Genus level

Genera	Animal Protein (%)	Genera	Plant Protein (%)
Muribaculaceae	21.93	*Muribaculaceae*	35.64
Faecalibaculum	10.01	*Faecalibaculum*	6.17
Akkermansia	8.70	*Akkermansia*	6.02
Lactococcus	7.34	*Allobaculum*	4.79
Ligilactobacillus	4.24	*Dubosiella*	4.79
Staphylococcus	3.25	*Bacteroides*	3.02
Bifidobacterium	3.21	*Ligilactobacillus*	2.99
Rikenellaceae_RC9_gut_group	2.82	*Alloprevotella*	2.15
HT002	2.49	*Alistipes*	1.84
Alistipes	2.41	*Desulfovibrio*	1.57
Others	20.44	Others	26.27