Deployment of endocytic machinery to periactive zones of nerve terminals is independent of active zone assembly and evoked release

Javier Emperador-Melero [1,2] *, Steven J. Del Signore [3] *, Kevin M. De León González [3], Pascal S. Kaeser [1] ** #, and Avital A. Rodal [3] ** #

[1]Department of Neurobiology, Harvard Medical School, Boston, USA

[2]Department of Neuroscience and Institute for Translational Neuroscience, New York University Grossman School of Medicine, New York, NY 10016, US

[3]Department of Biology, Brandeis University, Waltham, USA

*Equal contribution

**Equal contribution

correspondence: kaeser@hms.harvard.edu and arodal@brandeis.edu

Abstract

In presynaptic nerve terminals, the endocytic apparatus rapidly restores synaptic vesicles after neurotransmitter release. Many endocytic proteins localize to the periactive zone, a loosely defined area adjacent to active zones. A prevailing model posits that recruitment of these endocytic proteins to the periactive zone is activity-dependent. We here show that periactive zone targeting of endocytic proteins is largely independent of active zone machinery and synaptic activity. At periactive zones of mouse hippocampal synapses and *Drosophila* neuromuscular junctions, pharmacological or genetic silencing resulted in unchanged or increased levels of endocytic proteins including Dynamin, Amphiphysin, Nervous Wreck, PIPK1γ and AP-180. Similarly, disruption of active zone assembly via genetic ablation of active zone scaffolds at each synapse did not impair the localization of endocytic proteins. Overall, our work indicates that endocytic proteins are constitutively deployed to the periactive zone and supports the existence of independent assembly pathways for active zones and periactive zones.

Introduction

Evoked release of neurotransmitters relies on the coordinated function of two complementary protein machineries, each comprising many proteins. First, the active zone defines vesicular release sites and clusters voltage-gated Ca^{2+} channels of the Ca_V2 family for rapid fusion of synaptic vesicles in response to action potentials (Emperador-Melero and Kaeser, 2020; Sudhof, 2012). Second, the endocytic apparatus restores vesicles after exocytosis for subsequent rounds of refilling and release. The endocytic machinery is enriched at the periactive zone, a region typically localized adjacent to the active zone (Haucke et al., 2011; Saheki and De Camilli, 2012; Watanabe and Boucrot, 2017). Despite the functional relationships between exocytosis and endocytosis and the spatial coupling of their respective protein machineries, the mechanisms that determine the recruitment of the endocytic apparatus to the periactive zone remain uncertain.

A prevailing model posits that deployment of endocytic machinery to the periactive zone occurs in response to synaptic activity. Multiple lines of evidence support this model. First, depolarization causes some endocytic proteins (including Dap160/Intersectin, Endophilin and Dynamin) to partially shift from the cytoplasm or from synaptic vesicle pools to the plasma membrane (Bai et al., 2010; Jiang et al., 2024; Koh et al., 2007; Winther et al., 2015, 2013). Second, membrane uptake increases rapidly in response to exocytosis triggered by synaptic activity (Haucke et al., 2011; Saheki and De Camilli, 2012; Watanabe and Boucrot, 2017), indicating that the endocytic protein machinery is responsive to activity and might be recruited to the membrane in consequence. Third, activity-dependent mechanisms at synapses might parallel protein deployment for Clathrin-mediated endocytosis in non-neuronal cells, where cytosolic endocytic machinery is transiently and acutely recruited to discrete sites (Cocucci et al., 2014; Kaksonen and Roux, 2018; Taylor et al., 2011). Fourth, the distributions of Clathrin, the Clathrin adaptor γ-Adaptin and one Dynamin isoform become diffuse after blocking action

potentials in neurons with impaired endocytosis (Ferguson et al., 2007; Milosevic et al., 2011; Raimondi et al., 2011).

An alternative model in which the endocytic apparatus is constitutively recruited to the periactive zone is also supported by previous work. For example, unstimulated synapses contain endocytic machinery at the periactive zone (Bloom et al., 2003; Del Signore et al., 2023; Estes et al., 1996; Gerth et al., 2017; Imoto et al., 2022; Roos and Kelly, 1999; Wahl et al., 2013; Winther et al., 2015, 2013), raising the possibility that periactive zone assembly relies on active zone proteins or other activity-independent mechanisms. Furthermore, ultrafast endocytosis requires pre-deployment of Dynamin and pre-assembly of a ring of F-actin surrounding the active zone (Imoto et al., 2022; Ogunmowo et al., 2023). In addition, endocytic machinery is required to prevent rapid synaptic depression, which occurs at a timescale that might exceed the speed of activity-dependent protein recruitment (Hua et al., 2011; Jäpel et al., 2020; Kawasaki et al., 2000). Moreover, the endocytic machinery is involved in functions that span beyond endocytosis and are more likely to depend on predeployment of machinery to the plasma membrane. These include the regulation of the exocytic fusion pore, the fusion of dense core vesicles, and the trafficking of adhesion molecules, growth factors and extracellular vesicle cargoes (Anantharam et al., 2011; Bailey et al., 1992; Blanchette et al., 2022; Fu and Huang, 2010; Moro et al., 2021; Rodal et al., 2008; Samasilp et al., 2012). Overall, it has remained uncertain whether endocytic machinery is predeployed or recruited on an as-needed basis by activity.

Here, we combine genetic and pharmacological manipulations at two model synapses to test whether the recruitment of endocytic machinery to the periactive zone depends on synaptic activity or active zone scaffolds. Our work establishes that endocytic machinery is efficiently targeted to synapses and deployed to the periactive zone when evoked synaptic activity is

chronically inhibited either pharmacologically or genetically. Furthermore, multiple active zone

organizers, including RIM, ELKS and its homologue Brp, Liprin-α, and Rab3 were not required

for recruitment and positioning of endocytic proteins to the periactive zone. Our findings support

that endocytic machinery is constitutively deployed to presynaptic nerve terminals and localizes

to periactive zones independent of active zone assembly and evoked synaptic vesicle release.

Results

Localization of endocytic proteins after silencing or depolarization of mouse hippocampal neurons

We generated primary neurons from mouse hippocampi and modulated their activity either by silencing them chronically or by triggering exocytosis acutely (Fig. 1a). We then assessed the distribution of endocytic proteins in these conditions compared to untreated neurons. To chronically inhibit activity, we simultaneously blocked action potentials with the Na^+ channel blocker tetrodotoxin (TTX) and Ca^{2+} entry with blockers of $Ca_V2.1$ (ω-agatoxin IVA; ω-Aga) and $Ca_V2.2$ (ω-conotoxin GVIA; ω-Cono) channels. We added this drug cocktail at 3 days *in vitro* (DIV) and resupplied it every three days, which results in continuous inhibition, as established before (Held et al., 2020). To acutely trigger exocytosis in neurons that were not treated with blockers, we depolarized neurons with superfusion of 50 mM KCl for 30 seconds.

We fixed the neurons that were either treated with blockers (for 12 days) or depolarized acutely and used antibody staining to label components of the synaptic endocytic machinery. This machinery contains multiple proteins that can be distributed into several categories including (1) BAR-domain containing proteins that sense and generate membrane invaginations, (2) phosphoinositide enzymes that regulate lipid metabolism, (3) components of Clathrin coats that sort recycled proteins, and (4) Dynamin proteins that operate in membrane fission. We labeled the BAR-domain containing protein Amphiphysin, the phosphoinositide kinase PIPK1γ, the Clathrin adaptor AP-180, and Dynamin-1 (Di Paolo et al., 2002; Koo et al., 2015; Raimondi et al., 2011; Wenk et al., 2001). We co-stained for the synaptic vesicle marker Synaptophysin, and an active zone marker (either the scaffold Bassoon or the vesicle priming protein Munc13-1, as we did before (Wong et al., 2018)). We detected signals generated by these antibodies using two complementary approaches as we described before (Emperador-Melero et al., 2024,

2021b): confocal microscopy was used to quantify protein levels in presynaptic boutons, and

stimulated emission depletion (STED) microscopy was employed to assess subsynaptic protein

distributions and levels at the presynaptic plasma membrane.

To estimate synaptic levels in confocal images, we quantified the average signal intensities of

the proteins of interest within regions of interest (ROIs) defined by Synaptophysin staining using

a thresholding and segmentation method established before (Emperador-Melero et al., 2024,

2021b; Liu et al., 2018). The average Amphiphysin, PIPK1γ and AP-180 signal intensities were

increased by up to 40% after chronic inhibition of activity, and those of Dynamin-1 were

unaffected (Fig. 1 – figure supplement 1a+b). Enhanced levels in this condition were also

present for the vesicle protein Synaptophysin and for Munc13-1, and may reflect a homeostatic

increase of presynaptic material in response to chronic silencing. Hence, blocking action

potentials and Ca^{2+} entry did not decrease the targeting of endocytic proteins to synapses.

Acute depolarization enhanced the signal intensity of AP-180 by ~30%, while Amphiphysin,

PIPK1γ and Dynamin-1 were not increased. The increase in AP-180 may reflect that some

endocytic proteins can be recruited to synapses in response to strong depolarization, matching

previous reports (Bolz et al., 2023; Mueller et al., 2004).

To assess subsynaptic protein localizations, we analyzed side-view and en-face synapses after

acquiring images with STED microscopy (Fig. 1 – figure supplement 2a-e). Side-view analyses

enable the assessment of protein distributions axially from the plasma membrane to the inside

of the bouton. We identified side-view synapses as those containing an elongated active zone

marker (Bassoon or Munc13) at the edge of a Synaptophysin cloud. We then extracted the

distribution of the protein of interest by positioning a line profile perpendicular to the active zone

marker, as we did before (Chin and Kaeser, 2024; Emperador-Melero et al., 2024, 2021b; Held

et al., 2020; Tan et al., 2022). At untreated synapses, the distributions of Amphiphysin, PIPK1γ,

AP-180 and Dynamin-1 were broader than those of active zone proteins (Fig. 1b-i), consistent with localization at the periactive zone and in the presynaptic cytosol (Ganguly et al., 2021; Gerth et al., 2017; Wenk et al., 2001). The average peak signal intensities of these proteins were within ~100 nm of the peak of the active zone marker (Fig. 1j). To estimate levels at the presynaptic plasma membrane, we measured the signal intensities within a 135 nm region centered around the peak of the active zone marker (Wong et al., 2018), which likely includes the active zone and the periactive zone because the two regions overlap in side-view. The signal intensities for Amphiphysin and PIPK1γ in this region were increased upon chronic silencing, matching the higher synaptic levels quantified in confocal images, and the intensities of AP-180 and Dynamin-1 were similar to the untreated condition (Fig. 1k). The levels of AP-180, but not of the other three proteins, increased with KCl depolarization (Fig. 1k), matching the confocal data (Fig. 1 – figure supplement 1a, b). We observed a similar increase in AP-180 signals using an alternate antibody (Fig. 1 – figure supplement 3). We did not detect changes in the axial distribution of any of these proteins (Fig. 1j). Overall, these observations suggest that the localization of endocytic proteins to the presynaptic plasma membrane does not require action potential firing and presynaptic Ca^{2+} entry, but instead indicate activity-independent recruitment.

Next, we analyzed en-face synapses to assess the lateral protein distribution near the plane of the plasma membrane (Fig. 1 – figure supplement 2f-i). We identified en-face synapses as those that did not have a bar-like appearance of the marker; instead, the area of the marker was surrounded by Synaptophysin staining (Emperador-Melero et al., 2024). At untreated synapses, Amphiphysin, AP-180 and Dynamin were organized into multiple clusters adjacent to the active zone. On average, these clusters were located 200 to 300 nm lateral from the center of the active zone marker (Fig. 1l-q). The average distribution of PIPK1γ appeared closer to the active zone, possibly reflecting its roles in the synthesis of phosphatidylinositol 4,5-bisphosphate

(PI(4,5)P$_2$), a phospholipid important for both exo- and endocytosis (Bai et al., 2003; Bolz et al.,

2023; Di Paolo et al., 2004; Wenk et al., 2001). Neither chronic silencing nor strong

depolarization changed the lateral distribution of these proteins. Instead, they remained

distributed in clusters surrounding the active zone (Fig. 1l-q). There was a slight increase in the

number of PIPK1γ objects upon depolarization (Fig. 1p). Upon chronic silencing, the integrated

intensity of endocytic protein clusters (defined as the product of their size and average intensity)

increased or showed a positive trend for all proteins (Fig. 1 – figure supplement 4). Integrated

intensity also increased for AP-180 upon depolarization, matching with the higher levels

measured at side-view synapses and at confocal resolution (Fig. 1 and Fig. 1 – figure

supplement 4).

Together, these experiments indicate that while endocytic proteins are not impervious to

synaptic activity, they are efficiently deployed to presynaptic terminals and to the periactive zone

after pharmacological silencing of firing activity and Ca^{2+} entry.

Localization of endocytic proteins after chronic silencing or acute stimulation of

***Drosophila* NMJs**

We next asked whether recruitment of endocytic proteins to the presynaptic plasma membrane

follows a similar pattern at the *Drosophila* neuromuscular junction (NMJ), a widely used model

for studying synaptic mechanisms (Harris and Littleton, 2015). These synapses consist of 40 to

100 large boutons each containing tens of active zones surrounded by a honeycomb or mesh of

well-defined periactive zones (Del Signore et al., 2023, 2021; Winther et al., 2015). For most

experiments, we analyzed Type 1b terminals on muscles 4 or muscle 6/7, which have been

extensively characterized developmentally and electrophysiologically. For silencing

experiments, we analyzed Type 1b terminals at muscle 1 where a single-neuron GAL4 driver is

available (Jenett et al., 2012), since silencing NMJs broadly is embryonic lethal (Sweeney et al.,

1995). We specifically analyzed the endocytic proteins Dynamin and Nervous Wreck, the homologue of the mammalian F-BAR domain containing protein FCHSD2, because they have distinct molecular activities, functional contributions to endocytosis, and spatial patterning within the periactive zone (Coyle et al., 2004; Del Signore et al., 2023; Kawasaki et al., 2000; Salkoff and Kelly, 1978). These markers define a periactive zone in this model system as a discrete region surrounding active zones approximately 300 nm from the active zone center. As an active zone marker, we selected Bruchpilot (Brp), a homologue of ELKS (Kittel et al., 2006; Wagh et al., 2006).

To analyze the organization of active zone and periactive zone machineries, we made a series of measurements. First, we quantified the number of Brp objects per synaptic area, as well as the total intensity of Brp per object. Second, to determine recruitment or retention of endocytic machinery or active zones at the terminal, we measured the mean intensity within the entire 3D volume of the terminal. Third, to quantify deployment to the periactive zone, we used an established workflow to segment NMJs into single synaptic units composed of a periactive zone surrounding a center region containing ~0 to 3 Brp objects (Fig. 1 – figure supplement 2j-n) (Del Signore et al., 2023). Except where noted, we analyzed the ~60% of periactive zones that were associated with one Brp object. Each unit is divided into 'mesh' and 'core' regions, where the periactive zone mesh is a ~175 nm wide band localized on average ~330 nm from the center, and the 'core' region is interior to this periactive zone mesh (Del Signore et al., 2023). We estimated the levels of proteins in the periactive zone mesh as their mean intensity levels within this region and their distribution as the log ratio of the average intensity within the mesh over the core, with positive ratios indicating mesh enrichment and negative ratios indicating core enrichment (Fig. 1 – figure supplement 2j-n).

To assess the influence of activity, we silenced type 1b motor neurons on muscle 1 by

expressing Tetanus Neurotoxin (TeNT), which cleaves the SNARE protein Synaptobrevin (n-Syb) and disrupts evoked exocytosis (Sweeney et al., 1995). Control motor neurons expressed the GAL4 driver alone (Fig. 2a). We then used antibody staining followed by Airyscan confocal microscopy to visualize endocytic and active zone proteins, and analyzed the levels of these proteins at NMJs and their localization to the periactive zone. Chronic TeNT expression resulted in fewer Brp objects per μm^2 of NMJ and an increase in their integrated intensity, indicating a reorganization of active zone proteins (Fig. 2b-d), as recently observed (Akbergenova et al., 2025). However, TeNT expression did not alter the total levels of Nervous Wreck or Dynamin or their abundance in the periactive zone, and caused only a small redistribution of Dynamin (Fig. 2e-g).

We next tested the prediction that enhanced activity might increase the localization of endocytic proteins to the periactive zone mesh at the *Drosophila* NMJ. To do so, we electrically stimulated Type 1b motor neurons on muscles 4 and 6/7 with a 3-minute 40 Hz train (Fig. 2a), a stimulus which induces endocytosis and causes redistribution of PI(4,5)P2 at the presynaptic membrane (Li et al., 2020). If deployment were activity dependent, we would expect increased periactive zone targeting of endocytic proteins and a concomitant change increase in their polarization. However, stimulation did not change the mean levels of Dynamin, Nervous Wreck or Brp at boutons, though we noted a small decrease in the density of BRP objects. Similarly, the levels and polarization of Dynamin and Nervous Wreck at the periactive zone mesh also remained unchanged (Fig. 2h-m). Altogether, we conclude that evoked neurotransmission is not required to target Dynamin and Nervous Wreck to NMJs or to localize them within these boutons, nor does enhanced activity in the tested paradigms boost their periactive zone localization.

Efficient deployment of endocytic proteins after Ca$_V$2 ablation in mouse hippocampal neurons

Action potential-triggered fusion of synaptic vesicles depends on Ca^{2+} influx via voltage-gated Ca^{2+} channels of the Ca_V2 family (Cao et al., 2004; Cunningham et al., 2022; Held et al., 2020). As a complementary approach to study the role of synaptic activity in the recruitment of endocytic machinery, we assessed synapses lacking Ca_V2s (Fig. 3a). We cultured neurons from previously generated triple conditional Ca_V2 knockout mice ($Ca_V2.1$, $Ca_V2.2$ and $Ca_V2.3$) and used lentiviral transduction to express Cre recombinase to generate Ca_V2 triple knockout (cTKO[Cav2]) neurons This disrupts action potential-evoked exocytosis (Chin and Kaeser, 2024; Held et al., 2020). We expressed a recombination-deficient Cre enzyme that is truncated to generate control (control[Cav2]) neurons (Held et al., 2020). In confocal microscopic images, the average synaptic signal intensities of Amphiphysin, PIPK1γ, AP-180 and Dynamin-1 were intact at cTKO[Cav2] synapses (Fig. 3 – figure supplement 1), confirming efficient targeting of these proteins. No major changes in their subsynaptic localization were observed either. When assessed in STED microscopic images, the axial protein distributions of Amphiphysin, PIPK1γ, AP-180 and Dynamin-1 were unchanged at cTKO[Cav2] synapses, and the peak fluorescence intensities within the periactive zone region of cTKO[Cav2] synapses were within ~10% of the intensities measured at control[Cav2] synapses (Fig. 3b-k and Fig. 3 – figure supplement 2). At en-face synapses, the number and intensity of clusters positioned at 100 to 300 nm from the active zone center were unchanged (Fig. 3l-q and Fig. 3 – figure supplement 3).

We conclude that ablating Ca_V2s to disrupt action potential-induced synaptic vesicle exocytosis does not impair the localization of endocytic proteins, matching the outcomes of the pharmacological inhibition experiments.

Deployment of endocytic proteins after active zone disruption in mouse hippocampal neurons

The active zone generates sites for synaptic vesicle fusion and clusters Ca_V2 channels at the

presynaptic plasma membrane (Emperador-Melero and Kaeser, 2020; Sudhof, 2012). It is a

multiprotein machinery formed by RIM, RIM-BP, Liprin-α, Munc13, ELKS, Piccolo/Bassoon and

other proteins that is subdivided into distinct sub-machineries (Acuna et al., 2016; Aravamudan

et al., 1999; Biederer et al., 2017; Böhme et al., 2016; Emperador-Melero et al., 2024, 2021b;

Emperador-Melero and Kaeser, 2020; Graf et al., 2012; Kaufmann et al., 2002; Kittel et al.,

2006; Liu et al., 2011; Marcó de la Cruz et al., 2024; McDonald et al., 2020; Sudhof, 2012;

Wang et al., 2016; Zhen and Jin, 1999). Endocytic machineries are localized adjacent to active

zones, and the functions of these two machineries are coordinated. Hence, their assembly may

be linked with instructive roles of the active zone. To test this hypothesis, we used quadruple

conditional ablation of RIM1, RIM2, ELKS1 and ELKS2 at hippocampal synapses. At these

synapses, active zone assembly is strongly impaired with disrupted localization of Munc13,

Piccolo/Bassoon, RIM-BP and Ca_V2 (Emperador-Melero et al., 2024; Tan et al., 2022; Wang et

al., 2016; Wong et al., 2018). We cultured neurons from these mice and transduced them with

lentiviruses expressing Cre or inactive Cre to produce RIM+ELKS quadruple knockout

(cQKO[R+E]) or control neurons (control[R+E]), respectively (Fig. 4a). We antibody-stained these

cultures as in Figs. 1 and 3, but used the postsynaptic marker PSD-95 instead of an active zone

marker because the active zone is disrupted at cQKO[R+E] synapses. At side-view synapses,

PSD-95 is localized apposed to active zones at a distance of ~75 nm (Emperador-Melero et al.,

2024, 2021b; Held et al., 2020; Tan et al., 2022; Tang et al., 2016; Wong et al., 2018).

Altogether, endocytic machinery remained correctly localized after active zone disruption. In

confocal images (Fig. 4 – figure supplement 1), average intensities of fluorescence signals for

antibody staining for Amphiphysin, PIPK1γ, AP-180 and Dynamin-1 at presynapses were either

similar or higher in cQKO[R+E] neurons. At side-view synapses in STED microscopy (Fig. 4b-l and

Fig. 4 – figure supplement 2), the peak intensities were at a similar distance from PSD-95, and

the signals within the region corresponding to the periactive zone were either higher or

unchanged. Similarly, endocytic proteins at en-face synapses (Fig. 4 m-r and Fig. 4 – figure

307 supplement 3) from cQKO^{R+E} neurons were organized into clusters with comparable localization

308 and similar or greater integrated density. We conclude that the localization of endocytic proteins

309 to the periactive zone is not impaired after disrupting active zone assembly through knockout of

310 RIM and ELKS.

Deployment of endocytic proteins at *Drosophila* NMJs after disrupting active zone

assembly

The active zone of the *Drosophila* NMJ has a T-bar, a prominent electron-dense invagination

consisting mainly of Brp (Fouquet et al., 2009; Kittel et al., 2006; Wagh et al., 2006). Brp is

required for the normal clustering of Ca$_V$2 Ca^{2+} channels at the active zone, and its ablation

severely decreases evoked release (Fouquet et al., 2009; Kittel et al., 2006; Wagh et al., 2006).

To determine whether these defects perturb periactive zone protein recruitment, we analyzed

levels and localization of endocytic proteins in *brp* mutants (Fig. 5a). After genetic ablation of

brp, the levels of Dynamin and Nervous Wreck at NMJs were not decreased and these proteins

remained concentrated in the periactive zone mesh (Fig. 5b-e). These data indicate that Brp

and active zone T-bars are dispensable to localize these endocytic proteins to the periactive

zone at the *Drosophila* NMJ and corroborate the finding (Fig. 1b-g and 2b-g) that periactive

zone properties do not depend on evoked release.

Deployment of endocytic machinery after disruption of upstream active zone assembly

mechanisms

328 The observation that active zone disassembly does not reduce levels or disrupt localization of

329 endocytic proteins at periactive zones (Figs. 4 and 5) suggests that active zones and endocytic

330 machineries are organized independently of one another. Alternatively, shared upstream

331 assembly pathways might help co-organization of the active zone and the periactive zone.

Liprin-α is an upstream active zone organizer that maintains priming machineries and recruits additional presynaptic material (Böhme et al., 2016; Emperador-Melero et al., 2024, 2021b; Kaufmann et al., 2002; Marcó de la Cruz et al., 2024; McDonald et al., 2020; Wong et al., 2018; Zhen and Jin, 1999). We assessed the localization of endocytic proteins in Liprin-α mutants to assess whether these proteins might co-organize active zone and periactive zone assembly. In vertebrates, four Liprin-α proteins are expressed from four different genes (Zurner and Schoch, 2009). We previously generated quadruple Liprin-α mutant mice containing conditional Liprin-α1, -α2 and -α4 knockout alleles and constitutive Liprin-α3 knockout alleles (Emperador-Melero et al., 2024, 2021b; Wong et al., 2018). We cultured hippocampal neurons from these mice and transduced them with Cre lentivirus to produce quadruple knockout (cQKO^{L1-L4}) neurons, or a combination of two viruses to express Liprin-α3 and a truncated, inactive version of Cre, to generate control (control^{L1-L4}) neurons, as described before (Emperador-Melero et al., 2024). Synapses of cQKO^{L1-L4} neurons have altered active zone properties, with decreased levels of RIM and Munc13-1, accompanied by an impairment in the readily releasable pool of synaptic vesicles (Emperador-Melero et al., 2024). Endocytic proteins were overall efficiently deployed to presynaptic terminals and correctly localized to the periactive zone in cQKO^{L1-L4} neurons (Fig. 6 and Fig. 6 – figure supplements 1 and 2). There were only small decreases in signal intensities generated by Amphiphysin and AP-180 antibodies, a small increase in PIPK1y, and a modest shift in the lateral position of Amphiphysin. These results indicate that endocytic proteins are overall efficiently localized in the absence of Liprin-α proteins in mouse hippocampal neurons.

To disrupt *liprin-α* in *Drosophila*, we used a previously characterized heteroallelic mutant that results in loss-of-function of this protein (*liprin-α*^{EPexR60}/*liprin-α*^{F3ex15}) (Fouquet et al., 2009; Kaufmann et al., 2002; Owald et al., 2010) (Fig. 7a). Loss of *liprin-α* function reduced the overall levels of Brp at NMJs, and resulted in fewer Brp puncta with a greater integrated intensity (Fig. 7b-e) as described previously (Fouquet et al., 2009; Kaufmann et al., 2002; Owald et al., 2010)).

In contrast, the levels of Dynamin and Nervous Wreck were similar between controls and *liprin-α*

mutants, as was their polarization between periactive zone mesh and active zone, though we

did detect a small-magnitude decrease in Dynamin polarization (Fig. 7e-g). We conclude that

Liprin-α does not function as a major organizer of endocytic machinery at *Drosophila* NMJs, a

finding that further highlights the independence between active zone and periactive zone

assembly.

As an alternative strategy to address the independence between these two machineries at the

fly NMJ, we assessed the role of the synaptic vesicle-associated protein Rab3 (Fig. 8a) using an

established loss-of-function mutant (*rab3^{rup}*) (Graf et al., 2009; Peled and Isacoff, 2011). *rab3*

mutants had fewer but larger Brp objects (Fig. 8b-d) (Graf et al., 2009; Peled and Isacoff, 2011),

though by a mechanism distinct from TeNT expression (Akbergenova et al., 2025). Neither the

levels nor the distribution of Dynamin or Nervous Wreck were affected in *rab3* mutants (Fig. 8e-

g), showing that their periactive zone targeting is independent of *rab3*. Notably, the localization

of these endocytic proteins was similar between Brp-positive and Brp-negative periactive zones

in both controls and *rab3* mutants, with only modest shifts in the polarization of Dynamin in

controls and of Nervous Wreck in *rab3* mutants (Fig. 8h-k). Given the reduction in the number of

Brp objects, the percentage of periactive zones without detectable Brp objects in *rab3* mutants

was greater (Fig. 8l-n). Importantly, these Brp-absent periactive zones remained apposed to the

postsynaptic marker Pak (Fig. 8n), arguing that Brp-negative periactive zones are not simply

regions of membrane between adjacent periactive zones. We conclude that the recruitment of

endocytic machinery does not require *rab3* and that periactive zones can be formed without an

active zone.

383   Together, these data establish that the pathways organizing endocytic machinery are different

384   from those organizing active zones.

Discussion

We tested the hypotheses that synaptic activity or active zone scaffolds recruit and organize endocytic machinery at the periactive zone. In mouse hippocampal neurons, neither block of action potentials and presynaptic Ca^{2+} entry nor ablation of Ca_V2s disrupted the periactive zone localization of the endocytic proteins Amphiphysin, Dynamin-1, PIPK1γ or AP-180. Similarly, at the *Drosophila* NMJ, inhibiting evoked exocytosis did not change the levels or the subsynaptic distribution of Dynamin or Nervous Wreck. These endocytic proteins also remained clustered at the periactive zone in both types of synapses when active zone assembly was disrupted via ablation of the scaffolds RIM and ELKS or Brp, or when the upstream organizers Liprin-α or Rab3 were removed. Our data collectively argue that a periactive zone protein network, consisting of a high concentration of membrane-localized endocytic proteins, forms and is maintained independent of synaptic activity and active zone protein machinery.

Roles for activity in endocytic assemblies

Prior experiments tested the sufficiency of depolarization to induce recruitment of endocytic machinery to the periactive zone, but not whether it is necessary (Koh et al., 2007; Winther et al., 2015, 2013). We show that a significant fraction of endocytic proteins is already deployed to the membrane prior to stimulation. Neither continuous pharmacological silencing of neurons nor ablating Ca_V2 channels resulted in decreased levels of endocytic proteins at the periactive zone. Instead, the levels of some of these proteins were increased in these conditions, the opposite of the effect expected for activity-dependent recruitment. Our data do not exclude the possibility that a subset of endocytic proteins is acutely recruited to the periactive zone, for example Endophilin (Bai et al., 2010). In line with this, acute depolarization increased the clustering of AP-180 at periactive zones (Fig. 1), resembling previously reported increases in Clathrin using a similar experimental paradigm (Mueller et al., 2004). Importantly, Dynamin-1, Amphiphysin and

PIPK1γ did not increase upon depolarization with KCl. It is possible that, like the active zone

(Emperador-Melero et al., 2024), the endocytic machinery is formed by distinct submachineries

(Kaksonen and Roux, 2018), and that they respond to sustained stimulation differently.

Pathways for coordinating active zone and periactive zone assembly

Despite the functional coupling of exocytosis and endocytosis in nerve terminals, it has

remained unknown whether active zone proteins mediate the formation of the periactive zone.

We find that ablating active zone scaffolds, upstream organizers, or Ca^{2+} channels did not

disrupt the formation or structure of the periactive zone. What might be the pathways to

coordinate the assembly of these two adjacent machineries? One possibility is that the co-

organization of these machineries is instructed by a network of interconnected cell adhesion

proteins. LAR-RPTPs and Neurexins are localized at the active zone and organize this

machinery (Emperador-Melero et al., 2024, 2021a; Sclip and Südhof, 2023, 2020; Trotter et al.,

2019). Cell adhesion proteins localized at the edge of a synapse, such as SynCAM or Fasciclin-

II (Jiao et al., 2010; Perez de Arce et al., 2015), may play a similar role for endocytic machinery.

Presynaptic cell-adhesion complexes, for example LAR-RPTPs and Neurexins, might be

interlinked for the coordination of their intracellular interactors (Thivaios et al., 2024).

Presynaptic protein complexes might also be organized via transcellular interactions, a model

that is supported by the observation that periactive zones lacking Brp remain aligned with the

postsynaptic marker Pak, pointing to trans-synaptic mechanisms for periactive zone

organization.

Two additional contributing factors are the cytoskeleton and lipids. Multiple endocytic proteins

bind to and function in the nucleation and polymerization of actin filaments (Del Signore et al.,

2021; Saheki and De Camilli, 2012; Stanishneva-Konovalova et al., 2016). Interactions between

active zone proteins, such as Piccolo/Bassoon or Liprin-α, and actin regulators have also been

described (Brenig et al., 2015; Terry-Lorenzo et al., 2016) and these interactions may be

important for the early assembly of presynaptic compartments (Chia et al., 2012). Similarly,

$PI(4,5)P_2$ binds to RIM and this interaction is critical for synaptic strength (de Jong et al., 2018),

and the synthesis of this lipid is important for endocytosis (Di Paolo et al., 2004; Wenk et al.,

2001). Hence, actin and/or lipids may act as hubs around which active zone and endocytic

machineries are organized.

Finally, interactions between endocytic proteins may further contribute to the anchoring of this

apparatus. Most of these interactions are weak and transient, which may account for the high

degree of turnover or mobility of these proteins at synapses (Reshetniak et al., 2020). However,

their high concentration and their numerous interactions may suffice to maintain a stable

periactive zone structure (Saheki and De Camilli, 2012; Wilhelm et al., 2014). In support of this

point, perturbing interactions between Dynamin-1 and Endophilin-A1 increases the distance

between these proteins (Imoto et al., 2024), suggesting their binding has a scaffolding function.

Ultimately, it is plausible that several or all of these pathways organize the endocytic apparatus

in parallel. Redundancy is a guiding principle in the assembly of the active zone (Acuna et al.,

2016; Wang et al., 2016) and for synaptic cell adhesion (Sclip and Südhof, 2023), and it may

similarly guide endocytic assembly.

Importance of the constitutively deployed endocytic apparatus for synapse function

The constitutive deployment of endocytic machinery at periactive zones likely enables multiple

synaptic adaptations and functions of the endocytic machinery. First, ultrafast endocytosis,

which can occur within less than one second following stimulation (Watanabe et al., 2014),

depends on the pre-deployment of Dynamin-1 and assembly of a periactive zone-associated

ring of F-actin to facilitate membrane compression upon exocytosis (Imoto et al., 2022;

Ogunmowo et al., 2023). Pre-deployment may also facilitate vesicle recycling by limiting

diffusion of synaptic vesicle cargoes (Gimber et al., 2015), and may be important to prevent

synaptic depression (Hua et al., 2011; Kawasaki et al., 2000; Moro et al., 2021). Second,

proteins involved in endocytosis may contribute to additional functions, such as fusion of dense

core vesicles, regulation of the fusion pore, and trafficking of receptors, extracellular vesicles

and cell adhesion proteins (Anantharam et al., 2011; Bailey et al., 1992; Blanchette et al., 2022;

Fu and Huang, 2010; Moro et al., 2021; Rodal et al., 2008). Thus, constitutive deployment of the

endocytic machinery is likely an essential adaptation to meet the demands for fast, scalable,

and robust membrane internalization during synaptic vesicle recycling and to facilitate its

diverse functions at the synapse.

Given the constitutive deployment of endocytic machinery, why is it not constitutively active and

continuously retrieving presynaptic plasma membrane? Plausible mechanisms may include one

or more of the following factors, which might be relieved in response to Ca^{2+} entry or synaptic

vesicle fusion. First, the endocytic machinery may be maintained in an inactivated state via

autoinhibition of some of its components, for example Endophilin or Nervous Wreck (Del

Signore et al., 2021; Rao et al., 2010; Stanishneva-Konovalova et al., 2016). Second, endocytic

proteins deployed to the periactive zone might be segregated into distinct periactive zone

protein pools and thus not be able to functionally interact. In support of this model, we observed

that the clustering pattern of PIPK1γ in silenced hippocampal synapses was different from that

of Dynamin-1, Amphiphysin and AP-180 (Fig. 1). Furthermore, our previous work revealed low

colocalization between Dynamin, Nervous Wreck, DAP160/Intersectin and Clathrin within

periactive zones at the *Drosophila* NMJs (Del Signore et al., 2023). Third, one or multiple

essential components of the endocytic machinery might not be predeployed to the membrane

and may be delivered by an alternative mechanism (Bai et al., 2010). This is supported by the

observation that synaptic vesicle proteins, such as Synaptotagmin-1, have a role in endocytosis

(Bolz et al., 2023; Haucke and De Camilli, 1999; Jorgensen et al., 1995; Poskanzer et al., 2006;

Zhang et al., 1994). Finally, the endocytic process may rely on a decrease in membrane tension (Ogunmowo et al., 2023; Watanabe et al., 2013) and thus require fusion for its initiation. Given the multiple mechanisms by which periactive zone proteins likely contribute to synaptic vesicle endocytosis, and considering their potential synaptic vesicle-independent functions, it is possible that constitutive deployment is a necessary synaptic adaptation of the conserved endocytic machinery. Testing whether this is the case will require identifying the mechanisms that are essential for constitutive deployment of the endocytic machinery to the periactive zone.

Acknowledgments

We thank C. Qiao, V. Charles, G. Handy, and M. Quiñones-Frías for technical support, and members of the Kaeser and Rodal laboratories for insightful discussions. We thank P. De Camilli for anti-Dynamin-1, anti-PIPK1γ and anti-AP-180 antibodies, D. Dickman for anti-Dynamin antibodies, N. Harden for anti-Pak antibodies, A.M.J.M. van den Maagdenberg for Ca$_V$2.1 floxed mice, T. Schneider for Ca$_V$2.3 floxed mice, and Troy Littleton for *brp* mutant flies. This work was supported by grants from the NIH (R01MH113349 and R01NS083898 to PSK, R01NS116375 to AAR, T32007292 to KMDLG, and K99NS129959 to JE-M), from the NSF (Brandeis Materials Research and Engineering Center NSF-DMR 2011846), and from Harvard Medical School (Harvard/MITJoint Research Grant in Basic Neuroscience to PSK and AAR). JE-M was supported by an Alice and Joseph E. Brooks postdoctoral fellowship from Harvard Medical School. We used fly stocks from the Bloomington Drosophila Stock Center (NIH P40OD018537) and antibodies from the Developmental Studies Hybridoma Bank, generated by the NICHD of the NIH. We acknowledge the Harvard Neurobiology Imaging Facility and Brandeis University Light Microscopy Core Facility Imaging Facility (RRID:SCR_025892).

Author contributions

Conceptualization: JE-M, SJD, PSK and AAR; Methodology: JE-M, SJD; Investigation: JE-M, SJD, and KMDLG; Formal Analysis: JE-M, SJD and KMDLG; Writing-Original Draft: JE-M, SJD, PSK, AAR; Writing - Review and Editing: JE-M, SJD, KMDLG, PSK, AAR; Supervision: PSK, AAR; Funding Acquisition: JE-M, PSK, AAR. Experiments in mouse hippocampal neurons were conducted at Harvard Medical School and experiments in *Drosophila* were conducted at Brandeis University.

Declaration of interests

The authors declare no conflicts of interest.

Materials availability

Plasmids generated for this study will be shared upon request. Mouse lines and previously

generated plasmids will be shared upon request within the limits of the respective material

transfer agreements. Requests for resources and reagents should be directed to the

corresponding authors.

Data availability

Data points generated for this study are included in the figures. Additional data are available

from the corresponding authors upon request.

 MATERIALS AND METHODS

 Key resource table

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Genetic reagent (*Mus musculus*)	*Cacna1a* conditional knockout	(Todorov et al., 2006)		
Genetic reagent (*Mus musculus*)	*Cacna1b* conditional knockout	(Held et al., 2020)	IMSR_KOMP: CSD34514-1a-Wtsi	
Genetic reagent (*Mus musculus*)	*Cacna1e* conditional knockout	(Pereverzev et al., 2002)		
Genetic reagent (*Mus musculus*)	*Rims1* conditional knockout	(Kaeser et al., 2008)	IMSR_JAX:015832	
Genetic reagent (*Mus musculus*)	*Rims2* conditional knockout	(Kaeser et al., 2011)	IMSR_JAX:015833	
Genetic reagent (*Mus musculus*)	*Erc1* conditional knockout	(Liu et al., 2014)	IMSR_JAX:015830	
Genetic reagent (*Mus musculus*)	*Erc2* conditional knockout	(Kaeser et al., 2009)	IMSR_JAX:015831	
Genetic reagent (*Mus musculus*)	*Ppfia1* conditional knockout	(Emperador-Melero et al., 2024)	IMSR_EUMMCR: 25506	
Genetic reagent (*Mus musculus*)	*Ppfia2* conditional knockout	(Emperador-Melero et al., 2021b)	IMSR_HAR:6799	
Genetic reagent (*Mus musculus*)	*Ppfia3* constitutive knockout	(Wong et al., 2018)		
Genetic reagent (*Mus musculus*)	*Ppfia4* conditional knockout	(Emperador-Melero et al., 2024)	IMSR_EUMMCR: 3103	
Cell line (*Homo sapiens*)	HEK 293T cells	ATCC	Cat# CRL- 3216; RRID:CVCL_0063	
Plasmid	pFSW HA-Liprin-α3	(Wong et al., 2018)	p526	
Plasmid	pFSW	(Nyitrai et al., 2020)	p008	
Plasmid	pFSW GFP Cre	(Kaeser et al., 2011)	p009	
Plasmid	pFSW GFP inactive Cre	(Kaeser et al., 2011)	p010	
Antibody	Mouse anti-Dynamin-1	(Milosevic et al., 2011)		1:50 (IF, immunofluorescence)
Antibody	rabbit anti-Amphiphysin	(Di Paolo et al., 2002)		1:200 (IF)
Antibody	rabbit anti-PIPK1γ	(Wenk et al., 2001)		1:500 (IF)
Antibody	mouse anti-AP180	(Koo et al., 2015)		1:500 (IF)
Antibody	rabbit anti-AP180	SySy	AB_887691	1:500 (IF)
Antibody	rabbit anti-	SySy	AB_887733	1:500 (IF)

	Munc13-1			
Antibody	mouse anti-Synaptophysin	SySy	AB_887824	1:500 (IF)
Antibody	mouse anti-Bassoon	Enzo	AB_11181058	1:500 (IF)
Antibody	mouse anti-PSD-95	Neuromab	AB_10698024	1:500 (IF)
Antibody	guinea pig anti-PSD-95	SySy	AB_ 2232546	1:500 (IF)
Antibody	mouse anti-Gephyrin	SySy	AB_ 2232546	1:500 (IF)
Antibody	rabbit anti-Synapsin-1	SySy	AB_2200097	1:500 (IF)
Antibody	rabbit anti-Nervous Wreck 970	(Coyle et al., 2004)	RRID:AB_2567353	1:1000 (IF)
Antibody	guinea pig anti-Dynamin	other		1:1000 (IF)
Antibody	mouse anti-Brp nc82	Developmental Studies Hybridoma Bank (DSHB)	RRID:AB_2314866	1:1000 (IF)
Antibody	rabbit anti-Pak	(Harden et al., 1996)		1:1000 (IF)
Pharmacological agent	ω-Agatoxin IVA	Alomone labs	Cat#: STA-500	200 nM
Pharmacological agent	ω-Conotoxin GVIA	Alomone labs	Cat#: C-300	250 nM
Pharmacological agent	Tetrodotoxin	Tocris Bioscience	Cat # 1078	1 µM
Genetic reagent (*Drosophila melanogaster*)	*GMR94G06-GAL4*	Bloomington *Drosophila* Stock Center (BDSC); (Jenett et al., 2012)	RRID:BDSC_40701	
Genetic reagent (*Drosophila melanogaster*)	*UAS-TeTxLC; UAS-TeNT*	BDSC	RRID:BDSC_28838	
Genetic reagent (*Drosophila melanogaster*)	*brpDf/CyOGFP*	(Akbergenova et al., 2018)		
Genetic reagent (*Drosophila melanogaster*)	*brp69*	(Kittel et al., 2006)		
Genetic reagent (*Drosophila melanogaster*)	*rab3rup*	(Graf et al., 2009)	RRID:BDSC_78045	
Genetic reagent (*Drosophila melanogaster*)	*liprinR60*	BDSC; (Kaufmann et al., 2002)	RRID:BDSC_8561	
Genetic reagent (*Drosophila melanogaster*)	*liprinF3ex15*	BDSC; (Kaufmann et al., 2002)	RRID:BDSC_8563	

Mouse lines

The triple mutant mice for ablating Ca$_V$2.1 (targeting *Cacna1a* (Todorov et al., 2006)), Ca$_V$2.2

(targeting *Cacana1b* (Held et al., 2020); RRID: IMSR_KOMP: CSD34514-1a-Wtsi) and Ca$_V$2.3

(targeting *Cacana1e* (Pereverzev et al., 2002)) have been previously described (Chin and

Kaeser, 2024; Held et al., 2020). Quadruple mutant mice for ablating RIM1 (targeting *Rims1* to

remove RIM1α and RIM1β (Kaeser et al., 2008); RRID: IMSR_JAX:015832), RIM2 (targeting

Rims2 to remove RIM2α, RIM2β and RIM2γ (Kaeser et al., 2011); RRID: IMSR_JAX:015833),

ELKS1α (targeting *Erc1* to remove ELKS1αA and ELKS1αB (Liu et al., 2014); RRID:

IMSR_JAX:015830) and ELKS2α (targeting *Erc2* to remove ELKS2αA and ELKS2αB (Kaeser et

al., 2009); RRID: IMSR_JAX:015831) were previously described (Emperador-Melero et al.,

2024; Tan et al., 2022; Wang et al., 2016)**.** Quadruple mutant mice for ablating Liprin-α1

(targeting *Ppfia1* (Emperador-Melero et al., 2024); RRID: IMSR_EUMMCR:25506), Liprin-α2

(targeting Ppfia2 (Emperador-Melero et al., 2021b); RRID: IMSR_HAR:6799), Liprin-α3

(targeting *Ppfia3* (Wong et al., 2018)) and Liprin-α4 (targeting *Ppfia4* (Emperador-Melero et al.,

2024); RRID: IMSR_EUMMCR:3103) were previously described (Emperador-Melero et al.,

2024). Mouse lines were maintained as homozygotes for the respective mutations, and

offsprings were weaned at 21 to 28 days. Mice were kept separated by sex or housed as

breeding pairs in a room with a regular dark light cycle and set to 22 °C (range 20 °C to 24 °C)

and 50% humidity (range 35% to 70%). Experiments were approved by the Harvard University

Animal Care and Use Committee.

***Drosophila* strains**

Flies were cultured using standard media and techniques. All flies were raised at 25 °C. Fly

strains used in this work were: GMR94G06-GAL4 RRID:BDSC_40701, UAS-TeTxLC (aka UAS-

TeNT, RRID:BDSC_28838), *brp*[Df] (Akbergenova 2018), *brp*[69] (Kittel 2006), *rab3*[rup]

(RRID:BDSC_78045 (Graf 2007)), *liprin*[R60] (RRID:BDSC_8561, (Kaufmann, 2002)), *liprin*[F3ex15]

(RRID:BDSC_8563 (Kaufmann, 2002)).

Primary mouse hippocampal cultures

Primary hippocampal cultures were prepared using previously established protocols (Emperador-Melero et al., 2024; Held et al., 2020; Nyitrai et al., 2020; Wang et al., 2016; Wong et al., 2018). Hippocampi dissected from newborn pups (P0-P1) were digested in papain and dissociated. Dissociated cells were plated onto 12 mm, #1.5 glass coverslips in "plating medium" that contained Minimum Essential Medium (MEM) supplemented with 0.5% glucose, 0.02% NaHCO3, 0.1 mg/mL transferrin, 10% Fetal Select bovine serum (Atlas Biologicals FS-0500-AD), 2 mM L-glutamine, and 25 mg/mL insulin. Twenty-four hours after plating, the medium was exchanged with "growth medium" that contained MEM with 0.5% glucose, 0.02% NaHCO3, 0.1 mg/mL transferrin, 5% Fetal Select bovine serum (Atlas Biologicals FS-0500-AD), 2% B-27 supplement, and 0.5 mM L-glutamine. At 48 to 60 hours after plating, cytosine β-D-Arabinofuranoside (AraC) was added at a final concentration of 2-6 µM depending on glial growth. Cells were kept in an incubator at 37 °C until DIV15 to DIV16. For experiments modulating activity, a cocktail of drugs containing Tetrodotoxin (TTX), ω-Agatoxin IVA (ω-Aga) and ω-Conotoxin GVIA (ω-Cono) was added at DIV3 to a final concentration of 1 µM, 250 nm and 200 nM, respectively, and supplemented every three days. KCl was added to a final concentration of 50 mM from a 10X stock solution for 30 seconds before fixation.

Cell lines

HEK293T cells, an immortalized cell line of female origin, were purchased from ATCC (CRL-3216, RRID: CVCL_0063) as a mycoplasma-free cell line. These cells were maintained using established methods (Emperador-Melero et al., 2024; Held et al., 2020; Nyitrai et al., 2020; Wang et al., 2016; Wong et al., 2018). They were expanded and stored in liquid nitrogen until use. After thawing, cells were grown in Dulbecco's Modified Eagle Medium (DMEM) with 10% Fetal bovine serum (Atlas Biologicals F-0500-D) and 1% Penicillin-Streptomycin. HEK293T cells

were passaged at ratios between 1:3 and 1:10 up to ~25 times, when they were replaced with a

freshly thawed batch of cells.

Production of lentiviruses and transduction of primary mouse neurons

Lentiviruses were prepared using established methods (Emperador-Melero et al., 2024; Held et

al., 2020; Nyitrai et al., 2020; Wang et al., 2016; Wong et al., 2018). HEK293T cells were

transfected using the Ca^{2+} phosphate method with a combination of three packaging plasmids

(REV, RRE and VSV-G), and a lentiviral plasmid encoding either GFP-tagged Cre recombinase

(Kaeser lab plasmid code p009), an inactive version of Cre (p010), Liprin-α3 (p526) or a

lentivirus without an insert in the multiple cloning site (p008) at a molar ratio 1:1:1:1 and with a

total amount of ~4 µg DNA per T25 flask. At 20 to 30 hours after transfection, the medium was

changed to "growth medium" and 48 to 60 hours after transfection the supernatant was

collected. Viral transductions were done either with freshly collected viral supernatant or with

snap-frozen supernatant stored at -80 °C. Neuronal cultures were transduced with Cre or

inactive Cre at DIV2 for Ca_V2 ablation (Held et al., 2020), DIV5 for RIM+ELKS ablation

(Emperador-Melero et al., 2024; Tan et al., 2022; Wang et al., 2016), or DIV7 for Liprin-α

ablation (Emperador-Melero et al., 2024) following protocols established before. Transduction

efficiency was monitored via the presence of nuclear GFP fluorescence, and only cultures in

which no neurons without nuclear green fluorescence were readily detected were used for

experiments. Liprin-α mutant neuronal cultures were additionally transduced at DIV1 with either

lentivirus expressing either HA-tagged Liprin-α3 (p526; for control[L1-L4] neurons) or lentivirus

without an insert in the multiple cloning site (p008; for cQKO[L1-L4] neurons). pFSW HA-Liprin-α3

(p526) corresponds in sequence and numbering to NCBI Reference Sequence:

NP_001257914.1 with the addition of an N-terminal HA-tag and a short linker (sequence: M-

YPYDVPDYA-GAPS-C_3…C_{1192}) and has been described before (Emperador-Melero et al.,

2024, 2021b; Wong et al., 2018).

Immunofluorescence staining for STED and confocal microscopy of cultured

hippocampal neurons

Neurons were antibody-stained using previously established protocols (Emperador-Melero et al.,

2024, 2021b; Held et al., 2020; Wong et al., 2018). They were fixed at DIV15 to DIV16 in PBS

containing 2% PFA for 10 minutes at room temperature. For cultures stimulated with KCl, 50

mM KCl was also included in the fixation solution after the 30 s depolarization in described

below. Neurons were blocked and permeabilized in blocking solution (PBS, 3% BSA, 0.1%

Triton X-100) for 1 hour at room temperature and incubated with primary and secondary

antibodies, each overnight at 4 °C, in blocking solution. Three 5-minute washes with PBS were

performed after each antibody incubation. The cells were post-fixed in 4% PFA for 10 min,

washed in PBS, and mounted using mounting medium. Primary antibodies used were: mouse

anti-Dynamin-1 (Kaeser lab antibody code A242, 1:50, obtained from P. De Camilli, knockout-

validated for immunostaining in (Milosevic et al., 2011)), rabbit anti-Amphiphysin (A244, 1:200,

obtained from P. De Camilli, knockout-validated for immunostaining in (Di Paolo et al., 2002);

rabbit anti-PIPK1γ (A168, 1:500, obtained from P. De Camilli (Wenk et al., 2001)); mouse anti-

AP-180 (A246, 1:500, obtained from P. De Camilli, knockout-validated for immunostaining in

(Koo et al., 2015); rabbit anti-AP-180 (A219, 1:500; RRID: AB_887691); rabbit anti-Munc13-1

(A72, 1:500; RRID: AB_887733); mouse anti-Synaptophysin (A100, 1:500; RRID: AB_887824);

mouse anti-PSD-95 (A152, 1:500; RRID: AB_10698024); mouse anti-Bassoon (A85, 1:500;

RRID: AB_11181058); guinea pig anti-PSD-95 (A5, 1:500, RRID: AB_2619800); rabbit anti-

Synapsin-1 (A30, 1:500, RRID: AB_2200097); mouse anti-Gephyrin (A8; 1:500, RRID: AB_

2232546). Secondary antibodies used were: goat anti-guinea pig Alexa Fluor 633 (S34; 1:250,

RRID: AB_2535757), goat anti-mouse IgG2a Alexa Fluor 555 (S20; 1:250, RRID: AB_2535776),

goat anti-rabbit Alexa Fluor 488 (S5; 1:250, RRID: AB_2576217), goat anti-rabbit Alexa Fluor

555 (S22; 1:250, RRID: AB_2535849), goat anti-mouse Alexa Fluor 488 (S4; 1:250, RRID:

AB_2534088), goat anti-rabbit Alexa Fluor 633 (S33; 1:250, RRID: AB_2535731) and goat anti-

guinea pig Alexa Fluor 555 (S23; 1:250, RRID: AB_2535856).

Image acquisition and analyses for confocal and STED microscopy of cultured

hippocampal neurons

Images were acquired as previously established (Chin and Kaeser, 2024; Emperador-Melero et

al., 2024, 2021b, 2021a). A Leica SP8 Confocal/STED 3X microscope equipped with an oil-

immersion 100x objective (1.44 NA), a white laser, STED gated detectors, and 592, 660 and

770 nm depletion lasers was used to acquire 2048 pixels x 2048 pixels large images (pixel size

of 22.7 nm x 22.7 nm) that contained hundreds of synapses. Triple confocal scans for

Synaptophysin or Synapsin, an active zone (Munc13-1 or Bassoon) or postsynaptic density

(PSD-95) marker and a protein of interest were followed by double-color STED scans for the

active zone or postsynaptic density marker and the protein of interest. The exceptions were (1)

combinations containing Amphiphysin and PIPK1γ in Fig. 1, where Synaptophysin was also

imaged in STED, and (2) combinations containing Amphiphysin, PIPK1γ or AP-180 in Fig. 6,

where an antibody against Gephyrin was also added and imaged in confocal and STED modes

as well. Acquisition settings for a given staining and channel were identical for all images within

a batch of culture in which all conditions from one experiment were compared. To quantify

STED images, side-view synapses were selected by an experimenter blind to the protein of

interest. Side-view synapses were defined as those containing a vesicle cloud of 250 nm or

more in width from the marker to the inside of the presynaptic terminal and with a bar-like

Munc13-1, Bassoon or PSD-95 structure along its edge. A 750-nm long, rectangular area of

interest with a width exceeding that of the active zone or postsynaptic marker by up to five

pixels on each side was drawn perpendicular to the marker and across its center. After applying

a 5-pixel rolled average to the protein of interest and marker, line profiles of individual synapses

were aligned to the peak of the active zone or postsynaptic marker and averaged. Next, the

position of the maximum value of the endocytic protein relative to the maximum value of the

marker was calculated and plotted for each synapse. The maximum fluorescence value of the

endocytic protein within 136 nm relative to the peak fluorescence of the marker was also

measured and plotted. For Figs. 1 and 3, this region spanned 68 nm at each side of the peak of

the active zone marker. For Figs. 4 and 6, it spanned 136 nm from the PSD-95 peak towards

the presynaptic bouton. These distances were chosen to match the periactive zone area based

on the position of synaptic proteins at STED resolution (Wong et al., 2018). In each culture, line

profiles and peak intensities were normalized to the average signal in the condition used for

comparison (defined in the corresponding figure legend). In line profile plots, peaks are below

100% because peak intensities for proteins of interest are not always at the same position. En-

face synapses were selected as synapses that did not have a bar-like appearance of the active

zone or postsynaptic marker; instead, the area of the marker was surrounded by Synaptophysin

or Synapsin staining. The experimenter was blind to the protein of interest during en-face

synapse selection. For any given en-face synapse, only signals of the protein of interest that fell

within 136 nm of the edges of the synaptic vesicle cloud were included. This area was defined

by creating a binary mask for Synaptophysin or Synapsin and expanding it by 6 pixels (each

22.7 nm). Individual channels containing the marker or the protein of interest were next

thresholded. For any given experiment, thresholds were defined by an experimenter blind to the

condition through visual inspection of approximately ten images, and the same thresholds were

then applied to all images within an experiment. After thresholding, the "analyze particles"

function of Fiji was applied to detect individual objects and to measure their size, position and

intensity (the intensity was measured in the original, non-thresholded image). For quantification

of confocal signals, individual channels were analyzed with an automatic detection algorithm

(available at https://github.com/kaeserlab/3DSIM_Analysis_CL). With this algorithm, Otsu

thresholding was used to generate a presynaptic mask based on the Synaptophysin or

Synapsin signals. The synaptic mask was subsequently used to quantify the fluorescence

intensity levels of the protein of interest. To avoid detection artifacts, areas with somata and out-

of-focus areas were not included in the areas of interest. Confocal data were normalized to the

average in the condition that was used for comparison (noted in each figure legend) per culture.

For representative images, a smooth filter was added in some cases, brightness and contrast

were linearly adjusted, and images were interpolated. Identical adjustments were applied to

representative images of the same channel within an experiment. Quantifications were

performed on original images without brightness and contrast adjustments and without

background subtraction or resampling. Data were acquired and analyzed by an experimenter

blind to genotype.

Immunostaining of *Drosophila* NMJs

For analyses of NMJ morphology and protein localization, flies were maintained at low density

at 25 °C. Wandering third instar larvae were dissected in Ca^{2+}-free HL3.1 saline (70 mM NaCl, 5

mM KCl, 10 mM MgCl2, 10 mM NaHCO3, 5 mM trehalose, 5 mM HEPES, 115 mM sucrose

(Feng et al., 2004)) and fixed for 20 min in HL3.1 containing 4% PFA. Fixed larvae were

incubated in "blocking solution" containing 3% BSA, 0.1% Triton X-100 in PBS for 30 to 60 min

and incubated with primary antibody in blocking solution either overnight at 4 °C or for 2 hours

at room temperature. Samples were rinsed three times in a "washing buffer" containing 0.1%

Triton in PBS followed by three ten-minute washes in washing buffer. Samples were incubated

for one hour at room temperature with dye-conjugated secondary antibodies diluted to 1:250 in

washing buffer, followed by washing as after primary antibody incubation. Larvae were mounted

in Abberior Mount Liquid mounting medium. Primary antibodies used were rabbit anti-Nwk 970

(RRID:AB_2567353, knockout validated for immunostaining in (Coyle et al., 2004)), guinea pig

anti-Dynamin (gift from D. Dickman), mouse anti-Dynamin, (RRID:AB_397640, BD Biosciences

Clone 41, validated for immunostaining by RNAi knockdown), mouse anti-BRP

(RRID:AB_2314866, DSHB clone nc82, validated for immunostaining by RNAi in (Wagh et al.,

2006)), rabbit anti-Pak (gift of N. Harden, mutant-validated for immunostaining (Harden et al.,

1996)). Triple-labeling of Pak, FasII, and Brp was performed as follows: Samples were

incubated with anti-Pak and anti-FasII primary antibodies overnight at 4 °C, followed by

washing, species specific secondary antibody incubation, and further washing using the same

buffers and procedures as described above. Next, Brp was labeled using anti-Brp (nc82;

conjugated directly to Alexafluor-568, using a commercial antibody labeling kit (Thermofisher))

for four hours at room temperature, followed by three ten-minute washes.

Image acquisition and analyses of NMJs

Confocal images of NMJs were acquired at room temperature with a Zeiss 880FAS microscope

in SR mode, using a 63X (NA1.4) oil immersion objective and Zen Black software. All raw image

stacks were processed in Zen Blue to construct Airyscan images using 3D Airyscan processing

with automatic settings. Lateral and axial resolution were estimated to be ~175 nm and ~400

nm, respectively, by imaging Tetraspeck beads (Invitrogen) at 560 nm. To prepare images for

analyses, we excluded regions that were obscured by axon bundles or contained Type 1s

terminals. To measure intensities in whole boutons, a 3D presynaptic mask was generated as

follows. We produced a normalized sum image of Nwk, Dynamin, and Brp signals (except Fig.

5, which lacks Brp staining) by dividing each channel by its mean and summing the channels.

Sum images were then gaussian filtered (sigma = 5 pixels for all experiments except Fig. 8c-g,

where sigma = 4) and thresholded by intensity by a previously established algorithm (Li and

Tam, 1998) (except Fig. 8c-g, which used Otsu thesholding (Otsu, 1979)). The binary mask was

eroded by 4 pixels (1 pixel for Fig. 8c-g). Mask settings were manually confirmed to accurately

represent the 3D volume across images, and identical settings were used within an experiment.

For all images, the background was subtracted using the rolling ball method with a radius of 50

pixels, and signal intensities were measured in 3D using a custom FIJI script (available at

https://github.com/rodallab/nmj-measurement). For measurements of protein levels and

 polarization at the periactive zone, we used custom FIJI and Python scripts described in detail

(Del Signore et al., 2023) and available at https://github.com/rodallab/paz-analysis. First,

maximum intensity projections were made of the upper half of NMJ terminals, to analyze a

single plasma membrane surface. Second, a 2D mask of the total presynaptic area was

generated by summing all channels in each image and then thresholded by intensity using a

previously established algorithm (Li and Tam, 1998). Third, a periactive zone mesh composite

image was created by subtracting the Brp signal (except for Figs. 5 and 8) from the sum of Nwk

and Dyn signals. For Fig. 5, only the Nwk was used. For Fig. 8l-n, Brp and PAK signals were

subtracted from FasII to create the composite mask. Fourth, periactive zone units were detected

as local intensity minima and then expanded by the seeded region growing algorithm (via the IJ-

Plugins Toolkit). The thresholds for minima detection were computed automatically and

periactive zones detected at the edge of boutons (defined as having a mean Euclidean distance

map score of less than 7.5 pixels) were excluded from analysis as these are not planar (Del

Signore et al., 2023). As above, settings were applied identically within an experiment. Image

analyses settings were computed without manual user intervention and data acquisition and

analyses were not blinded. For representative images, brightness and contrast were linearly

adjusted and applied identically to representative images of the same channel within an

experiment.

Electrical stimulation of *Drosophila* NMJs

Control *white*[1118] animals were dissected at room temperature in HL3.1 with 0 mM extracellular

Ca^{2+}, and motor axons were cut close to the ventral ganglion. Prior to electrical stimulation, the

tissue sample was washed three times with 1 mL of HL3.1 solution containing 2 mM Ca^{2+} and 7

mM of glutamate (70 mM NaCl, 5 mM KCl, 10 mM $MgCl_2$, 10 mM $NaHCO_3$, 5 mM trehalose, 5

mM HEPES, 115 mM sucrose, 2 mM $CaCl_2$, 7 mM monosodium glutamate). The axon bundle

innervating segment A3 was sucked into a pipette and 40 Hz stimulation was administered for 3

minutes (5 V for 0.5 ms per stimulus) at room temperature. The stimulus was administered

through an A-M system 2100 stimulator using the ADinstruments lab chart software. Stimulation

was monitored by observing muscle contraction of the desired abdominal segment through the

eyepiece. For stimulation experiments, muscles 6/7 and muscle 4 were imaged and analyzed

together. As controls, comparisons were made to NMJs from the contralateral, unstimulated

muscles. After stimulation, the preparation was stretched and fixed within 30 seconds with 4%

PFA for 15 minutes, and samples were processed for immunohistochemistry as described

above.

Statistics

Data are shown as mean ± SEM. For analyses of STED images of hippocampal synapses, the

sample size is the number of analyzed synapses. For analyses of confocal images of

hippocampal synapses, the sample size is the number of analyzed images. For analyses of

Drosophila NMJs, the sample size is the number of NMJ terminals analyzed. Sample sizes and

statistical tests are included in each figure legend. Significance is reported as * $p < 0.05$, ** $p <$

0.01, or *** $p < 0.001$ and was assessed using parametric (t-test or one-way ANOVA) or non-

parametric (Mann-Whitney U or Kruskal-Wallis) tests depending on whether assumptions of

normality and homogeneity of variances were met (assessed using Shapiro or Levene's tests,

respectively). Tukey-Kramer or Holm corrections for multiple testing were applied. In Fig. 1 and

Fig. 1 – figure supplement 1, post-hoc comparisons between all groups were performed, and

only significance relative to the untreated condition is reported. For STED images, statistical

analyses were performed on the peak values of the line profiles. Statistical analyses were

performed in R.

Bibliography

Acuna C, Liu X, Südhof TC. 2016. How to Make an Active Zone: Unexpected Universal Functional Redundancy between RIMs and RIM-BPs. *Neuron* **91**:792–807. doi:10.1016/j.neuron.2016.07.042

Akbergenova Y, Cunningham KL, Zhang Y V., Weiss S, Littleton JT. 2018. Characterization of developmental and molecular factors underlying release heterogeneity at *Drosophila* synapses. *Elife* **7**. doi:10.7554/ELIFE.38268

Akbergenova Y, Matthias J, Littleton JT. 2025. Active zone maturation state controls synaptic output and release mode and is differentially regulated by neuronal activity. *bioRxiv* 2025.02.03.636302. doi:10.1101/2025.02.03.636302

Anantharam A, Bittner MA, Aikman RL, Stuenkel EL, Schmid SL, Axelrod D, Holz RW. 2011. A new role for the dynamin GTPase in the regulation of fusion pore expansion. *Mol Biol Cell* **22**:1907–1918. doi:10.1091/MBC.E11-02-0101

Aravamudan B, Fergestad T, Davis WS, Rodesch CK, Broadie K. 1999. *Drosophila* Unc-13 is essential for synaptic transmission. *Nature Neuroscience 1999 2:11* **2**:965–971. doi:10.1038/14764

Bai J, Hu Z, Dittman JS, Pym ECG, Kaplan JM. 2010. Endophilin functions as a membrane-bending molecule and is delivered to endocytic zones by exocytosis. *Cell* **143**:430–441. doi:10.1016/j.cell.2010.09.024

Bai J, Tucker WC, Chapman ER. 2003. PIP2 increases the speed of response of synaptotagmin and steers its membrane-penetration activity toward the plasma membrane. *Nature Structural & Molecular Biology 2003 11:1* **11**:36–44. doi:10.1038/nsmb709

Bailey CH, Chen M, Keller F, Kandel ER. 1992. Serotonin-Mediated Endocytosis of apCAM: an Early Step of Learning-Related Synaptic Growth in Aplysia. *Science (1979)* **256**:645–649. doi:10.1126/SCIENCE.1585177

Biederer T, Kaeser PS, Blanpied TA. 2017. Transcellular Nanoalignment of Synaptic Function. *Neuron* **96**:680–696. doi:10.1016/j.neuron.2017.10.006

Blanchette CR, Scalera AL, Harris KP, Zhao Z, Dresselhaus EC, Koles K, Yeh A, Apiki JK, Stewart BA, Rodal AA. 2022. Local regulation of extracellular vesicle traffic by the synaptic endocytic machinery. *Journal of Cell Biology* **221**. doi:10.1083/JCB.202112094/213086

Bloom O, Evergren E, Tomilin N, Kjaerulff O, Löw P, Brodin L, Pieribone VA, Greengard P, Shupliakov O. 2003. Colocalization of synapsin and actin during synaptic vesicle recycling. *Journal of Cell Biology* **161**:737–747. doi:10.1083/JCB.200212140

Böhme MA, Beis C, Reddy-Alla S, Reynolds E, Mampell MM, Grasskamp AT, Lützkendorf J, Bergeron DD, Driller JH, Babikir H, Göttfert F, Robinson IM, O'Kane CJ, Hell SW, Wahl MC, Stelzl U, Loll B, Walter AM, Sigrist SJ. 2016. Active zone scaffolds differentially accumulate Unc13 isoforms to tune Ca(2+) channel-vesicle coupling. *Nat Neurosci* **19**:1311–1320. doi:10.1038/nn.4364

Bolz S, Kaempf N, Puchkov D, Krauss M, Russo G, Soykan T, Schmied C, Lehmann M, Müller R, Schultz C, Perrais D, Maritzen T, Haucke V. 2023. Synaptotagmin 1-triggered lipid signaling facilitates coupling of exo- and endocytosis. *Neuron* **111**:3765-3774.e7. doi:10.1016/j.neuron.2023.08.016

Brenig J, De Boor S, Knyphausen P, Kuhlmann N, Wroblowski S, Baldus L, Scislowski L, Artz O, Trauschies P, Baumann U, Neundorf I, Lammers M. 2015. Structural and biochemical basis for the inhibitory effect of liprin-α3 on mouse diaphanous 1 (mDia1) Function. *Journal of Biological Chemistry* **290**:14314–14327. doi:10.1074/JBC.M114.621946

Cao YQ, Piedras-Rentería ES, Smith GB, Chen G, Harata NC, Tsien RW. 2004. Presynaptic Ca2+ Channels Compete for Channel Type-Preferring Slots in Altered

Neurotransmission Arising from Ca2+ Channelopathy. *Neuron* **43**:387–400. doi:10.1016/J.NEURON.2004.07.014

Chia PH, Patel MR, Shen K. 2012. NAB-1 instructs synapse assembly by linking adhesion molecules and F-actin to active zone proteins. *Nature Neuroscience 2012 15:2* **15**:234–242. doi:10.1038/NN.2991

Chin M, Kaeser PS. 2024. The intracellular C-terminus confers compartment-specific targeting of voltage-gated calcium channels. *Cell Rep* **43**:114428. doi:10.1016/J.CELREP.2024.114428

Cocucci E, Gaudin R, Kirchhausen T. 2014. Dynamin recruitment and membrane scission at the neck of a clathrin-coated pit. *Mol Biol Cell* **25**:3595–3609. doi:10.1091/mbc.E14-07-1240

Coyle IP, Koh YH, Lee WCM, Slind J, Fergestad T, Littleton JT, Ganetzky B. 2004. Nervous Wreck, an SH3 Adaptor Protein that Interacts with Wsp, Regulates Synaptic Growth in *Drosophila. Neuron* **41**:521–534. doi:10.1016/S0896-6273(04)00016-9

Cunningham KL, Sauvola CW, Tavana S, Troy Littleton J. 2022. Regulation of presynaptic Ca2+ channel abundance at active zones through a balance of delivery and turnover. *Elife* **11**. doi:10.7554/ELIFE.78648

de Jong APH, Roggero CM, Ho M-R, Wong MY, Brautigam CA, Rizo J, Kaeser PS. 2018. RIM C2B Domains Target Presynaptic Active Zone Functions to PIP2-Containing Membranes. *Neuron* **98**:335-349.e7. doi:10.1016/j.neuron.2018.03.011

Del Signore SJ, Kelley CF, Messelaar EM, Lemos T, Marchan MF, Ermanoska B, Mund M, Fai TG, Kaksonen M, Rodal AA. 2021. An autoinhibitory clamp of actin assembly constrains and directs synaptic endocytosis. *Elife* **10**. doi:10.7554/eLife.69597

Del Signore SJ, Mitzner MG, Silveira AM, Fai TG, Rodal AA. 2023. An approach for quantitative mapping of synaptic periactive zone architecture and organization. *Mol Biol Cell* **34**. doi:10.1091/MBC.E22-08-0372

Di Paolo G, Moskowitz HS, Gipson K, Wenk MR, Voronov S, Obayashi M, Flavell R, Fitzsimonds RM, Ryan TA, De Camill P. 2004. Impaired PtdIns(4,5)P2 synthesis in nerve terminals produces defects in synaptic vesicle trafficking. *Nature 2004 431:7007* **431**:415–422. doi:10.1038/nature02896

Di Paolo G, Sankaranarayanan S, Wenk MR, Daniell L, Perucco E, Caldarone BJ, Flavell R, Picciotto MR, Ryan TA, Cremona O, De Camilli P. 2002. Decreased synaptic vesicle recycling efficiency and cognitive deficits in amphiphysin 1 knockout mice. *Neuron* **33**:789–804. doi:10.1016/S0896-6273(02)00601-3

Emperador-Melero J, Andersen JW, Metzbower SR, Levy AD, Dharmasri PA, de Nola G, Blanpied TA, Kaeser PS. 2024. Distinct active zone protein machineries mediate Ca2+ channel clustering and vesicle priming at hippocampal synapses. *Nature Neuroscience 2024* 1–15. doi:10.1038/s41593-024-01720-5

Emperador-Melero J, de Nola G, Kaeser PS. 2021a. Intact synapse structure and function after combined knockout of PTPδ, PTPσ, and LAR. *Elife* **10**. doi:10.7554/eLife.66638

Emperador-Melero J, Kaeser PS. 2020. Assembly of the presynaptic active zone. *Curr Opin Neurobiol* **63**:95–103. doi:10.1016/j.conb.2020.03.008

Emperador-Melero J, Wong MY, Wang SSH, de Nola G, Nyitrai H, Kirchhausen T, Kaeser PS. 2021b. PKC-phosphorylation of Liprin-α3 triggers phase separation and controls presynaptic active zone structure. *Nat Commun* **12**:3057. doi:10.1038/s41467-021-23116-w

Estes PS, Roos J, Van Der Bliek A, Kelly RB, Krishnan KS, Ramaswami M. 1996. Traffic of Dynamin within Individual *Drosophila* Synaptic Boutons Relative to Compartment-Specific Markers. *Journal of Neuroscience* **16**:5443–5456. doi:10.1523/JNEUROSCI.16-17-05443.1996

Feng Y, Ueda A, Wu CF. 2004. A modified minimal hemolymph-like solution, HL3.1, for physiological recordings at the neuromuscular junctions of normal and mutant *Drosophila* larvae. *J Neurogenet* **18**:377–402. doi:10.1080/01677060490894522

Ferguson SM, Brasnjo G, Hayashi M, Wölfel M, Collesi C, Giovedi S, Raimondi A, Gong L-W, Ariel P, Paradise S, O'toole E, Flavell R, Cremona O, Miesenböck G, Ryan TA, De Camilli P. 2007. A selective activity-dependent requirement for dynamin 1 in synaptic vesicle endocytosis. *Science* **316**:570–574. doi:10.1126/science.1140621

Fouquet W, Owald D, Wichmann C, Mertel S, Depner H, Dyba M, Hallermann S, Kittel RJ, Eimer S, Sigrist SJ. 2009. Maturation of active zone assembly by *Drosophila* Bruchpilot. *Journal of Cell Biology* **186**:129–145. doi:10.1083/JCB.200812150

Fu Y, Huang ZJ. 2010. Differential dynamics and activity-dependent regulation of α- and β-neurexins at developing GABAergic synapses. *Proc Natl Acad Sci U S A* **107**:22699–22704. doi:10.1073/PNAS.1011233108

Ganguly A, Sharma R, Boyer NP, Wernert F, Phan S, Boassa D, Parra L, Das U, Caillol G, Han X, Yates JR 3rd, Ellisman MH, Leterrier C, Roy S. 2021. Clathrin packets move in slow axonal transport and deliver functional payloads to synapses. *Neuron* **109**:2884-2901.e7. doi:10.1016/j.neuron.2021.08.016

Gerth F, Jäpel M, Pechstein A, Kochlamazashvili G, Lehmann M, Puchkov D, Onofri F, Benfenati F, Nikonenko AG, Fredrich K, Shupliakov O, Maritzen T, Freund C, Haucke V. 2017. Intersectin associates with synapsin and regulates its nanoscale localization and function. *Proc Natl Acad Sci U S A* **114**:12057–12062. doi:10.1073/pnas.1715341114

Gimber N, Tadeus G, Maritzen T, Schmoranzer J, Haucke V. 2015. Diffusional spread and confinement of newly exocytosed synaptic vesicle proteins. *Nature Communications 2015 6:1* **6**:1–11. doi:10.1038/ncomms9392

Graf ER, Daniels RW, Burgess RW, Schwarz TL, DiAntonio A. 2009. Rab3 Dynamically Controls Protein Composition at Active Zones. *Neuron* **64**:663–677. doi:10.1016/j.neuron.2009.11.002

Graf ER, Valakh V, Wright CM, Wu C, Liu Z, Zhang YQ, DiAntonio A. 2012. RIM Promotes Calcium Channel Accumulation at Active Zones of the *Drosophila* Neuromuscular Junction. *Journal of Neuroscience* **32**:16586–16596. doi:10.1523/JNEUROSCI.0965-12.2012

Harden N, Lee J, Loh H-Y, Ong Y-M, Tan I, Leung T, Manser E, Lim L. 1996. A *Drosophila* Homolog of the Rac- and Cdc42-Activated Serine/Threonine Kinase PAK Is a Potential Focal Adhesion and Focal Complex Protein That Colocalizes with Dynamic Actin Structures. *Mol Cell Biol* **16**:1896–1908. doi:10.1128/MCB.16.5.1896

Harris KP, Littleton JT. 2015. Transmission, Development, and Plasticity of Synapses. *Genetics* **201**:345–375. doi:10.1534/GENETICS.115.176529

Haucke V, De Camilli P. 1999. AP-2 Recruitment to Synaptotagmin Stimulated by Tyrosine-Based Endocytic Motifs. *Science (1979)* **285**:1268–1271. doi:10.1126/SCIENCE.285.5431.1268

Haucke V, Neher E, Sigrist SJ. 2011. Protein scaffolds in the coupling of synaptic exocytosis and endocytosis. *Nat Rev Neurosci* **12**:127–138. doi:10.1038/nrn2948

Held RG, Liu C, Ma K, Ramsey AM, Tarr TB, De Nola G, Wang SSH, Wang J, van den Maagdenberg AMJM, Schneider T, Sun J, Blanpied TA, Kaeser PS. 2020. Synapse and Active Zone Assembly in the Absence of Presynaptic Ca(2+) Channels and Ca(2+) Entry. *Neuron* **107**:667-683.e9. doi:10.1016/j.neuron.2020.05.032

Hua Y, Sinha R, Thiel CS, Schmidt R, Hüve J, Martens H, Hell SW, Egner A, Klingauf J. 2011. A readily retrievable pool of synaptic vesicles. *Nature Neuroscience 2011 14:7* **14**:833–839. doi:10.1038/NN.2838

Imoto Y, Raychaudhuri S, Ma Y, Fenske P, Sandoval E, Itoh K, Blumrich EM, Matsubayashi HT, Mamer L, Zarebidaki F, Söhl-Kielczynski B, Trimbuch T, Nayak S, Iwasa JH, Liu J, Wu B, Ha T, Inoue T, Jorgensen EM, Cousin MA, Rosenmund C, Watanabe S. 2022. Dynamin is primed at endocytic sites for ultrafast endocytosis. *Neuron* **110**:2815-2835.e13. doi:10.1016/J.NEURON.2022.06.010

Imoto Y, Xue J, Luo L, Raychaudhuri S, Itoh K, Ma Y, Craft GE, Kwan AH, Ogunmowo TH, Ho A, Mackay JP, Ha T, Watanabe S, Robinson PJ. 2024. Dynamin 1xA interacts with Endophilin A1 via its spliced long C-terminus for ultrafast endocytosis. *EMBO Journal*. doi:10.1038/S44318-024-00145-X

Jäpel M, Gerth F, Sakaba T, Bacetic J, Yao L, Koo SJ, Maritzen T, Freund C, Haucke V. 2020. Intersectin-Mediated Clearance of SNARE Complexes Is Required for Fast Neurotransmission. *Cell Rep* **30**:409-420.e6. doi:10.1016/j.celrep.2019.12.035

Jenett A, Rubin GM, Ngo TTB, Shepherd D, Murphy C, Dionne H, Pfeiffer BD, Cavallaro A, Hall D, Jeter J, Iyer N, Fetter D, Hausenfluck JH, Peng H, Trautman ET, Svirskas RR, Myers EW, Iwinski ZR, Aso Y, DePasquale GM, Enos A, Hulamm P, Lam SCB, Li HH, Laverty TR, Long F, Qu L, Murphy SD, Rokicki K, Safford T, Shaw K, Simpson JH, Sowell A, Tae S, Yu Y, Zugates CT. 2012. A GAL4-Driver Line Resource for *Drosophila* Neurobiology. *Cell Rep* **2**:991–1001. doi:10.1016/J.CELREP.2012.09.011

Jiang A, Kudo K, Gormal RS, Ellis S, Guo S, Wallis TP, Longfield SF, Robinson PJ, Johnson ME, Joensuu M, Meunier FA. 2024. Dynamin1 long- and short-tail isoforms exploit distinct recruitment and spatial patterns to form endocytic nanoclusters. *Nature Communications 2024 15:1* **15**:1–21. doi:10.1038/s41467-024-47677-8

Jiao W, Shupliakov A, Shupliakov O. 2010. A semi-correlative technique for the subcellular localization of proteins in *Drosophila* synapses. *J Neurosci Methods* **185**:273–279. doi:10.1016/J.JNEUMETH.2009.10.010

Jorgensen EM, Hartwieg E, Schuske K, Nonet ML, Jin Y, Horvitz HR. 1995. Defective recycling of synaptic vesicles in synaptotagmin mutants of Caenorhabditis elegans. *Nature 1995 378:6553* **378**:196–199. doi:10.1038/378196a0

Kaeser PS, Deng L, Chávez AE, Liu X, Castillo PE, Südhof TC. 2009. ELKS2α/CAST Deletion Selectively Increases Neurotransmitter Release at Inhibitory Synapses. *Neuron* **64**:227–239. doi:10.1016/j.neuron.2009.09.019

Kaeser P S, Deng L, Wang Y, Dulubova I, Liu X, Rizo J, Sudhof TC. 2011. RIM proteins tether Ca2+ channels to presynaptic active zones via a direct PDZ-domain interaction. *Cell* **144**:282–295. doi:10.1016/j.cell.2010.12.029

Kaeser P S, Kwon HB, Chiu CQ, Deng L, Castillo PE, Sudhof TC. 2008. RIM1alpha and RIM1beta are synthesized from distinct promoters of the RIM1 gene to mediate differential but overlapping synaptic functions. *J Neurosci* **28**:13435–13447. doi:28/50/13435 [pii] 10.1523/JNEUROSCI.3235-08.2008

Kaksonen M, Roux A. 2018. Mechanisms of clathrin-mediated endocytosis. *Nat Rev Mol Cell Biol* **19**:313–326. doi:10.1038/nrm.2017.132

Kaufmann N, DeProto J, Ranjan R, Wan H, Van Vactor D. 2002. *Drosophila* liprin-alpha and the receptor phosphatase Dlar control synapse morphogenesis. *Neuron* **34**:27–38. doi:10.1016/s0896-6273(02)00643-8

Kawasaki F, Hazen M, Ordway RW. 2000. Fast synaptic fatigue in shibire mutants reveals a rapid requirement for dynamin in synaptic vesicle membrane trafficking. *Nat Neurosci* **3**:859–860. doi:10.1038/78753

Kittel RJ, Wichmann C, Rasse TM, Fouquet W, Schmidt M, Schmid A, Wagh DA, Pawlu C, Kellner RR, Willig KI, Hell SW, Buchner E, Heckmann M, Sigrist SJ. 2006. Bruchpilot promotes active zone assembly, Ca2+ channel clustering, and vesicle release. *Science (1979)* **312**:1051–1054. doi:10.1126/SCIENCE.1126308

Koh TW, Korolchuk VI, Wairkar YP, Jiao W, Evergren E, Pan H, Zhou Y, Venken KJT, Shupliakov O, Robinson IM, O'Kane CJ, Bellen HJ. 2007. Eps15 and Dap160 control synaptic vesicle membrane retrieval and synapse development. *Journal of Cell Biology* **178**:309–322. doi:10.1083/JCB.200701030

Koo SJ, Kochlamazashvili G, Rost B, Puchkov D, Gimber N, Lehmann M, Tadeus G, Schmoranzer J, Rosenmund C, Haucke V, Maritzen T. 2015. Vesicular Synaptobrevin/VAMP2 Levels Guarded by AP180 Control Efficient Neurotransmission. *Neuron* **88**:330–344. doi:10.1016/j.neuron.2015.08.034

Li CH, Tam PKS. 1998. An iterative algorithm for minimum cross entropy thresholding. *Pattern Recognit Lett* **19**:771–776. doi:10.1016/S0167-8655(98)00057-9

Li TN, Chen YJ, Lu TY, Wang YT, Lin HC, Yao CK. 2020. A positive feedback loop between flower and pi(4,5)p2 at periactive zones controls bulk endocytosis in *Drosophila*. *Elife* **9**:1–33. doi:10.7554/ELIFE.60125

Liu C., Bickford LS, Held RG, Nyitrai H, Sudhof TC, Kaeser PS. 2014. The Active Zone Protein Family ELKS Supports Ca2+ Influx at Nerve Terminals of Inhibitory Hippocampal Neurons. *Journal of Neuroscience* **34**:12289–12303. doi:10.1523/JNEUROSCI.0999-14.2014

Liu C, Kershberg L, Wang J, Schneeberger S, Kaeser PS. 2018. Dopamine Secretion Is Mediated by Sparse Active Zone-like Release Sites. *Cell* **172**:706-718.e15. doi:10.1016/j.cell.2018.01.008

Liu KSY, Siebert M, Mertel S, Knoche E, Wegener S, Wichmann C, Matkovic T, Muhammad K, Depner H, Mettke C, Bückers J, Hell SW, Müller M, Davis GW, Schmitz D, Sigrist SJ. 2011. RIM-binding protein, a central part of the active zone, is essential for neurotransmitter release. *Science (1979)* **334**:1565–1569. doi:10.1126/SCIENCE.1212991

Marcó de la Cruz B, Campos J, Molinaro A, Xie X, Jin G, Wei Z, Acuna C, Sterky FH. 2024. Liprin-α proteins are master regulators of human presynapse assembly. *Nature Neuroscience 2024 27:4* **27**:629–642. doi:10.1038/s41593-024-01592-9

McDonald NA, Fetter RD, Shen K. 2020. Assembly of synaptic active zones requires phase separation of scaffold molecules. *Nature 2020 588:7838* **588**:454–458. doi:10.1038/s41586-020-2942-0

Milosevic I, Giovedi S, Lou X, Raimondi A, Collesi C, Shen H, Paradise S, O'Toole E, Ferguson S, Cremona O, De Camilli P. 2011. Recruitment of endophilin to clathrin-coated pit necks is required for efficient vesicle uncoating after fission. *Neuron* **72**:587–601. doi:10.1016/j.neuron.2011.08.029

Moro A, van Nifterick A, Toonen RF, Verhage M. 2021. Dynamin controls neuropeptide secretion by organizing dense-core vesicle fusion sites. *Sci Adv* **7**. doi:10.1126/sciadv.abf0659

Mueller VJ, Wienisch M, Nehring RB, Klingauf J. 2004. Monitoring Clathrin-Mediated Endocytosis during Synaptic Activity. *Journal of Neuroscience* **24**:2004–2012. doi:10.1523/JNEUROSCI.4080-03.2004

Nyitrai H, Wang SSH, Kaeser PS. 2020. ELKS1 Captures Rab6-Marked Vesicular Cargo in Presynaptic Nerve Terminals. *Cell Rep* **31**:107712. doi:10.1016/j.celrep.2020.107712

Ogunmowo TH, Jing H, Raychaudhuri S, Kusick GF, Imoto Y, Li S, Itoh K, Ma Y, Jafri H, Dalva MB, Chapman ER, Ha T, Watanabe S, Liu J. 2023. Membrane compression by synaptic vesicle exocytosis triggers ultrafast endocytosis. *Nature Communications 2023 14:1* **14**:1–16. doi:10.1038/s41467-023-38595-2

Otsu N. 1979. THRESHOLD SELECTION METHOD FROM GRAY-LEVEL HISTOGRAMS. *IEEE Trans Syst Man Cybern* **SMC-9**:62–66. doi:10.1109/TSMC.1979.4310076

Owald D, Fouquet W, Schmidt M, Wichmann C, Mertel S, Depner H, Christiansen F, Zube C, Quentin C, Körner J, Urlaub H, Mechtler K, Sigrist SJ. 2010. A Syd-1 homologue

regulates pre- and postsynaptic maturation in *Drosophila*. *J Cell Biol* **188**:565–579. doi:10.1083/jcb.200908055

Peled ES, Isacoff EY. 2011. Optical quantal analysis of synaptic transmission in wild-type and rab3-mutant *Drosophila* motor axons. *Nature Neuroscience 2011 14:4* **14**:519–526. doi:10.1038/NN.2767

Pereverzev A, Mikhna M, Vajna R, Gissel C, Henry M, Weiergra¨ber M, Weiergra¨ber W, Ju¨ J, Hescheler J, Smyth N, Schneider T. 2002. Disturbances in Glucose-Tolerance, Insulin-Release, and Stress-Induced Hyperglycemia upon Disruption of the Cav2.3 (α1E) Subunit of Voltage-Gated Ca2+ Channels. *Molecular Endocrinology* **16**:884–895. doi:10.1210/MEND.16.4.0801

Perez de Arce K, Schrod N, Metzbower SWR, Allgeyer E, Kong GK-W, Tang A-H, Krupp AJ, Stein V, Liu X, Bewersdorf J, Blanpied TA, Lucić V, Biederer T. 2015. Topographic Mapping of the Synaptic Cleft into Adhesive Nanodomains. *Neuron* **88**:1165–1172. doi:10.1016/j.neuron.2015.11.011

Poskanzer KE, Fetter RD, Davis GW. 2006. Discrete Residues in the C2B Domain of Synaptotagmin I Independently Specify Endocytic Rate and Synaptic Vesicle Size. *Neuron* **50**:49–62. doi:10.1016/J.NEURON.2006.02.021

Raimondi A, Ferguson SM, Lou X, Armbruster M, Paradise S, Giovedi S, Messa M, Kono N, Takasaki J, Cappello V, O'Toole E, Ryan TA, De Camilli P. 2011. Overlapping role of dynamin isoforms in synaptic vesicle endocytosis. *Neuron* **70**:1100–1114. doi:10.1016/j.neuron.2011.04.031

Rao Y, Ma Q, Vahedi-Faridi A, Sundborger A, Pechstein A, Puchkov D, Luo L, Shupliakov O, Saenger W, Haucke V. 2010. Molecular basis for SH3 domain regulation of F-BAR-mediated membrane deformation. *Proc Natl Acad Sci U S A* **107**:8213–8218. doi:10.1073/PNAS.1003478107

Reshetniak S, Ußling J, Perego E, Rammner B, Schikorski T, Fornasiero EF, Truckenbrodt S, Köster S, Rizzoli SO. 2020. A comparative analysis of the mobility of 45 proteins in the synaptic bouton. *EMBO J* **39**. doi:10.15252/EMBJ.2020104596

Rodal AA, Motola-Barnes RN, Littleton JT. 2008. Nervous Wreck and Cdc42 Cooperate to Regulate Endocytic Actin Assembly during Synaptic Growth. *Journal of Neuroscience* **28**:8316–8325. doi:10.1523/JNEUROSCI.2304-08.2008

Roos J, Kelly RB. 1999. The endocytic machinery in nerve terminals surrounds sites of exocytosis. *Current Biology* **9**:1411–1414. doi:10.1016/S0960-9822(00)80087-1

Saheki Y, De Camilli P. 2012. Synaptic vesicle endocytosis. *Cold Spring Harb Perspect Biol* **4**:a005645. doi:10.1101/cshperspect.a005645

Salkoff L, Kelly L. 1978. Temperature-induced seizure and frequency-dependent neuromuscular block in a ts mutant of *Drosophila*. *Nature 1978 273:5658* **273**:156–158. doi:10.1038/273156a0

Samasilp P, Chan SA, Smith C. 2012. Activity-Dependent Fusion Pore Expansion Regulated by a Calcineurin-Dependent Dynamin-Syndapin Pathway in Mouse Adrenal Chromaffin Cells. *Journal of Neuroscience* **32**:10438–10447. doi:10.1523/JNEUROSCI.1299-12.2012

Sclip A, Südhof TC. 2023. Combinatorial expression of neurexins and LAR-type phosphotyrosine phosphatase receptors instructs assembly of a cerebellar circuit. *Nature Communications 2023 14:1* **14**:1–14. doi:10.1038/s41467-023-40526-0

Sclip A, Südhof TC. 2020. LAR receptor phospho-tyrosine phosphatases regulate NMDA-receptor responses. *Elife* **9**. doi:10.7554/eLife.53406

Stanishneva-Konovalova TB, Kelley CF, Eskin TL, Messelaar EM, Wasserman SA, Sokolova OS, Rodal AA. 2016. Coordinated autoinhibition of F-BAR domain membrane binding and WASp activation by Nervous Wreck. *Proc Natl Acad Sci U S A* **113**:E5552–E5561. doi:10.1073/PNAS.1524412113

Sudhof TC. 2012. The presynaptic active zone. *Neuron* **75**:11–25.
  doi:10.1016/j.neuron.2012.06.012

Sweeney ST, Broadie K, Keane J, Niemann H, O'Kane CJ. 1995. Targeted expression of
  tetanus toxin light chain in *Drosophila* specifically eliminates synaptic transmission
  and causes behavioral defects. *Neuron* **14**:341–351. doi:10.1016/0896-
  6273(95)90290-2

Tan C, Wang SSH, de Nola G, Kaeser PS. 2022. Rebuilding essential active zone
  functions within a synapse. *Neuron* **110**:1498-1515.e8.
  doi:10.1016/J.NEURON.2022.01.026

Tang A-H, Chen H, Li TP, Metzbower SR, MacGillavry HD, Blanpied TA. 2016. A trans-
  synaptic nanocolumn aligns neurotransmitter release to receptors. *Nature* **536**:210–
  214. doi:10.1038/nature19058

Taylor MJ, Perrais D, Merrifield CJ. 2011. A high precision survey of the molecular
  dynamics of mammalian clathrin-mediated endocytosis. *PLoS Biol* **9**:e1000604.
  doi:10.1371/journal.pbio.1000604

Terry-Lorenzo RT, Torres VI, Wagh D, Galaz J, Swanson SK, Florens L, Washburn MP,
  Waites CL, Gundelfinger ED, Reimer RJ, Garner CC. 2016. Trio, a Rho Family GEF,
  Interacts with the Presynaptic Active Zone Proteins Piccolo and Bassoon. *PLoS One*
  11:e0167535. doi:10.1371/JOURNAL.PONE.0167535

Thivaios S, Schwenk J, Brechet A, Boudkkazi S, Henneken P, Kocylowski MK, Haupt A,
  Kaminski D, Broucke J-B van den, Schreiner D, Schulte U, Sterky FH, Yuzaki M,
  Scheiffele P, Fakler B. 2024. A ternary Neurexin-T178-PTPR complex represents a
  core-module of neuronal synapse organization. *bioRxiv* 2024.07.16.603715.
  doi:10.1101/2024.07.16.603715

Todorov B, Van De Ven RCG, Kaja S, Broos LAM, Verbeek SJ, Plomp JJ, Ferrari MD,
  Frants RR, Van Den Maagdenberg AMJM. 2006. Conditional inactivation of the
  Cacna1a gene in transgenic mice. *genesis* **44**:589–594. doi:10.1002/DVG.20255

Trotter JH, Hao J, Maxeiner S, Tsetsenis T, Liu Z, Zhuang X, Südhof TC. 2019. Synaptic
  neurexin-1 assembles into dynamically regulated active zone nanoclusters. *J Cell Biol*
  218:2677–2698. doi:10.1083/jcb.201812076

Wagh DA, Rasse TM, Asan E, Hofbauer A, Schwenkert I, Dürrbeck H, Buchner S,
  Dabauvalle MC, Schmidt M, Qin G, Wichmann C, Kittel R, Sigrist SJ, Buchner E.
  2006. Bruchpilot, a protein with homology to ELKS/CAST, is required for structural
  integrity and function of synaptic active zones in *Drosophila*. *Neuron* **49**:833–844.
  doi:10.1016/j.neuron.2006.02.008

Wahl S, Katiyar R, Schmitz F. 2013. A Local, Periactive Zone Endocytic Machinery at
  Photoreceptor Synapses in Close Vicinity to Synaptic Ribbons. *Journal of
  Neuroscience* **33**:10278–10300. doi:10.1523/JNEUROSCI.5048-12.2013

Wang SSH, Held RG, Wong MY, Liu C, Karakhanyan A, Kaeser PS. 2016. Fusion
  Competent Synaptic Vesicles Persist upon Active Zone Disruption and Loss of Vesicle
  Docking. *Neuron* **91**:777–791. doi:10.1016/J.NEURON.2016.07.005

Watanabe S, Boucrot E. 2017. Fast and ultrafast endocytosis. *Curr Opin Cell Biol* **47**:64–
  71. doi:10.1016/j.ceb.2017.02.013

Watanabe S, Rost BR, Camacho-Pérez M, Davis MW, Söhl-Kielczynski B, Rosenmund C,
  Jorgensen EM. 2013. Ultrafast endocytosis at mouse hippocampal synapses. *Nature*
  504:242–247. doi:10.1038/nature12809

Watanabe S, Trimbuch T, Camacho-Pérez M, Rost BR, Brokowski B, Söhl-Kielczynski B,
  Felies A, Davis MW, Rosenmund C, Jorgensen EM. 2014. Clathrin regenerates
  synaptic vesicles from endosomes. *Nature* **515**:228–233. doi:10.1038/nature13846

Wenk MR, Pellegrini L, Klenchin VA, Di Paolo G, Chang S, Daniell L, Arioka M, Martin TF, De Camilli P. 2001. PIP kinase Igamma is the major PI(4,5)P(2) synthesizing enzyme at the synapse. *Neuron* **32**:79–88. doi:10.1016/s0896-6273(01)00456-1

Wilhelm BG, Mandad S, Truckenbrodt S, Kröhnert K, Schäfer C, Rammner B, Koo SJ, Claßen GA, Krauss M, Haucke V, Urlaub H, Rizzoli SO. 2014. Composition of isolated synaptic boutons reveals the amounts of vesicle trafficking proteins. *Science* **344**:1023–1028. doi:10.1126/science.1252884

Winther ÅME, Jiao W, Vorontsova O, Rees KA, Koh TW, Sopova E, Schulze KL, Bellen HJ, Shupliakov O. 2013. The dynamin-binding domains of Dap160/intersectin affect bulk membrane retrieval in synapses. *J Cell Sci* **126**:1021–1031. doi:10.1242/JCS.118968

Winther ÅME, Vorontsova O, Rees KA, Näreoja T, Sopova E, Jiao W, Shupliakov O. 2015. An Endocytic Scaffolding Protein together with Synapsin Regulates Synaptic Vesicle Clustering in the *Drosophila* Neuromuscular Junction. *J Neurosci* **35**:14756–14770. doi:10.1523/JNEUROSCI.1675-15.2015

Wong MY, Liu C, Wang SSH, Roquas ACF, Fowler SC, Kaeser PS. 2018. Liprin-alpha3 controls vesicle docking and exocytosis at the active zone of hippocampal synapses. *Proc Natl Acad Sci U S A* **115**:2234–2239. doi:10.1073/pnas.1719012115

Zhang JZ, Davletov BA, Südhof TC, Anderson RGW. 1994. Synaptotagmin I is a high affinity receptor for clathrin AP-2: Implications for membrane recycling. *Cell* **78**:751–760. doi:10.1016/S0092-8674(94)90442-1

Zhen M, Jin Y. 1999. The liprin protein SYD-2 regulates the differentiation of presynaptic termini in C. elegans. *Nature* **401**:371–375. doi:10.1038/43886

Zurner M, Schoch S. 2009. The mouse and human Liprin-alpha family of scaffolding proteins: genomic organization, expression profiling and regulation by alternative splicing. *Genomics* **93**:243–253. doi:10.1016/j.ygeno.2008.10.007

Figure 1. Deployment of endocytic proteins after chronic silencing or acute depolarization of mouse hippocampal neurons

(**a**) Schematic of the experiment in mouse hippocampal neurons. For chronic silencing, a cocktail ("blockers") with tetrodotoxin (TTX, 1 μM final concentration), ω-agatoxin IVA (ω-Aga, 250 nM) and ω-conotoxin GVIA (ω-Cono, 200 nM) was added every three days starting DIV3. For acute depolarization ("KCl"), neurons were stimulated with 50 mM KCl for 30 seconds before fixation.

(**b-i**) Example side-view synapses and average line profiles for neurons stained for Bassoon, Synaptophysin, and Amphiphysin (b+c), PIPK1γ (d+e), or AP-180 (f+g), or Synaptophysin, Munc13-1, and Dynamin-1 (h+i). Neurons were stained for a protein of interest (Amphiphysin, PIPK1γ, AP-180 or Dynamin-1; imaged in STED), an active zone marker (Bassoon or Munc13-1; imaged in STED), and Synaptophysin (imaged in confocal). An area of interest was positioned perpendicular to the center of the active zone marker, and synapses were aligned via the peak fluorescence of the active zone marker in the average profiles (c, e, g, i). Line profile plots were normalized to the average signal in the untreated condition. Grey dotted lines in c, e, g and i mark average levels in the untreated condition and grey bars represent the active zone and periactive zone area; n in b, c (synapses/cultures): untreated 38/3, blockers 40/3, KCl 45/3; d, e: untreated 51/3, blockers 49/3, KCl 42/3; f, g: untreated 58/3, blockers 61/3, KCl 50/3; h, i:, untreated 45/3, blockers 43/3, KCl 42/3.

(**j, k**) Quantification of the peak-to-peak distance of the active zone marker and the protein of interest (j), and of the peak levels in the periactive zone area (k). The periactive zone area is defined as an area within 68 nm on each side of the peak of the active zone marker (grey shaded areas in c, e, g and i); n as in b-i.

(**l-q**) Example en-face synapses (l-o) and quantification of the number of objects (p) and distance of these objects to the center of the active zone marker (q) of the experiment shown in a-l, n in p, q (synapses/cultures); Amphiphysin untreated 20/3, blockers 21/3, KCl 21/3; PIPK1γ, untreated 23/3, blockers 21/3, KCl 20/3; AP-180, untreated 21/3, blockers 24/3, KCl 18/3; Dynamin-1, untreated 22/3, blockers 20/3, KCl 22/3.

Data are shown as mean ± SEM; $*p < 0.05$, $**p < 0.05$, $***p < 0.001$ shown compared to the

untreated condition determined by Kruskal-Wallis followed by Holm post hoc tests (j for

Amphiphysin, PIPK1γ and AP-180; k, p, and q), or by one-way ANOVA followed by a Tukey-

Kramer post hoc test (j for Dynamin-1). For quantification of confocal images, see Fig. 1 - figure

supplement 1; for a workflow of STED analyses, see Fig. 1 - figure supplement 2; for

assessment of AP-180 using and independent antibody, see Fig. 1 - figure supplement 3, for

additional analyses of en-face synapses, see Fig. 1 – figure supplement 4.

1218

1219

Figure 2. Deployment of endocytic proteins after chronic silencing or acute stimulation of

***Drosophila* NMJs**

(**a**) Schematic of the experiment at the *Drosophila* neuromuscular junction (NMJ). To silence neurons chronically, tetanus neurotoxin light chain ("TeNT") was expressed in Type 1b motor neurons on muscle 1 using the GAL4 UAS system. These neurons were compared to those from larvae expressing the M11b-GAL4 driver alone ("control"). To activate neurons, electrical stimulation was applied at 40 Hz for three minutes ("40 Hz") to Type 1b motor neurons on muscles 4 and 6/7, and comparisons were made to the contralateral, unstimulated side ("no stim").

(**b-g**) Example maximum intensity projections of ventral half of individual boutons with or without TeNT-expression (b), and quantification of the number of Brp objects per μm^2 of NMJ (c) and their integrated intensity (d). For Brp, Dynamin and Nervous Wreck, the average fluorescence intensity per bouton (e), the average intensity at the periactive zone mesh (f) and the polarization within periactive zone units (g) were quantified. Data in d, e and g are normalized to the average of the control condition; n in c, d (NMJs/larvae): control 28/5, TeNT 27/6; e: control 28/5, TeNT 28/6; f, g: control 28/5, TeNT 22/6.

(**h-m**) Same as b-g but comparing acutely stimulated ("40 Hz") and unstimulated terminals ("no stim") boutons; n in I, j: no stim 26/14, 40 Hz 24/14; k: no stim 26/14, 40 Hz 25/14; l, m: no stim 26/14, 40 Hz 23/14.

Data are mean ± SEM; *p < 0.05 determined by two-sided Student's t-tests (e for Nervous Wreck; g for Brp and Nervous Wreck; i; j; k; m) or two-sided Mann-Whitney U tests (c; d; e for Brp and Dynamin; f; g for Dynamin; l).

Figure 3. Deployment of endocytic proteins after Ca$_V$2 ablation in mouse hippocampal

neurons

(**a**) Schematics of the *Cacna1a*, *Cacna1b* and *Cacna1e* mutant alleles that constitute the conditional Ca$_V$2 triple knockout mouse line as described in (Held et al., 2020).

(**b-i**) Example side-view synapses and average line profiles of Amphiphysin and Bassoon (b+c), PIPK1γ (d+e), AP-180 (f+g), and Dynamin-1 and Munc13-1 (h+i). Neurons were stained for a protein of interest (Amphiphysin, PIPK1γ, AP-180 or Dynamin-1; imaged in STED), an active zone marker (Bassoon or Munc13-1; imaged in STED), and Synaptophysin (imaged in confocal). An area of interest was positioned perpendicular to the center of the active zone marker, and synapses were aligned via the peak fluorescence of the active zone marker in the average profiles (c, e, g, i). Line profiles plots were normalized to the average signal in the control^{Cav2} condition. Turquoise dotted lines in c, e, g and i mark average levels in the control^{Cav2} condition and grey bars represent the active zone and periactive zone area; n in b, c (synapses/cultures): control^{Cav2} 58/3, cTKO^{Cav2} 50/3; d, e: control^{Cav2} 50/3, cTKO^{Cav2} 49/3; f, g: control^{Cav2} 50/3, cTKO^{Cav2} 47/3; h, i: control^{Cav2} 48/3, cTKO^{Cav2} 50/3.

(**j, k**) Quantification of the peak-to-peak distance of the active zone marker and the protein of interest (j), and of the peak intensity in the periactive zone area (k). The periactive zone area is defined as an area within 68 nm on each side of the peak of the active zone marker (grey shaded areas in c, e, g and i); n as in b-i.

(**l-q**) Example en-face synapses (l-o) and quantification of the number of objects (p) and distance of these objects to the center of the active zone marker (q) of the experiment shown in b-l, n in p, q: Amphiphysin, control^{Cav2} 23/3, cTKO^{Cav2} 24/3; PIPK1γ, control^{Cav2} 22/3, cTKO^{Cav2} 25/3; AP-180, control^{Cav2} 19/3, cTKO^{Cav2} 20/3; Dynamin-1, control^{Cav2} 23/3, cTKO^{Cav2} 20/3.

Data are mean ± SEM; * $p < 0.5$ determined by two-sided Student's t-tests (j for Dynamin-1; k for Dynamin-1; p for PIPK1γ and Dynamin; q for Amphiphysin, Ap-180 and Dynamin-1) or two-sided Mann-Whitney U tests (j for Amphiphysin, PIPK1γ, AP-180; k for Amphiphysin, Bassoon, PIPK1γ, AP-180 and Munc13-1; p for Amphiphysin and AP-180; q for PIPK1γ). For

1271   quantification of confocal signals, see Fig. 3 - figure supplement 1; for assessment of AP-180

1272   using and independent antibody, see Fig. 3 - figure supplement 2; for additional assessment of

1273   en-face synapses, see Fig. 3 – figure supplement 3.

1274

1276 **Figure 4. Recruitment of endocytic proteins to the periactive zone after active zone**

disruption in mouse hippocampal neurons

(**a**) Schematics of the *Rims1*, *Rims2*, *Erc1* and *Erc2* mutant alleles that constitute the conditional RIM+ELKS quadruple knockout mouse line as described in (Tan et al., 2022; Wang et al., 2016).

(**b-i**) Example side-view synapses and average line profiles of Amphiphysin and PSD-95 (b+c), PIPK1γ (d+e), AP-180 (f+g), and Dynamin-1 (h+i). Neurons were stained for a protein of interest (Amphiphysin, PIPK1γ, AP-180 or Dynamin-1; imaged in STED), a postsynaptic marker (PSD-95; imaged in STED), and a synaptic vesicle marker (Synaptophysin or Synapsin; imaged in confocal). An area of interest was positioned perpendicular to the center of the PSD-95 object, and synapses were aligned via the peak fluorescence of PSD-95 in the average line profiles (c, e, g, i). Line profile plots were normalized to the average signal in the control^{R+E} condition. Orange dotted lines in c, e, g and i mark average levels in the control^{R+E} condition and grey bars represent the active zone and periactive zone area; n in b, c (synapses/cultures): control^{R+E} 55/3, cQKO^{R+E} 53/3; d, e: control^{R+E} 55/3, cQKO^{R+E} 54/3; f, g: control^{R+E} 53/3, cQKO^{R+E} 54/3; h, i: control^{R+E} 54/3, cQKO^{R+E} 53/3.

(**j-l**) Quantification of the peak-to-peak distance of PSD-95 and the protein of interest (j), peak intensity in the periactive zone area (k), and peak intensity of PSD-95 (l). The periactive zone area is defined as an area between -136 nm and the peak of PSD-95 (grey shaded areas in c, e, g and i); n as in b-i.

(**m-r**) Example en-face synapses (m-p) and quantification of the number of objects (q) and distance of these objects to the center of the PSD-95 object (r) of the experiment shown in b-I, n in q, r: Amphiphysin, control^{R+E} 22/3, cQKO^{R+E} 23/3; PIPK1γ control^{R+E} 15/3, cQKO^{R+E} 18/3; AP-180 control^{R+E} 20/3, cQKO^{R+E} 19/3; Dynamin-1 control^{R+E} 17/3, cQKO^{R+E} 16/3.

Data are mean ± SEM; ***p < 0.001 as determined by two-sided Student's t-tests (l; q for Amphiphysin, AP-180 and Dynamin-1; r for PIPK1γ and Dynamin-1) or two-sided Mann-Whitney U tests (j; k; q for PIPK1γ; r for Amphiphysin and, AP-180). For quantification of confocal

1303   signals, see Fig. 4 - figure supplement 1; for assessment of AP-180 using and independent

1304   antibody, see Fig. 4- figure supplement 2; for additional assessment of en-face synapses, see

1305   Fig. 4 – figure supplement 3.

1306

Figure 5. Recruitment of endocytic proteins to the periactive zone after disrupting active zone assembly at *Drosophila* NMJs

(a) Schematic of the *brp69* and *brpDef* alleles; dashed lines indicate the extent of deletions.

(**b-e**) Example maximum intensity projections of individual boutons of muscle 4 Type 1b

terminals from control *white* and *brp^{69}/brp^{Def}* larvae (b) and quantification of the average

fluorescence intensity per bouton (c), average intensity at the periactive zone mesh (d) and

polarization within periactive zone units (e) for Dynamin and Nervous Wreck. Data in c and d

are normalized to the average control condition; n in c (NMJs/larvae): *brp* control 31/3, *brp$^{69/Def}$*

26/3; d+e, *brp* control 31/3, *brp$^{69/Def}$* 24/3.

Data are mean ± SEM; statistical significance assessed using two-sided Student's t-tests (c for

Nervous Wreck; d for Nervous Wreck; e for Nervous Wreck) or two-sided Mann-Whitney U tests

(c for Dynamin; d for Dynamin; e for Dynamin).

Figure 6. Deployment of endocytic proteins after Liprin-α ablation in mouse hippocampal

neurons

(**a**) Schematics of the *Ppfia1*, *Ppfia2*, *Ppfia3* and *Ppfia4* mutant alleles that constitute the conditional Liprin-α quadruple knockout mouse line as described in (Emperador-Melero et al., 2024).

(**b-I**) Example side-view synapses and average line profiles of Amphiphysin and PSD-95 (b+c), PIPK1γ (d+e), AP-180 (f+g), and Dynamin-1 (h+i). Neurons were stained for a protein of interest (Amphiphysin, PIPK1γ, AP-180 or Dynamin-1; imaged in STED), a postsynaptic marker (PSD-95; imaged in STED), and a synaptic vesicle marker (Synaptophysin or Synapsin; imaged in confocal). An area of interest was positioned perpendicular to the center of the PSD-95 object, and synapses were aligned via the peak fluorescence of PSD-95 in the average profiles (c, e, g, i). Line profile plots were normalized to the average signal in the control[L1-L4] condition. Orange dotted lines in c, e, g and i mark average levels in the control[L1-L4] condition and grey bars represent the active zone and periactive zone area; n in b (synapses/cultures), c: control[L1-4] 54/3, cQKO[L1-4] 64/3; d, e: control[L1-4] 55/3, cQKO[L1-4] 50/3; f, g: control[L1-4] 59/3, cQKO[L1-4] 59/3; h, i: control[L1-4] 45/3, cQKO[L1-4] 46/3.

(**j-I**) Quantification of the peak-to-peak distance of PSD-95 and the protein of interest (j), peak intensity in the periactive zone area (k), and peak intensity of PSD-95 (l). The periactive zone area is defined as an area between -136 nm and the peak of PSD-95 (grey shaded areas in c, e, g and i); n as in b-i.

(**m-r**) Example en-face synapses (m-p) and quantification of the number of objects (q) and distance of these objects to the center of the PSD-95 object (r) of the experiment shown in b-I, n in q, r: Amphiphysin, control[L1-4] 20/3, cQKO[L1-4] 18/3; PIPK1γ control[L1-4] 18/3, cQKO[L1-4] 16/3; AP-180 control[L1-4] 20/3, cQKO[L1-4] 15/3; Dynamin-1 control[L1-4] 19/3, cQKO[L1-4] 21/3.

Data are mean ± SEM; $*p < 0.05$, $***p < 0.001$ determined by two-sided Student's t-tests (j for PIPK1γ; k for Amphiphysin; q for Amphiphysin, AP-180 and Dynamin-1, r for Amphiphysin, AP-180 and Dynamin-1) or two-sided Mann-Whitney U tests (j for Amphiphysin, AP-180 and

1351 Dynamin-1; k for PIPK1γ, AP-180 and Dynamin-1; l; q for PIPK1γ; r for PIPK1γ). For

1352 quantification of confocal signals, see Fig. 6- figure supplement 1; for additional assessment of

1353 en-face synapses, see Fig. 6 – figure supplement 2.

1354

1355

1356

a

Liprin-α

Liprin-α^{R60}

Liprin-α^{F3-ex15}

1 kb

b

control   *Liprin-α^{R60/F3-ex15}*

Brp

Dynamin

Nervous Wreck

merged

1μm

c

Brp objects per μm²

***

d

Brp integreted intensity (% control)

***

control   *Liprin-α^{R60/F3-ex15}*

e

Brp   Dynamin   Nervous Wreck

levels at bouton (% control)

***

f

levels at the peracitive zone mesh (% control)

n/a

g

polarization (log$_2$)

**   **

control   *Liprin-α^{R60/F3-ex15}*   control   *Liprin-α^{R60/F3-ex15}*   control   *Liprin-α^{R60/F3-ex15}*

1357

Figure 7. Endocytic proteins are recruited to periactive zones of *Drosophila* NMJs in

Liprin-α mutants

(**a**) Schematic of the _Liprin-α_^{R60} and _Liprin-α_$^{F3-ex15}$ alleles; dashed lines indicate the extent of deletions.

(**b-g**) Example maximum intensity projections of muscle 4 Type 1b terminals from _Liprin-α_$^{R60/F3-ex15}$ mutant and _white_ control larvae (b), and quantification of the number of Brp objects per μm^2 of NMJ (c) and of their integrated intensity (d). For Brp, Dynamin and Nervous Wreck, the average fluorescence intensity per bouton (e), the average intensity at the periactive zone mesh (f) and the polarization within periactive zone units (g) were quantified. Data in d, e and f are normalized to the average of the control condition; n in c-g (NMJs/larvae): control 27/5, _Liprin-α_$^{R60/F3-ex15}$ 26/5.

Data are mean ± SEM; * $p<0.05$ ** $p<0.01$ *** $p<0.001$ determined two-sided Student's t-tests (e for Dynamin and Nervous Wreck; f, g for Dynamin) or two-sided Mann-Whitney U tests (c, d, e for Brp; g for Brp and Nervous Wreck).

Figure 8. Recruitment of endocytic proteins at *Drosophila* NMJs of *rab3* mutants

(**a**) Schematic of the *rab3*^{rup} allele; red triangle indicates site of 5 bp deletion and frameshift that eliminates the final 35 amino acids (highlighted in red).

(**b-g**) Example maximum intensity projections of muscle 4 Type 1b terminals from *rab3*^{rup} mutants and *white* larvae controls (b), and quantification of the number of Brp objects per μm^2 of NMJ (c) and of their integrated intensity (d). For Brp, Dynamin and Nervous Wreck, the average fluorescence intensity per bouton (e), the average intensity at the periactive zone mesh (f) and the polarization within periactive zone units (g) were quantified. Data in d, e and f are normalized to the average of the control condition; n in c, d (NMJs/larvae): control 22/5, *rab3*^{rup} 28/6; e: control 25/5, *rab3*^{rup} 30/6; f, g: control 23/5, *rab3*^{rup} 23/6.

(**h, i**) Quantification of the levels of Dynamin (k) and of its polarization (l); n in k, l: control Brp+, 23/5, control Brp- 21/6, *rab3*^{rup} Brp+ 23/6, *rab3*^{rup} Brp- 28/6;

(**j, k**) Same as h,i, but for Nervous Wreck; n as in h, i.

(**l-n**) Example maximum intensity projections (l) of muscle 4 Type 1b terminals co-stained for the active-zone marker Brp, the postsynaptic density marker Pak, and the periactive zone marker Fasciclin-II (including the periactive zone segmentation, bottom image), and quantification of the percentage of individual periactive zone segments that contain Brp (m) or are apposed to Pak (n); n control 11/3, *rab3*^{rup} 9/3.

Data are mean ± SEM; * p<0.05 ** p<0.01 *** p<0.001 determined by two-sided Mann-Whitney U tests (c, d, e, f, g, m, n) or by Kruskal-Wallis tests followed by Holm post hoc tests (h-k). In h-k, data are compared to the control Brp+ condition.

Figure supplements

Figure 1 - figure supplement 1. Confocal microscopic analyses of synapses after chronic silencing or acute depolarization of mouse hippocampal neurons

(**a, b**) Example confocal images (a) and quantification (b) of the average intensities of Amphiphysin, PIPK1γ, AP-180, Dynamin-1, Bassoon and Munc13-1 at synapses identified as Synaptophysin puncta. Intensities are normalized to the average signals in the untreated conditions per culture; n in b (images/cultures): Amphiphysin, 14/3; PIPK1γ, untreated 14/3, blockers 14/3, KCl 13/3; AP-180, 15/3; Dynamin-1, 14/3; Synaptophysin, untreated 57/3, blockers 57/3, KCl 56/3; Bassoon, untreated 43/3, blockers 43/3, KCl 42/3; Munc13-1, 14/3. The increase in Munc13-1 upon chronic silencing, which we previously reported (Held et al., 2020), and of Synaptophysin, may reflect a homeostatic adaptation. The decrease in Amphiphysin

1410   upon KCl stimulation may reflect a redistribution of this protein during prolonged stimulation.

1411   Data are mean ± SEM; *$p < 0.05$, **$p < 0.05$, ***$p < 0.001$ compared to the untreated condition

1412   determined by one-way ANOVA followed by a Tukey-Kramer post hoc tests for Bassoon or

1413   Kruskal-Wallis followed by a Holm post hoc tests for Amphiphysin, PIPK1γ, AP-180, Dynamin-1,

1414   Synaptophysin and Munc13-1.

1415

analyses of side-view synapses in mouse neurons

a Synaptophysin Bassoon

synapse selection and line scan position

b Synaptophysin Bassoon Amphiphysin

protein of interest channel activation

c Amphiphysin

generation of line profile

d average profiles

e peak intensities and distance of proteins of interest to the marker

analyses of en-face synapses in mouse neurons

f Synaptophysin Bassoon

synapse selection

g Synaptophysin Bassoon Amphiphysin

protein of interest channel activation

h Bassoon Amphiphysin

thresholding and detection of objects

i number of objects per synapse, their lateral distance to the active zone, and their integrated intensity

analyses of boutons in Drosophila neuromuscular junctions

j Nervous Wreck Brp

generation of half maximum intensity images for selected boutons

k quantification of average intensities

l segmentation

m filtering and measurement of intensities in core and mesh regions

n levels at the endocytic mesh and ratio calculation

Figure 1 - figure supplement 2. Workflows for STED analyses in mouse hippocampal neurons and for confocal analyses at _Drosophila_ neuromuscular junctions

(**a-e**) Workflow for the analyses of side-view synapses of mouse hippocampal neurons, showing an example synapse immunostained for the active zone marker Bassoon (imaged in STED), Amphiphysin (imaged in STED) and Synaptophysin (imaged in confocal). Synapse selection and placement of an area of interest (white transparent rectangle with line profile direction indicated by black arrow) perpendicular to the marker (Bassoon, in this example) is done by an experimenter blind for the protein of interest (Amphiphysin, in this example). Next, the protein of

interest channel is activated (b), and the profile is generated (c). Then, the average from many

synapses is calculated and plotted aligned to the peak of the marker protein (d). The distance of

the peak to the peak of the marker protein and the maximum intensity within the region

corresponding to the periactive zone are also calculated and plotted (e). The example

corresponds to the untreated condition in Fig. 1b+c.

(**f-i**) Workflow for the analyses of en-face synapses of mouse hippocampal neurons, showing an

example synapse immunostained for Bassoon (imaged in STED), Amphiphysin (imaged in

STED) and Synaptophysin (imaged in confocal). First, synapse selection is done by an

experimenter blind for the protein of interest (f). Next, the channel of the protein of interest is

activated (g), and objects containing endocytic proteins and the marker are identified in the

respective channels using an algorithm (h). Finally, the number, integrated intensites, and the

distances of the objects to the center of the marker are quantified (i). The example corresponds

to the untreated condition in Fig. 1l.

(**j-n**) Workflow for analyses of *Drosophila* neuromuscular junctions. Terminals are analyzed both

in 3D (k) and in 2D half-maximum intensity projections (j, l-n). The average intensities of the

active zone marker (Brp in this example) and the endocytic protein (Nervous Wreck in this

example) are quantified in the full 3D volume of the terminal (k). Next, the periactive zone levels

and degree of polarization are analyzed in 2D half-maximum intensity projections. Polarization

of each protein is quantified as the ratio between its average intensity at the mesh over its

average intensity in the core. To conduct this analysis, segmentation into mesh and core is

performed based on the difference in signal between proteins enriched in the periactive zone

mesh (e.g. Nwk and Dynamin) vs. proteins enriched in the core region (e.g. Brp and Pak) as

described in the methods. The resulting ROIs are used to measure average intensities within

the mesh and the core. Levels are plotted as normalized intensities in the mesh, and the

mesh:core ratios are calculated to determine polarization (n). Ratios greater than zero indicate

enrichment of the protein at the periactive zone, while those less than zero indicate de-

enrichment.

1453

Figure 1 - figure supplement 3. Assessment of AP-180 with alternate antibody after

chronic silencing or acute depolarization of mouse hippocampal neurons

(**a, b**) Example side-view synapses (a) and average line profiles of AP-180 and Munc13-1 (b).

Neurons were stained for AP-180 (imaged in STED), Munc13-1 (imaged in STED), and the

synaptic vesicle marker Synaptophysin (imaged in confocal). An area of interest was positioned

perpendicular to the center of the Munc13-1 object, and synapses were aligned via the peak

fluorescence of Munc13-1 in the average profiles. Line profiles were normalized to the average

signal in the untreated condition. Grey dotted lines mark average levels in the untreated

condition and grey bars represent the active zone area; n in b (synapses/cultures): untreated,

58/3; blockers, 61/3; KCl, 50/3.

(c, d) Quantification of the peak-to-peak distance of the active zone marker and the protein of

interest (c), and of the peak levels in the periactive zone area (d). The periactive zone area is

defined as an area within 68 nm on each side of the peak of the active zone marker (grey

shaded areas in b)(Held et al., 2020), n as in b.

Data are mean ± SEM; *p < 0.05, ***p < 0.001 compared to the untreated condition determined

by Kruskal-Wallis followed by Holm post-hoc tests.

1470

1471

Figure 1 - figure supplement 4. Additional analyses of en-face synapses after chronic

silencing or acute depolarization of mouse hippocampal neurons

Quantification of the average integrated intensities (calculated as the object area multiplied by

its average fluorescence intensity) of the Amphiphysin, PIPK1γ, AP-180 and Dynamin-1 objects

detected in en-face synapses from Fig. 1l-q. Intensities are normalized to the average signals in

the untreated conditions per culture; n as in Fig. 1p.

Data are mean ± SEM; **p < 0.05, ***p < 0.001 compared to the untreated condition determined

by one-way ANOVA followed by a Tukey-Kramer post hoc test for AP-180 or Kruskal-Wallis

followed by Holm post hoc tests for Amphiphysin, PIPK1γ and Dynamin-1.

1481

Figure 3 - figure supplement 1. Confocal microscopic analyses of synapses after Ca$_V$2 ablation in mouse hippocampal neurons

(**a, b**) Example confocal images (a) and quantification (b) of the average intensities of Amphiphysin, PIPK1γ, AP-180 and Dynamin-1 at synapses identified as Synaptophysin puncta. Intensities are normalized to the average signals in the control[Cav2] condition per culture; n in b (images/cultures): Amphiphysin, 20/3; PIPK1γ, 20/3; AP-180, 17/3; Dynamin-1, 18/3; Bassoon, 57/3; Munc13-1, 18/3; Synaptophysin, 75/3.

Data are mean ± SEM; *$p < 0.05$ determined by a two-sided Student's t-tests for Amphiphysin, PIPK1γ, Bassoon and Munc13-1, or two-sided Mann-Whitney U tests for AP-180, Dynamin-1 and Synaptophysin.

1495

Figure 3 - figure supplement 2. Assessment of AP-180 with an alternate antibody after

Ca_V2 ablation in mouse hippocampal neurons

(**a, b**) Example side-view synapses (a) and average line profiles (b) of AP-180 and Munc13-1.

Neurons were stained for AP-180 (imaged in STED), Munc13-1 (imaged in STED), and the

synaptic vesicle marker Synaptophysin (imaged in confocal). An area of interest was positioned

perpendicular to the center of the Munc13-1 object, and synapses were aligned via the peak

fluorescence of Munc13-1 in the average profiles. Line profiles were normalized to the average

signal in control[Cav2] condition. Green dotted lines mark average levels in the control[Cav2] and grey

bars represent the active zone area; n in b (synapses/cultures): control[Cav2], 50/3; cTKO[Cav2],

48/3.

(**c, d**) Quantification of the peak-to-peak distance of Munc13-1 and AP-180 (c) and of their peak

levels in the periactive zone area (d). The periactive zone area is defined as an area within 68

nm on each side of the peak of the active zone marker (grey shaded areas in b); n as in b.

Data are mean ± SEM; *p < 0.05, shown compared to the control[Cav2] determined by a two-sided

Student's t-test (d for AP-180) or two-sided Mann-Whitney U tests (c; d for Munc13-1).

1511

1512

1513

Figure 3 - figure supplement 3. Additional analyses of en-face synapses after Ca$_V$2

ablation in mouse hippocampal neurons

Quantification of the average integrated intensities (calculated as the object area multiplied by

its average fluorescence intensity) of the Amphiphysin, PIPK1γ, AP-180 and Dynamin-1 objects

detected in the en-face synapses quantified in Fig. 3l-q. Intensities are normalized to the

average signals in control^{Cav2} per culture; n as in Fig. 3p.

Data are mean ± SEM; *p < 0.05 determined by Mann-Whitney U tests.

1521

1522

Figure 4 - figure supplement 1. Additional analyses of endocytic proteins after active

zone disruption in mouse hippocampal neurons

(**a, b**) Example confocal images (a) and quantification (b) of the average intensities of

Amphiphysin, PIPK1γ, AP-180 and Dynamin-1 at synapses identified as Synaptophysin or

Synapsin puncta. Intensities are normalized to the average signals in control[R+E] per culture; n in

b (images/cultures): Amphiphysin, control[R+E] 15/3, cQKO[R+E] 16/3; PIPK1γ, 16/3; AP-180, 16/3;

Dynamin-1, control[R+E] 18/3, cQKO[R+E] 19/3; PSD-95, control[R+E] 65/3, cQKO[R+E] 67/3;

Synaptophysin, control[R+E] 47/3, cQKO[R+E] 48/3; Synapsin, control[R+E] 18/3, cQKO[R+E] 19/3.

Data are mean ± SEM; *p < 0.05, ***p < 0.001 determined by two-sided Student's t tests for

Dynamin-1 and Synapsin or by two-sided Mann-Whitney U tests for Amphiphysin, PIPK1γ, AP-

180, PSD-95 and Synaptophysin.

1534

1535

 Figure 4 - figure supplement 2. Assessment of AP-180 with an alternate antibody after

active zone disruption in mouse hippocampal neurons

(**a, b**) Example side-view synapses (a) and average line profiles (b) of AP-180 and PSD-95.

Neurons were stained for AP-180 (imaged in STED), PSD-95 (imaged in STED), and the

synaptic vesicle marker Synaptophysin (imaged in confocal). A line profile was positioned

perpendicular to the center of the PSD-95 object, and synapses were aligned via the peak

fluorescence of PSD-95 in the average profiles. Line profiles were normalized to the average

signal in control[R+E]. Green dotted lines mark average levels in the control[R+E] condition and grey

bars represent the active zone area, n in b (synapses/cultures): control[R+E], 52/3 cultures;

cQKO[R+E], 46/3.

(c, d) Quantification of the peak-to-peak distance of PSD-95 and AP-180 (c) and of their peak

levels in the periactive zone area (d). The periactive zone area is defined as the area -136 nm

from the PSD-95 peak towards the presynaptic bouton (grey shaded areas in b); n as in b.

Data are mean ± SEM; *$p < 0.05$, shown compared to the control[Cav2] condition determined by a

two-sided Student's t-test (d for AP-180) or two-sided Mann-Whitney U tests (c; d for Munc13-

1).

1552

1553

Figure 4 - figure supplement 3. Additional analyses of en-face synapses after active zone disruption in mouse hippocampal neurons

Quantification of the average integrated intensities (calculated as the object area multiplied by its average fluorescence intensity) of the Amphiphysin, PIPK1γ, AP-180 and Dynamin-1 objects detected in the en-face synapses quantified in Fig. 4m-r. Intensities are normalized to the average signal in control[R+E] per culture, n as in Fig. 4q.

Data are mean ± SEM; **$p < 0.01$ ***$p < 0.001$ determined by two-sided Mann-Whitney U tests.

1561

1562

Figure 6 - figure supplement 1. Confocal microscopic analyses of synapses after Liprin-α

ablation in mouse hippocampal neurons

(**a, b**) Example confocal images (a) and quantification (b) of the average intensities of

Amphiphysin, PIPK1γ, AP-180 and Dynamin-1 at synapses identified as Synaptophysin or

Synapsin puncta. Intensities are normalized to the average signal in control[L1-4] per culture; n in b

(images/cultures): Amphiphysin 26/3 ; PIPK1γ, 16/3; AP-180, 20/3; Dynamin-1, 18/3; PSD-95,

80/3; Synaptophysin, 62/3, Synapsin, 18/3.

Data are mean ± SEM; *p < 0.05 determined by two-sided Student's t-tests for Synapsin and

Dynamin-1 or two-sided Mann-Whitney U tests for Amphiphysin, PIPK1γ, AP-180, PSD-95 and

Synaptophysin.

 Figure 6 - figure supplement 2. Additional analyses of en-face synapses after Liprin-α

 ablation in mouse hippocampal neurons

 (c) Quantification of the average integrated intensities (calculated as the object area multiplied

 by its average fluorescence intensity) of the Amphiphysin, PIPK1γ, AP-180 and Dynamin-1

 objects detected in the en-face synapses quantified in Fig. 6m-r. Intensities are normalized to

 the average signals in control[L1-4] per culture; n as in Fig. 6q.

 Data are mean ± SEM; statistical significance was assessed by two-sided Mann-Whitney U

 tests.