A GWAS–machine learning framework reveals protein-synthesis pathway signals for yield in *Theobroma cacao* after population-structure correction

Insuck Baek[1,†], Jishnu Bhatt[2,†], Seunghyun Lim[2], Dongho Lee[3], Jae Hee Jang[2], Stephen P. Cohen[2], Amelia H. Lovelace[2], Moon S. Kim[1], Lyndel W. Meinhardt[2], Sunchung Park[2], Ezekiel Ahn[2,*]

[1] Environmental Microbial and Food Safety Laboratory, Agricultural Research Service, United States, Department of Agriculture, Beltsville, MD, 20705, USA

[2] Sustainable Perennial Crops Laboratory, Agricultural Research Service, United States, Department of Agriculture, Beltsville, MD, 20705, USA

[3] Soybean Genomics & Improvement Laboratory, Agricultural Research Service, United States, Department of Agriculture, Beltsville, MD, 20705, USA

[†] The authors contributed equally to this work.
* Author to whom correspondence should be addressed (email: Ezekiel.ahn@usda.gov).

Abstract

Improving cacao yield, a key objective in post-domestication crop improvement, remains a primary goal for breeders, but progress is often hindered by the confounding effects of population structure. To overcome this, we analyzed 346 diverse cacao accessions using an ML-based association mapping framework (with and without population structure adjustment) and a phenotype-only ML prediction of yield. By correcting for population structure, our Bootstrap Forest-based GWAS revealed association signals that showed consistent enrichment for ribosome and protein-synthesis functions, and a recurrent subset of SNPs with high importance appeared across multiple yield components, including pod index and seed number. In parallel, a Neural Network model was utilized to identify cotyledon mass and length as the most powerful predictors for total wet bean mass ($R^2 = 0.715$ by repeated five-fold cross-validation), suggesting a practical, low-cost screening proxy for breeding). Collectively, this study delivers a robust genetic framework and a novel predictive tool to accelerate the development of high-yielding cacao varieties through the early identification of elite clones.

Keywords: *Theobroma cacao*, Yield, GWAS, Machine learning, Protein synthesis, Genomic prediction, Population structure, Plant breeding, ribosome

Introduction

Improving the genetic potential of high-value horticultural crops is a central goal of modern agricultural science. *Theobroma cacao* L., the source of cocoa beans, is a globally significant

crop that underpins a multi-billion-dollar chocolate industry and supports the livelihoods of
millions of smallholder farmers[1,2].These smallholders are responsible for approximately 80%
of global cocoa production[3]. Originating in the upper Amazon basin[4], cacao exhibits rich
genetic diversity, contributing to a wide range of flavors, aromas, and other desirable traits[5].
However, cacao production faces increasing challenges from emerging diseases and pests,
and a growing global demand for chocolate[6,7]. These challenges necessitate the development
of improved cacao varieties with enhanced yield potential, disease resistance, and
environmental resilience, while maintaining and enhancing quality attributes. This continuous
drive for crop improvement, from early domestication to modern breeding, requires a deep
understanding of the genetic basis of key agronomic traits.

Efforts to understand genetic diversity and identify genes controlling important traits from the
cacao genome have been ongoing for decades. Studies have indicated that genetic
differentiation within cacao lineages results from spatial variation in selection pressures,
genetic drift, or a combination of both processes[8]. Yield stability over time is a critical trait in
cacao breeding[3], yet studies focusing on temporal stability remain limited[9]. A study
evaluating 34 cacao hybrid families over six years in Costa Rica revealed substantial
variability in annual yield, with coefficient of variation values ranging from 113% to 642%
for different years, particularly notable in years two and three, when 80% of trees produced
no pods[9]. Cacao yield was contingent on interrelated factors such as pod health and disease
incidence, shown through a positive correlation between the number of healthy pods and
overall yield efficiency[3]. When low yield variability is optimized for in breeding programs,
newly developed cultivars are susceptible to reduced yield efficiencies, necessitating broadly
informed selection of genetic traits to sustain productivity over time[3]. Genetic factors
significantly influence pod development and quality traits[10]. The selection of progeny based
on pod index, calculated as the number of pods required to produce one kilogram of dried
cocoa[11], and weight can improve overall tree productivity and efficiency in production
processes[10]. Monitoring genetic identity is essential as progenies from mislabeled or pollen-
contaminated clones confound breeding efforts towards the development of superior
varieties[12].

Traditional breeding methods, relying on phenotypic selection and controlled crosses, have
contributed to significant improvements in cacao varieties[13,14]. However, these methods are
unable to take into account the complex interplay and inheritance of many traits[15]. The advent
of genomic tools, such as genome-wide association studies (GWAS), offers new
opportunities to understand inherited trait complexity and accelerate cacao improvement[16,17].
Previous studies in cacao have employed GWAS to identify loci associated with disease
resistance, yield components, and bean quality attributes[11,16,17]. Despite the use of standard
statistical corrections, a critical limitation of these studies is that the strong population
structure in cacao often results in associations with general 'vigor' rather than specific,
actionable genes. Furthermore, translating these genetic findings into practical tools for
breeders remains a significant hurdle. Consequently, a comprehensive understanding of the
genetic architecture of complex traits in cacao, such as those related to yield potential, has
remained elusive. Only a few studies have integrated advanced machine learning techniques

to enhance the prediction and understanding of complex traits in cacao, primarily using inputs
such as image analysis for bean quality and canopy architecture[18,19], sensor data for
fermentation level[20], or agronomic and climate data for yield potential[21]. However, studies
that directly combine machine learning frameworks with high-throughput genomic data for
trait discovery and prediction remain rare.

This study addresses these limitations by leveraging a large and diverse collection of 346
cacao accessions, utilizing publicly available phenotypic and genotypic data from the ICGT
Cacao Germplasm Database[11]. These accessions, conserved *ex-situ* at the International Cocoa
Genebank, Trinidad (ICGT), represent a broad range of genetic and phenotypic variation[11].
We employed a multifaceted approach, combining detailed phenotypic evaluations of 27
traits, encompassing flower, fruit, and seed characteristics, with genome-wide SNP
genotyping, phylogenetic analysis, and advanced statistical methods. Specifically, as a key
innovation of this study, we utilized a dual-model approach, performing the GWAS both with
and without correction for population structure, to robustly identify significant marker-trait
associations and explored the relationships among traits using hierarchical clustering[22].
Furthermore, based on various phenotypic data, we trained a machine-learning model to
predict wet bean mass, a direct indicator of yield potential.

The primary objectives of this study were to: (1) investigate the relationship between genetic
relatedness and phenotypic variation in a diverse collection of cacao accessions; (2) identify
genomic regions and potential candidate genes associated with key traits using a Bootstrap
Forest-based GWAS approach; (3) explore the relationships among traits through hierarchical
clustering; and (4) develop and evaluate a machine learning model for predicting wet bean
mass. We tested several key hypotheses: first, that phenotypic variation in this diverse
collection would be complex and only partially explained by broad phylogenetic
relationships; second, that our GWAS and machine learning models would successfully
identify novel candidate genes and accurately predict yield potential, respectively; and
finally, that a corrected GWAS model would provide a more robust approach for
disentangling broad ancestral signals from loci with direct, trait-specific functions, which
would in turn resolve into core biological pathways upon Gene Ontology (GO) analysis.
114

Results
Correlation between patristic distances and phenotypic traits

To assess whether more distantly related accessions differ more in phenotype, we computed
Mantel correlations between patristic distances (from the phylogeny) and phenotypic distance
matrices across 344 traits out of 346 accessions with both genotypes and phenotypes
(intersection n= 342). Correlations were generally small (median $r \approx 0.10$). After false
discovery rate (FDR) correction (q< 0.05), 12 traits were significant, led by pod index ($r=$
0.25, q= 0.0019) and cotyledon width ($r=$ 0.20, q= 0.0019) (Fig. S1). The Mantel correlations
were generally small (median $r=$ ~0.10). The remaining traits were not significant after
correction
125

Visualization of phenotypic and genotypic diversity using PCA

Principal Component Analysis (PCA) was used to visualize and compare the patterns of
variation within the phenotypic and genotypic datasets (Fig. 1).
The PCA plot based on the phenotypic data reveals a single, diffuse cloud of points,
indicating that the measured traits vary continuously across the cacao collection (Fig. 1a).
The first two principal components explained 14.8% and 10.2% of the total phenotypic
variation, respectively. This lack of distinct clustering suggests that no clear-cut "types" of
trees can be delineated based on their physical traits alone. In contrast, the PCA based on the
genotypic SNP data shows clear evidence of population structure (Fig. 1b). The accessions
are organized into several distinct clusters, demonstrating that the collection is composed of
different underlying ancestral groups. The first two principal components for the genotype
data explained 16% and 11.8% of the total genetic variation. This visual evidence of genetic
structuring compared to the continuous phenotypic variation powerfully justifies the necessity
of correcting for population structure in the subsequent GWAS to avoid spurious findings.
140

141
Fig. 1
Principal component analysis of phenotypic and genotypic diversity in the cacao collection.
Each point represents a single cacao accession. (a) PCA based on all measured phenotypic
traits. The axes show the first two principal components, which account for 14.8% and 10.2%
of the phenotypic variance, respectively. The plot reveals a continuous distribution with no
distinct clusters. (b) PCA based on SNP genotype data. The axes show the first two principal

components, which account for 16% and 11.8% of the genetic variance, respectively. The
plot reveals clear clustering, indicating significant population structure within the collection.
150

Genome-wide association study of key yield and morphological traits
To identify genomic regions associated with phenotypic variation, we employed a machine
learning-based GWAS approach using Bootstrap Forest models. To specifically assess and
control for the impact of the strong population structure within the collection, two parallel
analyses were conducted: a naive model and a structured association model. The naive model
utilized the raw phenotypic data to provide an exploratory view of all potential genetic
associations, including those potentially confounded with ancestry. In contrast, the structured
association model was performed on ancestry-corrected residuals to identify robust loci with
a direct influence on traits, independent of the accessions' genetic background. This dual
approach allows for a direct comparison to distinguish robust, trait-specific associations from
broader effects linked to the collection's underlying genetic structure. The findings presented
below are from the structured association model, which controls for the effects of population
structure to identify direct marker-trait associations. This is a critical step in identifying the
genetic targets of past and future selection for crop improvement.
165

166   The analysis revealed numerous significant SNP-trait associations. The top-ranking SNPs for
167   seven key traits related to yield, morphology, and quality are summarized in Table 1 and Fig.
168   2. A list of the most significant associations (Portion > 0.005) for all 27 traits is available in
169   Supplementary Data 1.
170

Genetic loci for yield and its core components
Key yield traits, including pod index, wet bean mass, and seed number, shared a common set
of associated genetic loci, highlighting a core genetic architecture for productivity. Notably,
the SNPs TcSNP597 (encoding a 21 kDa seed protein) and TcSNP281 (encoding a 60S
ribosomal protein L31) appeared as top associations for all three of the aforementioned key
yield traits. Furthermore, TcSNP1366 (encoding a Heparan-alpha-glucosaminide N-
acetyltransferase) was a top locus for both pod index and wet bean mass. These findings
strongly link cacao yield potential to the genetic machinery governing protein synthesis
(ribosomal proteins), energy metabolism (heparan-alpha-glucosaminide N-acetyltransferase),
and the production of essential seed storage proteins.
181

Associations for key predictors of yield and fruit morphology
To identify direct targets for crop improvement, GWAS also identified loci controlling
important fruit and seed characteristics that contribute to overall yield potential, such as
cotyledon and fruit size. For cotyledon mass, top SNPs included TcSNP139 (within the gene
for a Putative Cytochrome B5 isoform D) and TcSNP17 (within a gene for a RING/U-box
superfamily protein). For cotyledon length, the most significant association was TcSNP1102,
within the gene for a mitochondrial cytochrome c oxidase subunit. Together, these results
suggest that the size and mass of cotyledons are influenced by genes involved in cellular
respiration, energy metabolism, and protein regulation. Additionally, for fruit length, a key
component of pod size, the top SNP was TcSNP1011, located within a gene encoding an

aspartyl protease family protein. Plant proteases are known to be involved in tissue
remodeling, making this a strong candidate gene for influencing fruit development.
194
Internal validation via pigmentation
For fruit surface anthocyanin level, the most significant SNP was TcSNP174. This SNP is
located within the gene encoding 6-phosphogluconate dehydrogenase. This enzyme is a key
component of the pentose phosphate pathway, which is directly responsible for producing the
precursors required for flavonoid and anthocyanin biosynthesis, providing a direct link
between an association and a known biochemical pathway controlling pigmentation in cacao.
201

Trait	SNP ID	Chr	Putative Gene	Importance Score (Portion)
Pod index	TcSNP597	2	*Tc02v2_p004390* 21 kDa seed protein	0.0632
	TcSNP281	1	*Tc01v2_p034810* 60S ribosomal protein L31	0.0226
	TcSNP1366	1	*Tc01v2_p025790* Heparan-alpha-glucosaminide N-acetyltransferase	0.0197
Wet bean mass	TcSNP1366	1	*Tc01v2_p025790* Heparan-alpha-glucosaminide N-acetyltransferase	0.0442
	TcSNP597	2	*Tc02v2_p004390* 21 kDa seed protein	0.038
	TcSNP281	1	*Tc01v2_p034810* 60S ribosomal protein L31	0.0297
Seed number	TcSNP281	1	*Tc01v2_p034810* 60S ribosomal protein L31	0.0638
	TcSNP1370	1	*Tc01v2_p008280* Histone H3.3	0.0611
	TcSNP597	2	*Tc02v2_p004390* 21 kDa seed protein	0.0483
Cotyledon length	TcSNP1102	1	*Tc01v2_p006960* Cytochrome c oxidase subunit 6a, mitochondrial	0.0545
	TcSNP1366	1	*Tc01v2_p025790* Heparan-alpha-glucosaminide N-acetyltransferase	0.0535
	TcSNP281	1	*Tc01v2_p034810* 60S ribosomal protein L31	0.0475
Cotyledon mass	TcSNP139	8	*Tc08v2_p007790* Putative Cytochrome B5 isoform D	0.0221
	TcSNP17	2	*Tc02v2_p029690* RING/U-box superfamily protein, putative	0.0204
	TcSNP32	4	*Tc04v2_p022290* Pyruvate, phosphate dikinase, chloroplastic	0.0144
Fruit length	TcSNP1011	1	*Tc01v2_p028610* Aspartyl protease family protein 2	0.0386
	TcSNP1258	2	*Tc02v2_p004580* Probable aquaporin PIP2-8	0.0109
	TcSNP189	8	*Tc08v2_p000820* Succinate dehydrogenase subunit 5, mitochondrial	0.0108
Fruit surface anthocyanin	TcSNP174	4	*Tc04v2_p009180* 6-phosphogluconate dehydrogenase, decarboxylating 3	0.0244
	TcSNP401	4	*Tc04v2_p010060* ATP-dependent Clp protease proteolytic subunit 3, chloroplastic	0.0218
	TcSNP591	1	*Tc01v2_p009870* 60S ribosomal protein L4-1	0.0199

202
Table 1.

Putative candidate genes near top SNPs associated with key cacao traits from corrected
GWAS. Summary of the most significant SNP associations identified through the Bootstrap
Forest-based GWAS after correcting for population structure. The table lists the top-ranking
SNPs for seven key traits related to yield, fruit morphology, and color. For each SNP, the
table provides its chromosomal location (Chr), the nearest annotated putative gene, and the
importance score (Portion) from the model. The importance score reflects the contribution of
each SNP to the prediction of the trait in the Bootstrap Forest model. SNPs that could not be
mapped to the reference genome are excluded.
212

213
Fig. 2
Manhattan plots from the corrected GWAS for key cacao traits. Each panel displays the
results from the ancestry-adjusted Bootstrap Forest GWAS for a specific key trait: (a) Pod
index, (b) Wet bean mass, (c) Seed number, (d) Cotyledon length, (e) Cotyledon mass, (f)
Fruit length, and (g) Fruit surface anthocyanin. The x-axis shows the SNP position across the
10 cacao chromosomes, while the y-axis represents the importance score (Portion) from the
model, where higher values indicate a stronger association. Top-ranking SNPs, corresponding
to those in Table 1, are labeled with their respective IDs. SNPs with unmapped chromosomal
locations are not shown.
223

Genomic hotspots of trait association
To visualize the global distribution of the most significant marker-trait associations and
identify genomic regions potentially under selection for multiple traits, the top three SNPs for
each of the 27 traits were plotted according to their physical position on the cacao
chromosomes (Fig. S2). This approach revealed several genomic regions, or "hotspots," that
contained significant associations for multiple, diverse traits. A notable pattern emerged
where these hotspots were often concentrated near the boundaries of the chromosomes. This
was particularly evident on chromosomes 1, 2, 5, 9, and 10 which all showed significant
clusters of associations near their starting positions, while chromosomes 1, 2, 4, and 9 also
showed clusters near their ends.
234

Hierarchical clustering reveals functional trait groups

To understand how selection might act on suites of related traits, we conducted hierarchical
clustering on the ancestry-adjusted SNP importance scores derived from the Bootstrap Forest
GWAS. The resulting dendrogram visualizes the relationships between traits based on their
shared genetic architecture (Fig. 3).

A number of expected, functionally related traits were successfully grouped. For instance, the
four measured anthocyanin intensity traits (in the filament, pedicel, ligule, and mature fruit
ridges) were found in close proximity within the same parent cluster. Similarly, various fruit
and seed shape traits formed logical, though not always adjacent, groupings.

The analysis revealed a distinct cluster composed of key yield-related traits. Key productivity
metrics such as pod index and wet bean mass formed a tight pair, which in turn clustered with
another group containing seed number, ovule number, and unexpectedly cotyledon length.
Cotyledon width grouped with fruit characteristics and cotyledon length-to-width ratio
grouped with seed shape traits and fruit width. Cotyledon mass was placed on a distant
branch of the dendrogram, positioned in proximity to traits including pod wall hardness, fruit
length, and several fruit apex form traits. The grouping of these distinct but related traits

indicates a shared genetic network may govern overall yield potential in cacao.

Fig. 3
Hierarchical clustering of cacao traits based on corrected genetic association profiles.
The dendrogram illustrates the relationships among traits, generated using Ward's method on
the SNP importance scores (Portion) from the corrected Bootstrap Forest GWAS. This
analysis groups traits that share similar underlying genetic associations, revealing distinct
functional clusters such as those related to overall yield and pigmentation.

Comparison of gene ontology enrichment between naive and structured models
To understand the biological functions of the genes associated with key traits, we performed
a GO enrichment analysis. We used pod index as a case study to demonstrate the importance
of correcting for population structure in a GWAS.

The GO analysis of the top candidate genes from the naive (uncorrected) model revealed a
broad mix of plausible biological pathways (Fig. 4a). The results were enriched for terms

related to general energy metabolism (e.g., "respiratory chain complex," "photosynthesis light
harvesting"), protein management (e.g., "protein processing in endoplasmic reticulum"), and
stress response (e.g., "response to hydrogen peroxide," "response to heat"). This represents a
classic "vigor" signal, likely reflecting the broad, fundamental differences in overall health
and metabolism between the different ancestral groups in the cacao collection. Detailed
KEGG pathway diagrams for key findings from this uncorrected analysis, such as protein
processing in the endoplasmic reticulum and photosynthesis, are shown in Supplementary
Figs. S3 and S4, respectively.

In contrast, the GO analysis of the top genes from the structured association (corrected)
model tells a much more specific and statistically significant story (Fig. 4b). After correcting
for population structure, the broad metabolic and stress signals disappeared, and the result
became overwhelmingly dominated by a single theme: the ribosome (FDR = 1.0E-08).
Multiple enriched terms, such as "cytosolic ribosome," "ribosomal subunit," and
"translation," all point to the core machinery of protein synthesis. A detailed KEGG pathway
diagram showing the specific ribosomal components identified is provided in Fig. S5. This
direct comparison demonstrates the power of the statistical correction, moving the analysis
from a broad signal to a specific and robust biological insight that links pod index to the
plant's capacity for protein production.

Fig. 4
**Comparison of go enrichment analyses for uncorrected and corrected pod index GWAS
results.** The plots show significantly enriched GO terms for the top candidate genes
identified for the pod index trait. (a) GO enrichment plot for genes from the naive
(uncorrected) analysis. The analysis reveals a broad mix of pathways related to energy
metabolism, protein processing, and stress response. Detailed KEGG pathways for protein
processing in the endoplasmic reticulum and photosynthesis are shown in Figs. S2 and S4,
respectively. (b) GO enrichment plot for genes from the structured association (corrected)
analysis. After correcting for population structure, the result is a significant and specific
enrichment for the ribosome and protein synthesis pathways. A detailed KEGG pathway for
the ribosome is shown in Fig. S5. This comparison illustrates how the statistical correction
filtered out confounding signals to reveal the core machinery of protein synthesis as the
primary biological process associated with pod index.

Combined-trait analysis reveals core biological processes

To identify the fundamental pathways underpinning groups of related traits, we performed combined GO enrichment analyses on our corrected GWAS results. First, identify the fundamental biological pathways underpinning this group of traits, we combined the candidate genes from nine size- and yield-related traits (ovule number, fruit length, fruit width, seed number, wet bean mass, cotyledon length, cotyledon width, cotyledon mass, and pod index). The GO enrichment analysis revealed a highly significant and specific enrichment for the ribosome and protein synthesis (Fig. 5a). This result indicates that the primary genetic drivers of yield are linked to the plant's capacity for protein production.

To further validate our analytical process, we investigated the genetic basis of pigmentation across tissues. We performed a combined analysis of five color-related traits (flower anthocyanin intensity in the pedicel, ligule, and filament; fruit surface anthocyanin intensity; and seed cotyledon color). This analysis revealed a strong enrichment for pathways related to primary energy metabolism, including glycolysis and ATP generation (Fig. 5b), indicating that producing the anthocyanin pigments is an energetically expensive process. A key KEGG pathway identified in this analysis was "photosynthesis - antenna proteins," and a detailed diagram showing the specific components identified, such as *Lhcb4*, is provided in Fig. S6.

A deeper look at the top KEGG pathway results provided a more nuanced story (Fig. 6). For the "yield & size" group, numerous components of both the small and large ribosomal subunits were identified (Fig. 6a). This suggests that overall yield is linked to the general capacity and throughput of the entire protein factory. In contrast, for the "color & pigmentation" group, only a few, specific ribosomal proteins were identified, such as L4e, L7e, and L24e (Fig. 6b). This suggests that the genetic control of pigmentation may be linked to these specific ribosomal proteins having specialized regulatory roles, rather than affecting the overall rate of protein synthesis.

328
Fig. 5
Comparison of GO enrichment plots for combined trait groups. (a) GO enrichment plot for
the combined "yield & size" traits. The analysis is overwhelmingly dominated by terms
related to the ribosome and protein synthesis. (b) GO enrichment plot for the combined "color
& pigmentation" traits. This analysis reveals a strong enrichment for pathways related to
primary energy metabolism.
335
336

Fig. 6

Comparison of ribosome KEGG pathway enrichment for the combined trait groups. Genes

highlighted in red represent candidate genes identified from the ancestry-corrected GWAS.

The figure displays a standardized KEGG reference map for the ribosome, which includes

examples from different domains of life. (a) Ribosome pathway for the "yield & size" group.

Numerous components of both the small (e.g., S9e, S16e) and large (e.g., L23Ae, L31e). (b)

Ribosome pathway for the "color & pigmentation" group. A smaller, more specific set of

ribosomal proteins (e.g., L4e, L7e, L22e) is highlighted, suggesting that pigmentation may be

linked to specific regulatory functions of the ribosome.

Corrected GWAS links distinct morphological traits to specific biological pathways

To demonstrate the breadth of biological mechanisms identified by our corrected GWAS, we

highlighted the GO enrichment results for three distinct morphological traits (Fig. 7).

The analysis for fruit shape - elliptic revealed an overwhelming enrichment for pathways

related to nucleotide and ATP transport across mitochondrial membranes (Fig. 7a). This

specific finding suggests that the developmental program creating an elliptic shape is strongly

linked to the efficient transport and delivery of cellular energy (ATP) and genetic building

blocks (nucleotides) to support localized growth.

In contrast, the analysis for Fruit Husk Hardness pointed to the fundamental machinery of

protein synthesis and cellular energy production (Fig. 7b). The most significant terms were

related to the tricarboxylic acid (TCA) cycle, which is the central hub of cellular respiration,
and numerous terms related to translation and peptide biosynthesis.
361
Finally, the analysis for fruit apex form ('attenuate', a shape where the fruit's tip gradually
tapers to a long, slender point) implicated pathways involved in stress response and
regulation (Fig. 7c). Enriched terms included cellular response to oxidative stress, response to
wounding, and various peptidase inhibitor activities. This suggests that the development of
this specific shape is tied to the genetic pathways that manage cellular stress and regulate
protein turnover.
368
369

Fig. 7
GO enrichment trees for notable individual morphological traits from the corrected GWAS.
Each panel displays the top significantly enriched GO terms for a specific trait. (a) GO
enrichment for fruit shape - elliptic, dominated by pathways for mitochondrial ATP and
nucleotide transmembrane transport. (b) GO enrichment for fruit husk hardness, showing a
strong association with the TCA cycle and the machinery of translation and protein synthesis.
(c) GO enrichment for fruit apex form - attenuate, highlighting pathways related to cellular
stress response and the regulation of protein activity.
379
380

381 **Predictive modeling of wet bean mass using a Neural Network model**
382 The pod index is calculated using the following formula:
383

$$Pod\ Index = \frac{1000}{average\ dried\ cotyledon\ mass\ in\ grams\ \times\ average\ number\ of\ seeds\ per\ pod}$$

384
This calculation determines the number of pods required to produce one kilogram (1000
grams) of dried cocoa beans. A lower pod index value indicates a higher yield potential, as
fewer pods are needed to produce the same amount of cocoa. Wet bean mass provides a more
direct assessment of yield, so we employed a machine learning approach using Neural
Network model to identify predictive phenotypic traits while excluding obvious correlating
traits such as seed number, fruit length, fruit width, and fruit length to width ratio. The
performance of the trained Neural Boosted model is shown in Fig. 8. The model exhibited
strong predictive accuracy on both the training set (R-squared = 0.790 +/− 0.08) with and the
validation set (R-squared = 0.715 +/− 0.12). The scatter plots of actual vs. predicted wet bean
mass values show a close alignment of the points with the diagonal line representing perfect
prediction, indicating that the model accurately captures the relationship between the
predictor variables and wet bean mass. Further analysis using the Neural Boosted model
identified cotyledon mass as the primary driver of wet bean mass, with a remarkably high
importance score of 6 (Table S1). Cotyledon length emerged as the second most influential
predictor, receiving an importance score of 2. Cotyledon length-to-width ratio showed slight
predictive power with an importance score of 1, whereas cotyledon width and other
phenotypic traits showed little to no predictive influence (Table S1). Cotyledon mass
displayed a substantial positive influence on wet bean mass, evidenced by its main effect of
0.484 and total effect of 0.5198. This suggests that cotyledon mass impacts wet bean mass
both directly and indirectly through interactions with other variables. Cotyledon mass
(portion = 0.520) and cotyledon length (portion = 0.213) accounted for the vast majority
(73.3%) of the predictive power in the model, highlighting their dominant roles in
determining wet bean mass.
408
409

410
Fig. 8
Evaluation of the Neural Boosted model for predicting wet bean mass in cacao, based on
phenotypic data. The model was evaluated on the full dataset (n=346) using a robust 5-fold
cross-validation procedure, which was repeated 20 times. (a) Actual vs. predicted values
aggregated from the training folds. (b) Actual vs. predicted values aggregated from the
validation folds. The black line in each plot represents a perfect prediction (actual =
predicted). The R-squared values shown are the average performance across all validation
runs, demonstrating the model's generalized predictive accuracy.

Discussion

Correlation between patristic distances and phenotypic traits
We first examined how ancestry affected phenotypic traits in 344 diverse cacao accessions.
The results indicate that broad ancestry accounts for a limited fraction of phenotypic
divergence in this panel: significant traits show small positive patristic–phenotypic
correlations, with the strongest for pod index ($r= 0.25$, $q= 0.0019$), followed by cotyledon
width ($r= 0.20$, $q= 0.0019$). While genetic divergence has increased trait difference for these
two traits, the weak overall correlation between patristic distances and phenotypic traits
suggests that broad phylogenetic relationships are not the primary determinants of cacao
phenotypic variation in this collection.

Correcting for population structure reveals the core genetic basis of yield
A key methodological finding of this study was the impact of correcting for population
structure in the GWAS, best illustrated by comparing the naive (uncorrected) and structured
(corrected) models for the pod index trait. The naive analysis highlighted a top SNP in a
peroxidase gene, a plausible but likely confounding signal for general plant vigor[23–25] that
could lead to spurious associations if population structure is not controlled[26,27]. In contrast,
the corrected model elevated TcSNP597 to the top-ranked position, a locus within a gene for
a 21 kDa proteinase inhibitor seed protein, a class of proteins known to be major
determinants of seed quality and nutritional value[28]. This shift is significant, as it moves the
focus from a general stress-response gene to a gene class with a specific role in seed
development and defense, making it a more direct and plausible candidate. Similarly, the

correction refined the associated pathway: the uncorrected results were enriched for the broad
but relevant pathway of "protein processing in the endoplasmic reticulum," while the
corrected results pinpointed the core of this process, becoming overwhelmingly dominated by
the "ribosome" pathway itself (Fig. 6).Together, these comparisons illustrate the utility of a
dual-model approach, where the exploratory findings of a naive analysis are refined by a
structured model to pinpoint the specific biological machinery, in this case, protein synthesis
directly linked to the trait.
449
The genetic architecture of cacao is centered on pleiotropic hubs for protein synthesis
Building on these robust results, our corrected GWAS revealed a core genetic network
associated with cacao productivity, showing that a small set of loci are central to multiple
yield-component traits. This phenomenon, known as pleiotropy, is a key feature of complex
traits in major crops, where single loci often influence multiple, correlated yield
components[29,30]. Specifically, TcSNP281, TcSNP597, and TcSNP1366 were all identified as
top associations for combinations of pod index, wet bean mass, and seed number (Table 1).
The putative genes at these loci are involved in fundamental biological processes: protein
synthesis (60S ribosomal protein L31), seed development and defense (21 kDa proteinase
inhibitor seed protein, known as a negative regulator for seed quality), and energy
metabolism (heparan-alpha-glucosaminide N-acetyltransferase). The convergence of these
crucial yield traits on the same genetic loci strongly suggests that they are regulated by a
central hub controlling the plant's capacity for protein production and energy allocation for
seed development. This suggests that this network represents a key target of selection, where
breeders have, either intentionally or unintentionally, selected for alleles that optimize the
overall capacity for protein production to enhance yield. This finding aligns with a growing
body of evidence highlighting that the synthesis of storage proteins and the regulation of
amino acid and energy metabolism are tightly controlled, central processes that determine
final seed composition and yield[31–34].
469
The validity of our analytical approach is demonstrated by an internal positive control. The
analysis for fruit surface anthocyanin recovered 6-phosphogluconate dehydrogenase
(TcSNP174), a key enzyme biochemically upstream of flavonoid precursors. This finding,
combined with the identification of a locus (TcSNP401) that converges with an
independently reported finding for cacao pigmentation, provides strong support that our
structured model successfully identifies pathway-specific, non-vigor associations. 6-
phosphogluconate dehydrogenase catalyzes a rate-limiting step in the oxidative pentose
phosphate pathway (OPP)[35]. The OPP pathway is critical as it generates carbon precursors for
the shikimate pathway, a direct link that has recently been genetically demonstrated through
the activity of 6-phosphogluconate dehydrogenase[36]. The shikimate pathway, in turn, is the
well-established route for the biosynthesis of aromatic amino acids and subsequently the
entire class of flavonoids, including the anthocyanin pigments that determine color[37,38]. Our
ability to link a top statistical hit from our corrected model directly to a key, rate-limiting
enzyme at the start of the relevant biochemical network provides strong validation for our
findings.

From predictive models to biological mechanisms

Our findings can be contextualized by comparing them to the initial analysis of this germplasm by Bekele et al.[11]. It is noteworthy that both studies, despite using different corrected GWAS models (Bootstrap Forest vs. MLM), independently identified TcSNP401 as a key locus associated with fruit surface anthocyanin. This independent validation strongly supports the importance of this genomic region in cacao pigmentation. Beyond this convergence, our machine learning-based approach identified a largely distinct set of top candidate genes for most traits, suggesting our analysis has captured different facets of the genetic architecture and provides a novel and complementary view of cacao yield genetics.

Furthermore, this study successfully created a bridge between predictive modeling and genetic markers. Our Neural Network model independently identified cotyledon mass and length as the most powerful predictors of final wet bean mass. Our GWAS then provided a genetic basis for these predictive traits. For instance, cotyledon length was strongly associated with a SNP in a gene for a mitochondrial cytochrome c oxidase subunit; this link is highly significant, as the activity of cytochrome c oxidase is known to be indispensable for proper plant embryogenesis and seed viability[39,40]. Cotyledon mass was associated with a SNP in a RING/U-box superfamily protein. This finding is well-supported by studies in other crops showing that this class of E3 ubiquitin ligases directly regulates seed and organ size by targeting key proteins for degradation[41,42].

Our combined-trait GO analysis revealed the fundamental processes underpinning major phenotypic groups. By combining all nine yield- and size-related traits, the analysis converged on a single, overwhelmingly significant biological theme: the ribosome and protein synthesis. A deeper look revealed that numerous components of both the small and large ribosomal subunits were associated with these traits, suggesting that overall yield potential is linked to the general capacity and throughput of the entire protein synthesis machinery. This conclusion is well-supported by studies showing that the disruption of individual ribosomal protein genes leads to significant defects in plant growth, development, and seed size[43–45]. In parallel, the combined analysis of five color- and pigmentation-related traits pointed directly to primary energy metabolism, including glycolysis and ATP generation. A key pathway identified in this group was "photosynthesis - antenna proteins," highlighting the specific gene *Lhcb4*. Antenna proteins regulate how much light energy reaches photosystem II and variations in these proteins can directly impact chloroplast redox balance which is a primary trigger for anthocyanins as photoprotective agents. This provides a direct mechanistic link between the machinery of light capture and the energetically expensive process of pigment production[46,47].

Beyond these broad themes, the analysis of individual corrected traits revealed a diversity of specific biological mechanisms controlling morphology. For instance, the genetics of fruit shape - elliptic were not linked to general growth but to the specific machinery of ATP and nucleotide transport across mitochondrial membranes. This suggests a key role for "energy logistics" in developmental patterning, a concept supported by findings that disrupting mitochondrial transporters can severely hamper organ development and morphology[48,49]. In

another example, fruit husk hardness was strongly associated with the Citrate (TCA) cycle
and the machinery of translation. This provides a direct link between a physical property and
the cell's core metabolic and biosynthetic capacity, as studies have shown that restricting
TCA cycle activity directly impairs the synthesis of secondary cell wall components like
cellulose and lignin that confer structural rigidity[50,51]. These findings showcase the
complexity of developmental genetics and demonstrate our approach's ability to uncover
novel and highly specific hypotheses for a wide range of traits.

Hierarchical clustering of traits based on the ancestry-adjusted GWAS results provided a
global view of the genetic architecture, revealing several functionally significant groupings
(Fig. 3). The analysis partitioned the traits into distinct clusters, most notably grouping key
productivity metrics such as pod index, wet bean mass, seed number, and cotyledon length.
This supports the concept of a shared genetic foundation for overall yield and corroborates
the pleiotropic effects suggested by the single-trait GWAS, where the same SNPs were
associated with multiple of these traits. Clustering of trait–gene association profiles
underscored the modular nature of yield components. Key productivity metrics, pod index
and wet bean mass, formed a tight pair and clustered with seed number (and ovule number,
cotyledon length), consistent with a shared genetic backbone for overall yield. By contrast,
cotyledon mass lay on a distant branch of the dendrogram, proximate to pod wall hardness,
fruit length, and several apex‑form traits. This separation indicates that, although
phenotypically related, the genetic factors governing individual bean mass are partly distinct
from those shaping bean count and pod‑level characteristics. Practically, this suggests that
bean size and bean number can be improved as semi‑independent targets in selection
schemes, rather than assuming one will automatically track the other.

Beyond the clustering of traits, our global analysis of SNP locations revealed a striking
pattern in the underlying genetic architecture. By mapping the top three associated SNPs for
all 27 traits, we identified several genomic "hotspots" containing associations for multiple,
diverse traits (Fig. S2), a phenomenon that has been observed for important agronomic traits
in other major crops[52,53]. A distinct trend emerged where many of these hotspots were
concentrated in the subtelomeric regions of the chromosomes, particularly on Chromosomes
1, 2, 4 and 9. These subtelomeric regions are known to be dynamic parts of the genome, often
characterized as being gene-rich and exhibiting high rates of recombination[54,55]. This unique
combination of features has led to the hypothesis that subtelomeres act as "hotbeds" for new
gene origination and are potential centers for rapid evolutionary adaptation[56,57]. This
concentration of trait-associated loci suggests these dynamic genomic regions may have been
key targets of selection during both the initial domestication of cacao and its subsequent,
geographically diverse improvement.

Implications for cacao breeding and future directions
The successful prediction of wet bean mass using a Neural Network model (Fig. 8), well-
suited for capturing complex, non-linear relationships often found in biological systems,
demonstrates the potential of leveraging phenotypic data for developing predictive tools for
complex traits in cacao[58]. The model's high accuracy (R-squared = 0.790 for training, 0.715

for validation) highlights the strong relationships between the predictor variables and wet
bean mass. Notably, cotyledon mass and cotyledon length were identified as the two most
important predictors (Table S1) of wet bean mass, a direct measure of yield. Larger
cotyledons on young cacao seedlings may be indicators of vigor and higher photosynthetic
capacity, which over the long-term could result in more productive cacao trees, a principle
supported in other crops where early-stage growth traits have been shown to be significantly
predictive of final yield[59]. While this suggests that simple measurement of these two traits
could provide an accurate and cost-effective method for predicting productivity, further
longitudinal evidence is required to demonstrate early screening utility of these key-traits.
Nonetheless, the identification of these early-screening traits may have significant
implications for cacao breeding programs that are often hampered by long generation times
and an urgent need for new, high-yielding varieties[13]. Targeting specific combinations of
traits, guided by the identified clusters (Fig. 3) and the machine learning model (Fig. 8),
could lead to even more efficient selection strategies for improving yield potential.
Furthermore, these results, combined with our GWAS findings, suggest that genomic
selection, incorporating information from multiple loci across the genome, could be a
powerful tool for accelerating cacao improvement, an approach particularly impactful for
perennial crops with long breeding cycles[60].

The findings of this study, particularly the successful application of a Neural Network model
for predicting wet bean mass, provide a springboard for several promising avenues of future
research in cacao. Given the potential impact on breeding strategies, prioritizing the
validation of candidate genes identified through GWAS would be beneficial. This could
involve gene expression studies and functional analyses using transgenic approaches or, more
directly, CRISPR-Cas9-mediated gene editing, which is a critical step for validating the
causal effect of candidate genes identified through association studies[61]. Second, expanding
the association mapping and genomic selection studies to larger and more diverse populations
of cacao, including different genetic groups and geographic origins, would help to confirm
the identified associations and identify additional loci contributing to trait variation across a
wider range of germplasm. This could also involve exploring genotype-by-environment
interactions by evaluating the performance of diverse accessions in multiple locations in
collaboration with international partners. Third, investigating the role of other types of
genetic variants, such as copy number variations and structural variants, could improve
predictive power. Fourth, integrating genomic data with other omics data, such as
transcriptomic, proteomic, and metabolomic profiles, could further elucidate the molecular
mechanisms underlying trait variation. Both the inclusion of structural variants and the use of
multi-omics data are considered key future directions for enhancing genomic prediction
models[60]. Moreover, exploring different Neural Network architectures, such as the deep
learning methods that are increasingly being harnessed for genomic prediction, could provide
even greater accuracy when modeling complex traits from high-dimensional data[62].

This study provides a comprehensive analysis of the phenotypic and genotypic variation in a
diverse cacao collection, dissecting the complex relationships between population structure,
genetic markers, and key agronomic traits. A central outcome was the demonstration of the

profound impact of population structure on association mapping. By applying an ancestry-
adjusted model, our GWAS successfully filtered out misleading signals related to general
plant 'vigor' to instead identify that the core genetic drivers of yield are robust, trait-specific
pathways for ribosome and protein synthesis. This corrected approach led to the identification
of a small set of loci which were responsible for influencing multiple yield components like
pod index, wet bean mass, and seed number. The hierarchical clustering of these corrected
results further illuminated the genetic landscape, partitioning traits into distinct functional
groups related to yield and pigmentation. Moreover, a highly accurate machine-learning
model identified cotyledon mass and length as the most powerful predictors of wet bean
mass, a direct measure of yield. While this work delivers a simple and much needed early-
stage screening tool for cacao breeding programs, further validation of cotyledon mass as a
predictor of yield is required. Collectively, these findings enhance our understanding of the
genetic architecture of yield in cacao and provide a powerful, statistically robust framework
for developing superior varieties through modern breeding strategies.
626

Materials and Methods

627
Phenotype and genotype data
Phenotypic and genotypic data used in this study were obtained from a publicly available,
large-scale assessment of cacao germplasm detailed in Bekele et al.[11]. The study utilized 421
cacao accessions from the ICGT, a diverse collection that includes 263 wild genotypes, 23
accession groups (e.g., wild cacao types such as AMAZ, GU, IMC, MO, NA, PA, POUND,
RB, SCA and SPEC accession groups)[11,63].

Phenotypic evaluations for 27 traits, encompassing flower, fruit, and seed characteristics,
were derived from the dataset by Bekele et al.[11]. Flower traits included anthocyanin intensity
in the column of the pedicel, sepal length, anthocyanin intensity on the ligule, ligule width,
anthocyanin intensity in the filament, style length, and ovule number. Fruit traits included
shape, basal constriction, apex form, surface texture, surface anthocyanin intensity in mature
ridges, ridge disposition, primary ridge separation, pod wall hardness, fruit length, fruit
width, and fruit length to width ratio. Seed traits included number, shape, cotyledon color,
total wet bean mass, cotyledon length, cotyledon width, cotyledon mass, and cotyledon length
to width ratio. Pod index, a key agronomic trait that reflects yield potential in cacao, was also
included as a composite trait[11].

Complete phenotypic data were available for 346 of the 421 accessions[11]. For the genotypic
data, Bekele et al.[11] targeted a total of 836 SNPs located in coding sequences, selected based
on their similarity to known protein sequences. After excluding SNPs with low call rates, 671
high-quality SNPs were retained. Consequently, the final dataset used for analysis in this
study consisted of 671 SNPs and phenotypic data for 346 accessions.

Correlation between patristic distances and phenotypic traits
We tested whether more distantly related accessions are more phenotypically different by
performing Mantel tests (Pearson, 9,999 permutations) between the patristic distance matrix
and pairwise phenotypic distances for each trait. Geneious software (V 2025.2.1) was used to

calculate patristic distances and GU305 and NA58 were omitted due missing overlaps
resulting in a total of 344 out of 346 accessions. Natural log transformations were applied for
normality on traits. FDR was controlled across traits.
656
Analysis of population structure and phenotypic variation using PCA
To visualize the underlying structure of the phenotypic and genotypic variation among the
cacao accessions, and to generate covariates for population structure correction, PCA was
performed using JMP Pro 17. Prior to analysis, the categorical SNP data was processed. First,
missing SNP genotypes (coded as 'N') were imputed using the Multivariate SVD Imputation
method within the 'Explore Missing Values' platform of JMP Pro 17, using 6 singular vectors
and a maximum of 10 iterations. Following imputation, the complete categorical SNP matrix
was converted into a numerical format based on allele counts (e.g., 0, 1, 2) using JMP Pro's
built-in functionalities to serve as input for the analysis. Two separate PCAs were then
conducted. The first was performed on the matrix of all measured phenotypic traits to
visualize phenotypic relationships. The second was performed on the final imputed and
numerically-coded SNP genotype matrix to assess population structure. From this genotypic
PCA, the resulting Principal Component Scores for the first three components were saved
back to the main data. To create the corrected phenotype for the GWAS, these three PCs were
then used as predictors in a standard least squares linear regression model for each trait, and
the resulting residuals were saved as the new, ancestry-corrected target variable.
673
Machine learning-based GWAS to associate markers with traits
To identify genomic regions associated with phenotypic variation in the cacao accessions, we
employed a machine learning GWAS approach based on Bootstrap Forest models
implemented in JMP Pro 17. Unlike traditional linear models, which primarily test for
additive effects of single markers, ensemble ML methods like Bootstrap Forest can capture
complex, non-linear relationships and potential gene-gene interactions (epistasis), which is
advantageous for dissecting polygenic traits.
681
To create a comprehensive view of the genetic architecture, two parallel analyses were
conducted: a naive model without population structure covariates, and a structured
association model that included the first three principal components as fixed-effect
covariates. The naive model based analysis provides an exploratory view of the entire genetic
landscape, identifying all associations including broad effects potentially confounded with
population structure. In contrast, the structured association model based analysis statistically
removes the ancestral background noise to pinpoint loci with a direct and independent effect
on the trait. Comparing these two results allows for the identification of the most robust
candidate genes and provides insight into how much of a trait's genetic signal is tied to the
underlying population structure.
692
The GWAS was conducted for all phenotypic traits, including those related to accession
group, flower morphology, fruit morphology, seed characteristics, cotyledon and bean mass,
and pod index. Pod index is a key agronomic trait that reflects yield potential, calculated as
1000/(average dried cotyledon mass × average number of seeds per pod)[11]. Importantly, it is a

composite trait that takes into account both pod size and seed number, with lower pod index
values indicating higher yield potential[11]. For the naive model based analysis, separate
Bootstrap Forest models were trained to predict each raw phenotypic trait using the full set of
SNP markers as predictors.

For the structured association model based, we first accounted for confounding due to
population structure using a two-step method. A linear model was used to predict each trait
using the first three principal components (PC1, PC2, and PC3) from the genotypic PCA as
predictors. The model can be represented by the formula:

$Y = \beta_0 + \beta_1(PC1) + \beta_2(PC2) + \beta_3(PC3) + \varepsilon$

where Y is the vector of phenotypic values for a given trait, β_0 is the overall mean, PC1,
PC2, and PC3 are the vectors of the principal component scores, β_1-β_3 are their
corresponding fixed-effect regression coefficients, and ε is the vector of residuals.
The residuals from this model (ε), which represent the phenotypic variation independent of
ancestry, were saved. Subsequently, separate Bootstrap Forest models were trained to predict
these ancestry-corrected residuals for each trait using the full set of SNP markers as
predictors.

For all Bootstrap Forest models in both analyses, default settings included the following
parameters: number of trees in the forest was set to 100, the number of terms sampled per
split was 1, the bootstrap sample rate was 1, the minimum number of splits per tree was 10,
the maximum number of splits per tree was 2,000, and the minimum size split was 5. The
top-ranking markers from both the corrected and uncorrected analyses were identified based
on their importance scores, calculated as the "Portion" in JMP Pro. Given the high mapping
resolution, with each marker representing a genomic region of approximately 1000 bp, we
proceeded to identify the most likely candidate genes. To explore these potential candidates,
we identified all annotated genes located within the genomic block represented by each top-
ranking marker using the Criollo v2 reference genome (GCF_000208745.1).

Gene ontology enrichment analysis
A GO enrichment analysis was conducted to investigate the potential biological functions and
pathways overrepresented among the candidate genes associated with key agronomic traits.
To compare the effects of population structure correction, this analysis was performed on
gene lists generated from both the naive (uncorrected) and structured association (corrected)
models for a subset of important traits, including pod index, wet bean mass, and anthocyanin
intensity on the ligule.

For each analysis, a list of unique gene identifiers was compiled from all markers with a
Portion score greater than 0.005, as identified by the Bootstrap Forest models. In cases where
a marker could not be mapped to the reference genome, it was omitted from this downstream
analysis. The curated list of unique *Theobroma cacao* gene IDs was submitted to ShinyGO
v0.82[64] (available at http://bioinformatics.sdstate.edu/go/). The enrichment analysis was

conducted using the *Theobroma cacao* Belizean Criollo B97-61 B2 genome database. Default
settings were applied, including a FDR cutoff of 0.05 to determine statistical significance.
Pathway size filters were set to a minimum of 2 and a maximum of 5,000 genes per GO term.
744

Hierarchical clustering of trait association profiles
To visualize the global genetic relationships between traits, we performed hierarchical
clustering analysis. This analysis used the Ward method based on the importance scores
(portion) derived from the structured Bootstrap Forest models for all traits. This approach
groups traits that exhibit similar genetic association profiles, suggesting shared underlying
genetic pathways or pleiotropic effects.
751

Development of machine learning model for predicting wet bean mass
To further investigate the complex relationships between the various measured traits and wet
bean mass, a machine learning approach employing Neural Network models was utilized[65].
The primary goal was to train a predictive model for wet bean mass, a key agronomic trait in
cacao, based on the suite of phenotypic characteristics used in this study. Model selection and
training were performed using JMP Pro 17. An initial model screening step was conducted
using the built-in model screening function to evaluate the performance of various algorithms
and identify the most promising model type for predicting wet bean mass. The Neural
Boosted model emerged as the best-performing model and was therefore selected for further
analysis. The model was configured with a single hidden layer containing three neurons with
a hyperbolic tangent (TanH) activation function (NTanH(3)) and built using 20 boosting
iterations (NBoost(20)).
764

To robustly assess the model's predictive power, a repeated 5-fold cross-validation scheme
was implemented on the full dataset (n=346). This procedure was repeated 20 times to ensure
a stable and reliable estimate of the model's performance. The Neural Boosted model, with
the specified NTanH(3) and NBoost(20) settings, was trained and validated during each run
of the cross-validation. Obvious correlating phenotypic traits that would lead to spurious
associations were excluded such as seed number, fruit length, fruit width, and fruit length to
width ratio, as to identify non-obvious phenotypic predictors. The performance of the trained
model was assessed using the coefficient of determination (R-squared), averaged across all
validation folds from the repeated runs. The importance of individual features (traits) in the
Neural Boosted model was assessed to gain insights into the factors most strongly influencing
wet bean mass prediction.
776

Acknowledgements

We are also grateful to the reviewers for their constructive feedback. Mention of any trade
names or commercial products in this article is solely for the purpose of providing specific
information and does not imply recommendation or endorsement by the U. S. Department of
Agriculture. USDA is an equal opportunity provider and employer, and all agency services
are available without discrimination.

Funding

This work is supported by the U.S. Department of Agriculture, Agricultural Research
Service, In-House Projects No. 8042-21220-258-000-D and 8042-21000-303-000-D.
787

788 ## Conflicts of interest
789 The authors declare no conflicts of interest.
790

Author Contributions
E.A. conceptualized and designed the study. The investigation was performed by J.B., S.P.,
and E.A. Data analysis, validation, and visualization were conducted by J.B., D.L., S.P.C.,
S.P., and E.A. The methodology and resources were provided with contributions from J.B.,
I.B., S.L., J.H.J., S.P.C., A.H.L., and L.W.M. E.A. wrote the original manuscript draft. All
authors reviewed, edited, and approved the final manuscript.
797

Data availability
All phenotypic and genotypic data analyzed in this study are publicly available. These data
were originally generated and published by Bekele et al. (2022) and can be found within the
Supporting Information files associated with that publication (PLoS One 17(1): e0260907),
accessible via: https://doi.org/10.1371/journal.pone.0260907.

References

1. Díaz-Montenegro, J. Livelihood strategies and risk behavior of cacao producers in

Ecuador: effects of national policies to support cacao farmers and specialty cacao

landraces. (2019).

2. Hall, J. N. Applying a One Health approach to study the livelihoods of cocoa farming

communities in Bougainville. (2022).

3. Bekele, F. & Phillips-Mora, W. Cacao (Theobroma cacao L.) breeding. *Advances in*

Plant Breeding Strategies: Industrial and Food Crops: Volume 6 409–487 (2019).

4. Alden, D. The significance of cacao production in the Amazon region during the late

colonial period: an essay in comparative economic history. *Proceedings of the American*

Philosophical Society **120**, 103–135 (1976).

5. Gardea, A. A. *et al.* Cacao (Theobroma cacao L.). *Fruit and Vegetable Phytochemicals:*

Chemistry and Human Health, 2nd Edition 921–940 (2017).

6. Kongor, J. E., Owusu, M. & Oduro-Yeboah, C. Cocoa production in the 2020s:

challenges and solutions. *CABI Agriculture and Bioscience* **5**, 102 (2024).

7.   Walters, D. *Chocolate Crisis: Climate Change and Other Threats to the Future of Cacao.*

(University Press of Florida, 2020).

8.   Boza, E. J. *et al.* Genetic characterization of the cacao cultivar CCN 51: its impact and

significance on global cacao improvement and production. *Journal of the American*

Society for Horticultural Science **139**, 219–229 (2014).

9.   Mustiga, G. M. *et al.* Phenotypic description of Theobroma cacao L. for yield and vigor

traits from 34 hybrid families in Costa Rica based on the genetic basis of the parental

population. *Frontiers in plant science* **9**, 808 (2018).

10. Izzah, N. K. *et al.* Improvement of Cacao Pod Characteristics and its Molecular

Characterization in 4 F1 Cacao Populations. (2024).

11. Bekele, F. L. *et al.* Genome-wide association studies and genomic selection assays

made in a large sample of cacao (Theobroma cacao L.) germplasm reveal significant

marker-trait associations and good predictive value for improving yield potential. *Plos*

one **17**, e0260907 (2022).

12. Bediako, K. A., Padi, F. K., Obeng-Bio, E. & Ofori, A. Genetic diversity and parentage of

cacao (Theobroma cacao L.) populations from Ghana using single nucleotide

polymorphism (SNP) markers. *Plant Genetic Resources* 1–8 (2024).

13. Gutiérrez, O. A., Campbell, A. S. & Phillips-Mora, W. Breeding for disease resistance in

cacao. *Cacao diseases: a history of old enemies and new encounters* 567–609 (2016).

14. Rodriguez-Medina, C. *et al.* Cacao breeding in Colombia, past, present and future.

Breeding Science **69**, 373–382 (2019).

15. Wickramasuriya, A. M. & Dunwell, J. M. Cacao biotechnology: current status and future

prospects. *Plant biotechnology journal* **16**, 4–17 (2018).

16. McElroy, M. S. *et al.* Prediction of cacao (Theobroma cacao) resistance to

Moniliophthora spp. diseases via genome-wide association analysis and genomic

selection. *Frontiers in plant science* **9**, 343 (2018).

17. Romero Navarro, J. A. *et al.* Application of genome wide association and genomic

prediction for improvement of cacao productivity and resistance to black and frosty pod

diseases. *Frontiers in Plant Science* **8**, 1905 (2017).

18. Duarte-Carvajalino, J. M., Paramo-Alvarez, M., Ramos-Calderón, P. F. & González-

Orozco, C. E. Estimation of canopy attributes of wild cacao trees using digital cover

photography and machine learning algorithms. *iForest-Biogeosciences and Forestry* **14**,

517 (2021).

19. Omas-as, A. M., ARBOLEDA, E. R. & DAANG, J. A. M. Machine Learning as a Strategic

Tool: A Comprehensive Literature Review for Advancing Agricultural Analysis, with

Emphasis on the Cocoa Bean Quality Assessment. *International Journal of Scientific*

Research and Engineering Development. **7**, (2024).

20. Tan, J., Balasubramanian, B., Sukha, D., Ramkissoon, S. & Umaharan, P. Sensing

fermentation degree of cocoa (Theobroma cacao L.) beans by machine learning

classification models based electronic nose system. *Journal of Food Process*

Engineering **42**, e13175 (2019).

21. Lamos-Díaz, H., Puentes-Garzón, D.-E. & Zarate-Caicedo, D.-A. Comparison between

machine learning models for yield forecast in cocoa crops in santander, Colombia.

Revista Facultad de Ingeniería **29**, (2020).

22. Breiman, L. Random forests. *Machine learning* **45**, 5–32 (2001).

23. Akbudak, M. A., Filiz, E., Vatansever, R. & Kontbay, K. Genome-wide identification and

expression profiling of ascorbate peroxidase (APX) and glutathione peroxidase (GPX)

genes under drought stress in sorghum (Sorghum bicolor L.). *Journal of Plant Growth*

Regulation **37**, 925–936 (2018).

24. Kamruzzaman, M. *et al.* Pinpointing genomic loci for drought-induced proline and

hydrogen peroxide accumulation in bread wheat under field conditions. *BMC Plant*

Biology **22**, 584 (2022).

25. Zhou, Z. *et al.* Identification of genomic regions affecting grain peroxidase activity in

bread wheat using genome-wide association study. *BMC Plant Biology* **21**, 523 (2021).

26. Richardson, K. & Jones, M. C. Why genome-wide associations with cognitive ability

measures are probably spurious. *New Ideas in Psychology* **55**, 35–41 (2019).

27. Sullivan, P. F. Spurious genetic associations. *Biological psychiatry* **61**, 1121–1126

(2007).

28. Clemente, A. *et al.* Eliminating anti-nutritional plant food proteins: the case of seed

protease inhibitors in pea. *PLoS One* **10**, e0134634 (2015).

29. Schulthess, A. W. *et al.* The roles of pleiotropy and close linkage as revealed by

association mapping of yield and correlated traits of wheat (Triticum aestivum L.).

Journal of Experimental Botany **68**, 4089–4101 (2017).

30. Liu, C. *et al.* Multi-trait genome-wide association studies reveal novel pleiotropic loci

associated with yield and yield-related traits in rice. *Journal of Integrative Agriculture*

(2024).

31. Galili, G., Avin-Wittenberg, T., Angelovici, R. & Fernie, A. R. The role of photosynthesis

and amino acid metabolism in the energy status during seed development. *Frontiers in*

plant science **5**, 447 (2014).

32. Amir, R., Galili, G. & Cohen, H. The metabolic roles of free amino acids during seed

development. *Plant Science* **275**, 11–18 (2018).

33. Borisjuk, L. *et al.* Seed development and differentiation: a role for metabolic regulation.

Plant biology **6**, 375–386 (2004).

34. Shrestha, V. *et al.* Multiomics approach reveals a role of translational machinery in

shaping maize kernel amino acid composition. *Plant physiology* **188**, 111–133 (2022).

35. Kruger, N. J. & Von Schaewen, A. The oxidative pentose phosphate pathway: structure

and organisation. *Current opinion in plant biology* **6**, 236–246 (2003).

36. Tang, Q. *et al.* 6-Phosphogluconate dehydrogenase 2 bridges the OPP and shikimate

pathways to enhance aromatic amino acid production in plants. *Science China Life*

Sciences 1–11 (2024).

37. Herrmann, K. M. & Weaver, L. M. The shikimate pathway. *Annual review of plant biology*

50, 473–503 (1999).

38. Maeda, H. & Dudareva, N. The shikimate pathway and aromatic amino acid biosynthesis

in plants. *Annual review of plant biology* **63**, 73–105 (2012).

39. Dahan, J. *et al.* Disruption of the CYTOCHROME C OXIDASE DEFICIENT1 Gene

Leads to Cytochrome c Oxidase Depletion and Reorchestrated Respiratory Metabolism

in Arabidopsis. *Plant Physiology* **166**, 1788–1802 (2014).

40. Mansilla, N., Garcia, L., Gonzalez, D. H. & Welchen, E. AtCOX10, a protein involved in

haem o synthesis during cytochrome c oxidase biogenesis, is essential for plant

embryogenesis and modulates the progression of senescence. *Journal of Experimental*

Botany **66**, 6761–6775 (2015).

41. Lv, Q. *et al.* Wheat E3 ubiquitin ligase TaGW2-6A degrades TaAGPS to affect seed size.

Plant Science **320**, 111274 (2022).

42. Xia, T. *et al.* The Ubiquitin Receptor DA1 Interacts with the E3 Ubiquitin Ligase DA2 to

Regulate Seed and Organ Size in Arabidopsis. *The Plant Cell* **25**, 3347–3359 (2013).

43. Popescu, S. C. & Tumer, N. E. Silencing of ribosomal protein L3 genes in N. tabacum

reveals coordinate expression and significant alterations in plant growth, development

and ribosome biogenesis. *The Plant Journal* **39**, 29–44 (2004).

44. Tian, S. *et al.* Ribosomal protein NtRPL17 interacts with kinesin-12 family protein NtKRP

and functions in the regulation of embryo/seed size and radicle growth. *Journal of*

Experimental Botany **68**, 5553–5564 (2017).

45. Weis, B. L., Kovacevic, J., Missbach, S. & Schleiff, E. Plant-Specific Features of

Ribosome Biogenesis. *Trends in Plant Science* **20**, 729–740 (2015).

46. Pescador-Dionisio, S. *et al.* Contribution of the regulatory miR156-SPL9 module to the

drought stress response in pigmented potato (Solanum tuberosum L.). *Plant Physiology*

and Biochemistry **217**, 109195 (2024).

47. Werner, C., Fasbender, L., Romek, K. M., Yáñez-Serrano, A. M. & Kreuzwieser, J. Heat

Waves Change Plant Carbon Allocation Among Primary and Secondary Metabolism

Altering CO2 Assimilation, Respiration, and VOC Emissions. *Frontiers in Plant Science*

Volume 11-2020, (2020).

48. Brown, D. C. W. & Thorpe, T. A. Mitochondrial activity during shoot formation and growth

in tobacco callus. *Physiologia Plantarum* **54**, 125–130 (1982).

49. Jia, F. *et al.* Overexpression of Mitochondrial Phosphate Transporter 3 Severely

Hampers Plant Development through Regulating Mitochondrial Function in Arabidopsis.

PLOS ONE **10**, e0129717 (2015).

50. van der Merwe, M. J. *et al.* Tricarboxylic Acid Cycle Activity Regulates Tomato Root

Growth via Effects on Secondary Cell Wall Production. *Plant Physiology* **153**, 611–621

(2010).

51. Verbančič, J., Lunn, J. E., Stitt, M. & Persson, S. Carbon Supply and the Regulation of

Cell Wall Synthesis. *Molecular Plant* **11**, 75–94 (2018).

52. Bus, A. *et al.* Species- and genome-wide dissection of the shoot ionome in Brassica

napus and its relationship to seedling development. *Frontiers in Plant Science* **Volume**

5-2014, (2014).

53. Wang, P., Zhou, G., Cui, K., Li, Z. & Yu, S. Clustered QTL for source leaf size and yield

traits in rice (Oryza sativa L.). *Molecular Breeding* **29**, 99–113 (2012).

54. Aguilar, M. & Prieto, P. Sequence analysis of wheat subtelomeres reveals a high

polymorphism among homoeologous chromosomes. *The Plant Genome* **13**, e20065

(2020).

55. Fan, C. *et al.* The Subtelomere of Oryza sativa Chromosome 3 Short Arm as a Hot Bed

of New Gene Origination in Rice. *Molecular Plant* **1**, 839–850 (2008).

56. Brown, C. A., Murray, A. W. & Verstrepen, K. J. Rapid Expansion and Functional

Divergence of Subtelomeric Gene Families in Yeasts. *Current Biology* **20**, 895–903

(2010).

57. Saint-Leandre, B. & Levine, M. T. The Telomere Paradox: Stable Genome Preservation

with Rapidly Evolving Proteins. *Trends in Genetics* **36**, 232–242 (2020).

58. Hesami, M., Naderi, R., Tohidfar, M. & Yoosefzadeh-Najafabadi, M. Development of

support vector machine-based model and comparative analysis with artificial neural

network for modeling the plant tissue culture procedures: effect of plant growth

regulators on somatic embryogenesis of chrysanthemum, as a case study. *Plant*

methods **16**, 1–15 (2020).

59. Wu, D. *et al.* Combining high-throughput micro-CT-RGB phenotyping and genome-wide

association study to dissect the genetic architecture of tiller growth in rice. *Journal of*

Experimental Botany **70**, 545–561 (2019).

60. Seyum, E. G. *et al.* Genomic selection in tropical perennial crops and plantation trees: a

review. *Molecular Breeding* **42**, 58 (2022).

61. Sallam, A., Alqudah, A. M., Baenziger, P. S. & Rasheed, A. Editorial: Genetic validation

and its role in crop improvement. *Frontiers in Genetics* **Volume 13-2022**, (2023).

62. Crossa, J. *et al.* Expanding genomic prediction in plant breeding: harnessing big data,

machine learning, and advanced software. *Trends in Plant Science* **30**, 756–774 (2025).

63. Motamayor, J. C. *et al.* Geographic and genetic population differentiation of the

Amazonian chocolate tree (Theobroma cacao L). *PloS one* **3**, e3311 (2008).

64. Ge, S. X., Jung, D. & Yao, R. ShinyGO: a graphical gene-set enrichment tool for animals

and plants. *Bioinformatics* **36**, 2628–2629 (2020).

65. Schwenk, H. & Bengio, Y. Boosting neural networks. *Neural computation* **12**, 1869–1887

(2000).

Supplementary Figures

977
Fig. S1
Bars show the Mantel correlation coefficient (r) between genetic distance and pairwise
phenotypic difference for each trait. Only traits with q < 0.05 are displayed; bars are ordered
by q and then r. Text above each bar reports r and q. Positive r indicates that more distantly
related accessions tend to differ more in the trait. The categorical label "Acc Group" was
excluded a priori. FR_BEW is cotyledon width, FL_LIG_COL is flower ligule anthocyanin
intensity, FL_PCOL_C is flower pedicel column anthocyanin intensity, FL_STY_LEN is
flower sepal length, FR_BW is cotyledon mass, FR_HH is pod wall hardness,
FR_BES_oblong is seed shape oblong, FL_FIL_COL is flower style anthocyanin intensity,
and PODL_W is pod length to width ratio.

988
989
990

991
Fig. S2
Genomic hotspot map of top associations across all traits. The plot shows the physical
locations (Mbp) of the top 3 most significant SNPs for each of the 27 traits analyzed. Clear

clusters, or "hotspots," of associated loci are visible on several chromosomes. A complete list
of all SNPs shown, their associated traits, and putative candidate genes can be found in
Supplementary Data 1.

999
1000

1001
Fig. S3
KEGG pathway for protein processing in the endoplasmic reticulum. This pathway was
significantly enriched in the uncorrected GWAS for the pod index trait. The analysis
identified several key genes within this pathway (highlighted in red), including components
of the N-glycosylation machinery (*OST4*) and molecular chaperones involved in protein
folding and the ER-associated degradation (ERAD) pathway (*Hsp90*, *Hsp40*). This suggests
that protein quality control is a significant component of the broad "vigor" signal detected in
the naive analysis.
1010

FIG. S4

KEGG pathway for photosynthesis - antenna proteins. This pathway was identified as significantly enriched in the analysis of the pod index trait for both the naive (uncorrected) model and the structured association (corrected) model, which highlighted a key gene, *Lhcb1* (highlighted in red), encoding a major light-harvesting chlorophyll-binding protein of photosystem II. The consistent significance of this pathway across both analytical models suggests that the efficiency of light capture is a robust component of the genetic architecture for yield-related traits in this cacao collection.

Ribosomal proteins

Small subunit

SAe	S2e	S3e	S3Ae	S4e	S5e	S6e	S7e	S8e	S9e	S10e	S11e	S12e	S13e	S14e	S15e	S15Ae	S16e	S17e	S18e	S19e
S20e	S21e	S23e	S24e	S25e	S26e	S27e	S27Ae	S28e	S29e	S30e	RACK1	Asc1								

S1	S2	S3	S4	S5	S6	S7	S8	S9	S10	S11	S12	S13	S14	S15	S16	S17	S18	S19	S20	S21

MRFS18B	MRPS22	MRPS23	MRPS24	MRPS25	MRPS26	MRPS27	MRPS28	MRPS29	MRPS30	MRPS31	MRPS33	MRPS34	MRPS35	MRPS36	MRP10	MRP21	YMR31	GCD1	PSRP3	PSRP4
							Dap3													

Large subunit

L3e	L4e	L5e	L6e	L7e	L7Ae	L8e	L9e	L10e	L10Ae	L11e	L12e	L13e	L13Ae	L14e	L15e	L17e	L18e	L18Ae	L19e	L21e
L22e	L23e	L23Ae	L24e	L26e	L27e	L27Ae	L28e	L29e	L30e	L31e	L32e	L34e	L35e	L35Ae	L36e	L37e	L37Ae	L38e	L39e	L40e
L41e	L44e	LP0	LP1	LP2	LP3	LX														
	L36Ae																			

L1	L2	L3	L4	L5	L6	L7A	L9	L10	L11	L7/L12	L12	L13	L14	L15	L16	L17	L18	L19	L20	L21
L22	L23	L24	L25	L27	L28	L29	L30	L31	L32	L33	L34	L35	L36							

MRPL37	MRPL38	MRPL39	MRPL40	MRPL41	MRPL42	MRPL43	MRPL44	MRPL45	MRPL46	MRPL47	MRPL48	MRPL49	MRPL50	MRPL51	MRPL52	MRPL53	MRPL54	MRPL55
MRPL15	MRPL25	MRPL35	MRP49	PSRP5	PSRP6													

Fig. S5

KEGG pathway for the ribosome. The pathway was the most significantly enriched result from the structured association (corrected) GWAS for the pod index trait. The analysis identified multiple genes encoding specific ribosomal proteins (highlighted in red), including components of the small subunit (*S9e*, *S16e*) and the large subunit (*L23Ae*, *L31e*, *L18e*). This finding strongly suggests that the genetic variation controlling yield potential, after correcting for population structure, is directly linked to the core machinery of protein synthesis.

Fig. S6
KEGG pathway for photosynthesis - antenna proteins. The pathway was identified as
significantly enriched in the corrected, combined-trait analysis for the "color & pigmentation"
group. The analysis highlighted the gene *Lhcb4* (highlighted in red), which encodes a key
component of the light-harvesting chlorophyll protein complex. This finding suggests a
specific link between the machinery of light capture and the energy-intensive process of
producing the pigments responsible for color.

Supplementary Tables

Trait	Main effect	Total effect	Importance score
Cotyledon mass	0.4843	0.5198	6
Cotyledon length	0.1864	0.2133	2
Cotyledon length to width ratio	0.0867	0.1101	1
Cotyledon width	0.04815	0.06895	0

Table S1
Most important traits for wet bean mass prediction based on the Neural Boosted model
analysis. Relative importance of the top four predictor traits as determined by the Neural
Network model. Feature importance was assessed using a repeated 5-fold cross-validation
procedure on the full dataset of 346 cacao accessions. The table lists each trait, its main
effect, total effect, and importance score. The importance score reflects the contribution of
each trait to the model's predictive accuracy, with higher scores indicating a stronger
influence. Cotyledon mass was identified as the most important predictive variable.