Structural Genomic Variation and Its Potential Role In Deer Speciation

Faezeh Azimi Chetabi[1], Aaron Shafer[1]

[1] 2140 East Bank Drive, Environmental and Life Sciences Graduate Program, Trent University, Peterborough, Ontario, Canada K9J 7B8

Correspondence: faezehazimichetabi@trentu.ca

Abstract

Speciation is a key driver of biodiversity and understanding its genomic underpinnings is important for predicting and managing biodiversity. Structural variants (SVs) are large-scale (>50 bp) changes in the genome and have been implicated in adaptive divergence and reproductive isolation. We investigated the role of SVs in the speciation and divergence of two deer species (*Odocoileus* spp.) across their North American range. Using multiple long-read and a short-read datasets, our bioinformatics workflow revealed SVs and genomic features that were unique to each species. The majority of species-specific SVs were deletions and insertions suggesting that these variants may show higher likelihoods of fixation within populations. Further, while most SVs were intergenic, some genes found to be impacted by species-specific SVs were under positive selection inferred from dN/dS. We also observed an increased number of enhancer motifs found in SVs compared to the rest of the genome. The SV-affected genes were associated with reproduction, sensory adaptation, and several metabolic pathways. Many of these functions are relevant to fertility and deer biology and therefore may provide insights into potential mechanisms leading to reproductive divergence.

Keywords: Comparative genomics, short-read, long-read, adaptation, genomic divergence, *Odocoileus*

1. Introduction

Speciation is a fundamental process driven most notably by ecological divergence and geographic isolation (Sobel, 2010), with the ultimate mode(s) of divergence leaving tell-tale signals on the genome (Shang et al., 2023). Structural variants (SVs) have long been considered important in driving divergence, with early cytogenetic work pointing to chromosomal differences as barriers to reproduction (Rieseberg, 2001). For example, two distinct clades of Brocket deer (*Mazama* spp.) show karyotypic differences that contribute to reproductive isolation (Abril et al., 2010). More recently, comprehensive reviews have highlighted that SVs, ranging from inversions to translocations, can influence speciation across diverse taxa (Wellenreuther et al., 2019; Berdan et al., 2023). The advent of high-throughput sequencing and novel bioinformatics tools has made it possible to assay SVs at base-pair resolution, opening a new line of inquiry into their role in adaptation and speciation.

While single nucleotide polymorphisms have historically dominated evolutionary genomics, SVs by definition affect larger stretches of DNA and can encompass entire genes and regulatory regions and may have stronger fitness consequences (Wellenreuther et al. 2019). Structural variants are generally considered features >50 bp and fall into several main types: deletions, duplications, inversions, insertions and translocations. Different sequencing and analytical techniques can be employed to detect SVs (Escaramís et al., 2015). Short-read (SR) data with paired read and split read analyses offer high-resolution breakpoint detection, though their effectiveness may depend on factors like insert size and coverage depth (Escaramís et al., 2015). Read depth analysis, often used with both SR and long-read (LR), provides information into SV presence by examining read density, but is less precise with breakpoint resolution (Escaramís et al., 2015). For LR data, de novo assembly and genome alignments are frequently employed, and offers the advantage of identifying novel SVs without relying on reference genomes, though it is more resource-intensive (Escaramís et al., 2015). Long-read sequencing is generally preferable than SR sequencing for detecting complex and large SVs, as SR struggle with novel insertions and complex rearrangements (Mahmoud et al., 2019). More broadly, LR sequencing is

proving transformative, revealing the prevalence and complexity of SVs across genomes and reshaping how we study divergence (Berdan et al., 2023).

Structural variants such as inversions and translocations can promote speciation through mechanisms like reproductive isolation (Homolka et al., 2007). Recombination suppression by SVs can couple adaptive alleles and incompatibilities, reinforcing divergence even in the face of gene flow (Berdan et al., 2023; Wellenreuther et al., 2019). Autosomal translocations have been shown to cause male sterility in laboratory mice due to disrupted synapsis and meiotic silencing (Homolka et al., 2007), while Yang et al., (2024) showed convergent evolutionary signals of SVs associated with domestication, adaptation, and production traits. Further, Gompert et al. (2025) demonstrated that complex structural variants, including inverted translocations, can independently arise in separate populations and repeatedly drive adaptive divergence by suppressing recombination and clustering functionally important genes, revealing a predictable mechanism of local adaptation and evolutionary change. Structural variants can also drive phenotypic diversity by influencing gene expression, with enhancers often connecting genetic changes to species-specific phenotypes (Kaplow et al., 2023). Collectively, these findings highlight SVs and enhancer regions as potential drivers of evolutionary divergence, shaping unique phenotypes and enabling evolution of traits across species.

The *Cervidae* (Deer) family offers an interesting case study for exploring SVs and genetic history. This family encompasses over 60 species with its origins dating back nearly 20 million years (Gonzalez and Duarte, 2020). The red brocket deer (*Mazama americana*) exhibits significant chromosomal diversity, with at least six distinct karyotypes across South America, suggesting the presence of multiple cryptic species (Cursino et al., 2014). Studies on hybrids between these karyotype lineages reveal post-zygotic reproductive isolation, with hybrids showing sterility or subfertility due to chromosomal imbalances impacting ovarian structure, oocyte development, and embryonic genome activation (Cursino et al., 2014). Two closely related species, white-tailed deer (WTD; *Odocoileus virginianus*) and mule deer (MD; *O. hemionus*), have a relatively recent speciation history, appearing to have split in allopatry just

over 1 million years ago (Ma) but also undergoing contemporary hybridization throughout their range (Kessler et al., 2023). Kessler et al., (2023) suggested potential incompatibilities given the lack of nuclear introgression and historical gene flow. Structural variations, such as inversions that suppress recombination in heterozygotes, might help maintain distinct genetic clusters between two species (Stevison et al., 2011). Barriers to gene flow could further be reinforced by genetic incompatibilities caused by SVs, suggesting that SVs might have play a role in species divergence (e.g. Martin et al., 2017 & Baurens et al., 2019).

In this study, we integrated multiple LR and SR sequencing datasets from MD and WTD across their geographic ranges to identify and characterize species-specific SVs. This comparative framework, which combines diverse sequencing approaches and detection tools, enables a comprehensive assessment of the evolutionary impact of SVs. We hypothesize that fixed SVs are important to deer speciation between the two deer species through two, non-mutually exclusive mechanisms: by driving adaptive divergence through changes in genic and regulatory regions, and by generating genomic incompatibilities that reduce hybrid fitness and reinforce reproductive isolation. By identifying consistent, species-specific SVs and examining their overlap with functional genomic regions, this study aims to better understand how SVs shapes adaptation, divergence, and the formation of species boundaries in *Odocoileus*.

2. Materials and Methods

2.1. Data generation and retrieval

We collated publicly available Illumina whole-genome short-read datasets from 79 deer (57 WTD and 22 MD; BioProject PRJNA830519); DNA was extracted from tissue samples (Qiagen DNeasy Blood and Tissue Kit) and sequenced at the Centre for Applied Genomics (Toronto, Canada) on an Illumina HiSeqX platform (see Kessler et al., 2023 for more information). We also retrieved two LR datasets from NCBI (SRR6668252 for WTD and SRR15720448 for MD).

We generated new LR data from five deer semen samples (3 WTD and 2 MD); samples were first thawed on ice and cells were pelleted in an isotonic sperm wash solution to remove debris and reduce somatic cell contamination. DNA was extracted using the NEB Monarch® HMW DNA Extraction Kit for Cells & Blood (#T3050) UHMW protocol, with some modifications to the cell lysis step. Specifically, cells were digested at 56°C for 1 hour at 300 rpm with 100 μL Nuclei Prep Buffer, 100 μL Nuclei Lysis Buffer, 10 μL Proteinase K (NEB #P8107, 20 mg/ml), and 10 μL 1M DTT (GoldBio #DTT in dH2O, final concentration ~50 mM). An additional 20-minute digestion was then performed with 5 μL RNase A (NEB #T3018, 20 mg/ml). DNA quality control, CCS library preparation, and sequencing on the PacBio Revio sequencing instrument was then conducted. Collectively, there are four LR samples for WTD and three LR samples for MD. General characteristics and quality of the long-read datasets used in our analyses were assessed using SeqKit (Shen et al., 2016). For each LR sample, we summarized the total number of reads, as well as the minimum, average, and maximum read lengths (Table 2).

We acquired two reference genome assemblies from the National Center for Biotechnology Information (NCBI) database: one of WTD (JAJQKH000000000) and the other of MD (JAJLRB000000000). To assess the completeness of these assemblies, we evaluated them using BUSCO (Simão et al, 2015), with the score of 98.3% for WTD and 98.0% for MD. We used the GENESPACE v1.1.4 R package (Lovell et al., 2022), that required OrthoFinder (Emms et al, 2019), DIAMOND (Buchfink et al, 2021) and MCScanX (Wang et al, 2012), to identify syntenic blocks and visualize macrosynteny relationships between WTD (*Odocoileus virginianus*), MD (*Odocoileus hemionus*), Caribou (*Rangifer tarandus*, CATKSN000000000), and domestic cattle (*Bos taurus*, NKLS00000000). To visualize chromosomal relationships, we generated riparian (macrosynteny) plots using the plot_riparian() function in GENESPACE. Color palettes and themes were customized using R language. The WTD genome was used as the reference in the visualization.

2.2. SV detection pipeline

We employed two distinct analytical approaches to detect SVs. We used both LR datasets and SR data in conjunction with highest quality genome per species to detect species specific SVs (Figure S1-S2). Because reference genome assemblies can produce artefacts (Howe et al, 2021), we recognized that there are multiple data and reference genome scenarios that can generate species-specific SVs (Table 1). The SVs detected in Scenarios 3 and 6 (both con-specific mapping) are likely only to arise via an assembly artefact and thus were removed form downstream analyses. All SV indels are defined relative to the reference genome, meaning each reported indel represents reference-relative classifications rather than absolute evolutionary events.

For the SR pipeline, we removed adaptors from the raw fastq files using Trimmomatic (Bolger et al., 2014) with a MINLEN threshold of 36. We employed Bowtie2 (Langmead et al., 2012) to align all samples to both the reference genomes of WTD and MD. We used SAMtools (Danecek et al., 2021) to convert, sort, and index the SAM files. To ensure data integrity, duplicate reads were eliminated using Picard (Broad Institute, 2018) with REMOVE_SEQUENCING_DUPLICATES set to TRUE. We employed three different SV calling tools for SR data: Delly (Rausch et al., 2012), Manta (Chen et al., 2016), and Lumpy (Layer, 2014), all with default settings. Further refinement and comparative analysis of VCFs were performed using BCFtools (Danecek et al., 2021), retaining only structural variants with SUPP > 5, representing the number of sequencing reads supporting a given SV. We merged the SV call sets from Delly, Manta, and Lumpy using Jasmine (Kirsche et al, 2023), configured to confirm SVs found in at least two tools. Using BCFtools (Danecek et al., 2021), we filtered the structural variants to extract specific information for comparison, such as the number of SVs, their sizes in base pairs (bp), scores, and zygosity status.

Our LR analysis pipeline followed a workflow similar to that described by Mérot et al. (2023). First, we mapped the LR datasets to the WTD and MD reference genomes using Winnowmap (Jain et al, 2020). The aligned SAM files were then converted to BAM format, sorted, and indexed with SAMtools (Danecek et al., 2021). Structural variants were identified using a combination of Sniffles (Sedlazeck et

al., 2018), SVIM (Heller et al, 2019), and NanoVar (Tham et al, 2020). Finally, we employed Jasmine (Kirsche et al, 2023) to merge the SV calls into a single VCF file. We filtered the merged VCF files based on read support greater than 5.

Both LR and SR pipelines produced four mapping categories: MD2MD, MD2WTD, WTD2WTD, and WTD2MD (Figure S1-S2). For each individual, SVs were classified according to whether they were detected in: (1) both WTD2WTD and WTD2MD; (2) both MD2WTD and MD2MD; (3) only WTD2WTD; (4) only WTD2MD; (5) only MD2WTD; or (6) only MD2MD (Table 1). This classification framework enabled us to identify assembly artefacts and species-specific SVs (See Table 1, Figure S1-S2). The SVs from scenarios 3 and 6 = were not included in further analysis,

To allow cross-referencing between species, we used MUMmer (Delcher et al., 2002) to align the WTD and MD reference genomes. Using the *show-coords* utility, we extracted the corresponding aligned intervals and identified regions shared between the two genomes. This allowed us to accurately lift over scaffold coordinates. To further refine our dataset and identify SVs with the potential relevance to speciation, we retained only those SVs present in 100% of the individuals within each scenario. In practice, this meant requiring SVs be detected in all 22 MD SR samples, or in all 57 WTD SR samples, or in all 4 WTD LR samples, or in all 3 MD LR samples, depending on the scenario. This filtering generated species-specific VCF datasets for both WTD and MD across the LR and SR data (see Figures 1–2 and Supplementary Data). Finally, we quantified the overlap of these species-specific SVs between LR and SR analyses within each scenario.

2.3. Gene Enrichment, Selection Analysis and Enhancer Motifs

We annotated the MD and WTD genome for enhancer motifs using Homer (Heinz et al., 2010). The findMotifsGenome, annotatePeaks, and scanMotifGenomeWide tools were employed and filtered based on the average motif scores. We extracted the number of motifs detected and compared the number in intergenic regions to the species-specific SVs; here, we normalized motif density, by dividing the

number of motifs by the length of each region to produce a value per kilobase pair (kbp) (Table S4). The resulting impact of SVs from both LR and SR analysis was annotated using SnpEff (Cingolani et al., 2012). To enable this, we created custom SnpEff databases using the genome FASTA files and their corresponding annotation GFF files. As with cross-referencing genome, we transferred annotations between assemblies using Mummer (Delcher et al., 2002) and custom Python scripts.

We performed GO term enrichment analyses only on those SV-affected genes that were fixed and detected by both SR and LR data (i.e. overlap); we extracted the coding sequences of genes directly impacted by these variants, with each sequence was queried using BLAST searches restricted to *Odocoileus* entries in UniProt and NCBI. For each gene, the best scoring hit was used to assign GO terms. We then applied DAVID (Sherman et al., 2024) to test for functional enrichment relative to the genome background. This approach allowed us to assess whether the SV-affected genes were statistically overrepresented in particular biological categories. We also calculated the ratio of nonsynonymous to synonymous substitution rates (dN/dS) for SV-specific genes overlapping SV results (Table S2-S3); to do so we collated additional genomes from Caribou (*Rangifer tarandus*), Elk (*Cervus canadensis*, JACTMD000000000), Moose (*Alces alces*, CAXWQY000000000), and domestic cattle (*Bos taurus*). After identifying homologous gene sequences in each species using BLAST (Camacho et al., 2008) the nucleotide sequences were aligned using MAFFT (Katoh et al., 2005) and converted to a codon alignment using PRANK (Löytynoja, 2014). The aligned codon sequences, along with a user-specified species tree (cow, (elk, (caribou, ((MD, WTD), moose)))), were analyzed using the adaptive Branch-Site Random Effects Likelihood (aBSREL) (Smith et al., 2015) method implemented in HyPHy (Kosakovsky et al., 2005). Evidence of selection was considered dN/dS > 1 or dN/dS < 1, and p-value < 0.05.

3. Results

3.1. SV exploration

Average sequencing coverage for the SR datasets was approximately 6X; the corresponding LR datasets showed variation in coverage, but all were > 7X (Table 2). Across all individuals in the SR merged SV datasets (i.e. 2/3 methods detected the SV) are shown in Figure 3. The average SV size for deletion was ≈19483 bps, duplication was ≈1477563 bps, insertion was ≈105 bps, inversion was ≈550336 bps and translocation was ≈469 bps; average individual SV zygosity was X%, Y, Z, A, B respectively. In the LR merged SV dataset across all individuals (i.e. 2/3 methods detected the SV) are shown in Figure 4. The average SV size for deletion was ≈605 bps, duplication was ≈12854 bps, insertion was ≈384 bps, inversion was ≈50411 bps and translocation was ≈417 bp; In the SR results, an average of 78.07% of deletions, 2.97% of duplications, 72.12% of insertions, 28.68% of inversions, and 3.75% of translocations were homozygous (Table S1). In the LR results, an average of 50.86% of deletions, 30.07% of duplications, 48.73% of insertions, 39.79% of inversions, and 10.84% of translocations were homozygous (Table S1).

3.2. Species specific SVs, Genome and Enhancer Impact

We next focused on species-specific SVs that were present in 100% of samples within each species in Scenarios 1,2,4,5 (Table 1). In total we identified 56 unique SVs in 100% of SR samples (Table 3). In contrast, we identified 123,384 unique SVs in 100% of LR samples (Table 4). The majority of gene-impacts were intronic (Table 3 and 4). We assessed the overlap between SR- and LR-derived SVs by considering only those variants located within exonic, intronic, or gene-level regions (Table 5); intergenic SVs were excluded from this comparison. Of note two of genes had dNdS values significantly deviating from 1 (Supplementary Data tables S2 and S3). Further, we identified notable impacted genes as it pertains to deer biology; SPMAP2 involved in Sperm microtubule associated protein and LOC110132624 involved in putative odorant cell receptor. For both MD and WTD, SVs were enriched for motif densities (Table S4).

Gene-level and gene-set enrichment analyses identified five clusters in MD and six in WTD (Tabel S4-S5). In MD, enriched categories included WD40-repeat proteins, TPR-repeat domains, ATP-binding/nucleotide-binding proteins, membrane-associated genes, and a smaller cluster associated with G-protein-coupled receptor (GPCR) signaling. In WTD, enriched categories included GTP-binding/P-loop NTPase proteins, homeobox transcription factors, Ig-like domains, membrane proteins, transcriptional regulators, and a small cluster related to GPCR signaling. Both species share some broad functional themes, but MD is more enriched for protein-interaction repeat domains whereas WTD show enrichment for regulatory gene families.

4. Discussion

The genomes of recent and hybridizing species provide a window into the mechanisms that drive biodiversity, adaptation, and reproductive isolation. We compared both short-read and long-read data on two closely related and hybridizing deer species, the white-tailed deer and mule deer, to identify SVs fixed between species. By integrating data from multiple LR and SR sequencing, we identified SVs within and between these species, their impact on genes, and general relationship to enhancer motifs. Genome analyses has shown negligible historical introgression between these species, despite contemporary hybridization (Kessler et al., 2023), suggesting that much of the observed genomic divergence reflects drift and isolation rather than ongoing exchange. This lack of gene flow complicates the identification of barrier loci, since patterns of divergence cannot be readily distinguished from neutral processes. While the presence of species-specific SVs reflects major evolutionary events, we acknowledge that fixed differences alone cannot demonstrate a causal role in reproductive isolation, as divergence also accumulates neutrally in the absence of gene flow. Nonetheless, the enrichment of SVs in genic and regulatory regions provides testable hypotheses and clues to species divergence and reproductive isolation.

Characteristics of SVs in *Odocoileus*

Our analysis revealed a predominance of insertions and deletions among SVs in deer. This finding is consistent with observations reported in other studies (Shi et al., 2023; Zhang et al., 2023), notably, Shi et al. (2023) identified deletions as the most frequent SV type in human populations. This pattern could be attributed to various mechanisms, including replication errors or unequal homologous recombination (Wold et al., 2023). Simmons et al. (2012) showed homozygous deletions often exhibit high penetrance; furthermore, the potential for homozygous deletions and insertions to stabilize in the genome could be linked to their lessened impact on recombination resulting in their tolerance and increased frequency. Berdan et al., (2024) showed that SVs, particularly when heterozygous, can suppress recombination, but homozygous deletions and insertions, by their nature, do not inherently disrupt the linear arrangement of genetic material, minimizing their impact on recombination. Duplications and translocations, particularly in their heterozygous form, can result in complex genomic rearrangements that disrupt gene order and heighten the risk of recombination errors. Zhang et al. (2021) discussed how gene duplications can drive speciation by disrupting gene function or fostering novel interactions. While the species-specific duplications were among the least common SV we detected (Table 3 and 4), these are worth further exploration, especially those impacting genic regions.

Most of the SV impacts we observed were intergenic, and similar trends were reported by Yang et al. (2024) and David et al. (2024). These studies, like ours, demonstrated that a significant portion of SVs localized to intergenic regions in sheep, goats, and wild bird populations. This dominance of intergenic impacts can be attributed to the vast amount of non-coding DNA within the genome; however, these regions also contain important regulatory elements, which comprise a significant portion of the mammalian genome (Villar et al, 2015). David et al. (2024) observed that a high percentage of deletions and inversions in intergenic regions were classified as 'modifiers,' suggesting they could have phenotypic effects. Our findings provide a similar perspective where 80.11% of deletions and 72.99% of inversions occurred in intergenic regions and 18.44% of intergenic deletions and 15.99% of intergenic inversions were annotated as modifiers, reiterating the potential consequential impacts SVs can have

outside genic regions. Moreover, the increased density of enhancers in SVs was seen regardless of location to a gene (Table S4), supporting the potential influence of intergenic space in the genome.

Species-species SVs, adaptation and potential incompatibilities

The identification of fixed SVs highlights potential connections to adaptive divergence. Recent studies in Eucalyptus species (Ferguson et al., 2024) showed the role of shared SVs in adaptation and speciation with significant number of SVs being enriched for genes involved in adaptive traits. Likewise, work on stick-insects showed clustering functionally important genes (Gompert et al. 2025). In deer, patterns were nuanced but did align with predictions and previous work. In mule deer, the SV impacted genes played a role in cellular differentiation, gene regulation, protein interactions, and metabolic processes, which could influence tissue development, immune responses, and overall cellular function, consistent with what Kessler et al. (2023) observed using genome outlier scans. These findings also align with previous studies demonstrating that SVs can drive evolutionary divergence by altering gene function rather than enhancer activity (Quan et al., 2021). Structural variants have been shown to impact gene expression through changes in chromatin organization and regulatory interactions, potentially facilitating adaptive traits without necessarily modifying enhancer density (Razin et al., 2023), though again we observed increase enhancer motifs in SVs (Table S4).

Several gene impacts were identified that align with known aspects of deer biology and presumed speciation (Kessler et al., 2024). Notably, a MD specific deletion impacted gene SPMAP2, which is involved in in male gonad development and sperm product; notably, hybrid deer show evidence of reduced sperm function (Derr et al., 1991). Similarly, both Bracewell et al. and Dowle et al. (2017) showed how SVs on neo-sex chromosomes drive hybrid male sterility, demonstrating that chromosomal rearrangements can underpin the genetic basis of speciation via genes directly involved in fertility. We also observed a WTD specific deletion impacting gene LOC110132624 which is linked to olfactory function, similar processes also were identified in the SNP based approach of Kessler et al. (2024). Differences in olfactory genes might facilitate ecological partitioning between WTD and MD,

contributing to their speciation. The role of olfactory genes in mate recognition and behavioral adaptation has been seen in house mice (North et al., 2020), where copy number variations in olfactory receptor genes influenced mate choice and reinforced reproductive isolation. This is particularly relevant in deer, as differences in olfactory responses may not only dictate ecological adaptations but also impact mate recognition, creating a dual mechanism for speciation through both environmental and sexual selection. Collectively, such fixed SVs and the corresponding phenotypic variation might implicate possible Dobzhansky-Muller incompatibilities in maintaining species boundaries in deer.

Further, several genes associated with species-specific SVs exhibited evidence of positive selection in both MD and WTD. These positively selected genes appear to be involved in key cellular and physiological processes (Table S2 and S3). The SV impacted genes with positive dN\dS values (2 total), were all either deletion or insertions; here we note a limitation in evaluating the evolutionary consequences of duplications lies in SV detection, as current SR-based tools identify but do not reconstruct the SV, with the absence of the SV in the assembly reflective of the common practice of purging duplications (Guan et al, 2020). As a result, dN/dS calculations are confined to the sequence of the initial (homologous) duplicated region, though LR approaches will allow for assess the evolution of duplicated regions. Further, some dN/dS values (Table S2 and S3) appear to be impacted by short-branches and limited substitutions, causing inflated values (Romiguer et al. 2012).

Differences between SV detection methods and future directions

The accurate estimation of SV size can be influenced by the specific tools and methodologies employed (Sedlazeck et al., 2018; Jeffares et al., 2017). The accuracy and sensitivity of SV callers vary significantly, as some tools are better suited for detecting larger SVs, while others excel at identifying smaller changes (Sedlazeck et al., 2018; Sudmant et al., 2015; Jeffares et al., 2017). This inherent bias across tools shapes the type of SV and the observed size distribution. To account for this, we employed two distinct data streams for SV detection to evaluate genome differences between MD and WTD, both reflective of commonly available datasets. Our comparison of SR and LR SV datasets shows differences

that shape how structural variation is represented in genomic studies. As demonstrated in earlier benchmarking and evolutionary studies (Pei et al., 2024; Mahmoud et al., 2019), LR technologies detect substantially more SVs across nearly all classes, and our results reflect this same pattern. This discovery is expected, as long reads traverse repetitive and low-complexity regions where most SV breakpoints occur, allowing callers to resolve thousands of smaller and intermediate events that SR platforms fail to detect. In contrast, the SR datasets captured markedly fewer SVs across all classes and exhibited patterns consistent with well-known short-read constraints (Mérot et al, 2022). The large SR sample size and requirement of 100% SV detection in all samples (both LR and SR), might be overly stringent, but provided confidence in the observed calls, especially those overlapping between data sets (Table 5). While the absolute counts between SR and LR calls differed (Table 3 and 4), any non-overlapping SV can be presumed a false-positive or false-negative, mostly likely occurring in the SR data. The varied scenarios (Table 1) also allowed for assembly artefacts to be removed, where the only explanation for SVs detected only in either WTD2WTD or MD2MD comparison is that assembly differs from the resequencing data. Ultimately, the combined tally of SVs (Table 5), reflect high-confidence calls, and are likely evolutionarily relevant.

Future research can build on our findings through several approaches aimed at confirming the role of SVs in speciation. Mouse models would offer experimental approaches to validate, but analyzing known hybrids and backcrosses in deer should allow assessing the distribution and inheritance patterns of SVs. RNA-seq combined with chromatin accessibility assays could allow us to measure how SVs and the enhancer-rich regions alter gene regulation in both species, with a particular focus on duplications found in one species and not the other. Lastly, expanding the scope to comparative analyses across other cervid species will further clarify whether similar classes of SVs repeatedly contribute to divergence and allow for reconstructing the ancestral state of species-specific SVs in *Odocoileus*.

Acknowledgements: We thank Alan Cain, Steeve Côté, Anh Dao, Marco Festa-Bianchet, Brad Fulk, Eric Hoffman, Levi Jaster, Daniel Koelsch, Emily Latch, Brent Patterson, Joe Nocera, Don Stewart, Russell Easy, Jonathan Shaw, David Walter, and Jon Wheeler for providing samples. We thank two reviewers for providing valuable feedback.

Data availability: All SR and some of LR sequencing samples are publicly available in the NCBI database under BioProject accession PRJNA830519. New LR data has been deposited in PRJAXXXXX. The code supporting this work is available in our lab's GitLab repository:

https://gitlab.com/WiDGeT_TrentU/graduate_theses/-/tree/master/AzimiChetabi/Chapter%20one

Author Contribution: We had no external contributions.

Funding: We were supported by Trent University's International Graduate Scholarship, an Ontario Graduate Scholarship, and Dean's PhD Scholarship. This work was supported by the Natural Sciences and Engineering Research Council of Canada Discovery Grant (grant number: RGPIN-2017-03934), ComputeCanada Resources for Research Groups (grant number: RRG gme-665-ab), Canada Foundation for Innovation: John R. Evans Leaders Fund, and Ontario Early Researcher Award (grant number: #36905).

Conflict of interest statement: The authors declare no competing interests.

References

Abril, V. V., E. A. G. Carnelossi, S. González, and J. M. B. Duarte. 2010. Elucidating the evolution of the red brocket deer Mazama americana complex (Artiodactyla; Cervidae). Cytogenetic and Genome Research 128:177–187.

Anaconda Software Distribution. (2023). Miniconda (Version 23.5.2) [Computer software]. Anaconda, Inc. Available at: https://docs.conda.io/en/latest/miniconda.html

Baurens, F.-C., G. Martin, C. Hervouet, et al. 2019. Recombination and large structural variations shape interspecific edible bananas genomes. *Molecular Biology and Evolution* 36(1):97–111.

Berdan, E. L., T. G. Aubier, S. Cozzolino, R. Faria, J. L. Feder, M. D. Giménez, M. Joron, J. B. Searle, and C. Mérot. 2024. Structural Variants and Speciation: Multiple Processes at Play. Cold Spring Harb Perspect Biol. 16:a041446.

Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics. 2014 Aug 1;30(15):2114-20. doi: 10.1093/bioinformatics/btu170. Epub 2014 Apr 1. PMID: 24695404; PMCID: PMC4103590.

Buchfink B, Reuter K, Drost HG. (2021). Sensitive protein alignments at tree-of-life scale using DIAMOND. Nature Methods, 18, 366-368. https://doi.org/10.1038/s41592-021-01101-x

Broad Institute. 2018. Picard Toolkit [GitHub repository]. Retrieved from http://broadinstitute.github.io/picard/

Bracewell, R. R., B. J. Bentz, B. T. Sullivan, and J. M. Good. 2017. Rapid Neo-Sex Chromosome Evolution and Incipient Speciation in a Major Forest Pest. Nat. Commun. 8:1593.

Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. BLAST+: architecture and applications. BMC Bioinformatics. 2009 Dec 15;10:421. doi: 10.1186/1471-2105-10-421. PMID: 20003500; PMCID: PMC2803857.

Chen, X., O. Schulz-Trieglaff, R. Shaw, B. Barnes, F. Schlesinger, M. Källberg, et al. 2016. Manta: Rapid detection of structural variants and indels for germline and cancer sequencing applications. Bioinformatics 32:1220–1222.

Cingolani, P., A. Platts, L. L. Wang, M. Coon, T. Nguyen, L. Wang, S. J. Land, X. Lu, and D. M. Ruden. 2012. A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster strain w1118; iso-2; iso-3. Fly 6:80–92.

Cursino, M. S., M. B. Salviano, V. V. Abril, et al. 2014. The role of chromosome variation in the speciation of the red brocket deer complex: The study of reproductive isolation in females. BMC Evolutionary Biology 14:40.

Danecek, P., J. K. Bonfield, J. Liddle, J. Marshall, V. Ohan, M. O. Pollard, A. Whitwham, T. Keane, S. A. McCarthy, R. M. Davies, and H. Li. 2021. Twelve years of SAMtools and BCFtools. GigaScience 10:giab008.

David, G., A. Bertolotti, R. Layer, D. Scofield, A. Hayward, T. Baril, H. A. Burnett, E. Gudmunds, H. Jensen, and A. Husby. 2024. Calling structural variants with confidence from short-read data in wild bird populations. Genome Biology and Evolution 16:evae049.

Delcher, A. L., A. Phillippy, J. Carlton, and S. L. Salzberg. 2002. Fast algorithms for large-scale genome alignment and comparison. Nucleic Acids Research 30:2478–2483.

Derežanin, L., J. Fickel, R. Tiedemann, and P. C. Alves. 2023. Contribution of structural variation to adaptive evolution of mammalian genomes. Universität Potsdam.

Derr, J. N., D. W. Hale, D. L. Ellsworth, and J. W. Bickham. 1991. Fertility in an F1 male hybrid of white-tailed deer (*Odocoileus virginianus*) x mule deer (*O. hemionus*). J Reprod Fertil. 93:111-7.

Dowle, E. J., R. R. Bracewell, M. E. Pfrender, K. E. Mock, B. J. Bentz, and G. J. Ragland. 2017. Reproductive Isolation and Environmental Adaptation Shape the Phylogeography of Mountain Pine Beetle (Dendroctonus Ponderosae). Mol. Ecol. 26:6071–6084.

Emms D.M., Kelly S. (2019). OrthoFinder: phylogenetic orthology inference for comparative genomics. Genome Biology, 20:238. https://doi.org/10.1186/s13059-019-1832-y

Escaramís, G., E. Docampo, and R. Rabionet. 2015. A decade of structural variants: Description, history, and methods to detect structural variation. Briefings in Functional Genomics 14:367–379.

Ferguson, S., A. Jones, K. Murray, R. L. Andrew, B. Schwessinger, H. Bothwell, J. Borevitz, 2024. Exploring the role of polymorphic interspecies structural variants in reproductive isolation and adaptive divergence in Eucalyptus. GigaScience 13:giae029.

Gompert, Z., J. L. Feder, T. L. Parchman, et al. 2025. Adaptation repeatedly uses complex structural genomic variation. *Science* 388(6744):eadp3745.

Gonzalez, S., and J. M. B. Duarte. 2020. Speciation, evolutionary history and conservation trends of Neotropical deer. *Mastozoología Neotropical* 27(SI):35–46.

Guan D, McCarthy SA, Wood J, Howe K, Wang Y, Durbin R. Identifying and removing haplotypic duplication in primary genome assemblies. Bioinformatics. 2020 May 1;36(9):2896-2898. doi: 10.1093/bioinformatics/btaa025. PMID: 31971576; PMCID: PMC7203741.

Heller, D., & Vingron, M. (2019). SVIM: Structural variant identification using mapped long reads. *Bioinformatics, 35*(17), 2907–2915. https://doi.org/10.1093/bioinformatics/btz041

Heinz, S., C. Benner, N. Spann, E. Bertolino et al. 2010. Simple Combinations of Lineage-Determining Transcription Factors Prime cis-Regulatory Elements Required for Macrophage and B Cell Identities. Mol Cell 38:576-589.

Homolka, D., R. Ivánek, J. Capkova, P. Jansa, and J. Forejt. 2007. Chromosomal rearrangement interferes with meiotic X chromosome inactivation. Genome Research 17:1431–1437

Howe, K., W. Chow, J. Collins, S. Pelan, D.-L. Pointon, Y. Sims, J. Torrance, A. Tracey, and J. Wood. 2021. Significantly improving the quality of genome assemblies through curation. *GigaScience* 10:1–9.

Jain, C., Rhie, A., Zhang, H., Chu, C., Walenz, B., Koren, S., & Phillippy, A. (2020). Weighted minimizer sampling improves long read mapping. Bioinformatics, 36(Suppl_1), i111–i118.

Jeffares, D. C., C. Jolly, M. Hoti, D. Speed, L. Shaw, C. Rallis, et al. 2017. Transient structural variations have strong effects on quantitative traits and reproductive isolation in fission yeast. Nature Communications 8:14061.

Kaplow, I. M., M. T. Maurano, A. Kundaje, et al. 2023. Relating enhancer genetic variation across mammals to complex phenotypes using machine learning. *Science* 380(6648):eabm7993.

Katoh, K., and D. M. Standley. 2013. MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. *Molecular Biology and Evolution* 30(4):772–780.

Kessler, C., and A. B. A. Shafer. 2023. Genomic analyses capture the human-induced demographic collapse and recovery in a wide-ranging cervid. bioRxiv. https://doi.org/10.1101/2023.07.19.

Kessler, C., E. Wootton, and A. B. A. Shafer. 2023. Speciation without gene-flow in hybridizing deer. Mol Ecol. 32:1117-1132.

Kirsche M, Prabhu G, Sherman R, Ni B, Battle A, Aganezov S, Schatz MC. Jasmine and Iris: population-scale structural variant comparison and analysis. Nat Methods. 2023 Mar;20(3):408-417. doi: 10.1038/s41592-022-01753-3. Epub 2023 Jan 19. PMID: 36658279; PMCID: PMC10006329.

Kosakovsky Pond, S. L., S. D. W. Frost, and S. V. Muse. 2005. HyPhy: Hypothesis testing using phylogenies. *Bioinformatics* 21(5):676–679.

Kosugi, S., & Terao, C. (2024). Comparative evaluation of SNVs, indels, and structural variations detected with short- and long-read sequencing data. *Human Genome Variation*, 11(1), 18. https://doi.org/10.1038/s41439-024-00276-x

Langmead, B., and S. L. Salzberg. 2012. Fast gapped-read alignment with Bowtie 2. Nature Methods 9:357–359.

Layer, R. M., C. Chiang, A. R. Quinlan, and I. M. Hall. 2014. LUMPY: A probabilistic framework for structural variant discovery. Genome Biology 15:R84.

Lovell, J. T., Sreedasyam, A., Schranz, M. E., Wilson, M., Carlson, J. W., Harkess, A., Emms, D., Goodstein, D. M., & Schmutz, J. (2022). GENESPACE tracks regions of interest and gene copy number variation across multiple genomes. *eLife*, 11, e78526.

Mahmoud, M., N. Gobet, D. I. Cruz-Dávalos, et al. 2019. Structural variant calling: the long and the short of it. Genome Biology 20:246.

Martin, G., F. Carreel, O. Coriton, C. Hervouet, C. Cardi, P. Derouault, D. Roques, F. Salmon, M. Rouard, J. Sardos, K. Labadie, F. C. Baurens, and A. D'Hont. 2017. Evolution of the banana genome (Musa acuminata) is impacted by large chromosomal translocations. Molecular Biology and Evolution 34:2140–2152.

Mérot, C., Stenløkk, K. S. R., Venney, C., Laporte, M., Moser, M., Normandeau, E., Árnyasi, M., Kent, M., Rougeux, C., Flynn, J. M., Lien, S., & Bernatchez, L. (2023). Genome assembly, structural variants, and genetic differentiation between lake whitefish young species pairs (Coregonus sp.) with long and short reads. *Molecular Ecology, 32*(6), 1458–1477.

North, H. L., P. Caminade, D. Severac, K. Belkhir, and C. M. Smadja. 2020. The role of copy-number variation in the reinforcement of sexual isolation between the two European subspecies of the house mouse. Phil. Trans. R. Soc. B 375:20190540.

Löytynoja A. Phylogeny-aware alignment with PRANK. 2014. Methods Mol Biol. 2014;1079:155-70. doi: 10.1007/978-1-62703-646-7_10. PMID: 24170401.

Pei, Y., Tanguy, M., Giess, A., Dixit, A., Wilson, L. C., Gibbons, R. J., Twigg, S. R. F., Elgar, G., & Wilkie, A. O. M. (2024). A comparison of structural variant calling from short-read and nanopore-based whole-genome sequencing using optical genome mapping as a benchmark. *Genes*, 15(7), 925. https://doi.org/10.3390/genes15070925

Pertea, M., G. M. Pertea, C. M. Antonescu, T.-C. Chang, J. T. Mendell, and S. L. Salzberg. 2015. StringTie enables improved reconstruction of a transcriptome from RNA-seq reads. Nature Biotechnology 33:290–295.

Quan C, Li Y, Liu X, Wang Y, Ping J, Lu Y, Zhou G. Characterization of structural variation in Tibetans reveals new evidence of high-altitude adaptation and introgression. Genome Biol. 2021 May 25;22(1):159. doi: 10.1186/s13059-021-02382-3. PMID: 34034800; PMCID: PMC8146648.

Quinlan, A. R., and I. M. Hall. 2010. BEDTools: a flexible suite of utilities for comparing genomic features. Bioinformatics 26:841–842.

Rausch, T., T. Zichner, A. Schlattl, A. M. Stutz, V. Benes, and J. O. Korbel. 2012. DELLY: structural variant discovery by integrated paired-end and split-read analysis. Bioinformatics 28:i333–i339.

Razin, S. V., S. V. Ulianov, and O. V. Iarovaia. 2023. Enhancer function in the 3D genome. *Genes* 14(6):1277.

Rieseberg, L. H. (2001). Chromosomal rearrangements and speciation. Trends in Ecology & Evolution, 16(7), 351–358. https://doi.org/10.1016/S0169-5347(01)02187-5

Romiguier, J., Figuet, E., Galtier, N., Douzery, E. J. P., Boussau, B., Dutheil, J. Y., & Ranwez, V. (2012). Fast and robust characterization of time-heterogeneous sequence evolutionary processes using substitution mapping. *PLOS ONE*, 7(3), 1–10. https://doi.org/10.1371/journal.pone.0033852

Sedlazeck, F. J., P. Rescheneder, M. Smolka, et al. 2018. Accurate detection of complex structural variations using single-molecule sequencing. Nature Methods 15:461–468.

Shang, H., D. L. Field, O. Paun, M. Rendón-Anaya, J. Hess, C. Vogl, J. Liu, P. K. Ingvarsson, C. Lexer, and T. Leroy. 2023. Drivers of genomic landscapes of differentiation across a Populus divergence gradient. Molecular Ecology 32:4348–4361.

Shen W, Le S, Li Y, Hu F (2016) SeqKit: A Cross-Platform and Ultrafast Toolkit for FASTA/Q File Manipulation. PLOS ONE 11(10): e0163962. https://doi.org/10.1371/journal.pone.0163962

Sherman, B. T., G. Panzade, T. Imamichi, and W. Chang. 2024. DAVID ortholog: An integrative tool to enhance functional analysis through orthologs. Bioinformatics. Advance online publication.

Shi, J., Z. Jia, J. Sun, et al. 2023. Structural variants involved in high-altitude adaptation detected using single-molecule long-read sequencing. Nat Commun 14:8282.

Simão F.A., Waterhouse R.M., Ioannidis P., Kriventseva E.V., Zdobnov E.M. (2015). BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics*, 31(19), 3210–3212. https://doi.org/10.1093/bioinformatics/btv351

Simmons, A. D., C. M. Carvalho, and J. R. Lupski. 2012. What have studies of genomic disorders taught us about our genome? Methods Mol Biol 838:1-27.

Smith, M. D., J. O. Wertheim, S. Weaver, et al. 2015. Less is more: An adaptive branch-site random effects model for efficient detection of episodic diversifying selection. *Molecular Biology and Evolution* 32(5):1342–1353.

Sobel, J. M., G. F. Chen, L. R. Watt, and D. W. Schemske. 2010. The biology of speciation. *Evolution* 64:295–315. https://doi.org/10.1111/j.1558-5646.2009.00877.x

SRA Toolkit Development Team. 2024. SRA Toolkit. Available from: https://trace.ncbi.nlm.nih.gov/Traces/sra/sra.cgi?view=software

Stevison, L. S., K. B. Hoehn, and M. A. Noor. 2011. Effects of inversions on within- and between-species recombination and divergence. *Genome Biology and Evolution* 3:830–841. https://doi.org/10.1093/gbe/evr081

Sudmant, P. H., T. Rausch, E. J. Gardner, R. E. Handsaker, A. Abyzov, J. Huddleston, Y. Zhang, et al. 2015. An integrated map of structural variation in 2,504 human genomes. *Nature* 526:75–81.

Tham, C., Tirado-Magallanes, R., Goh, Y. *et al.* NanoVar: accurate characterization of patients' genomic structural variants using low-depth nanopore sequencing. *Genome Biol* **21**, 56 (2020). https://doi.org/10.1186/s13059-020-01968-7

Villar, D., C. Berthelot, S. Aldridge, et al. 2015. Enhancer evolution across 20 mammalian species. *Cell* 160(3):554–566.

Wang Y, Tang H, Debarry JD, Tan X, Li J, Wang X, Lee T-H, Jin H, Marler B, Guo H, Kissinger JC, Paterson AH. (2012). MCScanX: a toolkit for detection and evolutionary analysis of gene synteny and collinearity. Nucleic Acids Research, 40(7), e49. https://doi.org/10.1093/nar/gkr1293

Wellenreuther M, Mérot C, Berdan E, Bernatchez L. Going beyond SNPs: The role of structural genomic variants in adaptive evolution and species diversification. Mol Ecol. 2019; 28: 1203–1209. https://doi.org/10.1111/mec.15066

Wold, J. R., J. G. Guhlin, P. K. Dearden, A. W. Santure, and T. E. Steeves. 2023. The promise and challenges of characterizing genome-wide structural variants: A case study in a critically endangered parrot. Mol Ecol Resour.

Yang L. A Practical Guide for Structural Variation Detection in the Human Genome. Curr Protoc Hum Genet. 2020 Sep;107(1):e103. doi: 10.1002/cphg.103. PMID: 32813322; PMCID: PMC7738216.

Yang, J., D. F. Wang, J. H. Huang, et al. 2024. Structural variant landscapes reveal convergent signatures of evolution in sheep and goats. Genome Biology 25:148.

Zhang, L., S. Chaturvedi, C. C. Nice, L. K. Lucas, and Z. Gompert. 2023. Population genomic evidence of selection on structural variants in a natural hybrid

Figure caption:

Figure 1. General workflow of this study depicted on the right. The map shows the geographical range of the two species under investigation. The light grey area represents the range of white-tailed deer, while the grey area indicates the range of mule deer. The dark grey region highlights the overlap between the two species' habitats. Each point indicates the specific locations where samples were collected.

Figure 2. Macrosynteny comparison among white-tailed deer (*Odocoileus virginianus*), mule deer (*Odocoileus hemionus*), caribou (*Rangifer tarandus*), and domestic cattle (*Bos taurus*). Chromosome segments are connected by colored ribbons, showing conservation of chromosome structure and highlighting the close genomic relationship between white tailed and mule deer.

Figure 3. Comparison of the number of structural variants (SVs) detected across SV types in the entire short read dataset. Each panel represents a different mapping strategy, showing the relative counts of deletions, duplications, insertions, inversions, and translocations on a log scale.

Figure 4. Comparison of the number of structural variants (SVs) detected across SV types in the entire long read dataset. Each panel represents a different mapping strategy, showing the relative counts of deletions, duplications, insertions, inversions, and translocations on a log scale.

Table captions :

Table 1. Summary of the mapping scenarios used to identify species-specific structural variants (SVs) in both short-read and long-read datasets. WTD and MD refer to white-tailed deer and mule deer, respectively. The first four columns indicate the mapping strategy for each dataset—that is, which species' samples were aligned to which reference genome, with 1 meaning all samples have the SV. For example, "WTD2WTD" represents white-tailed deer samples mapped to the white-tailed deer reference genome. Last column indicates if the SVs derived from that strategy were considered species-specific (Y) or no (N).

	WTD2WTD	**WTD2MD**	**MD2WTD**	**MD2MD**	**SPECIES_SV**
SCENARIO1	1	1	0	0	Y
SCENARIO2	0	0	1	1	Y
SCENARIO3	1	0	0	0	N
SCENARIO4	0	1	0	0	Y
SCENARIO5	0	0	1	0	Y
SCENARIO6	0	0	0	1	N

Table 2. Raw sequencing statistics for all long-read samples from white-tailed deer (WTD) and mule deer (MD). For each dataset, the total number of reads (No. reads), minimum read length (Min. length), average read length (Avg. Length), and maximum read length (Max length) are reported.

Sample	**Species**	**No. reads**	**Min. length**	**Avg. Length**	**Max length**
bc2001	WTD	468,805	215	11,831.10	38,337
bc2065	MD	1,820,410	51	11,292.10	52,827
bc2066	WTD	1,144,838	76	13,229.60	47,420
bc2069	WTD	1,735,556	64	12,179.30	41,157
bc2070	MD	1,909,554	77	12,406.30	42,022
SRR15720448	MD	15,003,446	50	19,657.10	256,018
SRR6668252	WTD	89,770,713	11	5,093	217,085

Table 3. Species-specific structural variants (SVs) detected in short-read dataset across the scenarios defined in Table 1, using the criterion that each SV must be present in 100% of samples. For each scenario, the table reports: (i) the total number of SVs; (ii) counts of each SV class (DEL, DUP, INS, INV, TRA); (iii) the number of genes overlapping those SVs; and (iv) the distribution of gene impacts, categorized as exonic, intronic or gene-level. SV types are abbreviated as follows: DEL = deletion, DUP = duplication, INS = insertion, INV = inversion, TRA = translocation.

		SR - 100% of samples		
	# SVs	**SV Types**	**# Genes**	**Gene Impact**
Scenario1	11	DEL:0 DUP:0 INS:3 INV:0 TRA:8	11	0\3\8
Scenario2	10	DEL:1 DUP:0 INS:0 INV:1 TRA:8	9	0\0\9
Scenario4	4	DEL:2 DUP:0 INS:1 INV:0 TRA:1	3	0\0\3
Scenario5	31	DEL:13 DUP:6 INS:0 INV:2 TRA:10	17	4\1\12

Table 4. Species-specific structural variants (SVs) detected in the long-read dataset across the scenarios defined in Table 1, using the criterion that each SV must be present in 100% of samples. For each scenario, the table reports: (i) the total number of SVs; (ii) counts of each SV class (DEL, DUP, INS, INV, TRA); (iii) the number of genes overlapping those SVs; and (iv) the distribution of gene impacts, categorized as exonic, intronic, or gene-level. SV types are abbreviated as follows: DEL = deletion, DUP = duplication, INS = insertion, INV = inversion, TRA = translocation.

LR - 100% of samples

	# SVs	SV Types	# Genes	Gene Impact
Scenario1	19280	DEL:8341 DUP:0 INS:10801 INV:138 TRA:0	4371	241\4127\3
Scenario2	14552	DEL:7911 DUP:99 INS:4156 INV:217 TRA:2169	6220	426\3570\2224
Scenario4	23074	DEL:14646 DUP:0 INS:8381 INV:47 TRA:0	2316	79\2050\187
Scenario5	66478	DEL:37002 DUP:0 INS:29295 INV:181 TRA:0	12134	1307\10716\111

Table 5. Species-specific structural variants (SVs) detected in both long and short read datasets across the scenarios defined in Table 1. For each scenario, the table reports: (i) the total number of SVs; (ii) counts of each SV class (DEL, DUP, INS, INV, TRA); (iii) the number of genes overlapping those SVs; and (iv) the distribution of gene impacts, categorized as exonic, intronic, or gene-level. SV types are abbreviated as follows: DEL = deletion, DUP = duplication, INS = insertion, INV = inversion, TRA = translocation. The number in parenthesis is the number of genes with significant dNdS values.

Overlap structural variants

	# SVs	SV Types	# Genes	Gene Impact
Scenario1	4	INS:1 TRA:3	4	0\1\3
Scenario2	5	TRA:5	5 (1)	0\0\5
Scenario4	3	DEL:2 TRA:1	3	0\0\3
Scenario5	15	DEL:3 DUP:2 INV:2 TRA:8	15 (1)	4\1\10