Optimised genome editing for precise DNA insertion and substitution using Prime
Editors in zebrafish

Yosuke Ono[1], Martin Peterka[2], Michael Love[3], Ashish Bhandari[1], Euan Gordon[2], Jonathan S.
Ball[3], Charles R. Tyler[3], Steve Rees[4], Mohammad Bohlooly-Y[2], Marcello Maresca[2], and
Steffen Scholpp[1*]

[1]Living Systems Institute, Biosciences, University of Exeter, United Kingdom; [2]Discovery
Sciences, BioPharmaceuticals R&D, AstraZeneca, Gothenburg, Sweden; [3]Biosciences,
University of Exeter, United Kingdom; [4]Discovery Sciences, BioPharmaceuticals R&D,
AstraZeneca, Cambridge, United Kingdom

12
*Correspondence: s.scholpp@exeter.ac.uk

Abstract

CRISPR/Cas9-mediated genome editing has rapidly become a popular tool for studying
gene functions and generating genetically modified organisms. However, using this system,
stochastic integration of random insertions and deletions restricts precise genome
manipulation. Advanced CRISPR/Cas9 technologies using Prime Editors (PEs), Cas9
proteins fused with reverse transcriptase, enable programmed integration of short DNA
modifications into the genome. However, its application in precise genome editing in animal
models is challenging. Here, we utilise a nickase- and a nuclease-based PE to perform
programmed short DNA substitution and insertion in various loci in the zebrafish genome.
Whereas the nickase-based PE is advantageous for nucleotide substitutions, we find that the
nuclease-based PE can be used to insert short DNA fragments precisely with high efficiency.
To further evaluate our approach, we inserted a nuclear localisation signal into a reporter
transgene to incorporate longer fragments by prime editing. These gene modifications were
transmitted to the next generation. We demonstrate that PE-mediated prime editing can
efficiently manipulate genome information in zebrafish without using exogenous donor DNA.

Keywords

Prime editing, CRISPR/Cas9, zebrafish, PE2, PEn

Introduction

Genetically modified animals serve as powerful tools for studying ontogenesis and disease mechanisms in complex living systems. The zebrafish, *Danio rerio,* is a well-established model vertebrate species in cell and developmental biology due to attributes including its high fecundity in laboratory environments, transparency of the embryos, and amenability to the application of tools for transgenesis[1–3]. Zebrafish have also emerged as a powerful vertebrate model organism for studying human diseases, as they exhibit a substantial degree of genetic similarity with humans (around 70% of human genes having a corresponding gene in zebrafish)[4]. Several technologies for targeted mutagenesis, such as zinc finger nuclease (ZFN) and transcription activator-like effector nuclease (TALEN), have been employed in zebrafish to establish genetic loss-of-function mutants[5–8]. Most recently, CRISPR/Cas9-mediated genome editing has been introduced, significantly advancing target mutagenesis in nearly every model system, including zebrafish[9–13]. The CRISPR/Cas9 system has the potential to mimic disease phenotypes more accurately, allowing for the creation of identical genetic alterations corresponding to disease-causing mutations in humans. However, it has the disadvantage of stochastically generating insertions and deletions (so-called indels). This is because this technology relies on the DNA double-strand break (DSB) induced by the endonuclease at the target sequence in the genome and subsequent DNA repair. To allow precise control of genome editing, homology-directed repair (HDR)-mediated knock-in approaches using exogenous donor DNA have been utilised in zebrafish[14–16]. However, HDR-mediated precise genome editing occurs less efficiently compared with random mutagenesis.

Due to the need for precise genome editing with high efficiency, new strategies have been employed and applied in model organisms. For example, CRISPR/Cas9-mediated precise editing using Prime Editor has recently been developed, which does not rely on exogenous DNA donors. The original Prime Editor is a fusion protein of Cas9-nickase and reverse transcriptase[17]. By combining the Prime Editor with a prime editing guide RNA (pegRNA), the resulting Prime Editor ribonucleoprotein (RNP) complex generates a single-strand break (SSB) or nick at the target site in the genome. The reverse transcriptase domain elongates an additional DNA sequence at the 3' end of cleaved DNA, following the reverse-transcription (RT) template sequence incorporated in the pegRNA. The additional short DNA fragment that contains the intended edit is integrated into the genomic DNA through the ligation of the 3' flap and the excision of the competing 5' flap from the original DNA sequence[17]. The

Cas9-nickase-based Prime Editor enables programmed short DNA modification (insertion,

deletion and substitution), as has been shown in cultured cells, mouse and zebrafish

embryos[17–20]. An alternative to the Cas9-nickase-based Prime Editor is prime editing with a

Cas9-nuclease-based Prime Editor (PEn)[21–23]. PEn facilitates programmed short DNA

modification through DNA repair based on homology annealing and non-homologous end

joining (NHEJ) after DSB[22]. However, evidence for the successful use of Prime Editors in

creating genetically modified animals is still limited. This is due to various technical

challenges, including high variability in editing modes, low editing efficiency, and different

delivery methods for prime editing components in living animals.

Here, we assess PE2, a nickase-based Prime Editor, and PEn-mediated prime editing for

establishing genetically modified zebrafish for programmed DNA modification in their

genome. We tested nucleotide substitution and the insertion of different lengths of DNA

sequences, including a stop codon and nuclear localisation signal (NLS). We performed

morphological assessments of these genetically modified embryos. We demonstrate that PE2-

mediated prime editing is a favourable tool for base pair substitution in zebrafish embryos,

whereas editing with PEn can more efficiently insert nucleotides up to 30 bp for modulating

protein function and behaviour, all without using exogenous donor DNA. We further show

that stable genetically modified zebrafish lines can be generated that inherit these

programmed short DNA modifications.

Results

Prime editing in nucleotide substitution

First, we tested the use of Prime Editors for substituting specific base pairs in the zebrafish

genome. To do this, we focused on the Cereblon (CRBN) gene, which has been associated

with developing resistance to thalidomide-based treatments, particularly in conditions like

multiple myeloma[24,25]. Here, we first compared the functioning of nuclease-based PEn and

nickase-based PE2 in nucleotide substitution in the zebrafish *crbn* gene (Fig. 1a), targeting

the sequence encoding the amino acid residue I378 associated with the sensitivity to

thalidomide and related drugs[26,27]. A mixture of Prime Editor proteins and chemically

synthesised pegRNAs for substituting 2 single nucleotides in the target sequence (+3 G to C

and +10 A to G; Supplementary Fig. 1a) was injected into zebrafish embryos at the early 1-

cell stage, after which the embryos were incubated at 32°C. Genomic DNA was then

extracted from these embryos at 96 hours post-fertilisation (hpf) to analyse genome edits at

the target site. Amplicon sequencing of the target region showed that both PEn and PE2

integrated the prime editing substitution into the genome (Fig. 1b). We found that PE2 had a

higher efficiency in precise substitutions (8.4%) compared to PEn (4.4%, Fig. 1c).

Furthermore, PEn induced a higher amount of indels (Fig. 1d). The precision score, defined

as the ratio of precise prime edits to the total edits, including imprecise prime edits and indels,

was significantly higher with PE2 (40.8%) compared to PEn (11.4%, Supplementary Fig. 1b).

We also tested the refolding procedure of pegRNA to prevent misfolding between the

complementary sequences in the spacer sequence and primer binding site (PBS) and RT

template sequence[28]. Refolding of the *crbn* pegRNA did not enhance prime editing

substitution with both PEn and PE2 (Fig. 1c). These results indicate that PE2 is the more

effective for very short nucleotide substitution compared with PEn in zebrafish.

Prime editing insertion of 3 bp DNA fragment

Next, we tested the functioning of the Prime Editors in the precise insertion of a 3 bp stop

codon into the coding sequence of the endogenous target gene to generate a mutant allele that

produces a precisely truncated protein. As a proof-of-concept, we chose the cognate receptor

gene *ror2*. Genetic mutations in the *ROR2* gene in humans can cause the autosomal recessive

Robinow syndrome, leading to short stature, distinctive facial features, and skeletal

abnormalities, including short limbs and a curved spine (*scoliosis*)[29–31]. In zebrafish, the

inhibition of Ror2 signalling function affects the convergence and extension of axial cells

during gastrulation and the elongation of the embryo body, leading to a broader and shorter

body axis[32–35], as well as the patterning of cranial tissues[36], and thus would be an excellent

model for studying many aspects of the Robinow syndrome.

Sequence comparison suggested that the zebrafish W722X allele corresponds to the

human disease-related W720X mutation generating a premature stop codon (TGA) in the

tyrosine kinase domain (Fig. 2a). Therefore, we aimed to establish the zebrafish Robinow

W722X model by designing pegRNA and single primed insertion gRNA (springRNA)[22]

generating a similar stop codon into the sequence of exon 9 in zebrafish *ror2*. The pegRNA

contained a 3 nt RT template sequence for the stop codon to be integrated, 13 nt PBS

sequence, and 13 nt sequence to extend the homology arm for the DNA integration via

homology annealing (Fig. 2b and c). We compared this strategy to using a springRNA

combined with PEn, without the template sequence for the homology arm, to insert the stop

codon via NHEJ (Fig. 2b and c). We microinjected combinations of Prime Editor mRNA and

guide RNA, that is, PE2/pegRNA, PEn/pegRNA and PEn/springRNA at the one-cell stage in

zebrafish embryos. The sequence in *ror2* exon 9 was amplified by PCR from a pool of

genomic DNA obtained from 10 injected embryos, and the induction of DNA modification

was assessed by a T7 endonuclease I (T7E1) assay. We observed the cleavage of PCR

products in the samples injected with PEn/pegRNA and PEn/springRNA combinations,

indicating sequence modification in the target site (Fig. 2d). In contrast to the base pair

substitution, PE2 was less effective.

The PCR products were also cloned into a cloning vector, and the target sequences in

randomly selected clones were analysed. Various types of edits in the target site were

observed in the clones from the PEn injected samples, including precise insertion of the stop

codon and/or random indels (Fig. 2e), although no sequence edits in the target site were

identified from the PE2 injected sample (0/20 clones; Supplementary Fig. 2). Subsequently,

we performed amplicon sequencing to quantify the proportion of each genome edit. We

found that the proportion of precise prime editing (only TGA insertion without any indels)

was higher when using PEn/pegRNA combination (10.3%) compared to PEn/springRNA

(4.0%) and PE2/pegRNA (0.4%, Fig. 2f). In addition, springRNA showed a higher proportion

of imprecise prime edit (TGA insertion with random indels; 37.5%) than pegRNA (11.1%

and 0.2% with PEn and PE2, respectively; Fig. 2f). The precision scores for the PE2/pegRNA,

PEn/pegRNA, and PEn/springRNA combinations were 58.9%, 32.6%, and 7.8%, respectively.

This suggests that prime editing via homology annealing with pegRNA was more accurate

than via NHEJ using springRNA.

Comparison of the mode of Prime Editor delivery as mRNA or RNP complex

Next, we compared PEn and PE2 in combination with different strategies of Prime Editor

delivery. Here, we performed amplicon sequencing with genomic DNAs extracted from

individual embryos injected with Prime Editor mRNA or RNP complex with the *ror2*

pegRNA (Fig. 3a). We found that the proportion of precise prime edit is significantly

increased with PE2 RNP complex (11.5%) compared to PE2 mRNA (1.3%), whereas the

there is no significant difference between PEn RNP complex (20.5%) compared to PEn

mRNA (18.8%; Fig. 3b). However, with both delivery methods of the Prime Editors, the

proportion of precise prime edit was significantly higher with PEn compared to PE2,

supporting our previous results (Fig. 2f and 3b). Furthermore, PEn also integrated a notably

higher number of indels that are unrelated to prime editing (Fig. 3c), resulting in lower

precision scores with PEn (34.3% and 28.2% for mRNA and RNP delivery, respectively)

compared to PE2 (71.6% and 43.4% for mRNA and RNP delivery, respectively;

Supplementary Fig. 3a). These findings suggest that the PEn-induced DNA DSBs are more

like to undergo repair events (both precise and imprecise) compared to the nickase-based PE2

method.

Induction of abasic spacer into springRNA to prevent gRNA scaffold insertion

In the analysis of the sequence edits in the *ror2* gene by PEn-mediated prime editing, we

found that a substantial proportion of the amplicons derived from embryos injected with the

PEn/springRNA combination contained imprecise prime edits with additional nucleotides

that correspond to the sequence of gRNA scaffold adjacent to the RT template (Fig. 3d and

Supplementary Fig. 3d), as reported previously[18]. This finding suggests that the readthrough

of RT from the template sequence into the gRNA scaffold causes unintended insertions, as

has also been reported in human cell lines[17,37]. Aiming to control the termination of RT in

prime editing, we synthesised a modified springRNA in which an abasic RNA spacer was

inserted between the RT template and gRNA scaffold (Fig. 3e). Microinjection of the abasic

springRNA together with PEn mRNA or RNP complex significantly reduced the unintended

insertion of gRNA scaffold sequence into the target site (Fig. 3f). However, the proportion of

precise prime edits did not show a significant improvement (Fig. 3g). Due to the significantly

reduced number of the imprecise prime edits, the precision scores increased from 9.6% to

21.7% and from 10.6% to 14.2% with PEn mRNA and RNP complex, respectively, through

the blocking of scaffold sequence incorporation (Supplementary Fig. 3b). These results

suggest that the read-through process is independent of precise genome editing events in

zebrafish embryos, unlike in human cultured cells[37].

Prime editing in adgrf3b gene using PEn

Next, we tested the PEn-mediated prime editing targeting at a different locus, here the

adgrf3b gene. A previously published study has reported that the efficiency of prime editing

in the *adgrf3b* gene using PE2 and pegRNA was decreased from 18.0% to 0.1% by extending

the length of prime editing insertion from 3 bp to 12 bp[18]. Therefore, we tested the efficiency

of our nuclease-based PE by injecting PEn mRNA with pegRNA or springRNA targeting the

same sequence in the *adgrf3b* gene to integrate the 3 bp or 12 bp insertion. Both

PEn/pegRNA and PEn/springRNA combinations successfully integrated the programmed 3

bp and 12 bp insertion into the target site in the injected embryos. Here, the mean proportion

of precise prime edit with pegRNA and springRNA in amplicon sequencing was 27.3% and

4.0% for 3 bp insertion and 1.9% and 0.7% for 12 bp insertion, respectively (Fig. 3h and i).

The precision scores with pegRNA and springRNA were 42.1% and 4.6% for a 3 bp insertion

and 8.6% and 1.1% for a 12 bp insertion, respectively (Supplementary Figure 3c), suggesting

that both the efficiency and accuracy of prime editing declined with an increase in insertion

length. Both PEn/pegRNA and PEn/springRNA combinations performed more efficiently

with the 12 bp insertion compared to that reported previously using the PE2 system, this

suggests that PEn-generated DNA DSB accepts longer insertion than a nickase-based Prime

Editor in zebrafish embryos.

Establishing stable ror2[W722X] *mutant zebrafish*

Having demonstrated the programmed insertion via PEn-mediated prime editing in

injected embryos, we wanted to establish if these precise edits are passed on to the next

generation, as this is a prerequisite for establishing a disease model. We, therefore, aimed to

develop stable *ror2*[W722X] mutant zebrafish. Therefore, first, we injected embryos with PEn

RNP complex and the *ror2* springRNA and raised the embryos to adults, and then we

outcrossed them with wild-type fish to collect F1 embryos. Through genotyping 16 embryos

from each F0 adult fish, we obtained F1 embryos harbouring the *ror2*[W722X] mutation. Six out

of 10 F0 adults were founders of the allele, with the germline transmission rate varying

between 6.3% (1/16) and 31.3% (5/16; Fig. 4a). We raised the F1 heterozygous *ror2*[W722X]

fish and inbred them to generate homozygous *ror2*[W722X] mutant embryos in the F2 generation.

The morphology of the zygotic *ror2*[W722X] mutant embryo and larvae was similar to wild-type

siblings, and the mutant animals were both viable and fertile (Fig. 4b, c and e). Subsequently,

we generated maternal-zygotic (MZ) mutants by in-crossing homozygous *ror2*[W722X] male and

female fish. The MZ *ror2*[W722X] mutant larvae at 5 dpf showed a mild curvature of the tail tip

and shorter body length (Fig. 4d and f), which are similar phenotypes to the MZ mutant

larvae of the loss-of-function alleles in *ror2* reported previously[32,36]. This suggests that the

human disease-related *ror2*[W722X] mutation in the tyrosine kinase domain affected the

Wnt/PCP signalling similarly to the observation in patients with Robinow syndrome. This set

of data suggests that PEn-based editing can be used to establish stable genetically modified

fish lines.

Insertion of nuclear localisation signal into transgenic reporter gene

After the successful insertion of the 3 bp stop codon into the *ror2* gene to establish a

zebrafish model of the Robinow syndrome and successful 3 and 12 bp insertions into the

adgrf3b gene, we tested whether PEn-mediated prime editing can be used to integrate up to

30 bp. Since we anticipated that these events would be rare, we employed a strategy that

enabled us to screen for prime edit events and distinguish between imprecise prime edits and

random indels at the cellular level in living zebrafish embryos, all without the need for time-

consuming sorting and genetic analysis. This was achieved using an eGFP-transgenic fish

line with an introduced nuclear localisation sequence (NLS) of 27 bp using PEn. We

anticipated three phenotypes: cytosolic GFP localisation for unsuccessful editing, no GFP

expression for imprecise edits, and nuclear GFP expression for precise prime editing insertion

(Fig. 5a). We subsequently mapped these effects in individual cells using high-resolution

confocal microscopy in living zebrafish larvae. Specifically, we tested the insertion of the

NLS sequence into the *smyhc1:gfp* transgene by prime editing. The *smyhc1:gfp* reporter gene

is expressed explicitly in the superficial slow-twitch muscle fibres in the body trunk

(Supplementary Fig. 4a), which are mononuclear during embryonic development[38]. We

designed pegRNAs that target the N-terminal end of the eGFP coding sequence in the

smyhc1:gfp[i104] transgenic allele and contain RT template sequence for inserting c-myc

NLS[39–41]. We expected that precise integration of the 30 bp sequence, including 27 bp c-myc

NLS and 3 bp spacer, translocates the eGFP expression from the cytoplasm to the nucleus in

the muscle fibres, whereas reporter expression is lost in the cells with imprecise prime edits

or random indels by frameshifting in the eGFP coding sequence (Fig. 5a). The embryos

injected with PEn RNP complex showed mosaic expression of eGFP in the cytoplasm of

muscle fibres, lack of GFP expression in some fibres along with the accumulation of eGFP

signals in a few cell nuclei (Fig. 5b). Amplicon sequencing of the target region in the genome

showed that the average efficiency of precise prime editing insertion in the whole injected

embryo was 8.5% with a precision score of 9.7% (Fig. 5c). We also compared the prime

editing efficiency to that using a longer pegRNA in which PBS sequence and template

sequence of the homology arms were extended from 10 nt to 13 nt and 14 nt to 36 nt,

respectively. However, the prime editing efficiency of the longer pegRNA was significantly

decreased, possibly due to increased RNA complexity leading to altered secondary structures

(Fig. 5c).

Next, to establish stable genetically modified zebrafish lines with the NLS insertion, we

raised F0 injected fish to adults, outcrossed them with wild-type, and screened the F1

embryos for their GFP expression. We successfully isolated embryos showing pronounced

eGFP expression in the nuclei of slow-twitch fibres (Fig. 5d and Supplementary Fig. 4b). We

confirmed the integration of the 30 bp sequence into the eGFP coding sequence (Fig. 5e and

Supplementary Fig. 4c). Notably, multiple types of edits were also identified in the amplicons

obtained from the single embryo with nuclear eGFP expression, suggesting that the

smyhc1:gfp[i104] transgenic allele contains at least 4 copies in this locus. Each copy was

targeted and edited individually by PEn/pegRNA complex (Fig. 5e and Supplementary Fig.

4c). Here, 4 F0 adult fish out of 6 screened were identified as founder fish of the prime

editing insertion in the target transgene. The proportion of the embryos showing nuclear

eGFP expression varied between 1.4 to 20.2% (Fig. 5f). Our findings indicate that PEn-

mediated precise DNA insertion is a practical tool for the accurate integration of functional

motifs with a length of up to 30 bp within a target gene in zebrafish.

Discussion

The need for precise genome editing

Target sequence-specific mutagenesis and transgenesis are essential tools for analysing

gene functions, and labelling expressed proteins, cellular components and specific cell types

in vivo. Induction of DSB in the genome DNA by ZFN, TALEN and CRISPR/Cas9

technologies enable random mutagenesis and donor DNA-based knock-in in the target

sequence. In addition to the conventional HDR-mediated knock-in, multiple systems for

targeted integration into the genome in zebrafish have been reported[42,43]. Adding further to

this, Cas9-nickase fusion proteins with cytidine or adenine deaminase, so-called Base

Editors[44,45], enable single nucleotide substitution in zebrafish to generate human disease

models[46–51]. Despite advancements in gene editing technologies, however, the application of

Prime Editors for the generation of genetically modified animals remains restricted[18,52,53].

This limitation can be attributed to several technical hurdles, including substantial variability

in editing modes, suboptimal editing efficiency, and complications associated with the

delivery of prime editing components *in vivo*.

In this study, we established programmed short DNA modification in the genome using

Prime Editors and generated stable genetically modified zebrafish lines with 3 and 30 bp

insertions. Firstly, we utilised PE2 for successful nucleotide substitution in a target sequence

associated with drug sensitivity and PEn to generate a zebrafish model for human genetic

disease. Specifically, we established a stable $ror2^{W722X}$ mutant line, which genocopies the

disease-related W720X mutation in the human *ROR2* gene. Various mutations have been

identified in the human *ROR2* gene as the causes of Robinow syndrome, and different types

of mutation lead to distinct phenotypes and inheritance patterns, possibly due to nonsense-

mediated decay of mutated mRNA[54], the mechanism which is also essential for embryonic

development and tissue homeostasis in zebrafish[55,56]. Establishing genetically modified

zebrafish carrying mutations related to genetic disease in humans will enable a more precise

analysis of the mechanisms of genetic diseases. Using prime editing and base editing by Base

Editors will open a wider window for precise genome editing in zebrafish for both drug

discovery and human disease models.

Secondly, we inserted a functional motif in existing lines to control the features of a

protein. Here, specifically, we inserted an NLS sequence into the *smyhc1:gfp*[i104] transgene

and assessed the prime editing *in vivo* by translocating eGFP expression. The same approach

can be applied to tag endogenous proteins and control their localisation and transport. The

pegRNAs we used to insert NLS were designed in the coding sequence of the N-terminal

region of eGFP, and therefore, the strategy can be used for other established eGFP reporter

zebrafish lines to convert their GFP expression from the cytoplasm to the cell nuclei and as a

control to assess prime editing *in vivo* morphologically.

Previous studies have reported a low efficiency of prime editing insertion. In those studies

it is proposed that the length negatively correlates with prime editing efficiency, while the

MLH1 protein influences editing outcomes for short sequences up to 13 bp in cultured

cells[57–59]. Enhancing pegRNA stability and preventing degradation can further improve

editing efficiency[28,60,61]. Finally, co-expressing effector proteins with Prime Editor to

suppress undesired DNA repair pathways or protect pegRNA from degradation has been

suggested to enhance editing[23,58,62].

The underlying mechanism of PEn-mediated precise genome editing

Here, we demonstrated that a nuclease-based PE can efficiently integrate DNA insertions

between 3–30 bp in zebrafish embryos. By utilising pegRNA and springRNA, we show that

the integration is based on both homology annealing and NHEJ, as suggested in cell culture[22].

Although pegRNA was superior to springRNA in both efficiency and accuracy at the tested

target loci, springRNA could be advantageous when longer PBS and/or RT template

sequences are necessary, as it lacks a template for homology arms. We also observe that the

repair outcome varies depending on the delivery mode of the PEn. Microinjection of PEn

mRNA leads to fewer indels than supplying the PEn as an RNP complex, although there was

no significant difference in the level of precise prime edit (Fig. 3). We hypothesise that the

RNP complex acts faster than the mRNA and a potential explanation could be that the

relatively rapid integration of random indels is more effective than the prime editing reaction

in the early zebrafish embryo (0–3 hpf), when cell cycle time is significantly shorter

(approximately 25 min). However, the precise DNA repair mechanism and accessibility to a

target site in the genome for the prime editing reaction in relation to the cell cycle duration

need to be explored in the future.

We also show that nuclease-based PE has its limitations. For example, we found that the

nickase-based PE2 seems to be a favourable tool for nucleotide substitution, as shown for

crbn (Fig. 1). Further analyses will be required to compare nuclease-based and nickase-based

Prime Editors, focusing on the distance between the substitution sites and Cas9

nicking/cleavage site. Furthermore, as PEn functions via DSB of genomic DNA, insertion

can potentially cause off-target effects, large deletions, and chromosomal rearrangements[63–65],

whereas the level of large deletions in PEn-mediated prime editing is comparable to that with

the nickase-based PE2 in cultured cells[22]. It will be interesting to see whether these adverse

effects that occurred in injected zebrafish can be neutralised by outcrossing against wild-type

fish in the next generations.

In conclusion, we have established genetically modified zebrafish with programmed short

DNA modifications by microinjecting PE mRNA or RNP complex with gRNAs that integrate

DNA modification via different DNA repairing pathways. Prime editing insertions were

germline-transmitted to the next generation with high efficiency to establish stable
genetically modified zebrafish lines. Our PE-mediated insertion method in zebrafish embryos
offers a highly efficient, low-error approach for editing the genome *in vivo*. The method can
be employed to manipulate coding sequences in endogenous genes and to insert functional
DNA motifs for protein tagging and modulating gene expression. Thus, the findings suggest
broad applicability for Prime Editors across model organisms by offering donor DNA-free
precise gene editing capabilities.

Materials and Methods

Zebrafish strains and husbandry

Wild-type zebrafish strains WIK and AB and the *Tg(smyhc1:gfp)*[i104] transgenic line[38] were used to provide embryos. The collected embryos were raised at 28°C. Adult fish stocks were reared and maintained in the Aquatic Resources Centre at the University of Exeter under the conditions stated in the previous study[66]. The experimental procedures involving the animals were reviewed and approved by the United Kingdom Home Office under the Animals Scientific Procedures Act 1986 (project license: PP3975835) and received ethical approval from the Animal Welfare and Ethical Review Body at the University of Exeter.

Preparation of pegRNA and springRNA

The sequences of the pegRNA and springRNA for prime editing in the *ror2* and *crbn* genes, and for the *smyhc1:gfp* transgene, are listed in Supplementary Table 1. The pegRNAs for prime editing insertion in the *adgrf3b* gene have been reported in a previous study[18]. The first three nucleotides and inter-nucleotide linkages at both ends of the gRNA were chemically modified with 2'-O-methylation and phosphorothioate, respectively. Chemically modified synthetic gRNAs were obtained from Integrated DNA Technologies. The *ror2* springRNA with an abasic RNA spacer was sourced from Horizon Discovery. Synthesised gRNAs were dissolved in nuclease-free duplex buffer (Integrated DNA Technologies) at 100 µM and stored at −20°C.

Microinjection

Microinjection into one-cell stage embryos was carried out under a stereomicroscope using a FemtoJet 4x microinjector (Eppendorf). To prepare an injection mixture for prime editing, 1.25 µl of 600 ng/µl Prime Editor mRNA or 0.6 µl of 10 mg/ml purified Prime Editor protein was combined with 2.5 µl of 12 µM gRNA and 0.5 µl of 0.5% phenol red (Sigma), and the volume was adjusted to 5 µl with 2 M potassium chloride. PEn and PE2 mRNA were synthesised from the linearised plasmids[22] using the mMESSAGE mMACHINE T7 kit (Invitrogen), following the manufacturer's instructions. Purified Prime Editor proteins were prepared according to the protocol in the previous study[37]. A mixture of Prime Editor protein

and gRNA was incubated at 37°C for 5 minutes prior to microinjection to form an RNP

complex. Glass needles for microinjection were prepared from glass capillaries (TW100F-4,

World Precision Instruments) using a P-1000 micropipette puller (Sutter Instrument).

Approximately 1–1.7 nl of the injection mixture was injected per embryo. After

microinjection, the injected embryos were incubated in E3 embryo medium (5 mM sodium

chloride, 0.17 mM potassium chloride, 0.33 mM calcium chloride, and 0.4 mM magnesium

chloride; pH 7.2) at 32°C, following the protocol by Petri et al[18].

DNA extraction, PCR analyses and molecular cloning

Genomic DNA was extracted from individual embryos and larvae using the modified

HotSHOT method[67]. The target region for prime editing in *ror2*, *adgrf3b*, and *smyhc1:gfp*

was amplified by PCR using the following primer sets: *ror2* (5'-

AAACTTATGGGTGCCAGTCC-3' and 5'-ATGGACACAAACTGAGGCTG-3'), *adgrf3b*

(5'-TGATTGCATACACACCTGACC-3' and 5'-AGGCACCTGCAGGAAAATTA-3'), and

smyhc1:gfp (5'-TGCAGTTACAAGGTACAGAGGTC-3' and 5'-

CGTCCTTGAAGAAGATGGTGCG-3'). The PCR mixture was prepared as follows: 10 µl

of 2x PCRBIO Taq Mix red (PCR Biosystems), 0.8 µl each of 10 µM forward and reverse

primers, 1 µl of genomic DNA pooled from 10 individual samples, and 7.4 µl of water,

totalling a volume of 20 µl. The PCR settings were as follows: 95°C for 1 min; 28 cycles of

95°C for 15 s, 58°C for 15 s, and 72°C for 15 s, followed by 72°C for 1 min. The mutation in

the target sequence of the PCR product was assessed by the T7 endonuclease I (T7E1) assay.

PCR products were denatured at 95°C for 5 min and then cooled to room temperature to form

heteroduplex DNAs. Subsequently, 5 µl of PCR product was mixed with 1 µl of NEBuffer 2

(New England Biolabs), 0.15 µl of T7E1 enzyme (10 U/µl, New England Biolabs), and 3.85

µl of water to a total volume of 10 µl and incubated at 37°C for 15 min. The PCR and T7E1-

digested products were assessed on a 3% agarose gel alongside a 100 bp DNA ladder (New

England Biolabs). PCR products were also cloned into the pGEM-T Easy vector (Promega)

and used for transformation with One Shot TOP10 chemically competent cells (Invitrogen)

following the manufacturer's instructions. Bacterial colonies grown on an LB agar plate with

100 µg/ml ampicillin were randomly selected and used for plasmid extraction using a

GeneJET plasmid miniprep kit (Thermo Scientific). Sanger sequencing of PCR products and

plasmids was performed by Eurofins Genomics.

Amplicon sequencing and bioinformatics

PCR amplicons for next-generation sequencing (NGS) of *ror2*, *adgrf3b*, *crbn*, and *smyhc1:gfp* were prepared using Phusion Flash High-Fidelity 2x Mastermix (F548, Thermo Scientific) in a reaction volume of 15 μl, which contained 1.5 μl of genomic DNA extract and 0.2 μM of specific primers with barcodes and adapters for NGS. All primer sequences are shown in Supplementary Table 2. The PCR protocol with Phusion Flash High-Fidelity 2x Mastermix included an initial step at 98°C for 3 min, followed by 30 cycles at 98°C for 10 s, 60°C for 5 s, and 72°C for 5 s. The resulting PCR amplicons were purified using the HighPrep PCR Clean-up System (MagBio Genomics). The characterisations of size, purity, and concentration of the amplicons were conducted using a fragment analyser (Agilent). A second PCR was carried out to add Illumina indexes to the amplicons, utilising KAPA HiFi HotStart Ready Mix (Roche) in a total volume of 25 μl, containing 0.067 ng of PCR template and 0.5 μM indexed primers (Illumina). The PCR conditions were set at 72°C for 3 min, 98°C for 30 s, followed by 10 cycles at 98°C for 10 s, 63°C for 30 s, and 72°C for 3 min, with a 5-minute final extension at 72°C. Amplicons were purified with the HighPrep PCR Clean-up System (MagBio Genomics) and analysed using a fragment analyser (Agilent). Quantification was performed using a Qubit 4 Fluorometer (Life Technologies), with sequencing carried out on the Illumina NextSeq system in accordance with the manufacturer's guidelines. Demultiplexing of the amplicon sequencing data was executed with bcl2fastq software. The resulting fastq files were processed with CRISPResso2 V2.2.12 in prime editing mode[68]. Detailed parameters of the CRISPResso analysis can be found in Supplementary Table 3.

Live imaging

Stereomicroscope images were obtained using an Olympus DP73 microscope camera, and measurements were conducted with cellSens imaging software (Olympus). Confocal images of *smyhc1:gfp* transgene expression were captured with a Leica TCS SP8 laser scanning microscope using a HC Fluotar L 25×/0.95 water objective and LAS X imaging software (Leica). For imaging, live zebrafish larvae were affixed to glass slides with 1% low melting agarose (5806A, Takara) in an E3 embryo medium containing 200 μg/ml tricaine (A5040, Sigma).

Acknowledgements

Research in the S.S. lab, including Y.O. and A.B., is supported by the Biotechnology and
Biological Sciences Research Council Industrial Partnership Awards with AstraZeneca
(BBSRC-IPA, BB/X008401/1; awarded to S.S., Y.O., and C.R.T) and the Living Systems
Institute Open Innovation Platform Fund at the University of Exeter (awarded to Y.O.). M.L.
receives support from the National Centre for the Replacement, Refinement and Reduction of
Animals in Research (NC3Rs) PhD studentship (NC/X001407/1; awarded to J.S.B and
C.R.T). We thank the Aquatic Resources Centre at the University of Exeter for the care of
zebrafish resources.

Author contributions

Y.O., M.P., M.M., and S.S. conceived the study and designed the experimental strategy. Y.O.,
M.L., and A.B. performed the animal experiments and Y.O. and M.P. analysed the data with
support from E.G., J.S.B., C.R.T., S.R., and M.B. The manuscript was prepared by Y.O. and
S.S.

Competing interests

M.P., E.G., S.R., M.B., and M.M. are employees of AstraZeneca and may be AstraZeneca
shareholders.

References

1. Kwan, K. M. et al. The Tol2kit: a multisite gateway-based construction kit for Tol2
transposon transgenesis constructs. *Dev. Dyn.* **236**, 3088–3099 (2007). 10.1002/dvdy.21343.

2. Grabher, C., Joly, J. S. & Wittbrodt, J. Highly efficient zebrafish transgenesis mediated by
the meganuclease I-SceI. *Methods Cell Biol.* **77**, 381–401 (2004). 10.1016/s0091-
679x(04)77021-1.

3. Thermes, V. et al. I-SceI meganuclease mediates highly efficient transgenesis in fish. *Mech.*
Dev. **118**, 91–98 (2002). 10.1016/s0925-4773(02)00218-6.

4. Patton, E. E., Zon, L. I. & Langenau, D. M. Zebrafish disease models in drug discovery:
from preclinical modelling to clinical trials. *Nat. Rev. Drug Discov.* **20**, 611–628 (2021).
10.1038/s41573-021-00210-8.

5. Meng, X., Noyes, M. B., Zhu, L. J., Lawson, N. D. & Wolfe, S. A. Targeted gene
inactivation in zebrafish using engineered zinc-finger nucleases. *Nat. Biotechnol.* **26**, 695–
701 (2008). 10.1038/nbt1398.

6. Doyon, Y. et al. Heritable targeted gene disruption in zebrafish using designed zinc-finger
nucleases. *Nat. Biotechnol.* **26**, 702–708 (2008). 10.1038/nbt1409.

7. Huang, P. et al. Heritable gene targeting in zebrafish using customized TALENs. *Nat.*
Biotechnol. **29**, 699–700 (2011). 10.1038/nbt.1939.

8. Sander, J. D. et al. Targeted gene disruption in somatic zebrafish cells using engineered
TALENs. *Nat. Biotechnol.* **29**, 697–698 (2011). 10.1038/nbt.1934.

9. Varshney, G. K. et al. High-throughput gene targeting and phenotyping in zebrafish using
CRISPR/Cas9. *Genome Res.* **25**, 1030–1042 (2015). 10.1101/gr.186379.114.

10. Jao, L. E., Wente, S. R. & Chen, W. Efficient multiplex biallelic zebrafish genome editing
using a CRISPR nuclease system. *Proc. Natl. Acad. Sci. USA.* **110**, 13904–13909 (2013).
10.1073/pnas.1308335110.

11. Hwang, W. Y. et al. Efficient genome editing in zebrafish using a CRISPR-Cas system.
Nat. Biotechnol. **31**, 227–229 (2013). 10.1038/nbt.2501.

12. Burger, A. et al. Maximizing mutagenesis with solubilized CRISPR-Cas9
ribonucleoprotein complexes. *Development* **143**, 2025–2037 (2016). 10.1242/dev.134809.

13. Wu, R. S. et al. A rapid method for directed gene knockout for screening in G0 zebrafish.
Dev. Cell **46**, 112–125 (2018). 10.1016/j.devcel.2018.06.003.

14. Hoshijima, K., Jurynec, M. J. & Grunwald, D. J. Precise editing of the zebrafish genome
made simple and efficient. *Dev. Cell* **36**, 654–667 (2016). 10.1016/j.devcel.2016.02.015.

15. DiNapoli, S. E. et al. Synthetic CRISPR/Cas9 reagents facilitate genome editing and
homology directed repair. *Nucleic Acids Res.* **48**, e38 (2020). 10.1093/nar/gkaa085.

16. Levic, D. S., Yamaguchi, N., Wang, S., Knaut, H. & Bagnat, M. Knock-in tagging in
zebrafish facilitated by insertion into non-coding regions. *Development* **148**, dev199994
(2021). 10.1242/dev.199994.

17. Anzalone, A. V. et al. Search-and-replace genome editing without double-strand breaks or
donor DNA. *Nature* **576**, 149–157 (2019). 10.1038/s41586-019-1711-4.

18. Petri, K. et al. CRISPR prime editing with ribonucleoprotein complexes in zebrafish and
primary human cells. *Nat. Biotechnol.* **40**, 189–193 (2022). 10.1038/s41587-021-00901-y.

19. Davis, J. R. et al. Efficient prime editing in mouse brain, liver and heart with dual AAVs.
Nat. Biotechnol. **42**, 253–264 (2024). 10.1038/s41587-023-01758-z.

20. Ely, Z. A. et al. A prime editor mouse to model a broad spectrum of somatic mutations in
vivo. *Nat. Biotechnol.* **42**, 424–436 (2024). 10.1038/s41587-023-01783-y.

21. Adikusuma, F. et al. Optimized nickase- and nuclease-based prime editing in human and
mouse cells. *Nucleic Acids Res.* **49**, 10785–10795 (2021). 10.1093/nar/gkab792.

22. Peterka, M. et al. Harnessing DSB repair to promote efficient homology-dependent and -
independent prime editing. *Nat. Commun.* **13**, 1240 (2022). 10.1038/s41467-022-28771-1.

23. Li, X. et al. Development of a versatile nuclease prime editor with upgraded precision.
Nat. Commun. **14**, 305 (2023). 10.1038/s41467-023-35870-0.

24. Kortüm, K. M. et al. Targeted sequencing of refractory myeloma reveals a high incidence
of mutations in CRBN and Ras pathway genes. *Blood* **128**, 1226–1233 (2016).
10.1182/blood-2016-02-698092.

25. Ito, T. et al. Identification of a primary target of thalidomide teratogenicity. *Science* **327**,
1345–1350 (2010). 10.1126/science.1177319.

26. Donovan, K. A. et al. Thalidomide promotes degradation of SALL4, a transcription factor
implicated in Duane Radial Ray syndrome. *eLife* **7**, e38430 (2018). 10.7554/eLife.38430.

27. Fink, E. C. et al. CrbnI391V is sufficient to confer in vivo sensitivity to thalidomide and
its derivatives in mice. *Blood* **132**, 1535–1544 (2018). 10.1182/blood-2018-05-852798.

28. Zhang, W. et al. Enhancing CRISPR prime editing by reducing misfolded pegRNA
interactions. *eLife* **12**, RP90948 (2024). 10.7554/eLife.90948.

29. van Bokhoven, H. et al. Mutation of the gene encoding the ROR2 tyrosine kinase causes
autosomal recessive Robinow syndrome. *Nat. Genet.* **25**, 423–426 (2000). 10.1038/78113.

30. Afzal, A. R. et al. Recessive Robinow syndrome, allelic to dominant brachydactyly type B,
is caused by mutation of ROR2. *Nat. Genet.* **25**, 419–422 (2000). 10.1038/78107.

31. Lima, A. R. et al. Phenotypic and mutational spectrum of ROR2-related Robinow
syndrome. *Hum. Mutat.* **43**, 900–918 (2022). 10.1002/humu.24375.

32. Zhang, C., Brunt, L., Ono, Y., Rogers, S. & Scholpp, S. Cytoneme-mediated transport of
active Wnt5b-Ror2 complexes in zebrafish. *Nature* **625**, 126–133 (2024). 10.1038/s41586-
023-06850-7.

33. Mattes, B. et al. Wnt/PCP controls spreading of Wnt/β-catenin signals by cytonemes in
vertebrates. *eLife* **7**, e36953 (2018). 10.7554/eLife.36953.

34. Brunt, L. et al. Vangl2 promotes the formation of long cytonemes to enable distant Wnt/β-
catenin signaling. *Nat. Commun.* **12**, 2058 (2021). 10.1038/s41467-021-22393-9.

35. Bai, Y. et al. Ror2 receptor mediates Wnt11 ligand signaling and affects convergence and
extension movements in zebrafish. *J. Biol. Chem.* **289**, 20664–20676 (2014).
10.1074/jbc.M114.586099.

36. Dranow, D. B., Le Pabic, P. & Schilling, T. F. The non-canonical Wnt receptor Ror2 is
required for cartilage cell polarity and morphogenesis of the craniofacial skeleton in zebrafish.
Development **150**, dev201273 (2023). 10.1242/dev.201273.

37. Antoniou, P. et al. Modified pegRNAs mitigate scaffold-derived prime editing by-
products. *Nat. Commun.* **16**, 3374 (2025). 10.1038/s41467-025-58653-1.

38. Elworthy, S., Hargrave, M., Knight, R., Mebus, K. & Ingham, P. W. Expression of
multiple slow myosin heavy chain genes reveals a diversity of zebrafish slow twitch muscle
fibres with differing requirements for Hedgehog and Prdm1 activity. *Development* **135**, 2115-
2126 (2008). 10.1242/dev.015719.

39. Dang, C. V. & Lee, W. M. Identification of the human c-myc protein nuclear translocation
signal. *Mol. Cell. Biol.* **8**, 4048–4054 (1988). 10.1128/mcb.8.10.4048-4054.1988.

40. Hodel, M. R., Corbett, A. H. & Hodel, A. E. Dissection of a nuclear localization signal. *J.*
Biol. Chem. **276**, 1317–1325 (2001). 10.1074/jbc.M008522200.

41. Ray, M., Tang, R., Jiang, Z. & Rotello, V. M. Quantitative tracking of protein trafficking
to the nucleus using cytosolic protein delivery by nanoparticle-stabilized nanocapsules.
Bioconjug. Chem. **26**, 1004–1007 (2015). 10.1021/acs.bioconjchem.5b00141.

42. Kuroki-Kami, A. et al. Targeted gene knockin in zebrafish using the 28S rDNA-specific
non-LTR-retrotransposon R2Ol. *Mob. DNA* **10**, 23 (2019). 10.1186/s13100-019-0167-2.

43. Lalonde, R. L. et al. pIGLET: Safe harbor landing sites for reproducible and efficient
transgenesis in zebrafish. *Sci. Adv.* **10**, eadn6603 (2024). 10.1126/sciadv.adn6603.

44. Komor, A. C., Kim, Y. B., Packer, M. S., Zuris, J. A. & Liu, D. R. Programmable editing
of a target base in genomic DNA without double-stranded DNA cleavage. *Nature* **533**, 420–
424 (2016). 10.1038/nature17946.

45. Gaudelli, N. M. et al. Programmable base editing of A•T to G•C in genomic DNA without
DNA cleavage. *Nature* **551**, 464–471 (2017). 10.1038/nature24644.

46. Zhang, Y. et al. Programmable base editing of zebrafish genome using a modified
CRISPR-Cas9 system. *Nat. Commun.* **8**, 118 (2017). 10.1038/s41467-017-00175-6.

47. Rees, H. A. et al. Improving the DNA specificity and applicability of base editing through
protein engineering and protein delivery. *Nat. Commun.* **8**, 15790 (2017).
10.1038/ncomms15790.

48. Zhao, Y., Shang, D., Ying, R., Cheng, H. & Zhou, R. An optimized base editor with
efficient C-to-T base editing in zebrafish. *BMC Biol.* **18**, 190 (2020). 10.1186/s12915-020-
00923-z.

49. Rosello, M. et al. Precise base editing for the in vivo study of developmental signaling
and human pathologies in zebrafish. *eLife* **10**, e65552 (2021). 10.7554/eLife.65552.

50. Qin, W. et al. ABE-ultramax for high-efficiency biallelic adenine base editing in zebrafish.
Nat. Commun. **15**, 5613 (2024). 10.1038/s41467-024-49943-1.

51. Zhang, Y. et al. Cytosine base editors with increased PAM and deaminase motif flexibility
for gene editing in zebrafish. *Nat. Commun.* **15**, 9526 (2024). 10.1038/s41467-024-53735-y.

52. Kim-Yip, R. P. et al. Efficient prime editing in two-cell mouse embryos using PEmbryo.
Nat. Biotechnol. **42**, 1822–1830 (2024). 10.1038/s41587-023-02106-x.

53. Salem, A. R. et al. Prime editing in mice with an engineered pegRNA. *Vascul. Pharmacol.*
154, 107269 (2024). 10.1016/j.vph.2023.107269.

54. Ben-Shachar, S. et al. Dominant versus recessive traits conveyed by allelic mutations - to
what extent is nonsense-mediated decay involved? *Clin. Genet.* **75**, 394–400 (2009).
10.1111/j.1399-0004.2008.01114.x.

55. Wittkopp, N. et al. Nonsense-mediated mRNA decay effectors are essential for zebrafish
embryonic development and survival. *Mol. Cell. Biol.* **29**, 3517–3528 (2009).
10.1128/MCB.00177-09.

56. Lai, S. et al. Effect of nonsense-mediated mRNA decay factor SMG9 deficiency on
premature aging in zebrafish. *Commun. Biol.* **7**, 654 (2024). 10.1038/s42003-024-06356-6.

57. Koeppel, J. et al. Prediction of prime editing insertion efficiencies using sequence
features and DNA repair determinants. *Nat. Biotechnol.* **41**, 1446–1456 (2023).
10.1038/s41587-023-01678-y.

58. Chen, P. J. et al. Enhanced prime editing systems by manipulating cellular determinants
of editing outcomes. *Cell* **184**, 5635–5652.e29 (2021). 10.1016/j.cell.2021.09.018.

59. Ferreira da Silva, J. et al. Prime editing efficiency and fidelity are enhanced in the
absence of mismatch repair. *Nat. Commun.* **13**, 760 (2022). 10.1038/s41467-022-28442-1.

60. Nelson, J. W. et al. Engineered pegRNAs improve prime editing efficiency. *Nat.*
Biotechnol. **40**, 402–410 (2022). 10.1038/s41587-021-01039-7.

61. Ponnienselvan, K. et al. Reducing the inherent auto-inhibitory interaction within the
pegRNA enhances prime editing efficiency. *Nucleic Acids Res.* **51**, 6966–6980 (2023).
10.1093/nar/gkad456.

62. Yan, J. et al. Improving prime editing with an endogenous small RNA-binding protein.
Nature **628**, 639–647 (2024). 10.1038/s41586-024-07259-6.

63. Kosicki, M., Tomberg, K. & Bradley, A. Repair of double-strand breaks induced by
CRISPR-Cas9 leads to large deletions and complex rearrangements. *Nat. Biotechnol.* **36**,
765–771 (2018). 10.1038/nbt.4192.

64. Adikusuma, F. et al. Large deletions induced by Cas9 cleavage. *Nature* **560**, E8–E9
(2018). 10.1038/s41586-018-0380-z.

65. Leibowitz, M. L. et al. Chromothripsis as an on-target consequence of CRISPR-Cas9
genome editing. *Nat. Genet.* **53**, 895–905 (2021). 10.1038/s41588-021-00838-7.

66. Paull, G. C. et al. Variability in measures of reproductive success in laboratory-kept
colonies of zebrafish and implications for studies addressing population-level effects of
environmental chemicals. *Aquat. Toxicol.* **87**, 115–126 (2008).
10.1016/j.aquatox.2008.01.008.

67. Meeker, N. D., Hutchinson, S. A., Ho, L. & Trede, N. S. Method for isolation of PCR-
ready genomic DNA from zebrafish tissues. *Biotechniques* **43**, 610–614 (2007).
10.2144/000112619.

68. Clement, K. et al. CRISPResso2 provides accurate and rapid genome editing sequence
analysis. *Nat. Biotechnol.* **37**, 224–226 (2019). 10.1038/s41587-019-0032-3.

Figure legends

Fig. 1 Prime editing substitution in the zebrafish *crbn* gene, comparing Cas9-nickase-based and nuclease-based Prime Editors. a, Schematic illustration of the functioning of prime editing by the Cas9-nickase-based Prime Editor (PE2) and the nuclease-based Prime Editor (PEn). b-d, Comparison between PEn and PE2 in prime editing substitution in the *crbn* gene with the pegRNA refolding procedure. Proportions of editing outcomes in individual injected embryos are shown by amplicon sequencing (b), alongside quantitative analyses of precise prime edits (c) and indels (d) comparing experimental conditions (n = 10 per group). P-values were determined using one-way ANOVA with Tukey's multiple comparison test in (c), and the Kruskal-Wallis test with Dunn's multiple comparison test in (d). Error bars in the bar graphs represent the mean and standard deviation, and individual data points indicate values from single injected embryos.

Fig. 2 Prime editing insertion in zebrafish *ror2* gene using Cas9-nuclease-based Prime Editor. a, Schematic illustration of the functional domains in the Ror2 protein and alignment of partial amino acid sequences within the tyrosine kinase domain. Sequences from multiple species, including those related to Robinow syndrome (RS) in humans W720X and the corresponding zebrafish W722X mutant, are aligned. The conserved tyrosine residue is highlighted. b, Schematic illustration of guide RNA (gRNA) designs for prime editing insertion in *ror2*. c, Schematic illustration of prime editing insertion by Cas9-nuclease-based Prime Editor (PEn). An additional DNA fragment, reverse-transcribed at the target cleavage site, containing the programmed insertion, is integrated into the genome via homology-directed repair or non-homologous end joining. d, Agarose gel images of genomic PCR products from embryos injected with Prime Editor mRNA and pegRNA/springRNA. PCR products of the *ror2* target region (top) and those after digestion with T7 endonuclease I (T7E1, bottom). e, Sequence alignment of the edits in the *ror2* target site obtained from embryos injected with PEn/springRNA. Prime editing insertion (TGA) is outlined, and the gRNA target sequence is underlined. f, Quantitative comparison of editing outcomes using different combinations of Prime Editor and gRNA. The proportion of sequence reads with each type of edit in amplicon sequencing is presented.

Fig. 3 Comparing different prime editing approaches in zebrafish embryos. a-c,
Comparison between PEn and PE2 in prime editing insertion into *ror2* with various delivery
methods. The proportion of editing outcomes in individual injected embryos was assessed
through amplicon sequencing (a), along with a quantitative analysis of precise prime edit (b)
and indels (c) across experimental conditions (n = 10 per group). d, Sequence alignment of
edits in prime editing insertion in *ror2* via the PEn mRNA/springRNA combination. e,
Schematic illustration of springRNA design featuring an abasic RNA spacer for prime editing
insertion in *ror2*. f and g, Proportion of reads with scaffold incorporation (f) and precise
prime edit (g) as determined by amplicon sequencing of prime editing insertion in *ror2* using
control and abasic springRNA (n = 10 per group; one sample of the PEn RNP/abasic
springRNA combination was excluded from analysis due to low read count). h and i, Prime
editing insertion in the *adgrf3b* gene using PEn. The proportion of reads with a precise 3 bp
(h) and 12 bp insertion (i) was evaluated using pegRNA or springRNA (n = 10 per group). P-
values were determined by Welch's one-way ANOVA with Dunnett T3 multiple comparison
test in b and c, and Kruskal-Wallis test with Dunn's multiple comparison test in f and g. Error
bars in bar graphs represent the mean and standard deviation, and each individual data point
indicates the value from a single injected embryo.

Fig. 4 Generation and characterisation of stable *ror2*^{W722X} mutant. a, Summary of the
screening of F1 embryos. Sixteen embryos were genotyped per F0 founder. Each circle
represents the genotype of a single embryo. The embryos were obtained by outcrossing the
injected founder with wild-type fish; thus, all embryos with the mutation are heterozygous. b-
d, Lateral images of wild type (b), zygotic *ror2*^{W722X} mutant (c), and maternal-zygotic (MZ)
ror2^{W722X} mutant (d) larvae at 5 days post-fertilisation (dpf). e, Sanger sequencing
chromatogram of wild type (top) and *ror2*^{W722X} mutant (bottom) at the prime editing target
site in ror2. Prime editing insertion (TGA) is highlighted. The target sequence of the guide
RNA is underlined, and the cleavage site of Cas9 is indicated by a dotted line. f, Quantitative
analysis of total length comparing wild type, zygotic *ror2*^{W722X} mutant, and MZ *ror2*^{W722X}
mutant at 5 dpf. Each data point on the graph represents the value from a single larva (n = 10
per group). P-values were determined by one-way ANOVA with a Tukey multiple
comparisons test. Error bars represent the mean and standard deviation.

Fig. 5 Prime editing to insert a nuclear localisation signal sequence into the *smyhc1:gfp* transgene. a, Schematic representation of the prime editing insertion of the nuclear localisation signal (NLS) sequence into the *smyhc1:gfp* transgene and the expected eGFP expression in slow-twitch muscle fibres. b, Confocal microscopy image of the trunk muscle in *smyhc1:gfp* larvae at 4 days post-fertilisation (dpf) that were injected with the PEn RNP complex for the prime editing NLS insertion. Putative nuclear GFP expression is indicated by an arrowhead. Anterior is positioned to the left. c, Quantitative analysis of the efficiency of precise NLS insertion via amplicon sequencing. Two pegRNAs of differing lengths are compared (n = 10 per group, with one sample excluded from the analysis due to low read count). The p-value was determined using the Mann-Whitney U test. Error bars represent the mean and standard deviation, while individual data points on the graph indicate values from single injected larvae. d, Confocal microscopy images of the trunk muscle of F1 larvae exhibiting nuclear GFP expression at 2 dpf, obtained from founder 1 (see panel f). The GFP fluorescence channel (left) and pseudo-colour (right) images are shown. Anterior is to the left. e, Sequence alignment of the edits in the target site obtained from F1 embryos exhibiting nuclear GFP expression (founder 1). The NLS sequence is outlined in red. f, Summary of F1 embryo screening. F1 embryos obtained from 6 founders are categorised based on the GFP expression pattern. The embryos were produced by outcrossing the heterozygous *smyhc1:gfp* founder injected with the PEn RNP complex to wild type; thus, half of the embryos are expected to be negative for the *smyhc1:gfp* transgene.

 Supplementary Figure legends

 Supplementary Fig. 1 Prime editing substitution in the zebrafish *crbn* gene. a, Schematic
illustration of the strategy for prime editing substitution in *crbn* gene. The nucleotides for
substitution are indicated in red. The guide RNA (gRNA) target sequence is underlined. b,
Quantitative analyses of precision scores comparing between PEn and PE2 in prime editing
substitution in the *crbn* gene with the pegRNA refolding procedure (n = 10 per group). P-
values were determined using Welch's one-way ANOVA with Dunnett T3 multiple
comparison test. Error bars in the bar graphs represent the mean and standard deviation, and
individual data points indicate values from single injected embryos.

Supplementary Fig. 2 Analysis of prime editing insertion in zebrafish *ror2* gene by
sequencing the clones. a, Sequence alignment of the edits in the *ror2* target site obtained
from 20 randomly selected bacterial clones comparing prime editing outcomes using
PE2/pegRNA (top) and PEn/pegRNA (bottom) combinations. The guide RNA (gRNA) target
sequence is indicated by green arrows. b, Quantitative comparison of editing outcomes using
different combinations of Prime Editor and guide RNA. The proportion of the clones with
each type of edit is presented.

Supplementary Fig. 3 Amplicon sequencing analysis of prime editing insertion in
zebrafish *ror2* and *agdrf3b* genes. a-c, Quantitative analyses of precision scores comparing
various experimental conditions for prime editing insertion in *ror2* gene using pegRNA (a)
and springRNA (b), and in *adgrf3b* gene (c). The sample size is n = 10 per group and one
sample of the PEn RNP/abasic springRNA combination was excluded from analysis in b due
to low read count. P-values were determined by Welch's one-way ANOVA with Dunnett T3
multiple comparison test in a, and Kruskal-Wallis test with Dunn's multiple comparison test
in b and c. Error bars in bar graphs represent the mean and standard deviation, and each
individual data point indicates the value from a single injected embryo. d, Sequence
alignment of edits in prime editing insertion in *ror2* via the PE2 mRNA/pegRNA (top) and
PEn mRNA/pegRNA (bottom) combinations.

Supplementary Fig. 4 Prime editing insertion of a nuclear localisation signal sequence

into the *smyhc1:gfp* **transgene.** a, The expression of *smyhc1:gfp* transgene in 2 days post-

fertilisation (dpf) larvae. Lateral images of whole larvae (left) and the trunk muscle (right).

Anterior is to the left. b, Confocal microscopy images of the trunk muscle of F1 larvae

exhibiting nuclear GFP expression at 2 dpf, obtained from founder 2, 3, and 4 (see Fig. 5f). c,

Sequence alignment of the edits in the target site obtained from F1 embryos exhibiting

nuclear GFP expression (founder 2, 3, and 4). The nuclear localisation signal (NLS)

sequences are outlined in red.

Figure 1

Figure 2

a

Ror2 — Ig-like — CRD — Kringle — TM — Tyrosine kinase — PRD — Ser/Thr — Ser/Thr

Human	LPCPDDCPAWVYALMIECWNEFPSRRPRFKDIHSRLRAWG
RS (W720X)	LPCPDDCPA*
Mouse	LPCPDDCPAWVYALMIECWNEFPSRRPRFKDIHSRLRSWG
Xenopus	LLCPDDCPAWIYTLMLECWSEFPARRPRFKDIHTRLRTWE
Zebrafish	LSCPDDCPAWIYTMMLECWNEFPSRRPRFKDIHARLRTWE
Zebrafish (W722X)	LSCPDDCPA*

b

pegRNA
| 20 nt | 76 nt | 13 nt | 3 nt | 13 nt |
5' | Spacer | Scaffold | Homology arm | PBS | 3'
UGA

springRNA
5' | Spacer | Scaffold | PBS | 3'
UGA

c

Insert — Homology annealing
Homology arm

Non-homologous end joining (NHEJ)

d

PE2 | PEn

M | Control | pegRNA | Control | pegRNA | springRNA

PCR

PCR +T7E1

e

C C A G C C - - - - T G G A T A T A C A C G A T G A T G C T   Reference genome

C C A G C C - - - - T G G A T A T A C A C G A T G A T G C T   Wild type (10/20)
C C A G C C - - T G A T G G A T A T A C A C G A T G A T G C T   Precise prime edit (1/20)
C C A G C C G C T G A T G G A T A T A C A C G A T G A T G C T
C C A G C C - C T G A T G G A T A T A C A C G A T G A T G C T   Prime edit +indel (7/20)
C C A A C - - T G A T G G A T A T A C A C G A T G A A G C T
C C A G C C - - - T G A T G G A T A T A C A C G A T G A T G C T
C C A G C C - - - - - - - - G A T A T A C A C G A T G A T G C T   Indel (2/20)
C C A G C C - - - - - - - - - - - A T A C A C G A T G A T G C T

▲ Cleavage site

f

ror2 3 bp insertion

Sequence (%)

- Precise prime edit
- Imprecise prime edit
- Indels
- Wild type

PE2+pegRNA | PEn+pegRNA | PEn+springRNA

Figure 3

a *ror2* 3 bp insertion

Precise prime edit
Imprecise prime edit
Indels
Wild type

b Precise prime edit

c Indels

d PEn + *ror2* 3 bp insertion springRNA

e *ror2* 3 bp insertion abasic springRNA

f Scaffold incorporation

g Precise prime edit

h *adgrf3b* 3 bp insertion

i *adgrf3b* 12 bp insertion

Figure 4

a
Founder
1 2 3 4 5 6 7 8 9 10

F1 embryo
16
1

● W722X (TGA insertion)
● Other mutations
○ Wild type

b Wild type

5 dpf

c *ror2*^W722X

d MZ *ror2*^W722X

1 mm

e
C C A G C C T G G A T A T A C A C G A T G A T G C T
Wild type

C C A G C C T G A T G G A T A T A C A C G A T G A T
ror2^W722X

f

P < 0.0001
P < 0.0001

Total length (µm)

Wild type   *ror2*^W722X   MZ *ror2*^W722X
n = 10   n = 10   n = 10

Figure 5