Ubiquitous systems drift in the evolution of development

P.L. van der Jagt [1,2]**, S. Oud** [1,2]**, and R.M.A. Vroomans** [1,2]

[1]Sainsbury Laboratory, University of Cambridge, Bateman Street, Cambridge CB2 1LR, UK

[2]Department of Genetics, Downing Site, University of Cambridge, Cambridge CB2 3EJ, UK

Statement of significance

A key open question in evo-devo research is the evolvability of complex phenotypes: to which extent is neutral or beneficial change hindered by deleterious mutations? We investigated the potential for developmental systems drift (DSD) in plant development using a computational evo-devo model. We found that the regulatory interactions between genes changed extensively, resulting in the continual neutral rewiring of the gene regulatory network underpinning development. Even regulatory interactions that were essential for correct development were replaced over long evolutionary time scales. Using plant genome and gene expression data from two publicly available datasets, we found high turnover of cis-regulatory elements without consistent change in gene expression, confirming the widespread occurrence of DSD as predicted by our model.

Developmental systems drift (DSD) is a process where a phenotypic trait is conserved over evolutionary time, while the genetic basis for the trait changes. DSD has been identified in models with simpler genotype-phenotype maps (GPMs), however the extent of DSD in more complex GPMs, such as developmental systems, is debated. To investigate the occurrence of DSD in complex developmental GPMs, we constructed a multi-scale computational model of the evolution of gene regulatory networks (GRNs) governing plant meristem (stem cell niche) development. We found that, during adaptation, some regulatory interactions became essential for the correct expression of stem cell niche genes. These regulatory interactions were subsequently conserved for thousands of generations. Nevertheless, we observed that these deeply conserved regulatory interactions could be lost over the extended period of neutral evolution. These losses were compensated by changes elsewhere in the GRN, which then became conserved as well. This gain and loss of regulatory interactions resulted in a continual *cis*-regulatory rewiring in which accumulated changes caused changes in the expression of several genes. Using two publicly available datasets we confirmed the prevalence of *cis*-regulatory changes across six evolutionary divergent plant species, and showed that these changes do not necessarily impact gene expression patterns, demonstrating the occurrence of DSD. These findings align with the results from our computational model, showing that DSD is pervasive in the evolution of complex developmental systems.

Developmental Systems Drift | Evolution | Development | Cis-regulation

Correspondence: *plv21@cam.ac.uk*

Introduction

Phenotypic traits are often conserved between related species, even when the developmental process that generates them diverged significantly (Cotterell and Sharpe, 2010; Brun-Usan et al., 2022). This phenomenon is called developmental systems drift (DSD) (True and Haag, 2001; McColgan and DiFrisco, 2024), and occurs when multiple genotypes which are separated by few mutational steps encode the same phenotype, forming a neutral path (Kimura, 1979; Wagner, 2008a). The length of the neutral path, and thus the potential for DSD, depends on how many of such genotypes are present and how they are mutationally connected to each other – this is also referred to as the genotype-phenotype map (GPM) (Huynen, 1996; Hogeweg, 2000). DSD is found to drive speciation (Schiffman and Ralph, 2022), explain the developmental hourglass model (Wotton et al., 2015; Crombach et al., 2016), and accelerate adaptation (Zheng et al., 2019, 2020).

However, it remains poorly understood how complex phenotypes are distributed throughout genotype space within highly complex GPMs: do neutral paths continue to percolate through genotype space, or do complex phenotypes occur in isolated genotype islands, limiting the extent of DSD? Computational models of high-dimensional GPMs, such as RNA folding (Schuster et al., 1994; Huynen, 1996; Wagner, 2008b; Van Nimwegen et al., 1999), protein folding (Lipman and Wilbur, 1991), gene regulatory networks (GRNs) (Ciliberti et al., 2007; Payne et al., 2014) have revealed extensive neutral paths that allow for significant genotype variation while maintaining the same phenotype. For other GPMs, such as those resulting from multicellular development, it has been suggested that complex phenotypes are sparsely distributed in genotype space, and have low potential for DSD because neutral mutations anti-correlate with phenotypic complexity (Orr, 2000; Hagolani et al., 2021). Others have found that increased complexity introduces more degrees of freedom, allowing for a greater number of genotypes to produce the same phenotype and potentially allowing for more DSD (Schiffman and Ralph, 2022; Greenbury et al., 2022). The question therefore remains: how much DSD actually occurs in species undergoing multicellular development?

One multicellular structure which displays evidence of DSD is the shoot apical meristem (SAM) in plants. SAMs are multicellular structures which contain a stem cell niche that generates all above-ground plant tissues (Szövényi et al., 2019). Some key regulators of SAM stem cell regulation, such as the CLAVATA3/Embryo Surrounding Region-Related (CLE) peptide family, are deeply conserved among land plants (Whitewoods et al., 2018). However, expression of these conserved regulators in the SAM has diverged significantly between major vascular plant lineages such as lycophytes, monilophytes and angiosperms (Frank et al., 2015). Even within angiosperms, the precise CLE peptides governing SAM function differ between species, and their expression patterns also vary (Somssich et al., 2016). Thus, even similar-looking SAMs are likely regulated by subtly different mechanisms, suggesting that DSD plays a role in SAM evolution.

To investigate the potential for DSD in plant SAMs, we developed a computational model of gene regulatory network evolution, extending an evo-devo approach previously applied to animal development ten Tusscher and Hogeweg (2011); Vroomans et al. (2016, 2018). We used this approach to generate GRNs that reproduce the expression patterns of the central zone (CZ, containing the stem cells), and the supporting organizing center (OC) of the *Arabidopsis thaliana* SAM, in a 2D cellularized tissue. Our evolutionary simulations show that a small number of regulatory interactions become highly conserved in the GRNs that govern the genetic patterning of the SAM. These interactions are typically very important for generating the CZ and OC pattern in the tissue, and disrupting them leads to lethal phenotypes. Surprisingly, we found that even deeply conserved, essential regulatory interactions can diverge over longer evolutionary timescales. Moreover, the expression pattern of genes that were not under direct selection changed over evolutionary time, demonstrating that the extensive neutral paths in genotype space also resulted in developmental systems drift at the level of tissue patterning.

Finally, we validated these theoretical findings using two publicly available datasets: one on conserved non-coding sequences (CNSs) in plants (Hendelman et al., 2021), and another on organ-specific RNA expression across plant species at different evolutionary distances (Schuster et al., 2024). Analysis of CNSs confirmed the existence of highly conserved regulatory sequences and supported our finding that gene regulatory interactions can be lost at very disparate rates over evolutionary time. Combining CNS data with cross-species RNA expression profiles further revealed a strong correlation between conserved sequences associated with a gene and its expression pattern. Interestingly, we found that entirely different sets of conserved regulatory sequences can result in similar gene expression patterns, providing empirical evidence for a many-to-one GP mapping underlying plant gene expression and widespread regulatory rewiring. These findings highlight the prevalence of DSD and its role in shaping developmental evolution.

Results

Model overview

We developed a computational model of gene expression evolution in shoot apical meristems (SAMs). We modelled a population of SAMs, each undergoing a developmental process encoded by a heritable genome. A genome consists of a string of genetic elements representing genes (each of which encodes a transcription factor (TF)), and transcription factor binding sites (TFBSs) which regulate the expression of the downstream gene (Fig. 1A) (Crombach and Hogeweg, 2008). The genome therefore encodes a gene regulatory network (GRN), which governs gene expression in the cells of a two-dimensional tissue representing a longitudinal cut through the SAM (Fig. 1B-D). A subset of TFs exhibit specific properties, such as the ability to diffuse, form dimers, or mediate cell-cell communication with directly neighbouring cells. Gene expression and protein production are subject to a small amount of molecular noise, modelled with stochastic differential equations (Methods, section D). Development therefore consists of spatiotemporal changes in protein distribution within the tissue due to gene expression, diffusion and noise (Fig. 1D).

The development of each individual begins with one TF uniformly distributed throughout the tissue, while a second diffusible TF is constitutively expressed in the epidermal layer (L1) (Fig. 1C). From this initial condition, individuals have a fixed amount of time to express other genes based on the interactions encoded by the genome. At the end of this period, each individual is assigned a fitness score based on the protein concentration of two target genes in specific regions of the SAM: one in the central zone (CZ), and one in the organizing center (OC) (Fig. 1C). This fitness score determines the probability that the individual will produce offspring in the next generation (Fig. 1E). During reproduction, the parent's genome is inherited by the offspring with random mutations (Fig. 1F). This cycle of development and reproduction with mutation is repeated for 50,000 generations.

Figure 1. Overview of the computational evo-devo model. A. A schematic representation of an *in silico* genome including genes and Transcription Factor Binding Sites (TFBSs). **B**. Such a genome can be translated into a gene regulatory network (GRN) which is used to simulate the development. **C**. The different functional areas in the simulated tissues. The L1 layer is shown as red-lined cells, the CZ is shown in the tip of the tissue, the OC is in the center, and their regions overlap in a single row of cells. **D**. An example of the expression pattern of the CZ gene at the end of development in a fit individual. **E**. Selection is based on the expression of fitness genes, shown here are examples of a high fitness, medium fitness and low fitness CZ gene expression pattern (left-to-right). **F**. The different types of structural genome mutations possible during the simulations. In addition to these mutations, mutations are possible in the binding constants of TFBSs, and the maximum transcription rates of genes.

Conservation of regulatory interactions during evolution of developmental programs

We ran 20 simulations, each with a constant population size of 1000 individuals. Out of the 20 populations, 15 evolved to correctly express both the OC and CZ genes, resulting in a fitness (f) score of at least 75 out of 100 (Fig. 2A, examples of high ($\geq$75) and low (<75) fitness patterns in B). To investigate how the developmental programs evolved that generated the pattern, we tracked how long each regulatory interaction between TFs was conserved in the ancestral lineage of each high-fitness population. Most interactions persisted for only a few hundred to a few thousand generations (Fig. 2C), in the same pattern as a control simulation (C) run with random reproduction and no selection for a pattern. However, a small but significant subset of interactions, which we call "conserved interactions," remained consistently present for more than 5000 generations, which did not occur in the control simulation (Fig.

Figure 2. Fitness increase is correlated with GRN conservation. **A**. Fitness of all individuals in the 20 populations at generation 50,000 (outliers not shown). Dotted line denotes the cutoff fitness of 75. **B**. Protein pattern of the two fitness genes for the fittest individuals of simulations 10 and 11, with a fitness of 51.9 and 89.4, respectively. **C**. The number of generations each regulatory interaction was conserved in the ancestor trace of a population. The box plot indicated with C is a control simulation without selection, resulting in the absence of highly conserved interactions. Dotted line indicates the maximum conservation time of interactions within the control simulation (7100 generations). We rounded this down to 5000 as 'conservation cutoff' for the rest of this work. **D**. The sorted conservation times of all regulatory interactions within the ancestry trace of simulation 2. In red are shown all interactions with a conservation time greater than the conservation cutoff of >5000 generations. **E**. Number of conserved (>5000 generations) interactions and fitness of ancestor trace for the first 50,000 generations of simulation 2. **F**. Fitness distribution of 10,000 randomly generated offspring of the fittest individual from simulation 2 at generation 50,000. Orange: offspring inheriting genomes without any mutations $n = 7{,}503$ IQR=5.69; Blue: offspring with mutation(s) $n = 2{,}497$ IQR=13.24.

---

¹¹⁵ 2C, D). These conserved interactions emerged as individuals in a population achieved higher fitness (Fig. 2E, S1,
¹¹⁶ control in S2). This suggests that deeply conserved interactions are a consequence of selection.
¹¹⁷ To assess the potential for neutral evolution and DSD after the target expression pattern evolved, we created
¹¹⁸ 10,000 offspring of the highest-fitness individual of population 2 at generation 50,000 (Fig. 2F). As mutations are
¹¹⁹ probabilistic, the majority of offspring inherit the parental genome without any mutations. Any variation in fitness of
¹²⁰ these non-mutated offspring results from variation in cell connectivity in the SAM (Methods, section C) and noisy
¹²¹ gene expression (Methods, section D). We found that fitness variation in these non-mutated individuals follows a tight
¹²² distribution around a high mean fitness, showing that the developmental mechanism is robust to these sources of
¹²³ noise (Fig. 2F, orange). In offspring which inherited a genome with mutations, the fitness distribution instead followed
¹²⁴ a U shape (Fig. 2F, blue). The majority of mutations was (near) neutral, indicating a high degree of mutational
¹²⁵ robustness and redundancy in the GRN, while a smaller set was very deleterious, resulting in a near-zero fitness.
¹²⁶ Mutations resulting in an 'in between' fitness were more rare, consistent with previous findings on fitness landscapes
¹²⁷ (Hogeweg, 2012). Overall this shows that the evolved genotypes are both developmentally and mutationally robust
¹²⁸ which indicates the existence of neutral areas within the GPM around this complex expression phenotype.

Developmental systems drift in evolved gene regulatory networks

¹³⁰ DSD occurs when the phenotype remains conserved along the ancestral lineage while the underlying regulatory
¹³¹ architecture changes. Our simulations matched the target phenotype by 30,000 generations, after which they entered
¹³² a fitness plateau where evolution was driven by stabilising selection rather than adaptation (Fig. S1). To study DSD
¹³³ at this fitness plateau, we selected the eight populations which reached the highest maximum fitness, created five
¹³⁴ clones of each, and evolved these for an additional 50,000 generations (Fig. 3A). The median fitness increase
¹³⁵ remained indistinguishable from background variability during this period (Fig. S3), indicating predominantly neutral
¹³⁶ evolution (we excluded the clones of 2/8 populations due to more significant fitness increase, Fig. S3).
¹³⁷ We quantified genetic divergence of individuals from their ancestor before the cloning at generation 50,000, using
¹³⁸ a divergence score based on the adjacency matrices of their GRNs (Methods I). We found that the full GRNs
¹³⁹ diverged rapidly from the common ancestor (CA) (Fig. 3B, S4, orange line), which was likely due to changes in
¹⁴⁰ the redundant or non-functional parts in the GRN, which can be mutated without any phenotypic consequence (see
¹⁴¹ fig. 2F). Therefore, we also measured divergence by including only the conserved interactions in the adjacency matrix
¹⁴² (Fig. 3B, S4, blue line). Strikingly, these conserved interactions gradually diverged from the CA, indicating neutral

Figure 3. Neutral evolutionary change in regulatory mechanisms. A. At generation 50,000, populations that reached a fitness plateau are cloned into 5 separate populations that continue to evolve independently but share a common ancestor before generation 50,000. **B**. Divergence of interactions between lineages of each clonal population. Divergence is shown for lineages of the full GRNs with that of the common ancestor of all lineages (Full-CA); only the conserved set of interactions with the common ancestor (Core-CA, > 5000 generations); and the conserved set of interactions between the populations (Core). Shown are the medians and IQRs. Divergence between networks is calculated as in Methods I. **C**. Functional networks (Methods G) and expression patterns of the five fittest individuals at generation 100,000 from simulations 5_0 to 5_4, and their common ancestor (CA) at generation 49,800. The expression patterns of genes $1, 2, 4$ and 11 are shown to illustrate different levels of genotypic and phenotypic divergence.

turnover despite their initial conservation. Since conserved interactions emerge during adaptation (Fig. 2E) and therefore are likely functionally important, their turnover may indicate that the regulatory dynamics that generate the target pattern have changed as well. To assess whether conserved interactions follow similar evolutionary trajectories across independent lineages, we performed a pairwise comparison of evolved GRNs between cloned populations (Fig. 3B, S4, green line). We found that conserved regulatory interactions diverged between populations at a similar rate as each lineage diverged from the CA. This indicates that overall, the different lineages follow different paths of divergence from the CA.

To observe the divergence in GRNs more closely, we pruned the full GRNs to remove non-functional and redundant genes and interactions, and compared these pruned GRNs between individuals from different clonal populations, as well as with their CA (Fig. 3C). As expected, some regulatory interactions in these networks were highly conserved, e.g., the regulation of gene 3, which is consistently activated by gene 11 and inhibited by gene 1 across all GRNs. In contrast, other interactions were rewired, e.g., the activation of gene 8, which is driven by gene 0 in networks $CA, 5_1, 5_3$ but by gene 1 in 5_0 and 5_4, and gene 5 in 5_2. Since these GRNs were pruned to eliminate redundancy and all descend from the same CA, differences in their regulatory interactions reflect functional divergence through neutral evolution, indicating the occurrence of DSD.

To understand the effects of this regulatory turnover on development, we examined the pattern of the proteins that were not subject to selection for a specific pattern. Indeed, we found that the patterns of some free proteins remained conserved, but the expression of other genes underwent significant change (Fig. 3C, S5, S6). For example, the expression of protein 1 was nearly identical among different populations, while proteins 2, 8, and 11 displayed varying degrees of divergence in pattern compared to the CA. Interestingly, protein 2 had varying patterns, but was absent in the pruned networks of $CA, 5_1, 5_3$. In line with our findings in figure 3B, this shows that different lineages explored different parts of the neutral evolutionary space, where in some cases TFs were recruited to regulate the expression of the genes under selection, whereas in others the TF remained nonfunctional and accumulated neutral changes. Taken together, we found that DSD can drive functional divergence resulting in novel spatial expression dynamics.

Network redundancy creates space for rewiring

To understand how conserved and functional interactions can diverge without disrupting fitness, we traced conserved regulatory interactions over evolutionary time in two of the cloned populations. Consistent with our earlier observations, we found that over time, interactions were lost and new conserved interactions arose (Fig. 4A,B). To investigate the conditions under which an interaction can be rewired, we examined a single interaction which is conserved in the lineage of one population but lost in the lineage of a related population, as indicated by the ◀ in Fig. 4A. We measured in each generation how much its removal affected fitness: an importance value of 1 indicates a total loss of fitness after removal, while a value of 0 indicates no change in fitness (see Methods H).

In both lineages, the importance of the interaction $(0 \rightarrow 8)$ remained high for the first 60,000 generations. However, in lineage 5_2, a new interaction $(5 \rightarrow 8)$ emerged that activated gene 8, which coincides with a drop in the importance of $0 \rightarrow 8$. This reduction in importance provided an opportunity for the deletion of $0 \rightarrow 8$ without a significant loss of fitness. This example shows the general mechanism by which functional redundancy enables the turnover of regulatory interactions, causing neutral GRN divergence over long evolutionary timescales.

We even observed this process among interactions regulating the fitness genes (Fig. 4C), which could be rewired multiple times in relatively quick succession. The TF taking over regulation did not necessarily have the same expression pattern as the original: e.g. gene 8, the new regulator of gene 12 at generation 60,000, has a different pattern from gene 4. However, they were both expressed around the center, which is where gene 4's regulation of gene 12 was active. This shows how the evolution of GRNs exploits overlaps in expression patterns to generate redundancies, allowing the rewiring of regulatory interactions.

DSD in plant gene regulation: Conservation, loss and emergence of deeply conserved non-coding sequences

Our computational simulations suggest that *cis*-regulatory rewiring plays a significant role in the divergence of developmental programs. To test this hypothesis, we used two available data sets: the Conservatory Project (Hendelman et al., 2021), and Schuster *et al.* Schuster et al. (2024). The Conservatory Project collects conserved non-coding sequences across plant genomes, which we used to investigate the extent of GRN rewiring in flowering plants (Hendelman et al., 2021). Schuster *et al.* measured gene expression in different tissues of several species via bulk RNAseq Schuster et al. (2024), which we used to test for gene expression changes that correspond to GRN rewiring. We constrained our search for regulatory rewiring to the six species shared by these two data sets (*Arabidopsis thaliana, Arabidopsis lyrata, Capsella rubella, Eutrema salsugineum, Medicago truncatula,* and *Brachypodium distachyon*).

Figure 4. Rewiring of conserved interactions through loss of function. A. Presence of conserved interactions in two related lineages, every bar representing a specific conserved interaction. The green bars indicated with a ◄ show the interaction 0 activates 8. The lineages are 5_0 and 5_2, respectively. **B.** Importance of conserved interactions that upregulate gene 8 expression over evolutionary time of the respective simulations shown in A. The legend shows the TF regulating 8. Indicated under the plot is the persistence time of each interaction. **C.** Importance of conserved interactions upregulating gene 12 (fitness gene). **D.** Expression of subset of genes $(1, 4, 7, 8)$ upregulating expression of gene 12 at generations 20,000, 40,000, 60,000, 80,000 and 100,000. Pattern is transparent if there is no interaction between the respective gene and gene 12 at that time point.

We used conserved non-coding Sequences (CNSs) from the Conservatory Project dataset as a proxy for *cis*-regulatory elements (Hendelman et al., 2021). A CNS is defined as a non-coding sequence conserved within an orthogroup, which is a collection of genes descending from a single ancestral gene (Emms and Kelly, 2019). The dataset contains 2.3 million CNSs identified across 284 species. Starting with the orthogroup of a gene in the CNS dataset, we mapped all CNSs to their corresponding genes across the six species (Fig. 5A); we focused on 3 orthogroups containing genes relevant to meristem function (see Methods J). One orthogroup of interest includes the MADS box transcription factors SEPALLATA (SEP) 1–4, which are important for organ specification in the floral meristem (Pelaz et al., 2000; Ditta et al., 2004). Within this orthogroup, we identified 1658 CNSs: 3 were conserved across all six species, 6 were conserved in eudicots, and 366 were conserved within Brassicaceae (Fig. 5B). The stark difference in shared CNSs between Brassicaceae and more distantly related species highlights a high turnover rate of CNSs, a pattern observed across all genes investigated (Fig. S7). Despite this high turnover, a small subset of nine CNSs was deeply conserved (Fig. 5B). Additionally, the data revealed likely cases of loss of otherwise deeply conserved CNSs in particular species. For example, 1 CNS was lost only in *A. lyrata* and another in *E. salsugineum*. This pattern – CNSs lost in one lineage but conserved in others – resembles the rewiring dynamics we observed in our simulations.

Next, we examined the CNS composition of individual SEP genes to gain a more fine-grained view of regulatory rewiring. We found that CNSs associated with a gene, e.g. SEP4, closely resemble the set of CNSs of SEP4 orthologs in related species (Fig. 5C,D). However, more distantly related genes, such as SEP4 and SEP3, tend to have disjoint CNS sets, likely due to *cis*-regulatory rewiring after the duplication event that created the paralogs. Interestingly, partial overlaps in CNS sets were observed between certain gene pairs, such as *SEP1* and *SEP2*. This overlap may reflect a more recent duplication event, with the shared CNSs present before duplication. Based on our simulations, we predict that the CNSs still shared between these genes have a higher regulatory importance.

Finally, we explored the relationship between CNS composition and gene expression patterns using RNA expression data from the Schuster dataset (Schuster et al., 2024). The SEP genes with similar CNS sets tend to exhibit similar expression patterns (Fig. 5E), which we quantified by calculating the Pearson distance between the expression patterns of all gene pairs, and then comparing this distance to the similarity in CNS set for each pair (Fig. 5F, S9, S10). Notably, genes with any CNS set similarity (> 0) rarely display high Pearson distances, suggesting that shared CNSs strongly correlate with similar expression patterns. However, dissimilar CNS sets do not necessarily imply different expression patterns (Fig. 5F). Most gene pairs with disjoint CNS sets (similarity = 0) exhibit a wide range of expression divergences, including pairs with very similar expression patterns. Assuming a shared and continuous ancestral function, this is evidence of regulatory rewiring while preserving phenotype — a hallmark of DSD.

Figure 5. **A**. Schematic depicting the relation between orthogroup, different species, different genes and CNSs. **B**. Counts of CNS presence in the six angiosperm species. The filled circles indicate in which species the CNSs are present, the barplot shows the count of number of CNSs with that profile. **C**. Gene tree of primary transcript protein sequences of SEP genes of the four Brassicaceae species, Es: *Eutrema salsugineum*; Cr: *Capsella rubella*; Al: *Arabidopsis lyrata*; At: *Arabidopsis thaliana*. Note that the branch to the outgroups has been truncated for readability, for a full tree see supplementary figure S8. Only bootstrap values less than 90% are shown. **D**. Heatmap of CNS presence for each of the SEP genes. Presence of a CNS is shown in grey. **E**. Expression profile of SEP genes in different organs. The expression data shown are variance stabilized mRNA counts taken directly from Schuster et al. (2024). **F**. Pearson distance of expression against the similarity of the CNS set for each gene pair in **C**.

Discussion

In this study, we investigated developmental systems drift (DSD) in a complex genotype-phenotype map (GPM) of plant development. We constructed a model of gene regulation evolution in the plant shoot apical meristem (SAM) to analyze genotypic and phenotypic changes over long evolutionary timescales. In our simulations we observed that a subset of gene interactions was conserved over thousands of generations, while other interactions turned over quickly. These interactions were highly important for generating the target gene expression pattern; nevertheless, mutations could result in redundancies that lowered the importance of such interactions, allowing for their eventual removal from the genome. This finding was supported with data from the Conservatory Project (Hendelman et al., 2021), which showed that regulatory rewiring, through gain and loss of conserved non-coding sequences, is prevalent in plants. We also found that such rewiring preserved the expression pattern under selection while facilitating the exploration of diverse expression patterns in genes that were not under direct selection, revealing a highly interconnected genotype space – a finding that we confirmed with the Schuster dataset (Schuster et al., 2024)

Step-wise *cis*-regulatory rewiring

By tracking regulatory interactions over evolutionary time, we found that *cis*-regulatory rewiring of the GRN can render previously essential interactions redundant. This redundancy creates opportunities for further rewiring, resulting in a stepwise mechanism of regulatory evolution that can completely change the structure of the GRN over time. Our analysis also showed that the GRN is both mutationally robust (allowing for rewiring)(Wagner, 2008b) and sensitive to mutations at the sites of highly important interactions. This matches bioinformatic analysis of the rice genome, where it was found that while most promoter sites experience low selection pressure, a small but significant subset is under strong selection pressure (Joly-Lopez et al., 2020). The generally low selection pressure on promoter sites suggests that there is sufficient evolutionary space for neutral mutations to accumulate and drive the gradual rewiring of the GRN.

A concrete example of regulatory rewiring in the SAM can be found in two antagonistic peptide signaling pathways involving genes of the CLE family. The CLV3 pathway inhibits stem cell fate, whereas the CLE40 pathway promotes it (Schlegel et al., 2021). Interestingly, the CLV3 and CLE40 proteins are functionally similar, and a *clv3-2* mutant can be rescued by expression of *CLE40* under a *CLV3* promoter (Hobe et al., 2003). Similarly, opposite effects have been found for mutations in the CNS of *UNUSUAL FLOWER ORGANS* in tomato and *A. thaliana* (Lanctot et al., 2025). This demonstrates that rewiring the regulatory context of a functionally similar protein can lead to a completely opposite function.

Genetic diversity due to DSD

DSD necessitates the existence of multiple genotypes that map to the same trait. Such many-to-one mappings have been identified in various models of genotype-phenotype maps (GPMs). For example, Cotterell and Sharpe (2010) showed that a three-gene model of stripe formation can generate similar spatial patterns through many distinct regulatory interactions. In their model, alternative solutions clustered in a few disconnected regions in genotype space, corresponding to distinct regulatory dynamics. This suggests a limited range of genotype space in which DSD could occur (Jiménez et al., 2015). Here, we show that large shifts in gene expression can accompany GRN rewiring during stabilising selection, suggesting that even the underlying regulatory mechanism can change.

GRN rewiring through regulatory redundancy requires that genes can regulate and be regulated by multiple TFs. This many-to-many architecture is therefore key to DSD. A previous model Johnson and Porter (2007) that lacked this regulatory flexibility, displayed DSD under directional selection but not during stabilising selection. Our model's more permissive and realistic network architecture allows for a phenotype to be encoded by a wider range of genotypes, facilitating accumulation of neutral changes. In this way, our system is conceptually similar to RNA folding GPMs, where very different sequences can produce the same secondary structure (Huynen, 1996; Van Nimwegen et al., 1999).

Change in gene expression patterns due to DSD

The structure of the GPM not only determines the length of the neutral path, but also the number of novel phenotypes that can be reached — often called evolvability (Wagner and Altenberg, 1996; Wagner, 2012). For simpler GPMs, such as that of RNA folding and GRNs, neutral paths were shown to connect areas with different phenotypes (Wagner, 2014; Crombach and Hogeweg, 2008). In a more complex GPM, such as morphological development, complex phenotypes were rare and mutations were found to often reduce complexity, which led to the conclusion that neutral paths, and therefore DSD, could be less extensive Hagolani et al. (2021). However, evolution has been shown to increase mutational robustness over time, resulting in the possibility for more neutral change (Van Nimwegen et al., 1999). Furthermore, fitness in our model is dependent on the expression pattern of only two genes, leaving the other twelve genes free to evolve the expression pattern needed to support correct fitness gene expression. We find that

under DSD, the expression of these non-fitness genes changes, even the ones directly impacting the expression of fitness genes. This may have allowed for more degrees of freedom, while morphogenesis instead could add additional constraints to the types of regulatory changes allowed (Cano-Fernández et al., 2024). Nevertheless, in our model many mutations are still highly deleterious, as seen in Figure 2F, making the mutational landscape similar to that of complex phenotypes in (Hagolani et al., 2021). It takes multiple generations for the evolutionary process to find and build on those few mutations that are neutral or beneficial. In future work, it will be therefore be interesting to explore the length of neutral paths in models of morphogenesis such as (Hagolani et al., 2021) over extended evolutionary time scales.

Future experimental validation

The pattern of conservation and loss of CNSs in the Conservatory Project dataset show a remarkable similarity to our results. However, CNSs most likely do not translate perfectly to GRN connections, as they may or may not contain one or more TF binding sites. This makes it difficult to make precise claims on the rewiring of interactions besides complete loss. Additionally, the dataset focuses solely on sequences that are conserved between (the reference genomes of) a large collection of different species, which means that it could exclude sequences which are gained and lost at much shorter time scales or within species. This would be resolved by better and denser sampling of genomes across the plant phylogeny.

To test our predictions more quantitatively, we need to know which TFs bind the upstream region of a gene, as direct evidence of a regulatory interaction. A recent preprint by Baumgart et al. (2024) exemplifies this approach, presenting experimental validation of TF binding sites across ten different species using a specialized DNA affinity purification sequencing (DAP-seq) technique. While DAP-seq screens do not provide conclusive evidence that the given TF-regulatory site interaction is important in a given developmental context (Bartlett et al., 2017), they do provide more direct insight into the regulatory elements that are conserved or lost across species. Furthermore, experimental work investigating the developmental role of CREs has shown extensive epistasis — where the effect of a mutation depends on the genetic background — supporting DSD (Schaerli et al., 2018; Aguirre et al., 2023; Ciren et al., 2024). Extending the DAPseq and functional approach to multiple species and multiple CREs would allow us to quantify the extent of such DSD.

The gene expression data utilized in this study represents bulk expression at the organ level, such as the vegetative meristem (Schuster et al., 2024). This limits our analysis of the phenotypic effects of rewiring to comparisons between organs. However, our computational simulations predict differences in gene expression within the same organ, suggesting a finer scale of regulation. More fine-grained approaches, such as single-cell RNA sequencing or spatial transcriptomics, will provide a more detailed understanding of how gene expression is modulated spatially and temporally within complex tissues, allowing for a closer alignment between computational predictions and experimental observations.

Supplementary Note 1: Acknowledgments

We thank Madelaine Bartlett for helpful discussions and support with the conservatory.org dataset. We thank Elliot Meyerowitz and Christoph Schuster for the use of their dataset. We thank the Sainsbury Laboratory Evolution Journal Club and Enrico Sandro Colizzi for their comments and suggestions on the manuscript.

Funding

RMAV is supported by the Gatsby Charitable Foundation (G112566). The funding body had no role in the design of the study, collection and analysis of the data, or writing of the manuscript.

Methods

A. Code availability

All code for the evolutionary model, analysing simulation output, and the bioinformatics analysis is available at GitLab.

B. Computational model

We developed a computational model of an evolving population of organisms that undergoes cycles of development and selection. Each organism possesses a "beads-on-a-string" genome consisting of genes and upstream regulatory sites (Crombach and Hogeweg, 2008). This genome encodes a gene regulatory network (GRN) that governs gene expression dynamics in a 2D cellularized tissue. A single cycle, or generation, of the population proceeds through the following steps: each individual undergoes development governed by the dynamics of its GRN; this results in gene expression patterns in the tissue, which form the phenotype of the individual; this phenotype is given a fitness value by evaluating how well it meets a target expression pattern; individuals are selected for reproduction based on their fitness value, passing on their genome to the next generation; when a genome is passed on, it is randomly mutated.

C. 2D Tissue generation

Before running the evolutionary simulations, 1000 different tissues (representing the shoot apical meristem, SAM) are generated with different placements of cells. Each tissue contains 130 cells, each cell is initialized as a point in two dimensional space ($x, y \in \mathbb{R}^2$) within $y = 0$ and $y = -0.01x^2$ on the domain $x = [-61, 61]$. Cells are attached to their neighbours with springs. The potential energy of springs is $U_s = \frac{1}{2}(x - x_r)^2 k$ where x is the current state of the spring, x_r the relaxed state and k the spring constant. To prevent cells from moving out of the predefined tissue shape, a boundary energy is added $U_b = \frac{-10}{d}$ where d is the distance between a cell and the boundary. Cells are distributed further by relaxing the potential energy for 500 steps with a dt of 0.1 . Using these positions, a voronoi diagram is computed and cells are moved for a further 2000 steps towards their centroid using the Lloyds algorithm (Du et al., 1999).

D. Development of genotype into phenotype

A genome is composed of two components: transcription factor binding sites (TFBSs) and genes (Fig. S11A). Genes encode for transcription factors (TFs) which each bind a specific TFBS type (denoted with an integer identifier). The TFBSs in front of a gene determine which TFs regulate the expression of that gene. TFs of type 8 and 9 can also form dimers, which bind a TF of type 7. Based on these interactions the genome is translated into a reaction network (Fig. S11B). The reaction network is used to compute gene expression in the cells of a multicellular tissue (Fig. S11C).

The reaction network is modelled as a set of stochastic differential equations (SDEs) (Ahmad, 1988; Gillespie, 2007) governing mRNA, protein and TF dimer concentrations. The rate equations for mRNA ($g(t)$) and protein concentrations ($p(t)$) are given by

$$dg(t) = \big[\tau(t) - \alpha g(t)\big]dt + S_g(t) \tag{1}$$

$$dp(t) = \big[\upsilon(t) - \beta p(t) + \psi(t) + D\nabla^2 p(t)\big]dt + S_p(t). \tag{2}$$

With $\tau(t)$ the transcription rate, see Eq. 3; $\upsilon(t)$ translation rate; α and β decay rates; $\psi(t)$ dimerization; and $D\nabla^2$ diffusion. The transcription rate of a gene is determined by the concentrations of the TFs that match its upstream TFBSs (θ, see Eq. 6), and modelled using a Hill-Langmuir equation. A TF either enhances or inhibits transcription, which is determined by the sign $w \in \{-1, 1\}$ of the TFBS.

$$\tau(t) = \begin{cases} \left[\dfrac{(\sum \big[w_i \theta_i(p_k)\big])^2}{(\sum \big[w_i \theta_i(p_k)\big])^2 + 1}\right]\tau_{max}, & \text{if } \sum \big[w_i \theta_i(p_k)\big] > 0, \\ 0, & \text{else.} \end{cases} \tag{3}$$

The translation rate is linear with the amount of available mRNA.

$$\upsilon_k(t) = g_k(t) \tag{4}$$

The S terms, see Eq. 5, represent the stochastic part of the equations as a Wiener process (Ahmad, 1988). We only model transcription, translation and decay as stochastic processes, while dimer formation is always assumed to be in equilibrium.

$$S_g(t) = \sqrt{\tau(t)}dW_1 + \sqrt{\alpha g(t)}dW_2$$
$$S_p(t) = \sqrt{\upsilon(t)}dW_3 + \sqrt{\beta p(t)}dW_4. \tag{5}$$

The effect of a TF on the transcription rate of a gene is modelled using a Hill equation. A TFBS i of type k has its own Hill constant H_i and is bound by a specific type of TF with concentration p_k.

$$\theta_i(p_k) = \left[\frac{p_k^2}{p_k^2 + H_i^2} \right]. \tag{6}$$

E. Fitness function

During development the fitness of individuals is determined at time points within the intervals $[200, 250]$ and $[450, 500]$. Fitness depends on the correct protein concentration of the two fitness genes in each cell, p_{cz} and p_{oc} respectively. Each cell is assigned to one of the following four types, a CZ cell (type 1); a CZ and OC cell (type 2); an OC cell (type 3); other cell, see Figure 1C.

$$f_i = \begin{cases} \dfrac{100 \cdot p_{cz}}{(p_{cz} + 100)} - \dfrac{50 \cdot p_{oc}}{(p_{oc} + 200)}, & \text{type==1} \\[2ex] \dfrac{50 \cdot p_{cz}}{(p_{cz} + 100)} + \dfrac{50 \cdot p_{oc}}{(p_{oc} + 100)}, & \text{type==2} \\[2ex] \dfrac{100 \cdot p_{oc}}{(p_{oc} + 100)} - \dfrac{50 \cdot p_{cz}}{(p_{cz} + 200)}, & \text{type==3} \\[2ex] -\dfrac{50 \cdot p_{oc}}{(p_{oc} + 200)} - \dfrac{50 \cdot p_{cz}}{(p_{cz} + 200)}, & \text{else.} \end{cases} \tag{7}$$

In order to prevent overexpression of fitness genes, the positive term of the fitness equations is capped at a concentration of $p = 400$, after which the fitness of a cell is set to 100.

F. Selection and Mutation

Individuals are selected for reproduction proportional to their fitness f_i, with the probability p of selecting an individual i from population P:

$$p_i = \frac{f_i}{\sum_{j \in P} f_j}. \tag{8}$$

Offspring inherits their parent's genome with mutations. All implemented mutations are given in table 1. Besides mutations in the genome, we also allow for changes in the diffusion, association and dissociation constants of the diffusing and dimerising proteins.

Gene	TFBS
Deletion	Deletion
Insertion	Insertion
Transcription rate change (τ_{max})	Hill constant change (H)
mRNA decay rate change (α)	Sign change (w)
	Type change
Whole genome duplication	

Table 1. The different mutation types implemented in the model. Indicated between brackets is the changed parameter if applicable, see D.

G. Functional network analysis

The evolved GRNs contain redundant and/or non-functional interactions. We obtain a 'functional network' from a full evolved network through a pruning algorithm. In the pruning algorithm all possible single deletions are performed on a genome, development run for this set of genomes (5 times per genome to account for stochastic development), and fitness is computed for all resulting expression patterns. The genome with the highest average fitness is accepted and the procedure is iterated until no mutations maintain a fitness within 5 points from the original fitness.

H. Importance analysis

The importance I of a regulatory interaction is calculated as the fraction of fitness loss after the interaction is removed. Since the model is stochastic, development is repeated N times after removal of the interaction and the average fitness is taken.

$$I = \frac{1}{N} \sum^{N} \frac{f_0 - f_r}{f_0}. \tag{9}$$

Here, f_0 is the original fitness and f_r is the fitness when the regulatory interaction is removed. For all analysis N was set to 500.

I. Divergence analysis

The divergence between two networks is computed by rewriting each network as a Boolean adjacency matrix where all genes of the same type are merged: if there are two genes of type X, all TFBSs of those two genes are marked as input (1) in row X of the adjacency matrix. We compute the difference between two of these matrices, A and B, as follows

$$d(A,B) = \frac{\sum_i^N \sum_j^N \left[|A_{i,j} - B_{i,j}| \right]}{\sum_i^N \sum_j^N \left[A_{i,j} + B_{i,j} \right]}. \tag{10}$$

Using this difference we can define the divergence between the GRNs of two individuals, C and D as:

$$div(C,D) = \frac{1}{2} \left[d(C_{act}, D_{act}) + d(C_{inh}, D_{inh}) \right], \tag{11}$$

where the GRNs of C and D are transformed into 2 Boolean matrices containing all activating (C_{act}, D_{act}) and inhibiting (C_{inh}, D_{inh}) interactions.

J. Conserved Non-coding Sequence analysis

For the counts of CNS presence profiles analysis we created a set with all the CNS ids which belong to a specific orthogroup. For each CNS in this set we assessed its presence in the given orthogroup of each species.
The phylogeny was made with the approach described in Man et al. (2020). In short, annotated proteins of interest were aligned using MAFFT and conserved regions identified (Katoh and Standley, 2013). For each conserved region, a BLAST search was performed (Altschul et al., 1990), and a hidden Markov model (HMMs) was generated with HMMER version 3.4 (Potter et al., 2018). Hits from both the BLAST and HMMER searches were taken together and a cutoff was determined using a known outgroup. Finally, using the generated set of hits a maximum likelihood (ML) tree was build using IQ-TREE 2 (Minh et al., 2020) with 100 bootstraps.

K. Parameters of simulation

	Value	Unit	Use
Mechanical parameters			
x_bounds	[-61, 61]	[um, um]	Horizontal bounds of the tissue
y_bounds	[0, 40]	[um, um]	Vertical bounds of the system
F_bound	10	a.f.u.	Boundary force
F_spring	1	a.f.u.	Spring constant
eq_dist	0.9	um	Equillibrium distance of the spring
Molecular parameters			
decay mRNA	0.2 [0.05, 0.9]	t^{-1}	Initial decay rate [min, max]
decay protein	0.2	t^{-1}	Decay rate of proteins
connections	3		Initial avg nr of connections per gene
H	40 [5, 100]	a.m.u.	Initial Hill constant [min, max]
E	25 [5, 80]	a.m.u. t^{-1}	Initial max expression rate [min, max]
k_{on}	0.01 [0.00001, 0.1]		Initial k_{on} of the dimer gene [min, max]
k_{off}	0.001 [0.00001, 0.1]		Initial k_{off} of the dimer gene [min, max]
D	1 [0.5, 5]	um^2 t^{-1}	Diffusion coefficient [min, max]
Simulation parameters			
N_pop	1000		Population size
dev_steps	500		Developmental timesteps
generations	50000		Number of generations (initially)
N_genes	14		Number of unique genes
N_diff	2		Number of diffusing gene products
N_cellcell	2		Number of cellcell gene products
Genome parameters			
gene duplication probability	0.002		per gene
gene deletion probability	0.003		per gene
gene parameter change	0.0005		Probability of changing decay mRNA or E
TFBS sign change	0.0002		per TFBS
TFBS type switch	0.0002		Probability of changing TFBS type per TFBS
TFBS duplication probability	0.0005		per TFBS
TFBS deletion probability	0.00075		per TFBS
TFBS innovation probability	0.00004		Probability of adding a new TFBS (per genome)
TFBS H change probability	0.002		per TFBS

Table 2. Table of the standard parameters used. a.f.u., arbitrary force unit. a.m.u., arbitrary molar unit. t, developmental time step.

Bibliography

Aguirre, L., Hendelman, A., Hutton, S. F., McCandlish, D. M., and Lippman, Z. B. (2023). Idiosyncratic and dose-dependent epistasis drives variation in tomato fruit size. *Science*, 382(6668):315–320.

Ahmad, R. Introduction to stochastic differential equations, (1988).

Altschul, S. F., Gish, W., Miller, W., Myers, E. W., and Lipman, D. J. (1990). Basic local alignment search tool. *Journal of molecular biology*, 215(3):403–410.

Bartlett, A., O'Malley, R. C., Huang, S.-s. C., Galli, M., Nery, J. R., Gallavotti, A., and Ecker, J. R. (2017). Mapping genome-wide transcription-factor binding sites using dap-seq. *Nature protocols*, 12(8):1659–1672.

Baumgart, L. A., Morales-Cruz, A., Greenblum, S. I., Wang, P., Zhang, Y., Yang, L., Chen, C., Dilworth, D. J., Garretson, A. C., Grosjean, N., et al. (2024). An atlas of conserved transcription factor binding sites reveals the cell type-resolved gene regulatory landscape of flowering plants. *bioRxiv*, pages 2024–10.

Brun-Usan, M., Zimm, R., and Uller, T. (2022). Beyond genotype-phenotype maps: Toward a phenotype-centered perspective on evolution. *BioEssays*, 44(9):2100225.

Cano-Fernández, H., Brun-Usan, M., Tissot, T., and Salazar-Ciudad, I. (2024). A morphospace exploration using a general model of development reveals a basic set of morphologies for early animal development and evolution. *Journal of Experimental Zoology Part B: Molecular and Developmental Evolution*.

Ciliberti, S., Martin, O. C., and Wagner, A. (2007). Innovation and robustness in complex regulatory gene networks. *Proceedings of the National Academy of Sciences*, 104(34):13591–13596.

Ciren, D., Zebell, S., and Lippman, Z. B. (2024). Extreme restructuring of cis-regulatory regions controlling a deeply conserved plant stem cell regulator. *PLoS Genetics*, 20(3):e1011174.

Cotterell, J. and Sharpe, J. (2010). An atlas of gene regulatory networks reveals multiple three-gene mechanisms for interpreting morphogen gradients. *Molecular systems biology*, 6(1):425.

Crombach, A. and Hogeweg, P. Evolution of evolvability in gene regulatory networks, (2008). ISSN 1553734X.

Crombach, A., Wotton, K. R., Jiménez-Guri, E., and Jaeger, J. (2016). Gap gene regulatory dynamics evolve along a genotype network. *Molecular biology and evolution*, 33(5):1293–1307.

Ditta, G., Pinyopich, A., Robles, P., Pelaz, S., and Yanofsky, M. F. (2004). The sep4 gene of arabidopsis thaliana functions in floral organ and meristem identity. *Current biology*, 14(21):1935–1940.

Du, Q., Faber, V., and Gunzburger, M. (1999). Centroidal voronoi tessellations: Applications and algorithms. *SIAM review*, 41(4): 637–676.

Emms, D. M. and Kelly, S. (2019). Orthofinder: phylogenetic orthology inference for comparative genomics. *Genome biology*, 20: 1–14.

Frank, M. H., Edwards, M. B., Schultz, E. R., McKain, M. R., Fei, Z., Sørensen, I., Rose, J. K., and Scanlon, M. J. (2015). Dissecting the molecular signatures of apical cell-type shoot meristems from two ancient land plant lineages. *New Phytologist*, 207(3):893–904.

Gillespie, D. T. (2007). Stochastic simulation of chemical kinetics. *Annu. Rev. Phys. Chem.*, 58(1):35–55.

Greenbury, S. F., Louis, A. A., and Ahnert, S. E. (2022). The structure of genotype-phenotype maps makes fitness landscapes navigable. *Nature Ecology & Evolution*, 6(11):1742–1752.

Hagolani, P. F., Zimm, R., Vroomans, R., and Salazar-Ciudad, I. (2021). On the evolution and development of morphological complexity: A view from gene regulatory networks. *PLoS computational biology*, 17(2):e1008570.

Hendelman, A., Zebell, S., Rodriguez-Leal, D., Dukler, N., Robitaille, G., Wu, X., Kostyun, J., Tal, L., Wang, P., Bartlett, M. E., et al. (2021). Conserved pleiotropy of an ancient plant homeobox gene uncovered by cis-regulatory dissection. *Cell*, 184(7): 1724–1739.

Hobe, M., Müller, R., Grünewald, M., Brand, U., and Simon, R. (2003). Loss of cle40, a protein functionally equivalent to the stem cell restricting signal clv3, enhances root waving in arabidopsis. *Development genes and evolution*, 213:371–381.

Hogeweg, P. (2000). Shapes in the shadow: evolutionary dynamics of morphogenesis. *Artificial Life*, 6(1):85–101.

Hogeweg, P. (2012). Toward a theory of multilevel evolution: long-term information integration shapes the mutational landscape and enhances evolvability. *Evolutionary systems biology*, pages 195–224.

Huynen, M. A. (1996). Exploring phenotype space through neutral evolution. *Journal of molecular evolution*, 43:165–169.

Jiménez, A., Cotterell, J., Munteanu, A., and Sharpe, J. (2015). Dynamics of gene circuits shapes evolvability. *Proceedings of the National Academy of Sciences*, 112(7):2103–2108.

Johnson, N. A. and Porter, A. H. (2007). Evolution of branched regulatory genetic pathways: directional selection on pleiotropic loci accelerates developmental system drift. *Genetica*, 129:57–70.

Joly-Lopez, Z., Platts, A. E., Gulko, B., Choi, J. Y., Groen, S. C., Zhong, X., Siepel, A., and Purugganan, M. D. (2020). An inferred fitness consequence map of the rice genome. *Nature plants*, 6(2):119–130.

Katoh, K. and Standley, D. M. (2013). Mafft multiple sequence alignment software version 7: improvements in performance and usability. *Molecular biology and evolution*, 30(4):772–780.

Kimura, M. (1979). The neutral theory of molecular evolution. *Scientific American*, 241(5):98–129.

Lanctot, A., Hendelman, A., Udilovich, P., Robitaille, G. M., and Lippman, Z. B. (2025). Antagonizing cis-regulatory elements of a conserved flowering gene mediate developmental robustness. *Proceedings of the National Academy of Sciences*, 122(8): e2421990122.

Lipman, D. J. and Wilbur, W. J. (1991). Modelling neutral and selective evolution of protein folding. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 245(1312):7–11.

Man, J., Gallagher, J. P., and Bartlett, M. (2020). Structural evolution drives diversification of the large lrr-rlk gene family. *New Phytologist*, 226(5):1492–1505.

McColgan, Á. and DiFrisco, J. (2024). Understanding developmental system drift. *Development*, 151(20).

Minh, B. Q., Schmidt, H. A., Chernomor, O., Schrempf, D., Woodhams, M. D., Von Haeseler, A., and Lanfear, R. (2020). Iq-tree 2: new models and efficient methods for phylogenetic inference in the genomic era. *Molecular biology and evolution*, 37(5): 1530–1534.

Orr, H. A. (2000). Adaptation and the cost of complexity. *Evolution*, 54(1):13–20.

Payne, J. L., Moore, J. H., and Wagner, A. (2014). Robustness, evolvability, and the logic of genetic regulation. *Artificial life*, 20
(1):111–126.
Pelaz, S., Ditta, G. S., Baumann, E., Wisman, E., and Yanofsky, M. F. (2000). B and c floral organ identity functions require
sepallata mads-box genes. *Nature*, 405(6783):200–203.
Potter, S. C., Luciani, A., Eddy, S. R., Park, Y., Lopez, R., and Finn, R. D. (2018). Hmmer web server: 2018 update. *Nucleic acids*
research, 46(W1):W200–W204.
Schaerli, Y., Jiménez, A., Duarte, J. M., Mihajlovic, L., Renggli, J., Isalan, M., Sharpe, J., and Wagner, A. (2018). Synthetic
circuits reveal how mechanisms of gene regulatory networks constrain evolution. *Molecular Systems Biology*, 14(9):e8102.
Schiffman, J. S. and Ralph, P. L. (2022). System drift and speciation. *Evolution*, 76(2):236–251.
Schlegel, J., Denay, G., Wink, R., Pinto, K. G., Stahl, Y., Schmid, J., Blümke, P., and Simon, R. G. (2021). Control of arabidopsis
shoot stem cell homeostasis by two antagonistic cle peptide signalling pathways. *Elife*, 10:e70934.
Schuster, C., Gabel, A., Drost, H.-G., Grosse, I., Leyser, O., and Meyerowitz, E. M. (2024). Rapid evolution of gene expression
patterns in flowering plants. *bioRxiv*, pages 2024–07.
Schuster, P., Fontana, W., Stadler, P. F., and Hofacker, I. L. (1994). From sequences to shapes and back: a case study in rna
secondary structures. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 255(1344):279–284.
Somssich, M., Je, B. I., Simon, R., and Jackson, D. (2016). Clavata-wuschel signaling in the shoot meristem. *Development*
(Cambridge), 143:3238–3248. doi: $10.1242/\mathrm{dev}.133645$.
Szövényi, P., Waller, M., and Kirbis, A. Chapter one - evolution of the plant body plan. In Grossniklaus, U., editor, *Plant*
Development and Evolution, volume 131 of *Current Topics in Developmental Biology*, pages 1 – 34. Academic Press, (2019).
doi: $\mathrm{https://doi.org/10.1016/bs.ctdb.2018.11.005}$.
ten Tusscher, K. H. and Hogeweg, P. (2011). Evolution of networks for body plan patterning; interplay of modularity, robustness
and evolvability. *PLoS Computational Biology*, 7. doi: $10.1371/\mathrm{journal.pcbi.1002208}$.
True, J. R. and Haag, E. S. (2001). Developmental system drift and flexibility in evolutionary trajectories. *Evolution & development*,
3(2):109–119.
Van Nimwegen, E., Crutchfield, J. P., and Huynen, M. (1999). Neutral evolution of mutational robustness. *Proceedings of the*
National Academy of Sciences, 96(17):9716–9720.
Vroomans, R. M., Hogeweg, P., and Ten Tusscher, K. H. (2016). In silico evo-devo: reconstructing stages in the evolution of
animal segmentation. *EvoDevo*, 7:1–20.
Vroomans, R. M., Hogeweg, P., and Ten Tusscher, K. H. (2018). Around the clock: gradient shape and noise impact the evolution
of oscillatory segmentation dynamics. *EvoDevo*, 9:1–20.
Wagner, A. (2008). Neutralism and selectionism: a network-based reconciliation. *Nature Reviews Genetics*, 9(12):965–974.
Wagner, A. (2008). Robustness and evolvability: a paradox resolved. *Proceedings of the Royal Society B: Biological Sciences*,
275(1630):91–100.
Wagner, A. (2012). The role of robustness in phenotypic adaptation and innovation. *Proceedings of the Royal Society B: Biological*
Sciences, 279(1732):1249–1258.
Wagner, A. (2014). Mutational robustness accelerates the origin of novel rna phenotypes through phenotypic plasticity.
Biophysical journal, 106(4):955–965.
Wagner, G. P. and Altenberg, L. (1996). Perspective: complex adaptations and the evolution of evolvability. *Evolution*, 50(3):
967–976.
Whitewoods, C. D., Cammarata, J., Nemec Venza, Z., Sang, S., Crook, A. D., Aoyama, T., Wang, X. Y., Waller, M., Kamisugi,
Y., Cuming, A. C., Szövényi, P., Nimchuk, Z. L., Roeder, A. H., Scanlon, M. J., and Harrison, C. J. (2018). Clavata was
a genetic novelty for the morphological innovation of 3d growth in land plants. *Current Biology*, 28(15):2365–2376.e5. doi:
$\mathrm{https://doi.org/10.1016/j.cub.2018.05.068}$.
Wotton, K. R., Jiménez-Guri, E., Crombach, A., Janssens, H., Alcaine-Colet, A., Lemke, S., Schmidt-Ott, U., and Jaeger, J.
(2015). Quantitative system drift compensates for altered maternal inputs to the gap gene network of the scuttle fly megaselia
abdita. *Elife*, 4:e04785.
Zheng, J., Payne, J. L., and Wagner, A. (2019). Cryptic genetic variation accelerates evolution by opening access to diverse
adaptive peaks. *Science*, 365(6451):347–353.
Zheng, J., Guo, N., and Wagner, A. (2020). Selection enhances protein evolvability by increasing mutational robustness and
foldability. *Science*, 370(6521):eabb5962.