1

# *PACS1 syndrome variant alters proteomic landscape of developing cortical organoids*

Ximena Gomez Maqueo[1], Lauren E Rylaarsdam [1], Ashley Woo [1], Annika L Schroder [1], Jennifer Rakotomamonjy [1], Shelby Ruiz [2], Jordan Gilardi[2], Lambertus Klei [2], Bernie Devlin [2], Matthew L MacDonald [2,*], Alicia Guemez-Gamboa[1,*]

[1]*Department of Neuroscience, Feinberg School of Medicine, Northwestern University, Chicago IL, USA*

[2]*Department of Psychiatry, University of Pittsburgh, Pittsburgh, PA, USA*

*Correspondence: alicia.guemez@northwestern.edu, mlmacdonald@gmail.com*

**Abstract**

PACS1 syndrome is a neurodevelopmental disorder resulting from a unique *de novo* p.R203W variant in Phosphofurin Acidic Cluster Sorting protein 1 (PACS1). *PACS1* encodes a multifunctional sorting protein required for localizing furin to the *trans*-Golgi network. Although few studies have started to investigate the impact of the PACS1 p.R203W variant, the mechanisms by which the variant affects neurodevelopment are still poorly understood. In recent years, ASD patient-derived brain organoids have been increasingly used to identify pathogenic mechanisms and possible therapeutic targets. While most of these studies investigate the mechanisms by which ASD-risk genes affect the transcriptome, studies investigating the proteome are limited. Here, we investigated the effect of PACS1 p.R203W on the proteomic landscape of brain organoids using tandem mass tag (TMT) mass-spectrometry. Time series analysis between PACS1$^{(+/+)}$ and PACS1$^{(+/R203W)}$ organoids uncovered several proteins with dysregulated abundance or phosphorylation status, including known PACS1 interactors. Although we observed low overlap between proteins with altered expression and phosphorylation, the resulting dysregulated processes converged. The presence of PACS1 p.R203W variant putatively accelerated synaptogenesis and impaired vesicle loading and recycling, putatively leading to defective and/or incomplete synaptic function. Finally, the key dysregulated proteins observed in PACS1$^{(+/R203W)}$ organoids were also enriched in ASD-risk genes and have been associated with other neurological diseases. Our results highlight that proteomic analyses not only enhance our understanding of general NDD mechanisms by complementing transcriptomic studies, but also could uncover additional targets, facilitating therapy development.

**Introduction**

Neurodevelopmental disorders (NDD) are heterogenous and usually present with complex etiology. Several large-scale genetic studies into the etiology of NDDs such as autism spectrum disorder (ASD), intellectual disability (ID), and epilepsy have been conducted, leading to hundreds of risk genes being identified. However, our understanding of the mechanisms by which the observed genetic variants act has lagged, limiting translation of genetic findings into clinical treatments (1, 2). Characterizing the pathogenic mechanisms of different monogenic NDDs may reveal convergent therapy targets. PACS1 syndrome (MIM615009) is a monogenic NDD characterized by ASD, ID, craniofacial abnormalities, and epilepsy (3-5). PACS1 syndrome results from a recurrent *de novo* c.607C>T variant in the gene encoding P̲hosphofurin A̲cidic C̲luster S̲orting protein 1 (PACS1). This missense variant is present in all reported patients and results in an arginine to tryptophan (p.R203W) substitution at residue 203, with only one known arginine to glutamine p.R203Q exception (5).

*PACS1* encodes a multifunctional sorting protein required for furin localization to the *trans*-Golgi network (tGN) (6, 7). PACS1 binds furin and other targets via acidic cluster sorting motifs (8-11). In addition to tGN retrieval, PACS1 mediates the recycling of endocytosed furin from early endosomes to the cell surface and the delivery of proteins to the primary cilium (12-14). PACS1 has been shown to interact with over 50 other proteins (10, 14-17). Although the role of PACS1 has primarily been studied in the cytoplasm, several reports show that it also accumulates in the nucleus (18, 19) where it mediates DNA replication repair (20, 21), contributes to the nuclear export of viral transcripts (18), and regulates chromatin stability through interaction with histone deacetylase (HDAC) proteins (20).

Recent reports have begun to explore the pathogenesis of the PACS1 p.R203W variant. Mice overexpressing PACS1 p.R203W exclusively in cortical neurons present with increased dendrite arborization and decreased postsynaptic currents, possibly due to a pathogenic interaction between PACS1 p.R203W and HDAC6. Moreover, antisense oligonucleotides (ASO)

directed towards both PACS1 and HDAC6 ameliorate these defects (22). Transcriptomic analysis

in patient-derived brain organoids revealed a dysregulation in cell projection organization in

immature PACS1 p.R203W neurons, while mature neurons exhibited an upregulation in synaptic

signaling processes when compared to isogenic controls. Subsequent characterization of

spontaneous electrical activity revealed that induced pluripotent stem cell (iPSC)-derived neurons

harboring the p.R203W variant display asynchrony and an increased network burst duration (23).

Finally, a study using *C. elegans* showed that the PACS1 syndrome-associated variant altered

PACS1 localization but had no effect on neuron morphology and synaptic patterns (24). Although

these studies have started to investigate the impact of the PACS1 p.R203W variant, the

mechanisms by which the variant affects neurodevelopment are still poorly understood. This in

turn results in few therapeutic for patients, which include only non-invasive behavioral approaches

and sensory integration therapies to manage the symptoms (25, 26).

In recent years, ASD patient-derived brain organoids have been increasingly used to

identify pathogenic mechanisms and possible therapeutic targets (27). While most of these

studies investigate the mechanisms by which ASD-risk genes affect the transcriptome (28, 29),

studies investigating the proteome are limited. However, mRNA levels do not always predict

protein levels, as protein expression and degradation are regulated by mechanisms beyond

transcription (30). Here we used tandem mass tag (TMT) mass-spectrometry to provide further

understanding into the pathogenic mechanisms of PACS1 syndrome by uncovering whether brain

organoids carrying the unique PACS1 p.R203W variant present with a dysregulated proteomic

landscape during maturation.

**Methods**

***Generation of forebrain organoids from iPSCs***

Three isogenic pairs of PACS1$^{(+/+)}$ and PACS1$^{(+/R203W)}$ iPSC lines were previously generated (23)

and differentiated to brain organoids using the kit based on Sloan et al. protocol (31) (Stem Cell

Technologies, 08620). See Supplementary Methods for further details on organoid generation

and immunostaining.

**Sample preparation for proteomic analysis**

Total protein was extracted from the organoids by sonication in lysis buffer (1% sodium dodecyl

sulfate (SDS), 125 mM triethylammonium bicarbonate (TEAB), 75 mM NaCl, and HaltTM protease

and phosphatase inhibitors). Lysate samples from two additional control cell lines were included

to improve protein detection. See Supplemental Methods and Supplementary Table 1 for detailed

description of the mass spectrometry sample preparation, TMT labeling and phosphopeptide

enrichment processing steps.

**MS data acquisition and raw data processing**

For peptide/protein quantification, TMT labeled peptides (~1 µg) were loaded onto a heated

PepMap RSLC C18 2 µm, 100 angstrom, 75 µm x 50 cm column (Thermo Scientific) and eluted

over 180 min gradients optimized for each high pH reverse-phase fraction (32). Sample eluate

was electrosprayed (2000 V) into a Thermo Scientific Orbitrap Eclipse mass spectrometer for

analysis. MS1 spectra were acquired at a resolving power of 120,000. MS2 spectra were acquired

in the Ion Trap with CID (35%) in centroid mode. Real-time search (max search time = 34 s; max

missed cleavages = 1; Xcorr = 1; dCn = 0.1; ppm = 5) was used to select ions for synchronous

precursor selection for MS3. MS3 spectra were acquired in the Orbitrap with HCD (60%) with an

isolation window = 0.7 m/z and a resolving power of 60,000, and a max injection time of 400ms.

For phosphopeptide quantification, 4 µl (out of 20) of the TMT labeled phosphopeptide

enrichments were loaded onto a heated PepMap RSLC C18 2 µm, 100 angstrom, 75 µm x 50 cm

column (Thermo Scientific) and eluted over a 180 min gradient: 1 min 2% B, 5 min 5% B, 160 min

25% B, 180 min 35% B. Sample eluate was electrosprayed (2000 V) into a Thermo Scientific

Orbitrap Eclipse mass spectrometer for analysis. MS1 spectra were acquired at a resolving power

of 120,000. MS2 spectra were acquired in the Orbitrap with HCD (38%) in centroid mode with an

isolation window = 0.4 m/z, a resolving power of 60,000, and a max injection time of 350 ms.

The raw MS files were processed using Proteome Discoverer version 2.4 (Thermo Scientific,

Waltham, MA). MS spectra were searched against the *Homo sapiens* Uniprot/SwissProt

database. SEQUEST search engine was used (enzyme = trypsin, max. missed cleavage = 4, min.

peptide length = 6, precursor tolerance = 10 ppm). Static modifications include carbamidomethyl

(C, +57.021 Da), and TMT labeling (N-term and K,+304.207 Da for TMTpro16). Dynamic

modifications include oxidation (M, +15.995 Da), Phosphorylation (S, T, Y,+79.966 Da, only for

phosphopeptide dataset), acetylation (N-term,+ 42.011 Da), Met-loss (N-term, –131.040 Da), and

Met-loss + Acetyl (N-term, –89.030 Da). PSMs were filtered by the Percolator node (max Delta

Cn = 0y .05, target FDR (strict) = 0.01, and target FDR (relaxed)=0.05). Proteins were identified

with a minimum of 1 unique peptide and protein-level combined q values < 0.05. Reporter ion

quantification was based on corrected S/N values with the following settings: integration tolerance

= 20 ppm, method = most confident centroid, co-isolation threshold = 70, and SPS mass matches

= 65.

***Bioinformatic processing***

Protein intensity data values were first normalized to adjust each TMT experiment to an equal

signal per channel. A global scaling value and the internal reference scaling factor were then

calculated (33), after which a median normalization was performed and then log2-transformed.

Further analysis of the data indicated a strong batch effect associated to the multiplex labeling,

which we ameliorated with the function removeBatchEffect in the R package limma (v3.58.0) (34).

The experimental variability associated to the genotype, day, isogenic cell line, and differentiation

variables were retained in the design matrix. Only the proteins (Supplementary Table 2 and 3)

and phosphopeptides (Supplementary Table 4 and 5) present in all plexes were used for

downstream analysis. This stringent filter was implemented to better control batch effects and

addressed the impossibility of differentiating the missing values from non-expressed proteins.

Despite this stringent filter, we captured a diverse pool of proteins and phosphopeptides similar

to other publicly available datasets (27).

*Time series clustering*. We used the Short Time-series Expression Miner software (STEM,

v1.3.13;) (35), which compares and assigns each expression profile from the dataset to a set of

distinct and representative models of temporal expression profiles based on the total timepoints.

These model profiles are independent of the data and correspond to possible profiles of a protein's

change in expression over time. For its implementation, the batch-corrected normalized protein

abundances in the organoids were first log2-transformed and averaged by isogenic line and

timepoint. We also used a fetal brain proteome previously reported by Djuric *et al*. (36) and

selected the prenatal samples from all reported brain regions (ventricular zone, intermediate zone,

subplate zone and cortex). The fetal brain samples were processed using each region as a

replicate for clustering. The STEM algorithm then transforms all trends to start at 0, generating a

vector $(0, v_1 — v_0, v_2 — v_0, …, v_n — v_0)$ for each protein. Proteins are then assigned to the model

profile to which their time series most closely matches based on correlation coefficient. The

number of proteins assigned to each model profile is then tested for statistical significance against

the number of proteins expected to be assigned to a profile. This number is estimated by randomly

permuting the original time points, renormalizing the protein abundances, and then assigning

them to the most closely matching model profiles. The average number of proteins assigned to a

model profile over all permutations was used as the estimate of the expected number of genes

assigned to the profile. This process was performed using a significance level of 0.05 of the

Bonferroni-corrected p-value. The values used for each parameter are presented in

Supplementary Table 6. Several preliminary runs were performed to select the values that both

offered clustering of most proteins and diversity of profiles, while also controlling for the number

of highly similar profiles. Protein overlap among analyzed datasets were performed using the

packages GeneOverlap (v.1.40.0) (37) and UpSetR (v.1.4.0) (38).

*Functional processes – Gene enrichment analysis.* Testing for enrichment was performed with

the online tool Metascape (v3.5.20240901) (39), using a hypergeometric test followed by the

Benjamini-Hochberg *p*-value correction, selecting statistically significant terms at a cutoff value

below -1.3 Log(q-value) (39), and then simplifying redundancy in the resulting lists using rrvgo in

R (v1.16) (40). Enrichment tests were performed using the Gene Ontology (GO) biological

processes (41, 42) and the gene-disease association DisGeNET (43) databases. Visualizations

were performed using webr (v.0.1.5) (44) and ggplot2 functions (v3.5.1) (45). The GO-term

network was performed in Metascape. In brief, the 20 top-score clusters (lowest p-values) were

selected from each dysregulated expression trend and then used the Kappa similarity statistic

above 0.3 to connect similar terms (39). The resulting network was then edited in Cytoscape

(v3.10.2) (46). Additionally, we queried the SFARI Gene database (2024 Q3 release) (47) to

detect dysregulated genes that have also been linked to ASD.

**Results**

***Brain organoids capture proteomic developmental trajectories and processes observed in***

***fetal brain***

Proteins are the ultimate effectors of the cellular processes, thus, here we characterized the

proteome of iPSC-derived brain organoids to better understand how p.R203W PACS1 impacts

neurodevelopment. Organoids were collected at distinct differentiation time points: day 21, when

proliferating neural precursors predominate; day 40, when some progenitors have transitioned

into intermediate progenitors and newly born neurons; and day 90, when post-mitotic neurons are

the most abundant population (Fig. 1A).

We utilized TMT mass-spectrometry to detect 3,766 unique proteins in our organoids

across stages, with ~60% overlapping with proteins present in fetal brains (Fig. S1A). Most

overlap was observed within organoids and brain samples from post-conception week 16 through

26, with protein abundance correlation increasing with organoid maturation (Fig. S1B). We first

aimed to understand the proteomic landscape of maturing organoids derived from neurotypical

iPSCs. We identified groups of proteins with common temporal trajectories by performing time

series and clustering analysis. We classified proteins into three temporal patterns: upregulation,

downregulation, and stable abundance over time. The up and downregulated patterns were

further clustered into two main profiles: 1) continuous protein change, with steady

increase/decrease in abundance (clusters "steady-up" and "steady-down"); and 2) delayed trend,

where protein abundance remained stable from day 21 to day 40, and then increased/decreased

at day 90 (clusters "delayed-up" and "delayed-down") (Fig. 1B; Supplementary Table 6).

Next, we used gene ontology (GO) analysis to determine the biological processes involved

within these distinct patterns (Supplementary Table 7). As expected, many developmental

processes changed with time: cell migration, neuron projection, and synaptic function were

upregulated (Fig. 1C, outer circle), while neurogenesis was downregulated (Fig. 1D, outer circle).

This is consistent with the differentiation from neural precursor to neurons as the organoids

mature. Within the upregulated clusters, the steady-up contained most of the neurodevelopmental

processes, while the delayed-up was mostly associated with signaling. Likewise, clusters steady-

down and delayed-down shared a downregulation of cell cycle and differentiation (Fig.1D). Stable

proteins were associated with basic functions like primary metabolism, regulation of gene

expression, and ribosome/translation-related processes (Fig. S1C).

Finally, we compared the observed organoid processes with the developing fetal brain

dynamics obtained from a reference dataset (36) (Fig. S1D and Supplementary Table 8). We

found significant overlaps between clusters in organoids with the same temporal patterns in the

fetal brain (Supplementary Table 9). In addition, most of the enriched processes during brain

development were captured in the corresponding pattern by the organoids. These results confirm

that our brain organoids recapitulate early stages of brain development, as previously reported

(27), supporting their use to model protein dysregulation in PACS1 syndrome.

***PACS1 p.R203W pathogenic variant putatively accelerates axonal and synaptic maturation***

We then profiled the proteome of PACS1 syndrome patient-derived brain organoids to evaluate

the effects of the PACS1 p.R203W pathogenic variant on protein regulation during development

(Fig. 1A). Maturation time was the strongest driver of variance when comparing PACS1$^{(+/+)}$ and

PACS1$^{(+/R203W)}$ organoids at days 21, 40, and 90 using principal component analysis (PCA, 42%

of variation) (Fig. S2). We investigated whether the presence of the PACS1 variant impacts

groups of proteins that share common temporal trajectories. PACS1$^{(+/R203W)}$ organoids also

presented with the three major patterns (upregulation, downregulation, and stable) and profiles

(steady vs. delayed) observed in controls (Fig. 2A, Supplementary Table 10). However, several

proteins switched abundance patterns (Fig. 2B). The profile rearrangement resulted in four

additional clusters, for a total of eight clusters found in PACS1$^{(+/R203W)}$ organoids. These clusters

included proteins that were up or down-regulated when compared to the stable pattern in controls

("stable-up" and "stable-down" clusters), or that lost the regulation observed in controls and

remained stable over maturation ("up-stable" and "down-stable" clusters; Supplementary Table

11).

The stable-up cluster from PACS1$^{(+/R203W)}$ organoids had an enrichment in processes like

axon projection, neuron projection guidance, and post-synapse organization (Fig. 2C). Proteins

that gain upregulation included the microtubule-associated protein MAP1S, required for neurite

extension (48); the transmembrane ligand ephrin B3, involved in synapse and spine formation

(49); and the histone deacetylase HDAC5, a potential PACS1 interactor involved in axon growth

and regeneration (50) (Supplementary Table 12). On the other hand, the stable-down cluster

showed enrichment of known PACS1 functions associated to viral transcripts trafficking (18), as

well as translation at synapses and RNA processing. Proteins that gain downregulation included

ribosomal subunit proteins such as RPL15 and RPS10, and spliceosomal complex assembly

proteins. In this cluster we also detected known PACS1 interactors including the phosphatase

subunit PPP1CA, essential for cell division and a regulator of neuronal differentiation and synaptic

transmission (51); and the IGF2R/CI-MPR receptor, involved in intracellular trafficking of

lysosomal enzymes (52) and recently implicated in neurodegenerative disorders (53).

The down-stable cluster was also enriched in translational elongation and alternative

splicing, with the broader "protein translation" category overlapping with the stable-down cluster.

However, the nested terms in this category suggest that there could be a shift in the location

where active translation is occurring in PACS1$^{(+/R203W)}$ organoids, decreasing at the synapse while

retaining it at the mitochondria. Proteins that lost their temporal downregulation and remained

stable over time included known PACS1 interactor CSNK2A2, the catalytic subunit alpha 2 of

casein kinase 2 (CK2). Interestingly, CSNK2A2 is not only activated by PACS1 but also enhances

PACS1 cargo-shuttling function by direct phosphorylation (52).

Finally, the up-stable cluster was enriched in processes related to vesicle recycling and

cargo loading at synapses, and Golgi organization and transport. Proteins that lost their temporal

upregulation and remained stable in PACS1$^{(+/R203W)}$ organoids included the secreted glycoprotein

WNT7A, involved in regulating protein localization at the pre-synapse and preferentially

stimulating excitatory synapse formation and function (54); as well as the transmembrane

trafficking proteins TMED2, TMED3, and TMED4, required for Golgi apparatus organization and

localization of proteins to the plasma membrane (55). As in the other clusters, we found a known

PACS1 interactor, the Golgi-associated gamma-adaptin GGA3. Notably, GGA3-mediated

trafficking is regulated not only by its interaction with PACS1 but also with CK2 (that lost its

temporal regulation and remained stable in PACS1[(+/R203W)] organoids) (52). Other proteins that

lost regulation over time, including the kinesin KIF5C, the ATPase TOR1A, and a component of

the coat protein complex II (COPII) SEC31A, are implicated in synaptic maturation through vesicle

loading and recycling.

Our results uncovered dysregulation of known interactors of PACS1 on the protein level

that were not resolved on the transcriptome level (23). Results suggest that the PACS1[(+/R203W)]

organoids could present with accelerated development, with the upregulation of processes related

to establishing cell polarity, axon growth, and synapse maturation. However, even if the

PACS1[(+/R203W)] organoids present with early synaptic formation signatures, they also lost

regulation of many synaptic processes, like vesicle loading and recycling, that could lead to

defective and/or incomplete synaptic function (Supplementary Table 13).

***Temporally regulated phosphorylation of proteins related to axonal and synaptic***

***processes are also impacted in PACS1[(+/R203W)] brain organoids***

Post-translational phosphorylation is a main regulator of protein function. Thus, we analyzed

whether the PACS1[(+/R203W)] organoids also presented with alterations in phosphorylation during

maturation. Quantification of phospho-peptides in organoids resolved a total of 4,238

phosphorylation sites. The time series and clustering analysis uncovered the same eight clusters

and stable patterns that resulted from the protein abundance analysis (Fig. 3A, Supplementary

Tables 14-16). However, an additional distinct cluster composed of peptides that acquired

phosphorylated sites between days 21 and 40 ("early-up" cluster) was observed in PACS1[(+/R203W)]

organoids (Fig. 3B).

A total of 554 phosphorylation sites on 347 unique proteins were dysregulated in

PACS1[(+/R203W)] organoids (Supplementary table 17). Examples include two PACS1 known

interactors, PACS2 (PACS1 paralog) and HDAC1. PACS1[(+/R203W)] organoids presented with

increased PACS2-S437 phosphorylation, which is required for PACS2-mediated trafficking and

repression of apoptosis (56). HDAC1 had decreased phosphorylation at residues S421 and S423

which is essential for its repressive function (57). Interestingly, both serine residues are also

phosphorylated by CK2 (found to have lost its temporal regulation and remaining stable in

PACS1$^{(+/R203W)}$ organoids).

Because phosphorylation could result either in an induction or inhibition of protein function,

and the activation status of a protein itself could have diverse effects on pathway activity, we used

the Phosphosite database to understand the functional impact of the observed dysregulated

phosphorylation sites (58). Phosphosite contained annotations on 51 out of 554 phosphorylation

sites across 41 unique proteins, with few proteins having multiple phosphorylated sites (Fig. S3,

Supplementary Table 19). To perform the functional enrichment analysis, we grouped

phosphorylated proteins that gained (clusters: stable -up and -down) or lost (clusters up- and

down-stable) phosphorylation regulation in contrast to the controls. Although the early-up cluster

also gained upregulation, due to its uniqueness to this dataset, we performed the functional

enrichment analysis separately (Fig. 3C, Supplementary Table 18).

The phosphorylated proteins that gained up/down- regulation in the PACS1$^{(+/R203W)}$

organoids over time while remaining stable in controls are involved in axonogenesis, as well as

synaptic function and organization processes (Fig. 3C, orange pie plot). Examples of these

proteins include synapsin1 (SYN1), implicated in both synaptogenesis and neurotransmitter

release, whose phosphorylation at residues S568/S570 causes conformational changes that

prevents its binding to actin or synaptic vesicles (59); synaptojanin1 (SYNJ1) which has a

prominent role in synaptic vesicle recycling (60); the beta subunit of the AP-3 protein complex, a

known PACS1 interactor, that helps form clathrin-coated synaptic vesicles (60); and DNM1-S774,

which regulates clathrin-mediated endocytosis through its activation by dephosphorylation (61).

Similarly, as observed in the other gain of regulation clusters, the early-up cluster had

enrichment for axon-related function and organization, but showed novel terms related to

signaling through semaphorin 3 (SEMA3E) (Fig. 3C, green pie plot). This result is in consensus

with single-cell RNA sequencing analysis, which identified *SEMA3E* to be upregulated in

PACS1$^{(+/R203W)}$ brain organoids (23). Our proteomic dataset shows that while there are no changes

in SEMA3E, several proteins involved in its signaling pathway are phosphorylated, including

CRMP1, DPYSL2 and DPYSL3. DPYSL2-S522 phosphorylation by CDK5 is involved in axonal

guidance and spine development (62).

Meanwhile, intracellular protein transport was the most enriched process in the clusters

that lost regulation (Fig. 3C, brown pie plot). Proteins that remained stable in PACS1$^{(+/R203W)}$

organoids include GLI3, which has a dual function as a transcriptional activator and a repressor

of the sonic hedgehog (Shh) pathway dependent on phosphorylation (63); and LIN28A, an RNA-

binding protein that is localized to the peri-endoplasmic reticulum and has been recently described

as a global suppressor of genes in the secretory pathway (64).

***Dysregulated protein abundance and phosphorylation status in PACS1$^{(+/R203W)}$ organoids***

***result in convergent processes and overlap with known NDD-related genes***

We next analyzed the overlap of protein identities with dysregulated abundance and

phosphorylation status in PACS1$^{(+/R203W)}$ organoids. Surprisingly, we observed a low overlap of

both the total and dysregulated proteins identified in both datasets (Fig. 4A, B). Nevertheless,

most affected processes converged (Fig. S4). For example, axon- and synapse- related

processes, as pointed out in the previous sections, are enriched due to dysregulated abundance

and phosphorylation status over time in PACS1$^{(+/R203W)}$ organoids when compared to controls

(purple and yellow). This is the case for most of the observed processes including protein

transport, cytoskeleton organization, and cell cycle. Some exceptions include processes like

energy metabolism that comes from the dysregulation of protein abundance (purple) and neuron

migration, which is composed of proteins showing a gain of upregulation in phosphorylation status

over time (yellow). These results suggest that PACS1$^{(+/R203W)}$ organoids development is disrupted

at different regulation levels.

Next, we investigated whether there are convergent dysregulated proteins in PACS1

syndrome compared to other NDDs. We performed gene-set enrichment analysis on the

dysregulated proteins using the DISGENET and the Simons Foundation ASD (SFARI) gene

databases. A total of 241 genes (20%) dysregulated in the PACS1$^{(+/R203W)}$ organoids compared to

controls are also reported in these databases. In DISGENET, we found 160 genes that showed

a significant enrichment in multiple nervous system disorders (Fig. 4A; Supplementary Table 20).

These disorders overlapped with phenotypes observed in PACS1 syndrome patients like ASD,

ID, seizures, and abnormal morphogenesis of the nervous system or facial features (3, 65), as

well as phenotypes yet to be reported like neurodegeneration. Finally, 117 of the genes that we

identified to be dysregulated in PACS1$^{(+/R203W)}$ organoids are also reported as ASD-risk genes in

the SFARI database (Fig. 4B, Supplementary Table 21). These results support the overlap of

PACS1 syndrome and dysregulated processes with ASD and expand on phenotypes that patients

could manifest later in life.

**Discussion**

NDDs are highly heterogenous and can arise from variants across hundreds of genes (57,

58). However, several limitations have hampered our understanding of the mechanisms by which

these variants act, limiting translation of genetic findings into the clinic (59, 60). Here we leveraged

the potential of brain organoids to model neurodevelopment and gain insights into the pathogenic

mechanisms of PACS1 syndrome. By using mass spectrometry with TMT, we identified 7,230

unique proteins, of which 3,677 were quantified with 100% present call in our samples, a yield

consistent with findings from other proteomic studies involving brain organoids (27, 61, 62).

Moreover, we observed temporal trajectories that reflected the transitions from proliferating

progenitor cells towards mature neuronal lineages, mimicking the temporal dynamics observed in

the fetal brain.

PACS1[(+/R203W)] organoids presented with dysregulated abundance and phosphorylation

status of temporally regulated proteins, including PACS1 known interactors. Despite the low

overlap between expression and phosphorylation alterations in PACS1[(+/R203W)], most dysregulated

processes converged. We observed upregulation of processes related to axonogenesis and

synapse maturation, suggesting accelerated development in PACS1[(+/R203W)] organoids. However,

although the PACS1[(+/R203W)] organoids presented with early synaptic formation signatures, the

regulation of multiple synapse and vesicle-related proteins are affected and many synaptic

maturation processes are lost, possibly disrupting proper synaptic function. This is consistent with

the observed phenotype in mice overexpressing PACS1 p.R203W (22); and the asynchrony and

an increased network burst duration reported in neurons harboring the p.R203W derived from

patient iPSC lines (23). Interestingly the synaptic terms obtained from protein abundance and

phosphorylation were also obtained from transcriptomic data, suggesting that the PACS1

p.R203W variant might affect multiple regulatory levels.

Proteomic approaches are usually bulk and lack single-cell resolution, our experimental

set up also lacks cells that emerge later in development (astrocytes and oligodendrocytes) or

arise from non-ectoderm layers (microglia). Other model systems should be explored to determine

how these cells contribute to PACS1 syndrome pathogenesis. Nevertheless, we identified multiple

proteins that displayed temporal changes in abundance and phosphorylation status in developing

PACS1[(+/R203W)] brain organoids with respect to controls. Interestingly, most proteins that we found

dysregulated in PACS1[(+/R203W)] organoids have also been linked to NDDs (63-65). Further

mechanistic analysis is required, however, to understand if these effects are direct or indirect.

Most proteins that lost their upregulation over time, and their resulting dysregulated

processes, could be linked to known PACS1 functions. For example, PACS1's role in protein

sorting explains the overrepresentation of vesicle-related processes in the up-stable cluster.

Moreover, many established PACS1 interactors were uncovered as dysregulated proteins were

not only related to known PACS1 functions but also to several known PACS1 interactors, such

as its close paralog, PACS2. PACS1 and PACS2 evolutionarily emerged as a duplication in higher

metazoans from an ancient PACS gene (66). PACS1 function primarily involves sorting from

endosomes to the tGN and is dependent on AP-1 and AP-3 clathrin adaptors (which also showed

decreased abundance over time in PACS1$^{(+/R203W)}$ organoids) (15). Meanwhile, PACS2 evolved

cargo-sorting functions from the Golgi to ER and vesicular coat preference for COPII. We detected

a gain of PACS2 regulation through increased phosphorylation at S437, which favors PACS2

trafficking function (47). Thus, this increased phosphorylated PACS2 could explain the

dysregulated vesicle trafficking and upregulation of the COPII-coated vesicle budding in the Golgi

transport observed in PACS1$^{(+/R203W)}$ organoids. On the other hand, it could also explain the

observed loss of upregulation of apoptotic proteins (Figure 2C). Interestingly, a recurrent *de novo*

missense mutation in the regulatory middle region of PACS2 is associated with PACS2

Syndrome, another NDD (67). Further research aimed to understand the interconnection of

PACS1 and PACS2 function, as well as their dysregulation and compensatory effects in disease,

will help shed light on the pathogenesis of both PACS1 and PACS2 syndromes.

The processes that lost downregulation over time were related to recently described

PACS1 functions, including proteins related to the maintenance of genome stability and DNA

repair via interaction with HDAC2, HDAC3, and HDAC6 (20, 22). Although we did not observe

protein abundance and phosphorylation status changes in HDAC2, 3, or 6, we found that HDAC5

gain upregulation and HDAC1 had decreased phosphorylation over maturation. This is consistent

with the possible pathogenic interaction between PACS1 p.R203W and HDAC6 (22). HDAC1

interacts with RB1 and SMARCA4 (both dysregulated at the phosphorylation level) in a repressive

complex in resting neurons that is abolished in a calcium-dependent manner to enable

transcription (68). This suggests that the PACS1 p.R203W variant could affect transcription

regulation by destabilizing the repressive complex formed by HDAC1, RB1, and SMARCA4.

Processes unrelated to known PACS1 functions also emerged, like the enrichment in

cluster down-stable associated to regulating protein translation. Although there is growing

evidence of the impact of protein translation on the pathogenesis of neurodevelopmental

disorders including ASD and fragile X syndrome (69), this is the first time it has been implicated

in PACS1 syndrome pathogenesis. Moreover, PACS1$^{(+/R203W)}$ organoids presented with changes

in genes reported to also be dysregulated in other neurological diseases. The presence of the

PACS1 p.R203W variant, at the proteomic level, seems to affect processes related to the clinical

features that patients manifest, including ASD, ID, and seizures (3, 56), as well as phenotypes

yet to be reported like neurodegeneration. Additional models are required to test mechanisms of

PACS1 syndrome outside the cerebral cortex, like the observed hypoplasia of the cerebellum and

the corpus callosum, or the dysmorphic facial features.

The processes that seem unrelated to PACS1 function but with observed upregulation in

the PACS1$^{(+/R203W)}$ organoids should be further explored. Because current evidence suggests that

the PACS1 p.R203W pathogenic variant could result in a gain of function (22, 23, 70), there is

increased interest in generating targeted therapeutic strategies aimed to suppress its toxic effects

(22, 26). Thus, the processes that seem unrelated to PACS1 function but with observed

upregulation in the PACS1$^{(+/R203W)}$ organoids should be further explored. Understanding whether

the proteomic dysregulation results from the toxicity associated to the variant or as a

compensatory mechanism to maintain cellular homeostasis would be especially relevant. Finally,

the neurological phenotypes associated with dysregulated protein abundance and

phosphorylation in PACS1 p.R203W organoids are consistent with clinical characteristics

observed in PACS1 syndrome patients. Further downstream analyses into understanding the

converging dysregulated processes could offer novel therapeutic strategies for PACS1 syndrome.

Our results highlight that proteomic analyses not only enhance our understanding of general NDD

mechanisms by complementing transcriptomic studies, but also could uncover additional targets,

facilitating therapy development.

***Acknowledgements***

This research was supported in part through the computational resources and staff contributions provided for the Quest high performance computing facility at Northwestern University which is jointly supported by the Office of the Provost, the Office for Research, and Northwestern University Information Technology.

***Authors' contributions***

This work was conceptualized by AGG. XGM performed proteomics analysis with help from SR, BD, JG, and MLM, and generated the figures with the help of AW. LR cultured the organoids, collected samples, and prepared them for proteomic analysis. AW, ALS, and JR performed wet lab experiments (immunohistochemistry, imaging) and data analysis. Writing was done by XGM and edited by AGG, MLM, LR, JR and ALS. All authors read and approved the final manuscript.

***Funding***

This work was funded by the PACS1 Syndrome Research Foundation (AGG) and by National Institutes of Health NINDS R01NS123163 (AGG) as well as R01MH125235 (M.L.M and B.D.), R01MH118497 (MLM). Confocal microscopy was done at the Northwestern University Center for Advanced Microscopy funded by NCI CCSG P30 CA060553.

***Competing interests***

The authors have nothing to disclose.

***Availability of data and materials***

478    Data is provided within the manuscript or supplementary information files and from the

479    corresponding author on reasonable request.

**References**

1.      Castellani CA, Arking DE. High-Risk, High-Reward Genetics in ASD. Neuron. 2020;105(3):407-10.
2.      Geschwind DH, Flint J. Genetics and genomics of psychiatric disease. Science. 2015;349(6255):1489-94.
3.      Schuurs-Hoeijmakers JH, Landsverk ML, Foulds N, Kukolich MK, Gavrilova RH, Greville-Heygate S, et al. Clinical delineation of the PACS1-related syndrome--Report on 19 patients. Am J Med Genet A. 2016;170(3):670-5.
4.      Schuurs-Hoeijmakers JH, Oh EC, Vissers LE, Swinkels ME, Gilissen C, Willemsen MA, et al. Recurrent de novo mutations in PACS1 cause defective cranial-neural-crest migration and define a recognizable intellectual-disability syndrome. Am J Hum Genet. 2012;91(6):1122-7.
5.      Seto MT, Bertoli-Avella AM, Cheung KW, Chan KY, Yeung KS, Fung JL, et al. Prenatal and postnatal diagnosis of Schuurs-Hoeijmakers syndrome: Case series and review of the literature. Am J Med Genet A. 2021;185(2):384-9.
6.      Thomas G, Aslan JE, Thomas L, Shinde P, Shinde U, Simmen T. Caught in the act - protein adaptation and the expanding roles of the PACS proteins in tissue homeostasis and disease. J Cell Sci. 2017;130(11):1865-76.
7.      Scott GK, Gu F, Crump CM, Thomas L, Wan L, Xiang Y, et al. The phosphorylation state of an autoregulatory domain controls PACS-1-directed protein traffic. EMBO J. 2003;22(23):6234-44.
8.      Wan L, Molloy SS, Thomas L, Liu G, Xiang Y, Rybak SL, et al. PACS-1 defines a novel gene family of cytosolic sorting proteins required for trans-Golgi network localization. Cell. 1998;94(2):205-16.
9.      Crump CM, Xiang Y, Thomas L, Gu F, Austin C, Tooze SA, et al. PACS-1 binding to adaptors is required for acidic cluster motif-mediated protein traffic. EMBO J. 2001;20(9):2191-201.
10.     Hinners I, Wendler F, Fei H, Thomas L, Thomas G, Tooze SA. AP-1 recruitment to VAMP4 is modulated by phosphorylation-dependent binding of PACS-1. EMBO Rep. 2003;4(12):1182-9.
11.     Gu F, Crump CM, Thomas G. Trans-Golgi network sorting. Cell Mol Life Sci. 2001;58(8):1067-84.
12.     Molloy SS, Thomas L, Kamibayashi C, Mumby MC, Thomas G. Regulation of endosome sorting by a specific PP2A isoform. J Cell Biol. 1998;142(6):1399-411.
13.     Schermer B, Hopker K, Omran H, Ghenoiu C, Fliegauf M, Fekete A, et al. Phosphorylation by casein kinase 2 induces PACS-1 binding of nephrocystin and targeting to cilia. EMBO J. 2005;24(24):4415-24.
14.     Jenkins PM, Zhang L, Thomas G, Martens JR. PACS-1 mediates phosphorylation-dependent ciliary trafficking of the cyclic-nucleotide-gated channel in olfactory sensory neurons. J Neurosci. 2009;29(34):10541-51.
15.     Kottgen M, Benzing T, Simmen T, Tauber R, Buchholz B, Feliciangeli S, et al. Trafficking of TRPP2 by PACS proteins represents a novel mechanism of ion channel regulation. EMBO J. 2005;24(4):705-16.
16.     Gorini G, Ponomareva O, Shores KS, Person MD, Harris RA, Mayfield RD. Dynamin-1 co-associates with native mouse brain BKCa channels: proteomics analysis of synaptic protein complexes. FEBS Lett. 2010;584(5):845-51.
17.     Taya S, Shinoda T, Tsuboi D, Asaki J, Nagai K, Hikita T, et al. DISC1 regulates the transport of the NUDEL/LIS1/14-3-3epsilon complex through kinesin-1. J Neurosci. 2007;27(1):15-26.
18.     Liu H, Hu PW, Budhiraja S, Misra A, Couturier J, Lloyd RE, et al. PACS1 is an HIV-1 cofactor that functions in Rev-mediated nuclear export of viral RNA. Virology. 2020;540:88-96.

19.    Atkins KM, Thomas LL, Barroso-Gonzalez J, Thomas L, Auclair S, Yin J, et al. The
multifunctional sorting protein PACS-2 regulates SIRT1-mediated deacetylation of p53 to
modulate p21-dependent cell-cycle arrest. Cell Rep. 2014;8(5):1545-57.
20.    Mani C, Tripathi K, Luan S, Clark DW, Andrews JF, Vindigni A, et al. The multifunctional
protein PACS-1 is required for HDAC2- and HDAC3-dependent chromatin maturation and
genomic stability. Oncogene. 2020;39(12):2583-96.
21.    Veena MS, Raychaudhuri S, Basak SK, Venkatesan N, Kumar P, Biswas R, et al.
Dysregulation of hsa-miR-34a and hsa-miR-449a leads to overexpression of PACS-1 and loss
of DNA damage response (DDR) in cervical cancer. J Biol Chem. 2020;295(50):17169-86.
22.    Villar-Pazos S, Thomas L, Yang Y, Chen K, Lyles JB, Deitch BJ, et al. Neural deficits in
a mouse model of PACS1 syndrome are corrected with PACS1- or HDAC6-targeting therapy.
Nat Commun. 2023;14(1):6547.
23.    Rylaarsdam L, Rakotomamonjy J, Pope E, Guemez-Gamboa A. iPSC-derived models of
PACS1 syndrome reveal transcriptional and functional deficits in neuron activity. Nat Commun.
2024;15(1):827.
24.    Byrd DT, Han ZC, Piggott CA, Jin Y. PACS-1 variant protein is aberrantly localized in
Caenorhabditis elegans model of PACS1/PACS2 syndromes. Genetics. 2024;228(2).
25.    Van Nuland A, Reddy T, Quassem F, Vassalli JD, Berg AT. PACS1-
Neurodevelopmental disorder: clinical features and trial readiness. Orphanet J Rare Dis.
2021;16(1):386.
26.    Rylaarsdam L, Reddy T, Guemez-Gamboa A. In search of a cure: PACS1 Research
Foundation as a model of rare disease therapy development. Trends Genet. 2022;38(2):109-12.
27.    Melliou S, Sangster KT, Djuric U, Diamandis P. The promise of organoids for unraveling
the proteomic landscape of the developing human brain. Mol Psychiatry. 2022;27(1):73-80.
28.    Li C, Fleck JS, Martins-Costa C, Burkard TR, Themann J, Stuempflen M, et al. Single-
cell brain organoid screening identifies developmental defects in autism. Nature.
2023;621(7978):373-80.
29.    Paulsen B, Velasco S, Kedaigle AJ, Pigoni M, Quadrato G, Deo AJ, et al. Autism genes
converge on asynchronous development of shared neuron classes. Nature.
2022;602(7896):268-73.
30.    Buccitelli C, Selbach M. mRNAs, proteins and the emerging principles of gene
expression control. Nat Rev Genet. 2020;21(10):630-44.
31.    Sloan SA, Andersen J, Pasca AM, Birey F, Pasca SP. Generation and assembly of
human brain region-specific three-dimensional cultures. Nat Protoc. 2018;13(9):2062-85.
32.    Dumrongprechachan V, Salisbury RB, Butler L, MacDonald ML, Kozorovitskiy Y.
Dynamic proteomic and phosphoproteomic atlas of corticostriatal axons in neurodevelopment.
Elife. 2022;11.
33.    Plubell DL, Wilmarth PA, Zhao Y, Fenton AM, Minnier J, Reddy AP, et al. Extended
Multiplexing of Tandem Mass Tags (TMT) Labeling Reveals Age and High Fat Diet Specific
Proteome Changes in Mouse Epididymal Adipose Tissue. Mol Cell Proteomics. 2017;16(5):873-
90.
34.    Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, et al. limma powers differential
expression analyses for RNA-sequencing and microarray studies. Nucleic Acids Res.
2015;43(7):e47.
35.    Ernst J, Bar-Joseph Z. STEM: a tool for the analysis of short time series gene
expression data. BMC Bioinformatics. 2006;7:191.
36.    Djuric U, Lam KHB, Kao J, Batruch I, Jevtic S, Papaioannou MD, et al. Defining Protein
Pattern Differences Among Molecular Subtypes of Diffuse Gliomas Using Mass Spectrometry.
Mol Cell Proteomics. 2019;18(10):2029-43.
37.    Shen L, Sinai M. GeneOverlap: Test and visualize gene overlaps. R package version.
2013;1(0):444.

38.     Gehlenborg N. UpSetR: a more scalable alternative to Venn and Euler diagrams for
visualizing intersecting sets. R package version. 2019;1(0).
39.     Zhou Y, Zhou B, Pache L, Chang M, Khodabakhshi AH, Tanaseichuk O, et al.
Metascape provides a biologist-oriented resource for the analysis of systems-level datasets. Nat
Commun. 2019;10(1):1523.
40.     Sayols S. rrvgo: a Bioconductor package for interpreting lists of Gene Ontology terms.
MicroPubl Biol. 2023;2023.
41.     Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, et al. Gene ontology:
tool for the unification of biology. The Gene Ontology Consortium. Nat Genet. 2000;25(1):25-9.
42.     Gene Ontology C, Aleksander SA, Balhoff J, Carbon S, Cherry JM, Drabkin HJ, et al.
The Gene Ontology knowledgebase in 2023. Genetics. 2023;224(1).
43.     Pinero J, Ramirez-Anguita JM, Sauch-Pitarch J, Ronzano F, Centeno E, Sanz F, et al.
The DisGeNET knowledge platform for disease genomics: 2019 update. Nucleic Acids Res.
2020;48(D1):D845-D55.
44.     Moon K, Martino T. R Package 'webr': Data and Functions for Web-Based Analysis. R
package version 01. 2020;5.
45.     Wickham H. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York;
2106.
46.     Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, et al. Cytoscape: a
software environment for integrated models of biomolecular interaction networks. Genome Res.
2003;13(11):2498-504.
47.     Banerjee-Basu S, Packer A. SFARI Gene: an evolving database for the autism research
community. Dis Model Mech. 2010;3(3-4):133-5.
48.     Anastasaki C, Wegscheid ML, Hartigan K, Papke JB, Kopp ND, Chen J, et al. Human
iPSC-Derived Neurons and Cerebral Organoids Establish Differential Effects of Germline NF1
Gene Mutations. Stem Cell Reports. 2020;14(4):541-50.
49.     Hruska M, Dalva MB. Ephrin regulation of synapse formation, function and plasticity. Mol
Cell Neurosci. 2012;50(1):35-44.
50.     Cho Y, Cavalli V. HDAC5 is a novel injury-regulated tubulin deacetylase controlling axon
regeneration. EMBO J. 2012;31(14):3063-78.
51.     Foley K, McKee C, Nairn AC, Xia H. Regulation of Synaptic Transmission and Plasticity
by Protein Phosphatase 1. J Neurosci. 2021;41(14):3040-50.
52.     Scott GK, Fei H, Thomas L, Medigeshi GR, Thomas G. A PACS-1, GGA3 and CK2
complex regulates CI-MPR trafficking. EMBO J. 2006;25(19):4423-35.
53.     Deng L, Ding L, Duan X, Peng Y. Shared molecular signatures between coronavirus
infection and neurodegenerative diseases provide targets for broad-spectrum drug
development. Sci Rep. 2023;13(1):5457.
54.     Ciani L, Boyle KA, Dickins E, Sahores M, Anane D, Lopes DM, et al. Wnt7a signaling
promotes dendritic spine growth and synaptic strength through Ca(2)(+)/Calmodulin-dependent
protein kinase II. Proc Natl Acad Sci U S A. 2011;108(26):10732-7.
55.     Aber R, Chan W, Mugisha S, Jerome-Majewska LA. Transmembrane emp24 domain
proteins in development and disease. Genet Res (Camb). 2019;101:e14.
56.     Aslan JE, You H, Williamson DM, Endig J, Youker RT, Thomas L, et al. Akt and 14-3-3
control a PACS-2 homeostatic switch that integrates membrane traffic with TRAIL-induced
apoptosis. Mol Cell. 2009;34(4):497-509.
57.     Pflum MK, Tong JK, Lane WS, Schreiber SL. Histone deacetylase 1 phosphorylation
promotes enzymatic activity and complex formation. J Biol Chem. 2001;276(50):47733-41.
58.     Tsukada M, Prokscha A, Ungewickell E, Eichele G. Doublecortin association with actin
filaments is regulated by neurabin II. J Biol Chem. 2005;280(12):11361-8.
59.     Cesca F, Baldelli P, Valtorta F, Benfenati F. The synapsins: key actors of synapse
function and plasticity. Prog Neurobiol. 2010;91(4):313-48.

60.    Nakatsu F, Ohno H. Adaptor protein complexes as the key regulators of protein sorting
in the post-Golgi network. Cell Struct Funct. 2003;28(5):419-29.
61.    Srinivasan S, Burckhardt CJ, Bhave M, Chen Z, Chen PH, Wang X, et al. A
noncanonical role for dynamin-1 in regulating early stages of clathrin-mediated endocytosis in
non-neuronal cells. PLoS Biol. 2018;16(4):e2005377.
62.    Gu Y, Hamajima N, Ihara Y. Neurofibrillary tangle-associated collapsin response
mediator protein-2 (CRMP-2) is highly phosphorylated on Thr-509, Ser-518, and Ser-522.
Biochemistry. 2000;39(15):4267-75.
63.    Andreu-Cervera A, Catala M, Schneider-Maunoury S. Cilia, ciliopathies and hedgehog-
related forebrain developmental disorders. Neurobiol Dis. 2021;150:105236.
64.    Cho J, Chang H, Kwon SC, Kim B, Kim Y, Choe J, et al. LIN28A is a suppressor of ER-
associated translation in embryonic stem cells. Cell. 2012;151(4):765-77.
65.    Tenorio-Castano J, Morte B, Nevado J, Martinez-Glez V, Santos-Simarro F, Garcia-
Minaur S, et al. Schuurs-Hoeijmakers Syndrome (PACS1 Neurodevelopmental Disorder):
Seven Novel Patients and a Review. Genes (Basel). 2021;12(5).
66.    Parikshak NN, Luo R, Zhang A, Won H, Lowe JK, Chandran V, et al. Integrative
functional genomic analyses implicate specific molecular pathways and circuits in autism. Cell.
2013;155(5):1008-21.
67.    Sanders SJ, Murtha MT, Gupta AR, Murdoch JD, Raubeson MJ, Willsey AJ, et al. De
novo mutations revealed by whole-exome sequencing are strongly associated with autism.
Nature. 2012;485(7397):237-41.
68.    Jourdon A, Scuderi S, Capauto D, Abyzov A, Vaccarino FM. PsychENCODE and
beyond: transcriptomics and epigenomics of brain development and organoids.
Neuropsychopharmacology. 2021;46(1):70-85.
69.    Zhao X, Bhattacharyya A. Human Models Are Needed for Studying Human
Neurodevelopmental Disorders. Am J Hum Genet. 2018;103(6):829-57.
70.    Nascimento JM, Saia-Cereda VM, Sartore RC, da Costa RM, Schitine CS, Freitas HR,
et al. Human Cerebral Organoids and Fetal Brain Tissue Share Proteomic Similarities. Front
Cell Dev Biol. 2019;7:303.
71.    Nascimento JM, Saia-Cereda VM, Zuccoli GS, Reis-de-Oliveira G, Carregari VC, Smith
BJ, et al. Proteomic signatures of schizophrenia-sourced iPSC-derived neural cells and brain
organoids are similar to patients' postmortem brains. Cell Biosci. 2022;12(1):189.
72.    Banerjee S, Zhao Q, Wang B, Qin J, Yuan X, Lou Z, et al. A novel in-frame deletion in
KIF5C gene causes infantile onset epilepsy and psychomotor retardation. MedComm (2020).
2024;5(4):e469.
73.    Granata A, Watson R, Collinson LM, Schiavo G, Warner TT. The dystonia-associated
protein torsinA modulates synaptic vesicle recycling. J Biol Chem. 2008;283(12):7568-79.
74.    Halperin D, Kadir R, Perez Y, Drabkin M, Yogev Y, Wormser O, et al. SEC31A mutation
affects ER homeostasis, causing a neurological syndrome. J Med Genet. 2019;56(3):139-48.
75.    Youker RT, Shinde U, Day R, Thomas G. At the crossroads of homoeostasis and
disease: roles of the PACS proteins in membrane traffic and apoptosis. Biochem J.
2009;421(1):1-15.
76.    Olson HE, Jean-Marcais N, Yang E, Heron D, Tatton-Brown K, van der Zwaag PA, et al.
A Recurrent De Novo PACS2 Heterozygous Missense Variant Causes Neonatal-Onset
Developmental Epileptic Encephalopathy, Facial Dysmorphism, and Cerebellar Dysgenesis. Am
J Hum Genet. 2018;103(4):631.
77.    Qiu Z, Ghosh A. A calcium-dependent switch in a CREST-BRG1 complex regulates
activity-dependent gene expression. Neuron. 2008;60(5):775-87.
78.    Jishi A, Qi X, Miranda HC. Implications of mRNA translation dysregulation for
neurological disorders. Semin Cell Dev Biol. 2021;114:11-9.

79.     Liu Y, Ding H, Yan T, Liu L, Yu L, Huang Y, et al. A Novel Multi-Exon Deletion of PACS1
in a Three-Generation Pedigree: Supplements to PACS1 Neurodevelopmental Disorder
Spectrum. Front Genet. 2021;12:690216.

**Figures**

**Figure 1. Cortical organoids recapitulate proteome dynamics during brain development.**

A) Expression of neural precursor protein SOX2 (red) and projection neuron marker CTIP2

(magenta) in day 21, 40, and 80 organoids derived from control (up) and PACS1 p.R203W

organoids (bottom). Scale bar = 100 μm. DAPI (blue) B) Total proteins detected in organoids were

clustered by abundance changes over maturation time into three major trends: upregulated (red),

downregulated (blue) or stable (gray) (graph on the left). Further classification detected four main

abundance trends, based on the timing of change: steady over time (clusters steady -up and -

down), or delayed to day 40 (clusters delayed -up and -down) (Right). Total number of proteins

assigned to a cluster are indicated in each graph. C, D) Top GO terms upregulated (red) and

downregulated (blue) by cluster over maturation time. The inner pie plots represent processes

common to the clusters in each panel. The nested pie charts show root and branch significant

GO enrichments with the width of the pie correlating to -log(p) of the GO term. Neurodevelopment-

related processes (bold). E) Top GO terms upregulated (red) and downregulated (blue) shared

between organoids (purple) and fetal brain (green) over time. Synapse and neurodevelopment-

related processes (box). Organoid profiles correspond to clusters 1 to 4, while the fetal brain

profiles correspond to clusters A to H. A full description of all enriched processes is included in

Supplementary Table 9.

**Figure 2. Organoids carrying the PACS1 p.R203W pathogenic variant present with altered protein dynamics during maturation.** A) Alluvial plot indicating the change of profile clustering in the PACS1 p.R203W organoids (right) in contrast to the clustering in controls (left). B) Four

clusters of dysregulated proteins were identified in PACS1 p.R203W organoids. The plots are

organized into a gain or loss of regulation categories: cluster stable-up (top, red), and stable-down

(bottom, blue) gained regulation while clusters up- and down-stable lost regulation (middle, grey

tones), C) Enriched biological functions per dysregulated cluster. The nested pie charts show root

and branch significant GO enrichments and the width of the pie correlating to -log(p) of the GO

term. Neurodevelopment-related processes (bold).

**Figure 3. Dysregulation of temporal phosphorylation dynamics in PACS1 p.R203W organoids.** A) Alluvial plot indicating the change of profile clustering in the PACS1 p.R203W organoids in contrast to controls. Breaks in the y axis were introduced to improve visualization of the dysregulated clusters by removing 3,000 modifications that remained stable over time. B) Five clusters of dysregulated phosphorylation trends were observed in PACS1 p.R203W organoids

that either gained a temporal regulation or lost the original pattern observed in the controls. The

plots are organized into a gain or loss of regulation categories: clusters stable- up and -down

gained temporal regulation in contrast to the controls (orange), while clusters up- and down-

stable lost temporal regulation (gray). Early-up is a new cluster exclusive to the phosphorylation

dataset that also gained temporal regulation (green). C) Enriched biological functions per

dysregulated cluster. The nested pie charts show root and branch significant GO enrichments

and the width of the pie correlating to -log(p) of the GO term. Neurodevelopment-related

processes (bold).

**Figure 4. Protein abundance and phosphorylation sites dysregulated by PACS1 p.R203W are enriched in neurological disease-related genes.** A) Overlapping identities between all detected proteins and proteins with identified phosphorylation sites. B) Overlapping identities between dysregulated proteins and phosphorylation sites observed in the PACS1 p.R203W organoids when compared to controls. C) Gene-disease association type ontologies enriched in the PACS1 p.R203W dysregulated proteome (yellow) and phosphorylation sites (purple). Selected top neuro-related categories at a corrected p-value of 0.01. D) ASD risk genes dysregulated in organoids carrying the p.R203W pathogenic variant. Genes are grouped into the

three risk level categories determined by the SFARI database (2024 Q3 release). For both panels,

the protein abundance and phosphorylation clusters are organized by major dysregulation trend

(gain or loss) with red representing upregulation and blue representing downregulation for the

protein abundance clusters. The phosphorylation clusters are grouped by major trend of

dysregulation presented in figure 3: gain of regulation (clusters stable- up and -down) or loss of

regulation (clusters up- and down- stable), as well as cluster early-up, exclusive of the

dysregulated phosphorylation sites.

**Supplemental Material**

**Supplementary information.** Supplementary Methods and supplementary figures 1-4 containing

additional data.

**Supplementary tables.** Supplementary Tables 1-20 containing the list of proteins and

phosphorylation sites, as well as accompanying statistical analysis performed on the data.