1

PACS1 syndrome variant alters proteomic landscape of developing cortical organoids

Ximena Gomez-Maqueo[1], Lauren E Rylaarsdam [1], Ashley Woo [1], Annika L Schroder [1], Jennifer Rakotomamonjy [1], Clare L Bossert [1], Tess A Smith [1], Shelby Ruiz [2], Jordan Gilardi[2], Lambertus Klei[2], Bernie Devlin[2], Matthew L MacDonald [2],*, Alicia Guemez-Gamboa[1],*

[1]Department of Neuroscience, Feinberg School of Medicine, Northwestern University, Chicago IL, USA

[2]Department of Psychiatry, University of Pittsburgh, Pittsburgh, PA, USA

*Correspondence: alicia.guemez@northwestern.edu, mlmacdonald@gmail.com

Running title: Proteomic landscape of PACS1 syndrome organoids

Abbreviations:

Antisense oligonucleotide (ASO)

Arginine to Tryptophan substitution (p.R203W)

Autism spectrum disorder (ASD)

Furin binding region (FBR)

Gene ontology (GO)

Gene-disease association ontologies (DISGENET)

Histone deacetylase (HDAC)

Induced pluripotent stem cell (iPSC)

Intellectual disability (ID)

Neurodevelopmental disorder (NDD)

Phosphofurin Acidic Cluster Sorting protein 1 (PACS1)

Post-conception week (PWC)

Principal component analysis (PCA)

Short Time-series Expression Miner (STEM)

Simons Foundation ASD gene database (SFARI)

Single cell RNA Sequencing (scRNASeq)

tandem mass tag mass-spectrometry (TMT mass spectrometry)

trans-Golgi network (tGN)

44
45

Abstract

PACS1 syndrome is a neurodevelopmental disorder (NDD) resulting from a unique *de novo* p.R203W variant in Phosphofurin Acidic Cluster Sorting protein 1 (PACS1). *PACS1* encodes a multifunctional sorting protein required for localizing furin to the *trans*-Golgi network. Although few studies have started to investigate the impact of the PACS1 p.R203W variant, the mechanisms by which the variant affects neurodevelopment are still poorly understood. In recent years, autism spectrum disorder (ASD) patient-derived brain organoids have been increasingly used to identify pathogenic mechanisms and possible therapeutic targets. While most of these studies evaluate the mechanisms by which ASD-risk genes affect the transcriptome, studies considering the proteome are limited. Here, we examine the effect of PACS1 p.R203W on the proteomic landscape of brain organoids using tandem mass tag (TMT) mass-spectrometry. Time series analysis between PACS1$^{(+/+)}$ and PACS1$^{(+/R203W)}$ organoids uncovered several proteins with dysregulated abundance or phosphorylation status, including known PACS1 interactors. Although we observed low overlap between proteins with altered expression and phosphorylation, the resulting dysregulated processes converged. The presence of the PACS1 p.R203W variant accelerated the emergence of proteins related to synaptogenesis and impaired vesicle loading and recycling. The earlier presence of these proteins and their related processes could lead to defective and/or incomplete synaptic function. Key dysregulated proteins observed in PACS1$^{(+/R203W)}$ organoids have been associated with several neurological diseases, and many are classified as NDD-causative and ASD-risk genes. Our results highlight that proteomic analyses not only enhance our understanding of general NDD mechanisms by complementing transcriptomic studies, but could also uncover additional targets, and therefore facilitate therapy development.

Introduction

Neurodevelopmental disorders (NDD) are heterogenous and usually present with complex

etiology. Several large-scale genetic studies investigating the cause of NDDs such as autism

spectrum disorder (ASD), intellectual disability (ID), and epilepsy have been conducted, and

have lead to the identification of hundreds of risk genes. However, our understanding of the

mechanisms by which the observed genetic variants act has lagged, limiting translation of

genetic findings into clinical treatments (1, 2). Characterizing the pathogenic mechanisms of

different monogenic NDDs may reveal convergent therapy targets. PACS1 syndrome

(MIM615009) is a monogenic NDD characterized by ASD, ID, craniofacial abnormalities, and

epilepsy (3-5). PACS1 syndrome results from a recurrent *de novo* c.607C>T variant in the gene

encoding P̲hosphofurin A̲cidic C̲luster S̲orting protein 1 (PACS1). This missense variant is

present in all reported patients and results in an arginine to tryptophan (p.R203W) substitution

at residue 203, with the exception of one known arginine to glutamine p.R203Q substitution (5).

PACS1 encodes a multifunctional sorting protein required for furin localization to the

trans-Golgi network (tGN) (6, 7). PACS1 binds furin and other targets via acidic cluster sorting

motifs (8-11). In addition to tGN retrieval, PACS1 mediates the recycling of endocytosed furin

from early endosomes to the cell surface and the delivery of proteins to the primary cilium (12-

14). PACS1 has been shown to interact with over 50 other proteins (10, 12, 15-17). Although

the role of PACS1 has primarily been studied in the cytoplasm, several reports show that it also

accumulates in the nucleus (18, 19) where it mediates DNA replication repair (20, 21),

contributes to the nuclear export of viral transcripts (19), and regulates chromatin stability

through interaction with histone deacetylase (HDAC) proteins (20).

Recent reports have begun to explore the pathogenesis of the PACS1 p.R203W variant.

Mice overexpressing PACS1 p.R203W exclusively in cortical neurons present with increased

dendrite arborization and decreased postsynaptic currents, possibly due to a pathogenic

interaction between PACS1 p.R203W and HDAC6. Moreover, antisense oligonucleotides (ASO)

directed towards both PACS1 and HDAC6 ameliorate these defects (22). Transcriptomic

analysis in patient-derived brain organoids revealed a dysregulation in cell projection

organization in immature PACS1 p.R203W neurons, while mature neurons exhibited an

upregulation in synaptic signaling processes when compared to isogenic controls. Subsequent

characterization of spontaneous electrical activity revealed that induced pluripotent stem cell

(iPSC)-derived neurons harboring the p.R203W variant display asynchrony and an increased

network burst duration (23). Finally, a study using *C. elegans* showed that the PACS1

syndrome-associated variant altered PACS1 localization but had no effect on neuron

morphology and synaptic patterns (24). Although these studies have started to investigate the

impact of the PACS1 p.R203W variant, the mechanisms by which the variant affects

neurodevelopment are still poorly understood. This in turn results in few therapeutic options for

patients, which at present, are limited to non-invasive behavioral approaches and sensory

integration therapies to manage symptoms (25, 26).

In recent years, ASD patient-derived brain organoids have been increasingly used to

identify pathogenic mechanisms and possible therapeutic targets (27). While most of these

studies investigate the mechanisms by which ASD-risk genes affect the transcriptome (28, 29),

studies considering the proteome are limited. However, mRNA levels do not always predict

protein levels, as protein expression and degradation are regulated by mechanisms beyond

transcription (30). Here, we uncover the proteomic landscape of brain organoids during

maturation to provide further understanding into the pathogenic mechanisms of PACS1

syndrome.

Methods

Experimental Design and Statistical Rationale

The objective of this investigation is to provide further understanding of the pathogenic

mechanisms of PACS1 syndrome by investigating proteomic landscape regulation during brain

organoid maturation. We used tandem mass tag (TMT) mass-spectrometry in brain organoids

generated from control and iPSC lines carrying the unique PACS1 p.R203W variant. Organoids

were collected on days 21, 40, and 90 to better understand the maturation process. Protein and

phosphorylated peptides were processed through mass-spectrometry. Outputs were normalized

and aggregated by isogenic pairs in R using the package limma (31). Processed outputs were

then fitted into a linear model and tested for differential expression over maturation time.

Processed data were imported into the short time series (STEM) (32) pipeline for expression

trend clustering (Figure S1).

Generation of forebrain organoids from iPSCs

Three isogenic pairs of PACS1$^{(+/+)}$ and PACS1$^{(+/R203W)}$ iPSC lines were previously generated (23)

and differentiated to brain organoids using the kit based on the protocol outlines by Sloan et al.

(33) (Stem Cell Technologies, 08620). See Supplementary Methods for further details on

experimental design, organoid generation, immunostaining and western blotting.

Sample preparation for proteomic analysis

Total protein was extracted from the organoids by sonication in lysis buffer (1% sodium dodecyl

sulfate (SDS), 125 mM triethylammonium bicarbonate (TEAB), 75 mM NaCl, and HaltTM

protease and phosphatase inhibitors). Lysate samples from two additional control cell lines were

included to improve protein detection. See Supplemental Methods and Supplementary Table 1

for a detailed description of mass spectrometry sample preparation, TMT labeling, and

phosphopeptide enrichment processing.

MS data acquisition and raw data processing

For peptide/protein quantification, TMT labeled peptides (~1 μg) were loaded onto a heated

PepMap RSLC C18 2 μm, 100 angstrom, 75 μm x 50 cm column (Thermo Scientific) and eluted

over 180 min gradients optimized for each high pH reverse-phase fraction (34). Sample eluate

were electrosprayed (2000 V) into a Thermo Scientific Orbitrap Eclipse mass spectrometer for

analysis. MS1 spectra were acquired at a resolving power of 120,000. MS2 spectra were

acquired in the Ion Trap with CID (35%) in centroid mode. Real-time search (max search time =

34 s; max missed cleavages = 1; Xcorr = 1; dCn = 0.1; ppm = 5) were used to select ions for

synchronous precursor selection for MS3. MS3 spectra were acquired in the Orbitrap with HCD

(60%) with an isolation window = 0.7 m/z and a resolving power of 60,000, and a max injection

time of 400ms. For phosphopeptide quantification, 4 µl (out of 20) of the TMT labeled

phosphopeptide enrichments were loaded onto a heated PepMap RSLC C18 2 µm, 100

angstrom, 75 µm x 50 cm column (Thermo Scientific) and eluted over a 180 min gradient: 1 min

2% B, 5 min 5% B, 160 min 25% B, 180 min 35% B. Sample eluate was electrosprayed (2000

V) into a Thermo Scientific Orbitrap Eclipse mass spectrometer for analysis. MS1 spectra were

acquired at a resolving power of 120,000. MS2 spectra were acquired in the Orbitrap with HCD

(38%) in centroid mode with an isolation window = 0.4 m/z, a resolving power of 60,000, and a

max injection time of 350 ms.

The raw MS files were processed using Proteome Discoverer version 2.4 (Thermo

Scientific, Waltham, MA). MS spectra were searched against the *Homo sapiens*

Uniprot/SwissProt database. SEQUEST search engine was used (enzyme = trypsin, max.

missed cleavage = 4, min. peptide length = 6, precursor tolerance = 10 ppm). Static

modifications include carbamidomethyl (C, +57.021 Da) and TMT labeling (N-term and K,

+304.207 Da for TMTpro16). Dynamic modifications include oxidation (M, +15.995 Da),

Phosphorylation (S, T, Y,+79.966 Da, only for phosphopeptide dataset), acetylation (N-term,+

42.011 Da), Met-loss (N-term, −131.040 Da), and Met-loss + Acetyl (N-term, −89.030 Da).

PSMs were filtered by the Percolator node (max Delta Cn = 0y .05, target FDR (strict) = 0.01,

and target FDR (relaxed)=0.05). Proteins were identified with a minimum of 1 unique peptide

and protein-level combined q values < 0.05. Reporter ion quantification was based on corrected

S/N values with the following settings: integration tolerance = 20 ppm, method = most confident

centroid, co-isolation threshold = 70, and SPS mass matches = 65.

Bioinformatic processing

Protein intensity data values were first normalized to adjust each TMT experiment to an equal

signal per channel. A global scaling value and the internal reference scaling factor were then

calculated (35), after which, a median normalization was performed and then log2-transformed.

Further analysis of the data indicated a strong batch effect associated with multiplex labeling,

which we ameliorated with the function removeBatchEffect in the R package limma (v3.58.0)

(31). The experimental variability associated with genotype, day, isogenic cell line, and

differentiation variables were retained in the design matrix to test for differential expression

across genotypes and maturation time. Only the proteins (Supplementary Table 2-4) and

phosphopeptides (Supplementary Table 5-7) present in all plexes were used for downstream

analysis. This stringent filter was implemented to better control batch effects and addressed the

impossibility of differentiating the missing values from non-expressed proteins. Despite this

stringent filter, we captured a diverse pool of proteins and phosphopeptides similar to other

publicly available datasets (27).

Time series clustering. We used the Short Time-series Expression Miner software (STEM,

v1.3.13;) (32), which compares and assigns each expression profile from the dataset to a set of

distinct and representative models of temporal expression profiles based on the total timepoints.

These model profiles are independent of the data and correspond to possible profiles of a

protein change in expression over time. For its implementation, the batch-corrected normalized

protein abundances in the organoids were first log2-transformed and averaged by isogenic line

and timepoint. We also used a fetal brain proteome previously reported by Djuric *et al.* (36) and

selected the prenatal samples from all reported brain regions (ventricular zone, intermediate

zone, subplate zone, and cortex). Fetal brain samples were processed using each region as a

replicate for clustering. The STEM algorithm then transforms all trends to start at 0, generating a

vector (0, $v_1 — v_0$, $v_2 — v_0$, …, $v_n — v_0$) for each protein. Proteins are then assigned to the model

profile to which their time series most closely matches based on correlation coefficient. The

number of proteins assigned to each model profile is then tested for statistical significance

against the number of proteins expected to be assigned to a profile. This number is estimated

by randomly permuting the original time points, renormalizing the protein abundances, and then

assigning them to the most closely matching model profiles. The average number of proteins

assigned to a model profile over all permutations was used as the estimate of the expected

number of genes assigned to the profile. This process was performed using a significance level

of 0.05 of the Bonferroni-corrected p-value. The values used for each parameter are presented

in Supplementary Table 8. Several preliminary runs were performed to select the values that

both offered clustering of most proteins and diversity of profiles, while also controlling the

number of highly similar profiles. Protein overlap among analyzed datasets were performed

using the packages GeneOverlap (v.1.40.0) (37) and UpSetR (v.1.4.0) (38).

Functional processes – Gene enrichment analysis. Testing for enrichment was performed with

the online tool Metascape (v3.5.20240901) (39) using a hypergeometric test followed by the

Benjamini-Hochberg *p*-value correction, selecting statistically significant terms at a cutoff value

below -1.3 Log(q-value) (39), and then simplifying redundancy in the resulting lists using rrvgo in

R (v1.16) (40). Enrichment tests were performed using the Gene Ontology (GO) biological

processes (41, 42) and the gene-disease association (DisGeNET) (43) databases.

Visualizations were performed using webr (v.0.1.5) (44) and ggplot2 functions (v3.5.1) (45). The

GO-term network was performed in Metascape. In brief, the 20 top-score clusters (lowest p-

values) were selected from each dysregulated expression trend and then used the Kappa

similarity statistic above 0.3 to connect similar terms (39). The resulting network was then edited

in Cytoscape (v3.10.2) (46). Additionally, we consulted the SFARI Gene database (2024 Q3

release) (47) and OMIM (48) to detect dysregulated proteins that have been annotated as ASD-

risk genes and/or pathogenic variants on such genes cause NDD.

Results

Brain organoids capture proteomic developmental trajectories and processes observed

in fetal brain

Proteins are the ultimate effectors of cellular processes. Thus, here we characterized the

proteome of iPSC-derived brain organoids to better understand how p.R203W PACS1 impacts

neurodevelopment. Organoids were collected at distinct differentiation time points: day 21, when

proliferating neural precursors predominate; day 40, when some progenitors have transitioned

into intermediate progenitors and newly born neurons; and day 90, when post-mitotic neurons

are the most abundant population (Fig. 1A).

We utilized TMT mass-spectrometry to detect 3,766 unique proteins in our organoids

across stages, with ~60% overlapping proteins present in fetal brains (Fig. S2A). Most overlap

was observed within organoids and brain samples from post-conception week 16 through 26,

with protein abundance correlation increasing with organoid maturation (Fig. S2B). First, we

aimed to understand the proteomic landscape of maturing brain organoids derived from

neurotypical iPSCs. We identified groups of proteins with common temporal trajectories by

performing time series and clustering analysis. We classified proteins into three temporal

patterns: upregulation, downregulation, and stable abundance over time. The up and

downregulated patterns were further clustered into two main profiles: 1) continuous protein

change, with steady increase/decrease in abundance (clusters "steady-up" and "steady-down");

and 2) delayed trend, where protein abundance remained stable during early differentiation

(from day 21 to day 40), and subsequently increased/decreased at day 90 (clusters "delayed-

up" and "delayed-down") (Fig. 1B; Supplementary Table 9).

Next, we used gene ontology (GO) analysis to determine the biological processes

involved within these distinct patterns (Supplementary Table 10). As expected, many

developmental processes changed with time, including neuron differentiation, axon extension,

dendrite morphogenesis, synaptic organization/function upregulation (Fig. 1C), and cell cycle

and stem cell population maintenance downregulation (Fig. 1D). This is consistent with the

differentiation from neural precursors to neurons during organoid maturation. Within the

upregulated clusters, the steady-up contained most of the neurodevelopmental processes, while

the delayed-up was mostly associated with signaling. Likewise, clusters steady-down and

delayed-down shared a downregulation of cell cycle and DNA replication or repair (Fig.1D).

Stable proteins were associated with basic functions like primary metabolism, regulation of gene

expression, and ribosome/translation-related processes (Fig. S2C).

Finally, we compared the observed organoid processes with the developing fetal brain

dynamics obtained from a reference dataset (36) (Fig. S2D and Supplementary Table 11). We

found significant overlap between clusters in organoids and the fetal brain (Supplementary

Table 12). The majority of enriched processes during fetal brain development were captured in

the corresponding organoid clusters. These results confirm that our brain organoids recapitulate

early stages of brain development, as previously reported (27), supporting their use to model

protein dysregulation in PACS1 syndrome.

PACS1 p.R203W pathogenic variant putatively accelerates neural maturation

We then profiled the proteome of PACS1 syndrome patient-derived brain organoids to evaluate

the effects of the PACS1 p.R203W pathogenic variant on protein regulation during development

(Fig. 1A). Organoids carrying the p.R203W variant developed similarly to control organoids by

day 21, with only five proteins (HMGB3, HTATSF1, HM13, TCEAL3 and TCEAL5) showing

upregulation (Fig. S3A). However, upon maturation, 329 proteins in PACS1$^{(+/R203W)}$ organoids

showed dysregulation when compared to controls (Supplementary Table 13). Oxidative

phosphorylation (ATP6V0A1, CYC1, NDUFA7) and intracellular calcium ion homeostasis

(ATP1B1, CALB2, LETM1, NPTN) were among the observed enriched processes resulting from

167 upregulated proteins in organoids carrying the p.R203W (Fig S3B, Supplementary Table

14). By day 40, most dysregulated proteins were related to established PACS1 functions

relating to endoplasmic reticulum (ER)-to-Golgi transport (ARF4, SEC23A, SNAPIN).

Upregulated proteins at day 40 were involved in neuron projection development, such as the

ligand of Eph-related receptor tyrosine kinases EFNB1 and the transcription factor FOXG1. This

was confirmed to be upregulated by western blot (Fig S3C). In contrast, neuron projection

development and synapse signaling related proteins like EFNB3, NF1, VDAC3, and SV2A were

enriched by day 90. Other upregulated processes include the metabolism of amino acids

(ASS1, GLO1, CTH, HSD17B10, PLOD2, SMS), and glycerolipids (PITPNM1, AGPAT4,

TMEM68). Interestingly, the PACS1 interactors PTBP1 (49), involved in mRNA processing, and

WDR37 (50), were also upregulated (Fig S3A,B). On the other hand, 173 down-regulated

proteins were enriched in processes including cell cycle division, DNA replication, and mRNA

splicing. This further included the RNA-binding protein PCBP4, involved in mRNA stability (51)

and histone deacetylase PTBP2, which is also involved in brain development. The intermediate

filament protein NES, which plays a role in proliferation and differentiation (52), was also found

to be among the downregulated proteins. Decreased abundance of NES was confirmed by

western blot (Fig S3).

Principal component analysis (PCA) indicated maturation time to be the strongest driver

of variance when comparing PACS1$^{(+/+)}$ and PACS1$^{(+/R203W)}$ organoids at days 21, 40, and 90 (42%

of variation) (Fig. S4A). Consequently, we investigated whether the presence of the PACS1

variant impacts groups of proteins that share common temporal trajectories. PACS1$^{(+/R203W)}$

organoids also presented with the three major patterns (upregulation, downregulation, and

stable) and profiles (steady vs. delayed) observed in controls (Fig. 2A, Supplementary Table

15). However, several proteins switched abundance patterns (Fig. 2B). The profile

rearrangement resulted in four additional clusters, for a total of eight clusters found in

PACS1$^{(+/R203W)}$ organoids. Additional clusters included proteins that were up or down-regulated

when compared to the stable pattern in controls ("stable-up" and "stable-down" clusters) and

those that lost the regulation observed in controls and remained stable over maturation ("up-

stable" and "down-stable" clusters; Supplementary Table 16).

The stable-up cluster from PACS1$^{(+/R203W)}$ organoids had a strong enrichment in oxidative

phosphorylation (23 out of 33 proteins), brain development, and neuron differentiation. Proteins

that gain upregulation include glucose transporter GLUT1, transcription factor FOXG1, and

mitochondrial genes UQCRQ, VDAC1, and ATP5PF (Fig. 2C, Supplementary Table 17). The

stable-down cluster was enriched in regulation of mRNA processing and translation. Proteins

that gain downregulation include ribosomal subunit protein RPL38, alpha subunit of the

elongation factor-1 complex EEF1A1, and developmental proteins Nestin (52), BRD2 (53), and

Wnt family member 7B WNT7B (54).

The down-stable cluster was enriched in the metabolism of the nucleotides process.

Proteins related to metabolism that lost their down regulation and remained stable over time

have also been associated with brain development and function. These proteins include,

ribokinase (RBKS), implicated in glucose metabolic disorders and recently associated to

psychiatric disorders (55), Phosphoribosylpyrophosphate synthetase 1 (PRPS1), implicated in

multiple syndromic and non-syndromic diseases causing hearing loss, hypotonia, and ataxia

(56), as well as catalase (CAT), implicated in oxidative stress-related diseases and ASD (57).

Finally, the up-stable cluster was comprised of eight proteins which lost their temporal

upregulation and remained stable in PACS1$^{(+/R203W)}$ organoids, with no significant enrichment. In

this group, however, several proteins have relevant functions in the brain, such as dendrite

morphogenesis (SS18L1/CREST) (58), autophagy and neuron projection regeneration after

injury (VIM) (59), and regulation of intracellular pH homeostasis (CA2) (60).

Our results uncovered the dysregulation of known PACS1 interactors at the protein level

that were not resolved on the transcriptome level (23). Results suggest that the PACS1$^{(+/R203W)}$

organoids undergo accelerated development, with the upregulation of processes related to

known PACS1 functions (i.e., Golgi to ER vesicle transport). However, even if the PACS1$^{(+/R203W)}$

organoids present with early neural maturation signatures, they also lost regulation of many

mRNA related processes, like transport and processing, which could lead to defective and/or

incomplete differentiation (Supplementary Table 17).

Temporally regulated phosphorylation of proteins related to axonal and synaptic

processes are also impacted in PACS1$^{(+/R203W)}$ brain organoids

Post-translational phosphorylation is a main regulator of protein function. Thus, we analyzed

whether the PACS1$^{(+/R203W)}$ organoids also presented with alterations in phosphorylation during

maturation. Quantification of phospho-peptides in organoids resolved a total of 4,238

phosphorylation sites (Supplementary Table 5). The time series and clustering analysis

uncovered the same eight clusters and stable patterns that resulted from the protein abundance

analysis (Fig. 3A, Supplementary Tables 18-20). However, an additional distinct cluster

composed of peptides that acquired phosphorylated sites between days 21 and 40 ("early-up"

cluster) was observed in PACS1$^{(+/R203W)}$ organoids (Fig. 3B).

A total of 530 phosphorylation sites on 325 unique proteins were dysregulated in

PACS1$^{(+/R203W)}$ organoids (Supplementary table 20). Examples include two PACS1-known

interactors, PACS2 (PACS1 paralog) and HDAC1. PACS1$^{(+/R203W)}$ organoids presented with

increased PACS2-S437 phosphorylation, which is required for PACS2-mediated trafficking and

repression of apoptosis (61). HDAC1 had decreased phosphorylation at residues S421 and

S423, which is essential for its repressive function (62). Interestingly, both serine residues are

also phosphorylated by CK2 (found to have lost its temporal regulation and remained stable in

PACS1$^{(+/R203W)}$ organoids).

Because phosphorylation could result either in an induction or inhibition of protein

function, and the activation status of a protein itself could have diverse effects on pathway

activity, we used the Phosphosite database to understand the functional impact of the observed

dysregulated phosphorylation sites (63). Phosphosite contained annotations on 51 out of 530

phosphorylation sites across 41 unique proteins, with few proteins having multiple

phosphorylated sites (Fig. S5, Supplementary Table 21). To perform the functional enrichment

analysis, we grouped phosphorylated proteins that gained (clusters: stable -up and -down) or

lost (clusters up- and down-stable) phosphorylation regulation in contrast to the controls.

Although the early-up cluster also gained upregulation, due to its uniqueness to this dataset, we

performed the functional enrichment analysis separately (Fig. 3C, Supplementary Table 22).

The phosphorylated proteins that gained up/down- regulation in the PACS1$^{(+/R203W)}$

organoids over time while remaining stable in controls are involved in axonogenesis, as well as

synaptic function and organization processes (Fig. 3C, orange pie plot). Examples of these

proteins include synapsin1 (SYN1), implicated in both synaptogenesis and neurotransmitter

release, whose phosphorylation at residues S568/S570 causes conformational changes that

prevents its binding to actin or synaptic vesicles (64); synaptojanin1 (SYNJ1), which has a

prominent role in synaptic vesicle recycling (65); the beta subunit of the AP-3 protein complex, a

known PACS1 interactor that helps form clathrin-coated synaptic vesicles (65); and DNM1-

S774, which regulates clathrin-mediated endocytosis through its activation by

dephosphorylation (66).

Similarly, as observed in the other gain of regulation clusters, the early-up cluster had

enrichment for axon-related function and organization, but showed novel terms related to

signaling through semaphorin 3 (SEMA3E) (Fig. 3C, green pie plot). This result is in consensus

with single-cell RNA sequencing analysis, which identified *SEMA3E* to be upregulated in

PACS1$^{(+/R203W)}$ brain organoids (23). Our proteomic dataset shows that while there are no

changes in SEMA3E, several proteins involved in its signaling pathway are phosphorylated,

including CRMP1, DPYSL2, and DPYSL3. DPYSL2-S522 phosphorylation by CDK5 is involved

in axonal guidance and spine development (67).

Meanwhile, intracellular protein transport was the most enriched process in the clusters

that lost regulation (Fig. 3C, brown pie plot). Proteins that remained stable in PACS1$^{(+/R203W)}$

organoids include GLI3, which has a dual function as a transcriptional activator and a repressor

of the sonic hedgehog (Shh) pathway dependent on phosphorylation (68) and LIN28A, an RNA-

binding protein that is localized to the peri-endoplasmic reticulum and has been recently

described as a global suppressor of genes in the secretory pathway (69).

Dysregulated protein abundance and phosphorylation status in PACS1$^{(+/R203W)}$ organoids

result in convergent processes and overlap with known NDD-related genes

We next analyzed the overlap of protein identities with dysregulated abundance and

phosphorylation status in PACS1$^{(+/R203W)}$ organoids. Surprisingly, we observed a low overlap of

both the total and dysregulated proteins identified in both datasets (Fig. 4A, B). Nevertheless,

several affected processes converged (Fig. S6). For example, neurogenesis and axon-related

processes are enriched due to both dysregulated protein abundance and phosphorylation status

over time in PACS1$^{(+/R203W)}$ organoids when compared to controls (Fig. S6 purple and yellow

circles). Other converging processes include mRNA transport, actin and microtubule

organization, and cell cycle dynamics. However, exceptions were also present, where

processes will be representing mostly protein abundance or phosphorylation status. These

exceptions included oxidative phosphorylation and energy production resulting from the

dysregulation of protein abundance (purple); and chromatin remodeling and synapse

organization, which are composed of proteins showing a gain of upregulation in phosphorylation

status over time (yellow) (Fig. S6). These results suggest that PACS1$^{(+/R203W)}$ organoid

development is disrupted at different regulation levels.

Next, we investigated whether there are convergent dysregulated proteins in PACS1

syndrome compared to other NDDs. We performed gene-set enrichment analysis on the

dysregulated proteins using the DISGENET, the Simons Foundation ASD (SFARI), and the

Online Mendelian Inheritance in Man (OMIM) gene databases. A total of 311 genes (26.8%)

dysregulated in the PACS1$^{(+/R203W)}$ organoids compared to controls are additionally reported in

these databases. In DISGENET, we found 212 genes that showed a significant enrichment in

multiple nervous system disorders (Fig. 4C; Supplementary Table 23). These disorders

overlapped with phenotypes observed in PACS1 syndrome patients like ASD, ID, seizures, and

abnormal morphogenesis of the nervous system or facial features (4, 70), as well as

phenotypes yet to be reported, such as neurodegeneration. Finally, 94 of the genes that we

identified to be dysregulated in PACS1$^{(+/R203W)}$ organoids are also reported as ASD-risk genes in

the SFARI database, while 105 proteins had reported pathogenic NDD-related phenotypes in

the OMIM database (Fig. 4D,E Supplementary Table 24). These results support the overlap of

PACS1 syndrome and dysregulated neurodevelopmental processes with ASD and expand on

phenotypes that patients could manifest later in life.

Discussion

NDDs are highly heterogenous and can arise from variants across hundreds of genes

(62, 63). Even so, several limitations have hampered our understanding of the mechanisms by

which these variants act, limiting translation of genetic findings into the clinic (64, 65). Here, we

leveraged the potential of brain organoids to model neurodevelopment and gain insights into the

pathogenic mechanisms of PACS1 syndrome. By using mass spectrometry with TMT, we

identified 7,230 unique proteins, of which 3,677 were quantified with 100% present call in our

samples, a yield consistent with findings from other proteomic studies involving brain organoids

(27, 66, 67). We further observed temporal trajectories that reflected the transitions from

proliferating progenitor cells towards mature neuronal lineages, mimicking the temporal

dynamics observed in the fetal brain.

PACS1$^{(+/R203W)}$ organoids presented with dysregulated abundance and phosphorylation

status of temporally regulated proteins, including PACS1 known interactors. Despite the low

overlap between expression and phosphorylation alterations in PACS1$^{(+/R203W)}$, most dysregulated

processes converged. We observed upregulation of processes related to axonogenesis and

synapse maturation, suggesting accelerated development in PACS1$^{(+/R203W)}$ organoids. Although

PACS1$^{(+/R203W)}$ organoids presented with early synaptic formation signatures, the regulation of

multiple synapse and vesicle-related proteins is affected. At the same time, many synaptic

maturation processes are lost, possibly disrupting proper synaptic function. This is consistent

with the observed phenotype in mice overexpressing PACS1 p.R203W (22); and the

asynchrony and an increased network burst duration reported in neurons harboring the

p.R203W derived from patient iPSC lines (23). Interestingly, the synaptic terms obtained from

protein abundance and phosphorylation were also obtained from transcriptomic data,

suggesting that the PACS1 p.R203W variant might affect multiple regulatory levels.

Proteomic approaches are usually bulk and lack single-cell resolution; our experimental

set up also lacks cells that emerge later in development (astrocytes and oligodendrocytes) or

arise from non-ectoderm layers (microglia). With this being the case, other model systems

should be explored to determine how these cells contribute to PACS1 syndrome pathogenesis.

Nevertheless, we identified multiple proteins that displayed temporal changes in abundance and

phosphorylation status in developing PACS1$^{(+/R203W)}$ brain organoids with respect to controls. Of

further interest, most proteins that we found dysregulated in PACS1$^{(+/R203W)}$ organoids have also

been linked to NDDs (68-70). Additional mechanistic analyses are required to understand if

these effects are direct or indirect.

Most proteins that gain upregulation over time, and their resulting dysregulated processes,

could be linked to known PACS1 functions. For example, PACS1's role in protein sorting

explains the overrepresentation of vesicle-related processes in the stable-up cluster. Moreover,

several established PACS1 interactors were uncovered as dysregulated proteins, such as its

close paralog, PACS2. PACS1 and PACS2 evolutionarily emerged as a duplication in higher

metazoans from an ancient PACS gene (66). PACS1 function primarily involves sorting from

endosomes to the tGN, which showed temporal dysregulation of over 20 proteins in

PACS1[(+/R203W)] organoids. Meanwhile, PACS2 evolved cargo-sorting functions from the Golgi to

ER and vesicular coat preference for COPII; both processes upregulated in PACS1[(+/R203W)]

organoids. The upregulation in these processes is consistent with the detected gain of PACS2

regulation through increased phosphorylation at S437, which favors PACS2 trafficking function

(47). This increased phosphorylated PACS2 could explain the dysregulated vesicular transport

observed in PACS1[(+/R203W)] organoids. On the other hand, it could also explain the downregulation

of apoptotic signaling and mitochondrial function (Figure 2C). Of interest, a recurrent *de novo*

missense mutation in the regulatory middle region of PACS2 is associated with PACS2

Syndrome, another NDD (71). Further research aimed at understanding the interconnection of

PACS1 and PACS2 function, as well as their dysregulation and compensatory effects in

disease, will help shed light on the pathogenesis of both PACS1 and PACS2 syndromes.

The processes that lost downregulation over time were related to recently described

PACS1 functions, including proteins related to the maintenance of genome stability and DNA

repair via interaction with HDAC2, HDAC3, and HDAC6 (20, 22). Although we did not observe

protein abundance and phosphorylation status changes in HDAC2, 3, or 6, we found

downregulation in HDAC5 at day 40, and HDAC1 had decreased phosphorylation over

maturation. This suggests PACS1 could also modulate DNA stability through these proteins.

HDAC1 interacts with RB1 and SMARCA4 (both dysregulated at the phosphorylation level) in a

repressive complex in resting neurons that is abolished in a calcium-dependent manner to

enable transcription (72). Indicating that the PACS1 p.R203W variant could affect transcription

regulation by destabilizing the repressive complex formed by HDAC1, RB1, and SMARCA4.

Processes unrelated to known PACS1 functions also emerged, such as the enrichment

associated with regulating nucleotide metabolism and protein translation in the down-stable

cluster. There is growing evidence of the impact of protein translation on the pathogenesis of

neurodevelopmental disorders, including ASD and fragile X syndrome (73). However, this is the

first time that either process has been implicated in the PACS1 syndrome pathogenesis.

Moreover, PACS1$^{(+/R203W)}$ organoids presented with changes in genes reported to also be

dysregulated in other neurological diseases. The presence of the PACS1 p.R203W variant, at

the proteomic level, seems to affect processes related to the clinical features that patients

manifest, including ASD, ID, and seizures (3, 61), as well as phenotypes yet to be reported,

such as neurodegeneration. Additional models are required to test mechanisms of PACS1

syndrome outside the cerebral cortex, like the observed hypoplasia of the cerebellum and the

corpus callosum, or the dysmorphic facial features.

Processes that seem unrelated to PACS1 function, but which have displayed

upregulation in the PACS1$^{(+/R203W)}$ organoids, should be further explored. Considering current

evidence suggests the PACS1 p.R203W pathogenic variant may result in a gain of function (22,

23, 74), there is increased interest in generating targeted therapeutic strategies aimed at

suppressing its toxic effects (22, 25). Thus, processes that seem unrelated to PACS1 function

but with observed upregulation in the PACS1$^{(+/R203W)}$ organoids should be a point of interest.

Understanding whether the proteomic dysregulation results from the toxicity associated with the

variant or as a compensatory mechanism to maintain cellular homeostasis would be especially

relevant. Finally, the neurological phenotypes associated with dysregulated protein abundance

and phosphorylation in PACS1 p.R203W organoids are consistent with clinical characteristics

observed in PACS1 syndrome patients. Additional downstream analyses that aim to understand

the converging dysregulated processes could offer novel therapeutic strategies for PACS1

syndrome. Our results highlight that proteomic analyses not only enhance our understanding of

general NDD mechanisms by complementing transcriptomic studies, but may also uncover

additional targets, facilitating therapy development.

Acknowledgements

This research was supported in part through the computational resources and staff contributions

provided for the Quest high performance computing facility at Northwestern University which is

jointly supported by the Office of the Provost, the Office for Research, and Northwestern

University Information Technology.

Authors' contributions

This work was conceptualized by AGG. XGM performed proteomics analysis with help from SR,

BD, JG, and MLM, and generated the figures with the help of AW. LR cultured the organoids,

collected samples, and prepared them for proteomic analysis. AW, ALS, JR, CLB, and TAS

performed wet lab experiments (immunohistochemistry, western blotting, imaging) and data

analysis. Writing was done by XGM and edited by AGG, MLM, LR, JR, ALS, TAS, and CLB. All

authors read and approved the final manuscript.

Funding

This work was funded by the PACS1 Syndrome Research Foundation (AGG) and by National

Institutes of Health NINDS R01NS123163 (AGG) as well as R01MH125235 (M.L.M and B.D.),

R01MH118497 (MLM). Confocal microscopy was done at the Northwestern University Center

for Advanced Microscopy funded by NCI CCSG P30 CA060553.

Competing interests

The authors have nothing to disclose.

Availability of data and materials

Data is provided within the manuscript or supplementary information files and from the

corresponding author on reasonable request.

References

1.   Castellani CA, Arking DE. High-Risk, High-Reward Genetics in ASD. Neuron.

2020;105(3):407-10.

2.   Geschwind DH, Flint J. Genetics and genomics of psychiatric disease. Science.

2015;349(6255):1489-94.

3.   Schuurs-Hoeijmakers JH, Landsverk ML, Foulds N, Kukolich MK, Gavrilova RH,

Greville-Heygate S, et al. Clinical delineation of the PACS1-related syndrome--Report on 19

patients. Am J Med Genet A. 2016;170(3):670-5.

4.   Schuurs-Hoeijmakers JH, Oh EC, Vissers LE, Swinkels ME, Gilissen C, Willemsen MA,

et al. Recurrent de novo mutations in PACS1 cause defective cranial-neural-crest migration and

define a recognizable intellectual-disability syndrome. Am J Hum Genet. 2012;91(6):1122-7.

5.   Seto MT, Bertoli-Avella AM, Cheung KW, Chan KY, Yeung KS, Fung JL, et al. Prenatal

and postnatal diagnosis of Schuurs-Hoeijmakers syndrome: Case series and review of the

literature. Am J Med Genet A. 2021;185(2):384-9.

6.   Scott GK, Gu F, Crump CM, Thomas L, Wan L, Xiang Y, et al. The phosphorylation state

of an autoregulatory domain controls PACS-1-directed protein traffic. EMBO J.

2003;22(23):6234-44.

7.   Thomas G, Aslan JE, Thomas L, Shinde P, Shinde U, Simmen T. Caught in the act -

protein adaptation and the expanding roles of the PACS proteins in tissue homeostasis and

disease. J Cell Sci. 2017;130(11):1865-76.

8. Crump CM, Xiang Y, Thomas L, Gu F, Austin C, Tooze SA, et al. PACS-1 binding to

adaptors is required for acidic cluster motif-mediated protein traffic. EMBO J. 2001;20(9):2191-

201.

9. Gu F, Crump CM, Thomas G. Trans-Golgi network sorting. Cell Mol Life Sci.

2001;58(8):1067-84.

10. Hinners I, Wendler F, Fei H, Thomas L, Thomas G, Tooze SA. AP-1 recruitment to

VAMP4 is modulated by phosphorylation-dependent binding of PACS-1. EMBO Rep.

2003;4(12):1182-9.

11. Wan L, Molloy SS, Thomas L, Liu G, Xiang Y, Rybak SL, et al. PACS-1 defines a novel

gene family of cytosolic sorting proteins required for trans-Golgi network localization. Cell.

1998;94(2):205-16.

12. Jenkins PM, Zhang L, Thomas G, Martens JR. PACS-1 mediates phosphorylation-

dependent ciliary trafficking of the cyclic-nucleotide-gated channel in olfactory sensory neurons.

J Neurosci. 2009;29(34):10541-51.

13. Molloy SS, Thomas L, Kamibayashi C, Mumby MC, Thomas G. Regulation of endosome

sorting by a specific PP2A isoform. J Cell Biol. 1998;142(6):1399-411.

14. Schermer B, Hopker K, Omran H, Ghenoiu C, Fliegauf M, Fekete A, et al.

Phosphorylation by casein kinase 2 induces PACS-1 binding of nephrocystin and targeting to

cilia. EMBO J. 2005;24(24):4415-24.

15. Gorini G, Ponomareva O, Shores KS, Person MD, Harris RA, Mayfield RD. Dynamin-1

co-associates with native mouse brain BKCa channels: proteomics analysis of synaptic protein

complexes. FEBS Lett. 2010;584(5):845-51.

16. Kottgen M, Benzing T, Simmen T, Tauber R, Buchholz B, Feliciangeli S, et al. Trafficking

of TRPP2 by PACS proteins represents a novel mechanism of ion channel regulation. EMBO J.

2005;24(4):705-16.

17.   Taya S, Shinoda T, Tsuboi D, Asaki J, Nagai K, Hikita T, et al. DISC1 regulates the

transport of the NUDEL/LIS1/14-3-3epsilon complex through kinesin-1. J Neurosci.

2007;27(1):15-26.

18.   Atkins KM, Thomas LL, Barroso-Gonzalez J, Thomas L, Auclair S, Yin J, et al. The

multifunctional sorting protein PACS-2 regulates SIRT1-mediated deacetylation of p53 to

modulate p21-dependent cell-cycle arrest. Cell Rep. 2014;8(5):1545-57.

19.   Liu H, Hu PW, Budhiraja S, Misra A, Couturier J, Lloyd RE, et al. PACS1 is an HIV-1

cofactor that functions in Rev-mediated nuclear export of viral RNA. Virology. 2020;540:88-96.

20.   Mani C, Tripathi K, Luan S, Clark DW, Andrews JF, Vindigni A, et al. The multifunctional

protein PACS-1 is required for HDAC2- and HDAC3-dependent chromatin maturation and

genomic stability. Oncogene. 2020;39(12):2583-96.

21.   Veena MS, Raychaudhuri S, Basak SK, Venkatesan N, Kumar P, Biswas R, et al.

Dysregulation of hsa-miR-34a and hsa-miR-449a leads to overexpression of PACS-1 and loss

of DNA damage response (DDR) in cervical cancer. J Biol Chem. 2020;295(50):17169-86.

22.   Villar-Pazos S, Thomas L, Yang Y, Chen K, Lyles JB, Deitch BJ, et al. Neural deficits in a

mouse model of PACS1 syndrome are corrected with PACS1- or HDAC6-targeting therapy. Nat

Commun. 2023;14(1):6547.

23.   Rylaarsdam L, Rakotomamonjy J, Pope E, Guemez-Gamboa A. iPSC-derived models of

PACS1 syndrome reveal transcriptional and functional deficits in neuron activity. Nat Commun.

2024;15(1):827.

24.   Byrd DT, Han ZC, Piggott CA, Jin Y. PACS-1 variant protein is aberrantly localized in

Caenorhabditis elegans model of PACS1/PACS2 syndromes. Genetics. 2024;228(2).

25.   Rylaarsdam L, Reddy T, Guemez-Gamboa A. In search of a cure: PACS1 Research

Foundation as a model of rare disease therapy development. Trends Genet. 2022;38(2):109-12.

26.   Van Nuland A, Reddy T, Quassem F, Vassalli JD, Berg AT. PACS1-Neurodevelopmental

disorder: clinical features and trial readiness. Orphanet J Rare Dis. 2021;16(1):386.

27.   Melliou S, Sangster KT, Djuric U, Diamandis P. The promise of organoids for unraveling

the proteomic landscape of the developing human brain. Mol Psychiatry. 2022;27(1):73-80.

28.   Li C, Fleck JS, Martins-Costa C, Burkard TR, Themann J, Stuempflen M, et al. Single-

cell brain organoid screening identifies developmental defects in autism. Nature.

2023;621(7978):373-80.

29.   Paulsen B, Velasco S, Kedaigle AJ, Pigoni M, Quadrato G, Deo AJ, et al. Autism genes

converge on asynchronous development of shared neuron classes. Nature.

2022;602(7896):268-73.

30.   Buccitelli C, Selbach M. mRNAs, proteins and the emerging principles of gene

expression control. Nat Rev Genet. 2020;21(10):630-44.

31.   Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, et al. limma powers differential

expression analyses for RNA-sequencing and microarray studies. Nucleic Acids Res.

2015;43(7):e47.

32.   Ernst J, Bar-Joseph Z. STEM: a tool for the analysis of short time series gene

expression data. BMC Bioinformatics. 2006;7:191.

33.   Sloan SA, Andersen J, Pasca AM, Birey F, Pasca SP. Generation and assembly of

human brain region-specific three-dimensional cultures. Nat Protoc. 2018;13(9):2062-85.

34.   Dumrongprechachan V, Salisbury RB, Butler L, MacDonald ML, Kozorovitskiy Y.

Dynamic proteomic and phosphoproteomic atlas of corticostriatal axons in neurodevelopment.

Elife. 2022;11.

35.   Plubell DL, Wilmarth PA, Zhao Y, Fenton AM, Minnier J, Reddy AP, et al. Extended

Multiplexing of Tandem Mass Tags (TMT) Labeling Reveals Age and High Fat Diet Specific

Proteome Changes in Mouse Epididymal Adipose Tissue. Mol Cell Proteomics. 2017;16(5):873-

90.

36.   Djuric U, Lam KHB, Kao J, Batruch I, Jevtic S, Papaioannou MD, et al. Defining Protein

Pattern Differences Among Molecular Subtypes of Diffuse Gliomas Using Mass Spectrometry.

Mol Cell Proteomics. 2019;18(10):2029-43.

37.   Shen L, Sinai M. GeneOverlap: Test and visualize gene overlaps. R package version.

2013;1(0):444.

38.   Gehlenborg N. UpSetR: a more scalable alternative to Venn and Euler diagrams for

visualizing intersecting sets. R package version. 2019;1(0).

39.   Zhou Y, Zhou B, Pache L, Chang M, Khodabakhshi AH, Tanaseichuk O, et al. Metascape

provides a biologist-oriented resource for the analysis of systems-level datasets. Nat Commun.

2019;10(1):1523.

40.   Sayols S. rrvgo: a Bioconductor package for interpreting lists of Gene Ontology terms.

MicroPubl Biol. 2023;2023.

41.   Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, et al. Gene ontology:

tool for the unification of biology. The Gene Ontology Consortium. Nat Genet. 2000;25(1):25-9.

42.   Gene Ontology C, Aleksander SA, Balhoff J, Carbon S, Cherry JM, Drabkin HJ, et al.

The Gene Ontology knowledgebase in 2023. Genetics. 2023;224(1).

43.   Pinero J, Ramirez-Anguita JM, Sauch-Pitarch J, Ronzano F, Centeno E, Sanz F, et al.

The DisGeNET knowledge platform for disease genomics: 2019 update. Nucleic Acids Res.

2020;48(D1):D845-D55.

44.   Moon K, Martino T. R Package 'webr': Data and Functions for Web-Based Analysis. R

package version 01. 2020;5.

45.   Wickham H. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York;

2106.

46.   Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, et al. Cytoscape: a

software environment for integrated models of biomolecular interaction networks. Genome Res.

2003;13(11):2498-504.

47.   Banerjee-Basu S, Packer A. SFARI Gene: an evolving database for the autism research

community. Dis Model Mech. 2010;3(3-4):133-5.

48.   Amberger JS, Bocchini CA, Schiettecatte F, Scott AF, Hamosh A. OMIM.org: Online

Mendelian Inheritance in Man (OMIM(R)), an online catalog of human genes and genetic

disorders. Nucleic Acids Res. 2015;43(Database issue):D789-98.

49.   Trothen SM, Zang RX, Lurie A, Dikeakos JD. PACS-1 contains distinct motifs for

nuclear-cytoplasmic transport and interacts with the RNA-binding protein PTBP1 in the nucleus

and cytosol. FEBS Lett. 2022;596(2):232-48.

50.   Sorokina EA, Reis LM, Thompson S, Agre K, Babovic-Vuksanovic D, Ellingson MS, et al.

WDR37 syndrome: identification of a distinct new cluster of disease-associated variants and

functional analyses of mutant proteins. Hum Genet. 2021;140(12):1775-89.

51.   Scoumanne A, Cho SJ, Zhang J, Chen X. The cyclin-dependent kinase inhibitor p21 is

regulated by RNA-binding protein PCBP4 via mRNA stability. Nucleic Acids Res.

2011;39(1):213-24.

52.   Bernal A, Arranz L. Nestin-expressing progenitor cells: function, identity and therapeutic

implications. Cell Mol Life Sci. 2018;75(12):2177-95.

53.   Gyuris A, Donovan DJ, Seymour KA, Lovasco LA, Smilowitz NR, Halperin AL, et al. The

chromatin-targeting protein Brd2 is required for neural tube closure and embryogenesis.

Biochim Biophys Acta. 2009;1789(5):413-21.

54.   Abu-Khalil A, Fu L, Grove EA, Zecevic N, Geschwind DH. Wnt genes define distinct

boundaries in the developing human brain: implications for human forebrain patterning. J Comp

Neurol. 2004;474(2):276-88.

55.   Zou S, Mendes-Silva AP, Dos Santos FC, Ebrahimi M, Kennedy JL, Goncalves VF. Multi-

omics Analysis of Energy Metabolism Pathways Across Major Psychiatric Disorders. Mol

Neurobiol. 2025;62(10):13272-85.

56.   Mittal R, Patel K, Mittal J, Chan B, Yan D, Grati M, et al. Association of PRPS1 Mutations

with Disease Phenotypes. Dis Markers. 2015;2015:127013.

57.   Bjorklund G, Meguid NA, El-Bana MA, Tinkov AA, Saad K, Dadar M, et al. Oxidative

Stress in Autism Spectrum Disorder. Mol Neurobiol. 2020;57(5):2314-32.

58.   Aizawa H, Hu SC, Bobb K, Balakrishnan K, Ince G, Gurevich I, et al. Dendrite

development regulated by CREST, a calcium-regulated transcriptional activator. Science.

2004;303(5655):197-202.

59.   Chen KZ, Liu SX, Li YW, He T, Zhao J, Wang T, et al. Vimentin as a potential target for

diverse nervous system diseases. Neural Regen Res. 2023;18(5):969-75.

60.   Ruusuvuori E, Kaila K. Carbonic anhydrases and brain pH in the control of neuronal

excitability. Subcell Biochem. 2014;75:271-90.

61.   Aslan JE, You H, Williamson DM, Endig J, Youker RT, Thomas L, et al. Akt and 14-3-3

control a PACS-2 homeostatic switch that integrates membrane traffic with TRAIL-induced

apoptosis. Mol Cell. 2009;34(4):497-509.

62.   Pflum MK, Tong JK, Lane WS, Schreiber SL. Histone deacetylase 1 phosphorylation

promotes enzymatic activity and complex formation. J Biol Chem. 2001;276(50):47733-41.

63.   Tsukada M, Prokscha A, Ungewickell E, Eichele G. Doublecortin association with actin

filaments is regulated by neurabin II. J Biol Chem. 2005;280(12):11361-8.

64.   Cesca F, Baldelli P, Valtorta F, Benfenati F. The synapsins: key actors of synapse

function and plasticity. Prog Neurobiol. 2010;91(4):313-48.

65.   Nakatsu F, Ohno H. Adaptor protein complexes as the key regulators of protein sorting in

the post-Golgi network. Cell Struct Funct. 2003;28(5):419-29.

66.   Srinivasan S, Burckhardt CJ, Bhave M, Chen Z, Chen PH, Wang X, et al. A noncanonical

role for dynamin-1 in regulating early stages of clathrin-mediated endocytosis in non-neuronal

cells. PLoS Biol. 2018;16(4):e2005377.

67.   Gu Y, Hamajima N, Ihara Y. Neurofibrillary tangle-associated collapsin response

mediator protein-2 (CRMP-2) is highly phosphorylated on Thr-509, Ser-518, and Ser-522.

Biochemistry. 2000;39(15):4267-75.

68.   Andreu-Cervera A, Catala M, Schneider-Maunoury S. Cilia, ciliopathies and hedgehog-

related forebrain developmental disorders. Neurobiol Dis. 2021;150:105236.

69.   Cho J, Chang H, Kwon SC, Kim B, Kim Y, Choe J, et al. LIN28A is a suppressor of ER-

associated translation in embryonic stem cells. Cell. 2012;151(4):765-77.

70.   Tenorio-Castano J, Morte B, Nevado J, Martinez-Glez V, Santos-Simarro F, Garcia-

Minaur S, et al. Schuurs-Hoeijmakers Syndrome (PACS1 Neurodevelopmental Disorder): Seven

Novel Patients and a Review. Genes (Basel). 2021;12(5).

71.   Sanders SJ, Murtha MT, Gupta AR, Murdoch JD, Raubeson MJ, Willsey AJ, et al. De

novo mutations revealed by whole-exome sequencing are strongly associated with autism.

Nature. 2012;485(7397):237-41.

72.   Jourdon A, Scuderi S, Capauto D, Abyzov A, Vaccarino FM. PsychENCODE and

beyond: transcriptomics and epigenomics of brain development and organoids.

Neuropsychopharmacology. 2021;46(1):70-85.

73.   Zhao X, Bhattacharyya A. Human Models Are Needed for Studying Human

Neurodevelopmental Disorders. Am J Hum Genet. 2018;103(6):829-57.

74.   Nascimento JM, Saia-Cereda VM, Sartore RC, da Costa RM, Schitine CS, Freitas HR,

et al. Human Cerebral Organoids and Fetal Brain Tissue Share Proteomic Similarities. Front

Cell Dev Biol. 2019;7:303.

Figures

Figure 1. Cortical organoids recapitulate proteome dynamics during brain development.

A) Expression of neural precursor protein SOX2 (red) and projection neuron marker CTIP2 (magenta) in day 21, 40, and 80 organoids derived from control (up) and PACS1 p.R203W organoids (bottom). Scale bar☐=☐100☐µm. DAPI (blue) B) Total proteins detected in organoids were clustered by abundance changes over maturation time into three major trends: upregulated (red), downregulated (blue) or stable (gray) (graph on the left). Further classification detected four main abundance trends, based on the timing of change: steady over time (clusters steady -up and -down), or delayed to day 40 (clusters delayed -up and -down) (Right). Total number of proteins assigned to a cluster are indicated in each graph. C, D) Top GO terms upregulated (red) and downregulated (blue) by cluster over maturation time. The inner pie plots represent processes common to the clusters in each panel. The nested pie charts show root and branch significant GO enrichments with the width of the pie correlating to -log(p) of the GO term. Processes in bold represent those mentioned in the text. E) Top GO terms upregulated (red) and downregulated (blue) shared between organoids (purple) and fetal brain (green) over time. Synapse and neurodevelopment-related processes (box). Organoid profiles correspond to clusters 1 to 4, while the fetal brain profiles correspond to clusters A to H. A full description of all enriched processes is included in Supplementary Table 10.

Figure 2. Organoids carrying the PACS1 p.R203W pathogenic variant present with altered protein dynamics during maturation. A) Alluvial plot indicating the change of profile clustering in the PACS1 p.R203W organoids (right) in contrast to the clustering in controls (left). Breaks in the y axis were introduced to improve visualization of the dysregulated clusters by removing 2,000 proteins that remained stable over time. B) Four clusters of dysregulated proteins were identified in PACS1 p.R203W organoids. The plots are organized into a gain or loss of regulation categories: cluster stable-up (top), and stable-down (bottom) gained regulation while clusters up- and down-stable lost regulation (middle), C) Enriched biological functions per

dysregulated cluster Stable-up (top, red), Down-stable (middle, greyish blue) and Stable-down

(bottom, blue). The Up-stable cluster had no significant enrichment due to cluster size. The

nested pie charts show root and branch significant GO enrichments and the width of the pie

correlating to -log(p) of the GO term. Neurodevelopment-related processes in (bold).

Figure 3. Dysregulation of temporal phosphorylation dynamics in PACS1 p.R203W

organoids. A) Alluvial plot indicating the change of profile clustering in the PACS1 p.R203W

organoids in contrast to controls. Breaks in the y axis were introduced to improve visualization

of the dysregulated clusters by removing 3,000 modifications that remained stable over time. B)

Five clusters of dysregulated phosphorylation trends were observed in PACS1 p.R203W

organoids that either gained a temporal regulation or lost the original pattern observed in the

controls. The plots are organized into a gain or loss of regulation categories: clusters stable- up

and -down gained temporal regulation in contrast to the controls (orange), while clusters up- and

down- stable lost temporal regulation (gray). Early-up is a new cluster exclusive to the

phosphorylation dataset that also gained temporal regulation (green). C) Enriched biological

functions per dysregulated cluster. The nested pie charts show root and branch significant GO

enrichments and the width of the pie correlating to -log(p) of the GO term. Neurodevelopment-

related processes (bold).

Figure 4. Protein abundance and phosphorylation sites dysregulated by PACS1 p.R203W

are enriched in neurological disease-related genes. A) Overlapping identities between all

detected proteins and proteins with identified phosphorylation sites. B) Overlapping identities

between dysregulated proteins and phosphorylation sites observed in the PACS1 p.R203W

organoids when compared to controls. C) Gene-disease association type ontologies enriched in

the PACS1 p.R203W dysregulated proteome and phosphorylation sites. Selected top neuro-

related categories at a corrected p-value of 0.01. D) NDD and ASD risk genes dysregulated in

organoids carrying the p.R203W pathogenic variant. Genes are grouped by proteins and

phosphorylations. The protein abundance group is colorized by dysregulation cluster while the

phosphorylation clusters are colorized by major dysregulation trend (gain or loss) as presented

in figure 3: gain of regulation (clusters stable- up and -down) or loss of regulation (clusters up-

and down- stable). E) Volcano plot representing the dysregulated proteins (upper panel) and

phosphorylations (lower panel) at days 40 and 90. The squares in green tones represent the

ASD genes, while the circles in purple tones represent the NDD genes.

Supplemental Material

Supplementary information. Supplementary methods and supplementary figures 1-6

containing additional data.

Supplementary tables. Supplementary tables 1-24 contain the list of proteins and

phosphorylation sites, as well as accompanying statistical analysis performed on the data.