TIGIT expression dictates the immunosuppressive reprogramming of myeloid

cells in glioblastoma.

Mohammad Asad[1], Julio Inocencio[1], Stefan Mitrasinovic[1], Minori Aoki[1], Celina Crisman[1],

Patrick Lasala[1], Emad Eskandar[1], Chandan Guha[2], XingXing Zang[3], Ian F. Parney[4], and

Benjamin T. Himes[1*]

[1]Department of Neurological Surgery, Albert Einstein College of Medicine/Montefiore
Medical Center; Bronx, NY, 10461, USA (Mohammad Asad, J.I., S. M., Minori Aoki, C.C.,
P.L., E.E.)

[2]Department of Radiation Oncology, Albert Einstein College of Medicine/Montefiore
Medical Center; Bronx, NY, 10461, USA (C.G.)

[3]Department of Microbiology & Immunology, Albert Einstein College of
Medicine/Montefiore Medical Center; Bronx, NY, 10461, USA (X.Z.)

[4]Department of Neurological Surgery, Mayo Clinic; Rochester, MN, 55905, USA (I.F.P.)

Running title: TIGIT drives immunosuppression in glioblastoma

*Corresponding author. Email: Dr. Benjamin T. Himes

Albert Einstein College of Medicine, Department of Neurological Surgery; Kennedy-530,
1410 Pelham Parkway S. Bronx, NY, 10461, USA

Email ID: bhimes@montefiore.org

Phone number: +1 (718) 430-2538

Manuscript word count (includes abstract, text, references, and figure legends):

TIGIT expression dictates the immunosuppressive reprogramming of myeloid

cells in glioblastoma.

Mohammad Asad, Julio Inocencio, Stefan Mitrasinovic, Minori Aoki, Celina Crisman,

Patrick Lasala, Emad Eskandar, Chandan Guha, XingXing Zang, Ian F. Parney, and

Benjamin T. Himes

Abstract

Glioblastoma (GBM) is a deadly brain cancer with near-universal recurrence despite

maximal treatment for which new innovations are sorely needed. Immunotherapy has

yet to make significant gains in GBM treatment despite revolutionizing other cancer

therapies, due in part to GBM-mediated immune suppression. This immune

derangement proceeds through several mechanisms, but increasing evidence points to

critical roles for tumor-derived extracellular vesicles (EVs) and immunosuppressive

myeloid cells as key factors in this process. In the present study, we demonstrate broad

expression of TIGIT across myeloid cell populations in the GBM microenvironment, a

finding recapitulated by conditioning healthy monocytes with GBM-derived EVs. Further,

knockdown of TIGIT expression reduced the immunosuppressive polarization of

monocytes, resulting in improvement in T cell function. This finding proceeded in an

NLRP3-dependent manner, with substantial co-localization of TIGIT and NLRP3

expression prior to knockdown. These findings point to a novel role for TIGIT expression

in diverse myeloid cells in the GBM microenvironment as a marker of

immunosuppressive activity and further indicate a hierarchy of immunomodulatory

protein activity in these myeloid cells, with TIGIT knockdown unmasking the pro-

inflammatory activity of NLRP3. This study bolsters understanding of the

immunosuppressive complexities of myeloid cells in the GBM microenvironment, while

lending further support to prevention or attenuation of immunosuppressive myeloid cell

activity as a means of restoring immune function in GBM.

Graphical abstract (Created in BioRender. Asad, M. (2025)
https://BioRender.com/euiljoq

Keywords

glioblastoma, TIGIT, extracellular vesicles, immunotherapy, myeloid cell, NLRP3

Key Points

1. Tumor-mediated immune suppression is a key barrier to the development of effective immunotherapies for GBM.

2. TIGIT is broadly expressed in myeloid cells within the GBM microenvironment and can be induced by GBM-derived extracellular vesicles.

3. Knockdown of TIGIT reduces immunosuppressive polarization of monocytes and enhances T cell activity via NLRP3 signaling, implicating TIGIT expression as a targetable modifier of immunosuppressive activity in GBM-associated myeloid cells.

Importance of the Study

Glioblastoma (GBM) remains a formidable clinical challenge, with poor prognosis and limited response to current immunotherapies. This study uncovers a novel immunosuppressive axis involving TIGIT expression in myeloid cells, which are key players in the GBM tumor microenvironment. By demonstrating GBM-derived extracellular vesicles induce TIGIT in healthy monocytes and TIGIT knockdown diminishes immunosuppressive polarization in an NLRP3-dependent manner, this study highlights TIGIT as both a marker and modulator of immune dysfunction in GBM. These findings introduce a functional hierarchy of immunoregulatory proteins in tumor-associated myeloid cells, positioning TIGIT as a potential checkpoint target. Restoring immune activity by disrupting this axis could enhance the efficacy of immunotherapy in GBM. Thus, this research not only advances our understanding of tumor-induced

immune suppression but also opens a promising therapeutic avenue to reinvigorate

anti-tumor immunity in a cancer type historically resistant to immunotherapeutic

approaches.

Introduction

Glioblastoma (GBM) is the most common and deadly intrinsic brain tumor in adults.

Current standard of care consists of maximal safe surgical resection followed by

radiation and temozolomide chemotherapy, in a protocol essentially unchanged for two

decades (*1*). Some recent innovations, such as tumor treating fields, have shown

modest promise but have not led to increased median overall survival beyond 15

months in population-based studies (*2*). Immunotherapy, including immune checkpoint

blockade, has been transformative in cancer care in recent years, but has failed to

improve outcomes in GBM (*3-6*). The profound immunosuppression found in GBM

patients plays a critical role in these failures, but its underlying mechanisms remain

poorly understood (*6, 7*).

Increasing evidence points to a crucial role for myeloid cells, which can comprise the

bulk of the tumor immune infiltrate, as mediators of immune suppression, preventing the

antitumor effects of T cells (*7, 8*). These myeloid cells are heterogenous with and

include a number of distinct populations, including myeloid-derived suppressor cells

(MDSCs) and non-classical monocytes (NCM) (*8-12*). These diverse populations are

united by the common trait of facilitating immune suppression, whether by release of

immunosuppressive cytokines or expression of surface ligands that facilitate exhaustion

of effector cells, including T cells (*7, 13*).

The mechanisms by which GBM induces immunosuppression are multiple and an active

field of study, but an increasing body of work points to a central role of tumor-derived

extracellular vesicles (EVs) (*7, 10, 14, 15*). These complex, multimodal signaling

molecules, are shed by all cells and play a critical role in cancer signaling, particularly in

immune evasion (*16*). In GBM, work by our group and others has pointed to a

potentially pivotal role of EVs in tumor-mediated immune suppression, including the

induction of immunosuppressive myeloid cells (*10, 14, 17*). Thus, the means by which

GBM induces the immunosuppressive polarization of myeloid cells represents a

potentially potent therapeutic axis for mitigation of GBM-mediated immune suppression.

In the present study, we examine the immune phenotype of myeloid cells in the GBM

tumor microenvironment at the time of surgical resection using ultrasonic aspirate

processed directly from the operating room to identify immunomodulatory markers

expressed across diverse myeloid cell populations that may represent therapeutically

significant targets across diverse cell populations. Interestingly, we found high rates of

TIGIT (T cell immunoreceptor with immunoglobulin and ITIM domain). TIGIT is

increasingly recognized as playing a critical role in limiting adaptive and innate immunity

and is canonically understood as a marker for T cell exhaustion. Recent data

demonstrates TIGIT expression across an array of myeloid cells in the *in- situ* GBM

microenvironment (*18-20*). These findings were recapitulated using our *in vitro* system

of immunosuppressive monocyte induction, whereupon healthy monocytes are

subjected to GBM-EVs collected from patient-derived cell lines. Upon knockdown of

TIGIT expression in monocytes, we found cells underwent a more pro-inflammatory

immune polarization in response to EV conditioning. TIGIT knockdown unmasked high

levels of co-expressed NLRP3 and increased levels of IL-13 expression. This resulted in

a rescue of T cell proliferation upon subsequent co-culture, indicating a level of

functional restoration of immune function. These results illustrate the potentially crucial

role of an immunomodulatory hierarchy in myeloid cells in the tumor microenvironment,

the understanding of which may prove critical for reversal of tumor-mediated immune

suppression.

Results

Immunophenotyping of myeloid cells in the GBM microenvironment reveals

broad TIGIT expression

GBM is an intrinsic infiltrative tumor, with the radiographic tumor bulk comprised of

heterogenous tumor cells, immune infiltrate, and normal brain cells. Surgical resection

of GBM entails removing as much of the lesion as safely possible with a range of

surgical techniques, frequently including liberal use of an ultrasonic aspirator, which

breaks up the bulk tumor and allows for resection without excess trauma due to heat or

retraction on the surrounding brain. The ultrasonic aspirate from this approach thus

consists of all of the constituent cells of the bulk tumor and is thus a valuable source of

material for study of the *in situ* tumor microenvironment, as well as tumor cells for

primary cell culture and establishment of patient-derived cell lines. Ultrasonic aspirate

from 13 GBM patients was collected and the immune cell infiltrate profiled via high

dimensional flow cytometry (Cytek Aurora, full panel and gating strategy available in

Table S1 and Fig. 1). Analysis of a number of key immunomodulatory proteins were

assessed across several known immunosuppressive monocyte populations, including

monocytic myeloid-derived suppressor cells (mMDSCs,

Fig. 1

Figure 1: Expression of immune modulatory molecules by suppressive monocytes of GBM

patients. For surveying the tumor-immune microenvironment in samples of ultrasonic aspirates collected

during surgery, samples were processed for flow cytometric studies and the single cell suspensions,

stained with various flow antibodies were run on Cytek aurora spectral flow cytometer. (**A**) UMAP analysis

of CD45+ cells from GBM patients show various infiltrating immune cells. (**B**) Showing the cellular

expression of immune modulatory molecules (TIGIT, PDL1, NLRP3, BTLA, CTLA4, ARG1) by infiltrating

immune suppressive monocyte subsets, monocytic Myeloid Derived Suppressor Cells, mMDSCs

($CD45^+CD11b^+CD15^-HLA-DR^{Lo}CD14$ cells) and nonclassical monocytes (NCM, $CD45^+CD11b^+CD15^-HLA-$

$DR^+CD14^-PD1^+CD16$ cells). (**C**) Histograms and a corresponding summary bar graph depicting the

immune modulatory molecules expressing cell population in mMDSCs and NCM. (**D**) Box plot obtained

from TCGA database illustrates the expression of TIGIT in normal and primary tumor samples obtained

from GBM patients. (**E**) Kaplan-Meier curve of TCGA and GTEx datasets of human GBM, demonstrating

the effect of TIGIT expression on patient survival in GBM. Data are presented as mean $\pm$ s.e.m. and

were analyzed by one-way analysis of variance (ANOVA) with Tukey's multiple comparisons.

$CD11b^+CD15^-CD14^+HLA-DR^{lo/-}$) and non-classical monocytes (NCM, $CD11b^+CD15^-$

$CD16^+HLA-DR^+PD-1^+$), that have been previously described as major populations of

immunosuppressive myeloid cells in the GBM microenvironment (**Fig. 1A**) (*10, 14, 21*).

Among the infiltrating immune cells, T cells constituted the largest population, followed

by immunosuppressive monocytes and B cells. Subsequent analysis of

immunomodulatory proteins in mMDSC and NCM populations revealed enriched

expression of a few key immune modulatory proteins, such as VISTA, PD-L1 and B7-H3

(**Fig. 1B, C**). Interestingly, we found that TIGIT, typically thought of as a T cell

exhaustion marker, was significantly elevated in elevated in immunosuppressive

myeloid cell populations relative to other immunomodulatory markers. Despite the

significant presence of PD-L1 (29.75±5.83%) and B7-H3 (50.34±9.093%) expressing

cells in NCM, $TIGIT^+$ cells predominated among the myeloid cell populations within the

various suppressive monocytic subsets (64.27±7.99% and 39.8±4.366% in mMDSCs

and NCM respectively). Analysis of The Cancer Genome Atlas (TCGA) data

demonstrated enrichment of TIGIT expression across several GBM samples. Further,

TCGA and GTEx datasets of human GBM was analyzed using Gene Expression

Profiling Interactive Analysis. The TIGIT high group (top 20%, N= 128) indicated in red

showed a significant worse overall survival than the TIGIT low group (the rest 80%, N=

130) in blue, p = 0.036 (**Fig. 1 D, E**). Thus, it appears that elevated TIGIT mRNA

expression is associated with poor survival in human GBM patients.

GBM-EVs recapitulate myeloid cell immunosuppressive polarization *in vitro*

We next sought to validate these findings outside of the highly heterogenous context of

the tumor microenvironment. Work by our group and others has pointed to a major role

for GBM-derived extracellular vesicles (EVs) in the formation of immunosuppressive

monocytes (*10*). We previously demonstrated that EVs collected from patient-derived

GBM cell cultures can induce the formation of immunosuppressive monocytes, including

mMDSCs and NCMs, when used to condition monocytes collected from healthy human

blood donors. Using these techniques, we collected GBM-EVs from three patient-

derived GBM cell lines (dBT114, dBT116, and dBT120). In order to collect EVs, GBM

cells were cultured in serum-free conditions for 72hr and subjected to staged

centrifugation as described above. Presence and sizing of EVs was confirmed by use of

the NanoSight Nanotracker system (ZetaView, Particle Matrix, Germany) as well as the

presence of the EV markers CD9 and CD63 via western blot (**Fig. 2A-C**). EV

quantification was performed by BCA assay to quantify total protein. To induce the

formation of immunosuppressive monocytes, 20ug of EVs was added to monocytes

Fig. 2

Figure 2: Isolation and characterization of tumor-derived extracellular vesicles (TEVs) and

induction of immune suppressive monocytes. Extra cellular vesicles from human GBM primary cells

lines were obtained by ultracentrifugation and characterized by Nano-sight and western blots. (**A**) Size

distribution analysis of tumor-derived extracellular vesicles (TEVs) isolated from conditioned medium post

ultracentrifugation. (**B**) Representative image depicting the shape and relative size of the heterogeneous

TEVs, obtained from Nano-sight. (**C**) Western blot analysis of TEVs and whole cell lysate (WCL) for the

presence of CD63, and CD9 as TEVs markers. (**D**) CD11b+ cells, isolated from healthy PBMCs and

induced with control or different TEVs isolated from primary GBM cell lines (dBT114, dBT116 and dBT120)

under hypoxic condition in serum free medium. After 72 hours, cells were collected and processed for

flow cytometry and stained with various fluorochorm tagged antibodies. Flow cytometric analysis of

monocytic Myeloid Derived Suppressor Cells (mMDSCs, $CD45^+CD11b^+CD15^-HLA-DR^{Lo}CD14$ cells and

nonclassical monocytes (NCM, $CD45^+CD11b^+CD15^-HLA-DR^+CD14^-PD1^+CD16$ cells) isolated from

healthy PBMCs and induced with control or different TEVs isolated from primary GBM cell lines. Data are

presented as mean ± s.e.m. and were analyzed by one-way analysis of variance (ANOVA) with Tukey's

multiple comparisons.

isolated from anonymous healthy blood donors (obtained from the New York Blood

Bank). Monocytes were conditioned with EVs for 72h in hypoxic (1% O2) conditions

prior to collection for flow cytometry analysis (panel included in **S2**). Consistent with our

previously published results, EVs derived from dBT114, dBT116, and dBT120 all

significantly induced the formation of both mMDSCs and NCM in this model system (**Fig.**

2D), which produce anti-inflammatory cytokines including IL-4 (**Fig. S2**).

Using this model, we next employed a high-dimensional flow cytometry approach as

employed for the ultrasonic aspirate samples to define the expression pattern of

immunomodulatory proteins in this model. We found substantial similarities in

immunomodulatory protein expression following EV-conditioning of monocytes,

including consistent increases in TIGIT and NLRP3 expression (**Fig. 3A**, **B**). TIGIT

expression was increased in both mMDSCs (73.53±6.27%, 68.33±8.66% and

63.99±8.11% respectively for the mentioned cell lines EVs, compared to unstimulated

control, 4.68±2.83%) and NCM (85.17±3.91%, 79.26±9.17% and 60.68±10.52%

respectively, compared to control, 11.3±3.3%) following treatment with EVs derived from

Fig. 3

224 CD11b⁺ cells,

223 **Figure 3: Expression of immune modulatory molecules on TEVs induced peripheral monocytic**

224 **Myeloid Derived Suppressor Cells (mMDSC) & non-classical monocytes (NCM).** CD11b⁺ cells,

isolated from healthy peripheral blood and under hypoxic condition in a serum free medium, co-cultured

for 72 hours with TEVs obtained from mentioned GBM cell lines. (**A**) TEVs induced CD11b$^+$ monocytes

were studied for the expression of various immune modulatory markers, TIGIT, PDL1, NLRP3, BTLA,

CTLA4 & ARG1 on suppressive monocyte subsets (mMDSCs and NCMs). (**B**) Histograms depicting the

expression levels of immune modulatory molecules on mMDSC and NCM populations cultured with

control or different TEVs (dBT114, dBT116 and dBT120). The percentage of cells expressing each

molecule within each subset is also shown correspondingly as a chart. (**C**) Immunofluorescence staining

of TIGIT in suppressive monocytes treated with control or TEVs. (**D**) Western blot analysis of GBM whole

cell lysate (WCL) for the presence of TIGIT. Data are presented as mean □±□ s.e.m. and were analyzed

by one-way analysis of variance (ANOVA) with Tukey's multiple comparisons.

all three GBM cell lines. We confirmed that TIGT expression in this model was present

in monocytes and not from co-localized EVs via western blot demonstrating TIGIT

expression in GBM whole cell lysate (WCL), but not in EVs (**Fig. 3C**). Further,

immunofluorescent staining following EV-monocyte co-culture for 72h demonstrated

robust TIGIT expression in monocytes conditioned with EVs, but not in untreated

controls, demonstrating that hypoxic culture conditions, which have previously been

shown to aid in MDSC induction, were insufficient to induce TIGIT expression (**Fig. 3D**).

TIGIT and key immunomodulatory proteins are co-expressed in EV-induced

immunosuppressive monocyte populations

While there is some recent work describing TIGIT expression in myeloid cells (*20*),

typically TIGIT has been viewed as a T cell exhaustion marker, and so it is not

necessarily intuitive that high levels of TIGIT expression on immunomodulatory myeloid

cells corresponds to downstream immunosuppressive functionality in myeloid cells. We,

therefore next interrogated the immunomodulatory proteins and cytokines co-expressed

on TIGIT+ immunosuppressive monocytes. Several other immune-checkpoint markers,

including PD-L1 and VISTA, demonstrated high levels of co-expression on TIGIT

positive mMDSCs and NCM following GBM-EV treatment (**Fig. 4A, B,** representative

dot plots for co-localization included in **Fig. S3**). Interestingly, NLRP3, typically

understood as a strong marker of pro-inflammatory activity important in the

inflammasome pathway, demonstrated the strongest amount of co-expression. It is

noteworthy that among all CD45$^+$ cells, TIGIT$^+$ cells showed a significantly higher

expression of NLRP3$^+$ cells (86.64±5.52%) compared to other immunomodulatory

molecules, including Arg1 (0.45±0.14%), BTLA (2.2±0.94%), CTLA-4 (27.34±11.12%),

and PD-L1 (48.09±4.41%) (Fig. S3c). This elevated NLRP3 expression was

accompanied by a significant increase in IL-1β-producing cells (55.88±5%), an

inflammatory cytokine associated with NLRP3, as well as IL-13-producing cells

(98.78±.3%). These levels were notably higher than those of the proinflammatory

cytokines IFNγ (7.3±2.65%) and IL-17 (1.63±0.63%), and the immunosuppressive

cytokines IL-4 (5.9±1.3%), IL-5 (16.98±5%), and IL-10 (12.49±2.35%) (**Fig. S3**). In

interrogating a panel of key cytokines, we found that the TH2 cytokine, IL-13 was co-

expressed with TIGIT expression on immunosuppressive monocytes (**Fig. 4C, D)**.

Taken together, these findings suggest that an immunosuppressive phenotype prevails

in the presence of co-expressed pro-inflammatory markers in EV-induced

immunosuppressive monocytes, suggestive of an

Fig. 4

Figure 4: Immune response by TIGIT+ suppressive monocytes. Isolated peripheral blood CD11b+

monocytes were induced with TEVs in hypoxia chamber in serum free medium. After 72 hours of

induction, cells were processed for flow cytometric studies. Single viable CD45+TIGIT cells were selected

for the analysis. (**A**) UMAP analysis of mMDSCs and NCM treated with control or different TEV

preparations (dBT114, dBT116, dBT120), highlighting the co-expression of various immune modulatory

molecules (VISTA, PDL1, NLRP3, CTLA4, BTLA and ARG1) within TIGIT+ cell population. (**B**) Dotplot

depicting the percentage of cells as a factor of size of the dot expressing different immune modulatory

molecules within the TIGIT pockets of mMDSCs and NCMs treated with control or different EV

preparations. Mean fluorescence intensity (MFI) shown corresponding to the color intensity of the dots for

each analyte. (**C**) UMAP analysis of mMDSCs and NCM treated with control or different TEVs,

highlighting the expression of various cytokines (IL17, IL13, IL10, IL5, IL4, IL18, IFNγ). (**D**) Dotplot

depicting the percentage and MFI of TIGIT+ cells co-expressing different cytokines in mMDSCs and NCM

treated with control or different EV preparations.

immunomodulatory protein hierarchy that may be leveraged to repolarize towards an

anti-tumor immune response.

TIGIT knockdown in EV-conditioned monocytes results in a pro-inflammatory

phenotype

To test if the TIGIT driven immunomodulatory hierarchy regulates the NLRP3-mediated

protective immune response against GBM, we next sought to determine if prevention of

TIGIT expression on monocytes was sufficient to 'unmask' a pro-inflammatory

phenotype in EV-induced immunosuppressive monocytes. Prior to EV conditioning,

monocytes isolated from PBMCs were treated with TIGIT-directed siRNA to knock down

TIGT expression. GBM-EVs were subsequently added for 72h of hypoxic co-culture as

described above. TIGIT knockdown demonstrated immunomodulatory reprogramming in

Fig. 5

Figure 5. Immune modulatory molecules expressing Immune suppressive monocytes post TIGIT

knockdown. Isolated CD11b[+] Monocytic cells from healthy peripheral blood were knockdown by

scramble or NLRP3-specific or TIGIT-specific siRNA and processed for flow cytometry post TEVs

treatment for 72 hours under hypoxia in serum free condition. (**A and B**) Uniform Manifold Approximation

and Projection (UMAP) plots showing the expression patterns of immune checkpoint molecules (Arg1,

B7H3, BTLA, CTLA4, PD-L1, NLRP3, TIGIT, and VISTA) across different clusters by mMDSC population

post TIGIT specific siRNA- (b) or scramble siRNA- (a) treated cells. Red color represents high expression,

whereas gray indicates the total cellular population of mMDSCs. Violin plots depicting the percentage of

mMDSCs expressing various immune modulatory molecules in TIGIT siRNA-treated and scramble

siRNA-treated peripheral monocytes. (**C and D**) UMAP showing the expression of the same molecules by

non-classical monocytes (NCM), after TIGIT siRNA-treated (d) and scramble siRNA-treated (c) peripheral

monocytes. Violin plots depicting the percentage of NCMs expressing various immune modulatory

molecules in TIGIT siRNA-treated and scramble siRNA-treated cells. Data are presented as

mean ± s.e.m. and were analyzed by one-way analysis of variance (ANOVA) with Tukey's multiple

comparisons.

both mMDSC and NCM immunosuppressive monocyte populations (**Fig. 5**). While

TIGIT knockdown in mMDSCs resulted in reduced expression of other key

immunomodulatory markers including B7-H3 (11.6±3.39), PD-L1 (11.04±6.59), VISTA

(12.3±2.05), and CTLA-4 (2.19±1.18, **Fig. 5B**) relative to scramble treated controls

(32.73±6, 20.52±9.4, 29.46±5.6 and 18.12±8.1 respectively, **Fig. 5A**), this effect in

NCMs did not reach a significant difference in any of the above mentioned immune

regulatory proteins (B7-H3 15.72±8.09, PD-L1 16.49±4.37, VISTA 34.82±9.7 and CTLA-

4 12.38±3.12, respectively) compared to scramble controls (41.55±5.19,37.04±10.35,

39.20±6.7 and 33.66±8.35, **Fig. 5C, D, Fig. S4**). Meanwhile, NLRP3 expressing cells

remained significantly elevated following TIGIT knockdown in both populations

(72.58±5.78 and 48.75±8.19 respectively compared to respective scrambles, 31.2±7.65

and 25.61 $p < 0.05$, **Fig. 5B, D**).

Fig. 6

Figure 6. Immune modulatory milieu by immune suppressive monocytes post-TIGIT knockdown.

CD11b[+] cells isolated from healthy peripheral blood mononuclear cells (PBMCs) were knockdown by

TIGIT-specific or scramble siRNA under hypoxic condition in serum free medium. These healthy

monocytes were co-cultured with TEVs for 72 hours and processed for flow cytometry. (**A and B**) UMAP

plots showing the cellular distribution of mMDSCs based on their expression of various cytokines (IL17,

IL13, IL10, IL5, IL4, IL1β, and IFNγ) producing cells in scramble-treated (a) and TIGIT-knockdown

monocytes (b). Red color represents high expression, whereas gray indicates the total cellular population

of mMDSC. Violin plots comparing the percentage of mMDSCs expressing these cytokines between the

scramble-treated and TIGIT-knockdown groups. (**C and D**) UMAP showing the expression of the same

cytokines producing cells, (IL17, IL13, IL10, IL5, IL4, IL1β, and IFNγ) among non-classical monocytes

(NCM), in scramble-treated and TIGIT-knockdown groups. Violin plots comparing the percentage of

NCMs expressing various given cytokines between the scramble-treated and TIGIT-knockdown groups.

Data are presented as mean□±□s.e.m. and were analyzed by one-way analysis of variance (ANOVA)

with Tukey's multiple comparisons.

We next examined the effect of TIGIT knockdown on cytokine expression in these

immunosuppressive monocyte populations. This revealed a marked increase in IL17-

expressing cells relative to scramble controls in both mMDSC and NCM populations

(**Fig. 6B, D**). However, this enhanced IL17 expression was significantly higher in

mMDSCs. Interestingly, increased IL17 expression was coupled with an attenuation of

IL13 expression, in TIGIT downregulated monocyte subsets, which as mentioned above

(**Fig. 4**) previously co-localized heavily with TIGIT[+] immunosuppressive monocytes (**Fig.**

6 A, C, Fig. S5). Interestingly, IL13 expressing cells were observed to co-expressed

with NLRP3[+] cells within the mMDSCs and NCMs (**Fig. 5** and **6**). Furthermore, IL1β

expressing cells also exhibited co-expression with NLRP3 in the NCM (**Fig. 5D** and **6D**).

Collectively, these data indicate that TIGIT-dependent IL13 may play a role in regulating

NLRP3-mediated inflammation.

TIGIT knockdown in EV-conditioned monocytes rescues T cell proliferation in an

NLRP3-dependent manner

To assess the impact of TIGIT knockdown on the immunosuppressive function of

immunomodulatory monocytes, we co-cultured donor-matched T cells with EV-

conditioned monocytes with or without siRNA-mediated TIGIT knockdown as described

above. Our previous work has demonstrated that EV-induced immunosuppressive

monocytes readily suppress T cell proliferation (*7, 10, 14*). In the present study, we

found that TIGIT knockdown prior to EV conditioning rescued T cell proliferation to

levels comparable to monocytes treated with scramble controls (p<0.0265, **Fig. 7A**).

This also resulted in reduction of the number of IL4$^+$ CD4$^+$ T cells following co-culture

(p<0.025, **Fig. 7D**).

Based on our earlier results demonstrating high level of NLRP3 co-expressed with

TIGIT cells, we wished to interrogate if this apparent rescue of TH1-type immune

function was dependent on NLRP3 expression. By treating PBMC-derived monocytes

with siRNA targeting NLRP3 and subsequently both TIGIT and NLRP3, we found that

NLRP3 knockdown was sufficient to eliminate the rescue of T cell proliferation observed

following TIGIT knockdown (**Fig. 7A**). NLRP3 and TIGIT double knockdown also

eliminated the increase in CD4$^+$ T cells observed following TIGIT knockdown (**Fig. 7B**)

as well as the reduction in IL-4+ CD4+ T cells observed with TIGIT knockdown alone.

Although there was not much effect on IFNγ level (**Fig. 7C**), however, the ratio of

IFNγ/IL-4+ T cells increased significantly with TIGIT knockdown in monocytes alone

(**Fig. 7E**), an effect that was eliminated with NLRP3 knockdown.

Fig. 7

371
Figure 7. Effect of TIGIT knockdown on T Cell Proliferation in the Presence of TEV-Induced

MDSCs.

CD11b+ and CD3+ cells were isolated from healthy volunteer peripheral blood monocytes. CD11b cells

were conditioned with TIGIT and NLRP3-specific siRNA in serum-free medium followed by priming with

TEVs for under hypoxic conditions (1%) for 72 hours. Then, CD11b and T cells (T cells stained with cell

trace yellow dye) were co-cultured for another 72 hours in RPMI-1640 medium with 10% serum at 37°C in

5% CO_2. Finally, cells were processed for flow cytometry. (**A**) Representative flow cytometry plots and

corresponding bar graphs showing CD3[+] T cell proliferation (Cell Trace Yellow) in the absence of T cell

activator, CD3/CD28 Dynbeads (inactive group), when T cells were cultured without conditioned

monocytes Only T cell group), scramble treated group, TIGIT specific siRNA treated group (TIGIT),

NLRP3 specific siRNA treated group (NLRP3) and both TIGIT along with NLRP3 siRNA treated group

(T&N). (**B**) Representative flow cytometry plots showing IFNγ and IL-4 producing CD4+ T cells in CD3[+] T

cells and CD11b[+] monocytes co-culture conditions, post scramble, TIGIT and/or NLRP3 knockdown. (**C**)

IFNγ producing CD4+ T cells in the above-mentioned co-culture conditions in the given treatment groups.

(**D**) IL-4 producing CD4+ T cells in CD3[+] T cells and CD11b[+] monocytes co-culture conditions, post

scramble, TIGIT and/or NLRP3 knockdown. (**E**) Ratio of IFNγ producing CD4[+] T cells and IL-4 producing

CD4[+] T cells in the given co-culture conditions. Data are presented as mean ± s.e.m. and were

analyzed by one-way analysis of variance (ANOVA) with Tukey's multiple comparisons.

Discussion

Anti-cancer immunotherapy, including immune checkpoint blockade, has revolutionized

the treatment of several solid tumors in recent years (*22, 23*). Metastatic melanoma,

which once carried a prognosis comparable to GBM, has seen a revolution in outcomes

with immune checkpoint therapy (*3, 24*). Several clinical trials have explored these

treatments in GBM, but the results have been disappointing (*5, 25, 26*). Several

mechanisms for these failures have been implicated, including the mutational burden of

GBM, the immune specialization of the central nervous system, but the profound

immune derangement found in GBM patients is a long-described and under explored

phenomena (*7, 27-29*). The mechanisms of systemic immune suppression generated by

a localized disease such as GBM remains an area of active study, but induction of distal

effector signals, including EVs and immunosuppressive myeloid cells, are potent

avenues by which such effects could be achieved.

Work by our group and others has contributed to a growing body of evidence that

myeloid cells in the GBM microenvironment are major intermediaries of tumor-mediated

immune suppression, and that GBM-EVs are an effective means of inducing the

formation of immunosuppressive myeloid cells (*10, 17, 30*). These myeloid cells are a

highly heterogenous milieu, comprised of as many as 14 subtypes of cells as recently

described (*8*), and the relative import of different immunosuppressive myeloid cell

populations to the overall milieu is an open question. In our study, we focused on

mMDSCs and NCMs as exemplar immunosuppressive monocyte populations, as our *in*

situ tumor microenvironment data demonstrates a robust presence in the GBM

microenvironment and previous work has demonstrated that these populations are

modifiable by GBM-EVs (*10, 14, 30*). A central question to unlocking the complexity of

the tumor-immune microenvironment is understanding the relative import and/or

functional redundancy of these immunosuppressive cell populations, in order to define

the critical therapeutic targets in order to effectively repolarize the tumor

microenvironment to allow for anti-tumor immunity.

In the present study, we were surprised to find broad TIGIT expression across a large

swath of cells in the GBM microenvironment, including immunosuppressive myeloid

cells such as MDSCs and NCMs. Further, TCGA data indicated an association of TIGIT

expression with reduced patient survival, indicating a potential role for TIGIT expression

in disease severity. TIGIT has been classically described as a T cell exhaustion marker

(*31, 32*), but some reports do describe a role for TIGIT expression in myeloid cells (*33-*

35), including a study that indicated a correlation with prognosis in GBM patients (*20*).

Anti-TIGIT antibodies have had some success in clinical trials as an anti-cancer

immunotherapy (*36*), but have yet to show benefit in GBM treatment (*37*).

We found in our study that TIGIT expression correlated with several pro-inflammatory

markers in otherwise phenotypically immunosuppressive myeloid cells, with particularly

strong co-expression of TIGIT and NLRP3. NLRP3 is typically associated with the

inflammasome and the production of pro-inflammatory cytokines such as IL-1B (*38-40*).

The role of NLRP3 in cancer is more controversial, with many studies pointing to an

immunosuppressive role, while others suggest a beneficial role (*41, 42*). The larger

tumor-immune context may well provide an explanation for these mixed findings. Indeed,

we found that knockdown of TIGIT resulted in a pro-inflammatory polarization of

mMDSCs and NCMs, and that this effect was lost upon concomitant knockdown of

NLRP3. These results are suggestive of a hierarchical arrangement of

immunomodulatory signals in myeloid cells in the GBM tumor microenvironment.

Inhibition of a specific immune checkpoint may be insufficient to eliminate the

immunosuppressive effects of these cells given the presence of redundant signals, but

prevention certain of these signals may signify a globally immunosuppressive

phenotype. Prevention of the expression of certain hierarchically important signals, such

as TIGIT in the present study, may be sufficient to repolarize these cells to a pro-

inflammatory phenotype, which could have major implications for reversal of tumor-

mediated immune suppression.

Materials and methods

Study design

To identify potential therapeutic targets within the glioblastoma (GBM) tumor microenvironment, we analyzed the immune phenotype of myeloid cells obtained from surgical resection specimens using ultrasonic aspirates. This direct approach allowed for the identification of immunomodulatory markers across diverse myeloid cell populations. These findings were then validated using an *in vitro* model of immunosuppressive monocyte induction, where healthy monocytes were exposed to GBM-derived extracellular vesicles (EVs) from patient-derived cell lines. TIGIT knockdown studies were then performed to interrogate its import on the immunosuppressive polarization of myeloid cells. Subsequent donor-matched T cell proliferation studies were then performed as a functional measure of immunosuppressive activity.

Ultrasonic aspirate collection

Patients were identified with a likely diagnosis of GBM based on preoperative magnetic resonance imaging (MRI) of the brain. Written informed consent was obtained from all surgical candidates prior to surgery, in accordance with the protocol approved by the Institutional Review Board of Montefiore Medical Center (IRB# 2022-14518). Study patients (n= 14) underwent surgery with the goal of maximal safe resection. Ultrasonic aspirate was collected for research purposes based upon frozen section preliminary diagnosis of high grade glioma and included for data analysis upon confirmed diagnosis of GBM based on final histopathology and genetic testing.

Ultrasonic aspirates were immediately placed on ice for processing following collection

of sample in the course of surgical resection. Fresh tumor tissue was mechanically

dissociated and then enzymatically digested with TrypLE Express (Gibco, 12605010) at

37°C for 10 minutes. The resulting cell suspension was filtered through 40 μm nylon

mesh, treated with red blood cell lysis buffer (BioLegend, 420301), and washed with

phosphate-buffered saline (PBS). Single-cell suspensions were then prepared for flow

cytometry analysis.

Cell Culture

Patient-derived GBM cell lines BT114, BT116, and BT120, were obtained from Mayo

Clinic and maintained as neurospheres in serum-free, HEPES-modified DMEM/F12

medium supplemented with EGF, FGF, and N2, as previously described.(*43*)

Subsequently, these cell lines were differentiated to monolayer culture for the purposes

of EV generation (dBT114, dBT116, and dBT120) in HEPES-modified EmbryoMax

DMEM/F12 medium (DF-042-B, Millipore, Burlington, MA) containing 1%

penicillin/streptomycin and 10% fetal calf serum. Cells were maintained in T75 flasks at

37°C with 5% CO2 in a humidified incubator, as previously described.(*10*) The cells

were grown as an adherent monolayer and passaged when confluency was around

90%. Early passages within 10 or lower were used for the study.

EV Isolation and protein estimation

For isolation of EVs, dBT114, dBT116, and dBT120 cells were transferred to serum-free

neurosphere culture, DMEM/F12 medium containing EGF and FGF growth factors in 5%

CO_2 incubator. After 72hrs, cultured supernatants were collected in 50 ml centrifuge

tubes and centrifuged at 1900 *g* for 10 min at room temperature. Then supernatants

were transferred to 14 ml propylene Ultracentrifuge tubes, 14 X 95 mm (Cat# 331374,

Beckman Coulter, Brea, CA) and centrifuged at 24000 rpm for 16 hrs at 4°C on Optima

XE-90 ultracentrifuge (Beckman Coulter, Brea, CA). After centrifugation, supernatant

was aspirated until ~200µl of EV-enriched concentrated remained, consistent with prior

protocols.(*10*) EVs were collected transferred to a cryovial, and protein concentration

was calculated by modified Pierce Rapid Gold BCA Protein Assay Kit (A53226, Thermo

Fisher, Waltham, MA) according to manufacturer's method and the read using an

Infinite 200Pro plate reader (Tecan, Mannedorf, Switzerland) using software Tecan I-

Control (version 2.0). The isolated EVs were stored at -80°C until used.

EV characterization

EVs were characterized using the ZetaView Nanotracker system (ZetaView, Particle

Metrix GmbH). Freshly isolated EVs were diluted 1000-fold with filtered (40 nm syringe

filter) phosphate-buffered saline (PBS) to achieve a particle concentration suitable for

analysis and to remove larger particles or aggregates, and 1µL was loaded into the

capillary system. The included software calculated EV size using the Stokes-Einstein

equation and histograms demonstrating the distribution of EV size generated, along with

movies demonstrating EV morphology.

Peripheral blood mononuclear cells (PBMC) isolation

Human blood samples were obtained from the New York Blood Center from healthy

donors (n=10). Either heparin or EDTA was used as an anticoagulant. Blood samples

were delivered at room temperature (~24°C) and processed immediately after the

delivery. All subsequent steps were performed at room temperature. PBMCs were

isolated from whole blood using standard density gradient centrifugation with Ficoll-

Paque PLUS (17-5442-03, density 1.077 ± 0.001 g/ml, GE Healthcare, Chicago, IL) as

described.(*30*) Briefly, blood was diluted 1:1 with 1X PBS and mixed gently. Ficoll-

Paque™ Premium was added to the bottom of a sterile 50 ml centrifuge tube. The

diluted blood was carefully layered on top of the Ficoll-Paque, maintaining a clear

interface between the two layers. The tube was centrifuged at 500g for 30 minutes

without brake. PBMCs were collected from the interface between the plasma and Ficoll

layers using a sterile transfer pipette. The collected PBMCs were washed with PBS to

remove any residual Ficoll or other contaminants and resuspended in PBS + 2% FBS.

Cells were counted using the trypan blue exclusion method on Countess 3 (Invitrogen,

Waltham, MA).

Isolation of myeloid cells

Myeloid cells were positively selected from healthy donor PBMCs, using CD11b

MicroBeads (130-049-601, Miltenyi Biotec, Bergisch Gladbach, Germany). Briefly, 5 µL

of CD11b MicroBeads were added per 10^7 total cells, and the mixture was incubated for

30 minutes at room temperature. Cells were then washed twice with PBS + 2% FBS

and resuspended in PBS + 2% FBS. The cell suspension was applied to an LS column

placed in a magnetic field of a MACS Separator (Miltenyi Biotec, Bergisch Gladbach,

Germany). Unlabeled cells were collected as the flow-through fraction. The column was

then removed from the magnetic field, and labeled monocytes were eluted by flushing

the column with PBS + 2% FBS. The purity of the isolated myeloid cells were assessed

by flow cytometry using anti-CD45 and anti-CD11b antibodies.

534 **Myeloid cell culture and EV treatment**

535 Isolated CD11b$^+$ cells were cultured in L-glutamine and HEPES-modified DMEM

536 medium supplemented with 1% penicillin/streptomycin under hypoxic conditions (1%

537 O2). Cells were seeded at a density of 1 million cells per well in a 96-well plate. Cells

538 were treated with EVs (20 μg/well) isolated from glioblastoma cell lines (dBT114,

539 dBT116 and dBT120) for 72hrs.

540 **Flow cytometry**

541 Cultured and EV-conditioned myeloid cells were washed twice with PBS and

542 resuspended in 50μl of PBS and processed for flow cytometric study as previously

543 described.(*44*) Briefly, cells were stained with LIVE/DEAD Fixable Blue Dead Cell Stain

544 (Thermo Fisher Scientific) for 30 minutes at room temperature in the dark. Following

545 staining, cells were washed twice with FACS buffer (containing 2% human serum) and

546 resuspended in 50μl of FACS buffer. To block non-specific binding, cells were incubated

547 with Fc blocking solution for 20 minutes at room temperature. Subsequently, cells were

548 washed twice with FACS buffer and resuspended in 50μl of staining buffer containing a

549 cocktail of fluorochrome-conjugated antibodies specific for surface markers of interest.

550 Cells were incubated for 60 minutes at 4°C in the dark. Following surface staining, cells

551 were washed twice with FACS buffer and resuspended in 50μl of FACS buffer

552 containing 2% human serum. Cells were fixed and permeabilized using

553 Cytofix/Cytoperm solution (BD Biosciences, San Jose, CA) for 20 minutes at 4°C. Cells

554 were then washed twice with permeabilization buffer and incubated with 50μl cocktail of

555 fluorochrome-conjugated antibodies specific for intracellular markers of interest for

overnight (12-14 hrs) at 4°C in the dark. After incubation, cells were washed twice with

permeabilization buffer, resuspended in 200µl of FACS buffer, and filtered through a 70-

micron nylon mesh to remove cell clumps. Finally, samples were acquired on a Cytek

Aurora 5 laser flow cytometer and analyzed using FlowJo software.

T cell isolation and proliferation assay

PBMCs were isolated from healthy human donors as described above. CD3+ T cells

were then purified from PBMCs via magnetic cell sorting using a commercially available

kit (130-050-101, Miltenyi Biotec, Bergisch Gladbach, Germany), following a similar

procedure to CD11b isolation described above. Isolated CD3+ T cells were cultured at a

density of 1 x 10^7 cells/ml in RPMI-1640 medium supplemented with 1% penicillin-

streptomycin and 10% fetal bovine serum for 3 days (or five days in the case of siRNA

knockdown experiments) at 37°C in 5% CO_2. CD3+ T cells were labeled with CellTrace

Yellow (Thermo Fisher) according to the manufacturer's instructions. Autologous donor-

matched labeled CD3+ T cells were co-cultured with EV-conditioned CD11b+ myeloid

cells at a 1:1 ratio (1 x 10^6 cells each) in RPMI-1640 medium supplemented with 1%

penicillin-streptomycin and 10% fetal bovine serum in the presence of aCD3/aCD28

DynaBeads (11131D Thermo Fisher, Waltham, MA) for 5 days at 37°C in 5% CO_2. After

5 days of co-culture, cells were harvested and processed for flow cytometry analysis.

siRNA knockdown in CD11b[+] myeloid cells

Purified CD11b+ myeloid cells (10^6 cells/well) were plated in a 96-well plate in 100µL of

serum-free medium (Dharmacon, Lafayette, CO) containing 1% penicillin-streptomycin

and transfected with Accell small interfering RNA (siRNA; Dharmacon, Lafayette, CO) at

a final concentration of 1µM. Cells were incubated under normoxic conditions (37°C, 5%

CO_2) for 48 hours. Following incubation, CD11b+ monocytes were stimulated with 20µg

of tumor-derived extracellular vesicles (TEVs) and cultured for an additional 72 hours

under hypoxic conditions (1% O_2, 5% CO_2, 37°C).

Immunofluorescence staining

CD11b$^+$ myeloid cells isolated from PBMCs as described above were cultured in 8-well

chamber slides at a density of 1×10^6 cells per well and allowed to adhere under

standard conditions (37°C, 5% CO_2). After adherence, the culture medium was

aspirated, and cells were gently rinsed with PBS. Fixation was performed using 4%

paraformaldehyde (PFA) in PBS for 15 minutes at room temperature (RT), followed by

three five minute PBS washes. Cells were permeabilized using 0.1% Triton X-100 in

PBS for 10 minutes at RT and washed three times with PBS. To block non-specific

binding, cells were incubated in blocking buffer (5% bovine serum albumin (BSA) in

PBS) for 1 hour at RT in a humidified chamber. Cells were incubated with primary

antibody diluted in blocking buffer at the recommended concentration overnight at 4°C

in a humidified chamber. After incubation, cells were washed three times with PBS (5

minutes each). A fluorophore-conjugated secondary antibody was diluted in blocking

buffer and applied to the cells for 1 hour at RT in the dark. Unbound antibody was

removed by washing three times with PBS (5 minutes each). Cells were incubated with

1µg/mL DAPI in PBS for 5 minutes at RT and subsequently washed three times with

PBS. The chamber walls were carefully removed, and a drop of mounting medium was

applied to each well. Coverslips were placed gently to avoid bubble formation. Slides

were allowed to set for at least 30 minutes at RT in the dark before imaging.

Immunofluorescence was visualized using a fluorescence microscope equipped with

appropriate filter sets.

Western blotting

Myeloid cells, cultured with or without EVs stimulation were harvested and washed with

ice cold PBS twice. Cells were then lysed in RIPA buffer supplemented with protease

and phosphatase inhibitors and incubated on ice for 30 min with intermittent vortexing.

Lysates were centrifuged at 12,000g for 7 min at 4°C, and the supernatant was

collected. Protein concentration was determined using a Pierce Rapid Gold BCA Protein

Assay Kit (A53226, Thermo Fisher, Waltham, MA) according to manufacturer's

instructions and read using an Infinite 200 Pro plate reader (Tecan, Mannedorf,

Switzerland) using software Tecan I-Control (version 2.0). Equal amounts of protein

(20–40 µg) were resolved on a 10–12% SDS-PAGE gel at 100V until the dye front

reached the bottom. Proteins were transferred to a PVDF or nitrocellulose membrane

using iBLOT3 (Invitrogen, Waltham, MA). Membranes were blocked with 5% BSA or 5%

nonfat milk in Tris-buffered saline with Tween 20 (TBST) for 1 hour at room temperature.

The membranes were incubated overnight at 4°C with anti-TIGIT primary antibody

(PA5-121546, Invitrogen, Waltham, MA) followed by three washes with TBST (5 min

each). Antibodies against HSP90 (Thermo Fisher, Waltham, MA) were used as loading

control. HRP-conjugated secondary antibody was applied for 1 hour at RT, followed by

three additional washes with TBST. Proteins were detected using an enhanced

chemiluminescence (ECL) reagent, and signals were visualized using a

chemiluminescence imaging system.

Statistics

For statistical analysis, GraphPad Prism V.10.0 software (La Jolla, CA) was used to

graph the data and perform necessary calculations. Statistical comparisons were

primarily conducted using one-way analysis of variance (ANOVA) with Tukey's multiple

comparisons. Data are presented as mean $\pm$ S.E.M. and a p value threshold of <0.05

was set for determining statistical significance across all experiments. In the summary

dot plots or violin plots, each dot represents an individual human value. The p values

and S.E.M are indicated on these plots through error bars.

Supplementary Materials

The PDF file includes:

Figs. S1 to S5

Tables S1 to S2

Funding: This work was supported by the Neurosurgery Research & Education

Foundation through the B*CURED & NREF Young Clinician Investigator Award, as well

as intramural K12 funding through the Institute for Clinical and Translational Research

and the NCI Paul Calabresi Career Development Award (1K12CA279871).

Author contributions: M.A. and B.T.H. conceptualized, designed the experiments and

wrote the manuscript. M.A., J.I. and Minori Aoki performed the experiments. C.C., P.L.

and E.E. provided the patient samples. M.A. and J.I. conducted data analysis. S.M. help

in preparing dot plot in figure 4. E.E., C.G. and X.Z. provided important advice. I.F.P.

provided the GBM cell lines. B.T.H. acquired funding and directed the research.

Competing interests: The authors declare that they have no competing interest.

Data and materials availability: All data associated with this study are present in the

paper or the Supplementary Materials.

References

1. R. Stupp *et al.*, Radiotherapy plus concomitant and adjuvant temozolomide for
glioblastoma. *N Engl J Med* **352**, 987-996 (2005).
2. L. Marenco-Hillembrand *et al.*, Trends in glioblastoma: outcomes over time and
type of intervention: a systematic evidence based analysis. *J Neurooncol* **147**,
297-307 (2020).
3. J. S. Weber *et al.*, Nivolumab versus chemotherapy in patients with advanced
melanoma who progressed after anti-CTLA-4 treatment (CheckMate 037): a
randomised, controlled, open-label, phase 3 trial. *Lancet Oncol* **16**, 375-384
(2015).
4. R. S. Herbst *et al.*, Pembrolizumab versus docetaxel for previously treated, PD-
L1-positive, advanced non-small-cell lung cancer (KEYNOTE-010): a randomised
controlled trial. *Lancet* **387**, 1540-1550 (2016).
5. J. F. Inocencio *et al.*, Immune checkpoint pathways in glioblastoma: a diverse
and evolving landscape. *Front Immunol* **15**, 1424396 (2024).
6. C. M. Jackson, J. Choi, M. Lim, Mechanisms of immunotherapy resistance:
lessons from glioblastoma. *Nat Immunol* **20**, 1100-1109 (2019).
7. B. T. Himes *et al.*, Immunosuppression in Glioblastoma: Current Understanding
and Therapeutic Implications. *Front Oncol* **11**, 770561 (2021).
8. C. Jackson *et al.*, Distinct myeloid-derived suppressor cell populations in human
glioblastoma. *Science* **387**, eabm5214 (2025).
9. S. Ostrand-Rosenberg, Myeloid derived-suppressor cells: their role in cancer and
obesity. *Curr Opin Immunol* **51**, 68-75 (2018).
10. B. T. Himes *et al.*, The role of extracellular vesicles and PD-L1 in glioblastoma-
mediated immunosuppressive monocyte induction. *Neuro Oncol* **22**, 967-978
(2020).
11. J. P. Antonios *et al.*, Immunosuppressive tumor-infiltrating myeloid cells mediate
adaptive immune resistance via a PD-1/PD-L1 mechanism in glioblastoma.
Neuro Oncol **19**, 796-807 (2017).
12. Y. Mi *et al.*, The Emerging Role of Myeloid-Derived Suppressor Cells in the
Glioma Immune Suppressive Microenvironment. *Front Immunol* **11**, 737 (2020).
13. M. P. Gustafson *et al.*, Systemic immune suppression in glioblastoma: the
interplay between CD14+HLA-DRlo/neg monocytes, tumor factors, and
dexamethasone. *Neuro Oncol* **12**, 631-644 (2010).
14. M. Y. Jung *et al.*, Superinduction of immunosuppressive glioblastoma
extracellular vesicles by IFN-gamma through PD-L1 and IDO1. *Neurooncol Adv*
4, vdac017 (2022).
15. F. L. Ricklefs *et al.*, Immune evasion mediated by PD-L1 on glioblastoma-derived
extracellular vesicles. *Sci Adv* **4**, eaar2766 (2018).
16. E. I. Buzas, The roles of extracellular vesicles in the immune system. *Nat Rev*
Immunol **23**, 236-250 (2023).

17.   N. Ludwig *et al.*, Characterization of systemic immunosuppression by IDH mutant
689   glioma small extracellular vesicles. *Neuro Oncol* **24**, 197-209 (2022).
690   18.   Y. Kong *et al.*, T-Cell Immunoglobulin and ITIM Domain (TIGIT) Associates with
691   CD8+ T-Cell Exhaustion and Poor Clinical Outcome in AML Patients. *Clin Cancer*
692   *Res* **22**, 3057-3066 (2016).
693   19.   L. Wu *et al.*, Blockade of TIGIT/CD155 Signaling Reverses T-cell Exhaustion and
694   Enhances Antitumor Capability in Head and Neck Squamous Cell Carcinoma.
695   *Cancer Immunol Res* **7**, 1700-1713 (2019).
696   20.   I. Raphael *et al.*, TIGIT and PD-1 Immune Checkpoint Pathways Are Associated
697   With Patient Outcome and Anti-Tumor Immunity in Glioblastoma. *Front Immunol*
698   **12**, 637146 (2021).
699   21.   J. C. Rodrigues *et al.*, Normal human monocytes exposed to glioma cells acquire
700   myeloid-derived suppressor cell-like properties. *Neuro Oncol* **12**, 351-365 (2010).
701   22.   M. A. Postow, M. K. Callahan, J. D. Wolchok, Immune Checkpoint Blockade in
702   Cancer Therapy. *J Clin Oncol* **33**, 1974-1982 (2015).
703   23.   C. H. June, R. S. O'Connor, O. U. Kawalekar, S. Ghassemi, M. C. Milone, CAR T
704   cell immunotherapy for human cancer. *Science* **359**, 1361-1365 (2018).
705   24.   A. Hauschild *et al.*, Dabrafenib in BRAF-mutated metastatic melanoma: a
706   multicentre, open-label, phase 3 randomised controlled trial. *Lancet* **380**, 358-365
707   (2012).
708   25.   P. Y. Wen *et al.*, Glioblastoma in adults: a Society for Neuro-Oncology (SNO) and
709   European Society of Neuro-Oncology (EANO) consensus review on current
710   management and future directions. *Neuro Oncol* **22**, 1073-1113 (2020).
711   26.   H. Y. Chan, J. Choi, C. Jackson, M. Lim, Combination immunotherapy strategies
712   for glioblastoma. *J Neurooncol* **151**, 375-391 (2021).
713   27.   P. E. Fecci *et al.*, Increased regulatory T-cell fraction amidst a diminished CD4
714   compartment explains cellular immune defects in patients with malignant glioma.
715   *Cancer Res* **66**, 3294-3302 (2006).
716   28.   C. A. Crane, B. J. Ahn, S. J. Han, A. T. Parsa, Soluble factors secreted by
717   glioblastoma cell lines facilitate recruitment, survival, and expansion of regulatory
718   T cells: implications for immunotherapy. *Neuro Oncol* **14**, 584-595 (2012).
719   29.   K. Ayasoufi *et al.*, Brain cancer induces systemic immunosuppression through
720   release of non-steroid soluble mediators. *Brain* **143**, 3629-3652 (2020).
721   30.   B. T. Himes *et al.*, Use of heparin to rescue immunosuppressive monocyte
722   reprogramming by glioblastoma-derived extracellular vesicles. *J Neurosurg*, 1-11
723   (2022).
724   31.   H. Harjunpaa, C. Guillerey, TIGIT as an emerging immune checkpoint. *Clin Exp*
725   *Immunol* **200**, 108-119 (2020).
726   32.   N. C. Blessin *et al.*, Patterns of TIGIT Expression in Lymphatic Tissue,
727   Inflammation, and Cancer. *Dis Markers* **2019**, 5160565 (2019).
728   33.   J. Yue *et al.*, Myeloid-derived suppressor cells inhibit natural killer cells in
729   myelodysplastic syndromes through the TIGIT/CD155 pathway. *Hematology* **28**,
730   2166333 (2023).
731   34.   X. Guan *et al.*, Anti-TIGIT antibody improves PD-L1 blockade through myeloid
732   and T(reg) cells. *Nature* **627**, 646-655 (2024).

733   35.   J. H. Han *et al.*, Effective Anti-tumor Response by TIGIT Blockade Associated
734   With FcgammaR Engagement and Myeloid Cell Activation. *Front Immunol* **11**,
735   573405 (2020).

736   36.   B. C. Cho *et al.*, Tiragolumab plus atezolizumab versus placebo plus
737   atezolizumab as a first-line treatment for PD-L1-selected non-small-cell lung
738   cancer (CITYSCAPE): primary and follow-up analyses of a randomised, double-
739   blind, phase 2 study. *Lancet Oncol* **23**, 781-792 (2022).

740   37.   E. Schonfeld, J. Choi, A. Tran, L. H. Kim, M. Lim, The landscape of immune
741   checkpoint inhibitor clinical trials in glioblastoma: A systematic review.
742   *Neurooncol Adv* **6**, vdae174 (2024).

743   38.   A. Grebe, F. Hoss, E. Latz, NLRP3 Inflammasome and the IL-1 Pathway in
744   Atherosclerosis. *Circ Res* **122**, 1722-1740 (2018).

745   39.   B. Z. Shao, Z. Q. Xu, B. Z. Han, D. F. Su, C. Liu, NLRP3 inflammasome and its
746   inhibitors: a review. *Front Pharmacol* **6**, 262 (2015).

747   40.   K. V. Swanson, M. Deng, J. P. Ting, The NLRP3 inflammasome: molecular
748   activation and regulation to therapeutics. *Nat Rev Immunol* **19**, 477-489 (2019).

749   41.   S. Hamarsheh, R. Zeiser, NLRP3 Inflammasome Activation in Cancer: A Double-
750   Edged Sword. *Front Immunol* **11**, 1444 (2020).

751   42.   B. R. Sharma, T. D. Kanneganti, NLRP3 inflammasome in cancer and metabolic
752   diseases. *Nat Immunol* **22**, 550-559 (2021).

753   43.   R. Kumar *et al.*, Modulating glioma-mediated myeloid-derived suppressor cell
754   development with sulforaphane. *PLoS One* **12**, e0179012 (2017).

755   44.   M. Asad *et al.*, EB1-3 Chain of IL-35 Along With TGF-beta Synergistically
756   Regulate Anti-leishmanial Immunity. *Front Immunol* **10**, 616 (2019).