An ERF transcription factor PTI5, a novel regulator of endophyte community

maintenance in potato

Tjaša Lukan[1*], Barbara Kraigher[2], Karmen Pogačar[1,3], Katja Stare[1], Teja Grubar Kovačič[1], Maja

Zagorščak[1], Marko Petek[1], Polonca Stefanic[2], Anže Vozelj[1,3], Valentina Levak[1,3], Tjaša Mahkovec

Povalej[1], Juan M. García[4], Maria J. Pozo[4], Ines Mandić-Mulec[2+], Kristina Gruden[1+*]

[1]Department of Biotechnology and Systems Biology, National Institute of Biology, Ljubljana, Slovenia

[2]Chair of Microbial Ecology and Physiology, Department of Microbiology, Biotechnical Faculty,
University of Ljubljana, Ljubljana, Slovenia

[3]Jožef Stefan International Postgraduate School, Jamova cesta 39, 1000 Ljubljana, Slovenia

[4]Department of Soil and Plant Microbiology, Estación Experimental del Zaidín (CSIC), Granada, Spain

[+]These authors share last authorship

*-corresponding authors
Kristina Gruden (kristina.gruden@nib.si)

ORCID IDs: Tjaša Lukan (0000-0002-6235-2816), Barbara Kraigher (0000-0002-3345-272X), Karmen
Pogačar (0000-0001-5190-7217), Katja Stare (0000-0002-3999-9400), Teja Grubar Kovačič (), Maja
Zagorščak (0000-0002-1669-6482), Marko Petek (0000-0003-3644-7827), Polonca Štefanič (0000-
0002-9444-597X), Anže Vozelj (0009-0003-8758-3244), Valentina Levak (0000-0001-5975-4032),
Tjaša Mahkovec Povalej (0000-0003-3706-2371), Juan M. García (0000-0002-4242-5440), Maria J.
Pozo (0000-0003-2780-9793), Ines Mandic Mulec (0000-0002-7629-3255), Kristina Gruden (0000-
0001-5906-8569)

Keywords: *Solanum tuberosum, Bacillus subtilis*, arbuscular mycorrhizal fungus *Rhizophagus*

irregularis, ERF transcription factor PTI5, endophyte colonisation mechanism, microbial triggered

immunity

ABSTRACT

We have recently identified ERF transcription factor PTI5 as a susceptibility factor, negatively

regulating immune response to diverse pathogens. Here we investigated the processes involved in

colonisation of potato with beneficial organisms. RNAseq showed that at the time of *Bacillus subtilis*

biofilm establishment, immune responses in interacting roots were attenuated, and complex

transcriptional network was triggered, with ethylene signalling being a central module and PTI5

strongly induced. Interestingly, the response is intensified if plants are inoculated by two antagonistic

B. subtilis strains. While PTI5 is not involved in the establishment of biofilm on roots, we show that

bacterial abundance increases in PTI5-silenced plants. Remarkably, root colonization by the

arbuscular mycorrhizal fungus *Rhizophagus irregularis* was also higher in the PTI5-silenced plants.

PTI5 is thus involved both in blocking defence against harmful and blocking colonisation with

beneficial microbes. Such mechanistic understanding of plant–microbe interaction paves the way for

sustainable crop management.

INTRODUCTION

Plant-microbe and microbe–microbe interactions in the rhizosphere determine plant health and

productivity.[1] Beneficial microorganisms are common in rhizosphere soils. Some can endophytically

colonize plants, frequently improving plant nutrition and plant growth, and protect plants from

disease and abiotic stresses through a wide variety of mechanisms.[2,3] They can be efficient and

environmentally friendly alternatives to chemical pesticides and fertilizers, due to their multiple

benefits for the plants and their long shelf life, which is comparable with that of agrochemicals.[4]

However, the current limitation for a wider adoption of microbial inoculants in agriculture is

reproducibility of the results under field conditions.[5] Additional knowledge, such as a deeper

understanding of the mechanisms involved in plant colonization and maintenance of the plant

microbiome is required to improve the use of microbes as plant protectants.[6]

The plant immune system is central in shaping the plant response to beneficial and pathogenic

microorganisms, regulating the extension and successful colonization of plant tissues by mutualistic

organisms and preventing the invasion of the parasitic ones. Recognition of pathogens leads to the

activation of interconnected plant hormonal signaling pathways mediating downstream

transcriptional reprogramming resulting in effective defense responses.[7,8] When plant interacts with

beneficial microbes, microbe triggered immunity (MTI) is activated similarly as in interaction with

pathogens.[9] Beneficial microbes developed several strategies to overcome this immune response and

colonize the plant. One option to avoid effective recognition is by modifying the microbe-associated-

pattern, recognized by immunity, for example flagellin. Alternatively, the microbes develop tolerance

to toxic compounds produced, e.g. reactive oxidative species (ROS) or they actively suppress MTI.[10]

They can also modify metabolism of plant to allow for easier penetration, for example suppress

synthesis of suberin.[11]

Successful colonization by beneficial microbes can induce resistance to pathogenic bacteria and

herbivores. Mechanisms of induced resistance (IR) are not well understood, but it has been shown to

involve jasmonic acid (JA), salicylic acid (SA) and ethylene signaling pathways.[11, 12, 13] Several

individual transcription factors have been shown also to be important for the establishment of the IR

state. As described for bacteria, root colonization by beneficial fungi can also boost plant immunity.

For example, root colonization by arbuscular mycorrhizal fungi is tightly regulated and the

modulation of immune responses during the symbiosis establishment can also leads to induced

resistance in systemic tissues.[15] Therefore, understanding the fine-tuned regulation of plant immune

responses during colonization by beneficials is essential for optimizing bioinoculant-based strategies

in agriculture, allowing their establishment and promoting their benefits in plant health.

In the case of bacteria, social behavior has been proven to be crucial for the success in colonizing

plant tissues. Bacteria often exist in multicellular groups of cells called biofilms,[16] where they engage

in competition for resources but also in cooperative (synergistic) interactions that enhance

community productivity.[17] These social engagements are most often mediated through secreted

molecules that are believed to have a profound impact on bacterial diversity, spatial organization,

and ecological functions.[16,18] *B. subtilis* species are capable of different forms of social interactions

besides biofilm formation, such as quorum sensing (QS), group motility (swarming), and kin

discrimination.[19] Non-kin *B. subtilis* strains form a striking boundary between approaching

swarms,[20,21] exclude a non-kin competitor from a common swarm,[22] and do not mix on *Arabidopsis*

thaliana roots, while kin strains merge on solid medium and cohabit root surfaces.[20] *B. subtilis* strains

also produce a range of antibiotics, surfactants, secondary metabolites, and bioactive volatiles

expected to modulate social interactions between bacteria and with the plant.[20,21,23,24]

B. subtilis QS system comprises the signalling peptide ComX, which positively regulates the synthesis

of surfactin (encoded by Srf operon), one of the most potent surfactants and a lipopeptide

antibiotic.[19,25] Surfactin acts as a signalling molecule during biofilm formation, promotor of horizontal

gene transfer and as an inducer of physiological changes in plants during colonization with

bacteria.[25,26,27,28,29] Despite evidence for *in situ* QS dependent expression of surfactin, the link

between QS and surfactin on plant roots and how QS affects plant colonisation and plant immune

response is less understood.

As potato, the third most important crop in the world, is exceptionally sensitive to a wide range of

environmental stresses,[31] studying its interaction with beneficial microbes is crucial to ensure an

efficient environmentally friendly system of plant protection based on modulation of its microbiome.

The aim of this study was to investigate mechanisms involved in the interaction between potato and

beneficial microbes. Different *B. subtilis* strains and mutants were used to investigate the principles

of bacterial social interactions and their interaction with the plant. We identified transcription factor

PTI5 as a potential regulatory hub for *B. subtilis* endophytic colonization and further investigated its

role in the plant–bacteria interaction using transgenic plants with silenced PTI5. We here show that

B. subtilis abundance is increased in systemic tissue of plants with silenced PTI5, suggesting that PTI5

maintains optimal *B. subtilis* abundance in colonized plants. Interestingly, a similar function was

confirmed also for the establishment of the mutualistic symbiosis between potato and arbuscular

mycorrhizal fungi.

RESULTS

Stable *B. subtilis* biofilm formation on potato roots coincides with first transcriptional responses in plant roots

We here studied the response of potato to colonization by two different *B. subtilis* strains (PS-216 and PS-218). Both formed biofilm on potato roots (Figure S1). As our aim was to capture both initial events in biofilm establishment as well as long-term effects of potato–*B. subtilis* interaction, we first determined timing of biofilm establishment on potato roots and transcriptional responses in the plant. We selected eight genes known to be involved in potato immune response as potential indicators of potato MTI to *B. subtilis* colonization[32,33] and followed their expression in potato roots and shoot tissue. The genes showed diverse expression profiles, which were for some genes tissue specific (Figure S2, Table S1). Two genes, an ERF transcription factor PTI5, which requires an active ethylene pathway for the activation and mediates crosstalk of ethylene and SA pathways[33] and a gene involved in JA synthesis, 13LOX,[32] were reproducibly upregulated in roots and shoots 24 hours post-bacterial inoculation (Figure 1, Figure S2, Table S1). The results were similar for both tested *B. subtilis* strains (Exp1 and Exp2 in Figure S2, Table S1).

121

Figure 1: Dynamics of biofilm formation and plant responses of potato to *B. subtilis*. A) Biofilm formation (YFP

fluorescence) was followed on potato roots after incubation with *B. subtilis* PS-218 YFP for a specified time.

Scale is 50 µm. B) Potato response to bacteria colonization is robustly followed by expression of PTI5 and 13-

LOX. Relative expression of PTI5 and 13-LOX in roots and shoots of *B. subtilis* PS-218 YFP inoculated and non-

inoculated plants, after overnight incubation. Relative gene expression was scaled to the average gene

expression of non-inoculated group. Welch's t-test was used to determine differences between treatments.

Individual measurements (circles), mean (squares) and standard error are shown. Asterisks (*) denote a

statistically significant difference (p-value < 0.05). Data from all experiments are given in Table S1. C) Dynamics

of PTI5 and 13-LOX expression (up- or downregulation) in roots and shoots of potato plants after inoculation

with high *B. subtilis* strain PS-218 culture density (10^7 CFU/mL) and incubation for a specified time. Data of all

experiments are available in Table S1. Up- or downregulation was determined by comparing expression in

inoculated plants with expression in non-inoculated plants. ↑: upregulation, ↓: downregulation, **↑**:

statistically significant upregulation determined by Welch's t-test, **↓**: statistically significant downregulation

determined by Welch's t-test. Exp: experiment.

To study the dynamics of biofilm formation, we set up an experiment with higher density bacteria

inoculum (10^7 CFU/mL). In such set-up, biofilm reproducibly began forming after the first 15 minutes

of incubation, and it was consistently present over the entire root surface after a 2-hour incubation

period (Figure 1). Interestingly, the induction of PTI5 and 13-LOX genes was also detected when the

biofilm was formed uniformly on the root (Figure 1, Figure S2, Table S1).

Complex regulatory network is triggered when potato is colonized with *B. subtilis*

To investigate plant response to *B. subtilis* colonization in more detail, we next performed RNA-seq

analysis of colonized roots and systemic shoot tissue. Our special interest was in the starting point of

colonization, as we intended to capture initial *B. subtilis* recognition related responses in the plant.

Therefore, roots were analyzed at the time of biofilm formation (2 hours post-inoculation (hpi)) and

24 hours later (26 hpi). The response of the plant was stronger at later time points (Figure 2A, Figure

S3). Moreover, shoots were analyzed at this later time to explore potential activation of MTI and/or

IR. For both root and shoot tissue, we identified a large overlap in differentially expressed genes for

plants inoculated with two different strains (Figure 2A, Figure S3).

Figure 2: Potato adjusts to colonization with *B. subtilis* by repression of immune signaling and cell wall synthesis. A) Number of genes, regulated by *B. subtilis* in plant roots and shoots after 2-h and 26-h incubation with PS-216 YFP and PS-218 mKate2. Blue: downregulated genes, red: upregulated genes. B) Visualization of results in the context of biological pathways. For each comparison between *B. subtilis*-inoculated and non-inoculated plants, gene sets enriched for up- or downregulated genes (q-value < 0.1) are shown and designated as red triangles or blue triangles for shoots and roots at different time points. MapMan ontology BINs were used to generate gene sets. Sample group comparisons of inoculated to non-inoculated (in columns) are designated with bacterial strain names used for inoculation. PS-216: PS-216 YFP, PS-218: PS-218 mKate2. See Table S2 for all results. C) *B. subtilis* morphology is changed after internalization in plants. Left: *B. subtilis* morphology 1 dpi (day post-inoculation). Right: *B. subtilis* morphology 3 dpi. See Figure S4 for more data. Scale is 25 μm. D) Electron micrograph of free-living (left) and *in planta* living *B. subtilis* PS-216 (right). Scale is 500 nm.

Many processes related to primary line of defense were downregulated in roots at the time of

biofilm formation, including diverse immune receptors, genes involved in cell wall modifications and

in synthesis of terpenoids and waxes (Figure 2B). Interestingly, among the genes that were

upregulated in roots already at the time of biofilm formation were in majority genes coding for

transcription factors, hormonal synthesis, signalling components and components of protein

degradation. Several ERF transcription factors were upregulated besides the already identified PTI5,

as well as many MYBs, WRKYs, bZIP and NAC transcription factors. We also identified several

lipoxygenases upregulated (JA synthesis), but also genes involved in ethylene and gibberellin

synthesis as well as in Ca^{2+} signalling (Table S3). The response of most of these genes was intensified

in later time point in both roots and shoots (Table S3).

Remarkably, more induced gene sets were detected in shoots. The processes activated in shoots

resemble IR as studied so far in other plants. Receptor kinases were upregulated as were MAP

kinases and executors of plant defense, such as beta-glucanases, alpha amylase inhibitors, PR1

proteins and PR5 proteins (Figure 2B, Table S2, Table S3). Among the genes associated with

transcriptional regulation, NPR1 was also specifically upregulated in shoots only.

In addition, we also followed the extent of bacterial internalization within the plant. We detected

bacteria in stem already 1 day post-inoculation (dpi) (Figure S4). Interestingly, while plant responds

to *B. subtilis* colonization, also *B. subtilis* adapts once internalized as its morphology starts to change

to more rounded form as early as 2 dpi (Figure 2C, Figure S4). Also using electron microscopy, we can

see that most free-living bacteria are multiplying, while no multiplication was detected in *in planta*

living bacteria (Figure 2D, Figure S5).

Biofilm formation and plant responses are attenuated in potato roots inoculated with *B. subtilis*

quorum sensing and surfactin-deficient mutants

To better understand the mechanisms that trigger plant responses, we inspected the role of the *B.*

subtilis ComX peptide mediated QS and surfactin production in potato–*B. subtilis* interaction. We

used two PS-216 mutant strains, one with deletion of the *comQXP* gene cluster, coding for the genes

involved in QS, which also regulate surfactin production (PS-216 ∆c*omQXP*),[25] and the other strain

with a mutation in the *srfA* operon, which is unable to produce surfactin (PS-216 ∆*srfA mutant*). Both

mutants are reported to still form biofilms in liquid media.[34] The biofilm formed by both mutants on

potato root was barely detectable at 26 hours after inoculation, while strong biofilm formation was

observed for wild-type (WT) bacteria (Figure 3A, Figure S6, Table S4). The reduced ability to form

biofilm was also reflected in the potato response to *B. subtilis*. The induction of JA synthesis and

ethylene signalling markers observed in response to the WT was attenuated in the plant response to

the mutants. Similarly, the colonization associated downregulation of apoplastic ROS generator

RBOHD and photosynthesis marker CAB occurred to a lesser extent in response to the mutants

(Figure 3B, Table S5).

Figure 3: Induction of plant responses is dependent on efficient *B. subtilis* biofilm formation on roots. A) Biofilm formation on potato roots after 26-h incubation with YFP-labelled *B. subtilis* PS-216 (left), quorum sensing-negative *B. subtilis* YFP-labelled mutant PS-216 Δ*comQXP* (middle) and PS-216 srfA mutant (right) culture. Scale is 500 μm. B) Expression of selected genes (RBOHD, 13-LOX, CAB, PTI5) in non-inoculated potato roots and after 26-h incubation with *B. subtilis* PS-216 YFP, PS-216 Δ*comQXP* YFP and PS-216 *srfA* YFP. Relative gene expression was scaled to the average gene expression of non-inoculated group. Welch's t-test was used to determine differences between groups. Individual measurements (circles), mean (squares) and standard error

are shown. Asterisks (*) denote statistically significant difference (p-value < 0.05). See Table S5 for the results

of all genes for both roots and shoots.

To confirm that biofilm establishment is required for successful activation of plant responses, and

that *B. subtilis* secondary metabolites are not sufficient, we treated plants with the conditioned

media in which bacteria were growing. Results showed that the conditioned media alone did not

trigger any response in treated plants, albeit containing surfactin (Figure S7, Table S6). Therefore, we

conclude that *B. subtilis* biofilm formation on the roots is required for induction of potato response.

Non-kin intraspecies interaction of *B. subtilis* strains on root intensifies potato responses

To understand plant responses to colonizing bacterial communities, which is relevant for application

in practice, we set up an experiment inoculating plants with two genetically different, categorized

previously as non-kin, and two isogenic (only differently fluorescently labelled) *B. subtilis* strains.

When roots were inoculated with PS-216 and PS-218 strains separately, both equally colonized the

roots (Figure 4A, first two columns). Similarly, when differently labelled isogenic strains were

combined, the resulting biofilm on the root contained similar abundances of each strain, with cells of

each strain spatially well mixed (Figure 4A, third column). On the other hand, when non-kin strains

PS-216 and PS-218 were combined, we observed biofilm with segregation of the two strains (Figure

4A, last column). Moreover, one of the strains dominated on the roots, with only a small amount of

biofilm formed by the other strain (Figure 4A, last column).

Figure 4: Social interactions of *B. subtillis* **strains on potato roots intensify potato immune response.** A)
Colonization of potato roots by isogenic and non-kin *B. subtilis* strains after 4-h incubation. From top to
bottom: YFP fluorescence, mKate2 fluorescence, brightfield, overlay of images, zoom in of overlay images.
From left to right: PS-216 YFP, PS-218 mKate2, PS-218 YFP and PS-218 mKate2 (isogenic – self interaction), PS-
216 YFP and PS-218 mKate2 (non-kin interaction). White color (overlay) shows colocalization of YFP and
mKate2. The scale is 25 μm for **zoom in** images and 100 μm for all others. B) Visualization of responses in the
context of biological pathways. For each comparison between *B. subtilis*-inoculated and non-inoculated plants,
gene sets enriched for up- or downregulated genes (q-value < 0.1) are shown and designated as red triangles or
blue triangles, respectively after 2-h incubation. MapMan ontology BINs were used to generate gene sets.
Sample group comparisons to non-inoculated ones (in columns) are designated with bacterial strain names
used for inoculation. PS-216: PS-216-YFP, PS-218: PS-218-mKate2, isogenic interaction: YFP-labelled PS-218 and
mKate2-labelled PS-218, non-kin interaction: YFP-labelled PS-216 and mKate2-labelled PS-218. See Table S2 for
all results. C) Examples of genes with intensified response when plant is interaction with two non-kin strains.
Log2 values are shown. Blue: downregulated genes, red: upregulated genes, according to scale on the left.

We next performed RNA-seq analysis of roots at the time of uniform establishment of *B. subtilis*
biofilm (2 hpi). In roots, the intensity of the transcriptional response to the non-kin strain mixture

was higher than that to the mixture of two isogenic, differentially labeled strains (Figure 4, Figure S3).

Regarding biological pathways, some receptor kinases (BINs 26.16 and 30.2.8.2) were specifically

upregulated in the non-kin interaction, as well as some enzymes from fatty acid biosynthesis (BIN

11.1.15) and heat shock proteins (BIN 20.2.1) (Figure 4B, Table S2). In addition, 422 genes had more

than 2-fold stronger response in the interaction with non-kin strains compared to monoculture or the

isogenic mixture. Among the more induced ones, are two MYB transcription factors, salicylate

carboxymethyltransferase, gibberellin 2-oxidase 2, peroxidase and beta-galactosidase genes (Table

S3).

B. subtilis **abundance is increased in plants with silenced PTI5**

We next overlayed data with plant stress signaling prior knowledge network, built from experimental

data on protein–protein interactions, protein–DNA interactions and metabolic pathways

(skm.nib.si).[35] The central signaling module that was regulated in the studied interaction was

ethylene signaling, linked to regulated JA synthesis, SA signaling and ROS signaling (Figure 5A; Figure

S8; Table S7). Thus, we focused on ERF transcription factors, as the role of ethylene signaling was so

far not in depth addressed in interaction of plants with beneficial microbes. Among ERFs, PTI5 was

found to be more than 50-fold induced when plants were exposed to *B. subtilis* monocultures or two-

strain mixtures, already early after colonization (after 2-hour incubation). This could indicate its role

as regulatory hub in plant colonization and/or biofilm formation. To test this role, we produced

transgenic potato with reduced expression of PTI5. First, we followed biofilm formation in transgenic

plants with silenced PTI5. Results showed that PTI5 is not involved in biofilm formation on roots

(Figure 5B, Table S8), as no difference in biofilm formation was detected between non-transgenic

(NT) and PTI5-silenced plants (Figure 5B, Figure S9, Table S8).

Figure 5: ERF transcription factor PTI5 is required for balanced colonization of potato plants. A) Plant immune

signalling modules that were regulated following incubation of roots with non-kin mixture of *B. subtilis* strains.

Genes and proteins are represented as nodes and interaction between them (protein–protein interaction,

protein–DNA interaction) as edges of the network. Differentially expressed genes from ethylene (blue), JA

(red), SA (violoet) and other (black) signalling modules are shown. Black edge connected two differentially
expressed genes. All non-differentially expressed genes and edges are shown in grey. The size of the node
indicates the difference in expression between PS-216 and PS-218 inoculated and non-inoculated roots
(log2FC). See Table S7 for the name and description of regulated genes, and Figure S8 for information on
central signalling modules that were regulated in potato roots in isogenic interaction (PS-218 YFP + PS-218
mKate2). B) *B. subtilis* PS-218 YFP root colonization (yellow fluorescence) of transgenic potato with silenced
PTI5 (shPTI5) and non-transgenic potato plant (NT) after 2-h incubation. Scale is 250 µm. C) Bacillus abundance
in shPTI5 and NT potato plants. The scheme shows leaves of four shPTI5 and four NT plants in which the
presence of *B. subtilis* was (+) or was not (-) detected by qPCR in plants from soil which were incubated for 2 h
in PS-218 YFP before being planted and grown in soil for one week. Leaves with no mark were not analyzed.
The table presents a statistically significant difference in the percentage of positive samples (leaves, nodes and
internodes) between NT and shPTI5 plants determined by Fisher's exact test. See Table S9 for details. The
results of line L6 are shown and were confirmed on another transgenic line shPTI5 L2 (Figure S9, Table S10).

PTI5 was also upregulated in potato shoots. To test if it has a role in maintenance of *B. subtilis*
endophytic community we followed *B. subtilis* abundance in shoots of NT and PTI5-silenced plants. *B.*
subtilis abundance in systemic tissue was higher in PTI5-silenced plants, revealing that the balance in
the abundance of endophytic *B. subtilis* community was perturbed (Figure 5C, Table S9, Table S10).

PTI5 is also regulating the extent of potato root colonization by arbuscular mycorrhizal fungi

Considering the role of PTI5 in regulating endophytic colonization of *B. subtilis*, we aimed to explore
if this role was specific for this interaction or could be extended to other beneficial symbiosis. We
decided to explore the potential role of PTI5 in regulating the extension of root colonization by
arbuscular mycorrhizal fungi. This symbiosis is known to be under plant control to keep the
interaction under mutualistic levels and avoid excessive colonization.[36,37] With this aim, we explored
the colonization levels and symbiosis functionality in NT and PTI5-silenced lines. As shown in Figure 6,
R. irregularis successfully colonized potato roots, with intercellular hyphae, arbuscules and vesicles
present in the cortex. Remarkably, the extension of colonization, including the abundance of
arbuscules and vesicles was higher in the PTI5-silenced lines (Figure 6A). Indeed, the percentage of

root length colonized by the fungus was more than 2-fold higher in the silenced lines (Figure 6B), in

agreement with the higher amount of fungal DNA in those plants. To check if the functionality of the

mycorrhizal symbiosis was altered in those lines, we checked the expression levels of plant and

fungal transporters involved in the nutrient exchange between the symbionts within the arbuscules:

the potato phosphate transporter StPT4, and the fungal monosacharide transporter RiMST2, both

transcriptionally upregulated in arbusculated cells and defined as markers for symbiotic functionality

(Figure 6C). Both markers were upregulated in the PTI5-silenced lines, confirming the functionality of

the symbiosis.

Figure 6: Mycorrhizal colonization of potato roots in non-transgenic potato plant (NT) and PTI5-silenced

genotype (shPTI5). A) Representative images of mycorrhizal colonization by *R. irregularis* in the different

genotypes. Fungal structures within the root cortex of potato plants are stained in blue (a: arbuscules, v:

vesicles). Scale is 200 µm. B) Percentage of root length colonized by the arbuscular mycorrhizal fungi *R.*

irregularis. C) Expression analysis of *R. irregularis* constitutive gene (RiEF) and symbiosis marker genes, the *R.*
irregularis monosaccharide transporter (RiMST2) and the potato phosphate transporter (StPT4), as well as
regression analysis using Dunn's pairwise comparisons were used to determine the differences between the
genotypes. Different letters show statistical differences. Permutation test was used to determine the
differences between the genotypes. Individual measurements (circles), mean (squares) and standard error are
shown. Asterisks denote a statistically significant difference (*: p-value < 0.05, **: p-value < 0.01). The results
of analysis of line L6 are shown and were confirmed on another transgenic line shPTI5 L2 (Figure S10).

DISCUSSION

Microbial inoculants have been proven as efficient and environmentally friendly alternatives to
chemical pesticides and fertilizers.[38,39] Current limitation in adopting such agricultural practices is
their variable efficiency in the field due to lack of understanding of mechanisms involved in the
regulation of plant colonization and maintenance of the plant microbiome.[6,40] Therefore, studying
the mechanisms regulating the interactions between crop plants and microbes is crucial to ensure
robustness of such plant protection approach. As the third most important crop in the world,
exceptionally sensitive to a wide range of stresses, potato is one of the most suitable candidates for
plant growth-promoting rhizobia (PGPR) based products.[31] We here established a model system for
such studies, exploring selected *B. subtilis* soil isolates individually or in mixtures. Furthermore, we
explored the potential extension of our findings by testing other widespread mutualistic interaction,
the arbuscular mycorrhizal symbiosis.

B. subtilis initially suppresses immune response in potato roots (e.g. downregulation of diverse
receptor kinases) thereby facilitating its own colonization, while it boosts induction of plant immunity
related genes in shoots at a later stage (Figure 2, Table S2), as it was shown also in other studies of
plants with beneficials as part of induced resistance phenomena.[13,41] Here we show that live bacteria
are required for triggering of the plant response, as the conditioned media in which bacteria were
growing did not trigger the response (Figure S7). Moreover, the intensity of the plant response was
dependent on the extent of biofilm formation on the potato root (Figure 3). Both, the mutant in

surfactin production and mutant in QS (also deficient in surfactin production) showed a reduced

ability to form biofilm on roots. This is interesting as the same strain was able to form floating

biofilms (pellicles),[42] suggesting that the ComX QS control of the biofilm development depends also

on plant factors. Besides residing on the plant roots, *B. subtilis* can also colonize the plant

endophytically. Internalised bacteria stop dividing and change the morphology (Figure 2, Figure S5).

The round bacterial form closely resembles L-form bacteria, which are characterized by modified or

absent cell walls.[42,43,44] We also tested the extent of bacteria internalisation with QS impaired strains,

which was interestingly not affected (Table S11).

To get insights into the complexity of plant–microbe interactions, we extended our studies to

application of two different *B. subtilis* strains to explore how bacterial diversity may influence the

plant response. We studied pairs of strains with different kin discrimination types of intraspecies

interactions — isogenic strains were predicted to be cooperative, and non-kin strains to be

antagonistic.[22] We observed that non-kin interaction of two *B. subtilis* strains invigorates the plant

response, especially in immune and abiotic signalling components (Figure 4, Table S3). These results

imply that even minor taxonomic changes in the phytobiome can lead to substantial differences in

the plant responses. Increasing evidence support that plants actively manage their microbiota for

balancing growth with immunity.[46] It has been reported that *B. subtilis* can promote plant growth via

direct pathways such as producing phytohormones (e.g., auxin, gibberellin, cytokinin, and abscisic

acid).[46,47,48] Both strains used in this study produce auxins (Table S12). Interestingly, we here show

that *B. subtilis* additionally regulates phytohormone metabolism *in planta*. Gibberellin 2-oxidase 2

and cytokinin oxidase, which catalyze the irreversible degradation of cytokinin and gibberellin

phytohormones, have been among the most induced genes by inoculation of non-kin *B. subtilis*

strains (Figure 4, Table S3). This could suggest that equilibrium between growth and immunity is

shifted towards immunity in plants inoculated by non-kin *B. subtilis* strains, as a functional relation

between lower activity of gibberellin signalling and reduced plant disease severity has previously

been confirmed in potato.[50] In accordance, indole-3-acetic acid-amido synthetase GH3.8 was in our

study induced in *B. subtilis* inoculated plants at late time point in both shoots and roots (Table S3).

GH3.8 functions as IAA-amino synthetase that prevents the accumulation of free indole acetic acid

(IAA), which increases plant disease resistance, growth and development.[51]

Not much is however known on the mechanisms that allow for successful colonisation of plants with

beneficial microbes while avoiding colonization by harmful pathogens. MTI is known to be induced

also when plant is exposed to beneficials.[11] In interaction of Arabidopsis with beneficial

Pseudomonas strain it was found that suberin biosynthesis is down regulated to allow for better

penetration of bacteria.[11] Similarly, in our study cell wall metabolism was downregulated in root

colonization (Figure 2, Figure 4). In addition, synthesis of toxic terpenoids was downregulated in

roots at the biofilm forming stage (Figure 2). Suppression of toxic ROS is crucial for successful

colonisation of plants by microbes.[24] We found ROS scavenger peroxidase and ROS generator RBOHD

induced and repressed, respectively, following inoculation by *B. subtilis* (Figure 3). Several studies

also reported the increase of peroxidase in different plant–*Bacillus* interactions, and in some cases

this induction have been associated to pathogen resistance.[51,52,53,54,55,56]

Plant colonisation is frequently linked to induction of SA, JA and ethylene signalling pathways and

transcriptional rewiring, of which MYB72, MYC2 and NPR1 transcription factors are the most studied

ones, and are proposed to be relevant for IR responses.[13,58] Here we found upregulation of the 13-

LOX pathway leading to JA synthesis, as previously described in other plant-PGPR interactions (Figure

1, Figure 3).[58,59,60] Similarly, several studies reported activation of SA signalling pathway.[14] We found

NPR1 upregulated in shoots (Table S2, Table S3). In corroboration with previous studies, we also

identified several MYB transcription factors induced in both roots and shoots (Table S3).

Modulation of ethylene signalling in plant microbiome interactions have been pinpointed as a central

element in their regulation and their impact on plant responses to stress.[62] For example, ethylene

have been proposed to regulate biofilm formation and root colonization by beneficial bacteria[63,64]

and the arbuscular mycorrhizal symbiosis,[65] and it is required for IR by beneficial bacteria and

fungi.[66,67] We also detected induction of the ethylene signalling pathway in *B. subtilis* inoculated

roots, including ETR, EIN3 and several ERF transcription factors (Figure 5, Figure S8, Table S3).

Moreover, 1-aminocyclopropane-1-carboxylate oxidase, which catalyzes the final step in ethylene

biosynthesis, was in our study induced in all tissue and time points, further suggesting involvement of

ethylene signalling pathway in potato response to *B. subtilis* (Figure 5, Figure S8, Table S3).

ERF transcription factors, the members of the AP2/ERF family active under ethylene signling, play

prominent roles in plant immunity, although the function of most of them remains unknown.[13,59,68,69]

A different set of ERFs was regulated in roots and shoots in our study. Among them, ERF transcription

factor PTI5 was strongly upregulated in all studied conditions. PTI5 was first identified in tomato as

interacting partner of the *R* gene *Pto* that confers resistance against the bacterial pathogen

Pseudomonas syringae pv. tomato.[70,71] On the other hand, we have previously shown that PTI5

negatively regulates potato defence response to pathogens potato virus Y and *Ralstonia*

Solanacearum, suggesting its role as susceptibility factor in potato immunity.[33] Interestingly, PTI5 is

transcriptionally regulated by EIN3 in the presence of SA only and on the protein level regulated by

autophagy degradation.[33] Results of this study suggest that PTI5 has a negative regulatory role also in

the maintenance of endophytic community in potato, as both *B. subtilis* abundance in systemic

tissues and mycorrhizal colonization of roots were increased in PTI5-silenced potato plants (Figure 5,

Figure 6). High PTI5 levels thus both increase susceptibility of potato to pathogens and reduce the

capability of plant to interact with beneficial organisms. PTI5 thus represents a target for future

breeding efforts leading towards sustainable potato production. Taking it altogether, we here set a

cornerstone in understanding how beneficial microbes and plant communicate to regulate

endophytic colonization and maintain mutualism and have identified a way for improved colonization

by beneficial through silencing of PTI5.

ACKNOWLEDGEMENTS

We thank Nastja Marondini, Anja Moškrič, Tina Koželj, Tim Godec, Ana Žuran and Olena Nesterenko

for technical support and laboratory assistance. This research was financially supported by the

Slovenian Research Agency grants P4-0165, P4-0116, J4-9302, J4-4550 and J4-3089, and grant

MCIN/AEI/ PID2021-124813OBC31, "ERDF A way of making Europe" financed by the European Union.

AUTHOR CONTRIBUTIONS

Concept and design: TL, MJP, IMM, KG. Acquisition, analysis or interpretation of data: TL, BK, KP, KS,

TGK, MZ, MP, PS, AV, VL, TMP, JMP, MJP, IMM, KG. Drafting of manuscript: TL, KG. Critical revision of

the manuscript for important intellectual content: MZ, MP, PS, MJP, IMM, KG.

DECLARATION OF INTERESTS

The authors declare no competing interests.

METHODS

Plant material

Potato plants cv. Rywal (NT) and shPTI5-Rywal were grown in stem node tissue culture for two weeks

after *in-vitro* micropropagation under controlled environmental conditions (22 ± 2°C in the light and

19 ± 2°C in the dark with 70–90 µmol/m2/s2 radiation (OSRAM L 58 W/77 FLUORA lamps, Germany)

and a 16-h photoperiod) as previously explained.[72] For studying systemic *B. sublilis* spread, plants

were planted in soil two weeks after node segmentation and grown under controlled environmental

conditions for one week as previously described.[73] shPTI5-Rywal transgenic lines were prepared by

introducing shPTI5 construct[33] into NT using *A. tumefaciens* LBA4404.[74] The transformed bacteria

were used for the transformation of NT stem internodes from in vitro plantlets as described in Lukan

et al.[73] using appropriate selection media.

B. subtilis strains

WT _B. subtilis_ strains PS-216 and PS-218[75] tagged with red fluorescent protein (mKate2) and yellow

fluorescent protein (YFP) gene linked to a constitutive hyperspank promotor were used, namely PS-

218 _amyE::$P_{hypercI03}$-YFP_ (Sp) (PS-218 YFP), PS-216 _amyE::$P_{hypercI03}$-YFP_ (Sp) (PS-216 YFP), PS-218

amyE::$P_{hyperspank}$-mKate2 (Chl) (PS-218 mKate2) and PS-216 _amyE::$P_{hyperspank}$-mKate2_ (Chl) (PS-216

mKate2).[20,22] PS-216 mutants with impaired surfactin production PS-216 _amyE::PhypercI03-YFP_ (Sp)

srfA::Tn917 (mls) (PS-216 _srfA_ YFP)[22] and QS mutant PS-216 Δ_comQXP::kan amyE::$P_{hypercI03}$-YFP_ (Sp)

(PS-216 Δ_comQXP_ YFP) (this work) were used. QS mutant PS-216 ΔcomQXP::kan amyE::PhypercI03-

YFP (Sp) was constructed by transformation of PS-216 amyE::PhypercI03-YFP (Sp)[22] with plasmid

pED302,[76] carrying a kanamycin cassette between _degQ_ and _comA_, resulting in a deletion of the

comQXP locus.

IAA production detection

The production of IAA was assessed following the protocol of Pramanik et al.[77] Fresh bacterial

cultures were prepared from single colonies in LB medium supplemented with 0.1 % tryptophan and

incubated in the dark at 30℃ with shaking at 120 rotations per minute (rpm) for 48 hours. After

incubation, cultures were centrifuged at 10,000 rpm for 5 minutes, and the supernatant was

transferred to a microtiter plate in triplicates. Equal volumes of Salkowski reagent (prepared by

adding 1 mL of 0.5 M ferric chloride to 50 mL of 35 % perchloric acid, mixed well, and stored in a

brown bottle) were added to each well. The mixture was incubated in the dark at room temperature

for 30 minutes. Blanks were prepared similarly, using only LB medium with 0,1 % tryptophan.

Absorbance was measured spectrophotometrically at 536 nm, and IAA concentrations were

calculated using a standard curve prepared with concentrations ranging from 10–100 µg/mL of IAA.

Potato inoculation with *B. subtilis*

On day one, *B. subtilis* from permanent culture was plated on a solid LB medium with appropriate

(see methods section *B. subtilis* strains) antibiotics (100 µg/mL spectinomycin or 5 µg/mL

chloramphenicol) and incubated overnight at 37°C. On day two, a single colony was resuspended in 3

mL of LB medium with antibiotics and grown overnight at 200 rpm and 37°C. On day three at noon,

30 µL of the bacteria culture was inoculated in 3 mL of LB medium without antibiotic and grown for 3

hours at 200 rpm at 37°C (OD_{600} between 0.4 and 0.6). Bacterial culture was then diluted in 3 mL of

MS30 (5 g of Murashige and Skoog Basal Medium with vitamins and 30 g of sucrose per 1 L of

bidistilled water, pH 5.8) medium to OD_{600} 0.02.

For studying marker genes expression after potato incubation in bacterial culture of low density, two

weeks old potato plants from tissue cultures were added to diluted bacterial culture. Note that

diluted bacterial culture was incubated 16 hours at 11 rpm before adding potato plants. Shoots were

separated from the culture by parafilm cover and incubated for 21 hours at 11 rpm in a growth

chamber prior sampling for quantitative PCR (qPCR) (Exp1 and Exp2 in Figure S2). When studying if

the conditioned media triggered any response in treated plants (Table S6), plants were added to a

culture (shoots were separated from the culture by parafilm cover) and discarded after 17-hour

incubation. New plants were added to conditioned medium and incubated for 26 hours.

For studying *B. subtilis* morphology after internalization in plants, two weeks old potato plants in

tissue cultures grown in the solid MS30 medium were inoculated with 300 µL of bacterial culture,

which was spread on the medium and after a specified time (Figure S4), stem and leaves were

observed under confocal microscope.

In all other experiments, on day four in the morning (after 16-hour incubation), bacterial cultures

were vortexed, except when studying the role of QS and surfactin (Table S5, Table S6). Cultures of

different strains were mixed in a ratio of 1:1 when interactions were studied. In 3 mL of bacterial

cultures, two-week old potato plants from tissue cultures were added (shoots were separated from

the culture by parafilm cover) and incubated for a specified time (15 min to 26 hours), depending on

the experiment. When studying *B. subtilis* abundance in shPTI5 transgenic plants with qPCR (Table

S9, Table S10), plants were planted in soil after a 2-hour incubation. In all experiments, shoots were

not in contact with bacterial culture.

Mycorrhizal symbiosis establishment and quantification

The mycorrhizal inoculum was maintained as monoxenic culture consisting in an *in vitro* culture of

transformed carrot roots colonized by the arbuscular mycorrhizal fungus *R. irregularis* (MUCL 57021)

(REKA Group. B.V., Bleiswijk, The Netherlands), produced and maintained growing in Gel-Gro (ICN

Biochemicals, Aurora, OH, USA) as previously described in Chabot et al.[78] Potato plantlets growing in

125 mL pots with a mixture of soil, sand and vermiculite (3:2:1, v:v:v) were inoculated by adding a 1

cm piece of the monoxenic culture, containing approximately 50 *R. irregularis* spores, fungal hyphae

and colonized carrot roots. In the non-inoculated treatment, a piece of medium containing only

uninfected carrot roots was applied to the non-mycorrhizal controls. Plants were grown in a growth

chamber (day/night cycle: 16 h, 24°C/8 h, 19°C; relative humidity: 50 %). The experiment was

repeated twice and eight independent plants per genotype were analyzed.

Mycorrhizal colonization of roots and symbiosis establishment was evaluated histochemically and

molecularly mycorrhizal structures within the roots were observed and quantified after clearing the

roots in 10 % KOH and staining the fungal structures with 5 % black ink in 2 % acetic acid solution as

described in García at al.[79] Percentage of mycorrhizal colonization of roots was determined following

the gridline intersection method[80] using a Nikon SMZ1000 stereomicroscope. To quantify the total

amount of fungus within the root and assess symbiotic functionality we analysed the expression

levels of marker genes by qPCR (see below).

Confocal microscopy

Stellaris 8 Confocal microscope with HC PL FLUOTAR 10x/0.30 DRY objective (Leica Microsystems) was used to detect emission of YFP and mKate2 on the roots of plants in the experiment for investigating timing of biofilm formation, *B. subtilis* morphology after internalization in plants, biofilm formation pattern when studying bacteria–plant social interactions, biofilm presence on the roots of plants sampled for RNA-seq, biofilm formation in transgenic plants with silenced PTI5 and prior all qPCR experiments to confirm roots colonization. Confocal microscope Leica TCS LSI macroscope with Plan APO 5 x objective (Leica Microsystems) was used to detect emission of YFP when investigating biofilm formation on the roots of mutants that have attenuated surfactin production. The emission of YFP was followed after excitation with 488 nm laser in the window 505–560 nm. The emission of mKate was followed after excitation with 589 nm laser in the window between 605–650 nm. Regions of interest (ROI) were scanned unidirectionally with scan speed of 400 Hz and frame average at least 2. The images were processed to obtain maximum projections from Z-stacks for all channels and merged using Leica LAS X software (Leica Microsystems). Mean intensity of fluorescence on confocal images was determined in LAS X software (Leica). Statistically significant differences were determined by Student's t-test.

Transmission electron microscopy (TEM)

Samples were prepared for examination with TEM using negative staining method. Pellets of centrifuged liquid culture or leaf homogenate infiltrated with *B. subtilis* were resuspended in 0.1 M phosphate buffer with 2 % polyvinylpyrrolidone. Six µL of resuspension was applied to copper grids with or without glow discharge (400 mesh, formvar-carbon coated) for 5 minutes. Afterwards, the remaining resuspension was dabbed with filter paper. This procedure was repeated two more times.

Afterwards, the grids were washed with bidistilled water and stained with one droplet of 1 % (w/v)

water solution of uranyl acetate. Two grids were prepared for each sample. The grids were observed

by transmission electron microscope TALOS L120 (ThermoFisher SCIENTIFIC, The Netherlands),

operating at 100 kV and representative micrographs were acquired (camera Ceta 16M) using Velox

software.

qPCR analysis

The shoots and elongation zones of roots of *B. subtilis*-inoculated and non-inoculated (see

description of inoculation chapter) potato plants were sampled in four or five biological replicates

(plants). RNA was isolated by RNeasy Plant Mini Kit (Qiagen) from shoots and RNeasy Plant Mini Kit

(Qiagen) or RNeasy Micro Kit (Qiagen) from roots according to the manufacturer's protocols.

Concentration and purity of DNase-treated (0.5 μL of DNase I per μg RNA; Qiagen) total RNA was

evaluated using NanoDrop ND1000 spectrophotometer (Nanodrop technologies) and agarose gel

electrophoresis and then reverse-transcribed using the High-Capacity cDNA Reverse Transcription Kit

(Thermo Fisher Scientific) or High-Capacity RNA-to-cDNA Kit (Thermo Fisher Scientific). To determine

B. subtilis abundance in systemic shoot tissue, genomic DNA was isolated by MagMAX™ Plant DNA

Isolation Kit (Thermo Fisher Scientific) according to the manufacturer's protocol. Concentration and

purity were evaluated as stated above.

Samples were analyzed in the setup for qPCR as previously described.[32] The expression of 8 genes

(RBOHD, HSP70, PTI5, 13-LOX, PR1B, BGLU2, CPI8, CAB) involved in different steps of immune

signaling (Table S13 for a full list of genes and their description) was determined and normalized to

the expression of two validated reference genes, COX and elongation factor 1 (EF-1), as described

previously.[81] See Lukan et al.[32] for primer and probe information. Relative copy numbers of *comQ*

from *B. subtilis* were determined and normalized to relative copy numbers of NPR1 (a single copy

gene, https://spuddb.uga.edu/). The standard curve method was used for relative gene expression

quantification using quantGenius (http://quantgenius.nib.si).[82]

For molecular assessment of mycorrhizal colonization, the expression of the constitutive *R.*

irregularis elongation factor 1 (RiEF) gene and the markers of symbiotic functionality, the fungal

RiMST2 and the plant StPT4, both involved in the nutrient exchange between the symbionts in the

arbuscules were determined in roots by qPCR using gene-specific primers (Table S13). Relative

quantification of specific mRNA levels was performed using the comparative $2-\Delta(\Delta Ct)$ method.[83]

Expression values were normalized using the normalizer gene EF-1α encoding the potato translation

elongation factor-1α. [84] Five independent biological replicates per treatment were analyzed.

Statistical analysis was conducted in R version 4.3.1.[85] Relative gene expression was scaled to the

average relative gene expression of the non-inoculated group.[86] Mean, standard deviation, standard

error of the mean and 95 % confidence intervals were calculated per gene, tissue type and genotype

and visualised using the ggplot2 package v3.3.6.[87] Pairwise Welch's t-test with holm correction for

multiple testing adjustment was employed to quantify the difference between the groups. Results

were visualised using rstatix v0.7.0[88] and ggpubr v0.4.0[89] R packages.

RNA-seq analysis

RNA was isolated from shoots and elongation zones of roots by RNeasy Plant Mini Kit (Qiagen) and

treated with RNA Clean & Concentrator-5 (DNase Included) (Zymo Research). In RNA-seq Exp 1, RNA-

seq was for roots performed after 2- and 26-hours incubations with *B. subtilis* culture and for shoots

after 26 hours of incubation, both for monocultures (PS-216 YFP, PS-218 YFP) and non-inoculated

samples (incubation in MS30 medium) as a control. In RNA-seq Exp 2, RNA-seq was for roots

performed after 2 hours of incubation with *B. subtilis* culture for monocultures (PS-216 YFP, PS-218

YFP, PS-218 mKate2), isogenic mixture (PS-218 mKate2 + PS-218 YFP), non-kin mixture (PS-218

mKate2 + PS-216 YFP) and non-inoculated samples (incubation in MS30 medium) as a control. Four

biological replicates (plants) were analyzed for each interaction (Table S14).

Raw paired-end Illumina reads were adapter-trimmed and reads below average PHRED basecall

quality score 20 were removed using Trim Galore (https://github.com/FelixKrueger/TrimGalore).[90]

Sequencing quality control was performed using FastQC v0.11.9.[91] Taxonomic classification of reads

was performed using Centrifuge v1.0.4[92] in conjunctions with the 2018 nt database. Reads were

mapped to *Solanum tuberosum* Phureja clone DM 1-3 genome with STAR v2.7.5c[93] using the merged

genome annotations GFF3.[94] Only uniquely mapped fragments were counted (--

outFilterMultimapNmax 1) and additional alignment parameters were set to restrict erroneous

mapping (--outFilterMismatchNoverReadLmax 0.02, --quantTranscriptomeBan Singleend, --

outFilterType BySJout, --alignSJoverhangMin 10, --alignSJDBoverhangMin 1, --alignIntronMin 20, --

alignIntronMax 10000, --alignMatesGapMax 10000). Centrifuge and mapping results showed

substantial contamination of sample BS472, therefore we excluded this sample from further analysis.

Gene Set Enrichment Analysis[95] was performed using MapMan[96] based gene sets (gmt file available

at https://github.com/nib-si/ptda)[97] and non-filtered normalized read counts, comparing different

Bacillus treatments to non-inoculated plants to identify significantly altered regulation of processes

and functionally related gene groups (FDR corrected q-value < 0.10). Results of all GSEA comparisons

were merged using R v4.3.1.[85] GSEA figure was generated using adjusted gseaFromStats function

from biokit v0.1.1 package.[98]

Differential gene expression analysis was performed in R using the *limma* package.[99] Low expressed

genes were filtered by retaining only genes with raw read counts above 100 in at least 11 samples for

RNA-seq Exp 2 and above 50 counts in at least 4 samples for RNA-seq Exp 1. Root and leaf samples

were analyzed separately. Normalized counts were transformed using voom function, followed by

linear model fit, and statistics calculation for defined contrasts using the eBayes function. Genes with

Benjamini-Hochberg FDR adjusted p-values < 0.05 and $|log_2FC| > 1$ were considered significantly

differentially expressed. Venn diagrams were plotted using a modified limma::vennDiagram function.

Raw and normalized read counts were deposited in GEO under accession number GSE232028.

Results of RNA-seq were validated by qPCR (Table S1).

Network analysis

Mechanistic Plant Stress Signalling (PSS) (https://skm.nib.si/)[35] reaction network was pre-processed

using DiNARs'[100] pre-processing app. Interactions among different molecular entities (such as genes,

proteins or transcripts) are defined on functional cluster level, mostly encompassed of *A. thaliana*

gene identifiers. Therefore, *Solanum tuberosum* gene identifiers[94] were first translated to Arabidopsis

gene identifiers using BLAST+ reciprocal best hit (RBH) search[101] on proteomes (file available at

https://github.com/NIB-SI/DiNAR/tree/master/TranslationTables). Top three potato identifiers per

Arabidopsis identifier were considered and further prioritised to the functinal cluster using DiNARs'

prioritisation script. Experimental data were superimposed onto the PSS network and visualised

using DiNAR. Separate condition specific networks, as defined by differential expression contrasts,

were exported as individual interactive .html files for further examination.

REFERENCES

1.   Mauchline, T.H., and Malone, J.G. (2017). Life in earth – the root microbiome to the rescue?
Curr Opin Microbiol *37*, 23–28. https://doi.org/10.1016/j.mib.2017.03.005.

2.   Timmusk, S., Behers, L., Muthoni, J., Muraya, A., and Aronsson, A.C. (2017). Perspectives and
challenges of microbial application for crop improvement. Front Plant Sci *8*.
https://doi.org/10.3389/fpls.2017.00049.

3.   Pozo, M.J., Zabalgogeazcoa, I., Vazquez de Aldana, B.R., and Martinez-Medina, A. (2021).
Untapping the potential of plant mycobiomes for applications in agriculture. Curr Opin Plant
Biol *60*. https://doi.org/10.1016/j.pbi.2021.102034.

4.   Kandel, S.L., Joubert, P.M., and Doty, S.L. (2017). Bacterial endophyte colonization and
distribution within plants. Microorganisms *5*.
https://doi.org/10.3390/microorganisms5040077.

5. O'Callaghan, M., Ballard, R.A., and Wright, D. (2022). Soil microbial inoculants for sustainable
agriculture: Limitations and opportunities. Soil Use Manag *38*, 1340–1369.
https://doi.org/10.1111/sum.12811.

6. Lutz, S., Bodenhausen, N., Hess, J., Valzano-Held, A., Waelchli, J., Deslandes-Hérold, G.,
Schlaeppi, K., and van der Heijden, M.G.A. (2023). Soil microbiome indicators can predict crop
growth response to large-scale inoculation with arbuscular mycorrhizal fungi. Nat Microbiol *8*,
2277–2289. https://doi.org/10.1038/s41564-023-01520-w.

7. Ngou, B.P.M., Ding, P., and Jones, J.D.G. (2022). Thirty years of resistance: Zig-zag through the
plant immune system. Plant Cell *34*, 1447–1478. https://doi.org/10.1093/plcell/koac041.

8. Baebler, Š., Witek, K., Petek, M., Stare, K., Tušek-Žnidarič, M., Pompe-Novak, M., Renaut, J.,
Szajko, K., Strzelczyk-Zyta, D., Marczewski, W., et al. (2014). Salicylic acid is an indispensable
component of the Ny-1 resistance-gene-mediated response against Potato virus y infection in
potato. J Exp Bot *65*, 1095–1109. https://doi.org/10.1093/jxb/ert447.

9. Tsai, H.H., Wang, J., Geldner, N., and Zhou, F. (2023). Spatiotemporal control of root immune
responses during microbial colonization. Curr Opin Plant Biol *74*.
https://doi.org/10.1016/j.pbi.2023.102369.

10. Teixeira, P.J.P.L., Colaianni, N.R., Law, T.F., Conway, J.M., Gilbert, S., Li, H., Salas-González, I.,
Panda, D., Risco, N.M. Del, Finkel, O.M., et al. (2021). Specific modulation of the root immune
system by a community of commensal bacteria. Proc Natl Acad Sci U S A *118*.
https://doi.org/10.1073/pnas.2100678118.

11. Verbon, E.H., Liberman, L.M., Zhou, J., Yin, J., Pieterse, C.M.J., Benfey, P.N., Stringlis, I.A., and
de Jonge, R. (2023). Cell-type-specific transcriptomics reveals that root hairs and endodermal
barriers play important roles in beneficial plant-rhizobacterium interactions. Mol Plant *16*,
1160–1177. https://doi.org/10.1016/j.molp.2023.06.001.

12. De Kesel, J., Conrath, U., Flors, V., Luna, E., Mageroy, M.H., Mauch-Mani, B., Pastor, V., Pozo,
M.J., Pieterse, C.M.J., Ton, J., et al. (2021). The Induced Resistance Lexicon: Do's and Don'ts.
Trends Plant Sci *26*, 685–691. https://doi.org/10.1016/j.tplants.2021.01.001.

13. Pieterse, C.M.J., Zamioudis, C., Berendsen, R.L., Weller, D.M., Van Wees, S.C.M., and Bakker,
P.A.H.M. (2014). Induced systemic resistance by beneficial microbes. Annu Rev Phytopathol
52, 347–375. https://doi.org/10.1146/annurev-phyto-082712-102340.

14. Yu, Y., Gui, Y., Li, Z., Jiang, C., Guo, J., and Niu, D. (2022). Induced Systemic Resistance for
Improving Plant Immunity by Beneficial Microbes. Plants *11*.
https://doi.org/10.3390/plants11030386.

15. Fiorilli, V., Martínez-Medina, A., Pozo, M.J., and Lanfranco, L. (2024). Plant Immunity
Modulation in Arbuscular Mycorrhizal Symbiosis and Its Impact on Pathogens and Pests. Annu
Rev Phytopathol *62*, 127–156. https://doi.org/10.1146/annurev-phyto-121423-042014.

16. Flemming, H.C., Wingender, J., Szewzyk, U., Steinberg, P., Rice, S.A., and Kjelleberg, S. (2016).
Biofilms: An emergent form of bacterial life. Nat Rev Microbiol *14*, 563–575.
https://doi.org/10.1038/nrmicro.2016.94.

17.   Mitri, S., and Richard Foster, K. (2013). The genotypic view of social interactions in microbial
communities. Annu Rev Genet *47*, 247–273. https://doi.org/10.1146/annurev-genet-111212-
133307.

18.   Asfahl, K.L., and Schuster, M. (2017). Social interactions in bacterial cell-cell signaling. FEMS
Microbiol Rev *41*, 92–107. https://doi.org/10.1093/femsre/fuw038.

19.   Kalamara, M., Spacapan, M., Mandic-Mulec, I., and Stanley-Wall, N.R. (2018). Social
behaviours by Bacillus subtilis: quorum sensing, kin discrimination and beyond. Mol Microbiol
110, 863–878. https://doi.org/10.1111/mmi.14127.

20.   Stefanic, P., Kraigher, B., Lyons, N.A., Kolter, R., and Mandic-Mulec, I. (2015). Kin
discrimination between sympatric Bacillus subtilis isolates. Proc Natl Acad Sci U S A *112*,
14042–14047. https://doi.org/doi.org/10.1073/pnas.1512671112.

21.   Lyons, N.A., Kraigher, B., Stefanic, P., Mandic-Mulec, I., and Kolter, R. (2016). A combinatorial
kin discrimination system in Bacillus subtilis. Current Biology *26*, 733–742.
https://doi.org/10.1016/j.cub.2016.01.032.

22.   Kraigher, B., Butolen, M., Stefanic, P., and Mandic Mulec, I. (2022). Kin discrimination drives
territorial exclusion during Bacillus subtilis swarming and restrains exploitation of surfactin.
ISME Journal *16*, 833–841. https://doi.org/10.1038/s41396-021-01124-4.

23.   Lareen, A., Burton, F., and Schäfer, P. (2016). Plant root-microbe communication in shaping
root microbiomes. Plant Mol Biol *90*, 575–587. https://doi.org/10.1007/s11103-015-0417-8.

24.   Liu, Y., Xu, Z., Chen, L., Xun, W., Shu, X., Chen, Y., Sun, X., Wang, Z., Ren, Y., Shen, Q., et al.
(2024). Root colonization by beneficial rhizobacteria. FEMS Microbiol Rev *48*.
https://doi.org/10.1093/femsre/fuad066.

25.   Oslizlo, A., Stefanic, P., Dogsa, I., and Mandic-Mulec, I. (2014). Private link between signal and
response in Bacillus subtilis quorum sensing. Proc Natl Acad Sci U S A *111*, 1586–1591.
https://doi.org/10.1073/pnas.1316283111.

26.   Ongena, M., Jourdan, E., Adam, A., Paquot, M., Brans, A., Joris, B., Arpigny, J.L., and Thonart,
P. (2007). Surfactin and fengycin lipopeptides of Bacillus subtilis as elicitors of induced
systemic resistance in plants. Environ Microbiol *9*, 1084–1090.
https://doi.org/10.1111/j.1462-2920.2006.01202.x.

27.   López, D., Fischbach, M.A., Chu, F., Losick, R., and Kolter, R. (2009). Structurally diverse natural
products that cause potassium leakage trigger multicellularity in Bacillus subtilis. Proc Natl
Acad Sci U S A *106*, 280–285. https://doi.org/doi.org/10.1073/pnas.0810940106.

28.   Danevčič, T., Dragoš, A., Spacapan, M., Stefanic, P., Dogsa, I., and Mandic-Mulec, I. (2021).
Surfactin Facilitates Horizontal Gene Transfer in Bacillus subtilis. Front Microbiol *12*.
https://doi.org/10.3389/fmicb.2021.657407.

29.   Rahman, F. Bin, Sarkar, B., Moni, R., and Rahman, M.S. (2021). Molecular genetics of surfactin
and its effects on different sub-populations of Bacillus subtilis. Biotechnology Reports *32*.
https://doi.org/10.1016/j.btre.2021.e00686.

30.   Chaudhry, V., Runge, P., Sengupta, P., Doehlemann, G., Parker, J.E., and Kemen, E. (2021).
Shaping the leaf microbiota: Plant-microbe-microbe interactions. J Exp Bot *72*, 36–56.
https://doi.org/10.1093/jxb/eraa417.

31. Vilvert, E., Stridh, L., Andersson, B., Olson, Å., Aldén, L., and Berlin, A. (2022). Evidence based
disease control methods in potato production: a systematic map protocol. Environ Evid *11*.
https://doi.org/10.1186/s13750-022-00259-x.

32. Lukan, T., Pompe-Novak, M., Baebler, Š., Tušek-Žnidarič, M., Kladnik, A., Križnik, M., Blejec, A.,
Zagorščak, M., Stare, K., Dušak, B., et al. (2020). Precision transcriptomics of viral foci reveals
the spatial regulation of immune-signaling genes and identifies RBOHD as an important player
in the incompatible interaction between potato virus Y and potato. Plant Journal *104*, 645–
661. https://doi.org/10.1111/tpj.14953.

33. Coll, A., Lukan, T., Stare, K., Zagorščak, M., Mahkovec Povalej, T., Baebler, Š., Prat, S., Coll,
N.S., Valls, M., Petek, M., et al. (2024). The StPti5 ethylene response factor acts as a
susceptibility factor by negatively regulating the potato immune response to pathogens. New
Phytologist *244*, 202–218. https://doi.org/10.1111/nph.20004.

34. Thérien, M., Kiesewalter, H.T., Auria, E., Charron-Lamoureux, V., Wibowo, M., Maróti, G.,
Kovács, Á.T., and Beauregard, P.B. (2020). Surfactin production is not essential for pellicle and
root-associated biofilm development of Bacillus subtilis. Biofilm *2*.
https://doi.org/10.1016/j.bioflm.2020.100021.

35. Bleker, C., Ramšak, Ž., Bittner, A., Podpečan, V., Zagorščak, M., Wurzinger, B., Baebler, Š.,
Petek, M., Križnik, M., van Dieren, A., et al. (2024). Stress Knowledge Map: A knowledge graph
resource for systems biology analysis of plant stress responses. Plant Commun *5*.
https://doi.org/10.1016/j.xplc.2024.100920.

36. Lidoy, J., López-García, Á., Amate, C., García, J.M., Flors, V., García-Garrido, J.M., Azcón-
Aguilar, C., López-Raez, J.A., and Pozo, M.J. (2023). Regulation of mycorrhizal colonization
under stress in tomato depends on symbiotic efficiency. Environ Exp Bot *215*.
https://doi.org/10.1016/j.envexpbot.2023.105479.

37. Ho-Plágaro, T., and García-Garrido, J.M. (2022). Molecular Regulation of Arbuscular
Mycorrhizal Symbiosis. Int J Mol Sci *23*. https://doi.org/10.3390/ijms23115960.

38. Alves de Andrade, L., Santos, C.H.B., Frezarin, E.T., Sales, L.R., and Rigobelo, E.C. (2023). Plant
Growth-Promoting Rhizobacteria for Sustainable Agricultural Production. Microorganisms *11*.
https://doi.org/10.3390/microorganisms11041088.

39. Minchev, Z., Ramírez-Serrano, B., Dejana, L., Lee Díaz, A.S., Zitlalpopoca-Hernandez, G., Orine,
D., Saha, H., Papantoniou, D., García, J.M., González-Céspedes, A., et al. (2024). Beneficial soil
fungi enhance tomato crop productivity and resistance to the leaf-mining pest Tuta absoluta
in agronomic conditions. Agron Sustain Dev *44*. https://doi.org/10.1007/s13593-024-00991-3.

40. Tsotetsi, T., Nephali, L., Malebe, M., and Tugizimana, F. (2022). Bacillus for Plant Growth
Promotion and Stress Resilience: What Have We Learned? Plants *11*.
https://doi.org/10.3390/plants11192482.

41. Blake, C., Christensen, M.N., and Kovacs, A.T. (2021). Molecular aspects of plant growth
promotion and protection by Bacillus subtilis. Molecular Plant-Microbe Interactions *34*, 15–
25. https://doi.org/10.1094/MPMI-08-20-0225-CR.

42. Špacapan, M., Danevčič, T., Štefanic, P., Porter, M., Stanley-Wall, N.R., and Mandic-Mulec, I.
(2020). The ComX quorum sensing peptide of Bacillus subtilis affects biofilm formation

negatively and sporulation positively. Microorganisms *8*.
https://doi.org/10.3390/microorganisms8081131.

43. Allan, E.J., Hoischen, C., and Gumpert, J. (2009). Chapter 1 Bacterial L-Forms
https://doi.org/10.1016/S0065-2164(09)01201-5.

44. Leaver, M., Domínguez-Cuevas, P., Coxhead, J.M., Daniel, R.A., and Errington, J. (2009). Life
without a wall or division machine in Bacillus subtilis. Nature *457*, 849–853.
https://doi.org/10.1038/nature07742.

45. Daulagala, P.W.H.K.P. (2021). Association of L-form Bacteria with Plants and their Application
in Biological Control of Phytopathogenic Fungi and Bacteria: A Review. J Adv Microbiol *21*, 77–
86. https://doi.org/10.9734/jamb/2021/v21i1030395.

46. Ku, Y.S., Liao, Y.J., Chiou, S.P., Lam, H.M., and Chan, C. (2024). From trade-off to synergy:
microbial insights into enhancing plant growth and immunity. Plant Biotechnol J *22*, 2461–
2471. https://doi.org/10.1111/pbi.14360.

47. Karadeniz, A., Topcuoǧlu, Ş.F., and Inan, S. (2006). Auxin, gibberellin, cytokinin and abscisic
acid production in some bacteria. World J Microbiol Biotechnol *22*, 1061–1064.
https://doi.org/10.1007/s11274-005-4561-1.

48. Hamdache, A., Lamarti, A., Aleu, J., and Collado, I.G. (2011). Non-peptide metabolites from
the genus Bacillus. J Nat Prod *74*, 893–899. https://doi.org/10.1021/np100853e.

49. Gamalero, E., and Glick, B.R. (2011). Chapter 2 Mechanisms Used by Plant Growth-Promoting
Bacteria D. K. Maheshwari, ed. (Springer) https://doi.org/10.1007/978-3-642-21061-7_2.

50. Križnik, M., Petek, M., Dobnik, D., Ramšak, Ž., Baebler, Š., Pollmann, S., Kreuze, J.F., Žel, J., and
Gruden, K. (2017). Salicylic acid perturbs sRNA-gibberellin regulatory network in immune
response of potato to potato virus Y infection. Front Plant Sci *8*.
https://doi.org/10.3389/fpls.2017.02192.

51. Xu, G., Zhang, Y., Li, M., Jiao, X., Zhou, L., and Ming, Z. (2021). Crystal structure of the acyl acid
amido synthetase GH3-8 from Oryza sativa. Biochem Biophys Res Commun *534*, 266–271.
https://doi.org/10.1016/j.bbrc.2020.11.098.

52. Chowdappa, P., Mohan Kumar, S.P., Jyothi Lakshmi, M., and Upreti, K.K. (2013). Growth
stimulation and induction of systemic resistance in tomato against early and late blight by
Bacillus subtilis OTPB1 or Trichoderma harzianum OTPB3. Biological Control *65*, 109–117.
https://doi.org/10.1016/j.biocontrol.2012.11.009.

53. Lian, L., Xie, L., Zheng, L., and Lin, Q. (2011). Induction of systemic resistance in tobacco
against Tobacco mosaic virus by Bacillus spp. Biocontrol Sci Technol *21*, 281–292.
https://doi.org/10.1080/09583157.2010.543667.

54. Guo, Q., Li, Y., Lou, Y., Shi, M., Jiang, Y., Zhou, J., Sun, Y., Xue, Q., and Lai, H. (2019). Bacillus
amyloliquefaciens Ba13 induces plant systemic resistance and improves rhizosphere
microecology against tomato yellow leaf curl virus disease. Applied Soil Ecology *137*, 154–166.
https://doi.org/10.1016/j.apsoil.2019.01.015.

55. Srivastava, S., Bist, V., Srivastava, S., Singh, P.C., Trivedi, P.K., Asif, M.H., Chauhan, P.S., and
Nautiyal, C.S. (2016). Unraveling aspects of Bacillus amyloliquefaciens mediated enhanced

production of rice under biotic stress of Rhizoctonia solani. Front Plant Sci *7*.
https://doi.org/10.3389/fpls.2016.00587.

56. Yasmin, S., Zaka, A., Imran, A., Zahid, M.A., Yousaf, S., Rasul, G., Arif, M., and Mirza, M.S.
(2016). Plant growth promotion and suppression of bacterial leaf blight in rice by inoculated
bacteria. PLoS One *11*. https://doi.org/10.1371/journal.pone.0160688.

57. Mehmood, S., Muneer, M.A., Tahir, M., Javed, M.T., Mahmood, T., Afridi, M.S., Pakar, N.P.,
Abbasi, H.A., Munis, M.F.H., and Chaudhary, H.J. (2021). Deciphering distinct biological control
and growth promoting potential of multi-stress tolerant Bacillus subtilis PM32 for potato stem
canker. Physiology and Molecular Biology of Plants *27*, 2101–2114.
https://doi.org/10.1007/s12298-021-01067-2.

58. Zamioudis, C., Korteland, J., Van Pelt, J.A., Van Hamersveld, M., Dombrowski, N., Bai, Y.,
Hanson, J., Van Verk, M.C., Ling, H.Q., Schulze-Lefert, P., et al. (2015). Rhizobacterial volatiles
and photosynthesis-related signals coordinate MYB72 expression in Arabidopsis roots during
onset of induced systemic resistance and iron-deficiency responses. Plant Journal *84*, 309–
322. https://doi.org/10.1111/tpj.12995.

59. Wu, L., Huang, Z., Li, X., Ma, L., Gu, Q., Wu, H., Liu, J., Borriss, R., Wu, Z., and Gao, X. (2018).
Stomatal closure and SA-, JA/ET-signaling pathways are essential for Bacillus
amyloliquefaciens FZB42 to restrict leaf disease caused by Phytophthora nicotianae in
Nicotiana benthamiana. Front Microbiol *9*. https://doi.org/10.3389/fmicb.2018.00847.

60. Cawoy, H., Mariutto, M., Henry, G., Fisher, C., Vasilyeva, N., Thonart, P., Dommes, J., and
Ongena, M. (2014). Plant defense stimulation by natural isolates of Bacillus depends on
efficient surfactin production. Molecular Plant-Microbe Interactions *27*, 87–100.
https://doi.org/10.1094/MPMI-09-13-0262-R.

61. Zebelo, S., Song, Y., Kloepper, J.W., and Fadamiro, H. (2016). Rhizobacteria activates (+)-δ-
cadinene synthase genes and induces systemic resistance in cotton against beet armyworm
(Spodoptera exigua). Plant Cell Environ *39*, 935–943. https://doi.org/10.1111/pce.12704.

62. Ravanbakhsh, M., Sasidharan, R., Voesenek, L.A.C.J., Kowalchuk, G.A., and Jousset, A. (2018).
Microbial modulation of plant ethylene signaling: ecological and evolutionary consequences.
Microbiome *6*. https://doi.org/10.1186/s40168-018-0436-1.

63. Berrabah, F., Balliau, T., Aït-Salem, E.H., George, J., Zivy, M., Ratet, P., and Gourion, B. (2018).
Control of the ethylene signaling pathway prevents plant defenses during intracellular
accommodation of the rhizobia. New Phytologist *219*, 310–323.
https://doi.org/10.1111/nph.15142.

64. Carlew, S.S., Brenya, E., Ferdous, M., Banerjee, I., Donnelly, L., Heinze, E., King, J., Sexton, B.,
Lacey, R.F., Bakshi, A., et al. (2025). Ethylene signals through an ethylene receptor to
modulate biofilm formation and root colonization in a beneficial plant-associated bacterium.
PLoS Genet *21*. https://doi.org/10.1371/journal.pgen.1011587.

65. Das, D., Varshney, K., Ogawa, S., Torabi, S., Hüttl, R., Nelson, D.C., and Gutjahr, C. (2025).
Ethylene promotes SMAX1 accumulation to inhibit arbuscular mycorrhiza symbiosis. Nature
Communications *16*. https://doi.org/10.1038/s41467-025-57222-w.

66. Knoester, M., Pieterse, C.M.J., Bol, J.F., and Van Loon, L.C. (1999). Systemic Resistance in
Arabidopsis Induced by Rhizobacteria Requires Ethylene-Dependent Signaling at the Site of

Application. Molecular Plant-Microbe Interactions *12*, 720–727.
https://doi.org/10.1094/MPMI.1999.12.8.720.

67.   Lidoy, J., Rivero, J., Ramšak, Ž., Petek, M., Križnik, M., Flors, V., Lopez-Raez, J.A., Martinez-
Medina, A., Gruden, K., and Pozo, M.J. (2025). Ethylene signaling is essential for mycorrhiza-
induced resistance against chewing herbivores in tomato. J Exp Bot.
https://doi.org/10.1093/jxb/eraf053.

68.   Huang, P.Y., Catinot, J., and Zimmerli, L. (2016). Ethylene response factors in Arabidopsis
immunity. J Exp Bot *67*, 1231–1241. https://doi.org/10.1093/jxb/erv518.

69.   Chowdhury, S.P., Uhl, J., Grosch, R., Alquéres, S., Pittroff, S., Dietel, K., Schmitt-Kopplin, P.,
Borriss, R., and Hartmann, A. (2015). Cyclic lipopeptides of Bacillus amyloliquefaciens subsp.
plantarum colonizing the lettuce rhizosphere enhance plant defense responses toward the
bottom rot pathogen Rhizoctonia solani. Molecular Plant-Microbe Interactions *28*, 984–995.
https://doi.org/10.1094/MPMI-03-15-0066-R.

70.   Zhou, J., Tang, X., and Martin, G.B. (1997). The Pto kinase conferring resistance to tomato
bacterial speck disease interacts with proteins that bind a cis-element of pathogenesis-related
genes. EMBO J *16*, 3207–3218. https://doi.org/10.1093/emboj/16.11.3207.

71.   He, P., Warren, R.F., Zhao, T., Shan, L., Zhu, L., Tang, X., and Zhou, J.-M. (2001).
Overexpression of Pti5 in Tomato Potentiates Pathogen-Induced Defense Gene Expression
and Enhances Disease Resistance to Pseudomonas syringae pv. tomato. Molecular Plant-
Microbe Interactions *14*, 1453–1457. https://doi.org/10.1094/MPMI.2001.14.12.1453.

72.   Lukan, T., Veillet, F., Križnik, M., Coll, A., Mahkovec Povalej, T., Pogačar, K., Stare, K., Chauvin,
L., Chauvin, J.E., and Gruden, K. (2022). CRISPR/Cas9-mediated fine-tuning of miRNA
expression in tetraploid potato. Hortic Res *9*. https://doi.org/10.1093/hr/uhac147.

73.   Lukan, T., Županič, A., Mahkovec Povalej, T., Brunkard, J.O., Kmetič, M., Juteršek, M., Baebler,
Š., and Gruden, K. (2023). Chloroplast redox state changes mark cell-to-cell signaling in the
hypersensitive response. New Phytologist *237*, 548–562. https://doi.org/10.1111/nph.18425.

74.   Lukan, T., Machens, F., Coll, A., Baebler, Š., Messerschmidt, K., and Gruden, K. (2018). Plant X-
tender: An extension of the AssemblX system for the assembly and expression of multigene
constructs in plants. PLoS One *13*. https://doi.org/10.1371/journal.pone.0190526.

75.   Stefanic, P., and Mandic-Mulec, I. (2009). Social interactions and distribution of Bacillus
subtilis pherotypes at microscale. J Bacteriol *191*, 1756–1764.
https://doi.org/10.1128/JB.01290-08.

76.   Tortosa, P., Logsdon, L., Kraigher, B., Itoh, Y., Mandic-Mulec, I., and Dubnau, D. (2001).
Specificity and genetic polymorphism of the Bacillus competence quorum-sensing system. J
Bacteriol *183*, 451–460. https://doi.org/10.1128/JB.183.2.451-460.2001.

77.   Pramanik, R.S., Chandwani, S., and Amaresan, N. (2023). Screening of Endophytes for Plant
Growth-Promoting Metabolites A. Sankaranarayanan, N. Amaresan, and M. K. Dwivedi, eds.
(Springer Protocols Handbooks). https://doi.org/10.1007/978-1-0716-2827-0_19.

78.   Chabot, S., Bécard, G., and Piché, Y. (1992). Life Cycle of Glomus Intraradix in Root Organ
Culture. Mycologia *84*, 315–321. https://doi.org/10.1080/00275514.1992.12026144.

79.   García, J.M., Pozo, M.J., and López-Ráez, J.A. (2020). Histochemical and Molecular
891   Quantification of Arbuscular Mycorrhiza Symbiosis M. Rodríguez-Concepción and R. Welsch,
892   eds. (Humana, New York, NY) https://doi.org/10.1007/978-1-4939-9952-1_22.

893   80.   GIOVANNETTI, M., and MOSSE, B. (1980). An evaluation of techniques for measuring vesicular
894   arbuscular mycorrhizal infection in roots. New Phytologist *84*, 489–500.
895   https://doi.org/10.1111/j.1469-8137.1980.tb04556.x.

896   81.   Petek, M., Rotter, A., Kogovšek, P., Baebler, Š., Mithöfer, A., and Gruden, K. (2014). Potato
897   virus Y infection hinders potato defence response and renders plants more vulnerable to
898   Colorado potato beetle attack. Mol Ecol *23*, 5378–5391. https://doi.org/10.1111/mec.12932.

899   82.   Baebler, Š., Svalina, M., Petek, M., Stare, K., Rotter, A., Pompe-Novak, M., and Gruden, K.
900   (2017). quantGenius: Implementation of a decision support system for qPCR-based gene
901   quantification. BMC Bioinformatics *18*. https://doi.org/10.1186/s12859-017-1688-7.

902   83.   Livak, K.J., and Schmittgen, T.D. (2001). Analysis of relative gene expression data using real-
903   time quantitative PCR and the 2-ΔΔCT method. Methods *25*, 402–408.
904   https://doi.org/10.1006/meth.2001.1262.

905   84.   Morcillo, R.J.L., Navarrete, M.I.T., Bote, J.A.O., Monguio, S.P., and García-Garrido, J.M. (2016).
906   Suppression of allene oxide synthase 3 in potato increases degree of arbuscular mycorrhizal
907   fungal colonization. J Plant Physiol *190*, 15–25. https://doi.org/10.1016/j.jplph.2015.11.003.

908   85.   R Core Team (2023). R: A Language and Environment for Statistical Computing. R Foundation
909   for Statistical Computing, Vienna. https://www.R-project.org/.

910   86.   R Core Team (2021). R: A Language and Environment for Statistical Computing. R Foundation
911   for Statistical Computing, Vienna. https://www.R-project.org/.

912   87.   Wickham, H. (2009). ggplot2 1st ed. (Springer New York). https://doi.org/10.1007/978-0-387-
913   98141-3.

914   88.   Kassambara, A. (2019). rstatix: Pipe-Friendly Framework for Basic Statistical Tests. Preprint,
915   https://doi.org/10.32614/CRAN.package.rstatix.

916   89.   Kassambara, A. (2016). ggpubr: "ggplot2" Based Publication Ready Plots. Preprint,
917   https://doi.org/10.32614/CRAN.package.ggpubr.

918   90.   Krueger, F. (2021). Trimgalore . GitHub repository.
919   https://github.com/FelixKrueger/TrimGalore.

920   91.   Andrews, S. (2010). FastQC: A Quality Control Tool for High Throughput Sequence Data
921   [Online]. Babraham Bioinformatics.
922   http://www.bioinformatics.babraham.ac.uk/projects/fastqc/.

923   92.   Kim, D., Song, L., Breitwieser, F.P., and Salzberg, S.L. (2016). Centrifuge: rapid and sensitive
924   classification of metagenomic sequences. Genome Res *26*, 1721–1729.
925   https://doi.org/10.1101/gr.210641.116.

926   93.   Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M.,
927   and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner. Bioinformatics *29*, 15–21.
928   https://doi.org/10.1093/bioinformatics/bts635.

94. Petek, M., Zagorščak, M., Ramšak, Ž., Sanders, S., Tomaž, Š., Tseng, E., Zouine, M., Coll, A., and
Gruden, K. (2020). Cultivar-specific transcriptome and pan-transcriptome reconstruction of
tetraploid potato. Sci Data *7*. https://doi.org/10.1038/s41597-020-00581-4.

95. Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A.,
Paulovich, A., Pomeroy, S.L., Golub, T.R., Lander, E.S., et al. (2005). Gene set enrichment
analysis: A knowledge-based approach for interpreting genome-wide expression profiles. Proc
Natl Acad Sci U S A *102*, 15545–15550. https://doi.org/10.1073/pnas.0506580102.

96. Rotter, A., Usadel, B., Baebler, S., Stitt, M., and Gruden, K. (2007). Adaptation of the MapMan
ontology to biotic stress responses: application in solanaceous species. Plant Methods *3*.
https://doi.org/10.1186/1746-4811-3-10.

97. Ramšak, Ž., Petek, M., and Baebler, Š. (2021). RNA Sequencing Analyses for Deciphering
Potato Molecular Responses. In Solanum tuberosum, D. Dobnik, K. Gruden, Ž. Ramšak, and A.
Coll, eds. (Humana, New York, NY), pp. 57–94. https://doi.org/10.1007/978-1-0716-1609-3_3.

98. Garrido Rodríguez-Córdoba, M. (2023). biokit: Toolkit for the automatic analysis of omics
data. github. https://github.com/martingarridorc/biokit.

99. Ritchie, M.E., Phipson, B., Wu, D., Hu, Y., Law, C.W., Shi, W., and Smyth, G.K. (2015). Limma
powers differential expression analyses for RNA-sequencing and microarray studies. Nucleic
Acids Res *43*, e47. https://doi.org/10.1093/nar/gkv007.

100. Zagorščak, M., Blejec, A., Ramšak, Ž., Petek, M., Stare, T., and Gruden, K. (2018). DiNAR:
Revealing hidden patterns of plant signalling dynamics using Differential Network Analysis in
R. Plant Methods *14*. https://doi.org/10.1186/s13007-018-0345-0.

101. Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., and Madden, T.L.
(2009). BLAST+: Architecture and applications. BMC Bioinformatics *10*.
https://doi.org/10.1186/1471-2105-10-421.

Supplemental information

Figure S1: Biofilm on potato roots after incubation in *B. subtilis* culture, imaged at different time
points.

Figure S2: Expression profiles of selected genes.

Figure S3: Number of genes, regulated by *B. subtilis* in plants incubated in *B. subtilis* culture of high
density (107 CFU/m^L).

Figure S4: Observation of *B. subtilis* cells in potato tissue after internalization.

Figure S5: B. subtilis cells as observed in liquid culture (A) and potato leaf homogenate (B).

Figure S6: Biofilm formation on potato roots inoculated with *B. subtilis* mutants that have attenuated
surfactin production.

Figure S7: *B. subtilis*-produced secondary metabolites in the medium do not contribute significantly
to the PTI5 and 13-LOX upregulation.

Figure S8: Regulated central signaling modules in potato roots in isogenic and non-kin interactions.

Figure S9: *B. subtilis* root colonization of transgenic potato lines with silenced PTI5 (shPTI5 L2 and L6)
and non-transgenic (NT) potato plant.

Figure S10: Mycorrhizal colonization of potato roots in non-transgenic potato plant (NT) and PTI5
silenced genotype line 2 (shPTI5).

Table S1: Standardised qPCR gene expression data.

Table S2: Results of gene set enrichment analysis.

Table S3: Differential gene expression table for comparisons between plants inoculated with *B.*
subtilis and non-inoculated plants.

Table S4: The amount of biofilm on potato roots.

Table S5: Relative gene expression in potato inoculated with *B. subtilis* mutants that have attenuated
surfactin production.

Table S6: Expression of selected potato genes induced by conditioned medium in which *B. subtilis*
was growing.

Table S7: Plant stress signaling (PSS) prior knowledge network node information with corresponding
RNA-seq differential expression.

Table S8: The amount of biofilm formation on potato roots for non-transgenic (NT) and shPTI5 plants.

Table S9: *B. subtilis* abundance in shPTI5 L6 and non-transgenic (NT) potato plants determined by
qPCR.

Table S10: *B. subtilis* abundance in shPTI5 L2 and non-transgenic (NT) potato plants determined by
qPCR.

Table S11: Abundance of *B. subtilis* PS-216 and QS impaired strain PS-216 *srfA* in Rywal non-
transgenic (NT) potato plants, determined by qPCR.

Table S12: The production of indole acetic acid (IAA) was assessed following the protocol of Pramanik
et al.[77]

Table S13: Selected genes and corresponding primers and probes sequences for expression analyses
with quantitative PCR.

Table S14: Sample information for RNA-seq experiments.