Motor deficits in the McGill-R-Thy1-APP transgenic rat model of Alzheimer's Disease

Kyle M Roddick[1,2]*, Heather M Schellinck[1], Richard E Brown[1]

[1]Department of Psychology and Neuroscience, Dalhousie University, Halifax, Nova Scotia
[2]Department of Psychology, Mount Allison University, Sackville, New Brunswick

*Corresponding author:
Dr. Kyle Roddick
Department of Psychology
Mount Allison University
Sackville, New Brunswick
Canada E4L 1E4

Phone: 902-240-0914
Email: kroddick@mta.ca

Abstract

The McGill-R-Thy1-APP rat is a transgenic model of Alzheimer's Disease (AD) which expresses APP with two mutations found in cases of familial AD, resulting in the development of amyloid pathology and cognitive deficits. Motor deficits are common symptoms of AD, emerging early in the disease, and are correlated with AD neuropathology and cognitive symptoms. This study evaluated heterozygous and homozygous McGill-R-Thy1-APP rats and their wildtype littermates for spontaneous alternation and locomotion in the T and Y mazes, and motor behaviour on an accelerating rotarod. Because rats often jumped off the rotarod, the maximum latency to fall across trials was examined. We found no genotype or sex effects in spontaneous alternation in either maze, nor a significant correlation of spontaneous alternation behaviour between the mazes. Female rats travelled greater distances than male rats in both mazes. While there was no genotype effect in the T maze on distance travelled, in the Y maze the hemizygous rats travelled shorter distances than the wildtype rats, while the homozygous rats travelled greater distances. There was a significant correlation between the distances travelled in each maze. Both heterozygous and homozygous rats performed worse than their wildtype littermates on the rotarod, while heavier rats performed worse than lighter rats, and female rats performed worse than male rats once their differences in weights were accounted for. These findings support the continued use of these rats as a model of AD and highlight the need to consider the possible confounding effect motor impairments have on other behavioural tests.

Key Words: Alzheimer's Disease, McGill-R-Thy1-APP rat, Motor behaviour, Spontaneous alternation, Rotarod

1.0 Introduction

Alzheimer's Disease (AD) is an age-related neurodegenerative disorder characterized by the development of Aβ-plaques and neurofibrillary tangles (1–3). Although most AD research is focused on cognitive deficits, patients with AD also have behavioural (4–6), neuropsychiatric (7,8), sensory (9) and motor dysfunctions (10). Motor behaviour disabilities are common symptoms of AD (11) and higher levels of motor dysfunction are correlated with increased severity of cognitive symptoms (12–14). Severity of motor behaviour impairments differentiates between healthy controls, patients with mild cognitive impairment, and patients with AD (15). Walking gait (14,16), finger dexterity (17–19), and handwriting (20) are all impaired in AD patients, and the degree of motor impairment is associated with the level of neuropathology (21–23). Disturbances in gait have been proposed as a method of differentiating different types of dementia (24,25).

Rodent models play an important role in understanding the neurobiology of AD (26–29) and in the search for new therapeutics (30–32). Many transgenic rodent models have been developed which display the neuropathological hallmarks of AD, as well as the associated cognitive deficits (33). However, it is also important to assess the neuro-behavioural, sensory, and motor functions of these models, particularly as many behavioural tests commonly used to evaluate cognitive function require the use of visual (34) or auditory (35) cues and locomotor behaviour (36) to perform these tests. Accordingly, motor behavioural deficits may confound the results of behavioural studies. Age related motor deficits have been shown in the APPswe/PS1dE9 (37) and 5xFAD mouse models of AD (36,38), while the 3xTg-AD mouse model has enhanced motor performance on the rotarod (39–41).

The McGill-R-Thy1-APP transgenic rat model of AD expresses human amyloid-β

precursor protein with both the Indiana and Swedish mutations (42). Both hemizygous and

homozygous transgenic rats develop intracellular accumulation of amyloid-β, with homozygous

animals also developing extracellular Aβ plaques (42,43), neuroinflammation (44), and

dysregulation of neurotrophins (45). Both hemizygous and homozygous rats show cognitive

deficits as early as 3-months of age in the Morris Water Maze (42,46). Changes in motor

behaviour have also been observed in this rat model. Homozygous rats show motor deficits in a

beam walking test, taking longer to cross the beam and making a greater number of foot slips

during crossings than wildtype controls (47). However, the study found no difference between

homozygous rats and wildtype controls in swim speed during the Morris Water Maze test, nor

in total distance traveled In the Elevated Plus Maze or the Open Field. Other studies, while not

explicitly evaluating motor behaviour, have reported no differences in swimming ability in the

Morris Water Maze in either hemizygous (46,48,49) or homozygous McGill-R-Thy1-APP rats

(50), nor differences in the total distance travelled in the open field in hemizygous (51) or

homozygous rats (52). These studies evaluated either hemizygous or homozygous rats, not

both. The present study is the first to evaluate the motor behaviour of both hemizygous and

homozygous McGill-R-Thy1-APP rats. Female and male transgenic rats, and their wildtype

littermates, were tested in the Y and T mazes for spontaneous alternation at 12 months of age.

In addition to their percent alternation, the total distance traveled was compared between

genotypes. At 13 months of age they were tested on an accelerating rotarod and their latency

to fall was compared. Based on the findings of impaired performance of homozygous rats on a

beam walking task (47), we hypothesised that the homozygous rats would display motor

impairments on the rotarod. The are no previous studies which explicitly examined hemizygous McGill-R-Thy1-APP rats for motor impairment, however, as previous studies using the Morris Water Maze reported no differences in swim speed (46,48,49), we predicted no motor deficits in the hemizygous rats.

2.0 Methods

2.1 Subjects:

McGill-R-Thy1-APP transgenic rats express human amyloid-β precursor protein with both the Indiana and Swedish mutations. Expression is under to control of the murine *Thy*1.2 promoter and they are on a HsdBrl:WH Wistar background (42). Hemizygous rats were obtained from the Cuello Laboratory (McGill University) and bred in our laboratory at Dalhousie University. After weaning, rats were housed with same sex littermates in groups of 2 or 3 in polyethylene cages (45cm x 24cm x 20cm) with woodchip bedding and a polyethylene tube for enrichment. Food (Purina rodent chow #5001) and water were available *ad libitum*. The colony room was under a reversed 12:12 light dark cycle and rats were tested during the dark phase of the light dark cycle. Rats from 4 litters were tested in this experiment. The rats were 12 months of age during the T and Y maze tasks, and 13 months of age during the Rotarod task. The rats were extensively handled prior to being run in these tests. They underwent testing in a neurodevelopmental test battery from PND 1 until PND 20, requiring daily handling, and were tested daily in an operant olfactometer from 3 to 7 months of age, also involving daily handling. Genotype was determined via histological examination of brain tissue for intracellular amyloid-β and amyloid-β plaques following the behavioural studies, which were done blind with respect

to genotype. There were 10 wildtype (WT) rats (5 females and 5 males), 15 hemizygous (HE)

transgenic rats (8 females and 7 males), and 10 homozygous (HO) transgenic rats (2 females

and 8 males; Table 1). All procedures were approved by the Dalhousie University's University

Committee on Laboratory Animals (#14-059), following the guidelines of the Canadian Council

on Animal Care.

Table 1: Number and weights of 12 to 13 month old male and female wildtype (WT), hemizygous (+/-), and heterozygous (+/+) McGill-R-Thy1-APP rats tested in spontaneous alternation and the rotarod.

	Female			Male			Overall		
	WT (N=5)	+/- (N=8)	+/+ (N=2)	WT (N=5)	+/- (N=7)	+/+ (N=8)	WT (N=10)	+/- (N=15)	+/+ (N=10)
Weight (g)									
Mean (SD)	368 (18.2)	365 (43.1)	365 (14.8)	625 (55.8)	617 (61.8)	593 (51.3)	496 (141)	483 (140)	548 (107)
Median [Min, Max]	364 [351, 393]	361 [308, 419]	365 [354, 375]	614 [567, 704]	610 [537, 683]	591 [526, 674]	480 [351, 704]	419 [308, 683]	575 [354, 674]

2.2 T and Y mazes:

2.2.1 Apparatus:

Both the T and Y mazes were constructed of transparent Plexiglas, with black paper

covering on the outside of the walls. The T maze had equal-sized stem and arms (40cm length,

12cm width, 19.5cm height) without any start box. The Y maze was symmetrical with its equal

sized arms (40cm length, 12cm width, 19.5cm height) forming 120° angles. The tops of the

mazes were covered with transparent Plexiglas lids to prevent the rats from climbing out of the

mazes. An overhead camera was used to record the rats' behaviour, and Limelight software

(Actimetrics, Lafayette, IN) was used to track the position of the rats within the mazes.

2.2.2 Procedure:

To test the rats for spontaneous alternation, the continuous (free-running) method was used. Animals were placed inside one of the arms of either of the T or Y maze and allowed to explore for 10 minutes without any reinforcement or punishment. All of the arms were available to enter. There were no training sessions for the animals prior to the experiment; this was the first time that they were exposed to either the T or Y maze. A week after being tested in one maze, the animals were tested in the other maze using the same procedure. In order to control for the effect of familiarity due to the similarity of the mazes, half of the animals were tested in the T maze first and the other half in the Y maze first.

2.2.3 Analysis:

The Limelight software recorded each arm entry that the animal made during the test. An arm entry was counted when the animal's body (the middle-point between its nose and the base of the tail) entered an arm. An alternation triplet was counted when the animal entered three different arms in succession. These triplet sets could overlap and still be counted as alternations. For example, entries of ACBABACBAB would be counted as: ACB, CBA, BAC, ACB, and CBA; therefore 5 alternations in total (53). The percentage of alternations was calculated by dividing the number of alternations by the number of total entries minus 2, all multiply by 100 (Equation 1).

Equation 1: Calculation of percent alternation.

$$\text{percent alternation} = \frac{\text{number of alternations}}{\text{total number of arm entries} - 2} \times 100$$

2.3 Rotarod:

2.3.1 Apparatus:

Rats were tested on an AccuRotor accelerating Rotarod (Accuscan Instruments Inc. Columbus Ohio) which had an acrylic rod (7cm in diameter), separated into four 11cm sections by 11cm high Plexiglass dividers. The Rotarod accelerated from 0 to 48 rotations per minute over the course of 360s. Latency to fall was recorded by sensors in holding chambers located 36cm below the rod. The Rotarod was housed in a 1.2m x 2.6m room lit by a 60W red light.

2.3.2 Procedure:

Rats were tested over 5 days, receiving 6 trials each day. During each trial, rats were tested until they fell from the rod or the maximum trial length of 360s was reached. Rats were tested two at a time so that they were not in adjacent sections. Rats were weighted each day before testing.

2.3.3 Statistical Analyses:

All statistical analyses were performed with the statistical program R (54). Because some rats jumped from the Rotarod into the holding chamber rather than fell, the maximum latency to fall that each rat achieved across all trials was analyzed. The mean weights of the rats during Rotarod testing were also analyzed. Linear regression models were compared using Akaike's Information Criterion (AIC) which compares the statistical models according to their complexity and how well they fit the data (55,56). Confidence intervals (95%) were calculated for all analyses.

2.4 Immunohistochemistry:

Rats were euthanized with an overdose of sodium pentobarbital, and perfused through the left ventricle, first with saline (0.9% NaCl in dH2O, pH 7.4), followed by 4% paraformaldehyde in 0.1M phosphate buffer (pH 7.4). Brains were removed and stored in 30% sucrose in 0.1M phosphate buffer at 4°C. The brains were frozen and, using a cryostat, cut into 40μm coronal sections. The sections were then collected and stored in 0.1M phosphate buffer at 4°C.

Four sections from each rat underwent immunolabeling. Sections were permeabilized with PBS with .02% Triton X-100 (PBS-T) for 20min, followed by 20min in PBS-T with 0.3% H_2O_2 to quench endogenous peroxidases. After 3x 5min PBS-T washes, the sections were blocked with 30min in 5% normal goat serum, followed by overnight incubation in the primary antibody, goat anti-Ab antibody McSA1 1:500 (MédiMabs; MM-0015-P, Montreal, QC, Canada). Sections were again washed (3x 5min PBS-T) followed by the secondary antibody, biotinylated goat anti-mouse IgG 1:500 (H&L Biotin; ab6788) for 1 hour, and another wash (3x 5min PBS-T). Sections were processed with a Vectastain Elite ABC-peroxidase kit (Vector Laboratories PK-6100) for 1 hour according to manufacturer specifications, followed by another wash (3x 5min PBS-T), 15min in 0.6% 3,3'-diaminobenzidine tetrahydrochloride, 4 to 6min with 0.01% H_2O_2, and a final wash (3x 5min PBS-T). Sections were mounted on slides the next day, allowed to air dry, dehydrated in graded EtOH (2min each in 50%, 70%, 90%, 95%, 100%, and 100%), cleared with Histo-Clear (2x 2min), and cover slipped with Permount.

Pictomicrographs were captured using a microscope-attached digital camera system (Leica, DM E), and used to genotype the rats based on the presence of intracellular and extracellular Aβ.

4.0 Results:

4.1 T and Y maze:

4.1.1 Quality checks:

To ensure the quality of the data, checks were performed to confirm an unbiased assessment of spontaneous alternation (57). A significant correlation between percent alternation and either total number of arm entries or total distance travelled could indicate that hyperdynamic locomotion had an impact on the results. Pearson correlations were run between the percent alternation, and total number of arm entries and total distance travelled, for both mazes. Percent alternation was not correlated with total number of arm entries for either the T maze ($r = -0.07$, $p = 0.69$), or the Y maze ($r = 0.029$, $p = 0.87$). Nor was percent alternation correlated with total distance travelled for either the T maze ($r = -0.042$, $p = 0.81$), or the Y maze ($r = 0.066$, $p = 0.7$). Therefore, there is no indication of hyperdynamic locomotion impacting the results.

ANOVAs were run on both mazes to examine if any of the arms were entered more times than the others, an indicator of potential environmental cues in the maze leading to biased behaviour (57). There were no significant differences in the number of times each arm was entered for either the T maze ($F_{(2,102)} = 0.745$, $p = 0.477$, $\eta_G^2 = 0.014$) or the Y maze

$(F_{(2,102)}$ = 0.28, p = 0.756, $\eta_G{}^2$ = 0.005). Therefore, there is no indication of environmental

cues biasing the behaviour of the rats.

4.1.2 Spontaneous alternation:

Linear regression models of percent alternation in the T and Y mazes were compared

using AIC to select the best model for each maze. For the T maze, the null model best fit the

data (Supplemental Table 1) indicating no significant effect of genotype or sex (Figure 1A).

Figure 1 Mean (± CI[95]) percent alternation in the A) T maze and the B) Y maze. There were no significant effects of genotype nor sex on percent alternation in either maze.

For the Y maze, the model with an effect of genotype best fit the model, however, this

model did not differ significantly from the null model (F = 2.5, p = 0.096; Supplemental Table 2)

again indicating no effects of genotype or sex (Figure 1B).

A Pearson correlation coefficient between the percent alternation by all the rats in the

two mazes was not significant (r = -0.24, p = 0.17; Supplemental Figure 1A).

4.1.3 Distance travelled:

Linear regression models of the distances travelled in the T and Y mazes were compared

using AIC to select the best model for each maze. For the T maze the model with just an effect

of sex was the best model (Supplemental Table 3), and this model differed significantly from

the null model (F = 11.00, p = 0.002). This model indicates that the male rats travelled less

distance (33.8 ± 8.11 m) than the female rats (43.6 ± 9.18 m) in the T maze (CI[95] of the

difference: -15.8 - -3.9 m; Figure 2A).

Figure 2: Mean (± CI^{95}) total distance travelled in A) the T maze and B) the Y maze. The female rats traveled greater distances than the male rats in both mazes. In the Y maze, the hemizygous (+/-) rats travelled less than the wildtype (WT) rats, and the homozygous (+/+) rats travelled more than the wildtype rats.

For the Y maze the model with effects of genotype and sex was the best model (Supplemental Table 4), and this model differed significantly from the null model (F = 5.2, p = 0.005). Again, the male rats (34.4 ± 8.3 m) travelled less distance than the female rats (43.9 ± 8.54 m; CI^{95} of the difference: -17 - -5.1 m; Figure 2B). This model also indicates that the hemizygous (+/-) rats travelled less than the wildtype (WT) rats (CI^{95} of the difference: -10.2 - 3.4), while the homozygous (+/+) rats travelled further than the WT rats (CI^{95} of the difference: -4.18 - 11).

4.2 Rotarod:

The time to fall from the rotarod for each rat on each of six trials per day for five days, with LOESS (locally estimated scatterplot smoothing; (58) trend lines and 95% confidence intervals (CI^{95}) for each genotype, is shown in Figure 3. Early in the study, the rats began to jump from the rotarod, resulting in frequent very low latencies and decreasing latencies to fall across trials. With the exception of one hemizygous female which achieved perfect performance by the final day of testing (Supplemental Figure 3A) rats failed to show increases in latency to fall across days as would be expected for motor learning on the rotarod (38). In order to best evaluate locomotor ability, the effects of sex, genotype, and body weight on the maximum latency to fall that each rat achieved across all trials were analyzed.

Figure 3: Latency(s) to fall from the Rotarod across the six trials on each of five days of testing. Points for each individual trial and loess lines, with CI^{95} for each genotype, are shown. Latencies for all genotypes began to fall in the second half of the first day of testing, and continued to decrease each day. This was because the rats jumped off the Rotarod rather than fell off.

Linear regression models of the mean weights of the rats (see Table 1) across testing

days were compared using AIC (Supplemental Table 5). The best model of the data (AIC = 372.4,

weight = 0.75) included no effect of genotype on body weight, but did include an effect of sex

with males weighing more than females (CI^{95} 209 – 274 g). This model differed significantly

from the null model (F = 228, p < 0.0001). Linear regression models of the maximum latency to

fall for each rat were compared using AIC (Supplemental Table 6). The additive model with

effects of mean body weight, sex, and genotype was found to best model the data (AIC =

378.13, weight = 0.286), and differed significantly from the null model (F = 8.3, p < 0.001). This

model indicated that body weight was negatively associated with the maximum latency to fall

with lighter rats having longer latencies to fall than heavier rats (CI^{95} -1.091 – -0.332 s/g) This

means that the model predicts a decrease in the latency to fall of approximately .7 seconds for

each gram of body weight (s/g). The model also shows males having longer latencies to fall

than females (CI^{95} 13.71 – 214.09 s), and both hemizygous (CI^{95} -69.90 – 11.61 s) and

homozygous rats (CI^{95} -98.04 – -4.70 s) having shorter latencies to fall than wildtype rats.

Supplemental Figure 2 shows raincloud plots for the genotype differences in maximum latency

to fall over the 30 trials, while Figure 4 shows that the lighter rats stayed on the rotarod longer

than heavier rats for both females (r = -0.60, p = 0.017) and males (r = -0.49, p = 0.028).

Figure 4: Maximum latency to fall (s) from the Rotarod for individual female and male rats according to their body weight (g). Linear regression lines (± CI^{95}) of the effect of weight on maximum latency to fall for females and males are shown separately because of the sex differences in body weight.

4.3 Cross test correlations:

Pairwise Pearson correlations were run between the three measures of locomotor

activity, distance travelled in the T and Y mazes and the maximum latency to fall from the

Rotarod. While the distance travelled in the T and Y mazes were positively correlated (r = 0.65,

$p < 0.0001$), the maximum latency to fall from the Rotarod was not significantly correlated with distance travelled in either the T (r = 0.3, p = 0.08) or Y maze (r = 0.061, p = 0.73; Figure 5).

Figure 5: Pearson correlation matrix of locomotor measures. The plots in the central diagonal show histograms of the data for each variable overlayed with density curves. The plots in the lower half scatterplots of the two variables with loesses lines of best fit, while the numbers in the upper half indicate the correlation coefficient with significance level (square for p < 0.1, *** for p < 0.001).

5.0 Discussion

This study found no evidence of differences in either the homozygous or hemizygous McGill-R-Thy1-APP transgenic rats on spontaneous alternation compared to wildtype rats. This is in contrast to Galeano et al. (46), who found spontaneous alternation deficits in hemizygous McGill-R-Thy1-APP rats at 6 and 12 months of age, but not at 3 months. Other rat models of AD have been tested for spontaneous alternation with mixed results. TgF344-AD transgenic rats, with human APP containing the Swedish mutation and Δ exon 9 mutant human presenilin-1 (59), showed impaired spontaneous alternation at 18 months of age (60). Aβ-injected rats had lower rates of spontaneous alternation than controls in some studies (61–65), but not in others (66). However, aluminum chloride induced AD model rats (67,68) and a streptozotocin induced AD rat model (69) had lower rates of spontaneous alternation than controls.

The present study found no significant correlation between spontaneous alternation in the T and Y mazes, but there was a significant, positive correlation between distance traveled in the two mazes. In both mazes the male rats travelled shorter distances than the female rats, but there was no clear genotype pattern between the mazes. Thus, there is little evidence of a motor impairment in either the hemizygous or homozygous McGill-R-Thy1-APP rats in the T or Y mazes.

However, there was clear evidence that homozygous McGill-R-Thy1-APP rats showed impaired motor behaviour on the rotarod compared to their wildtype littermates. The evidence also strongly suggested an impairment in motor behaviour on the rotarod in the hemizygous transgenic rats. The finding of impaired motor behaviour in the homozygous rats is in agreement with the results of Petrasek et al. (47), who found homozygous rats to be impaired on a beam walking task. While some previous studies have tested hemizygous rats in the Morris Water Maze and reported no differences in swimming speed (46,48,49), no prior studies have explicitly evaluated the motor behaviour of hemizygous rats. It has been argued that the hemizygous McGill-R-Thy1-APP rats are a good model of the early stages of AD (46) and this argument is supported by the present evidence of motor impairments on the Rotarod in the hemizygous rats. As motor dysfunction is a well established symptom of early AD (11), animal models of early AD should also be expected to display motor dysfunction.

We analyzed the single best trial for each rat, a measure which has been previously used to assess motor behaviour (70), because the rats frequently jumped off of the rod during testing rather than falling off. The data support this observation, with only one rat showing a clear improvement over the test (Supplemental Figure 3A). While there were 104 trials with a latency of less than 10 seconds to fall, including 23 trials with a latency of less than 2 seconds. All three genotypes began to show a decline in performance during the first day of testing (Figure 3), indicating that jumping from the rotarod started to occur very early during testing. This jumping behaviour means that in many cases the latency to fall is not an accurate measure of how long the rat was able to stay on the rod. By taking the longest latency for each individual rat we were able to get a better measure of the capacity of the rat to stay on the rod. While

54% of the rats had their maximum latency trial on the first day of testing, no rats had their

maximum latency on the first trial. It is possible that the advanced age of the rats when tested

(13 months) made the repeated trials of the task more tiring for them, and that this lead to the

jumping behaviour. Other studies have reported instances of passive rotation on the Rotarod

(71,72), where the rats grip the rod and rotate with it, but no instances of passive rotation were

observed in the present study. We found no other published data on rats jumping from the

rotarod, though it has been observed in mice (73), so there are no established methods for

dealing with this type of animal "misbehaviour" (74).

The method of analyzing the maximum latency to fall for each rat is not without fault.

The worst performing rat, a hemizygous male, had a maximum latency to fall of just 23.1 sec,

and three trials with a latency less than 1 second (Supplemental Figure 3B). While this rat was

the second heaviest rat tested, it is unlikely that the rotarod was able to measure the true

motor capacity of this animal. However, we found a clear effect of weight in this study, with

lighter rats performing better on the rotarod than heavier rats, which is in agreement with what

we have found when testing mouse models of AD (36–38,40,41). This suggests that using the

maximum latency each rat achieved is accurately measuring the effects of genotype and sex on

of motor function.

When corrected for body weight, the model of the latencies to fall indicated that males

stayed on the rod longer than females. This is in contrast with the uncorrected results of the

experiment which found that female rats had a greater maximum latency to fall (219 ± 54 s)

than males (153 ± 62 s). This apparent contradiction is due to the combination of the effect of

weight on the latency to fall, and the difference in weights between female and male rats. Our

model found that the greater the weight of the rat, the shorter the time it was able to stay on the rod. We also found a substantial difference in the weights of female and male rats (Table 1, Figure 4). There was no overlap in weights between the two sexes, with the heaviest female (417g) weighing 100g less than the lightest male (517g). So while the males did not stay on the rod as long as the females, they performed better than the females once their weight was taken into account. Studies of sex differences in rotarod performance in rat and mouse models of AD found that female rats performed better than male rats (75), though the effect of body weight was not accounted for in that study. When rats were matched by weight, female rats also performed better than males (76). We previously found no effect of sex on rotarod performance in 3xTg-AD mice at either 6 (41), or 16 months of age (40) once weight was taken into account. Female 5xFAD mice performed better than males (36), however this study did not account for differences in their body weight in the analysis of sex differences.

Other rat models of AD have been tested on the rotarod with mixed results. Some studies using a model of AD induced by Aβ injection found no deficits in latency to fall from the Rotarod (77,78), while other studies using an Aβ-induced model did find deficits in Rotarod performance (79–81). Deficits in motor performance on the rotarod have also been found in models of AD induced by aluminum chloride exposure (82,83), trimethyltin chloride (84), and scopolamine (85). Meanwhile, a study using a streptozotocin induced AD model (86), and a study using the App^{NL-G-F} transgenic rat (87), with three familiar App mutations (Swedish, Beyreuther/Iberian, and Arctic) and a humanized Aβ sequence knocked into the rat App gene, found no differences on the rotarod at five months of age. These findings highlight the need for studies to consider the possible impact motor deficits may have when performing cognitive

tests on AD models. Many tests, such as the Morris Water Maze, the T maze, and the Y maze, require the animal to move around the testing area. If they have motor impairments, a lower performance could be due to the motor impairments rather than a cognitive deficit. Conversely, if an experimental intervention restores cognitive function, but is not effective for motor function, these tests may falsely find a negative result if motor function impacts the animal's ability to complete the test.

Different genotype effects were found in distance travelled in the T and Y mazes versus the maximum latency to fall from the rotarod. There was also no significant correlation between the maximum latency to fall from the rotarod and the distance travelled in either maze. This is likely due to the distance travelled in the T and Y mazes being a measure of voluntary movement, while the maximum latency to fall from the rotarod is a measure of forced movement. Lambert et al. (88) found that performance of rats on a forced movement treadmill task did not predict their voluntary movement in a home-cage running wheel. Additionally, a study examining the motor performance of seven inbred strains of mice found no correlation between the strains performance on a forced treadmill task and voluntary wheel running (89).

Although the rotarod is a commonly used apparatus for measuring motor learning and coordination in mice and rats (73), there are a number of parameters which can affect the performance of rodents on the rotarod. The use of an accelerating vs fixed speed rotarod, and their rate of acceleration or speed, respectively, will influence the sensitivity of the test (90). The diameter of the rod (if it is small enough for the rodent to grab hold of and passively rotate) can also effect performance (91). Differences in performance can also be seen between

rotarods from different manufacturers (92), which could be due to differences in the rate of acceleration of the rod (93). The training schedule can also effect performance, with rodents showing a learning curve over trials (36,40,91).

Few experiments test aged rodents on the rotarod, and since AD is a disease of aging, it is important to examine the age-related decline in motor behaviour (36,40). Although mice gain weight with age, the increase is from 26-34 g at 6 months of age (41) up to 26-46 g at 16 months of age (40). This is a very small age-related increase in body weight compared rats, which increase from 179-349 g in young rats (94) to the 308-704 g found in the present study. As with mice (41), heavier rats fell faster from the rotarod than lighter rats (95).

Two behaviours which have resulted in animals being excluded from rotarod studies involve passive rotations and failure to learn over trials. Rodents can cling to the rod and show passive rotations (93,96) and Shan et al (97) excluded 10 of 58 C57BL/6 mice due to "bad performance in baseline testing" on the rotarod. Zieglowski et al. (98) excluded 14 of 30 Wistar Han rats due to "lack of learning ability" and Cartmell et al. (94) excluded 25 of 60 Sprague Dawley rats for failing to show improved over 10 trials on the rotarod. These problems came as a surprise to us as we had not had mice show passive rotations nor have we had to remove mice for failing to learn over repeated trials on the rotarod (36,40,41). We have even recommended the rotarod as one of the most reliable tests for measuring motor function in mice (36). The standard rotarod is not designed for these large rats, which easily jump down from the rod without waiting to fall. Although we modified the standard procedure by testing only two rats at a time, they quickly learned to jump off the rod and other modifications are necessary (73,93,94). Some rotarod systems are capable of delivering a mild electrical shock to

the rodents when they fall off the rod and land in the holding chambers below (95), it is possible that such a system would discourage such jumping behaviour, leading to a more typical increase in latency to fall across trials and more reliable measure of age-related motor dysfunction.

The findings of this study, particularly with regards to the effects of sex in the homozygous rats, should be interpreted with some caution due to the combination of the jumping off behaviour and the imbalanced sex ratio of the homozygous rats (2 females and 8 males). The jumping off behaviour prevents the examination of motor learning which is typically assessed in studies using the rotarod, and while the use of the maximum latency to fall as the main dependent measure appears to have been an effective method of assessing the motor ability of these rats, as shown by the detection of weight effects similar to those found in other studies, it remains a cruder measure. The analysis of the maximum latency to fall from the rotarod found main effects of genotype and sex, but no significant interaction between these two factors. However, due to the imbalanced sex ratio among the homozygous rats this lack of an interaction should be interpreted with particular caution.

In summary, this study found evidence of motor deficits in both the homozygous and hemizygous transgenic McGill-R-Thy1-APP rat models of AD. While Petrasek et al. (47) also found impairment in the homozygous rats, this is the first study to find such deficits in the hemizygous rats. These findings support the use of the McGill-R-Thy1-APP as a model of AD, though it is important to consider the potential effects of motor impairments when assessing other behaviours in this, or any other rodent AD model.

Data Availability

The data set used in this study are available on the Dalhousie University Dataverse at

Acknowledgments

This work was supported by a Discovery Grant from the Natural Sciences and Engineering

Research Council of Canada (http://nserc-crsng.gc.ca) (RG7441) to R.E.B. Funders did not play

any role in the study design, data collection and analysis, decision to publish, or preparation of

the manuscript. Thanks to Amirreza Shahisavandi for assistance with behavioural testing and

Paige Northrup for assistance with the immunohistochemistry.

CRediT Author contributions:

KMR: Conceptualization; Data curation; Formal analysis; Investigation; Methodology;

Visualization; Writing - original draft; Writing - review & editing. HMS: Conceptualization;

Project administration; Methodology; Resources; Supervision. REB: Conceptualization; Funding

acquisition; Project administration; Resources; Supervision; Writing - review & editing.

Declarations

Ethics statement

All procedures were approved by the Dalhousie University's University Committee on

Laboratory Animals (#14-059), following the guidelines of the Canadian Council on Animal Care.

Declaration of Interests

The authors have no competing interests to declare.

References

1. Duyckaerts C, Delatour B, Potier MC. Classification and basic pathology of Alzheimer disease. Acta Neuropathol (Berl). 2009 Jul;118(1):5–36.

2. Lane CA, Hardy J, Schott JM. Alzheimer's disease. Eur J Neurol. 2018;25(1):59–70.

3. Scheltens P, De Strooper B, Kivipelto M, Holstege H, Chételat G, Teunissen CE, et al. Alzheimer's disease. The Lancet. 2021 Apr 24;397(10284):1577–90.

4. El Haj M, Roche J, Jardri R, Kapogiannis D, Gallouj K, Antoine P. Clinical and neurocognitive aspects of hallucinations in Alzheimer's disease. Neurosci Biobehav Rev. 2017 Dec 1;83:713–20.

5. Victoroff J, Lin F V., Coburn KL, Shillcutt SD, Voon V, Ducharme S. Noncognitive behavioral changes associated with Alzheimer's disease: Implications of neuroimaging findings. J Neuropsychiatry Clin Neurosci. 2018;30(1):14–21.

6. Zvěřová M. Clinical aspects of Alzheimer's disease. Clin Biochem. 2019 Oct 1;72:3–6.

7. Banning LCP, Ramakers IHGB, Köhler S, Bron EE, Verhey FRJ, de Deyn PP, et al. The association between biomarkers and neuropsychiatric symptoms across the Alzheimer's disease spectrum. Am J Geriatr Psychiatry. 2020 Jul 1;28(7):735–44.

8. Lyketsos CG, Carrillo MC, Ryan JM, Khachaturian AS, Trzepacz P, Amatniek J, et al. Neuropsychiatric symptoms in Alzheimer's disease. Alzheimers Dement. 2011;7(5):532–9.

9. Murphy C. Olfactory and other sensory impairments in Alzheimer disease. Nat Rev Neurol. 2019;15(1):11–24.

10. Albers MW, Gilmore GC, Kaye J, Murphy C, Wingfield A, Bennett DA, et al. At the interface of sensory and motor dysfunctions and Alzheimer's disease. Alzheimers Dement. 2015;11(1):70–98.

11. Buchman AS, Bennett DA. Loss of motor function in preclinical Alzheimer's disease. Expert Rev Neurother. 2011 May 9;11(5):665–76.

12. Aggarwal NT, Wilson RS, Beck TL, Bienias JL, Bennett DA. Motor dysfunction in mild cognitive impairment and the risk of incident Alzheimer disease. Arch Neurol. 2006;63(12):1763–9.

13. Tian Q, Resnick SM, Studenski SA, Ferrucci L. Lap time variability from a 400-m walk is associated with future mild cognitive impairment and Alzheimer's disease. J Am Med Dir Assoc. 2019 Dec 1;20(12):1535-1539.e3.

14. Tian Q, Resnick SM, Mielke MM, Yaffe K, Launer LJ, Jonsson PV, et al. Association of dual decline in memory and gait speed with risk for dementia among adults older than 60 years: a multicohort individual-level meta-analysis. JAMA Netw Open. 2020 Feb 21;3(2):e1921636.

15. Ansai JH, Andrade LP de, Rossi PG, Nakagawa TH, Vale FAC, Rebelatto JR. Differences in timed up and go subtasks between older people with mild cognitive impairment and mild Alzheimer's disease. Motor Control. 2019 Jan 1;23(1):1–12.

16. Buracchio T, Dodge HH, Howieson D, Wasserman D, Kaye J. The trajectory of gait speed preceding mild cognitive impairment. Arch Neurol. 2010;67(8):980–6.

17. Bologna M, Guerra A, Colella D, Cioffi E, Paparella G, Di Vita A, et al. Bradykinesia in Alzheimer's disease and its neurophysiological substrates. Clin Neurophysiol. 2020 Apr;131(4):850–8.

18. Mollica MA, Tort-Merino A, Navarra J, Fernández-Prieto I, Valech N, Olives J, et al. Early detection of subtle motor dysfunction in cognitively normal subjects with amyloid-β positivity. Cortex. 2019;121:117–24.

19. Suzumura S, Osawa A, Maeda N, Sano Y, Kandori A, Mizuguchi T, et al. Differences among patients with Alzheimer's disease, older adults with mild cognitive impairment and healthy older adults in finger dexterity. Geriatr Gerontol Int. 2018;18(6):907–14.

20. Yu NY, Chang SH. Characterization of the fine motor problems in patients with cognitive dysfunction – A computerized handwriting analysis. Hum Mov Sci. 2019 Jun 1;65:71–9.

21. Koychev I, Galna B, Zetterberg H, Lawson J, Zamboni G, Ridha BH, et al. Aβ 42/Aβ 40 and Aβ 42/Aβ 38 ratios are associated with measures of gait variability and activities of daily living in mild Alzheimer's disease: a pilot study. J Alzheimers Dis. 2018 Jan 1;65(4):1377–83.

22. Nadkarni NK, Perera S, Snitz BE, Mathis CA, Price J, Williamson JD, et al. Association of brain amyloid-β with slow gait in elderly individuals without dementia influence of cognition and apolipoprotein e ε4 genotype. JAMA Neurol. 2017;74(1):82–90.

23. Schirinzi T, Di Lorenzo F, Sancesario GM, Di Lazzaro G, Ponzo V, Pisani A, et al. Amyloid-mediated cholinergic dysfunction in motor impairment related to Alzheimer's disease. J Alzheimers Dis. 2018 Jan 1;64(2):525–32.

24. Mc Ardle R, Galna B, Donaghy P, Thomas A, Rochester L. Do Alzheimer's and Lewy body disease have discrete pathological signatures of gait? Alzheimers Dement. 2019;15(10):1367–77.

25. Mc Ardle R, Del Din S, Galna B, Thomas A, Rochester L. Differentiating dementia disease subtypes with gait analysis: feasibility of wearable sensors? Gait Posture. 2020 Feb 1;76:372–6.

26. Do Carmo S, Cuello AC. Modeling Alzheimer's disease in transgenic rats. Mol Neurodegener. 2013;8(1):37.

27. Drummond E, Wisniewski T. Alzheimer's disease: experimental models and reality. Acta Neuropathol (Berl). 2017;133(2):155–75.

28. Götz J, Bodea LG, Goedert M. Rodent models for Alzheimer disease. Nat Rev Neurosci. 2018;19(10):583–98.

29. Mckean NE, Handley RR, Snell RG. A Review of the Current Mammalian Models of Alzheimer's Disease and Challenges That Need to Be Overcome. Int J Mol Sci. 2021 Dec 6;22(23):13168.

30. Cuello AC, Hall H, Do Carmo S. Experimental pharmacology in transgenic rodent models of Alzheimer's disease. Front Pharmacol. 2019 Mar 4;10:189.

31. Puzzo D, Gulisano W, Palmeri A, Arancio O. Rodent models for Alzheimer's disease drug discovery. Expert Opin Drug Discov. 2015 Jul 3;10(7):703–11.

32. Snyder HM, Shineman DW, Friedman LG, Hendrix JA, Khachaturian A, Le Guillou I, et al. Guidelines to improve animal study design and reproducibility for Alzheimer's disease and related dementias: For funders and researchers. Alzheimers Dement. 2016;12(11):1177–85.

33. Pádua MS, Guil-Guerrero JL, Prates JAM, Lopes PA. Insights on the use of transgenic mice models in Alzheimer's disease research. Int J Mol Sci. 2024 Jan;25(5):2805.

34. Brown RE, Wong AA. The influence of visual ability on learning and memory performance in 13 strains of mice. Learn Mem. 2007;14(3):134–44.

35. O'Leary TP, Shin S, Fertan E, Dingle RN, Almuklass A, Gunn RK, et al. Reduced acoustic startle response and peripheral hearing loss in the 5xFAD mouse model of Alzheimer's disease. Genes Brain Behav. 2017 Jun;16(5):554–63.

36. O'Leary TP, Mantolino HM, Stover KR, Brown RE. Age-related deterioration of motor function in male and female 5xFAD mice from 3 to 16 months of age. Genes Brain Behav. 2020;19(3):e12538.

37. O'Leary TP, Hussin AT, Gunn RK, Brown RE. Locomotor activity, emotionality, sensori-motor gating, learning and memory in the APPswe/PS1dE9 mouse model of Alzheimer's disease. Brain Res Bull. 2018 Jun 1;140:347–54.

38. O'Leary TP, Robertson A, Chipman PH, Rafuse VF, Brown RE. Motor function deficits in the 12 month-old female 5xFAD mouse model of Alzheimer's disease. Behav Brain Res. 2018 Jan 30;337:256–63.

39. Filali M, Lalonde R, Theriault P, Julien C, Calon F, Planel E. Cognitive and non-cognitive behaviors in the triple transgenic mouse model of Alzheimer's disease expressing mutated APP, PS1, and Mapt (3xTg-AD). Behav Brain Res. 2012;234(2):334–42.

40. Garvock-de Montbrun T, Fertan E, Stover K, Brown RE. Motor deficits in 16-month-old male and female 3xTg-AD mice. Behav Brain Res. 2019 Jan 1;356:305–13.

41. Stover KR, Campbell MA, Van Winssen CM, Brown RE. Analysis of motor function in 6-month-old male and female 3xTg-AD mice. Behav Brain Res. 2015;281:16–23.

42. Leon WC, Canneva F, Partridge V, Allard S, Ferretti MT, DeWilde A, et al. A novel transgenic rat model with a full alzheimer's-like amyloid pathology displays pre-plaque intracellular amyloid-β-associated cognitive impairment. J Alzheimers Dis. 2010 Jan 1;20(1):113–26.

43. Iulita MF, Allard S, Richter L, Munter LM, Ducatenzeiler A, Weise C, et al. Intracellular Aβ pathology and early cognitive impairments in a transgenic rat overexpressing human amyloid precursor protein: A multidimensional study. Acta Neuropathol Commun. 2014;2(1):1–17.

44. Hanzel CE, Pichet-Binette A, Pimentel LSB, Iulita MF, Allard S, Ducatenzeiler A, et al. Neuronal driven pre-plaque inflammation in a transgenic rat model of Alzheimer's disease. Neurobiol Aging. 2014;35(10):2249–62.

45. Iulita MF, Bistué Millón MB, Pentz R, Aguilar LF, Do Carmo S, Allard S, et al. Differential deregulation of NGF and BDNF neurotrophins in a transgenic rat model of Alzheimer's disease. Neurobiol Dis. 2017 Dec 1;108:307–23.

46. Galeano P, Martino Adami PV, Do Carmo S, Blanco E, Rotondaro C, Capani F, et al. Longitudinal analysis of the behavioral phenotype in a novel transgenic rat model of early stages of Alzheimer's disease. Front Behav Neurosci. 2014;8:321.

47. Petrasek T, Vojtechova I, Lobellova V, Popelikova A, Janikova M, Brozka H, et al. The McGill transgenic rat model of alzheimer's disease displays cognitive and motor impairments, changes in anxiety and social behavior, and altered circadian activity. Front Aging Neurosci. 2018;10:250.

48. Galeano P, Leal MC, Ferrari CC, Dalmasso MC, Martino Adami PV, Farías MI, et al. Chronic hippocampal expression of notch intracellular domain induces vascular thickening, reduces glucose availability, and exacerbates spatial memory deficits in a rat model of early Alzheimer. Mol Neurobiol. 2018 Nov 1;55(11):8637–50.

49. Martino Adami PV, Galeano P, Wallinger ML, Quijano C, Rabossi A, Pagano ES, et al. Worsening of memory deficit induced by energy-dense diet in a rat model of early-Alzheimer's disease is associated to neurotoxic Aβ species and independent of neuroinflammation. Biochim Biophys Acta BBA - Mol Basis Dis. 2017 Mar 1;1863(3):731–43.

50. Hall H, Iulita MF, Gubert P, Flores Aguilar L, Ducatenzeiler A, Fisher A, et al. AF710B, an M1/sigma-1 receptor agonist with long-lasting disease-modifying properties in a transgenic rat model of Alzheimer's disease. Alzheimers Dement. 2018 Jun;14(6):811–23.

51. Habif M, Do Carmo S, Báez MV, Colettis NC, Cercato MC, Salas DA, et al. Early long-term memory impairment and changes in the expression of synaptic plasticity-associated genes, in the McGill-R-thy1-APP rat model of Alzheimer's-like brain amyloidosis. Front Aging Neurosci. 2021;12:585873.

52. Orciani C, Do Carmo S, Foret MK, Hall H, Bonomo Q, Lavagna A, et al. Early treatment with an M1 and sigma-1 receptor agonist prevents cognitive decline in a transgenic rat model displaying Alzheimer-like amyloid pathology. Neurobiol Aging. 2023 Dec 1;132:220–32.

53. Hughes RN. The value of spontaneous alternation behavior (SAB) as a test of retention in pharmacological investigations of memory. Neurosci Biobehav Rev. 2004 Sep 1;28(5):497–505.

54. R Core Team. R: A Language and Environment for Statistical Computing. Vienna, Austria: R Foundation for Statistical Computing; 2019.

55. Akaike H. A new look at the statistical model identification. IEEE Trans Autom Control. 1974 Dec;19(6):716–23.

56. Burnham KP, Anderson DR. Multimodel Inference. Sociol Methods Res. 2004 Nov 30;33(2):261–304.

57. Miedel CJ, Patton JM, Miedel AN, Miedel ES, Levenson JM. Assessment of spontaneous alternation, novel object recognition and limb clasping in transgenic mouse models of amyloid-β and tau neuropathology. JoVE J Vis Exp. 2017 May 28;(123):e55523.

58. Cleveland WS. Robust locally weighted regression and smoothing scatterplots. J Am Stat Assoc. 1979;74(368):829–36.

59. Cohen RM, Rezai-Zadeh K, Weitz TM, Rentsendorj A, Gate D, Spivak I, et al. A transgenic Alzheimer rat with plaques, tau pathology, behavioral impairment, oligomeric Aβ, and frank neuronal loss. J Neurosci. 2013 Apr 10;33(15):6245–56.

60. Yang L, Wu C, Parker E, Li Y, Dong Y, Tucker L, et al. Non-invasive photobiomodulation treatment in an Alzheimer Disease-like transgenic rat model. Theranostics. 2022 Feb 14;12(5):2205–31.

61. Bagheri M, Joghataei MT, Mohseni S, Roghani M. Genistein ameliorates learning and memory deficits in amyloid β(1–40) rat model of Alzheimer's disease. Neurobiol Learn Mem. 2011 Mar 1;95(3):270–6.

62. Cioanca O, Hritcu L, Mihasan M, Hancianu M. Cognitive-enhancing and antioxidant activities of inhaled coriander volatile oil in amyloid β(1–42) rat model of Alzheimer's disease. Physiol Behav. 2013 Aug 15;120:193–202.

63. Hritcu L, Noumedem JA, Cioanca O, Hancianu M, Kuete V, Mihasan M. Methanolic extract of piper nigrum fruits improves memory impairment by decreasing brain oxidative stress in amyloid beta(1–42) rat model of Alzheimer's disease. Cell Mol Neurobiol. 2014 Apr 1;34(3):437–49.

64. Wang R, Chen P, Shen Z, Lin G, Xiao G, Dai Z, et al. Brain amide proton transfer imaging of rat with Alzheimer's disease using saturation with frequency alternating RF irradiation method. Front Aging Neurosci. 2019;11:217.

65. Yin Y, Liu Y, Huang L, Huang S, Zhuang J, Chen X, et al. Anti-apoptosis effect of astragaloside iv on Alzheimer's disease rat model via enhancing the expression of bcl-2 and bcl-xl. Scand J Lab Anim Sci. 2010 Dec 1;37(2):75–82.

66. Sipos E, Kurunczi A, Kasza Á, Horváth J, Felszeghy K, Laroche S, et al. β-Amyloid pathology in the entorhinal cortex of rats induces memory deficits: Implications for Alzheimer's disease. Neuroscience. 2007 Jun 15;147(1):28–36.

67. Zaher MA, Bendary MA, Aly AS. Effect of thymoquinone against aluminum chloride-induced Alzheimer-like model in rats: a neurophysiological and behavioral study. Med J Cairo Univ. 2020 Mar 1;88(1):355–65.

68. Olajide OJ, Yawson EO, Gbadamosi IT, Arogundade TT, Lambe E, Obasi K, et al. Ascorbic acid ameliorates behavioural deficits and neuropathological alterations in rat model of Alzheimer's disease. Environ Toxicol Pharmacol. 2017 Mar 1;50:200–11.

69. Shi L, Zhang Z, Li L, Hölscher C. A novel dual GLP-1/GIP receptor agonist alleviates cognitive decline by re-sensitizing insulin signaling in the Alzheimer icv. STZ rat model. Behav Brain Res. 2017 Jun 1;327:65–74.

70. Allard MJ, Brochu ME, Bergeron JD, Sebire G. Hyperactive behavior in female rats in utero-exposed to group B Streptococcus-induced inflammation. Int J Dev Neurosci. 2018;69(1):17–22.

71. Hamm RJ, Pike BR, O'Dell DM, Lyeth BG, Jenkins LW. The rotarod test: an evaluation of its effectiveness in assessing motor deficits following traumatic brain injury. J Neurotrauma. 1994 Jan;11(2):187–96.

72. Meconi A, Wortman RC, Wright DK, Neale KJ, Clarkson M, Shultz SR, et al. Repeated mild traumatic brain injury can cause acute neurologic impairment without overt structural damage in juvenile rats. PLoS ONE. 2018;13(5):1–24.

73. Schönfeld LM, Dooley D, Jahanshahi A, Temel Y, Hendrix S. Evaluating rodent motor functions: Which tests to choose? Neurosci Biobehav Rev. 2017 Dec 1;83:298–312.

74. Breland K, Breland M. The misbehavior of organisms. Am Psychol. 1961;16(11):681–4.

75. Hernandez AR, Truckenbrod LM, Campos KT, Williams SA, Burke SN. Sex differences in age-related impairments vary across cognitive and physical assessments in rats. Behav Neurosci. 2020;134(2):69–81.

76. O'Connor CA, Cernak I, Vink R. Interaction between anesthesia, gender, and functional outcome task following diffuse traumatic brain injury in rats. J Neurotrauma. 2003 Jun;20(6):533–41.

77. Huang TC, Lu KT, Wo YYP, Wu YJ, Yang YL. Resveratrol protects rats from Aβ-induced neurotoxicity by the reduction of iNOS expression and lipid peroxidation. PLoS ONE. 2011;6(12).

78. Navabi SP, Sarkaki A, Mansouri E, Badavi M, Ghadiri A, Farbood Y. The effects of betulinic acid on neurobehavioral activity, electrophysiology and histological changes in an animal model of the Alzheimer's disease. Behav Brain Res. 2018 Jan 30;337:99–106.

79. Kuo SC, Chio CC, Yeh CH, Ma JT, Liu WP, Lin MT, et al. Mesenchymal stem cell-conditioned medium attenuates the retinal pathology in amyloid-β-induced rat model of Alzheimer's disease: Underlying mechanisms. Aging Cell. 2021;20(5):e13340.

80. Lee JM, Shin MS, Ji ES, Kim TW, Cho HS, Kim CJ, et al. Treadmill exercise improves motor coordination through ameliorating Purkinje cell loss in amyloid beta23-35-induced Alzheimer's disease rats. J Exerc Rehabil. 2014;10(5):258–64.

81. Prakash A, Medhi B, Chopra K. Granulocyte colony stimulating factor (GCSF) improves memory and neurobehavior in an amyloid-β induced experimental model of Alzheimer's disease. Pharmacol Biochem Behav. 2013;110:46–57.

82. Justin-Thenmozhi A, Dhivya Bharathi M, Kiruthika R, Manivasagam T, Borah A, Essa MM. Attenuation of aluminum chloride-induced neuroinflammation and caspase activation through the AKT/GSK-3β pathway by hesperidin in Wistar rats. Neurotox Res. 2018 Oct 1;34(3):463–76.

83. Lakshmi BVS, Sudhakar M, Prakash KS. Protective effect of selenium against aluminum chloride-induced Alzheimer's disease: behavioral and biochemical alterations in rats. Biol Trace Elem Res. 2015 May 1;165(1):67–74.

84. Hosseini SA, Salehi OR, Farzanegi P, Farkhaie F, Darvishpour AR, Roozegar S. Interactive effects of endurance training and royal jelly consumption on motor balance and pain threshold in animal model of the Alzheimer disease. Arch Neurosci [Internet]. 2020 [cited 2023 Nov 15];7(2). Available from: https://brieflands.com/articles/ans-91857.html#abstract

85. Manral A, Meena P, Saini V, Siraj F, Shalini S, Tiwari M. DADS analogues ameliorated the cognitive impairments of Alzheimer-like rat model induced by scopolamine. Neurotox Res. 2016 Oct 1;30(3):407–26.

86. Stanojevic J, Dragic M, Stevanovic I, Ilic T, Stojanovic I, Zeljkovic M, et al. Intermittent theta burst stimulation ameliorates cognitive impairment and hippocampal gliosis in the Streptozotocin-induced model of Alzheimer's disease. Behav Brain Res. 2022 Sep 5;433:113984.

87. Pang K, Jiang R, Zhang W, Yang Z, Li LL, Shimozawa M, et al. An App knock-in rat model for Alzheimer's disease exhibiting Aβ and tau pathologies, neuronal death and cognitive impairments. Cell Res. 2022 Feb;32(2):157–75.

88. Lambert MI, Van Zyl C, Jaunky R, Lambert EV, Noakes TD. Tests of running performance do not predict subsequent spontaneous running in rats. Physiol Behav. 1996 Jul 1;60(1):171–6.

89. Lerman I, Harrison BC, Freeman K, Hewett TE, Allen DL, Robbins J, et al. Genetic variability in forced and voluntary endurance exercise performance in seven inbred mouse strains. J Appl Physiol. 2002 Jun;92(6):2245–55.

90. Rustay NR, Wahlsten D, Crabbe JC. Assessment of genetic susceptibility to ethanol intoxication in mice. Proc Natl Acad Sci. 2003 Mar 4;100(5):2917–22.

91. Rustay NR, Wahlsten D, Crabbe JC. Influence of task parameters on rotarod performance and sensitivity to ethanol in mice. Behav Brain Res. 2003 May 15;141(2):237–49.

92. Dreesen E, Riedel G. Comparison of mouse motor performance on different RotaRod systems. In: Grant RA, Allen T, Spink A, Sullivan M, editors. Measuring Behavior 2018. Manchester, UK; 2018. p. 389–91.

93. Bohlen M, Cameron A, Metten P, Crabbe JC, Wahlsten D. Calibration of rotational acceleration for the rotarod test of rodent motor coordination. J Neurosci Methods. 2009 Mar 30;178(1):10–4.

94. Cartmell SM, Gelgor L, Mitchell D. A revised rotarod procedure for measuring the effect of antinociceptive drugs on motor function in the rat. J Pharmacol Methods. 1991 Sep 1;26(2):149–59.

95. Rozas G, Guerra MJ, Labandeira-García JL. An automated rotarod method for quantitative drug-free evaluation of overall motor deficits in rat models of parkinsonism. Brain Res Protoc. 1997 Dec 1;2(1):75–84.

96. Deacon RMJ. Measuring motor coordination in mice. JoVE J Vis Exp. 2013 May 29;(75):e2609.

97. Shan HM, Maurer MA, Schwab ME. Four-parameter analysis in modified Rotarod test for detecting minor motor deficits in mice. BMC Biol. 2023 Aug 17;21(1):177.

98. Zieglowski L, Kümmecke A, Ernst L, Schulz M, Talbot SR, Palme R, et al. Severity assessment using three common behavioral or locomotor tests after laparotomy in rats: a pilot study. Lab Anim. 2020 Dec 1;54(6):525–35.

A T maze

B Y maze