Large-scale skin metagenomics reveals extensive prevalence, coordination, and functional adaptation of skin microbiome dermotypes across body sites

Chengchen Li[1,2*], Aarthi Ravikrishnan[1*], Indrik Wijaya[1*], Ahmad Nazri Mohamed Naim[1], Jean-Sebastien Gounot[1], Stephen Wearne[3], Yue Yuan On[1], Eliza Xin Pei Ho[1], Qi Yan Ang[1], Minghao Chia[1], Ng Hui Qi Amanda[1], Srinivas Ramaswami[3], Yik Weng Yew[4,5], Marie Loh[1,4,5,6], Ruo Yan Ong[3], Faith Ng[3], Sze Han Lee[3], James Chan[3,7], The HELIOS Study team, Viduthalai Rasheedkhan Regina[3,8], Thomas Dawson[3,9#], John Chambers[5#], John Common[3#], Niranjan Nagarajan[1,2#]

[1]*Genome Institute of Singapore (GIS), Agency for Science, Technology and Research (A*STAR), Singapore, Republic of Singapore*

[2]*Yong Loo Lin School of Medicine, National University of Singapore, Singapore, Republic of Singapore*

[3]*A*STAR Skin Research Labs (A*SRL), Agency for Science, Technology and Research (A*STAR) & Skin Research Institute of Singapore (SRIS), Singapore, Republic of Singapore*

[4]*National Skin Centre, National Healthcare Group, Singapore, Republic of Singapore*

[5]*Lee Kong Chian School of Medicine, Nanyang Technological University, Singapore, Republic of Singapore*

[6]*Department of Epidemiology and Biostatistics, Imperial College London, South Kensington, London, UK*

[7]*A*STAR Singapore Institute of Food and Biotechnology Innovation (SIFBI), Agency for Science, Technology and Research (A*STAR), Singapore, Republic of Singapore*

[8]*Singapore Centre for Environmental Life Sciences and Engineering, Nanyang Technical University, Singapore, Republic of Singapore*

[9]*Department of Drug Discovery, College of Pharmacy, Medical University of South Carolina, Charleston, SC, USA*

*Joint First Authors

#Corresponding Authors

Lead Contact: nagarajann@gis.a-star.edu.sg

Abstract

The human skin microbiome is increasingly recognized to influence skin health, immune function and disease susceptibility. However, large-scale, multi-site metagenomic studies in the general population remain scarce, limiting our understanding of the full diversity and variability of skin microbial communities across different individuals and body sites. Here, we present the largest-to-date skin microbiome dataset (>3,550 shotgun metagenomes across 18 sampling sites), generated using standardized cohort-wide protocols and rigorous contaminant removal. We define and characterize 17 robust, site-specific dermotypes—distinct microbial configurations—across seven out of nine body sites, whose coordination across disparate skin regions suggests systemic microbiome regulation. We also uncover striking bilateral concordance (median Pearson correlation>0.95 except for axilla) in microbial composition, further suggesting that host-intrinsic factors may contribute more than behavioral influences to shape skin microbiota. We trained near-perfect classifiers for dermotype labels for dermotype identification with as few as five signature species (AUC-ROC>0.98). Dermotype-dependent microbial interactions such as hypoxia-dependent inhibition of *S. hominis* by *S. epidermidis* and *M. luteus*, and metabolic resource utilization (e.g., galactose and histidine metabolism), reveal adaptive specialization within dermotypes, with implications for microbial assembly and host health. Notably, we associate a *S. hominis*-dominated elbow crease dermotype with skin irritation and itch, suggesting a potential link between microbial shifts and variations in barrier functions. By establishing dermotypes as a key organizing principle of skin microbial ecology, our findings provide a framework for microbiome-based prognostics and personalized therapeutic strategies.

Introduction

The human skin harbors a diverse community of bacterial, viral, and eukaryotic microbial species whose specific composition is known to vary substantially across body sites[1–4]. Within an individual and body site, community composition is however surprisingly stable over time[5]. Several factors could influence skin microbiome composition, including skin physiology[1,2], microbial ecology[6–9], lifestyle choices[10–13], and environmental influences[14–18], but the extent to which they shape microbiome heterogeneity across individuals within a population remains unknown.

The presence of distinct ecological community types is a key organizing principle and risk factor associated with host health, particularly in the gut and the vaginal tract[37–45]. For the gut microbiome, initial studies revealed three distinct so-called enterotypes—stable microbiome patterns within the gut—each characterized by the predominance of either *Bacteroides, Prevotella,* or *Ruminococcus* species*,* and further associated with long-term dietary patterns[37–39]. A recent study identified five entero-signatures (*Bacteroides, Firmicutes, Prevotella, Bifidobacterium,* and *Escherichia*), with combinations of these signatures accurately describing gut microbial composition in individuals across diverse human populations[22]. Similarly, vaginal microbiome studies have consistently identified five distinct community types dominated by various *Lactobacillus* species and other anaerobes (e.g., *Gardnerella*, *Prevotella,* and *Bifidobacterium*) and linked them with personal hygiene, lifestyle, and reproductive health factors[23–27].

Skin microbiome studies, in comparison, have primarily focused on disease-associated or single-site studies[17,18,28–30]. In previous work, we showed that atopic dermatitis patients have one of two distinct community compositions—so-called dermotypes—in the elbow crease, one of which is associated with more severe itch, higher IgE levels, and increased disease severity[28]. Similarly, a study on psoriasis identified a distinctive enrichment of *Firmicutes* and *Actinobacteria*-rich communities on the skin of affected individuals, independent of body site[29]. Finally, two distinct microbial communities—co-called cutotypes—on the face have been associated with air pollution exposure, and further linked to differences in skin physiological measurements and abundance of antimicrobial resistance genes[17,18]. Despite these initial studies, the heterogeneity of skin microbiome configurations across body sites in the general population has not been explored.

Here we address these knowledge gaps by comprehensively studying the skin microbiomes
in 18 different sampling sites of a multi-ethnic cohort of 200 adults from the general population
that is not focused on a specific disease. We captured a broad range of parameters by combining
shotgun metagenomic sequencing data (n=3,600 samples), demographic information (e.g., age,
ethnicity, etc.), skin physiological measurements (e.g., skin pH, hydration, etc.) and extensive skin
health questionnaire data (e.g., skin care and skin discomfort/disease related questions). Analysis
of body site-specific microbiome variability revealed strong bilateral concordance within most
individuals, but substantial inter-host variability. We uncover the existence of distinct ecological
configurations—dermotypes—in seven out of nine body sites, including five different dermotypes
in the axilla. Each dermotype was found to contain a core set of taxa that were not necessarily core
to its respective skin site, including species with median relative abundances <0.5%, together with
ancillary species that have dermotype-specific blooms in some individuals (e.g., *Pseudomonas*
stutzeri, Acinetobacter pittii/calcoaceticus/nosocomialis). Microbiome-intrinsic factors associated
with dermotypes include differences in species co-abundance patterns, interspecies interactions,
and microbial metabolic pathways within a site. Host factors such as demographics were found to
be informative for dermotype prediction in only a few body sites (e.g., elbow crease, mean AUC-
ROC=0.79), while intrinsic factors such as metabolic functions and dermotypes from other body
sites typically had greater predictive power. We trained near-perfect classifiers for dermotype
labels (AUC-ROC>0.98), with as few as five species in all cases, highlighting the potential for
dermotype identification with just a few signature species. Overall, this work establishes the
importance of accounting for population heterogeneity in skin microbiome investigations, and
highlights the diverse intrinsic and extrinsic factors that can shape skin microbiome variability and
its impact on host phenotypes. We further provide a clinically actionable framework for assigning
skin microbiome states to individuals, potentially facilitating the development of stratification and
personalized treatments.

Results

Analysis of inter-individual heterogeneity in the skin microbiome identifies robust
dermotype states across multiple body sites

To capture skin microbiome heterogeneity across individuals and different body sites in the general
population, we obtained a large collection of skin samples (n=3,600) from 18 different sampling

sites in 200 adults recruited from the Health For Life In Singapore (HELIOS) precision medicine cohort[31] (**Figure 1A**, **Table 1**; **Methods**). The samples included the left and right sides of nine skin sites: Scalp – Sc, Parietal scalp – Ps, Forehead – Fo, Cheek – Ch, Upper back – Ub, Axilla – Ax, Antecubital fossa (elbow crease) – Ac, Volar forearm – Vf, and Lower leg – Ll (**Figure 1A**). 99.6% of the samples (3,586/3,600) were successfully and consistently processed for the construction of metagenomic libraries and sequenced on an Illumina HiSeq platform to generate >70 billion reads and >10Tbp of data in total (2×150bp reads, a median of 21.6 million reads and 3.3Gbp of data per library). We processed the resulting metagenomic datasets to filter reads for quality, identify the non-human fraction (median of 7.4 million reads and 1.1Gbp per library), and conduct microbial taxonomic profiling. Data from negative controls (n=46; **Supplementary File 1**) were integrated to identify and exclude contaminant taxa (**Supplementary File 2**) and remove poor quality profiles (**Supplementary Figure 1**; **Methods**). Separately, we collected extensive participant data to investigate associations between inter-individual skin microbiome variations and host factors, including demographic information (age, gender, ethnicity, etc.), skin physiological measurements (sebum level, pH, etc.), skin-care practice (hygiene, personal-care product usage), and skin health data (skin discomfort and diseases; **Table 1**). This collection provides the largest consistently processed skin metagenome dataset for a population-wide cohort to date (n=3,572; **Supplementary File 3**), enabling in-depth comparisons of the skin microbiome across individuals and analysis of its implication for skin health.

Leveraging data availability from the left and right sides of different body sites of many individuals, we first investigated whether the skin microbiomes exhibited similarity between sides. While several skin physiological properties, including sebum level[32] and trans-epidermal water loss (TEWL)[33], are known to exhibit bilateral symmetry, human behaviors such as dominant hand use or skin product application are less so. Limited previous work suggested that in a few individuals[35] and some sites[34,28] bilateral consistency of microbial community composition was observed. However, our data suggests this is a strong and general phenomenon across body sites and for most individuals (median left-right Pearson correlation>0.95, except for axilla; **Figure 1B**). At all nine body sites studied, skin microbiomes from the left and right sides of the body were significantly more similar than across individuals (Wilcoxon test p-value<0.001; **Figure 1B**). This pattern is robust and seen with the Yue-Clayton similarity index as well, which factors in shared and non-shared species[36] (Wilcoxon test p-value<0.001, **Supplementary Figure 2A**). As we

found that intra-individual similarity is inversely correlated with community (within-sample)
diversity (Shannon index, **Figure 1B**), which in turn is influenced by *Cutibacterium acnes*
dominance[3], we reanalyzed the data after removing *C. acnes* from the profiles and confirmed that
bilateral concordance is a uniformly consistent property across body sites (Wilcoxon test p-
value<0.001, **Supplementary Figure 2B**; **Methods**). Despite the general concordance observed,
14% of sites (255/1779) exhibited discordance in 121 individuals across different body sites
(**Supplementary Figure 3A**; **Methods**). Intriguingly, bilateral discordance was enriched in a few
individuals (n=5, binomial test BH-adjusted p-value<0.05; **Supplementary Figure 3B**), and
multiple body sites exhibited correlated patterns of bilateral discordance (e.g., head sites and upper
back; **Supplementary Figure 3C**). Specific associations were detected between bilateral
microbiome discordance and various host attributes, including gender (cheek discordance in
women; **Supplementary Figure 3D**), personal habits (scalp discordance and daily hair covering;
Figure 1C), and skin conditions (forehead discordance and higher itch score; **Figure 1C**). Hence,
while the adult skin microbiome is typically strongly bilaterally concordant, discordance at specific
sites may be triggered by host-specific differences in the skin microenvironment arising out of
behavioral and/or skin condition changes.

Aggregating skin microbiome taxonomic profiles across body sites confirmed that skin
microbiome composition is influenced by skin microenvironment[1,2], with profiles from the same
site largely clustering together, and axilla (Ax) sites with their hypoxic and moist environments
being the most distinct (**Supplementary Figure 4A**). While sebaceous head and scalp sites (Sc,
Ps, Fo, and Ch) exhibited the highest similarity, even the two most similar scalp sites (Ps, Sc) had
statistically significant differences in beta (between-sample) diversity (PERMANOVA adonis
R^2=0.003, p-value<0.05; **Supplementary Figure 4B**). Site-specific differentially abundant
species included known enrichment in lipophilic *C. acnes* in sebaceous head sites[1], the fungus
Malassezia globosa in the elbow crease (Ac) and upper back[37] (Ub), and *Malassezia restricta* in
scalp sites, as well as new observations including *Cutibacterium avidum*, *Cutibacterium*
granulosum, and various *Staphylococcus* and *Corynebacterium* species in the axilla (consistent
with their preference for moist environments), and *Micrococcus luteus, Moraxella osloensis*, and
common oral taxa (*Streptococcus pneumoniae*, *Corynebacterium matruchotii*, and *Rothia*
mucilaginosa) in the forearm (Vf) and elbow crease (Ac; **Supplementary File 4**).

Despite these consistent differences across sites, we found that the individual skin
microbiomes from a given body site exhibit multimodal distributions across principal axes of
variation (e.g., Ac and Ax sites; **Supplementary Figure 4A**), suggesting substantial inter-
individual heterogeneity in the population. Exploiting the availability of a large number of
consistently processed taxonomic profiles for each site (n>395), we developed a systematic
clustering strategy to identify robust stratification in the data and assess stability of cluster
membership via bootstrapping (n=100; **Supplementary Figure 5**; **Methods**). Overall, seven out
of nine sites (i.e. Ax, Ac, Vf, Ub, Fo, Ps, Sc) showed clear evidence for multiple robust clusters
we term dermotypes, with all clusters having a median stability >90%. Of note, axilla sites
displayed the greatest diversity with five robust dermotypes, each represented by a high-density
region of samples (**Figure 1D**). Additionally, we identified two dermotypes at each of the
remaining sites except cheek and lower leg sites, where only a single robust cluster was detected
(**Supplementary Figure 6A-F**). In all seven sites with dermotypes identified, a *C. acnes*-enriched
dermotype with high prevalence (Ax-2, Ac-1, Vf-1, Ub-1, Fo-1, Ps-1, and Sc-1; **Figure 1E**) and
lower community diversity was observed (except Ac-1, Wilcoxon test p-value<0.05;
Supplementary Figure 7A). In the axilla, the remaining four dermotypes were dominated either
by combinations of *Corynebacterium* and *Dermabacter* species with higher α-diversity (Ax-1, Ax-
4) or two different *Staphylococcal* species (*Staphylococcus hominis* at Ax-3, *Staphylococcus*
epidermidis at Ax-5) with intermediate levels of community diversity. Intriguingly, the elbow
crease exhibited a *Staphylococcus*-dominant dermotype (Ac-2) seen in >15% of the general
population, similar to our previous observation in atopic dermatitis patients[28], while scalp sites
(Sc, Ps) exhibited an *M. restricta* enriched dermotype in ~30% of the population, with its known
association with dandruff[38] (**Supplementary Figure 7B**). In other body sites, while *M. luteus* and
M. osloensis were the most abundant species after *C. acnes* in the less-prevalent dermotypes, a
broader set of *Corynebacterium*, *Dermabacter, Streptococcus*, *Finegoldia*, *Brevibacterium,* and
Rothia species were found to be differentially abundant, highlighting their potential contributions
to dermotype structure and function (**Supplementary Figure 7B**; **Supplementary File 5**).
Species-level relative abundances enabled the training of near-perfect classifiers for dermotype
labels, with as few as five species in all cases, highlighting the potential for dermotype
identification with just a few signature species (AUC-ROC>0.98; **Supplementary Figure 7C**;
Supplementary File 6).

Building on the observation that skin microbiomes are bilaterally concordant, we
investigated whether dermotypes across different body sites exhibited consistent associations
within individuals (**Figure 1F**). Indeed, remarkably strong associations were detected between
dermotypes across various body sites, particularly those between physically proximal sites (e.g.,
scalp and parietal scalp), those with similar physiological characteristics (e.g., scalp and forehead),
but also more distant sites with different physiology (e.g., dry volar forearm and sebaceous
forehead sites; Fisher's exact test adjusted p-value<0.05; **Figure 1F**). These associations do not
seem to be fully explained just by preferences for specific species; for example, *Staphylococcus*-
dominant dermotypes in the elbow crease and axilla were negatively associated with each other
(**Supplementary Figure 8A**), even though *Staphylococcus* strains are known to be transmissible
via skin contact[39–41]. Furthermore, no strong correlations were observed for the abundance of
Staphylococcus species between these sites (**Supplementary Figure 8B**), indicating that other
systemic factors play a role in the observed dermotype associations. In summary, distinct
dermotype configurations exist both across individuals and body sites. Within individuals, these
exhibit high levels of bilateral and cross-site concordance, likely as a complex reflection of shared
host, environmental, and microbial factors that we explore next.

Dermotype-specific core and ancillary species exhibit coordinated patterns across body sites

To gain deeper insights into dermotype composition, we categorized species into dermotype-
specific core (with ≥50% prevalence) and ancillary (<10% prevalence, ≥3 occurrences) species
(**Figure 2A**; **Methods**). Across all dermotypes, this identified 15 core bacterial species from the
genera *Cutibacterium*, *Staphylococcus*, *Corynebacterium*, *Micrococcus,* and *Moraxella*, and two
core fungal species from the genus *Malassezia* (**Figure 2B**). No archaea or viruses were core
among the 560 species analyzed, emphasizing their sporadic distribution in the human skin
microbiome[5]. Each dermotype contained 2-12 core species, collectively accounting for >70% of
community abundance, with *C. acnes* being a universal core species across all dermotypes and
sites. Most taxa were, however, core in only some sites or dermotypes, e.g., *Dermabacter sp.*
HFH0086, which was categorized as a core species in the most common dermotype in the axilla
(Ax-1, prevalence>77%) but rarely seen in the other dermotype (Ax-2, prevalence<2%; **Figure**
2B). Even abundant species such as *S. epidermidis* and *S. hominis* were not universally core, while
some taxa with low relative abundances (median abundance<0.5%) were core in some dermotypes
(e.g., *Corynebacterium tuberculostearicum*, *Corynebacterium pseudogenitalium*, *Staphylococcus*

caprae/capitis, and *M. globosa*). This co-existence of high- and low-abundance core species within

communities suggests that their collective maintenance may arise from niche partitioning or

complementary functional roles, independent of an abundance effect[42] (**Figure 2B**).

Building on our observation of intra-individual dermotype correlation between different

body sites (**Figure 1F**), we analyzed intra-individual correlations in core species abundances

across the nine skin sites. Several taxa, such as *M. restricta*, *S. epidermidis,* and *C. acnes*, exhibited

modest positive correlations when aggregated across all sites spanning diverse microenvironments

(mean ρ=0.26), and higher correlations when excluding a few sites (e.g. *M. restricta* excluding Ax,

mean ρ=0.37; **Supplementary Figure 9A-D**). This suggests that core species shared across skin

sites are systemically coordinated within individuals, potentially mediated by global sebum

secretion favoring lipophilic taxa, by horizontal gene transfer disseminating antibiotic resistance

genes and facilitating strain dispersal[43], or by immune-driven niche filtering[44,45]. In contrast, *S.*

hominis showed positive correlations limited to the non-scalp region (**Supplementary Figure 9E**),

while *M. globosa* displayed divergent correlation patterns, with positive associations among facial

sites (ρ=0.28) but negative correlations between the cheek and upper back (ρ=-0.25;

Supplementary Figure 9F). These heterogeneous patterns imply that systemic host drivers might

interact with localized ecological factors, such as interspecies competition and

microenvironmental variation (e.g., pH, humidity gradients), further modulating cross-site

microbial assembly patterns.

The 25 ancillary species we identified were disproportionally abundant at various sites and

dermotypes (in some cases >50% relative abundance) despite being, by definition, not overall very

prevalent (<10%; **Figure 2C**; **Methods**). This group includes bacteria primarily from the

Pseudomonas, *Acinetobacter*, and *Gordonia* genera (often environmentally resilient and biofilm

forming), along with bacteriophages of common skin bacteria (e.g., *Cutibacterium*,

Staphylococcus), common skin viruses (e.g., polyomavirus and human papillomavirus), and a

fungus (*Malassezia slooffiae*; **Figure 2C**). Nearly half of ancillary taxa (12/25=48%) showed

specificity to a particular dermotype; for example, *Acinetobacter pittii/calcoaceticus/nosocomialis*

was disproportionally abundant (median abundance>50%) only in the less prevalent forehead

dermotype Fo-2, indicating that ancillary taxa may be adapted to specific niches available only in

some individuals and microbiome configurations. This was further seen in the form of significant

enrichment of ancillary species across multiple body sites of many individuals ($\geq$2 sites in 87

individuals, Poisson binomial test BH-adjusted p-value<0.05; **Figure 2C**, **Supplementary Figure**
10A). A striking example of this is *polyomavirus HPyV6,* which was abundantly present (>10%)
in many skin sites of some individuals (n=3) despite being lowly abundant in the general
population (<0.1%; **Figure 2D**). Interestingly, we also detected co-occurrence patterns in these
"blooms" of ancillary species across various body sites and species (**Supplementary 10B**),
including statistically significant coordination between head and limb sites for *polyomavirus*
HPyV6 (cheek and lower leg, scalp and forearm, Fisher's exact test p-value<0.05; **Figure 2E**). In
some cases, more frequent carriage of an ancillary species across body sites was also associated
with host attributes including ethnicity (Chinese and *Malassezia slooffiae*) and skin phenotypes
(itch severity and *Betapapillomavirus 2;* adjusted p-value<0.05; **Supplementary Figure 10C**).
These results highlight the dermotype-specific nature of core and ancillary species, their
coordinated distribution across body sites within individuals, and their unique associations with
host attributes.

Species co-abundance patterns reveal dermotype-dependent microbial interactions

To understand how dermotypes emerge at various skin sites, we analyzed microbial correlated
abundance (co-abundance) patterns with a view to gaining insights into resource competition and
metabolic cross-feeding interactions (**Methods**). The scale of our dataset enabled robust
identification of >100 species-level co-abundance relationships across the nine body sites and 16
distinct dermotypes (**Supplementary File 7**). These include known antagonism between *C. acnes*
and *S. epidermidis*[46–48], and cross-feeding interactions between *Corynebacterium* and
Staphylococcal species[49] (|r|≥0.3, CCREPE q-value<0.05; **Figure 3A**; **Supplementary Figure 11**).
Systematic comparison of co-abundance networks and hub species—highly connected taxa—
across dermotypes from the same site showed that while a core set of relationships are conserved,
8-48% of co-abundance relationships differed significantly between dermotypes, with highest
differences in the axilla and lowest in parietal scalp and forearm (Fisher's Z-test p-value<0.05;
Supplementary Figure 12A). For example, *C. acnes* and *M. luteus* abundances displayed a strong
negative correlation in the Ac-1 dermotype of the elbow crease, but positive correlation values
were obtained in the Ac-2 dermotype (**Figure 3A**).

This unexpected pattern extended to core species (significant changes in 10-68% of co-
abundance relationships; **Supplementary Figure 12B**), with *C. acnes* in particular frequently
exhibiting switched relationships across dermotypes (**Supplementary Figure 12C**). Given that *C.*

acnes is an aerotolerant anaerobe, its variable abundance and metabolism may reflect local oxygen

gradients. Therefore, we hypothesized that oxygen availability strongly (negatively) impacts *C.*

acnes growth and through its antagonistic interactions with other skin microbes plays a role in

shaping dermotype structure across many body sites. To explore this, we systematically conducted

pairwise antagonism assays among a subset of core skin bacteria (*C. acnes*, *S. hominis*, *S.*

epidermidis and *M. luteus*) under hypoxic, aerobic and mixed conditions (n=186 experiments, with

triplicates; **Supplementary File 8**; **Methods**). As expected, *C. acnes*, exhibited poor growth in

aerobic conditions but thrives under hypoxic conditions, while *M. luteus* (a strict aerobe) exhibited

the opposite pattern. Consistent with prior reports[50], we observed partial inhibition of *M. luteus* by

S. epidermidis (**Supplementary File 8**). Interestingly, we observed frequent oxygen-dependent

interspecies inhibitions, with *S. hominis*, a dominant species in the Ac-2 dermotype, being strongly

inhibited by *S. epidermidis* and *M. luteus* under hypoxic but not aerobic conditions (**Figure 3B**,

Supplementary Figure 12D-E). Further testing showed that the agar became alkaline (pH>7) in

aerobic conditions but turned acidic under hypoxic conditions (**Supplementary File 8**), indicating

that hypoxia-dependent inhibition could be mediated by acidification or fermentation-derived

short-chain fatty acids[51] that are unfavorable for *S. hominis* growth. Our findings suggest that the

common elbow crease dermotype (Ac-1) may represent a hypoxic niche where *C. acnes* flourishes,

but *S. hominis* remained suppressed due to inhibition by other skin species. In contrast, the less

common elbow crease dermotype (Ac-2) could provide a more aerobic environment enabling *S.*

hominis to evade inhibition and dominate, especially with reduced growth of *C. acnes*. Consistent

with this hypothesis, we noted that while *S. epidermidis* is a prevalent hub species in the Ac-1

dermotype, it is no longer so in the Ac-2 dermotype and is instead replaced by *S. hominis* as a hub

(**Figure 3C**), supporting the notion that *S. hominis* is more well-adapted to the environment of the

Ac-2 dermotype.

Further analysis of hub nodes identified key species such as *C. pseudogenitalium* and *C.*

tuberculostearicum in 4 out of 12 networks, highlighting that even low-abundance community

members can play a major role in determining community structure (**Figure 3C**). Lastly, we noted

that several bacterial species typically found as part of commensal oral flora were network hubs

for specific dermotypes in the forearm, upper back, and forehead (**Figure 3C**). Further

investigation of their species correlations revealed a guild of co-abundant oral bacteria such as

Veillonella parvula, *Rothia dentocariosa*, and *Corynebacterium matruchotii*, that were seen in

several dermotypes and were strongly co-abundant in some (e.g. Ac-1 & Vf-2; **Figure 3A**,

Supplementary Figure 11). These species are known to form a "hedgehog" structure in the

supragingival biofilm, which regulates pH in response to changes in nutrient availability associated

with food intake[52,53]. While the co-abundance patterns in these dermotypes are reminiscent of oral

interactions, it remains unknown whether they perform similar pH-stabilizing functions, act as

transient co-colonizers, or potentially exacerbate pathogenicity on skin—analogous to the role of

oral microbes in gastrointestinal diseases when translocating to the gut[54,55]. Overall, the analysis

of species co-abundance networks, supported by the scale of our datasets and experimental data

from pairwise antagonism assays, revealed dermotype-specific inter-species interactions and

switches in hub species that may be a function of the aerobicity of the environment and likely play

an essential role in dermotype assembly.

Metabolic variations between dermotypes reflect distinct host microenvironments and influence on skin phenotypes

Based on the observation that microbes residing on skin adapt to limited nutrient availability in

the form of carbohydrates, lipids, and proteins present in sweat, sebum, and the stratum

corneum[3,56], we hypothesized that differences in metabolic function between dermotypes of the

same body site could provide insights into how they assemble and interact with the host.

Systematic comparisons of metabolic pathways present in skin metagenomes (**Supplementary**

File 9) highlighted significant differences in pathway diversity across dermotypes

(**Supplementary Figure 13A; Supplementary File 10**), with a large fraction of pathways (>35%)

being differentially abundant in at least one dermotype of a body site, particularly relating to lipid,

carbohydrate, amino acid, nucleotide, and energy metabolism, as well as vitamin and cofactor

synthesis (**Supplementary Figure 13B**). We further observed a suite of metabolic pathways that,

while consistently present across distinct dermotypes of a given site, vary significantly in

abundance. For example, enrichments in arginine, tryptophan, and branched-chain amino acid

biosynthesis (e.g., L-isoleucine, L-valine; **Supplementary Figure 14**), that suggest potential

microbial compensation for amino acid auxotrophy under nutrient-limited conditions[57–59].

Focusing on key differences in microbial functions for the degradation of specific sugars

and amino acids, and oxidation of fatty acids, highlighted the strong enrichment of lactose and

galactose degradation pathways in the elbow crease dermotype Ac-2 (**Figure 3D**). Galactose is a

common component of human sweat[60] and complex stratum corneum glycoconjugates where it is

released by corneodesmosome degradation[61], and it is absorbed by skin commensals via diverse

phosphotransferase systems[62]. We therefore hypothesized that inter-individual differences may

play an important role in dermotype structure. Indeed, our data indicates that specific

Staphylococcus species, particularly *S. hominis*, contribute strongly to the enrichment of this

pathway (LACTOSECAT-PWY, **Figure 3E**), potentially due to a competitive advantage under

these conditions. To test this hypothesis, we compared growth kinetics of *S. hominis* and *S.*

epidermidis in mono- and co-culture, under baseline (BHI) and galactose-supplemented conditions

(**Supplementary Figure 15A**). While galactose enhanced overall growth in both species, *S.*

hominis exhibited disproportionally greater fitness gains in co-culture (>100% relative to *S.*

epidermidis; **Supplementary Figure 15B**), suggesting that preferential galactose metabolism by

S. hominis can provide it a competitive edge and that host-derived metabolites could shape

dermotype structure.

Similarly, histidine, a product from epidermal filaggrin breakdown and a common amino

acid detected on human skin[63], can be degraded by skin microbes[64]. We observed significant

variations in the L-histidine degradation pathway across dermotypes (**Supplementary Figure 14**).

For example, the *Cutibacterium* dominant dermotype Ac-1 in the elbow crease exhibited

enrichment in L-histidine degradation, with *M. luteus* being a key contributor (HISDEG-PWY;

Figure 3F; **Supplementary Figure 16A**). Further analysis of metagenomic and metabolomic data

collected at the elbow crease from a separate cohort (n=100; **Methods**), showed significant

differences in histidine derivatives, including enrichment for cis-urocanic acid (cis-UCA) and

depletion for trans-UCA in Ac-1 subjects (**Supplementary Figure 16B**), consistent with more

effective degradation of trans-UCA by *M. luteus* (urocanate hydrolase; **Supplementary Figure**

16A). *M. luteus* abundance on skin exhibited marginal correlation with the levels of histidine and

its derivates (ρ=0.21; **Supplementary Figure 16C**), as well as a notable growth advantage with

L-histidine supplementation *in vitro* (OD=1.8 vs 1.5; **Supplementary File 12**). Intriguingly, we

observed a negative correlation between lactose metabolism and histidine catabolism pathways

specific to Ac-2 subjects, potentially reflecting niche-specific usage of these energy sources

(**Supplementary Figure 16D**). Overall, enrichment of the histidine degradation pathway was

negatively associated with skin sensitivity (p-value<0.05; **Supplementary Figure 16E**;

Supplementary File 11), consistent with the known immunosuppressive effects of the histidine

derivative cis-urocanic acid[65]. These findings therefore highlight that dermotypes, beyond

representing mere taxonomic differences, are niches with distinct metabolic properties that can be
shaped by differences in host-derived nutrients, and in turn may impact host skin conditions.

Dermotype composition is linked to demographic factors, along with physiological,
behavioral, and skin condition attributes

Skin microbial composition can be shaped by a combination of endogenous and exogenous host
factors (e.g., genetics, skin physiology, lifestyle, and skin conditions) and in turn, influences some
of them. We thus leveraged the scale of our cohort to evaluate associations between dermotypes
and host demographics, skin physiological measurements, >15 personal care behaviors, and >25
skin conditions (**Methods**). Focusing on attributes relevant to each body site, we identified >35
significant associations (**Supplementary File 13)**.

Among demographic factors, ethnicity exhibited strong associations with axilla and elbow
crease dermotypes. In particular, the axilla dermotypes dominated by *C. acnes* (Ax-2), *S. hominis*
(Ax-3), and *C. kroppenstedii* (Ax-4) were significantly enriched in ethnic Malay, Chinese, and
Indian subjects, respectively (adjusted p-value<0.05; **Figure 4A**). In contrast, the *S. hominis*-
enriched elbow crease dermotype (Ac-2) was more frequent in Malay and less common in Chinese
subjects (adjusted p-value<0.05; **Figure 4A**). Beyond ethnicity, the *S. hominis*-enriched elbow
crease dermotype (Ac-2) and the *Cutibacterium*-depleted upper back dermotype (Ub-2) were
significantly associated with female and older subjects (adjusted p-value<0.05). Strikingly, no
significant demographic associations were observed for head sites (forehead, scalp, parietal scalp;
Supplementary File 13), indicating that other factors (e.g. localized skincare practices or
environmental exposures) may drive microbiome variation in the head region.

We next assessed if skin physiological measurements revealed distinct trends across
dermotypes. Of note, individuals with an *S. hominis*-enriched elbow crease dermotype (Ac-2)
exhibited lower forearm surface hydration (p-value<0.05; **Figure 4B**), consistent with our prior
report of dry skin in atopic dermatitis (AD) patients harboring a *Staphylococcus*-enriched
dermotype[28], and indicating that Ac-2 may arise from compromised skin barrier function.
Additionally, elevated cheek sebum levels, a proxy for overall sebum production in the body, were
observed in individuals with *C. acnes*-enriched dermotypes at the elbow crease and the volar
forearm (i.e., Ac-1, Vf-1), consistent with the robust growth of *C. acnes* in sebum-rich
environments. In contrast, higher cheek sebum levels were associated with the *C. kroppenstedii*-
enriched dermotype in the axilla (i.e., Ax-4; p-value<0.05; **Figure 4B**). While both species are

lipophilic[66], this divergence may reflect differences in local lipid composition and species-specific

substrate preferences. More broadly, to explore whether site-specific skin physiological

measurements reflect body-wide intrinsic states, we conducted site-agnostic analyses. Intriguingly,

we observed strong associations between lower trans-epidermal water loss (TEWL) levels on the

cheek and *C. acnes*-depleted dermotypes in the upper back (Ub-2) and forehead (Fo-2; p-

value<0.05; **Supplementary Figure 17A**). Similarly, lower TEWL on the forearm associated with

the *C. acnes*-depleted parietal scalp dermotype (Ps-2; p-value<0.05; **Supplementary Figure**

17A), highlighting the systemic connections between skin barrier functionality and skin microbial

community structure.

428   Leveraging skincare routine information based on questionnaire data, several noteworthy

associations with dermotypes were observed. For example, dermotypes in the axilla were found to

frequently exhibit statistically significant associations with underarm product use, even after

accounting for demographic factors. For example, routine deodorant use was positively associated

with Ax-2 (~60% prevalence), while antiperspirant use was linked to Ax-4 (~30% prevalence);

conversely, both product uses were negatively correlated with the most common axilla dermotype

Ax-1, where deodorant and antiperspirant use were notably lower (11% and 4.9%, respectively; p-

value<0.05; **Figure 4C**). Consistent with the antimicrobial effects of these products, Ax-2 and Ax-

4 exhibited reduced alpha diversity compared to Ax-1 (**Supplementary Figure 7A**). Similarly,

routine moisturizer use and hair treatments correlated with the *C. acnes*-depleted forehead

dermotype (Fo-2), while daily hair covering was associated with the *C. acnes*-depleted parietal

scalp dermotype (Ps-2), indicating that skincare practices can significantly influence facial,

parietal scalp, and axilla microbiomes.

441   Several skin condition-related attributes, including sensitivity, itch, and eczema history,

were associated with the *Staphylococcus*-enriched elbow crease dermotype while adjusting for

demographic factors (Ac-2; p-value<0.05; **Figure 4C**), consistent with our previous atopic

dermatitis-focused study[28]. While skin sensitivity and itch were more common in those with

eczema history, Ac-2 subjects showed more severe irritation and itch, even in the absence of

eczema (**Supplementary Figure 17B-C**). Treatment of human keratinocytes with supernatants

from *S. hominis* strains revealed that several of them (5/8; as well as *S. epidermidis* strains: 9/12)

induced significantly elevated pro-IL-33 levels (cytokine signal in itch[67,68]) compared to controls

(**Supplementary Figure 17D**), suggesting that enrichment of *Staphylococcus* species on skin (as

is seen in the Ac-2 dermotype) can play a role in modulating host immune signaling, heightened
skin irritation and itch.

Finally, we applied machine learning models to test whether the information in the
demographic data, skin physiological measurements, and questionnaire responses could predict
dermotype labels across body sites (**Supplementary Figure 18A**; **Methods**). Machine learning
models that were significantly better than random were obtained in all sites except for scalp sites,
with the highest classification performance observed in the elbow crease (Ac random forest AUC-
ROC=0.8; **Figure 4D**). Among various host attributes, demographic factors emerged as the most
predictive in four out of seven sites analyzed (axilla, scalp, upper back, elbow crease), with
ethnicity frequently being among the most important features (e.g. for elbow crease; **Figure 4D**,
Supplementary Figure 18B). In summary, our findings underscore the intricate interplay between
host demographics (particularly ethnicity), skin physiology, skincare practices, and skin microbial
composition, suggesting that dermotypes could serve as biomarkers for skin health, disease risk
and therapeutic eligibility in precision medicine.

Metabolic functions and dermotypes across body sites are more strongly predictive of
dermotypes

Having shown that certain host attributes can predict dermotype occurrence, we wondered if
metabolic functions could serve this purpose. To ensure predictions were not driven by species-
specific pathway signatures, we excluded pathways highly correlated with taxa abundances
(Spearman $\rho \geq 0.9$, p-value<0.05) and trained multiple classifiers to explore key metabolic
functions that could jointly predict dermotype structure; among them, ridge classifiers achieved
the highest performance for many sites (AUC-ROC>0.95, except for Ps and Sc; **Figure 4E**). This
analysis was further restricted to conserved pathways (prevalence$\geq$95% across all dermotypes;
Supplementary Figure 19A), which still retained strong predictive ability (**Figure 4E**). We
identified a small number of strongly predictive pathways (**Supplementary Figure 19B**), and their
aggregation into metabolic superclasses highlighted the role of carboxylate degradation, amino-
acid degradation, carbohydrate biosynthesis, fermentation, and secondary metabolite biosynthesis
in being predictive of dermotype labels (**Supplementary Figure 19C**). Hence, a small number of
microbial metabolic functions are more predictive of dermotype identity than most of the host
attributes analyzed here.

As dermotypes were strongly correlated across body sites (**Figure 1F**), we investigated if

dermotype labels could be predicted based on taxonomic profiles from other sites. Remarkably,

high predictive performance was obtained in many sites (AUC-ROC>0.8, except for Ax, Ac;

Supplementary Figure 20A), even when only data from a single site (**Supplementary Figure**

20B-C), or a small number of taxa were used (best AUC-ROC=0.89; **Supplementary Figure**

20D). Intriguingly, data from the lower leg (Ll) consistently ranked highest in feature importance

among top-performing classifiers, despite no robust dermotypes being discovered at this site.

Several species including *C. acnes*, *S. epidermidis, M. restricta*, *M. globosa,* and *M. luteus,*

consistently ranked high in terms of feature importance for dermotype classification across sites

(**Figure 4F**), reflecting the importance of their systemic coordination within the host for defining

dermotype structure. Restricting to just dermotype labels, we again noted that cross-site labels

were strongly predictive and in many cases more so than host attributes (AUC-ROC>0.84, except

for Ax, Ac; **Figure 4G**).

Ridge classifiers that integrated cross-site dermotype labels, metabolic pathway profiles,

and demographic factors were used to evaluate their relative utility for dermotype prediction

(**Figure 4H**), with uniformly high performance across all sites (AUC-ROC>0.9). Notably, cross-

site dermotype labels ranked highest by feature importance at the parietal scalp (Ps), upper back

(Ub), and volar forearm (Vf; **Figure 4H**), highlighting the importance of systemic microbial

assembly patterns underlying dermotype structure across body sites. Although metabolic pathways

were generally secondary to cross-site dermotype labels, they provided complementary

information and even ranked highest in terms of feature importance for the scalp (Sc), suggesting

that microbial metabolic capabilities may have greater influence on dermotype assembly for some

sites. Similarly, while demographic factors had lesser feature importance in general, they had

significant predictive value in select sites (e.g., elbow crease [Ac] and forehead [Fo]; **Figure 4H**),

showcasing the influence of host attributes in defining microbial community structure.

Overall, our findings emphasize the unexpected strength and predictive power of

dermotype coordination across body sites and the critical role of key microbial metabolic functions

and host demographic factors in shaping dermotypes on skin.

Discussion

Despite growing efforts to understand the role of the skin microbiome in skin health, disease-

agnostic studies in the general population to help define baseline variations and features have been

limited to a few early studies with small cohort sizes[1,5] (n<20). Here we present a metagenomic

characterization of skin microbiomes from multiple body sites (n=18) in an adult population cohort

that is an order of magnitude larger (n=200). This provides a well-powered, rich dataset that has

been consistently generated and processed (n=3,572 metagenomes), and complemented by

detailed phenotypic information about the subjects, creating a unique resource for investigating

skin microbial ecology and function in relation to host attributes and skin health.

517   We systematically identified the presence of two or more robust dermotypes in seven out

of nine body sites, revealing significant yet consistent heterogeneity in the general population,

even within a well-defined skin site. The dermotypes identified here seem to represent distinct

microbiome states in regions with higher density of support (**Figure 1D**, **Supplementary Figure**

6), unlike the continuum seen in gut enterotypes[69,70], and are defined by the abundances of multiple

species rather than just one signature species (**Supplementary File 6**). This allowed the training

of near-perfect classifiers for dermotype labels, with as few as five species in all cases. Our work

thus goes beyond prior studies which have identified skin microbial community types (i) on the

face in the context of air pollution[18] and elbow crease for eczema patients[28], or (ii) as site-agnostic

patterns that overlook key site-specific differences[17], by showing that there is clear intra-site

heterogeneity in multiple body sites (**Figure 1D-E**). For example, the axilla demonstrates striking

heterogeneity that has never been described before, with evidence for at least five robust

dermotypes dominated by multiple *Staphylococcus* and *Corynebacterium* species, providing a

basis for further studying their ecological interactions and associations with body odor[71] and

underarm product use[72]. Similarly, we identified a *Staphylococcus*-dominant elbow crease

dermotype (Ac-2) as a feature of the general population and not just atopic dermatitis patients[28],

defining an at-risk population for further study. Additional observations of *M. luteus*-enriched

dermotypes in dry sites, and *M. restricta*-enriched dermotypes in sebaceous regions, such as the

scalp, provide the foundation for further investigations into the skin microbiome's role in skin

conditions such as dryness and dandruff, potentially guiding new treatment approaches. Our

dermotype classifier models provide an accessible starting point for launching such studies that

could be based on cost-effective and clinically-applicable strategies (e.g. using qPCR panels) to

assign dermotypes to individuals.

540   Leveraging the multi-site design of our study, we revealed remarkable cross-site

correlations in dermotype patterns, even between skin sites that are physically and physiologically

distinct (e.g. dry forearm and sebaceous forehead sites; **Figure 1F**). This goes beyond the site

specificity of skin microbiome composition that has been previously demonstrated[1,2], as well as

the high bilateral concordance in skin microbiomes across all 9 sites and most individuals as we

establish here (**Figure 1B**, **Supplementary Figure 2**), and indicates a deeper systemic

coordination across the body within individual hosts. This coordination is also evident from the

ability to classify dermotypes at one site using information for as little as 10 species or a few

dermotype labels from another site (**Figure 4F**). Such patterns may partially result from correlated

abundances of a few core species shared across multiple sites (e.g., *C. acnes*, *S. epidermidis*, and

M. restricta; **Figure 2B**, **Supplementary Figure 9**), perhaps due to strain-specific ecological

functions[73–75]. However, the preference for specific species/strains alone does not fully explain the

patterns observed, as some taxa show no significant correlations between specific sites (e.g.,

Staphylococcus species in axilla and elbow crease; **Supplementary Figure 9**). This suggests that

other shared factors (e.g. host genetics, global sebum production levels) or behaviors (e.g.

showering habits) might dominantly shape the skin microbiome in a non-site-specific, bilaterally-

symmetric manner. The observation that loss of bilateral concordance can also be correlated across

body sites further suggests that such dysbiosis might also be shaped by coordinated influences

across the body (**Supplementary Figure 3**). Interestingly, we also observed body-wide

enrichment of ancillary species, particularly skin eukaryotic viruses (e.g., polyomaviruses or

papillomaviruses; **Figure 2C-D**), within some individuals, indicating that immune dysregulation

could also play a role[1,2] (**Figure 2E**). Overall, these observations provide the key hypothesis that

the colonization and adaptation of microbial species, and the corresponding emergence of

dermotypes, is a coordinated phenomenon across various skin sites that is strongly shaped by host

factors, and not purely a stochastic assembly process.

Systematically investigating this further based on dermotype-dependent co-abundance

patterns and pairwise antagonism assays between core species, we identified varying oxygen

availability (shaped by skin physiology[76,77]) as a key factor impacting interspecies interactions,

such as hypoxia-dependent inhibition of *S. hominis* by *S. epidermidis* and *M. luteus* (**Figure 3B**).

Such oxygen-dependent antagonism may arise from differential toxin production (e.g., *S.*

epidermidis phenol-soluble modulins produced under hypoxia[78]) or redox-sensitive metabolic

cross-feeding[79]. Similar modulation was also seen in a previous study of tri-species interaction

dynamics in healthy and dandruff scalp microbiomes[80], highlighting the role of oxygen availability

in defining microbial interactions on skin. While oxygen gradients explain part of this dynamic,

nutrient availability (lipids, carbohydrates and amino acids) may synergistically shape interactions.

For example, in sebaceous environments *C. acnes* may grow well by hydrolyzing sebum

triglycerides into free fatty acids and thus acidifying the skin surface. This acidification, combined

with lipid reduction and potential antimicrobial activities, likely contributes to unfavorable

conditions for *M. luteus* and other skin bacteria[81]. Conversely under lipid-depleted conditions, the

competitive dominance of *C. acnes* may diminish (**Supplementary File 8**), while species such as

S. hominis and *M. luteus* may competitively leverage availability of other nutrients such as

galactose and histidine to grow well (**Figure 3E-F**). Future work in human cohorts would benefit

from integration of metatranscriptomic[87], metabolomic[86], and lipidomic[84,82] data in a longitudinal

study design, while *in vitro* studies could use multi-species consortia co-cultured with skin models

to further investigate the relative contributions of oxygen gradients, nutrient availability, and pH

in shaping dermotype structures across individuals.

Several of our observations in the *S. hominis*-dominated elbow crease dermotype (Ac-2)

including associations with older age, reduced surface hydration, skin sensitivity, and itch, might

indeed reflect a bidirectional interplay between microbial ecology, metabolic activities, and barrier

dysfunction (**Supplementary Figure 21**). Compromised skin barrier, whether due to dryness,

inflammation, or mechanical damage from scratching, may enhance oxygen penetration, creating

aerobic microenvironments where *S. hominis* escapes inhibition by *S. epidermidis* and *M. luteus*.

Notably, *S. hominis* may further circumvent *S. epidermidis* antagonism via galactose metabolism,

which disproportionately enhances its growth relative to *S. epidermidis* (**Supplementary Figure**

15). Thriving under these niche conditions in the skin, *S. hominis* can exacerbate itch by inducing

IL-33 expression in keratinocytes, establishing an itch-scratch cycle that further damages the

barrier and reinforces the aerobic conditions that favor its dominance (**Supplementary Figure**

21). In contrast, enhanced *M. luteus*-mediated histidine degradation in Ac-1 dermotypes could

promote production of immunosuppressive cis-urocanic acid and thus contribute to reduced skin

sensitivity in Ac-1 individuals. Together, such metabolic cross-talk underscores the role of host-

derived nutrient availability and microbial substrate preferences in shaping distinct ecological

niches, and consequently, skin phenotypes (**Supplementary Figure 21**). Further work is thus

warranted to disentangle this complex interplay in contributing to skin itch and inflammatory skin

conditions such as atopic dermatitis. As a result, dermotype status could serve to stratify subjects

in terms of their disease risk, severity and treatment response, while targeting microbial pathways
(e.g., IL-33-inducing proteases), modulating oxygen gradients (e.g., via topical emollients or
occlusion therapy), or leveraging microbial metabolic pathways (e.g., through probiotic or
metabolite supplementation) may offer more effective personalized therapeutic avenues to
interrupt this pathogenic cycle.

Overall, our study also helps delineate the potential contributions of host demographics,
skin physiology, and lifestyle factors in shaping skin microbiome variability across the population.
Leveraging our multi-ethnic cohort spanning a broad age range, we demonstrate that ethnicity,
age, and gender have site-specific associations with dermotypes, particularly in moist sites such as
the axilla and elbow crease (**Figure 4A**). While the strong ethnicity-specific patterning of axillary
dermotypes may arise from genetic predispositions to lipid and sweat content on skin[85,86],
differences in skincare practices, or cultural norms for product usage could also play a role. The
absence of demographic associations at head sites on the other hand underscore the influence of
localized practices—such as washing, chemical treatment, or hair covering—in overriding broader
host factors in these regions (**Figure 4C**), aligning with growing evidence that personal care
products can shape microbial resilience on skin[87,88,11]. However, by integrating all these factors
into a machine learning framework we found that cross-site microbial coordination and core
microbial metabolic functions are stronger predictors of dermotype identity than host attributes in
most regions (**Figure 4H**). The prominence of shared pathways such as amino acid, carbohydrate
and nucleotide biosynthesis suggests that microbial metabolic function (shaped by nutrient
availability from the host), rather than taxonomic identity, drives niche specialization
(**Supplementary Figure 19**). What is even more striking is the predictive power of dermotype
information across sites, with AUC-ROC values as high as 0.9 for some sites (**Figure 4F-G**),
pointing to systemic coordination in the host that transcends local differences in ecology.

In conclusion, we demonstrate the use of a large-scale, multi-site skin microbiome resource
of >3,550 metagenomes across 18 sampling sites to establish key features including bilateral
symmetry, dermotype prevalence, and coordinated occurrence of dermotypes across body sites.
We show that beyond mere differences in species abundance, dermotypes represent distinct
ecological niches where microbial populations adapt and specialize in exploiting their unique
resource availability and environmental conditions, reflected in dermotype-dependent microbial
interactions (e.g., *M. luteus* and *S. hominis*) and metabolic activities (e.g., galactose and histidine

degradation pathways). Integrating the availability of extensive host phenotype data, we provide

new insights into how host attributes may shape dermotypes, which in turn can influence host skin

conditions. Our results, dermotype prediction models and metagenomic datasets lay the

groundwork for future research into therapeutic strategies for skin conditions based on

microbiome-informed stratification and personalized skin interventions.

Methods

Subject recruitment and microbiome sampling

Subjects were recruited as part of the Asian Skin Microbiome Programme (ASMP) sub-study for the Health For Life In Singapore (HELIOS) study (https://www.healthforlife.sg/), a large population cohort that is central to the National Precision Medicine program (NPM) in Singapore. The HELIOS cohort study specifically recruits subjects to ensure that their demographics accurately match that of Singapore's population, considering factors such as ethnicity, age, gender and socio-economic status. Informed consent was obtained from all subjects, and associated protocols for the study were approved by the NTU Institutional Review Board (IRB reference number: IRB-2016-11-030). Subjects were excluded from the study if they were currently pregnant, breastfeeding, had any ongoing acute illness, had recent major surgery, were under the age of 30, or had any mental incapacities that would prevent them from understanding and giving informed consent. Upon arrival to the HELIOS clinic, subjects were given time to acclimatize to the temperature of the clinic. Following this, subjects were asked to complete a skin health questionnaire. This extensive questionnaire has questions related to several topics including overall skin health, skin-related medical treatments, personal care routines, product usage, lifestyle habits, and finally a series of questions relating to menopause for all female subjects (**Supplementary File 14**). Subjects were then moved to a climate-controlled room for sampling by a trained research coordinator/clinician.

A diverse set of body sites were chosen for skin surface sampling, based on their importance for consumer care studies, and to capture the breadth of the human skin microbiome. These consisted of nine different body sites, sampled on both the left and right sides (**Figure 1A**), including: scalp (Sc), forehead (Fo), cheek (Ch), upper back, (Ub), antecubital fossa (elbow crease, Ac), volar forearm (Vf), and front of lower leg (Ll), for which an adhesive tape disc was used for sampling (D-Squame Standard Sampling Discs, D100, CuDerm). The tape was repeatedly placed on the same location with pressure and peeled 25 times to saturate discs with skin surface material. For the remaining two sites, sterile swabs (FLOQSwabs Flocked Swabs, 552C, Copan) were used to sample the skin microbiome on the parietal scalp (5cm above the ear, Ps) and axilla (armpit, Ax), by soaking in 0.9% saline and rubbing back and forth on the skin for 30 seconds. One tape disc or swab was collected from each side (left and right) of the skin sites, giving 14 tape discs and

670   4 swabs per subject. All collected samples were stored at -80°C before sample processing.

671   Negative control samples were collected in the form of blank tape discs (n=34) and swabs (n=12)

672   that were processed in the same way as skin samples.

Skin physiological measurements

Trans-epidermal water loss (TEWL) was measured using a VapoMeter (Delfin) on the forearm

(left or right) and on the cheek (just below the left eye). pH was measured using a pH meter (Delfin)

on the forearm (left or right). Stratum corneum surface hydration was measured using a

MoistureMeter (Delfin) on the forearm (left or right). Skin sebum secretion was measured using a

SebumScale (Delfin) on the cheek (just below the left eye). Three consistent consecutive readings

were taken for all measurements and averaged. Measurements were performed in a controlled

temperature (20-22°C) and humidity (relative humidity 40-60%) environment after at least 20

minutes of acclimatization.

DNA extraction

Genomic DNA was extracted from skin tapes and swabs using a previously established

protocol[35,28], with modifications to improve yield. Briefly, samples were transferred into Lysing

Matrix E tubes containing 500 µL of Buffer ATL (Qiagen) and subjected to bead-beating at 4 m/s

for 30 seconds, repeated twice, using a FastPrep-24 Automated Homogenizer (MP Biomedicals).

Following homogenization, tubes were centrifuged at maximum speed for 5 minutes, and 200 µL

of supernatant was transferred to 2 mL EZ1 sample tubes (Qiagen). Proteinase K (12 µL) was

added, and samples were vortexed and incubated at 56 °C for 15 minutes. DNA was then purified

using the EZ1 Advanced XL Instrument (Qiagen) with the EZ1 DNA Tissue Kit and eluted in

50 µL of EB buffer.

Construction and sequencing of metagenomic libraries

Purified genomic DNA underwent NGS library construction steps using NEBNext Ultra II FS

DNA Library Prep Kit according to manufacturer's instructions. 26 µl of DNA added to

fragmentation reagents were mixed well and placed in a thermocycler for 10 mins at 37°C,

followed by 30 mins at 65°C to complete enzymatic fragmentation. Post fragmentation, adaptor-

ligation was performed using Illumina-compatible adaptors, diluted 10-fold as per kit's

recommendations prior to use. Post-ligation, size selection or cleanup of adaptor-ligated DNA was

performed using Ampure XP beads in a 7:10 beads-to-sample volume ratio. Unique barcode

indexes were added to each sample and amplified for 12 cycles under recommended kit conditions

to achieve multiplexing within a batch of samples. Finally, each library sample was assessed for

quality based on fragment size and concentration using the Agilent D1000 ScreenTape system.

Samples that passed the quality-checks were adjusted to identical concentrations employing

dilution and volume-adjusted pooling. The multiplexed sample pool was paired-end (2×151bp)

sequenced on an Illumina HiSeq X Ten system. All libraries were sequenced at the Novogene

sequencing facility in the Genome Institute of Singapore following standard Illumina sequencing

protocols.

Taxonomic and functional profiling

Raw Illumina sequencing reads were processed using an in-house Nextflow pipeline

(https://github.com/CSB5/shotgunmetagenomics-nf). Quality control and adapter trimming were

performed with FastQC v0.20.0 (default parameters). Human reads were removed by mapping to

the hg19 human reference genome using BWA-MEM (v0.7.17-r1188, default parameters), and

filtering with SAMtools (v1.7). Taxonomic profiling of the remaining reads was conducted with

MetaPhlAn2 (v2.7.7, default parameters). During sequencing data generation (2020 January),

MetaPhlAn3 (v3.0, 2020 April release) was tested but frequently omitted eukaryotic taxa,

including fungal species commonly observed and cultured from human skin; additionally,

MetaPhlAn4 (v4.0, 2022 April release) outputs viral profiles separately, without integration into

the non-viral taxonomic profile. Due to these limitations, we opted to use MetaPhlAn2 consistently

in this long-running project. To address the limited representation of *Malassezia* species in

MetaPhlAn2 (a commonly found fungal member of the human skin microbiome[89,90]), we

integrated results from PathoScope[91] (v2.0) for *Malassezia* species. Briefly, after human read

removal, we aligned the remaining reads to 14 *Malassezia* genomes (*M. furfur, M. obtusa, M.*

yamatoensis, M. japonica, M. sympodialis, M. pachydermatis, M. restricta, M. globosa, M.

slooffiae, M. caprae, M. cuniculi, M. dermatis, M. equina, and *M. nana*) downloaded from NCBI

using bpipe (v0.9.8.6_rc), and normalized mapped read counts by the total number of reads to

estimate the relative abundance of *Malassezia* species. Lastly, MetaPhlAn2 taxonomic profiles

(excluding *Malassezia*) were rescaled by multiplying by the proportion of reads that were not

mapped to *Malassezia* (calculated from PathoScope results). The rescaled MetaPhlAn2 profile and

PathoScope-derived *Malassezia* abundances were merged and renormalized using Total Sum

Scaling (TSS) to generate final taxonomic profiles (n=3,586) for downstream analyses. In addition,

we reannotated *Enyhydrobacter aerosaccus* as *Moraxella osloensis*, consistent with a previous

study showing that 99.4% of sequences labeled as *E. aerosaccus* in the MetaPhlAn2 database were

identified as *M. osloensis* using BLASTN against the NCBI nt database[17].

Sequencing data from negative control samples (n=46; **Supplementary File 1**) and

information from existing databases (PubMed, DISBIOME[92], MicroPhenoDB[93]) were used to

identify potential artifacts in the taxonomic profiles, particularly *kitome* species derived from DNA

extraction kits and sequencing reagents (**Supplementary Figure 1**). In the first tier, species

commonly found in negative controls (median abundance$\geq$0.1%) were evaluated for negative

correlation with library concentration, a signature of a contaminant species[94] (negative Spearman

correlation, p-value<0.05). These kitome candidates were then used to identify additional taxa

highly correlated with them, potentially also arising from contamination (Spearman $\rho\geq$0.7, p-

value<0.05). The second tier refines this initial selection by restricting to genera not previously

reported on human skin or associated with skin disease in three different databases (PubMed,

DISBIOME[92], MicroPhenoDB[93]). Poor-quality profiles were removed (n=14, 0.4%; kitome total

abundance>50%). Kitome taxa (**Supplementary File 2**) and extremely low-abundance species

(relative abundance<0.1%) were excluded from taxonomic profiles (set to 0), and profiles were

TSS-renormalized (n=3,572; **Supplementary File 3**).

Functional profiles and detailed species-specific reconstructions of microbial metabolic

pathways (MetaCyc) were generated with HUMAnN2 (v2.8.1, default parameters). Pathways that

were unannotated, or highly correlated with kitome species (Spearman $\rho\geq$0.7) were filtered out.

TSS-renormalized pathway abundance profiles were used for subsequent analyses (n=3,550;

Supplementary File 9). Downstream data analysis was performed in R v4.4.0 unless stated

otherwise.

Identification of dermotypes

Dermotypes were identified via a three-step protocol (**Supplementary Figure 6**). Firstly,

Partitioning Around Medoid (PAM) clustering was performed using Bray-Curtis dissimilarity for

each site. The optimal number of clusters was determined using key indices (prediction strength,

silhouette index, and Calinski-Harabasz index). In cases of ambiguity (i.e., disagreement across

various indices), we employed Mean-Shift clustering (bandwidths determined by cross-validation)

to search the local maxima of kernel density estimated in PCoA space (PCoA 1-3). Next, we

assessed sample stability and cluster robustness via bootstrapping (n=100). Clusters were deemed

robust if over half of the samples were consistently grouped into the same cluster in over 90% of

iterations (median stability≥0.9). Only sites with at least 2 robust clusters detected were defined as

having distinct dermotypes for subsequent analyses. Finally, we relied on PAM to infer cluster

memberships for robust dermotypes, with samples with <90% stability excluded from further

analyses.

Diversity and differential abundance analysis

Metrics for α-diversity (Shannon index) and β-diversity (Bray-Curtis dissimilarity) were computed

using the vegan package (v2.6.4). Permutational multivariate analysis of variance

(PERMANOVA) was performed with Bray-Curtis dissimilarity values to quantify differences in

microbial compositions using the adonis2 function from vegan[2] (v2.6.4) with 1,000 permutations

and restrictive permutations within blocks of SubjectID. Multivariable analysis estimating which

microbiome attributes (species, pathways) differ significantly across comparison groups was

carried out using MaAsLin2 (linear model, "LOG" transformation, prevalence≥10%, BH-adjusted

p-value<0.05), including demographic covariates (age, gender, ethnicity) as fixed effects and

subject ID as a random effect when applicable.

Construction of species co-abundance networks

Species co-abundance networks were constructed with taxonomic profiles with high cluster

membership stability (≥90%) in each dermotype using the "netConstruct" function in NetCoMi[95]

(v1.1.0), and visualized using igraph[96]. Prevalence filtering was applied to focus on prevalent taxa

(≥30% for Vf, Ac, Ub, and Fo dermotypes, and ≥10% for Sc, Ps, and Ax dermotypes); this filtering

ensures that each network construction included at least 15 taxa, minimizing spurious findings and

allowing for biologically meaningful insights. Compositionally corrected Spearman ρ for each

species pair and corresponding p-values were obtained from CCREPE[97] (v1.34.0), and

downstream analysis were restricted to species pairs with strong co-abundant or co-exclusion

patterns in each dermotype (|ρ|>0.3, BH-adjusted p-value<0.05; **Supplementary Figure 12**).

Hubs were defined as species with correlation strength-weighted degree centrality above the 95[th]

percentile in each network. Differential edges between dermotypes for each skin site were

identified using Fisher's z-test[98] for correlation values.

Antagonism co-culture assays

The bacterial strains studied (*Cutibacterium acnes* 6919, *Micrococcus luteus* 10240, *Staphylococcus epidermidis* 12228, and *Staphylococcus hominis* 27844) were obtained from ATCC. *C. acnes* was cultured in half-strength BHI broth in an anaerobic chamber at 37°C overnight with agitation, and other species were cultured under the same condition aerobically. After measuring the optical density at 600 nm (OD600), each culture was diluted to OD600=0.1 with fresh broth (5mL total) and incubated at 37°C with agitation (220 rpm) until growing to an OD that corresponds to the mid-log phase of the bacteria (i.e., *C. acnes* OD=0.6; *M. luteus* OD=1; *S. epidermidis* OD=2; *S. hominis* OD=2).

Assay plates were prepared by dispensing 2mL of molten half-strength BHI agar into each well of a 12-well plate and allowing it to solidify and dry thoroughly. Each bacterial pair was assayed in triplicates, with one well per replicate. For hypoxic conditions, plates were placed in compact plastic pouches with anaerobic sachets (Thermo Fisher Scientific Oxoid AnaeroGen, Cat. No. AN0020D). A total of 40μL of test bacterial culture (bacteria being tested for inhibitory activity; 10μL per droplet, sequentially spotted four times) was applied to each well, allowing each droplet to dry before applying the next. The following day, cultures of target bacteria (bacteria to be assessed for inhibition) were diluted 100× in 1× PBS (10× for *C. acnes*). A 10μL aliquot of the diluted target culture was then spotted adjacent to the test bacteria in each well. The plates were incubated at 37°C overnight, or two nights if the target bacteria was *C. acnes*, and clearing zones were assessed the following day.

Effect of galactose on *Staphylococcus* species growth

The effect of galactose on the growth of *Staphylococcus* species was assessed by performing monoculture and co-culture experiments with *Staphylococcus epidermidis* 12228 (SE) and *Staphylococcus hominis* 27844 (SH) in Brain heart infusion (BHI) broth with and without 5% galactose supplementation. Briefly, six different experimental conditions were designed: 1) BHI + SE 2) BHI + SH 3) BHI + Galactose 5% + SE 4) BHI + Galactose 5% + SH 5) BHI + SE + SH 6) BHI + Galactose 5% + SE + SH, with two separate negative controls including, BHI and BHI + Galactose 5%. Growth curve experiments were performed at 37°C for 48h with high amplitude, fast speed shaking in a 100-well Honeycomb plate (Growth Curves USA) with a working volume of 200μl in each well. All the experiments were performed in quadruplicates. Optical Density (OD)

at 600nm measured using an automated Bioscreen C MBR (Growth Curves USA) reader at 30-
minute intervals was plotted to obtain growth curves.

Cell pellet was used for DNA extraction for qPCR analysis. Each sample was treated with
500µl Buffer ATL (Qiagen) in Lysing Matrix E tubes and homogenized twice at 4 m/s for 30
seconds using the FastPrep-24 (MP Biomedicals). After centrifugation at maximum speed for 5
minutes, 200µl of supernatant was transferred into 2ml EZ1 tubes (Qiagen), mixed with 12µl
Proteinase K, vortexed, and incubated at 56°C for 15 min. DNA was extracted in EZ1 Advanced
XL Instrument (Qiagen) using EZ1 DNA Tissue Kit (Qiagen) and eluted in 50µl EB buffer. DNA
concentration was determined using a Qubit 4 Fluorometer with the Qubit dsDNA BR Assay Kit
(Invitrogen). qPCR reaction mixture was prepared following the instructions in the PowerSYBR
Green MasterMix protocol (Applied Biosystems, Cat. No. 4368577) and 1µl of the extracted DNA
was used as the DNA template. Reactions were run in triplicates on a 96-well plate in ViiA 7 Real-
Time PCR System (Applied Biosystems) with the following cycling conditions: 95°C for 2 min;
40 cycles of 95°C for 15 s, 60°C for 1 min; followed by a dissociation step. Standard curves were
generated using gBlocks (IDT) diluted to 10^8–10^3 copies/µl. Primer and gBlock sequences used
for the qPCR are listed in **Supplementary File 17**.

Effect of histidine on *Micrococcus luteus* growth

The effect of histidine on *Micrococcus luteus* was assessed by culturing *M. luteus* in Brain heart
infusion (BHI) broth with and without L-histidine supplementation (50µg/ml concentration), with
BHI used as a negative control. Growth curve experiments were performed at 37°C for 6 days at
high amplitude, normal speed shaking in a 100-well Honeycomb plate (Growth Curves USA) with
a working volume of 200µl in each well. All the experiments were performed in triplicates. Optical
Density (OD) at 600nm measured using an automated Bioscreen C MBR (Growth Curves USA)
reader at 30-minute intervals was plotted to obtain growth curves.

Metabolomics sample collection and processing

An additional cohort of 100 subjects was recruited and skin samples collected in a separate sub-
study of the HELIOS study for skin metagenomic and metabolomic analysis (Ac, elbow crease).
For metabolomics, D-Squame (D100, Clinical and Derm) skin tapes were collected in the same
manner as for metagenomics, except that they were pushed deep into their collection tube and
processed by adding 1 ml of extraction buffer per tube, consisting of 0.1% formic acid in water,

and 0.1µg/ml heavy isotope labelled standards. The tubes were then water bath sonicated (Elma)

for 15 minutes at 4°C. Following sonication, 120 µl from each sample was loaded onto 96 well

chimney plates (Greiner Bio-One) prior to acquisition by liquid chromatography mass

spectrometry (LCMS). An additional 150µl was used to determine the protein concentration with

the Micro BCA kit (Thermo Scientific). 96 well plates holding the samples were loaded into the

Vanquish LC system (Thermo Scientific), utilizing the Atlantis Premier BEH C18 AX 1.7 µm,

2.1×100 mm column (Waters) before being acquired on the TSQ Quantis triple quadrupole MS

machine (Thermo Scientific). All data files were analysed via the software Skyline. No heavy

isotope labelled standard were available for trans-UCA, so the cis-UCA heavy isotope labelled

standard was used as a proxy. Output values from each sample were then normalized against their

respective Micro BCA derived protein concentrations.

Dermotype associations with host attributes

Samples with unstable dermotype assignments (stability<90%) were excluded for each skin site;

additionally, subjects with highly stable yet discordant dermotypes at the same site were excluded,

to enhance consistency. Associations between dermotypes and demographic or physiological data

was assessed in two stages. In the first stage, categorical variables were analyzed using Fisher's

exact test, while continuous variables were evaluated using Wilcoxon tests. Only attributes that

demonstrated overall statistical significance were carried forward to the second stage. In the

second stage, categorical variables were analyzed using Fisher's exact test in a one-versus-the-rest

framework, with odds ratios recorded as effect sizes; for continuous variables, pairwise Wilcoxon

tests were performed, and rank-biserial correlations were reported as effect sizes.

Associations between dermotypes and questionnaire responses were assessed using logistic

regression models with a one-versus-rest approach to accommodate multi-label cases (e.g., with

five axilla dermotypes), with Beta coefficients recorded as effect sizes. When dermotypes

exhibited significant associations with demographic factors in prior analyses (**Figure 4A**), these

factors were included as covariates in the regression model. We prioritized responses relevant to

the body site (e.g., underarm product usage for axilla dermotypes), and included response groups

with at least five observations to maintain statistical power. Questions related to specific product

name were excluded.

Machine learning-based dermotype classification

To ensure comparability across questionnaire responses, all continuous variables (e.g., itch
severity) were z-standardized, and categorical variables were one-hot encoded. Binary-class
machine learning classifiers were trained to predict the dermotype of the sample based on its own
metagenomic profile. A one-vs-rest strategy was employed to accommodate multi-label
classification (i.e., axilla dermotype classification). To address class imbalance, we employed the
Synthetic Minority Over-sampling Technique (SMOTE) to up-sample minority class instances[99].
Samples with unstable clustering membership (stability<90%) were excluded from each skin site.

We selected six machine learning methods based on their distinct properties, such as linear
and non-linear capabilities. These included decision trees, logistic regression, random forests,
support vector classifiers[100] (linear and gaussian kernel) and XGBoost[101]. Model training and
performance reporting utilized a nested cross-validation approach to provide an unbiased
performance estimate by accounting for hyperparameter selection—this involved 4-fold training
with 3-fold inner cross-validation for hyperparameter tuning. Tuned models were evaluated on a
withheld 1-fold test, and the process was repeated five times with different data splits. Mean
accuracy and AUC-ROC scores were reported (one-vs-rest, macro average). Using the same
framework as described above, we trained additional dermotype classifiers with alternative input
features, including: 1) the sample's metabolic pathway profile, 2) metagenomic profiles from other
skin sites within the same host. Further analyses were based on the best-performing methods:
random forests for models using metagenomic profiles, and logistic regression for models using
metabolic pathway profiles.

Random forest classifiers were further trained to assess associations between host factors
and dermotypes using host attributes (i.e., demographics, skin physiological parameters, and
questionnaire data), as they provided the best performance. Only subjects with concordant
dermotype labels between bilateral sites were included. Lastly, to determine the relative
informativeness of different feature sets for dermotype classification at a given skin site, we trained
an inclusive ridge classifier using three broad categories of features: demographic factors,
conversed metabolic pathway profiles, and dermotype labels from other skin sites. Categorical
values were one-hot encoded and continuous values z-standardized.

Commensal *Staphylococcus* strain isolation

Commensal *Staphylococcus* strains were isolated from skin swabs for 4 healthy donors across 4
separate body sites: antecubital fossa, axilla, cheek and scalp. Swabs were immersed in 2.5mL

BHI broth (Oxoid), incubated at 37°C at 210rpm for 2 hours before being spread onto TSA-SB

(Thermo Fisher Scientific) and Baird Parker (Oxoid) plates and incubated at 37°C for 24 hours.

Colonies were then picked and grown in 4ml BHI broth for 16 hours overnight at 230rpm and

37°C. 200μL of the cultures were spun down at 5000rpm for 5 minutes. To extract gDNA the pellet

was resuspended in 50μL QuickExtract (Lucigen). Bacterial suspensions were then heated to 65°C

for 6 minutes, briefly vortexed, then heated to 98°C for 4 minutes and briefly vortexed again. PCR

reactions were setup with 10μL KAPA SYBR FAST (Merck), 6.4μL nuclease free water

(Promega), 3μL extracted gDNA and 0.6μL primer mix. Primer mixes were specific for 4 specific

Staphylococcal species (**Supplementary File 15**). PCR was run on an AriaMx Real Time PCR

System (Agilent Technologies). PCR products were sent for Sanger sequencing to confirm product

sequences. The identified *Staphylococcal* colonies were inoculated into 4ml BHI broth and grown

overnight for 16 hours at 230rpm and 37°C. Optical density (OD) was measured with a

SpectraMax M5 Microplate Reader (Molecular Devices). Overnight cultures were then grown

from OD 0.1 in fresh BHI for 26 hours at 230rpm and 37°C. Final OD measurements for all strains

after 26 hours was normalized to OD 2.75. Bacterial cultures were spun down at 5000rpm for 5

minutes, pellets were discarded and supernatants stored at 4°C.

Human keratinocyte cell culture and Nano-Glo HiBiT

A genetically modified N/TERT keratinocyte cell-line containing a 33 nucleotide HiBiT tag

directly adjacent to the IL33 gene, was utilized for Nano-Glo HiBiT experiments. Keratinocytes

were cultured in keratinocyte serum free medium (KSFM; Gibco) supplemented with bovine

pituitary extract (BPE) at a final concentration of 20μg/mL, epidermal growth factor (EGF) at a

final concentration of 0.2ng/mL, calcium chloride at a final concentration of 300μM and 1:1000

penicillin-streptomycin (pen-strep; Gibco). Keratinocytes were seeded onto 96-well cell culture

plates (Greiner) at a density of 20,000 cells per well and grown for 24 hours in 100μL KSFM.

After 24 hours, KSFM was removed and the keratinocytes were cultured in 100μL KSFM with

either 0.2% TritonX (Merck), 5% BHI, or 5% bacterial supernatants. Keratinocytes were cultured

overnight for 16 hours in treatment conditions. After 16 hours of treatment on the HiBiT N/TERT

keratinocytes, 50μL of the KSFM was transferred to a white-bottom 96-well plate (Sigma-Aldrich)

and mixed with 50μL Nano-Glo HiBiT extracellular reagent. The remaining 50μL of KSFM was

retained on the cell culture plate with the cells and mixed with 50μL CellTitre-Glo reagent. Both

plates were mixed on an orbital shaker for 1 minute, and incubated for 10 minutes at room
temperature before luminescence was measured with a SpectraMax M5 Microplate Reader.

Data availability

Shotgun metagenomic sequencing data is available from the European Nucleotide Archive (ENA,
https://www.ebi.ac.uk/ena/browser/home) under project accession number PRJEB87291.
Demographic data and skin physiological measurements can be found in **Supplementary File 18**.
Large datasets and metabolomics data are available on Figshare at
https://figshare.com/projects/ASMP_Dermotype/243932. The authors of this study do not own the
rights to the HELIOS dataset, and this dataset is under controlled access to ensure good data
governance, responsible data use, and that the dataset is only used for the intended research
purposes in compliance with HELIOS study cohort IRB and ethics approval. Access to individual
level data for study participants should be submitted to the HELIOS Data Access Committee
(email: helios_science@ntu.edu.sg).

Code availability

Source code for scripts used to analyze the data are available in a GitHub project at
https://github.com/CSB5/ASMP_Dermotype.

Acknowledgments

The work is supported by funding for the Asian Skin Microbiome Programme (Industry Alignment
Fund Pre-Positioning – IAF-PP; H18/01/a0/16) and Asian Skin Microbiome Programme 2.0 (IAF-
PP H22/J1/a0/040) from the National Research Foundation, Singapore. Additional support is
provided by HELIOS study (IRB No. 2016-11-030), which is supported by Singapore Ministry of
Health's (MOH) National Medical Research Council (NMRC) under its OF-LCG funding scheme
(MOH-000271-00), Singapore Translational Research (STaR) funding scheme
(NMRC/StaR/0028/2017), the National Research Foundation (Singapore) through the Singapore
MOH NMRC and the Precision Health Research (PRECISE) under the National Precision
Medicine programme (NMRC/PRECISE/2020), National Cohorts Office through the Singapore
MOH NMRC (P2022-01-03 and P2022-02-03) and intramural funding from Nanyang
Technological University, Lee Kong Chian School of Medicine and the National Healthcare
Group. The HELIOS study is also supported by an outstanding team of administrative and

operational staff (**Supplementary File 16**). The computational work for this article was partially

performed on resources of the National Supercomputing Centre, Singapore (https://www.nscc.sg)

and the A*STAR Computational Resource Center. We thank all investigators, staff members and

study participants who made HELIOS and ASMP studies possible.

Competing interests

The authors declare no competing interests.

Table 1. Characteristics of study participants (n=200)

Demographics	
Age (years, median [IQR])	49.00 [40.00, 57.00]
Gender Male (%)	96 (48.0)
Ethnicity (%)	
Chinese	120 (60.0)
Indian	40 (20.0)
Malay	40 (20.0)
Monthly household income (SGD) (%)	
< 6K	79 (39.5)
≥ 6K	102 (51.0)
Unknown	19 (9.5)
Current smoker (%)	0 (0)
Skin physiological measurements	
Forearm	
Surface hydration (median [IQR])	30.02 [23.18, 39.00]
pH (median [IQR])	4.67 [4.38, 5.14]
TEWL[1] (median [IQR])	4.60 [3.67, 5.34]
Cheek	
TEWL[1] (median [IQR])	13.18 [11.64, 15.27]
Sebum (median [IQR])	7.00 [2.00, 19.25]
Skin health and conditions	
Itchy skin (%)	59 (29.5)
Sensitive skin (%)	69 (34.5)
Body odor (%)	44 (22.0)
Antibiotic usage in the last month (%)	12 (6.0)
Topical antimicrobial usage in the last month (%)	24 (12.0)
History of eczema (%)	44 (22.0)
Acne (%)	17 (8.5)
Psoriasis (%)	5 (2.5)
Vitiligo (%)	2 (1.0)
Skin infection[2] (%)	18 (9.0)
Skin cancer (%)	0 (0)
Alopecia areata (%)	3 (1.5)

976   [1]TEWL = Transepidermal water loss

977   [2]Skin infection includes bacterial infection, fungal infection (at feet, body, and groin), viral

978   warts, scabies, chronic hives and ulcer wounds

     Figure Legends

Figure 1 | Heterogeneity of site-specific skin microbial community compositions within and

across individuals. A. Overview of skin sampling strategy, illustrating bilateral sampling at nine

sites with swabs for Ps and Ax and tapes for others. **B.** Boxplots quantifying the similarity of the

skin microbiome within (left versus right) and between individuals across sites. Sites in the inset

figure for alpha diversity are presented in the same order. All p-values were calculated using the

one-sided Wilcoxon rank-sum test (* denotes BH-adjusted p-value<0.001). Boxplot whiskers

represent 1.5× interquartile range or the maximum/minimum data point within the range. **C.** Left

barplot comparing the percentage of subjects with their hair covered daily between those with

concordant and discordant parietal scalp microbiomes between left and right body sites; right

boxplot showing that individuals with discordant forehead microbiomes have significantly higher

itch severity. **D**. Perspective kernel density plot of Ax microbiome PCoA results with the five red

dots representing local maxima determined via Mean Shift analysis. **E.** Averaged relative

abundance plots of major species ranked by median abundance for each dermotype or site, with

the number of stable samples in each site and dermotype shown at the bottom. **F.** Heatmap showing

significant correlations for dermotype labels across sites. The color gradient denotes log10-

transformed BH-adjusted Fisher's exact test p-values, and odds ratios are annotated in the boxes

where appropriate.

Figure 2 | Dermotypes are characterized by distinct core and ancillary species. A. Overview

of the general trend relating species prevalence and abundances (lower leg microbiomes as an

example), and the schema to categorize species into *core species* (≥50% prevalence, in blue) and

ancillary species (≥3 occurrences, <10% prevalence, ≥95th percentile non-zero median abundance,

in orange). **B.** Core species distribution by site and dermotype, with sizes corresponding to

prevalences and color intensities corresponding to non-zero median abundance. Circles with red

boundaries mark species with <0.5% median abundance. **C.** Left: Ancillary species distribution

and non-zero median abundance at each site and dermotype. Right: Boxplots showing the intra-

individual occurrence and median abundance of each ancillary species, with the number of

individuals indicated on the right (each overlaid point represents one subject). The x-axis shows

the number of sites (out of nine) where a given ancillary species is observed, and the color indicates

its within-subject median abundance. **D.** Stacked barcharts depicting relative abundance profiles

for selected subjects with blooms of *HPyV6* across multiple sites. **E.** Significant inter-site co-
abundance patterns of *HPyV6,* with the color gradient denoting log10-transformed BH-adjusted
Fisher's exact test p-values.

Figure 3 | Dermotypes exhibit varied species co-abundance patterns and interactions, and
present different metabolic niches. A. Species co-abundance networks for dermotypes in the
axilla and volar forearm, respectively. Nodes represent microbial species, sized by relative
abundance and colored by microbial genera (names are bolded for hub species). Squares denote
species typically present in oral cavities. Edges denote significant Spearman correlations
(CCREPE $|r| \geq 0.3$, p-value<0.05): purple for positive and orange for negative correlations. Thin
lines indicate shared edges, while thick solid and thick dashed lines denote differential edges with
$|r| \geq 0.5$ and $0.3 \leq |r| < 0.5$, respectively. **B.** Representative agar-based antagonism co-culture assay
results showing *S. epidermidis* and *M. luteus* interactions with *S. hominis* under hypoxic and
aerobic conditions. **C.** Presence/absence patterns for hub species identified from species co-
abundance networks across dermotypes. **D.** Selected degradation-related metabolic pathways that
are differentially abundant across dermotypes. **E-F.** Boxplots depicting the variation in
community-level relative abundance of, (**E**) Lactose and galactose degradation I pathway, and (**F**)
L-histidine degradation I pathway between two elbow crease dermotypes (Ac-1 & Ac-2), with
corresponding HUMAnN2-derived species contributions summarized in stacked barcharts. P-
values for comparisons across dermotypes were calculated using a one-sided Wilcoxon rank-sum
test (∗∗∗∗ denotes p−value<0.0001), and boxplot whiskers represent 1.5× interquartile range or the
maximum/minimum data point.

Figure 4 | Comprehensive investigation of host attributes, same-site and cross-site intrinsic-
microbiome factors for their ability to predict dermotype labels. A. Heatmap depicting
associations between dermotypes and host demographic factors. Effect sizes are represented as
odds ratios (gender, ethnicity; Fisher's exact test) or rank-biserial correlations (age;
Wilcoxon test). **B.** Boxplots displaying variations in skin physiological measurements across
dermotypes. **C.** Heatmap of beta coefficients from one-versus-rest logistic regression models, with
corresponding p-value labeled on each tile, showing associations between dermotypes and selected
personal care behaviors and skin conditions. **D.** AUC-ROC values for dermotype classifiers using
random forest models trained on multi-domain host attributes. The left panel compares feature

importance of various demographic factors for the Ac dermotype classifier. **E.** Boxplots depicting AUC-ROC values for ridge classifiers trained on different categories of metabolic pathways. **F.** Top boxplots comparing AUC-ROC values for cross-site random forest classifiers trained on the full taxonomic profile versus top 10 abundant species from a single site, with the bottom heatmap showing feature importance for the top 10 species. **G.** Top barplot showing AUC-ROC values for the best-performing cross-site dermotype classifiers trained on dermotype labels from other skin sites, with the heatmap displaying combinations of dermotype labels used for training. **H.** Feature importance comparison in an inclusive ridge classifier trained on host demographic features, cross-site dermotype labels, and conserved metabolic pathway profiles. For inter-group comparisons of continuous variables (e.g., skin physiology, AUC-ROC, feature importance), p-values were calculated using a Wilcoxon rank-sum test, and boxplot whiskers represent 1.5× interquartile range or the maximum/minimum data point. For all subfigures *, **, *** and **** denote p−value<0.05, 0.01, 0.001 and 0.0001, respectively.

References

1. Oh, J. *et al.* Biogeography and individuality shape function in the human skin metagenome. *Nature* **514**, 59–64 (2014).

2. Perez, G. I. P. *et al.* Body Site Is a More Determinant Factor than Human Population Diversity in the Healthy Skin Microbiome. *PLOS ONE* **11**, e0151990 (2016).

3. Byrd, A. L., Belkaid, Y. & Segre, J. A. The human skin microbiome. *Nat Rev Microbiol* **16**, 143–155 (2018).

4. Roth, R. R. & James, W. D. Microbial Ecology of the Skin. *Annu. Rev. Microbiol.* **42**, 441–464 (1988).

5. Oh, J., Byrd, A. L., Park, M., Kong, H. H. & Segre, J. A. Temporal Stability of the Human Skin Microbiome. *Cell* **165**, 854–866 (2016).

6. Claesen, J. *et al.* A Cutibacterium acnes antibiotic modulates human skin microbiota composition in hair follicles. *Sci Transl Med* **12**, eaay5445 (2020).

7. Nakatsuji, T. *et al.* Antimicrobials from human skin commensal bacteria protect against Staphylococcus aureus and are deficient in atopic dermatitis. *Sci Transl Med* **9**, eaah4680 (2017).

8. Bomar, L., Brugger, S. D., Yost, B. H., Davies, S. S. & Lemon, K. P. Corynebacterium accolens Releases Antipneumococcal Free Fatty Acids from Human Nostril and Skin Surface Triacylglycerols. *mBio* **7**, e01725-01715 (2016).

9. Iwase, T. *et al.* Staphylococcus epidermidis Esp inhibits Staphylococcus aureus biofilm formation and nasal colonization. *Nature* **465**, 346–349 (2010).

10. Lee, H. J. *et al.* Effects of cosmetics on the skin microbiome of facial cheeks with different hydration levels. *Microbiologyopen* **7**, e00557 (2017).

11. Bouslimani, A. *et al.* The impact of skin care products on skin chemistry and microbiome

dynamics. *BMC Biol* **17**, 47 (2019).

12. Murphy, B. *et al.* Alteration of barrier properties, stratum corneum ceramides and

microbiome composition in response to lotion application on cosmetic dry skin. *Sci Rep* **12**,

5223 (2022).

13. Fyolek, J. *et al.* Modulation of the Skin Microbiome Induced by High-Fat Diet in Children

with Food Allergies and Eczema. *Journal of Allergy and Clinical Immunology* **153**, AB66

(2024).

14. Wang, L. *et al.* Facial Skin Microbiota-Mediated Host Response to Pollution Stress Revealed

by Microbiome Networks of Individual. *mSystems* **6**, e00319-21.

15. van Mierlo, M. M. F. *et al.* The influence of treatment in alpine and moderate maritime

climate on the composition of the skin microbiome in patients with difficult to treat atopic

dermatitis. *Clinical & Experimental Allergy* **49**, 1437–1445 (2019).

16. Leung, M. H. Y. *et al.* Changes of the human skin microbiota upon chronic exposure to

polycyclic aromatic hydrocarbon pollutants. *Microbiome* **8**, 100 (2020).

17. Li, Z. *et al.* Characterization of the human skin resistome and identification of two

microbiota cutotypes. *Microbiome* **9**, 47 (2021).

18. Leung, M. H. Y. *et al.* Skin microbiome differentiates into distinct cutotypes with unique

metabolic functions upon exposure to polycyclic aromatic hydrocarbons. *Microbiome* **11**,

124 (2023).

19. Arumugam, M. *et al.* Enterotypes of the human gut microbiome. *Nature* **473**, 174–180

(2011).

20. Wu, G. D. *et al.* Linking Long-Term Dietary Patterns with Gut Microbial Enterotypes.

Science **334**, 105–108 (2011).

21. Costea, P. I. *et al.* Enterotypes in the landscape of gut microbial community composition.

Nat Microbiol **3**, 8–16 (2018).

22. Frioux, C. *et al.* Enterosignatures define common bacterial guilds in the human gut

microbiome. *Cell Host Microbe* **31**, 1111-1125.e6 (2023).

23. Ravel, J. *et al.* Vaginal microbiome of reproductive-age women. *Proc Natl Acad Sci U S A*

108, 4680–4687 (2011).

24. Gajer, P. *et al.* Temporal Dynamics of the Human Vaginal Microbiota. *Sci Transl Med* **4**,

132ra52 (2012).

25. De Seta, F., Campisciano, G., Zanotta, N., Ricci, G. & Comar, M. The Vaginal Community

State Types Microbiome-Immune Network as Key Factor for Bacterial Vaginosis and

Aerobic Vaginitis. *Front Microbiol* **10**, 2451 (2019).

26. France, M. T. *et al.* VALENCIA: a nearest centroid classification method for vaginal

microbial communities based on composition. *Microbiome* **8**, 166 (2020).

27. Lebeer, S. *et al.* A citizen-science-enabled catalogue of the vaginal microbiome and

associated factors. *Nat Microbiol* **8**, 2183–2195 (2023).

28. Tay, A. S. L. *et al.* Atopic dermatitis microbiomes stratify into ecologic dermotypes enabling

microbial virulence and disease severity. *Journal of Allergy and Clinical Immunology* **147**,

1329–1340 (2021).

29. Alekseyenko, A. V. *et al.* Community differentiation of the cutaneous microbiota in

psoriasis. *Microbiome* **1**, 31 (2013).

30. Sun, C. *et al.* Integrated analysis of facial microbiome and skin physio-optical properties

unveils cutotype-dependent aging effects. *Microbiome* **12**, 163 (2024).

31. Wang, X. *et al.* The Health for Life in Singapore (HELIOS) Study: delivering Precision

Medicine research for Asian populations. 2024.05.14.24307259 Preprint at

https://doi.org/10.1101/2024.05.14.24307259 (2024).

32. Liu, Y. *et al.* An optimal method for quantifying the facial sebum level and characterizing

facial sebum features. *Skin Res Technol* **29**, e13454 (2023).

33. Akdeniz, M., Gabriel, S., Lichterfeld-Kottner, A., Blume-Peytavi, U. & Kottner, J.

Transepidermal water loss in healthy adults: a systematic review and meta-analysis update.

Br J Dermatol **179**, 1049–1055 (2018).

34. Grice, E. A. *et al.* Topographical and Temporal Diversity of the Human Skin Microbiome.

Science **324**, 1190–1192 (2009).

35. Chng, K. R. *et al.* Whole metagenome profiling reveals skin microbiome-dependent

susceptibility to atopic dermatitis flare. *Nat Microbiol* **1**, 16106 (2016).

36. Yue, J. C. & Clayton, M. K. A Similarity Measure Based on Species Proportions.

Communications in Statistics - Theory and Methods **34**, 2123–2131 (2005).

37. Theelen, B. *et al.* Malassezia ecology, pathophysiology, and treatment. *Medical Mycology*

56, S10–S25 (2018).

38. Gupta, A. K., Batra, R., Bluhm, R., Boekhout, T. & Dawson, T. L. Skin diseases associated

with *Malassezia* species. *Journal of the American Academy of Dermatology* **51**, 785–798

(2004).

39. Fritz, S. A. *et al.* Staphylococcus aureus Colonization in Children With Community-

Associated Staphylococcus aureus Skin Infections and Their Household Contacts. *Archives*

of Pediatrics & Adolescent Medicine **166**, 551–557 (2012).

40. Zhou, W. *et al.* Host-specific evolutionary and transmission dynamics shape the functional

diversification of Staphylococcus epidermidis in human skin. *Cell* **180**, 454-470.e18 (2020).

41. Chia, M. *et al.* Shared signatures and divergence in skin microbiomes of children with atopic

dermatitis and their caregivers. *Journal of Allergy and Clinical Immunology* **150**, 894–908

(2022).

42. Han, G., Luong, H. & Vaishnava, S. Low abundance members of the gut microbiome exhibit

high immunogenicity. *Gut Microbes* **14**, 2104086.

43. Zhou, W. *et al.* Host-Specific Evolutionary and Transmission Dynamics Shape the

Functional Diversification of *Staphylococcus epidermidis* in Human Skin. *Cell* **180**, 454-

470.e18 (2020).

44. Oh, J. *et al.* The altered landscape of the human skin microbiome in patients with primary

immunodeficiencies. *Genome Res* **23**, 2103–2114 (2013).

45. Zhou, X. *et al.* Longitudinal profiling of the microbiome at four body sites reveals core

stability and individualized dynamics during health and disease. *Cell Host & Microbe* **32**,

506-526.e9 (2024).

46. Christensen, G. J. M. *et al.* Antagonism between Staphylococcus epidermidis and

Propionibacterium acnes and its genomic basis. *BMC Genomics* **17**, 152 (2016).

47. Nakamura, K. *et al.* Short chain fatty acids produced by Cutibacterium acnes inhibit biofilm

formation by Staphylococcus epidermidis. *Sci Rep* **10**, 21237 (2020).

48. Ahle, C. M. *et al.* Interference and co-existence of staphylococci and Cutibacterium acnes

within the healthy human skin microbiome. *Commun Biol* **5**, 1–14 (2022).

49. Kwaszewska, A., Sobiś-Glinkowska, M. & Szewczyk, E. M. Cohabitation—relationships of

corynebacteria and staphylococci on human skin. *Folia Microbiol* **59**, 495–502 (2014).

50. Nakazono, K. *et al.* Complete sequences of epidermin and nukacin encoding plasmids from

oral-derived Staphylococcus epidermidis and their antibacterial activity. *PLoS One* **17**,

e0258283 (2022).

51. Huang, T. Y., Jiang, Y. E. & Scott, D. A. Culturable bacteria in the entire acne lesion and

short-chain fatty acid metabolites of *Cutibacterium acnes* and *Staphylococcus epidermidis*

isolates. *Biochemical and Biophysical Research Communications* **622**, 45–49 (2022).

52. Mark Welch, J. L., Rossetti, B. J., Rieken, C. W., Dewhirst, F. E. & Borisy, G. G.

Biogeography of a human oral microbiome at the micron scale. *Proceedings of the National*

Academy of Sciences **113**, E791–E800 (2016).

53. Radaic, A. & Kapila, Y. L. The oralome and its dysbiosis: New insights into oral

microbiome-host interactions. *Computational and Structural Biotechnology Journal* **19**,

1335–1360 (2021).

54. Gevers, D. *et al.* The Treatment-Naive Microbiome in New-Onset Crohn's Disease. *Cell*

Host & Microbe **15**, 382–392 (2014).

55. Yachida, S. *et al.* Metagenomic and metabolomic analyses reveal distinct stage-specific

phenotypes of the gut microbiota in colorectal cancer. *Nat Med* **25**, 968–976 (2019).

56. Chen, Y. E., Fischbach, M. A. & Belkaid, Y. Skin microbiota–host interactions. *Nature* **553**,

427–436 (2018).

57. Joglekar, P. *et al.* Integrated genomic and functional analyses of human skin–associated

Staphylococcus reveal extensive inter- and intra-species diversity. *Proc. Natl. Acad. Sci.*

U.S.A. **120**, e2310585120 (2023).

58. Kaiser, J. C. & Heinrichs, D. E. Branching Out: Alterations in Bacterial Physiology and

Virulence Due to Branched-Chain Amino Acid Deprivation. *mBio* **9**, e01188-18 (2018).

59. Kaiser, J. C. *et al.* Repression of branched-chain amino acid synthesis in Staphylococcus

aureus is mediated by isoleucine via CodY, and by a leucine-rich attenuator peptide. *PLOS*

Genetics **14**, e1007159 (2018).

60. Roux, P.-F., Oddos, T. & Stamatas, G. Deciphering the Role of Skin Surface Microbiome in

Skin Health: An Integrative Multiomics Approach Reveals Three Distinct Metabolite–

Microbe Clusters. *Journal of Investigative Dermatology* **142**, 469-479.e5 (2022).

61. Danzberger, J. *et al.* Glycan distribution and density in native skin's stratum corneum. *Skin*

Res Technol **24**, 450–458 (2018).

62. Li, Z. *et al.* Integrated Human Skin Bacteria Genome Catalog Reveals Extensive Unexplored

Habitat-Specific Microbiome Diversity and Function. *Advanced Science* **10**, 2300050 (2023).

63. Kezic, S. *et al.* Levels of filaggrin degradation products are influenced by both filaggrin

genotype and atopic dermatitis severity. *Allergy* **66**, 934–940 (2011).

64. Hug, D. H., Dunkerson, D. D. & Hunter, J. K. The degradation of l-histidine and *trans* and

cis-urocanic acid by bacteria from skin and the role of bacterial *cis*-urocanic acid isomerase.

Journal of Photochemistry and Photobiology B: Biology **50**, 66–73 (1999).

65. Walterscheid, J. P. *et al.* Cis-urocanic acid, a sunlight-induced immunosuppressive factor,

activates immune suppression via the 5-HT2A receptor. *Proceedings of the National*

Academy of Sciences **103**, 17420–17425 (2006).

66. Tauch, A., Fernández-Natal, I. & Soriano, F. A microbiological and clinical review on

Corynebacterium kroppenstedtii. International Journal of Infectious Diseases **48**, 33–39

(2016).

67. Liu, B. *et al.* IL-33/ST2 signaling excites sensory neurons and mediates itch response in a

mouse model of poison ivy contact allergy. *Proceedings of the National Academy of*

Sciences **113**, E7572–E7579 (2016).

68. Trier, A. M. *et al.* IL-33 signaling in sensory neurons promotes dry skin itch. *Journal of*

Allergy and Clinical Immunology **149**, 1473-1480.e6 (2022).

69. Jeffery, I. B., Claesson, M. J., O'Toole, P. W. & Shanahan, F. Categorization of the gut

microbiota: enterotypes or gradients? *Nat Rev Microbiol* **10**, 591–592 (2012).

70. Knights, D. *et al.* Rethinking "Enterotypes". *Cell Host Microbe* **16**, 433–437 (2014).

71. Lam, T. H. *et al.* Understanding the microbial basis of body odor in pre-pubescent children

and teenagers. *Microbiome* **6**, 213 (2018).

72. Urban, J. *et al.* The effect of habitual and experimental antiperspirant and deodorant product

use on the armpit microbiome. *PeerJ* **4**, e1605 (2016).

73. Spittaels, K.-J., Ongena, R., Zouboulis, C. C., Crabbé, A. & Coenye, T. Cutibacterium acnes

Phylotype I and II Strains Interact Differently With Human Skin Cells. *Front Cell Infect*

Microbiol **10**, 575164 (2020).

74. Conwill, A. *et al.* Anatomy promotes neutral coexistence of strains in the human skin

microbiome. *Cell Host & Microbe* **30**, 171-182.e7 (2022).

75. Severn, M. M. & Horswill, A. R. Staphylococcus epidermidis and its dual lifestyle in skin

health and infection. *Nat Rev Microbiol* **21**, 97–111 (2023).

76. Evans, S. M., Schrlau, A. E., Chalian, A. A., Zhang, P. & Koch, C. J. Oxygen Levels in

Normal and Previously Irradiated Human Skin as Assessed by EF5 Binding. *Journal of*

Investigative Dermatology **126**, 2596–2606 (2006).

77. Wong, W. J., Richardson, T., Seykora, J. T., Cotsarelis, G. & Simon, M. C. Hypoxia-

Inducible Factors Regulate Filaggrin Expression and Epidermal Barrier Function. *Journal of*

Investigative Dermatology **135**, 454–461 (2015).

78. Cheung, G. Y. C., Joo, H.-S., Chatterjee, S. S. & Otto, M. Phenol-soluble modulins – critical

determinants of staphylococcal virulence. *FEMS microbiology reviews* **38**, 698 (2014).

79. Ch'ng, J.-H. *et al.* Heme cross-feeding can augment *Staphylococcus aureus* and

Enterococcus faecalis dual species biofilms. *The ISME Journal* **16**, 2015–2026 (2022).

80. Regina, V. R. *et al.* Decoding scalp health and microbiome dysbiosis in dandruff.

2024.05.02.592279 Preprint at https://doi.org/10.1101/2024.05.02.592279 (2024).

81. Seo, Y., Sung, M., Hwang, J. & Yoon, Y. Minimum Inhibitory Concentration (MIC) of

Propionic Acid, Sorbic Acid, and Benzoic Acid against Food Spoilage Microorganisms in

Animal Products to Use MIC as Threshold for Natural Preservative Production. *Food Sci*

Anim Resour **43**, 319–330 (2023).

82. Pagac, M. P. *et al.* Life stage impact on the human skin ecosystem: lipids and the microbial

community. *NPJ Biofilms Microbiomes* **11**, 13 (2025).

83. Chia, M. *et al.* Skin metatranscriptomics reveals landscape of variation in microbial activity

and gene expression across the human body. 2024.12.02.626500 Preprint at

https://doi.org/10.1101/2024.12.02.626500 (2024).

84. Li, M. *et al.* Microbiome and lipidomic analysis reveal the interplay between skin bacteria

and lipids in a cohort study. *Front Microbiol* **15**, 1383656 (2024).

85. Martin, A. *et al.* A Functional ABCC11 Allele Is Essential in the Biochemical Formation of

Human Axillary Odor. *Journal of Investigative Dermatology* **130**, 529–540 (2010).

86. Pappas, A., Fantasia, J. & Chen, T. Age and ethnic variations in sebaceous lipids.

Dermatoendocrinol **5**, 319–324 (2013).

87. Pierard, G. E., Arrese, J. E., Pierard-Franchimont, C. & DE Doncker, P. Prolonged effects of

antidandruff shampoos - time to recurrence of Malassezia ovalis colonization of skin. *Int J*

Cosmet Sci **19**, 111–117 (1997).

88. Urban, J. *et al.* The effect of habitual and experimental antiperspirant and deodorant product

use on the armpit microbiome. *PeerJ* **4**, e1605 (2016).

89. Grice, E. A. & Dawson, T. L. Host–microbe interactions: *Malassezia* and human skin.

Current Opinion in Microbiology **40**, 81–87 (2017).

90. Ianiri, G., LeibundGut-Landmann, S. & Dawson, T. L. Malassezia: A Commensal, Pathogen,

and Mutualist of Human and Animal Skin. *Annual Review of Microbiology* **76**, 757–782

(2022).

91. Hong, C. *et al.* PathoScope 2.0: a complete computational framework for strain identification

in environmental or clinical sequencing samples. *Microbiome* **2**, 33 (2014).

92. Janssens, Y. *et al.* Disbiome database: linking the microbiome to disease. *BMC Microbiology*

18, 50 (2018).

93. Yao, G. *et al.* MicroPhenoDB Associates Metagenomic Data with Pathogenic Microbes,

Microbial Core Genes, and Human Disease Phenotypes. *Genomics Proteomics*

Bioinformatics **18**, 760–772 (2020).

94. de Goffau, M. C. *et al.* Recognizing the reagent microbiome. *Nat Microbiol* **3**, 851–853

(2018).

95. Peschel, S., Müller, C. L., Von Mutius, E., Boulesteix, A.-L. & Depner, M. NetCoMi:

network construction and comparison for microbiome data in R. *Briefings in Bioinformatics*

22, bbaa290 (2021).

96. Csárdi, G. *et al.* igraph for R: R interface of the igraph library for graph theory and network

analysis. Zenodo https://doi.org/10.5281/ZENODO.7682609 (2024).

97. Faust, K. *et al.* Microbial Co-occurrence Relationships in the Human Microbiome. *PLOS*

Computational Biology **8**, e1002606 (2012).

98. Gill, R., Datta, S. & Datta, S. A statistical framework for differential network analysis from

microarray data. *BMC Bioinformatics* **11**, 95 (2010).

99. Chawla, N. V., Bowyer, K. W., Hall, L. O. & Kegelmeyer, W. P. SMOTE: synthetic

minority over-sampling technique. *J. Artif. Int. Res.* **16**, 321–357 (2002).

100.   Pedregosa, F. *et al.* Scikit-learn: Machine Learning in Python. *Journal of Machine*

Learning Research **12**, 2825–2830 (2011).

101.   Chen, T. & Guestrin, C. XGBoost: A Scalable Tree Boosting System. in *Proceedings of*

the 22nd ACM SIGKDD International Conference on Knowledge Discovery and Data

Mining 785–794 (ACM, San Francisco California USA, 2016).

doi:10.1145/2939672.2939785.

1292

Figure 1

Figure 2

Figure 3

Figure 4