Title: Increased clonality, decreased allele diversity and high genetic structure in tetraploid sea

anemone *Aulactinia stella* populations from North Pacific to Atlantic across the Arctic Ocean

Authors: Ekaterina Bocharova[1,2], Aleksandr Volkov[3], Solenn Stoeckel[4,5]

Affiliations:

[1] Faculty of Life Sciences, University of Vienna, Vienna, Austria

[2] Vavilov Institute of General Genetics of Russian Academy of Sciences, Moscow, Russia

[3] Branch for the freshwater fisheries of VNIRO (VNIIPRKH), Moscow region, Russia

[4] DECOD, L'Institut Agro, IFREMER, INRAE, 35000, Rennes, France

[5] IGEPP, INRAE, Institut Agro, Université de Rennes, Le Rheu, France

Corresponding authors: bocharova.ekaterina@gmail.com and solenn.stoeckel@inrae.fr

Keywords: partial asexuality, population genetics, autopolyploidy, genetic structure, gene flow,

genetical endemism, animal parthenogenesis

Graphical Abstract (photo: Ekaterina Bocharova)

Abstract

Reproductive modes, migration, genetic diversity and local adaptation are important evolutionary traits that limit species' ranges. Understanding the interactions between evolutionary and ecological processes at species' edges provides clues to how species adapt to rapid environmental changes. In the Arctic Ocean, melting sea ice and increasing human activity are bridging coastal populations that were once separated by extreme polar conditions. *Aulactinia stella* is a keystone tetraploid circumpolar sea anemone species that is witnessing these changes. Here, we investigated how reproduction and genetic diversity structured in *A. stella* populations and between adults and juveniles sampled across the Arctic Ocean.

Outside Kamchatka, we only found females or individuals without gonads. Genetic indices and changes in genotype frequencies confirmed that this species reproduces partially by parthenogenesis. Populations on the Atlantic side were highly clonal, while those on the Pacific side reproduced more sexually. Allelic diversity was twice as high in Kamchatka and Kuril populations, suggesting North Pacific coasts being the main last glacial refugia of *A. stella*. Patterns of genetic differentiation between site pairs revealed gene flow routes extending from Kamchatka to Atlantic populations. These routes aligned with ocean currents and reduced sea ice, and not with the past maritime exchanges between Atlantic and Pacific Oceans.

Within each population, only a subset of juvenile genetic diversity, mostly of local origin, was represented among established adults, suggesting local adaptation or kin recognition mechanisms promoting endemism. Juveniles showed less genetic differentiation across the Arctic Ocean, possibly indicating that global changes are favoring juvenile dispersal.

Introduction

The way individuals reproduce is the most influential trait in governing genetic diversity and potentially evolutionary response to environmental changes (Duminil et al., 2007; Pierre et al., 2022). Reproductive mode governs the persistence of DNA variations through generations. It has therefore major impacts on the level of genome-wide genetic diversity and its structuration within and among populations (Ellegren & Galtier 2016). Reproductive modes also profoundly affect other biological traits that play important roles for ecological processes, including population growth, dispersal, colonization of new areas and the genetic basis for neo-adaptation (Yu et al., 2016; He et al., 2024).

Clonality is a dominant reproductive mode found in populations colonizing new area, especially when enduring harsh conditions at the edge of their living range (Cadotte et al., 2006; Roiloa, 2019; Franklin et al., 2021). All groups and kingdoms of Eukaryota domain, including most of the species that are essential for ecosystem functions, reproduce using partial clonality (Avise, 2015; Schön et al., 2009; Tibayrenc et al., 2015; Orive & Krueger-Hadfield, 2021). It consists for each parent to produce at least one descendant that is genetically identical to them at all loci in its genome, with the exception of a few somatic mutations and rarer recombinations (De Meeûs et al., 2007; Stoeckel et al., 2021a). In such species, the proportion of descendants produced by clonality, i.e. the rate of clonality, is a key biological factor to understand their genetic and trait evolution (Yu et al., 2016; Becheler et al., 2017). Clonality tends to increase in populations spreading at range margins by enabling pioneers to settle and maintain without mates (Baker & Stebbins, 1965; Liu et al., 2006; Wang et al., 2019). Moreover, clonality allows genotypes that are successful locally to maintain over generations without recombining with less adapted genotypes and hasten population growth (Fouqueau & Roze, 2021; Pierre et al., 2022). It also decreases the loss of allele diversity and of heterozygosity by genetic drift,

especially in small founding populations (Navascues et al., 2010; Stoeckel & Masson, 2014;

Reichel et al., 2016; Stoeckel et al., 2021b).

The Arctic Ocean has been previously considered as an insurmountable barrier for sessile and

semi-sessile marine species dispersal due to limited access, harsh environmental conditions and

inadequate food resources (Chan et al., 2019). Climate change is opening new routes across the

Arctic Ocean, linking Pacific and Atlantic biomes (Vermeij & Roopnarine, 2008). The

increasing ice melting period during summer may favor the recolonization of multiple species

from their Atlantic and Pacific last glacial refugia. Understanding genetic diversity and

population structure in such species provide useful insights on ecological and evolutionary

processes occurring during range expansion in real and complex conditions (Sax et al., 2007;

Moran & Alexender, 2014; Bridle & Hoffmann, 2022). It also provides a frame of reference to

managers and decision makers to identify priorities and mechanisms to mitigate adverse effects

of global changes on intraspecific diversity and to control future invasive populations (Chan et

al., 2019).

Sea anemones are sessile and semi-sessile organisms reproducing using partial clonality

(Bocharova & Kozevich, 2011; Bocharova, 2016). *Aulactinia stella* (Cnidaria, Anthozoa) is an

intertidal and sublittoral species belonging to Actiniidae family. It was first observed and

described on the eastern coast of the USA in the 19th century (Verrill, 1864), probably in the

southernmost part of its distribution area. Its geographic range is rather circumpolar, along the

coasts of the Arctic Ocean and of the northern parts of Atlantic and Pacific Oceans (Grebelnyi,

1980; Sanamyan & Sanamyan, 1998, 2006). *A. stella* mainly live fixed to rocky ground, even

if it can occasionally be found buried in sandy or muddy soils or hidden in crevices (Bocharova,

2012). In the benthic communities, during its larval stage, *A. stella* feed predating smaller

mollusks, crustaceans, fish larvae and other animals (Bocharova, 2012). First studies of

Aulactinia stella populations in the Arctic Ocean, on the Murman Coast, were carried on about

fifty years ago (Loseva, 1974). As many circumpolar marine species, direct observations on

how populations reproduce and disperse still remain unrealistic. Inferring its important

biological and ecological features using genetic analyses therefore remains the best option to

understand such species.

Developmental and histological evidence suggests that this species may reproduce through

parthenogenesis (Bocharova, 2012). Preliminary analyses based on a limited number of

mitochondrial and ribosomal sequences hypothesized that these populations are likely highly

clonal and exhibit strong genetic structuring across the Arctic Ocean despite the region's

increasing ice melt over recent decades (Bocharova & Mugue, 2012; Bocharova, 2015).

However, while these sequences were suitable for phylogenetic analyses, they lacked the

resolution necessary to confirm clonality and accurately assess the genetic structure of *A. stella*

populations. The recent development of microsatellite primer pairs now makes it possible to

investigate the biology of this circumpolar keystone species more precisely, using approaches

based on genetic diversity and its spatial patterns (Bocharova et al., 2018).

In this study, we aimed to confirm the occurrence of clonal reproduction in *A. stella* populations

in natural environments. We also sought to quantify the extent and variability of clonal

reproduction across the species' range. The low and spatially structured genetic diversity

previously observed from a limited number of sequences highlighted the need for a more robust

assessment of current genetic structure, using neutral and high-resolution markers. This would

also allow us to identify potential pathways of gene flow around the Arctic Ocean.

To investigate clonal reproduction in this species, we analyzed genotypic and genetic indices

across five geographically distinct sites spanning the Arctic Ocean, including locations in the

northern Atlantic and northern Pacific. A total of 315 individuals, representing two age cohorts,

juveniles and adults, were sampled to estimate the contribution of clonal reproduction and its

variation across the species' range. By examining temporal changes in genotypic frequencies

within each site, we aimed to infer the prevalence of clonal reproduction.

We hypothesized that clonal reproduction may facilitate the establishment and persistence of

populations under harsh polar coastal conditions. Conversely, we also considered that the

genetic recombination associated with sexual reproduction may promote the emergence of

genotypes better adapted to local environments. Our findings on reproductive modes are

interpreted in the context of the specific ecological challenges faced by species expanding into

previously unfavorable habitats. In addition, we used the genotyped populations to assess

genetic differentiation and coancestry between sampled sites. Sampling across the full

distribution range allowed us to pinpoint regions with higher genetic diversity, which may

correspond to historical refugia. By comparing genetic differentiation between juveniles and

adults, we aimed to identify present-day gene flow and corresponding ongoing dispersal routes.

Although *A. stella* was first described along the North Atlantic coasts, suggesting a possible

Atlantic origin, previous limited sequence data instead indicated higher genetic diversity in

North Pacific populations. If confirmed, this may suggest a more recent expansion into the

North Atlantic via the Arctic Ocean. This expansion could have been facilitated either by

increased ice melt in recent decades or through anthropogenic means such as maritime traffic.

Under the natural expansion scenario, we would expect a stepping-stone pattern of genetic

structure across the Arctic. In contrast, under the anthropogenic dispersal hypothesis, we would

expect low genetic differentiation between North Pacific and North Atlantic populations,

especially given that the Northern Sea Route has only been regularly used for about thirty years.

Finally, we compared our results with patterns observed in other circumpolar coastal marine

species and interpreted them in light of known ecological dynamics and environmental changes

in these regions.

Materials and methods

Sample collection

In September 2012, we sampled at the low tide in the littoral zone 23 *Aulactinia stella* in the White Sea Biological Station of Moscow State University (WSBS), Kandalaksha Bay of the White Sea and 30 individuals at Liinkhamari, in the Varanger Fjord of the Barents Sea (see Bocharova 2015). We also sampled using SCUBA diving at 5-18 m depth in the sublittoral zone 29 individuals at the Cape Broyle in the Admirals Cove area, Newfoundland, Canada, in June 2017, 14 individuals in the Avachinsky Bay and Viluchinsky Bay of the Pacific coast of Kamchatka Peninsula in June 2010 and June 2016 and 19 individuals in the intertidal zone of the Matua Island of the Kuril Islands in August 2016.

We checked that the phenotypic and biometric appearance of each individual belonged to *Aulactinia stella* species using descriptions in Loseva (1974), Fautin Dunn et al. (1980), Sanamyan and Sanamyan (2008), Sanamyan et al. (2019, 2020) and WoRMS (Rodríguez et al., 2025). In addition, we used nematocyst identification from tentacles, pharynx, body wall and mesentery (Manuel, 1981), and mitochondrial DNA sequence analysis to confirm the species. To analyze genetic diversity in two successive generations, we purposely genotyped large females and the juveniles they brooded in their gastrovascular cavity.

DNA extraction, mtDNA sequencing and genotyping by microsatellite loci

We extracted the total DNA of each sampled adult and juvenile polyp. For each sample, we desiccated and grinded the sampled tissues. We then used the Wizard SV Genomic DNA Purification System (Promega, USA) following the manufacturer's protocol. Extracted DNA was preserved in 96% ethanol before genotyping.

For mitochondrial gene (12S rRNA, 16S rRNA, cytochrome oxidase III) sequencing, we only used adult samples: 23 individuals from the White Sea, 30 from the Barents Sea, 29 from

Newfoundland, 14 from Kamchatka, 19 from the Kuril Islands. We applied the primer sets and

methodology thoroughly described for 16S rRNA and COIII (Geller & Walton, 2001) and 12S

rRNA mitochondrial fragments (Bocharova 2015). Sequencing was performed by 3500 Genetic

Analyzer (Applied Biosystems, USA) using BigDye Terminator v3.1 Cycle Sequencing Kit.

The sequences were analyzed by Geneious Prime software. By concatenating these three

mtDNA sequences (12S, 16S, COIII), we grouped the samples into haplotypes that were aligned

and perfectly matched the five previously described haplotypes (Bocharova, 2015). Including

these mitochondrial haplotypes (Pacific 1-5), we used geneHapR (Zhang et al., 2023) package

to produce the haplotype network (Figure 2). We then used the extracted nuclear DNA to

amplify 41 microsatellites using primer pairs previously developed in Bocharova et. al. (2018)

for this species and another sea anemone species of Actiniidae family, *Cribrinopsis*

albopunctata (Sanamyan & Sanamyan 2006, Sanamyan et al. 2019) with the corresponding

PCR conditions. PCR products were visualized in 3500 Genetic Analyzer (Applied Biosystems,

USA) using POP7 gel polymer. We analyzed the obtained chromatograms using GenMapper

Software (ThermoFisher Scientific, USA). Out of these 41 primer pairs, only ten pairs (Act007,

Act011, Act028, Act067, Act173, Act177, Act235, Act238, Act249, Act386,) successfully

amplified and were polymorphic in our samples. We then multiplexed the ten microsatellites

into four PCR pools using three specific dyes (TAM, ROX, FAM) for visualization: Pool 1)

Act007 (FAM), Act011 (TAM); Pool 2) Act238 (FAM), Act249 (TAM), Act173 (ROX); Pool

3) Act067 (FAM), Act177 (ROX), Act386 (TAM); Pool 4) Act28 (TAM), Act235 (ROX).

To estimate if this marker set was powerful enough to distinguish between individuals, we

computed the overall observed unbiased probability of identity in state under panmixia

(hereafter overall pidu) and the overall observed unbiased probability of identity in state

between sibs (overall pids) as recommended by Waits et al. (2001). To achieve allele dosage

within each individual per locus, we used the dose-effect method with some modifications

(Rodzen & May, 2002; Drauch Schreier et al., 2011). In cases of more than 1 allele, dose-effect

is counted following the instruction (Figure S1): two alleles of almost the same height were

considered as AABB (Figure S1A); One high allele and two low alleles as AABC (Figure S1B);

One very high allele and one low allele as AAAB (Figure S1C); Tree alleles of the same height

were considered as ABC + null allele (Figure S1D); One high allele and one allele half lower

were considered as AAB + null allele (Figure S1E).

Analyses of genetic diversity within spatio-temporal samples

Parents that produce descendants by clonality can result to observe multiple times the same

Multi-Locus Genotype (hereafter MLG) among the genotyped samples (Halkett et al., 2005;

Arnaud-Haond et al., 2007). However, these repeated MLGs may also come from genetic

marker lacking sufficient resolution to disentangle between descendants produced by sexual

reproduction, either under panmixia (the lower probability to observe two identical sexual

descendants) or between mating sibs (the higher probability to observe two identical sexual

descendants). To estimate the power of our marker set to distinguish different individuals and

assess the upper and lower bounds for the probability of observing identical MLG drawing a

pair of individuals sampled in a same population, we therefore computed the probabilities of

identity under panmixia (*pidu*) and considering only sib mating (*pids*; Jacquard, 1968, 1970;

Waits et al., 2001) using formula adapted for tetraploid populations implemented in

GenAPoPop software (Stoeckel et al., 2024).

To assess the level of genetic diversity within populations and compare between populations,

for each age cohort in one site, we reported the sample size (N) and computed the average

number of observed alleles $\bar{N}a$ and its variance Var(Na) over the ten microsatellites, the average

effective number of alleles $\bar{A}_E$ and its variance Var(A_E), the average gene diversity $\bar{H}_E$ and its

variance $Var(H_E)$, the average observed heterozygosity $\bar{H}_O$ and its variance $Var(H_O)$ and the

average value of the inbreeding coefficient $\overline{F_{IS}}$ and its variance $Var(F_{IS})$ over all loci.

We also computed genotypic indices including the number of unique genotypes (G), the

unbiased genotypic richness R (Dorken & Eckert, 2001), the genotypic evenness as Pareto β to

assess the potential uneven contribution of some clonal MLGs to the genotypic composition of

populations (Arnaud-Haond et al., 2007; Bailleul et al., 2016).

We searched for clonal signatures in genetic diversity by analyzing $\overline{F_{IS}}$ and its variance $Var(F_{IS})$

and the overall estimate of the linkage disequilibrium ($\bar{r}_d$, Agapow & Burt, 2001) as commonly

interpreted using one-shoot samples (Halket et al., 2005; Arnaud-Haond et al., 2020; Stoeckel

et al., 2021a). Finally, to assess rates of clonality within spatial sites, we used the Bayesian

estimate of rates of clonality using genotype frequency changes between adult and juvenile at

the same sample site (ClonEstiMate, Becheler et al., 2017 adapted to autpolyploids in

GenAPoPop Stoeckel et al., 2024).

To assess the genetic structure between populations and generations, we computed temporal

and spatial ρST (Ronfort et al., 1998) using GenAPoPop, and plotted the unrooted minimum

spanning tree between individuals using their pairwise number of shared alleles (identity-in-

state) using the classical equal-angle algorithm (Felsenstein, 2004) to picture the coancestry

between individuals across populations at our marker set. To assess the dynamics (speed and

direction) of genetic diversity along sampling sites, we compared juvenile and adult

subpopulations sampled at the same sites and between sites computing the corresponding ρST.

We also used a Discriminant Analysis of Principal Components (DAPC, Jombart et al., 2010)

to investigate the genetic structure of our five sites and tens cohorts. We choose to retain 200

principal component eigenvalues. We used a knee locator approach to find the minimal number

of clusters best structuring the differentiation between individuals and groups. We verify this

value by increasing and decreasing the number of clusters and checking that we obtained the

same groups and distribution of clusters among sites and cohorts.

Statistical analyses

To test if distribution of population genetic indices were significantly different between groups

of sites and groups of generations, we used Mann-Whitney test and reported the median values

of the two groups ($median_{group1}$ versus $median_{group2}$) in the order they are mentioned, the sum

of the rank of the first group (U), the number of measures in each group ($n1=4$ for the cohorts

sampled on the Pacific side of the Arctic Ocean and $n2=6$ for the cohorts sampled on the

Atlantic side) and the probability value (p) of the hypothesis that the two groups have identical

distribution of their genetic indices. To compare ρST between juveniles and adults among pairs

of the same sites, we used Wilcoxon signed rank test. We reported the median ρST value of the

two groups ($median_{group1}$ versus $median_{group2}$) in the order they are mentioned, the sum of the

ranks of the differences above zero (W) and the probability value (p) of the hypothesis that the

two age cohorts have identical distribution of ρST.

All these tests were performed using python 3.11 and Scipy.stats 1.13 (Virtanen et al., 2020).

255 **Results**

We genotyped 314 individuals of Aulactinia stella sampled on five sites, including a total of 75

adults, ranging from 6 to 21 with a median of 15 adults per site, and a total of 239 juveniles,

ranging from 6 to 93 with a median of 49 juveniles per site (Table 1). We found both females

and males in the Kamchatka population, but, while we found only females in the Kuril Islands,

Newfoundland, the Barents and the White Seas. We then only used the adult females sampled

in each site for the following genetic analysis.

We found 15 brooding females in White Sea, 21 in Barents Sea, 14 in Newfoundland, 6 in

Kamchatka and 19 in Kuril Islands sampled sites. The oral disc diameter of mature females

varied from 10 to 45 mm while the overall size of the juveniles released from the gastrovascular

cavities of the adults varied from 1 to 4 mm.

Gene, allele and genotypic diversities in nuclear microsatellites

We found a maximum of four different alleles for each locus in one individual over all our

samples and concluded that *A. stella* must be at least an autotetraploid species. Using our ten

microsatellites, the mean numbers of alleles per locus per site and developmental stage ranged

from 1.2 to 5.7, with a median of 2.75. The mean effective numbers of alleles per locus per site

and developmental stage ranged from 1.19 to 2.53 with a median of 1.69. Over all samples,

probability of identities under panmixia was 8.3×10^{-9} and probability of identity between sibs

was 4.5×10^{-3}. These low probabilities that two individuals generated by sexual reproduction

would result into similar genotypes advocated that the ten microsatellites we used were

powerful enough to genetically distinguish between sexual descendants, even resulting from a

sib mating (Waits et al. 2001). We thus considered afterward that individuals with identical

MLGs were clones (i.e., ramets of a same genet).

Over all sites and cohorts, we found 154 unique MLGs and 26 of these MLGs were found

repeated in more than one sampled individual, i.e. genets with more than one ramet (Table S1).

In Kuril Islands, we genotyped no repeated MLG. In Kamchatka, we found one repeated MLG

(genet) including three ramets. In Barents Sea, we found three repeated MLGs, including

respectively two, five and thirteen ramets. In White Sea, we found eight repeated MLGs

involving two to forty ramets, for a total of 88 ramets. In Newfoundland, we found fourteen

repeated MLGs involving two to eleven ramets, for a total of 49 ramets. All ramets of repeated

MLGs were exclusively sampled on the same site. In all populations with repeated MLGs,

287 repeated MLGs were found both in adults and in juveniles. Some of these genets were even

288 shared by adults and juveniles, except in the Newfoundland samples.

289 Genetic and genotypic diversities showed contrasted measures between the two sites on the

290 Pacific side (i.e., Kuril and Kamchatka) and the three sites on the Atlantic side of the Arctic

291 Ocean (i.e., Newfoundland, Barents and White Seas, Table 1). The four cohorts sampled in

292 Kuril Islands and in Kamchatka showed nearly twice more average gene diversity, ranging from

293 0.372 to 0.460, than found in the six cohorts sampled in Newfoundland, Barents and White

294 Seas, that ranged from 0.097 to 0.266 (0.427 versus 0.174, U=24, n1=4, n2=6, p=0.009), while

295 they both showed similar standard deviation of gene diversity over all loci (U=15, n1=4, n2=6,

296 p=0.610). Cohorts in the Pacific sites also showed significantly higher mean numbers of alleles

297 per locus (4.9 versus 2.0, U=23.5, n1=4, n2=6, p=0.019) and higher effective number of alleles

298 (2.43 versus 1.43, U=24, n1=4, n2=6, p=0.009).

299 Kamchatka and Kuril samples showed higher genotypic richness measured as R than

300 populations on the Atlantic side (1.00 versus 0.37, U=24, n1=4, n2=6, p=0.009) and higher

301 genotypic evenness measured as Pareto β values (3.64 versus 0.45, U=23.5, n1=4, n2=6,

302 p=0.019). Over all loci, Kamchatka and Kuril samples showed lower probabilities of identity

303 under panmixia ($<10^{-6}$ versus 0.003, U=0, n1=4, n2=6, p=0.009) and between sibs (0.008 versus

304 0.118, U=0, n1=4, n2=6, p=0.009) than found in samples from the Atlantic side.

306 **Heterozygosity and deviation from the genotypic proportions expected under Hardy-**

307 **Weinberg hypotheses**

308 We found no difference in observed heterozygosity between cohorts on the Pacific and Atlantic

309 sides (0.241 versus 0.181, U=15, n1=4, n2=6, p=0.61), though variance of observed

310 heterozygosity between loci was nearly two times higher in cohorts on the Atlantic side (0.389

311 versus 0.701, U=1, n1=4, n2=6, p=0.019). Mean Fis over all loci were significantly lower in

Newfoundland, the Barents and the White Seas than in Kamchatka and Kurils (0.406 versus -

0.066, U=24, n1=4, n2=6, p=0.009) and even negative in the juvenile cohort of Barents Sea,

and in adults and juveniles of White Sea. Variances of Fis values between loci were significantly

higher, three to ten times, in cohorts on the Atlantic side (1.08 versus 6.41, U=24, n1=4, n2=6,

p=0.009), than found in populations on the Pacific side. Though linkage disequilibrium seemed

higher in cohorts on the Atlantic side, they were not significantly different from values

measured in cohorts on the Pacific side (0.051versus 0.356, U=5, n1=4, n2=6, p=0.171).

Estimated rates of clonality using temporal genotype frequencies

Credible intervals including 99% of posterior probabilities of rates of clonality estimated from

temporal samples were quite sharp, spanning around their peaks on a range of 0.06 to maximum

0.25 (Figure 1). In Kuril Islands, rate of clonality was estimated between 0.47 and 0.65 with a

peak at 0.56. In Kamchatka, it ranged between 0.37 to 0.53 with a peak at 0.46; In Barents Sea,

between 0.75 and 1.0 with a peak at 1; In White Sea, between 0.9 and 0.96 with a peak at 0.94;

In Newfoundland, between 0.74 and 0.89 with a peak at 0.82.

Genetic structure between sites using mitochondrial gene sequences

Using three gene sequences (12S, 16S, COIII), we found that all the 82 adults sampled on the

Atlantic side (Newfoundland, the White and the Barents Seas) shared the same mitochondrial

haplotype, Pacific-3 (classification according to Bocharova, 2015; GenBank Accession

Numbers: 12S: KT310190, 16S: KT310200, COIII: KT310212). On the Pacific side, all the 19

individuals sampled in the Kuril islands belonged to another mitochondrial haplotype, Pacific-

1 (GenBank Accession Numbers: 12S: KT310188, 16S: KT310198, COIII: KT310210) while

Kamchatka's samples distributed among five mitochondrial haplotypes: eight individuals

matching Pacific-1 haplotype; two individuals matching Pacific-2 (GenBank Accession

Numbers: 12S: KT310189, 16S: KT310199, COIII: KT310211); one individual matching

Pacific-3; two individuals matching Pacific-4 (GenBank Accession Numbers: 12S: KT310191,

16S: KT310201, COIII: KT310213) and one individual matching Pacific-5 (GenBank

Accession Numbers: 12S: KT310192, 16S: KT310202, COIII: KT310214). Pacific samples

distributed among all the haplotypes known in this species, with Pacific-1 being the most

common haplotype. In contrast, all samples on the Atlantic side matched Pacific-3 haplotype

that was also found to be the rarest in Kamchatka (Figure 2).

Genetic structure between sites and cohorts using nuclear microsatellites

Among adults, genetic differentiation between sites ranged from 0.155 to 0.612 while among

juveniles, genetic differentiation between sites ranged from 0.105 to 0.530 (Table S2). Within

each sampled site, ρST between adults and juveniles ranged from 0.009 to 0.088, therefore

between six to eleven times lower than between sites. Between sites, adults were more

genetically structured than juveniles (0.366 versus 0.245, W=43, p=0.065; Figures 3, S2).

Considering the adults and juveniles together by site, genetic differentiation between sites

ranged from 0.096 to 0.521 (Table S2a). Globally, in juveniles, ρST between sites showed very

close values and followed ranks of geographical and along-the-coast distances (Table S2b,

Figure S2). In contrast, adults of White Sea appeared highly genetically isolated from the other

sites, excepted from juveniles in Newfoundland. ρST genetic differentiation hierarchically

structured the samples into one Pacific side group and one Atlantic side group then subdivided

by sites and then by age cohorts (Table S2b, Figure 3).

This hierarchical genetic structure was congruently supported by the DAPC results (Figure 4).

Interestingly, the Bayesian information criteria of the DAPC decreased continuously with the

increasing number of clusters, as expected in a spatial stepping-stone model (Figure S3). The

minimal number of clusters maximizing the distance between groups was 4. Whatever the

number of clusters, DAPC grouped together individuals of Barents Sea and Newfoundland into one common and compact group, while the other sites each stood for their own genetic groups with couples to handful individuals assigned to other groups. Looking at the genetic structure between cohorts and sites, juveniles from the Kuril Islands shared more genetic diversity with adults from Kamchatka than the inverse (Figure 2). Adults of Barents Sea shared more similar genetic diversity with juveniles of Newfoundland and of White Sea than the inverse. DAPC proportion of each individual assigned to one DAPC cluster also supported this dynamic tendency (Figure 4): one descendant sampled in Kuril was assigned to Kamchatka genetic cluster; Three adults sampled in Kamchatka were assigned to Barents-Newfoundland genetic cluster and one adult sampled in Barents Sea was assigned to Kamchatka genetic cluster; Four adults sampled in Barents Sea were assigned to white sea genetic cluster, and one adult and ten descendants sampled in White Sea were assigned to Barents-Newfoundland genetic cluster. Similarly, the minimum spanning tree of pairwise genetic distances between individuals showed two main groups, one including Kurils and Kamchatka (Pacific side group), and one including Newfoundland, Barents and White Seas (Atlantic side group, Figure 5). Atlantic side group showed more compact genetic distances between individuals than found in the Pacific side group. Atlantic side group also showed more rosary and chaplet subgraphs, commonly observed in clonal lineages (Stoeckel et al., 2024). In accordance with ρST, juveniles were more centrally connected in this network and less genetically distant to the other populations than the adults within each site.

Discussion

We report here the analyses of genetic diversity, genetic differentiation and reproductive mode of five populations of the autotetraploid sea anemone, *Aulactinia stella*, living on North Pacific, Arctic and North Atlantic coasts, at three mitochondrial sequences and ten nuclear

microsatellites. *A. stella* is commonly distributed in the north hemisphere and play a significant

role in shallow-water communities as a sessile predator for smaller animals while feeding larger

predators such as nudibranchs, fish and crabs (Ottaway, 1977). We sampled, genotyped and

analyzed two successive age cohorts within the five sampled sites to assess reproductive modes

in these sites, and to analyze their genetic diversity and its geographical structure across the

Arctic Ocean. By comparing changes in genetic differentiation between adults and juveniles,

we also assessed the potential routes of ongoing gene flow among these sites. The ten nuclear

microsatellites we used provided sufficient statistical power to disentangle between individuals,

enabling differentiating clones of a same genet from inbred offspring.

397   *Aulactinia stella* spreading across the Arctic Ocean from north Pacific?

A. stella presented a genetic pattern typical of one main source population expanding into new

areas (He et al., 2024). Allelic diversity was nearly twice as high in North Pacific populations

compared to those on the Atlantic side of the Arctic Ocean. Pairwise genetic distances revealed

more recent and compact coancestry among individuals from Newfoundland, the Barents Sea,

and the White Sea, in contrast to the more diverse and divergent genetic profiles observed in

populations from the Kuril Islands and Kamchatka. Consistently, mitochondrial sequences

showed the highest haplotype diversity in Kamchatka. Notably, all individuals from Atlantic

populations belonged to the Pacific-3 haplotype, one of the least common haplotypes in Pacific

populations. In the Kuril Islands, only one haplotype, Pacific-1, was identified, which was also

the most prevalent among Kamchatka individuals. This pattern of allelic and haplotype

diversity suggests that *A. stella* is likely native to the North Pacific, particularly the Kamchatka

region, or that the Kamchatka coast served as one of its last glacial refugia. In contrast,

populations on the Atlantic side of the Arctic Ocean may have been established more recently

or may have originated from smaller, more localized glacial refugia.

Genetic differentiation in *A. stella* revealed a hierarchical structure: first separating into a Pacific and an Atlantic group, each of which was further subdivided by sampling site, and then by age cohort. This type of structure closely resembles patterns seen in the intertidal macroalga *Fucus distichus* and in other Arctic and peri-Arctic sessile coastal marine species (Coyer et al., 2011; Bringloe et al., 2020). Environmental factors, rather than species-specific biological or behavioral traits, may likely be the primary drivers of genetic differentiation in this region.

When comparing genetic diversity between adults and juveniles, juveniles were less genetically structured, indicating ongoing gene flow across the Arctic Ocean following a stepping-stone model of dispersal. Pairwise ρST between sites and age cohorts, and DAPC indicated that genetic diversity from adult populations in Kamchatka seemed to flow into juvenile populations in the Kuril Islands and in the Barents Sea. Then, genetic diversity found in Barents Sea adults seemed to extend into juvenile populations in Newfoundland and the White Sea. These dynamics may point the expansion route of *A. stella* over recent centuries from North Pacific towards Atlantic coasts. Interestingly, these dispersal routes align with i). the coastal oceanic currents, i.e., the Anadyr current going from the North Pacific into the Arctic Ocean, then the Beaufort gyre and the transpolar drift, which split into the Baffin current to Newfoundland and into the Persey current to the Barents Sea relayed by the Murman coastal current to the White Sea (Rudels & Carmack, 2022; Timmermans & Marshall, 2020), ii). routes of lower ice concentration along a year (Constantine & Johnson, 2023) and iii). direct and along-coasts geographic distances.

Additionally, we may conjecture that *A. stella* may benefit from increasing maritime traffic across the Arctic. This could occur either through active transport in ballast water from cargo ships, fishing vessels, and tankers using the Northern Sea Route and the Northeast Passage on the Asian side, or the Northwest Passage on the American side (PAME, 2024; Melia et al., 2016; Boylan, 2021); or through passive dispersal, with ship wake currents enhancing the dispersal

of gametes and juveniles along these routes. Our results about *A. stella* foresee on the potential role of increasing Arctic maritime traffic in accelerating the spread of marine organisms between the North Pacific and North Atlantic, with important implications for Arctic biodiversity and ecosystem dynamics.

Reproductive modes in *Aulactinia stella* populations

We found strong genetic evidence that *A. stella* reproduced clonally in natural populations. These results were consistent with the absence of males in some populations despite the presence of juveniles and with previous developmental observations showing that some embryos can develop from unfertilized eggs within the mesenteries of parental sea anemones and be released at the mid-gastrula stage (Bocharova, 2012; 2015; Kaliszewicz et al., 2012). This genetic evidence, together with the absence of males in some populations despite the observation of juveniles argued that clonality may commonly occur in this species, probably via ameiotic parthenogenesis. We therefore assessed the quantitative significance of clonality within populations.

In the North Pacific populations, we identified only one trio of individuals with an identical MLG in Kamchatka and none in Kuril Islands. Yet, genetic indices, including interlocus F_{IS} variances measured around 1 and low levels of linkage disequilibrium, indicated that these populations reproduce with intermediate rates of clonality (Navascues et al., 2010; Stoeckel et al., 2021a). Using changes in genotype frequencies between adults and juveniles, maximum posterior probabilities and their associated 99% credible intervals estimated rates of clonality between 0.37 and 0.65 in these North Pacific populations. The absence of repeated genotypes, despite intermediate clonality, is common in samples from large populations or when clonal offspring are spatially dispersed before settling as adults. This results in high clonal evenness

and reduces the likelihood of sampling identical genotypes (Arnaud-Haond et al., 2007; Stoeckel et al., 2021a,b; 2024).

In contrast, populations from Newfoundland, the Barents Sea, and the White Sea, which are likely to be more recently established due to their lower allelic diversity, showed strong signals of clonal reproduction. All individuals sampled shared the same mitochondrial haplotype (Pacific-3) and many shared identical nuclear MLGs, despite the low probability of such identity by sexual reproduction, even between siblings, given the resolution of our microsatellite markers. Mean negative F_{IS}, large variance of F_{IS} between loci and significant linkage disequilibrium also indicated high rates of clonality (Halkett et al., 2005; De Meeûs et al., 2006; Navascues et al., 2010; Stoeckel et al., 2021a). Estimated clonality rates in these populations ranged from 0.74 to 1.0, nearly double the rates found in North Pacific populations.

The advantages of clonal reproduction at the limits of species' ranges

High levels of uniparental reproduction, including clonality, are commonly observed in species expanding into new or marginal environments (Barrett, 2015; Pannel et al., 2015; Burke & Bonduriansky, 2019, Pereyra et al., 2023), particularly when enduring harsh conditions (Klimeš 2008). The post-glacial Baltic sea provides several examples, including the emblematic case of *Fucus vesiculosus* (Johannesson & André 2006; Pereyra et al., 2009; 2023). In plant species, the persistence of connection between ramets (mainly by roots) offers ecological advantages that may alone explain the prevalence of clonality at range margins, including regeneration after injury, distribution of physiological functions among ramets, translocation of resources between interconnected ramets (clonal integration) and space monopolization using different strategies (including phalanx versus guerrilla) to address intra- and inter-specific competitions depending on the environmental context (Cadotte et al., 2006; Gonzalvez et al., 2008; Roiloa, 2019; Franklin et al., 2021).

In contrast, dispersed clones in marine species such as *A. stella* do not benefit from such clonal

integration. In this context, clonality should only favor kin cooperation (e.g., through brooding

or cooperative behavior Bocharova, 2012; Rudin & Briffa, 2012) but more importantly it should

favor the persistence of local populations under harsh conditions by avoiding the energetic and

biological costs of sexual reproduction (Klimeš, 2008; Pierre et al., 2022).

Baker (1955) postulated that uniparental reproduction (including clonality) provides a critical

advantage for colonization: individuals arriving in new areas can reproduce clonally to establish

a new area. First, clonality allows a population to maintain itself locally until compatible mates

become available (Baker & Stebbins, 1965, Daly et al., 2023). This hypothesis is particularly

relevant for gonochoristic (or dioecious) sessile organisms that require close proximity for

successful sexual reproduction (Barrett, 2015), such as in sea anemones. Second, in new harsh

environments, the more clonal individuals that manage to survive minimize the risks of genetic

reassortment associated with sexual reproduction for their offspring (Peniston et al., 2021;

Pierre et al., 2022). Third, clonality maintains higher effective population sizes than in identical

more sexual populations (Reichel et al. 2016; Stoeckel et al., 2021b). Clonality limits the

negative effects of genetic drift during founder events, including the fixation of deleterious

alleles in small populations (Navascues et al., 2010; He et al., 2024).

Our results in *A. stella*, which showed a twofold increase in clonality at the species' range edge,

highlight the key role of clonal reproduction in maintaining populations and higher effective

population size under unfavorable conditions while expanding into new areas (Baker's

conjecture, Pannel et al., 2015) without invoking for the benefits of clonal integration (as

proposed in plants) or habitat modification (as proposed in ecological engineering species) as

drivers of clonal success.

Restricted dispersal or local adaptation in *Aulactinia stella*?

Interestingly, we found no MLGs shared between sampling sites, even those in close geographic proximity, such as Barents and White Seas sampling sites (~360 kilometers), and the Kuril Islands and Kamchatka coasts (~680 kilometers). The presence of local, site-specific clones and high levels of genetic differentiation between populations appears to be a common feature of several cold-water sea anemone species. These include *Actinia tenebrosa* in New Zealand (Veale & Lavery, 2012) and on the Australia and Tasmania coasts (Sherman et al., 2007; Sherman & Ayre, 2008), *Anthothoe albocincta* on the south-eastern Australia coast (Billingham & Ayre, 1996), *Diadumene lineata* in the North Atlantic (Ryan et al., 2021), and the North Pacific species *Anthopleura elegantissima*, *Epiactis prolifera*, *Epiactis lisbethae* and *Epiactis ritteri* (Edmands & Potts, 1997) *Haliplanella lucia* (Shick & Lamb, 1977). This pattern of high genetic differentiation contrasts with tropical sea anemones, such as *Actinia bermudensis*, *Bunodosoma caissarum*, *Entacmaea quadricolor*, *Heteractis magnifica* and *Stichodactyla mertensii* which typically show much lower levels of genetic structure and often share genets across sites separated by hundreds of kilometers apart (Russo et al., 1994; Emms et al., 2020; but see Li et al., 2024 for *Paracondylactis sinensis*). These observations may support the hypothesis that the Arctic marine environments may have constrained and challenged the dispersal of sea anemones (Chan et al., 2019). However, other passively dispersing organisms in similar cold-water habitats, such as macroalgae, seagrasses, and bivalves, generally show much less population structure and often share genets between sites (Coolen et al., 2020; Pereyra et al., 2023; Krueger-Hadfield et al., 2021). In addition, juvenile sea anemones are capable of active movement in the direction of marine currents, enabling long-distance dispersal (Riemann-Zürneck, 1998). *Urticina eques*, another cold-water sea anemone, shows very low genetic differentiation beween populations distributed over hundreds of kilometers along the British coast (Cava-Solé et al., 1994).

In the case of *A. stella*, we found lower genetic differentiation between sites in juveniles and

pairwise genetic distances were more compact in juveniles than in adults. The DAPC results

supported a stepping-stone genetic structure. These results suggest that the lack of shared MLGs

between sites and the observed genetic structure may be better explained by local genetic

endemism or gregarious reproductive behavior promoting shared ancestry, rather than by

persistent founder effects or a complete lack of gene flow (Lane et al., 2020; Ryan et al., 2021).

We rather observed *A. stella* occupying diverse habitats across sampling sites. For example,

although the Barents and White Seas are separated by only about 100 km, the White Sea is

characterized by significantly lower salinity. In marine environments, factors such as water

temperature, depth, tidal exposure, and substrate disturbance are known to influence both

reproductive strategies and levels of genetic differentiation (Eriksson and Johansson 2005;

Krueger-Hadfield et al., 2013; Berković et al., 2014; Dudgeon et al., 2016; McMahon et al.,

2017; Ryan & Miller, 2019; Ryan et al., 2021; Williams et al., 2024). Similar to *A. stella*, *Actinia*

tenebrosa, a partially clonal brooding sea anemone, localized clonal distributions and endemic

genetic lineages have been linked to specific habitat characteristics, such as atmospheric

exposure, temperature, salinity, UV radiation, and local community composition, despite its

potential for wider dispersal (Sherman & Ayre, 2008). *Diadumene lineata* also reproduces with

various different rates of clonality depending on water temperature and shows similar

geographic genetic differentiation, particularly between northern and southern Atlantic

populations (Ryan et al., 2021).

In the case of *A. stella*, Arctic populations may have only narrow temporal windows of suitable

thermal conditions for successful sexual reproduction, particularly in the intertidal and shallow

sublittoral zones. These habitats are also characterized by extreme variability in salinity and

desiccation at low tide. In such habitats, *A. stella* is often found hiding under *Fucus* algae, in

rock crevices, or in tide pools, behaviors that indicate exposure to environmental stress. Our

results, showing approximately twice the level of genetic differentiation among adults

compared to juveniles, highlight the need for further ecological and behavioral studies. *A. stella*

presents a paradoxical tension between its current ability to disperse into new diverse areas and

an interesting tendency towards endemism, commonly observed in several other sea anemone

species. The spread of *A. stella* to the North Atlantic sites may provide an interesting

opportunity to study how endemism can arise in populations colonizing new areas from a

common genetic background. Experiments using common gardens and reciprocal

transplantations between sites could help to assess the role of local adaptation versus migration

in shaping the genetic endemism of *A. stella* and to identify which environmental conditions

are most influential for this species. If local adaptation is confirmed, an important question

remains: how has *A. stella* successfully colonized such a wide and environmentally diverse

range across the Arctic? Experimental studies that expose both clones and sexually-derived

offspring to novel habitats over multiple generations may reveal the pace and mechanisms of

adaptation in this species as it makes its odyssey through the Arctic Ocean.

Data Availability

576 MLG genotypes, sample GPS locations and clone sizes are openly available on Zenodo

577 https://doi.org/10.5281/zenodo.15102245

Supplementary Material

Supplementary material provides two tables and three figures. Table S1 reports distributions of

repeated MLGs within sites and MLGs shared between adults and juveniles; Table S2 reports

pairwise ρST between sites and between age cohorts; Figure S1 reports how allele dosage of

the microsatellites was achieved; Figure S2 reports the minimum spanning trees of genetic

differentiation of adults, juveniles and overall among sites; Figure S3 reports the change of the

Bayesian information criteria of the DAPC with increasing number of genetic clusters arguing
for a stepping-stone structure of demes.

Acknowledgements

The authors wish to thank N.P. Sanamyan, K.E. Sanamyan, Osadchenko B.V and Kosevich I.A.
for providing material and assistance in species identifications, N.S. Mugue, T.V. Neretina, A.E.
Barmintseva and S.D. Grebelnyi for professional consultations, A. Mercier, J.-F. Hamel, E.
Montgomery and the Ocean Sciences Field Services of Memorial University for their assistance
with the animal collections. We thank Patricia Nadan and Pascale Le Névé (INRAE IGEPP) for
their administrative support along this project.

Funding

This work was funded by the Russian Foundation for Basic Research (Grant No. 16-04-01685
to E.S. Bocharova), by the French National Research Agency (project Clonix2D ANR-18-
CE32-0001 to S. Stoeckel) and the French Embassy in the Russian Federation (Metchnikov
Stipend 2019 to E.S. Bocharova).

Author contributions

EB and SS laid the foundation of this work while EB visited SS lab in 2018 and 2019, conceived
the study and were responsible for funding applications. EB was responsible for data
acquisition. She supervised and achieved sample collection, determined the species and sexed
the sampled individuals. She also conceived, developed the sequences and microsatellites,
supervised and achieved sequencing and genotyping of all the samples. From sequencing
machines, she genotyped the samples. AV contributed to methodology, supervision and

administration of data acquisition. EB performed data analyses and produced figure on

mitochondrial sequences. SS performed data analyses and produced tables and figures on

microsatellites. SS and EB tracked and managed the bibliography, interpreted results, wrote the

manuscript and managed its edition. All authors read and approved this manuscript.

Conflict of interest disclosure

615 The authors declare that they comply with the PCI rule of having no financial conflicts of

616 interest in relation to the content of the article.

References

Agapow P-M, Burt A (2001) Indices of multilocus linkage disequilibrium. Molecular Ecology Notes,
1, 101–102. https://doi.org/10.1046/j.1471-8278.2000.00014.x
Arnaud-Haond S, Duarte CM, Alberto F, Serrão EA (2007) Standardizing methods to address
clonality in population studies. Molecular Ecology, 16, 5115–5139.
https://doi.org/10.1111/j.1365-294X.2007.03535.x
Arnaud-Haond S, Stoeckel S, Bailleul D (2020) New insights into the population genetics of partially
clonal organisms: When seagrass data meet theoretical expectations. Molecular Ecology, 29,
3248–3260. https://doi.org/10.1111/mec.15532
Avise JC (2015) Evolutionary perspectives on clonal reproduction in vertebrate animals.
Proceedings of the National Academy of Sciences, 112, 8867–8873.
https://doi.org/10.1073/pnas.1501820112
Bailleul D, Stoeckel S, Arnaud-Haond S (2016) RClone: a package to identify MultiLocus Clonal
Lineages and handle clonal data sets in R. Methods in Ecology and Evolution, 7, 966–970.
https://doi.org/10.1111/2041-210X.12550
Baker HG (1955) Self-Compatibility and Establishment After '"Long-Distance" Dispersal. Evolution,
9, 347–349. https://doi.org/10.1111/j.1558-5646.1955.tb01544.x
Baker HG, Stebbins GL (1965) The genetics of colonizing species. New York.
Barrett SCH (2015) Foundations of invasion genetics: the Baker and Stebbins legacy. Molecular
Ecology, 24, 1927–1941. https://doi.org/10.1111/mec.13014
Becheler R, Masson J-P, Arnaud-Haond S, Halkett F, Mariette S, Guillemin M-L, Valero M,
Destombe C, Stoeckel S (2017) ClonEstiMate, a Bayesian method for quantifying rates of
clonality of populations genotyped at two-time steps. Molecular Ecology Resources, 17, e251–
e267. https://doi.org/10.1111/1755-0998.12698
Berković B, Cabaço S, Barrio JM, Santos R, Serrão EA, Alberto F (2014) Extending the life history
of a clonal aquatic plant: Dispersal potential of sexual and asexual propagules of Zostera noltii.
Aquatic Botany, 113, 123–129. https://doi.org/10.1016/j.aquabot.2013.10.007
Billingham M, Ayre DJ (1996) Genetic subdivision in the subtidal, clonal sea anemone Anthothoe
albocincta. Marine Biology, 125, 153–163. https://doi.org/10.1007/BF00350769
Bocharova ES (2012) Osobennosti razmnozheniya Aulactinia stella (Verrill, 1864) (Hexacorallia,
Actiniidae) v Belom i Barentsevom moryakh (Reproduction characters of Au- lactinia stella

(Verrill, 1864) (Hexacorallia, Actiniidae) in the White and the Barents Seas). Moscow State University, Moscow.

Bocharova ES (2015) Reproductive biology and genetic diversity of the Sea Anemone Aulactinia stella (Verrill, 1864). Hydrobiologia, 759, 27–38. https://doi.org/10.1007/s10750-015-2250-0

Bocharova E (2016) Reproduction of Sea Anemones and Other Hexacorals. In: The Cnidaria, Past, Present and Future (eds Goffredo S, Dubinsky Z), pp. 239–248. Springer International Publishing, Cham. https://doi.org/10.1007/978-3-319-31305-4_15

Bocharova ES, Kozevich IA (2011) Modes of reproduction in sea anemones (Cnidaria, Anthozoa). Biology Bulletin, 38, 849–860. https://doi.org/10.1134/S1062359011090020

Bocharova ES, Mugue NS (2012) Sea anemones Aulactinia stella (Verrill, 1864) (Hexacorallia, Actiniidae) can brood offspring from other individuals of the same species. Doklady Biological Sciences, 444, 173–175. https://doi.org/10.1134/S0012496612030040

Bocharova ES, Sergeev AA, Volkov AA (2018) Identification of microsatellite loci in sea anemones Aulactinia stella and Cribrinopsis albopunctata (family Actiniidae). F1000Research, 7, 232. https://doi.org/10.12688/f1000research.13724.1

Boylan BM (2021) Increased maritime traffic in the Arctic: Implications for governance of Arctic sea routes. Marine Policy, 131, 104566. https://doi.org/10.1016/j.marpol.2021.104566

Bridle J, Hoffmann A (2022) Understanding the biology of species' ranges: when and how does evolution change the rules of ecological engagement? Philosophical Transactions of the Royal Society B: Biological Sciences, 377, 20210027. https://doi.org/10.1098/rstb.2021.0027

Bringloe TT, Verbruggen H, Saunders GW (2020) Unique biodiversity in Arctic marine forests is shaped by diverse recolonization pathways and far northern glacial refugia. Proceedings of the National Academy of Sciences, 117, 22590–22596. https://doi.org/10.1073/pnas.2002753117

Burke NW, Bonduriansky R (2018) The geography of sex: sexual conflict, environmental gradients and local loss of sex in facultatively parthenogenetic animals. Philosophical Transactions of the Royal Society B: Biological Sciences, 373, 20170422. https://doi.org/10.1098/rstb.2017.0422

Cadotte MW, Murray BR, Lovett-Doust J (2006) Ecological Patterns and Biological Invasions: Using Regional Species Inventories in Macroecology. Biological Invasions, 8, 809–821. https://doi.org/10.1007/s10530-005-3839-4

Cava-Solé AM, Thorpe JP, Todd CD (1994) High genetic similarity between geographically distant populations in a sea anemone with low dispersal capabilities. Journal of the Marine Biological Association of the United Kingdom, 74, 895–902. https://doi.org/10.1017/S0025315400090123

Chan FT, Stanislawczyk K, Sneekes AC, Dvoretsky A, Gollasch S, Minchin D, David M, Jelmert A, Albretsen J, Bailey SA (2019) Climate change opens new frontiers for marine species in the Arctic: Current trends and future invasion risks. Global Change Biology, 25, 25–38. https://doi.org/10.1111/gcb.14469

Constantin A, Johnson RS (2023) On the dynamics of the near-surface currents in the Arctic Ocean. Nonlinear Analysis: Real World Applications, 73, 103894. https://doi.org/10.1016/j.nonrwa.2023.103894

Coolen JWP, Boon AR, Crooijmans R, van Pelt H, Kleissen F, Gerla D, Beermann J, Birchenough SNR, Becking LE, Luttikhuizen PC (2020) Marine stepping-stones: Connectivity of Mytilus edulis populations between offshore energy installations. Molecular Ecology, 29, 686–703. https://doi.org/10.1111/mec.15364

Coyer JA, Hoarau G, Van Schaik J, Luijckx P, Olsen JL (2011) Trans-Pacific and trans-Arctic pathways of the intertidal macroalga Fucus distichus L. reveal multiple glacial refugia and colonizations from the North Pacific to the North Atlantic. Journal of Biogeography, 38, 756–771. https://doi.org/10.1111/j.1365-2699.2010.02437.x

Daly EZ, Chabrerie O, Massol F, Facon B, Hess MCM, Tasiemski A, Grandjean F, Chauvat M, Viard F, Forey E, Folcher L, Buisson E, Boivin T, Baltora-Rosset S, Ulmer R, Gibert P, Thiébaut G, Pantel JH, Heger T, Richardson DM, Renault D (2023) A synthesis of biological invasion hypotheses associated with the introduction–naturalisation–invasion continuum. Oikos, 2023, e09645. https://doi.org/10.1111/oik.09645

De Meeûs T, Lehmann L, Balloux F (2006) Molecular epidemiology of clonal diploids: A quick
overview and a short DIY (do it yourself) notice. Infection, Genetics and Evolution, 6, 163–170.
https://doi.org/10.1016/j.meegid.2005.02.004
De Meeûs T, Prugnolle F, Agnew P (2007) Asexual reproduction: Genetics and evolutionary
aspects. Cellular and Molecular Life Sciences, 64, 1355–1372. https://doi.org/10.1007/s00018-
007-6515-2
Dorken ME, Eckert CG (2001) Severely reduced sexual reproduction in northern populations of a
clonal plant, Decodonverticillatus (Lythraceae). Journal of Ecology, 89, 339–350.
https://doi.org/10.1046/j.1365-2745.2001.00558.x
Drauch Schreier A, Gille D, Mahardja B, May B (2011) Neutral markers confirm the octoploid origin
and reveal spontaneous autopolyploidy in white sturgeon, Acipenser transmontanus: Neutral
markers and ploidy in white sturgeon. Journal of Applied Ichthyology, 27, 24–33.
https://doi.org/10.1111/j.1439-0426.2011.01873.x
Dudgeon S, Kübler JE, West JA, Kamiya M, Krueger-Hadfield SA (2017) Asexuality and the cryptic
species problem. Perspectives in Phycology, 47–59. https://doi.org/10.1127/pip/2017/0070
Duminil J, Fineschi S, Hampe A, Jordano P, Salvini D, Vendramin GG, Petit RJ (2007) Can
Population Genetic Structure Be Predicted from Life-History Traits? The American Naturalist,
169, 662–672. https://doi.org/10.1086/513490
Edmands S, Potts DC (1997) Population genetic structure in brooding sea anemones (Epiactis
spp.) with contrasting reproductive modes. Marine Biology, 127, 485–498.
https://doi.org/10.1007/s002270050037
Ellegren H, Galtier N (2016) Determinants of genetic diversity. Nature Reviews Genetics, 17, 422–
433. https://doi.org/10.1038/nrg.2016.58
Emms MA, Saenz-Agudelo P, Giles EC, Gatins R, Nanninga GB, Scott A, Hobbs J-PA, Frisch AJ,
Mills SC, Beldade R, Berumen ML (2020) Comparative phylogeography of three host sea
anemones in the Indo-Pacific. Journal of Biogeography, 47, 487–500.
https://doi.org/10.1111/jbi.13775
Eriksson BK, Johansson G (2005) Effects of sedimentation on macroalgae: species-specific
responses are related to reproductive traits. Oecologia, 143, 438–448.
https://doi.org/10.1007/s00442-004-1810-1
Fautin Dunn DF, Devaney DM, Roth B (1980) Stylobates: A shell-forming sea anemone
(Coelenterata, Anthozoa, Actiniidae). Pacific Science, 34, 379–388.
Felsenstein J (2004) Inferring phylogenies. Sinauer Associates, Sunderland, Mass.
Fouqueau L, Roze D (2021) The evolution of sex along an environmental gradient. Evolution, 75,
1334–1347. https://doi.org/10.1111/evo.14237
Franklin S, Alpert P, Salguero-Gómez R, Janovský Z, Herben T, Klimešová J, Douhovnikoff V
(2021) Next-gen plant clonal ecology. Perspectives in Plant Ecology, Evolution and
Systematics, 49, 125601. https://doi.org/10.1016/j.ppees.2021.125601
Gonzalez A, Lambert A, Ricciardi A (2008) When does ecosystem engineering cause invasion and
species replacement? Oikos, 117, 1247–1257. https://doi.org/10.1111/j.0030-
1299.2008.16419.x
Grebelnyi SD Distribution of the sea anemones in the Arctic. In: The theoretical and practical
significance of Coelenterata , pp. 20–33. Publishing House of the Zoological Institute of the
USSR Academy of Sciences, Leningrad.
Halkett F, Simon J-C, Balloux F (2005) Tackling the population genetics of clonal and partially
clonal organisms. Trends in Ecology & Evolution, 20, 194–201.
https://doi.org/10.1016/j.tree.2005.01.001
He Z-Z, Stotz GC, Liu X, Liu J-J, Wang Y-G, Yang J, Li L-F, Zhang W-J, Nan P, Song Z-P (2024)
A global synthesis of the patterns of genetic diversity in endangered and invasive plants.
Biological Conservation, 291, 110473. https://doi.org/10.1016/j.biocon.2024.110473
Jacquard A (1968) Panmixie et consanguinite. Quelques precisions de langage. Population
(French Edition), 23, 1065. https://doi.org/10.2307/1528645
Jacquard A (1970) Structure génétiques des populations. Masson & cie, Paris.

Johannesson K, André C (2006) INVITED REVIEW: Life on the margin: genetic isolation and diversity loss in a peripheral marine ecosystem, the Baltic Sea. Molecular Ecology, 15, 2013–2029. https://doi.org/10.1111/j.1365-294X.2006.02919.x

Jombart T, Devillard S, Balloux F (2010) Discriminant analysis of principal components: a new method for the analysis of genetically structured populations. BMC Genetics, 11, 94. https://doi.org/10.1186/1471-2156-11-94

Kaliszewicz A, Panteleeva N, Olejniczak I, Boniecki P, Sawicki M (2012) Internal brooding affects the spatial structure of intertidal sea anemones in the Arctic-boreal region. Polar Biology, 35, 1911–1919. https://doi.org/10.1007/s00300-012-1232-1

Klimeš L (2008) Clonal splitters and integrators in harsh environments of the Trans-Himalaya. Evolutionary Ecology, 22, 351–367. https://doi.org/10.1007/s10682-007-9195-3

Krueger-Hadfield SA, Guillemin M-L, Destombe C, Valero M, Stoeckel S (2021) Exploring the Genetic Consequences of Clonality in Haplodiplontic Taxa. Journal of Heredity, 112, 92–107. https://doi.org/10.1093/jhered/esaa063

Krueger-Hadfield SA, Kübler JE, Dudgeon SR (2013) Reproductive effort of Mastocarpus papillatus (Rhodophyta) along the California coast1. Journal of Phycology, 49, 271–281. https://doi.org/10.1111/jpy.12034

Lane SM, Wilson AJ, Briffa M (2020) Analysis of direct and indirect genetic effects in fighting sea anemones. Behavioral Ecology, 31, 540–547. https://doi.org/10.1093/beheco/arz217

Li J, Yang X, Zhan Z, Feng J, Xie T, Li Y (2024) Development of microsatellite markers and evaluation of the genetic diversity of the edible sea anemone Paracondylactis sinensis (Cnidaria, Anthozoa) in China. Biodiversity Data Journal, 12, e134363. https://doi.org/10.3897/BDJ.12.e134363

Liu J, Dong M, Miao SL, Li ZY, Song MH, Wang RQ (2006) Invasive alien plants in China: role of clonality and geographical origin. Biological Invasions, 8, 1461–1470. https://doi.org/10.1007/s10530-005-5838-x

Loseva LM (1974) Polovoye sostoyanie Bunodactis stella (Sex composition of Bunodactis stella). In: Biologiya razvi- tiya morskikh organizmov (Developmental Biology of Marine Organisms) , pp. 133–142. Akademiya Nauk SSSR, Apatity.

Manuel RL (1981) British Anthozoa: Keys and notes for the identification of the species. Acad. Pr, London u.a.

McMahon KM, Evans RD, van Dijk K, Hernawan U, Kendrick GA, Lavery PS, Lowe R, Puotinen M, Waycott M (2017) Disturbance Is an Important Driver of Clonal Richness in Tropical Seagrasses. Frontiers in Plant Science, 8. https://doi.org/10.3389/fpls.2017.02026

Melia N, Haines K, Hawkins E (2016) Sea ice decline and 21st century trans-Arctic shipping routes. Geophysical Research Letters, 43, 9720–9728. https://doi.org/10.1002/2016GL069315

Moran EV, Alexander JM (2014) Evolutionary responses to global change: lessons from invasive species. Ecology Letters, 17, 637–649. https://doi.org/10.1111/ele.12262

Navascués M, Stoeckel S, Mariette S (2010) Genetic diversity and fitness in small populations of partially asexual, self-incompatible plants. Heredity, 104, 482–492. https://doi.org/10.1038/hdy.2009.159

Orive ME, Krueger-Hadfield SA (2021) Sex and Asex: A Clonal Lexicon. Journal of Heredity, 112, 1–8. https://doi.org/10.1093/jhered/esaa058

Ottaway JR (1977) Predators of sea anemones. Tuatara, 22, 213–220.

Pannell JR, Auld JR, Brandvain Y, Burd M, Busch JW, Cheptou P-O, Conner JK, Goldberg EE, Grant A-G, Grossenbacher DL, Hovick SM, Igic B, Kalisz S, Petanidou T, Randle AM, de Casas RR, Pauw A, Vamosi JC, Winn AA (2015) The scope of Baker's law. New Phytologist, 208, 656–667. https://doi.org/10.1111/nph.13539

Pereyra RT, Bergström L, Kautsky L, Johannesson K (2009) Rapid speciation in a newly opened postglacial marine environment, the Baltic Sea. BMC Evolutionary Biology, 9, 70. https://doi.org/10.1186/1471-2148-9-70

Pereyra RT, Rafajlović M, De Wit P, Pinder M, Kinnby A, Töpel M, Johannesson K (2023) Clones
on the run: The genomics of a recently expanded partially clonal species. Molecular Ecology,
32, 4209–4223. https://doi.org/10.1111/mec.16996

Pierre J-S, Stoeckel S, Wajnberg E (2022) The advantage of sex: Reinserting fluctuating selection
in the pluralist approach (T Friedlander, Ed,). PLOS ONE, 17, e0272134.
https://doi.org/10.1371/journal.pone.0272134

Protection of the Arctic Marine Environment (PAME) - Working Group of the Arctic Council (2024)
Arctic Ship Traffic Data (ASTD).

Reichel K, Masson J-P, Malrieu F, Arnaud-Haond S, Stoeckel S (2016) Rare sex or out of reach
equilibrium? The dynamics of F IS in partially clonal organisms. BMC Genetics, 17, 76.
https://doi.org/10.1186/s12863-016-0388-z

Riemann-Zürneck K (1998) How Sessile are Sea Anemones? A Review of Free-living Forms in the
Actiniaria Cnidaria: Anthozoa. Marine Ecology, 19, 247–261. https://doi.org/10.1111/j.1439-
0485.1998.tb00466.x

Rodríguez E, Fautin D, Daly M (2025) World List of Actiniaria. Accessed at
https://www.marinespecies.org/actiniaria on yyyy-mm-dd. https://doi.org/10.14284/568

Rodzen JA, May B (2002) Inheritance of microsatellite loci in the white sturgeon (Acipenser
transmontanus). Genome, 45, 1064–1076. https://doi.org/10.1139/g02-083

Roiloa SR (2019) Clonal traits and plant invasiveness: The case of Carpobrotus N.E.Br.
(Aizoaceae). Perspectives in Plant Ecology, Evolution and Systematics, 40, 125479.
https://doi.org/10.1016/j.ppees.2019.125479

Ronfort J, Jenczewski E, Bataillon T, Rousset F (1998) Analysis of Population Structure in
Autotetraploid Species. Genetics, 150, 921–930. https://doi.org/10.1093/genetics/150.2.921

Rudels B, Carmack E (2022) Arctic Ocean Water Mass Structure and Circulation. Oceanography,
35, 52–65. https://doi.org/10.5670/oceanog.2022.116

Rudin FS, Briffa M (2012) Is boldness a resource-holding potential trait? Fighting prowess and
changes in startle response in the sea anemone, Actinia equina. Proceedings of the Royal
Society B: Biological Sciences, 279, 1904–1910. https://doi.org/10.1098/rspb.2011.2418

Russo CAM, Solé-Cava AM, Thorpe JP (1994) Population structure and genetic variation in two
tropical sea anemones (Cnidaria, Actinidae) with different reproductive strategies. Marine
Biology, 119, 267–276. https://doi.org/10.1007/BF00349566

Ryan WH, Aida J, Krueger-Hadfield SA (2021) The Contribution of Clonality to Population Genetic
Structure in the Sea Anemone, Diadumene lineata. Journal of Heredity, 112, 122–139.
https://doi.org/10.1093/jhered/esaa050

Ryan WH, Miller TE (2019) Reproductive strategy changes across latitude in a clonal sea anemone.
Marine Ecology Progress Series, 611, 129–141. https://doi.org/10.3354/meps12862

Sanamyan NP, Sanamyan KE (1998) Some Actiniaria from the Commander Islands (Cnidaria:
Anthozoa). Zoosystematica Rossica, 1, 1–8.

Sanamyan NP, Sanamyan KE (2006) The genera Urticina and Cribrinopsis (Anthozoa: Actiniaria)
from the north-western Pacific. Journal of Natural History, 40, 359–393.
https://doi.org/10.1080/00222930600703532

Sanamyan NP, Sanamyan KE Melkovodniye aktinii (Cnidaria: Actiniaria) yugo-vostochnogo
poberezh'ya Kamchatki (Shallow water sea anemones (Cnidaria: Actiniaria) from south-eastern
coast of Kamchatka). Zoologia bespozvonochnikh (Invertebrate Zoology), 155–172.

Sanamyan NP, Sanamyan KE, McDaniel N, Martynov AV, Korshunova TA, Bocharova ES (2019)
A revision of sea anemones of the genus Cribrinopsis Carlgren, 1921(Actiniaria: Actiniidae) from
British Columbia with the description of a new species. Marine Biodiversity, 49, 1951–1969.
https://doi.org/10.1007/s12526-019-00956-w

Sanamyan NP, Sanamyan KE, Mercier A, Hamel J-F, Bocharova ES (2020) Morphological and
molecular assessment of large sea anemones (Actiniaria: Actiniidae) in Newfoundland (eastern
Canada). Polar Biology, 43, 495–509. https://doi.org/10.1007/s00300-020-02652-z

Sax DF, Stachowicz JJ, Brown JH, Bruno JF, Dawson MN, Gaines SD, Grosberg RK, Hastings A,
Holt RD, Mayfield MM, O'Connor MI, Rice WR (2007) Ecological and evolutionary insights from

species invasions. Trends in Ecology & Evolution, 22, 465–471.
https://doi.org/10.1016/j.tree.2007.06.009

Schön I, Martens K, Dijk P (Eds.) (2009) Lost Sex: The Evolutionary Biology of Parthenogenesis.
Springer Netherlands, Dordrecht. https://doi.org/10.1007/978-90-481-2770-2

Sherman CDH, Ayre DJ (2008) Fine-scale adaptation in a clonal sea anemone. Evolution, 62,
1373–1380. https://doi.org/10.1111/j.1558-5646.2008.00375.x

Sherman CDH, Hunt A, Ayre DJ (2008) Is life history a barrier to dispersal? Contrasting patterns
of genetic differentiation along an oceanographically complex coast. Biological Journal of the
Linnean Society, 95, 106–116. https://doi.org/10.1111/j.1095-8312.2008.01044.x

Sherman CDH, Peucker AJ, Ayre DJ (2007) Do reproductive tactics vary with habitat heterogeneity
in the intertidal sea anemone Actinia tenebrosa? Journal of Experimental Marine Biology and
Ecology, 340, 259–267. https://doi.org/10.1016/j.jembe.2006.09.016

Shick JM, Lamb AN (1977) Asexual reproduction and genetic population structure in the colonizing
sea anemone haliplanella luciae. The Biological Bulletin, 153, 604–617.
https://doi.org/10.2307/1540609

Solé-Cava AM, Thorpe JP, Kaye JG (1985) Reproductive isolation with little genetic divergence
between Urticina (=Tealia) felina and U. eques (Anthozoa: Actiniaria). Marine Biology, 85, 279–
284. https://doi.org/10.1007/BF00393248

Stoeckel S, Arnaud-Haond S, Krueger-Hadfield SA (2021) The Combined Effect of Haplodiplonty
and Partial Clonality on Genotypic and Genetic Diversity in a Finite Mutating Population. Journal
of Heredity, 112, 78–91. https://doi.org/10.1093/jhered/esaa062

Stoeckel S, Becheler R, Bocharova E, Barloy D (2024) GenAPoPop 1.0: A user-friendly software
to analyse genetic diversity and structure from partially clonal and selfed autopolyploid
organisms. Molecular Ecology Resources, 24, e13886. https://doi.org/10.1111/1755-
0998.13886

Stoeckel S, Masson J-P (2014) The Exact Distributions of FIS under Partial Asexuality in Small
Finite Populations with Mutation. PLOS ONE, 9, e85228.
https://doi.org/10.1371/journal.pone.0085228

Stoeckel S, Porro B, Arnaud-Haond S (2021) The discernible and hidden effects of clonality on the
genotypic and genetic states of populations: Improving our estimation of clonal rates. Molecular
Ecology Resources, 21, 1068–1084. https://doi.org/10.1111/1755-0998.13316

Tibayrenc M, Avise JC, Ayala FJ (2015) In the light of evolution IX: Clonal reproduction: Alternatives
to sex. Proceedings of the National Academy of Sciences, 112, 8824–8826.
https://doi.org/10.1073/pnas.1508087112

Timmermans M-L, Marshall J (2020) Understanding Arctic Ocean Circulation: A Review of Ocean
Dynamics in a Changing Climate. Journal of Geophysical Research: Oceans, 125,
e2018JC014378. https://doi.org/10.1029/2018JC014378

Turner V, Lynch S, Paterson L, León-Cortés J, Thorpe J (2003) Aggression as a function of genetic
relatedness in the sea anemone Actinia equina (Anthozoa: Actiniaria). Marine Ecology Progress
Series, 247, 85–92. https://doi.org/10.3354/meps247085

Veale A, Lavery S (2012) The population genetic structure of the waratah anemone (Actinia
tenebrosa) around New Zealand. New Zealand Journal of Marine and Freshwater Research,
46, 523–536. https://doi.org/10.1080/00288330.2012.730053

Vermeij GJ, Roopnarine PD (2008) The Coming Arctic Invasion. Science, 321, 780–781.
https://doi.org/10.1126/science.1160852

Verrill AE (1864) List of the polyps and corals sent by the Museum of Comparative Zoology to other
institutions in exchange, with annotations. Bulletin of the Museum of Comparative Zoology, 1,
29–60.

Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D, Burovski E, Peterson
P, Weckesser W, Bright J, Van Der Walt SJ, Brett M, Wilson J, Millman KJ, Mayorov N, Nelson
ARJ, Jones E, Kern R, Larson E, Carey CJ, Polat İ, Feng Y, Moore EW, VanderPlas J, Laxalde
D, Perktold J, Cimrman R, Henriksen I, Quintero EA, Harris CR, Archibald AM, Ribeiro AH,
Pedregosa F, Van Mulbregt P, SciPy 1.0 Contributors, Vijaykumar A, Bardelli AP, Rothberg A,

Hilboll A, Kloeckner A, Scopatz A, Lee A, Rokem A, Woods CN, Fulton C, Masson C, Häggström C, Fitzgerald C, Nicholson DA, Hagen DR, Pasechnik DV, Olivetti E, Martin E, Wieser E, Silva F, Lenders F, Wilhelm F, Young G, Price GA, Ingold G-L, Allen GE, Lee GR, Audren H, Probst I, Dietrich JP, Silterra J, Webber JT, Slavič J, Nothman J, Buchner J, Kulick J, Schönberger JL, De Miranda Cardoso JV, Reimer J, Harrington J, Rodríguez JLC, Nunez-Iglesias J, Kuczynski J, Tritz K, Thoma M, Newville M, Kümmerer M, Bolingbroke M, Tartre M, Pak M, Smith NJ, Nowaczyk N, Shebanov N, Pavlyk O, Brodtkorb PA, Lee P, McGibbon RT, Feldbauer R, Lewis S, Tygier S, Sievert S, Vigna S, Peterson S, More S, Pudlik T, Oshima T, Pingel TJ, Robitaille TP, Spura T, Jones TR, Cera T, Leslie T, Zito T, Krauss T, Upadhyay U, Halchenko YO, Vázquez-Baeza Y (2020) SciPy 1.0: fundamental algorithms for scientific computing in Python. Nature Methods, 17, 261–272. https://doi.org/10.1038/s41592-019-0686-2

Waits LP, Luikart G, Taberlet P (2001) Estimating the probability of identity among genotypes in natural populations: cautions and guidelines. Molecular Ecology, 10, 249–256. https://doi.org/10.1046/j.1365-294X.2001.01185.x

Wang Y-J, Chen D, Yan R, Yu F-H, van Kleunen M (2019) Invasive alien clonal plants are competitively superior over co-occurring native clonal plants. Perspectives in Plant Ecology, Evolution and Systematics, 40, 125484. https://doi.org/10.1016/j.ppees.2019.125484

Williams TM, Krueger-Hadfield SA, Hill-Spanik KM, Kosaki RK, Stoeckel S, Spalding HL (2024) The reproductive system of the cryptogenic alga Chondria tumulosa (Florideophyceae) at Manawai, Papahānaumokuākea Marine National Monument. Phycologia, 63, 36–44. https://doi.org/10.1080/00318884.2023.2284074

Yu F-H, Roiloa SR, Alpert P (2016) Editorial: Global Change, Clonal Growth, and Biological Invasions by Plants. Frontiers in Plant Science, 7. https://doi.org/10.3389/fpls.2016.01467

Zhang R, Jia G, Diao X (2023) geneHapR: an R package for gene haplotypic statistics and visualization. BMC Bioinformatics, 24, 199. https://doi.org/10.1186/s12859-023-05318-9

Tables and Figures

Figure 1: Posterior density probabilities of estimates of rates of clonality using changes of genotype frequencies over one generation for each sampled site using ClonEstiMate method for autopolyploids in GenAPoPop software. Kuril Islands in blue, Kamchatka in green, Newfoundland in red, White Sea in pink and Barents Sea in gold.

Figure 2: Haplotype network using three gene sequences (12S, 16S, COIII). The proportion of each sampled site per haplotype is represented in color. The circle sizes of each haplotype are proportional to their number in our dataset. We used haplotype names initially identified in Bocharova (2015), thus Pacific 1 to 5. We found no new haplotypes.

Figure 3: The oriented graph of the ρST values between sites and cohorts (vertices). Orange edges and boxes allow by comparison between pair of sites to understand the dynamics and directions of genetic spreading. Comparisons between blue boxes (adults) and green boxes (juveniles) between pairs of sites to understand the temporal genetic structure between sampled sites.

Figure 4: Discriminant Analysis of Principal Components of the genetic structure of *Aulactinia stella* across the Arctic Ocean. We found four clusters being the minimal number of clusters maximizing the distance between groups.

Figure 5: Minimum spanning tree of the genetic differentiation between sampled sites and cohorts of *A.stella* using pairwise ρ_{ST}. Kamchatka adults and juveniles respectively in green and pale green; Kuril adults and juveniles respectively in dark blue and blue; Barents sea adults and juveniles respectively in gold and pale gold colors; White sea adults and juveniles respectively in pink and pale pink; Newfoundland adults and juveniles respectively in dark red and red.

Site	Age cohort	N	G	R	Paretoβ	$\bar{r}_d$	Na	Var(Na)	$\bar{A}_E$	Var(A_E)	$\bar{H}_O$	Var(H_O)	$\bar{H}_E$	Var(H_E)	$\overline{F_{IS}}$	Var(F_{IS})	pidu	pids
Kamchatka	**Adults**	6	6	1.000	1.807	3.0	30.0	2.2	17.3	17.27	0.294	0.488	0.416	0.590	0.292	1.635	1.4E-06	0.010
	Juveniles	49	48	0.979	4.615	5.2	151.6	2.4	24.3	24.25	0.314	0.397	0.460	0.534	0.318	1.156	0.0E+00	0.006
	Both	55	53	0.963	2.648	5.4	180.4	2.4	29.1	29.06	0.312	0.397	0.459	0.544	0.321	1.140	0.0E+00	0.006
Kuril Islands	**Adults**	19	19	1.000	3.322	4.6	84.4	2.5	46.9	46.90	0.176	0.121	0.438	0.640	0.598	1.016	0.0E+00	0.006
	Juveniles	30	30	1.000	3.954	5.7	234.1	2.5	53.1	53.09	0.188	0.301	0.372	0.891	0.494	0.948	0.0E+00	0.013
	Both	49	49	1.000	4.644	7.8	315.6	2.7	69.7	69.69	0.184	0.131	0.454	0.580	0.595	0.722	0.0E+00	0.005
Barents Sea	**Adults**	21	10	0.450	0.302	2.5	14.5	1.4	1.0	0.98	0.197	0.425	0.223	0.287	0.115	3.124	1.8E-03	0.096
	Juveniles	6	2	0.200	0.591	1.2	1.6	1.2	1.4	1.42	0.125	0.628	0.097	0.376	-0.290	13.323	1.3E-01	0.364
	Both	27	10	0.346	0.265	2.5	14.5	1.3	1.0	1.00	0.181	0.448	0.198	0.275	0.088	3.357	4.6E-03	0.126
Newfoundland	**Adults**	14	12	0.846	1.807	2.1	18.9	1.4	6.5	6.47	0.165	0.656	0.170	0.512	0.024	4.971	3.8E-03	0.151
	Juveniles	61	28	0.450	0.716	1.8	9.6	1.4	6.6	6.63	0.158	0.747	0.178	0.596	0.112	4.875	5.6E-03	0.140
	Both	75	40	0.527	0.799	2.2	17.6	1.5	6.3	6.35	0.160	0.694	0.210	0.551	0.241	3.604	2.0E-03	0.100
White Sea	**Adults**	15	5	0.286	0.129	1.9	18.9	1.6	7.8	7.81	0.314	1.526	0.238	0.886	-0.322	10.807	8.8E-04	0.069
	Juveniles	93	28	0.293	0.261	3.0	48.0	1.7	8.5	8.53	0.308	1.447	0.266	0.912	-0.157	7.841	1.9E-04	0.048
	Both	108	28	0.252	0.224	3.0	48.0	1.7	8.5	8.47	0.309	1.457	0.264	0.912	-0.171	8.048	2.2E-04	0.049
Over All	**Both**	314	180	0.572	0.482	11.1	468.9	2.1	6.2	6.24	0.243	0.387	0.440	0.454	0.448	0.870	2.0E-07	0.007

Table1: Summary of population genetic indices in ten spatio-temporal populations of *Aulactinia stella*. For each age cohort at one site, are reported the sample size N, number of genotypes G, genotypic richness R, genotypic evenness *Paretoß*, overall estimate of linkage disequilibrium $\bar{r}_d$, average number of observed alleles Na and its variance *Var(Na)* over the ten microsatellites, average effective number of alleles $\bar{A}_E$ and its variance $Var(A_E)$, average observed heterozygosity $\overline{H}_O$ and its variance $Var(H_O)$, average gene diversity $\overline{H}_E$ and its variance $Var(H_E)$, average value of the inbreeding coefficient $\overline{F_{IS}}$ and its variance $Var(F_{IS})$ over all loci, unbiased probability of identity under panmixia *pidu* and between sibs *pids*.

Supplementary information

Table S1: Distribution of repeated MLGs within sites and MLGs shared between adults and juveniles. Genet ID, the number of ramets within the genet and if the considered genet has been found in adults, juveniles or both are reported. No repeated MLG was found in Kuril islands sampled genotypes.

Site	Genet ID	Number of ramet	In the two cohorts?
Kamchatka	MLG123	3	Both
Barents	MLG32	2	Adults
Barents	MLG54	5	Both
Barents	MLG26	13	Both
Newfoundland	MLG12	2	Adults
Newfoundland	MLG7	2	Juveniles
Newfoundland	MLG37	2	Juveniles
Newfoundland	MLG67	2	Juveniles
Newfoundland	MLG11	2	Adults
Newfoundland	MLG5	2	Juveniles
Newfoundland	MLG19	2	Juveniles
Newfoundland	MLG6	3	Juveniles
Newfoundland	MLG62	3	Juveniles
Newfoundland	MLG52	3	Juveniles
Newfoundland	MLG63	4	Juveniles
Newfoundland	MLG68	4	Juveniles
Newfoundland	MLG53	7	Juveniles
Newfoundland	MLG57	11	Juveniles
White Sea	MLG47	2	Juveniles
White Sea	MLG15	3	Juveniles
White Sea	MLG66	3	Both
White Sea	MLG44	4	Both
White Sea	MLG22	4	Both
White Sea	MLG38	8	Both
White Sea	MLG16	24	Juveniles
White Sea	MLG46	40	Both

Table S2: Pairwise ρST between sites (A) and between age cohorts within sites and among sites (B).

A).

	Barents	Kamchatka	Kuril	Newfoundland	White sea
Barents	#				
Kamchatka	0.411	#			
Kuril	0.364	0.168	#		
Newfoundland	0.096	0.352	0.315	#	
White sea	0.442	0.348	0.521	0.167	#

B).

		Adults	Juveniles	Adults	Juveniles	Adults	Juveniles	Adults	Juveniles	Adults	Juveniles
		Barents		**Kamchatka**		**Kuril**		**Newfoundland**		**White sea**	
Barents	**Adults**	#									
	Juveniles	0.088	#								
Kamchatka	**Adults**	0.381	0.337	#							
	Juveniles	0.383	0.133	0.014	#						
Kuril	**Adults**	0.361	0.482	0.155	0.221	#					
	Juveniles	0.422	0.290	0.122	0.183	0.103	#				
Newfoundland	**Adults**	0.238	0.470	0.489	0.285	0.312	0.465	#			
	Juveniles	0.191	0.105	0.224	0.290	0.278	0.432	0.054	#		
White sea	**Adults**	0.505	0.971	0.359	0.279	0.371	0.387	0.612	0.058	#	
	Juveniles	0.386	0.199	0.155	0.352	0.459	0.530	0.372	0.118	0.009	#

Figure S1: Counting of allele dosage of the microsatellite locus 249TAM (as an example) in *Aulactinia stella* (within each individual per locus): A). 129, 129, 133, 133 (AABB), B). 129, 129, 137, 141 (AABC), C). 137, 137, 137, 141 (AAAB), D). 129, 133, 137 (ABC+null allele), E). 133, 133, 137 (AAB+null allele).

Figure S2: Minimum spanning tree of A). the ρST of adults, B). the ρST of juveniles and C). the ρST of all age cohorts over all loci between sampled sites.

A).

B).

C).

Figure S3: Change of the Bayesian information criteria (BIC) with increasing number of genetic clusters used in the Discriminant Analysis of Principal Components to investigate the genetic differentiation between individuals and among sampled sites and cohorts. The continuously decreasing pattern observed is typical of a stepping-stone structure of the studied demes.